Environmental pH signals the release of monosaccharides from cell wall in coral symbiotic alga

Yuu Ishii[1, 2], Hironori Ishii[1], Takeshi Kuroha[1]†, Ryusuke Yokoyama[1], Ryusaku Deguchi[2], Kazuhiko
Nishitani[3], Jun Minagawa[4,5], Masakado Kawata[1], Shunichi Takahashi[6], Shinichiro Maruyama[1,7]*

[1] Department of Ecological Developmental Adaptability Life Sciences, Graduate School of Life
Sciences, Tohoku University, Sendai, Miyagi, Japan

[2] Department of Biology, Miyagi University of Education, Sendai, Miyagi, Japan

[3] Department of Biological Sciences, Faculty of Science, Kanagawa University, Yokohama,
Kanagawa, Japan

[4] Division of Environmental Photobiology, National Institute for Basic Biology, Okazaki, Aichi,
Japan

[5] Department of Basic Biology, School of Life Science, SOKENDAI (The Graduate University for
Advanced Studies), Okazaki, Aichi, Japan

[6] Tropical Biosphere Research Center, University of the Ryukyus, Motobu, Okinawa, Japan

[7] Graduate School of Humanities and Sciences, Ochanomizu University, Bunkyo-ku, Tokyo,
Japan.

† Present address: Division of Crop Genome Editing, Institute of Agrobiological Sciences,
National Agriculture and Food Research Organization (NARO), Tsukuba, Ibaraki, Japan.

*Corresponding author: Shinichiro Maruyama

Email: maruyama@tohoku.ac.jp

Abstract

Reef-building corals thrive in oligotrophic environments due to their possession of endosymbiotic algae. Confined to the low pH interior of the symbiosome within the cell, the algal symbiont provides the coral host with photosynthetically fixed carbon. However, it remains unknown how carbon is released from the algal symbiont for uptake by the host. Here we show, using cultured symbiotic dinoflagellate, *Breviolum* sp., that decreases in pH directly accelerates the release of monosaccharides, i.e., glucose and galactose, into the ambient environment. Under low pH conditions, the cell surface structures were deformed and genes related to cellulase were significantly upregulated in *Breviolum*. Importantly, the release of monosaccharides was suppressed by the cellulase inhibitor, glucopyranoside, linking the release of carbon to degradation of the agal cell wall. Our results suggest that the low pH signals the cellulase-mediated release of monosaccharides from the algal cell wall as an environmental response in coral reef ecosystems.

Main text

Introduction

Coral reef ecosystems are sustained by symbiosis between stony corals and marine dinoflagellates from the family Symbiodiniaceae, which are found in nature as free-living mixotrophs (Decelle et al., 2018; Jeong et al., 2012), as well as are primary producers in symbiotic relationships with various partners, including multicellular (e.g. Cnidaria, Mollusca, Porifera) and unicellular organisms (Foraminifera, Ciliates) (LaJeunesse et al., 2018). In oligotrophic oceans, transfer of atmospheric carbon photosynthetically fixed by the symbiotic algae to their hosts is a fundamental flux to sustain the growth and productivity of coral reef ecosystems.

Although it is generally accepted that Symbiodiniaceae algae provide photosynthates to their symbiotic partners, the molecular details are largely unknown (Falkowski et al., 1984; Ishii et al., 2019; Ishikura et al., 1999; Muscatine, 1990; Rahav et al., 1989; Stat et al., 2008; Whitehead and Douglas, 2003). Members of this family reside in an intracellular organelle called the 'symbiosome' within cnidarian host cells or in the extracellular 'symbiotic tube' systems of giant clams. These are thought to be special low pH environments that are acidified by V-type H^+-ATPase proton pumps (Armstrong et al., 2018; Barott et al., 2015; Davy et al., 2012). While low pH environments are stressors to algae in general, they can be beneficial when CO_2 uptake is encouraged by the hosts' carbon-concentrating functions, enhancing photosynthesis (Armstrong et al., 2018; Barott et al., 2015). A previous study has demonstrated a photosynthesis-dependent glucose transfer from Symbiodiniaceae to sea anemone hosts (Burriesci et al., 2012), and some sugar transporters

are proposed to be involved in glucose transfer (Lehnert et al., 2014; Sproles et al., 2018). Other studies

suggest that the amount of transfer is regulated by the C-N balance (Rädecker et al., 2021; Xiang et al.,

2020). Nevertheless, the mechanism of algal glucose secretion is not yet characterized.

Results

To investigate the physiological effects of low pH, a characteristic environmental factor in symbioses, on

algal intrinsic properties, a Symbiodiniaceae alga *Breviolum* sp. SSB01 (hereafter, *Breviolum*) was grown

in a host-independent manner and cell proliferation and photosynthetic activities were measured. By

comparing the growth rate of *Breviolum* in normal culture medium (pH 7.8) and acidic medium (pH 5.5,

hereafter called "low pH"), we showed that the low pH medium considerably suppressed algal growth

(Figure 1A) and culturing at low pH for one day resulted in significant declines in photosynthesis activity

(Figure 1B).

Contrary to our expectation, the amount of glucose secreted into the culture medium was higher at low pH

(Figure 1C) and the secreted galactose similarly showed an increasing trend (Figure 1D). These trends

suggest that *Breviolum* is capable of secreting monosaccharides autonomously without host signals, and

that low pH enhanced the secretion. On the addition of the photosynthesis inhibitor 3-(3,4-dichlorophenyl)-

1,1-dimethylurea (DCMU), the concentrations of glucose and galactose in the medium increased

(Supplementary Figure S1), suggesting the presence of a yet uncharacterized pathway (Burriesci et al.,

2012).

To investigate the response of *Breviolum* to acidic environments at the morphological level, cells cultured

in different media were examined by microscopy. *Breviolum* cells in low pH media were more spread out

and less clustered than the cells in normal media (Figure 2A, B). Scanning electron microscope (SEM)

observations revealed that many of the *Breviolum* cells cultured at low pH exhibited wrinkled structures on

their cell surfaces (Figure 2C, D). Furthermore, transmission electron microscopy (TEM) revealed that the

cell surface structures of the low pH media group were 'exfoliated' (Figure 2E, F). These suggest that low

pH affects the structures and properties of a cellulosic cell wall found in coccoid Symbiodiniaceae cells

(Colley and Trench, 1983; Markell et al., 1992).

To identify the mechanism involved in the monosaccharide secretion of *Breviolum*, we compared gene

expression changes between the 'control vs normal' and 'control vs low pH' comparisons, and identified 3

and 4527 differentially expressed genes (DEGs), respectively (Figure 3A, Supplementary Dataset S1). The

gene ontology (GO) term enrichment and KEGG pathway analysis of these two gene sets resulted in the

detection of 0 (control vs. normal) and 16 (control vs. low pH) terms (Supplementary Dataset S2), which

included categories related to carbon metabolism (Figure 3B, Supplementary Figure S2). The CAZy

database (Lombard et al., 2014) analysis showed that 12 DEGs (28 isoforms) were annotated with

Carbohydrate-Active enZymes (CAZymes) activity (Figure 3C, Supplementary Dataset S3). One of the

genes, TRINITY_DN40554_c2_g2, was shown to encode cellulase (exocrine cellulolytic enzyme)

harbouring a signal peptide (Figure 3C) with high similarity to dinoflagellate homologs (Kwok and Wong,

2010) (Supplementary Figure S3).

To confirm the effect of cellulase on monosaccharide secretion, we examined whether secretions were

inhibited by the cellulase inhibitor Para-nitrophenyl 1-thio-beta-d-glucopyranosid (PSG) (Figure 4). PSG

inhibited the secretion of both glucose and galactose in a dose-dependent manner, suggesting that cell wall

degradation by cellulase is involved in the secretion of monosaccharides from *Breviolum* cells.

Discussion

The transfer of photosynthetically fixed carbon from symbiotic algae to host cnidarians, including corals, is

a cornerstone of their mutual symbiotic relationship (Muscatine, 1990). Unlike the current accepted model

of monosaccharide secretion that assumes photosynthetically fixed carbon is directly exported from the

algal symbiont via unidentified glucose transporter(s) (Lehnert et al., 2014; Sproles et al., 2018), our results

suggest that stored carbon can be released from the algal cell wall as an environmental response. In the

present study, we showed that a decrease in ambient pH, consistent with the interior pH of the symbiosome,

accelerates the monosaccharide secretion from *Breviolum* (Figure 1). Importantly, this low pH-associated

secretion was suppressed by inhibition of cellulase (Figure 4), suggesting that algal symbionts release

monosaccharides into the symbiosome within host cnidarian cells by cell wall degradation. Previous studies

showed that cell wall degradation/rearrangement by cellulase is required for cell cycle progression (Kwok

and Wong, 2010) and cellulose synthesis is involved in morphogenesis (Chan et al., 2019) in

dinoflagellates. Indeed, wrinkle and exfoliation of the algal cell wall was observed under low pH conditions

using SEM and TEM, respectively (Figure 2). We need to note that our results do not deny the current

accepted model, but rather suggest a multi-pathway hypothesis supported by the following observations: (i)

the secretion of monosaccharides was not completely inhibited by cellulase inhibition, (ii) this new

pathway occurred over days, compared to previous reports, where exported glucose was detectable in the

host after only 30-60 minutes (Burriesci et al., 2012), (iii) in contrast to previous studies (Burriesci et al.,

2012), DCMU did not suppress, but rather increased the secretion of monosaccharides over a longer time

span (Supplementary Figure S1). Importantly, low pH upregulated the expression of genes associated with

not only cellulase (Figure 3) but glycolysis probably fuelled by degradation of storage compound like

starch (Supplementary Figure S2). This suggests the photosynthesis-limiting conditions trigger an

environmental response in the algae to sustain cell proliferation, through cell cycle progression by cellulase,

accompanying cell wall degradation and monosaccharide secretion.

Like Symbiodiniaceae, some freshwater green algae are known to be symbiotic with a range of hosts. A

number of *Chlorella* strains, with and without symbiotic ability, autonomously secrete monosaccharides

under low pH conditions via unknown mechanisms (Kessler et al., 1991; Mews and Smith, 1982). The

monosaccharides include maltose and, to a lesser extent, glucose (Arriola et al., 2018). Dinoflagellates are

also known to secrete viscous substances, including monosaccharides, as an environmental response, likely

for cell aggregation and biofilm formation (Mandal et al., 2011). Thus, secretion of monosaccharides

appears to be a fairly conserved environmental response and may be to compensate for retarded cell cycle

progression. Therefore, acidifying symbiosomes may be an evolutionary successful strategy for cnidarian

hosts to promote environmental responses of algal symbionts, which enables monosaccharides to be

efficiently secreted within the organelle.

Generally, within ecosystems energy is transferred from photosynthetic primary producer to consumer by

predation. Uniquely, in coral reef ecosystems energy is mainly transported from algae to corals after

establishing a symbiotic relationship (Davy et al., 2012). Thus, understanding its mechanism has wider

implications to understanding how energy is shared over the entire coral reef ecosystem. The multi-

pathway hypothesis we propose here entails the direct transfer of photosynthates via glucose transporter(s)

on their cell membrane (Lehnert et al., 2014; Sproles et al., 2018) as well as monosaccharide secretion

following cell wall degradation. It remains to be determined how much each pathway contributes to the

energy supply of host. However, since one pathway uses de novo photosynthates and the other uses stored

photosynthates, combined they might allow for a stable supply of energy to the host, e.g., over the entire

light/dark day cycle, and under photosynthesis-limiting conditions like environmental stress or cloudy days

where the cellulose-related pathway could be of substantial importance. Although genetic transformation

for Symbiodiniaceae is unavailable yet (Chen et al., 2019), knock-down of the cellulase gene will bring a

clue to test this. Overall, our study provides a new insight into how carbon is provided by symbiotic algae

to the coral reef ecosystem.

Materials and Methods

Strains and culture conditions

We obtained the *Breviolum* (formerly *Symbiodinium* clade B) strain SSB01, an axenic uni-algal strain

closely related to the genome-sequenced strain *B. minutum* Mf1.05b (clade B), as a generous gift from

Profs. John R. Pringle and Arthur R. Grossman (Shoguchi et al., 2013; Xiang et al., 2013). The *Breviolum*

was maintained according to previous study (Ishii et al., 2018). Stock cultures were incubated at 25°C in

medium containing 33.5 g/L of Marine Broth (MB) (Difco Laboratories, New Jersey, USA), 250 mg/L of

Daigo's IMK Medium (Nihon Pharmaceutical, Japan), and PSN (Gibco, Thermo Fisher Scientific,

Massachusetts, USA), with final concentrations of penicillin, streptomycin, and neomycin at 0.01, 0.01, and

0.02 mg/mL, respectively. Light was provided at an irradiance of approximately 100 µmol photons/m^{2}s in a

12 h light:12 h dark cycle. In experiments, IMK medium containing 33.5 g/L of sea salt (Sigma-Aldrich,

Merck Millipore, Germany), 250 mg/L of Daigo's IMK Medium, and PSN with final concentrations of

penicillin, streptomycin, and neomycin at 0.01, 0.01, and 0.02 mg/mL, respectively, was used. The pH of

this medium was adjusted to 5.5 using HCl for the low pH experiments.

Sample preparation for Ion chromatography, RNAseq, and electron microscopy

The incubation conditions were designed to allow detection of changes in the early phases of low pH

stimulus responses. Prior to exposure to low pH conditions, *Breviolum* cultures were incubated for one

week in IMK medium at densities of 8×10^6 cells/20 mL in T25 culture flasks with a filter cap (TrueLine

Cell Culture Flasks, TR6000) under a light-dark cycle. During experiments, *Breviolum* was cultured in two

kinds of media: normal (pH 7.8) and low pH (pH 5.5), for 24 h (12 h light:12 h dark), starting at densities

of 8×10^6 cells/20 mL per T25 culture flask, placed at randomized positions. DCMU was dissolved in

ethanol at the concentration of 20 mM and was added to cultures to a final concentration of 20 µM. The

cultured cells were collected by centrifugation at 2,000 × g for 5 min at room temperature. The cells were

then used for RNA-seq and electron microscopy, and the supernatants were used for ion chromatographic

determination.

Ion chromatography

Samples (n = 4 biological replicates) of the supernatant from the control (0 h), normal (24 h) and low pH

(24 h) cultures were filtered using a 0.22 µm PVDF filter (Merck Millipore, Germany). These samples were

loaded onto an OnGuard column (Dionex OnGuard II Cartridge) (Thermo Scientific) to remove the

sulphate and halogen, according to the manufacturer's instructions. The samples were quantified using

high-performance anion-exchange chromatography with pulsed amperometric detection (HPAEC-PAD)

using a Dionex ICS-5000 system equipped with a CarboPac PA1 column (Dionex) (Shinohara et al., 2017).

The column was operated at a flow rate of 1.1 mL/min with the following phases: (1) a linear gradient of 0–

100 mM NaOH from 0 to 31 min, (2) a linear gradient of 0–150 mM sodium acetate containing 100 mM

NaOH from 31 to 34 min, and (3) an isocratic 150 mM sodium acetate/100 mM NaOH from 34 to 41 min.

Myo-inositol (2 µg/mL) was added to each sample as an internal standard for quantification.

The concentrations of monosaccharides were calculated by comparing the peak ratios between the targets

of interest and standards. The secreted amounts were calculated by subtracting the concentrations on day 0

from those of day 1.

Growth rate, photosynthesis, and respiratory activity assay

To measure growth rate (n = 6 biological replicates), photosynthesis and respiratory activities (n = 4 biological replicates), *Breviolum* cultures were incubated in normal IMK media for one week and subsequently changed to fresh normal or low pH IMK media. Growth rate comparisons between the normal and low pH media conditions were conducted using 100 µL of media (625 cells/µL) in a 96-well plate. Cell growth was monitored by measuring the optical density at 730 nm (OD730) of the liquid cultures using a Multiskan GO microplate spectrophotometer (Thermo Fisher Scientific, Massachusetts, USA). Photosynthesis and respiratory activities were determined using cultures at densities of 1×10^6 cells/mL in fresh normal and low pH media, on days 0 and 1 after changing the medium, separate from the transcriptome analysis. Respiration rates were calculated using the dark-phase oxygen consumption rates and photosynthesis rates were calculated by subtracting the respiration rates from the light-phase oxygen evolution rates. Mean estimates with standard errors were calculated from single measurements of four different cultures per medium condition.

RNA extraction and sequencing

Four samples (n = 4 biological replicates) from each of the control (day 0), normal (day 1), or low pH (day 1) cultures were added to 500 µL of TRIZOL reagent (Thermo Fisher Scientific, Massachusetts, USA) and stored at -80 °C. The samples were ground with two sizes of glass beads (20 µL volume each of '≤106 μm' and '425–600 μm') (Sigma-Aldrich, Merck Millipore, Germany) using a vortex and performing 5 cycles of freezing and thawing with a -80˚C freezer. RNA extraction with TRIZOL reagent and a high salt solution for precipitation (plant) (Takara Bio, Japan) was conducted according to the manufacturer's instructions. The quality and quantity of the RNA was verified using an Agilent RNA 6000 Nano Kit on an Agilent Bioanalyzer (Agilent Technologies, California, USA) and a Nanodrop spectrophotometer (Thermo Fisher Scientific, Massachusetts, USA), respectively. Total RNA samples were subjected to library preparation using an NEB Next Ultra RNA Library Prep Kit (New England Biolabs, Ipswich, MA, USA) according to the manufacturer's protocol (NEB #E7530). These mRNA libraries were sequenced in an Illumina NovaSeq6000 (S2 flow cell) in dual flow cell mode with 150-mer paired-end sequences (Filgen Inc., Japan). The raw read data were submitted to DDBJ/EMBL-EBI/GenBank under the BioProject accession number PRJDB12295.

Transcriptome analysis

A total of 12 libraries were obtained, trimmed, and filtered using the trimmomatic option of the Trinity program. Paired output reads were used for analysis, and de novo assembly was performed using the Trinity program (Grabherr et al., 2011) to obtain the transcript sequences. The reads from each library were mapped onto the de novo assembly sequences and read count data, and the transcripts per million (TPM) were calculated using RSEM (Li and Dewey, 2011) with bowtie2 (Langmead and Salzberg, 2012).

In this study, RNA-seq produced 550,711,174 reads from the 12 samples (four independent culture flasks

under three conditions), yielding 239,047 contigs by de novo assembly using Trinity. The total number of

mapped reads of the quadruplicates in the de novo assembled transcriptome dataset were 77,160,394 reads

for samples taken before medium change (labelled as 'control'), 74,594,446 for the normal pH culture

condition (labelled as 'normal'), and 73,882,605 for the low pH culture condition (labelled as 'lowpH').

Overall, we obtained the count values of the genes in the transcriptome dataset under the three conditions.

Differential gene expression analysis was conducted using the count data as inputs for the R package TCC

(Sun et al., 2013) to compare the tag count data with robust normalization strategies, with an option using

edgeR (Robinson et al., 2010) to detect differential expressions implemented in TCC. To identify the

DEGs, a false discovery rate (FDR, or q-value) of 0.01, was used as the cutoff.

To annotate the de novo transcript sequences, BLASTp search was performed (E-value cutoff, 10^{-4}) against

the GenBank *nr* database using all the transcript sequences as queries, resulting in 51,833 orthologs. Gene

ontology (GO) term annotation of the de novo transcript sequences was performed using InterProScan

(Jones et al., 2014) ver 5.42-78.0, resulting in 7,336 genes with GO terms. GO term enrichment analysis

was performed using the GOseq (Young et al., 2010) package in R. Overrepresented p-values produced by

GOseq were adjusted using the Benjamini-Hochberg correction (Benjamini and Hochberg, 1995). An

adjusted p-value (q-value) of 0.05, was used to define enriched GO terms. In the KEGG pathway analysis,

the ortholog protein sequence obtained via BLASTp search of the DEGs was used as a query. Additionally,

KOID was added by blastKOALA (Kanehisa et al., 2016) (https://www.kegg.jp/blastkoala/) and mapped to

the KEGG pathway using KEGmappar (Kanehisa and Sato, 2020)

(https://www.genome.jp/kegg/mapper.html). For CAZome analysis, all isoform sequences of the DEGs

were analysed using CAZy (Lombard et al., 2014) (http://www.cazy.org/).

Cellulase inhibition experiment

Breviolum cells cultured in IMK for more than 1 week were transferred to 24 well plates at 4×10^6 cells/ml

density and incubated for 24 h (12 h light:12 h dark) in IMK medium containing 0, 0.1, and 1 mM PSG (n

= 4 biological replicates for each condition). The supernatant from each culture was collected following

centrifugation at $2,000 \times g$ for 2 min at room temperature. The medium samples were filtered using a 0.22

μm PVDF filter (Merck Millipore, Germany).

Glucose and galactose were quantified using a LC–MS/MS system in which a Shimadzu UPLC system

(Shimadzu, Kyoto, Japan) was interfaced to an AB Sciex QTrap 5500 mass spectrometer equipped with an

electrospray ionization source (AB SCIEX, Foster, CA, USA). A UK-Amino column (3 μm, 2.1 mm × 250

mm, Imtakt Corporation, Kyoto, Japan) was applied for analysis. Mobile phase A is 0.1% formic acid, and

mobile phase B is acetonitrile. Samples (1 μl) were injected and analyzed over a gradient of: 0–0.5 min

95% buffer B (isocratic); 0.5–10 min 85% buffer B (linearly decreasing); 10–15 min 40% buffer B (linear

decreasing). The column was equilibrated for 5 min before each sample injection. The flow rate was 0.3

ml/min. Under these analytical conditions, the retention times for glucose and galactose were 11.6 and 10.9

minutes, respectively. Mass spectrometric analysis employed electrospray ionization in the negative mode

with multiple reaction monitoring (MRM) at the transitions of m/z 179→89 for glucose and galactose. The

optimized MS parameters were as follows: ion spray voltage (-4500V), declustering potential (-90V),

entrance potential (-10V), collision energy (-12V), collision exit potential (-11V), collision gas (N2 gas)

and nebulizer temperature (450 ∘C). Raw data was analyzed using MultiQuant software (AB SCIEX,

Foster, CA, USA). Concentrations of monosaccharides were calculated by comparing the peak ratios

between the targets of interest and standards.

290 Microscopy

To compare the cell conditions between the normal and low pH media, *Breviolum* was cultured and

incubated for 3 weeks (12 h light:12 h dark) starting at densities of 1.6×10^7 cells/20 mL per T25 culture

flask. Cell photos were taken using a TC20 automated cell counter (Bio-Rad Laboratories, Hercules, CA),

and the numbers of cells adjacent to and isolated from other cells were randomly counted (853 and 664

cells pooled form n = 3 biological replicates were scored in the normal and low pH conditions,

respectively). The cells were fixed in a DMSO:KOH (1:1) solution, stained with 10 mg/mL calcofluor

white solution in Milli-Q water, and observed under an inverted microscope (Leica DMI6000 B).

For SEM, cells were fixed in 2% glutaraldehyde and 2% osmium (VIII) oxide, dehydrated with

ethanol, and dried using the critical point drying technique. The samples were coated with osmium plasma

and observed under a JSM-7500F microscope at 5 kV. The surface patterns of the cells were manually

scored and classified as 'smooth' or 'wrinkled' (73 and 129 cells pooled form n = 4 technical replicates

were blindly scored under the normal and low pH conditions, respectively). For TEM observation, the

samples were dehydrated with ethanol and embedded in EPON812 polymerized with epoxy resin. Sections

80–90 nm thick were cut, coated with evaporated carbon for stabilisation, and stained with uranyl acetate

and lead citrate. The sections were then imaged at 100 kV using a HITACHI H-7600 transmission electron

microscope. The cells were then categorized as NP (non-peeled), P (peeled) or CP (completely-peeled)

(181 and 166 cells pooled form n = 10 technical replicates were blindly scored under normal and low pH

conditions, respectively).

309
310

311 Acknowledgments
312
We thank Drs. Naoki Shinohara and Fukumatsu Iwahashi for their support in ion chromatography and mass

spectrometry, Ms Yuna Uchida for her assistance in maintaining the algal cultures, Dr. Sara E. Milward for

her critical reading of the manuscript, Prof. Hiromu Tanimoto and Dr. Vladimiros Thoma for their help in

preparing the manuscript, Drs. Kota Kera and Seiji Kojima for their contributions in the initial phase of the

project, as well as Profs. John R. Pringle and Arthur R. Grossman for their generous gift of algal cultures.

This work was supported by JSPS KAKENHI (Grant Numbers JP20J01658 [to Y.I.], JP17K15163,

JP19H04713 and JP19K06786 [to S.M.]), NIBB Collaborative Research Program 18-321 and 19-332 (to

S.M.), Institute for Fermentation, Osaka (to S.M.), Program for Creation of Interdisciplinary Research,

Frontier Research Institute for Interdisciplinary Sciences, Tohoku University (to S.M.), and the Gordon &

Betty Moore Foundation's Marine Microbiology Initiative #4985 (to J.M.). Computational resources were

provided by the Data Integration and Analysis Facility at the National Institute for Basic Biology and the

NIG supercomputer at the ROIS National Institute of Genetics.

325
326

327   **Competing Interests**

328   Authors declare that they have no competing interests.

330
331   **References**
332

333   Armstrong EJ, Roa JN, Stillman JH, Tresguerres M. 2018. Symbiont photosynthesis in giant clams is
334   promoted by V-type H+-ATPase from host cells. *J Exp Biol* **221**. doi:10.1242/jeb.177220
335   Arriola MB, Velmurugan N, Zhang Y, Plunkett MH, Hondzo H, Barney BM. 2018. Genome sequences of
336   *Chlorella sorokiniana* UTEX 1602 and *Micractinium conductrix* SAG 241.80: implications to
337   maltose excretion by a green alga. *Plant J* **93**:566–586. doi:10.1111/tpj.13789
338   Barott KL, Venn AA, Perez SO, Tambutté S, Tresguerres M. 2015. Coral host cells acidify symbiotic algal
339   microenvironment to promote photosynthesis. *Proc Natl Acad Sci USA* **112**:607–612.
340   doi:10.1073/pnas.1413483112
341   Benjamini Y, Hochberg Y. 1995. Controlling the false discovery rate: A practical and powerful approach to
342   multiple testing. *J R Stat Soc, Ser B, Methodol (Online)* **57**:289–300. doi:10.1111/j.2517-
343   6161.1995.tb02031.x
344   Burriesci MS, Raab TK, Pringle JR. 2012. Evidence that glucose is the major transferred metabolite in
345   dinoflagellate-cnidarian symbiosis. *J Exp Biol* **215**:3467–3477. doi:10.1242/jeb.070946
346   Chan WS, Kwok ACM, Wong JTY. 2019. Knockdown of Dinoflagellate Cellulose Synthase *CesA1*
347   Resulted in Malformed Intracellular Cellulosic Thecal Plates and Severely Impeded Cyst-to-
348   Swarmer Transition. *Front Microbiol* **10**. doi:10.3389/fmicb.2019.00546
349   Chen JE, Barbrook AC, Cui G, Howe CJ, Aranda M. 2019. The genetic intractability of *Symbiodinium*
350   *microadriaticum* to standard algal transformation methods. *PLOS ONE* **14**:e0211936.
351   doi:10.1371/journal.pone.0211936
352   Colley NJ, Trench RK. 1983. Selectivity in phagocytosis and persistence of symbiotic algae by the
353   scyphistoma stage of the jellyfish *Cassiopeia xamachana*. *Proceedings of the Royal Society of
354   London Series B Biological Sciences* **219**:61–82. doi:10.1098/rspb.1983.0059
355   Davy SK, Allemand D, Weis VM. 2012. Cell biology of cnidarian-dinoflagellate symbiosis. *Microbiol Mol
356   Biol Rev* **76**:229–261. doi:10.1128/MMBR.05014-11
357   Decelle J, Carradec Q, Pochon X, Henry N, Romac S, Mahé F, Dunthorn M, Kourlaiev A, Voolstra CR,
358   Wincker P, de Vargas C. 2018. Worldwide occurrence and activity of the reef-building coral
359   symbiont *Symbiodinium* in the open ocean. *Curr Biol* **28**:3625-3633.e3.
360   doi:10.1016/j.cub.2018.09.024
361   Falkowski PG, Dubinsky Z, Muscatine L, Porter JW. 1984. Light and the bioenergetics of a symbiotic
362   coral. *Bioscience* **34**:705–709. doi:10.2307/1309663

Grabherr MG, Haas BJ, Yassour M, Levin JZ, Thompson DA, Amit I, Adiconis X, Fan L, Raychowdhury
R, Zeng Q, Chen Z, Mauceli E, Hacohen N, Gnirke A, Rhind N, di Palma F, Birren BW, Nusbaum
C, Lindblad-Toh K, Friedman N, Regev A. 2011. Full-length transcriptome assembly from RNA-
Seq data without a reference genome. *Nat Biotechnol* **29**:644–652. doi:10.1038/nbt.1883
Ishii Y, Maruyama S, Fujimura-Kamada K, Kutsuna N, Takahashi S, Kawata M, Minagawa J. 2018.
Isolation of uracil auxotroph mutants of coral symbiont alga for symbiosis studies. *Sci Rep* **8**:3237.
doi:10.1038/s41598-018-21499-3
Ishii Y, Maruyama S, Takahashi H, Aihara Y, Yamaguchi T, Yamaguchi K, Shigenobu S, Kawata M, Ueno
N, Minagawa J. 2019. Global shifts in gene expression profiles accompanied with environmental
changes in cnidarian-dinoflagellate endosymbiosis. *G3-GENES GENOM GENET* **9**:2337–2347.
doi:10.1534/g3.118.201012
Ishikura M, Adachi K, Maruyama T. 1999. Zooxanthellae release glucose in the tissue of a giant clam,
Tridacna crocea. Mar Biol **133**:665–673. doi:10.1007/s002270050507
Jeong HJ, Yoo YD, Kang NS, Lim AS, Seong KA, Lee SY, Lee MJ, Lee KH, Kim HS, Shin W, Nam SW,
Yih W, Lee K. 2012. Heterotrophic feeding as a newly identified survival strategy of the
dinoflagellate *Symbiodinium. Proc Natl Acad Sci USA* **109**:12604–12609.
doi:10.1073/pnas.1204302109
Jones P, Binns D, Chang H-Y, Fraser M, Li W, McAnulla C, McWilliam H, Maslen J, Mitchell A, Nuka G,
Pesseat S, Quinn AF, Sangrador-Vegas A, Scheremetjew M, Yong S-Y, Lopez R, Hunter S. 2014.
InterProScan 5: genome-scale protein function classification. *Bioinformatics* **30**:1236–1240.
doi:10.1093/bioinformatics/btu031
Kanehisa M, Sato Y. 2020. KEGG Mapper for inferring cellular functions from protein sequences. *Protein*
Sci **29**:28–35. doi:10.1002/pro.3711
Kanehisa M, Sato Y, Morishima K. 2016. BlastKOALA and GhostKOALA: KEGG tools for functional
characterization of genome and metagenome sequences. *J Mol Biol* **428**:726–731.
doi:10.1016/j.jmb.2015.11.006
Kessler E, Kauer G, Rahat M. 1991. Excretion of sugars by *Chlorella* species capable and incapable of
symbiosis with *Hydra viridis. Bot Acta* **104**:58–63. doi:10.1111/j.1438-8677.1991.tb00194.x
Kwok ACM, Wong JTY. 2010. The activity of a wall-bound cellulase is required for and is coupled to cell
cycle progression in the dinoflagellate *Crypthecodinium cohnii. Plant Cell* **22**:1281–1298.
doi:10.1105/tpc.109.070243
LaJeunesse TC, Parkinson JE, Gabrielson PW, Jeong HJ, Reimer JD, Voolstra CR, Santos SR. 2018.
Systematic revision of Symbiodiniaceae highlights the antiquity and diversity of coral
endosymbionts. *Curr Biol* **28**:2570-2580.e6. doi:10.1016/j.cub.2018.07.008
Langmead B, Salzberg SL. 2012. Fast gapped-read alignment with Bowtie 2. *Nat Methods* **9**:357–359.
doi:10.1038/nmeth.1923
Lehnert EM, Mouchka ME, Burriesci MS, Gallo ND, Schwarz JA, Pringle JR. 2014. Extensive differences
in gene expression between symbiotic and aposymbiotic cnidarians. *G3-GENES GENOM GENET*
4:277–295. doi:10.1534/g3.113.009084
Li B, Dewey CN. 2011. RSEM: accurate transcript quantification from RNA-Seq data with or without a
reference genome. *BMC Bioinformatics* **12**:323. doi:10.1186/1471-2105-12-323
Lombard V, Golaconda Ramulu H, Drula E, Coutinho PM, Henrissat B. 2014. The carbohydrate-active
enzymes database (CAZy) in 2013. *Nucleic Acids Res* **42**:D490-495. doi:10.1093/nar/gkt1178
Mandal SK, Singh RP, Patel V. 2011. Isolation and characterization of exopolysaccharide secreted by a
toxic dinoflagellate, *Amphidinium carterae* Hulburt 1957 and its probable role in harmful algal
blooms (HABs). *Microb Ecol* **62**:518–527. doi:10.1007/s00248-011-9852-5
Markell D, Trench RK, Iglesias-Prieto R. 1992. Macromolecules associated with the cell walls of symbiotic
dinoflagellates. *Symbiosis* **12**:19–31.
Mews LK, Smith DC. 1982. The green hydra symbiosis. VI. What is the role of maltose transfer from alga
to animal? *Proc Royal Soc B* **216**:397–413. doi:10.1098/rspb.1982.0083
Muscatine L. 1990. The role of symbiotic algae in carbon and energy flux in reef corals. *Coral Reefs*
25:75–87.

Rädecker N, Pogoreutz C, Gegner HM, Cárdenas A, Roth F, Bougoure J, Guagliardo P, Wild C, Pernice M,
Raina J-B, Meibom A, Voolstra CR. 2021. Heat stress destabilizes symbiotic nutrient cycling in
corals. *Proc Natl Acad Sci USA* **118**:e2022653118. doi:10.1073/pnas.2022653118
Rahav O, Dubinsky Z, Achituv Y, Falkowski PG. 1989. Ammonium metabolism in the zooxanthellate
coral, *Stylophora pistillata*. *Proc Royal Soc B* **236**:325–337. doi:10.1098/rspb.1989.0026
Robinson MD, McCarthy DJ, Smyth GK. 2010. edgeR: a Bioconductor package for differential expression
analysis of digital gene expression data. *Bioinformatics* **26**:139–140.
doi:10.1093/bioinformatics/btp616
Shinohara N, Sunagawa N, Tamura S, Yokoyama R, Ueda M, Igarashi K, Nishitani K. 2017. The plant
cell-wall enzyme AtXTH3 catalyses covalent cross-linking between cellulose and cello-
oligosaccharide. *Sci Rep* **7**:46099. doi:10.1038/srep46099
Shoguchi E, Shinzato C, Kawashima T, Gyoja F, Mungpakdee S, Koyanagi R, Takeuchi T, Hisata K,
Tanaka M, Fujiwara M, Hamada M, Seidi A, Fujie M, Usami T, Goto H, Yamasaki S, Arakaki N,
Suzuki Y, Sugano S, Toyoda A, Kuroki Y, Fujiyama A, Medina M, Coffroth MA, Bhattacharya
D, Satoh N. 2013. Draft assembly of the *Symbiodinium minutum* nuclear genome reveals
dinoflagellate gene structure. *Curr Biol* **23**:1399–1408. doi:10.1016/j.cub.2013.05.062
Sproles AE, Kirk NL, Kitchen SA, Oakley CA, Grossman AR, Weis VM, Davy SK. 2018. Phylogenetic
characterization of transporter proteins in the cnidarian-dinoflagellate symbiosis. *Mol Phylogenet*
Evol **120**:307–320. doi:10.1016/j.ympev.2017.12.007
Stat M, Morris E, Gates RD. 2008. Functional diversity in coral-dinoflagellate symbiosis. *Proc Natl Acad*
Sci USA **105**:9256–9261. doi:10.1073/pnas.0801328105
Sun J, Nishiyama T, Shimizu K, Kadota K. 2013. TCC: an R package for comparing tag count data with
robust normalization strategies. *BMC Bioinform* **14**:219. doi:10.1186/1471-2105-14-219
Whitehead LF, Douglas AE. 2003. Metabolite comparisons and the identity of nutrients translocated from
symbiotic algae to an animal host. *J Exp Biol* **206**:3149–3157. doi:10.1242/jeb.00539
Xiang T, Hambleton EA, DeNofrio JC, Pringle JR, Grossman AR. 2013. Isolation of clonal axenic strains
of the symbiotic dinoflagellate *Symbiodinium* and their growth and host specificity. *J Phycol*
49:447–458. doi:10.1111/jpy.12055
Xiang T, Lehnert E, Jinkerson RE, Clowez S, Kim RG, DeNofrio JC, Pringle JR, Grossman AR. 2020.
Symbiont population control by host-symbiont metabolic interaction in Symbiodiniaceae-
cnidarian associations. *Nat Commun* **11**:108. doi:10.1038/s41467-019-13963-z
Young MD, Wakefield MJ, Smyth GK, Oshlack A. 2010. Gene ontology analysis for RNA-seq: accounting
for selection bias. *Genome Biol* **11**:R14. doi:10.1186/gb-2010-11-2-r14
448
449

List of figure supplements and source data files

Figure 1-figure supplement 1. Effect of photosynthesis inhibitor on glucose and galactose secretion.

Figure 1- source data 1. Raw data of growth rates.

Figure 1- source data 2. Raw data of photosynthesis activities.

Figure 1- source data 3. Raw data of glucose and galactose concentrations.

Figure 2- source data 1. Raw data of bright field cell counts.

Figure 2- source data 2. Raw data of SEM cell counts.

Figure 2- source data 3. Raw data of TEM cell counts.

Figure 3-figure supplement 1. Mapping of DEGs between the low pH and control groups on KEGG
pathways.

Figure 3-figure supplement 2. Phylogenetic tree of the cellulase proteins.

Figure 3-source data 1. Expression levels of the annotated low pH DEGs.

Figure 3-source data 2. GO enrichment analysis results.

Figure 3-source data 3. CAZome analysis results.

Figure 3-source data 4. Multiple alignment of the cellulase proteins.

Figure 4- source data 1. Raw data of glucose and galactose concentrations with PSG.

Figure 1. Physiological characterisation and monosaccharide secretion of cultured Breviolum. A. Growth rate (n = 6 per treatment, t-test). Asterisks indicate statistically significant differences (t-test, p < 0.005). B. Photosynthesis activity (n = 4 per treatment, t-test) C, D. Glucose (C) and galactose (D) concentrations secreted in normal and low pH media after 1 day incubation (n = 4 per treatment, t-test).

Figure 2. Cell structures under different pH condition. A. Bright field images of the cells under different conditions. The lower panels show high-magnification views of boxed areas in the upper panels. Scale bar = 50 μm. B. Quantification of the number of cells forming clusters (Fisher's exact test for "1 or 2" vs "3 or more", p = 1.727 × 10-07). C. SEM images of the representative cells. Scale bar = 1 μm. D. Quantification of the cell surface structures of the SEM images (Fisher's exact test, p < 2.2 × 10-16). E. TEM images of the representative cells. Scale bar = 2 μm. F. Quantification of the cell surface structures of the TEM images (Fisher's exact test for "P or CP" vs "NP", p = 4.621 × 10-11).

Figure 3. Differentially expressed genes under different pH conditions. A. Venn diagram showing the numbers of DEGs under different conditions. B. GO term enrichment analysis. Circles indicate the statistical significance (FDR) of the enriched GO terms, with the numbers of DEGs (numDEG) associated with each GO term. C. Isoform-level expression analysis of genes encoding CAZymes. Symbols indicate the statistical significance (FDR) of the differentially expressed isoforms associated with the DEGs, with the presence (triangle) or absence (rhombus) of signal peptide predicted in the amino acid sequence. Symbol colors represent the log fold-changes (logFC) of the expression levels of each isoform (low pH/control). Asterisks and dashed line indicate isoforms of the cellulase gene and a threshold for differential expression (FDR = 0.01), respectively.

Figure 4. Glucose and galactose secretion in cellulase inhibitor treatment. The concentrations of glucose (A) and galactose (B) in the medium after 1 day incubation with PSG (n = 3 per treatment, t-test).

Figure 1-figure supplement 1. Effect of photosynthesis inhibitor on glucose and galactose secretion

Glucose (A) and galactose (B) secreted after 1 day incubation with and without the addition of DCMU (n = 4 per treatment, t-test). Values of 'Normal' in A were obtained from the same data shown in Figure 1.

Figure 3-figure supplement 1. Mapping of DEGs between the low pH and control groups on KEGG pathways. The KEGG pathways for glycolysis/gluconeogenesis are shown. The colours indicate gene expression levels.

Figure 3-figure supplement 2. Phylogenetic tree of the cellulase proteins. A maximum likelihood tree of cellulase proteins labeled with species names and GenBank IDs is shown. Support values using an SH-like approximate likelihood ratio test (left) and ultrafast bootstrap approximation (right) are shown for each branch. For Breviolum sp. SSB01, the Trinity assembly contig IDs are shown.