The contrasted impacts of grasshoppers on soil microbial activities in function of primary production and herbivore diet

Sébastien Ibanez[1], Arnaud Foulquier[1], Charles Brun[1,2], Marie-Pascale Colace[1], Gabin Piton[1], Lionel
Bernard[1], Christiane Gallet[1], Jean-Christophe Clément[1,3]

1 Univ. Grenoble Alpes, Univ. Savoie Mont Blanc, CNRS UMR 5553, LECA, Chambéry, France

2 University of Applied Sciences and Arts Western Switzerland – Land, Nature, Environment
Institute, Hepia Geneva, Route de Presinge 150, CH-1254, Jussy, Switzerland

3 Univ. Savoie Mont Blanc, INRAE, CARRTEL, Thonon-Les-Bains, France

Abstract

Herbivory can have contrasted impacts on soil microbes and nutrient cycling, which has stimulated the development of conceptual frameworks exploring the links between below- and aboveground processes. The "productivity model" predicts that herbivores stimulate microbial activities and accelerate nutrient mineralization in productive ecosystems, while they have an opposite effect in less productive ecosystems. In parallel, the "diet model" predicts that herbivores feeding on conservative plants accelerate nutrient cycling while those feeding on exploitative plants decelerate nutrient cycling, due to changes in litter inputs. Since these two frameworks can lead to conflicting predictions in some cases, experimental evidence combining herbivore diet and plant productivity is required.

During two consecutive years, we conducted an experiment controlling the presence of three grasshopper species consuming either grasses, forbs or both in twelve natural and managed alpine grasslands with contrasted productivities. In order to assess the effects of herbivory on soil microbes, we measured their extracellular enzymatic activities, biomass and potential nitrogen mineralization (PNM). Soil and vegetation were also characterized to test how much they modulated the effects of herbivory on microbes.

Contrary to the predictions of the diet model, the effects of herbivory on microbial characteristics did not depend on the herbivores diet, but were influenced by primary production, though in a way that differed from the productivity model. The most productive sites were constituted by exploitative plant species which depleted N resources in the soil, and by microbes producing relatively few extracellular enzymes, leading to a lower PNM. Herbivory increased microbial biomass and decreased the production of extracellular enzymes in those sites, possibly through the stimulation of root exudates produced by exploitative species. The least productive sites were characterized by conservative plants, high soil C content, and by microbes having a resource acquisition strategy (more extracellular enzymes, higher PNM). Herbivory decreased microbial biomass and increased the production of extracellular enzymes in those sites. This pattern can be explained by the loss of carbon associated with insect respiration, which increases the resource requirements of microbes and by a lower production of root exudates by conservative species. Therefore, the effects of two years of herbivory on soil microbes were at odds with the productivity model, which focuses instead on longer term effects corresponding to herbivory-induced changes in plant species composition. This highlights the multidimensional feature of the impacts of herbivory on ecosystem functioning, both in space and time.

Introduction

During the last decades the influence of herbivory on terrestrial ecosystem functioning has been highlighted through its effects on matter and energy fluxes linking above and belowground communities (Parker et al. 2017; Kristensen et al. 2020; Sandén et al. 2020). Plant-herbivore interactions influence the quantity and quality of organic matter inputs to soil detrital food webs with important implications on the rate of microbial processes which regulate organic matter decomposition and nutrient recycling and ultimately control the maintenance of soil fertility and carbon sequestration (Bardgett and Wardle 2010). Previous studies have already reported a diversity of impacts of herbivory on soil carbon and nutrient cycling (Hunter 2001; Bakker et al. 2004), including positive (e.g. Frank et al. 2000; Belovsky and Slade 2000), negative (e.g. Ritchie et al. 1998) or no detectable effects (Singer and Schoenecker 2003). An overarching conceptual model of these contrasting effects is needed to understand which ecological variables control the direction and magnitude of the effects of herbivory on soil microbial communities and related ecosystem processes.

Among the different frameworks that have been proposed, one focuses on the contrasting effects of ecosystem productivity (Bardgett and Wardle 2003, 2010; Wardle et al. 2004), and one on the mitigation by herbivore diet (Ritchie et al. 1998; Belovsky and Slade 2000; Hunter 2001; Tuomi et al. 2019). The "productivity model" predicts that in productive ecosystems herbivores consume a high percentage of the net primary production, rapidly returning organic matter to the soil as easily decomposable fecal material enriched in nutrients ("fast cycle", McNaughton et al. 1988). Herbivores also promote compensatory plant growth (McNaughton 1983) or nutrient reallocation in leaf tissues of exploitative plant species (Potthast et al. 2021), which have high leaf nutrient content (low C/N) and grow faster. Herbivores also slow down the establishment of conservative plant species (high C/N, slow growing) which produce more recalcitrant litter (Reich 2014). The combination of these positive effects on the quality of detrital resources leads to an acceleration of nutrient cycling, which further induces a positive feedback loop. Instead, in less productive ecosystems herbivores consume a smaller proportion of the net primary production, favoring the accumulation of recalcitrant plant litter ("slow cycle", McNaughton et al. 1988). This comes along with the promotion of conservative plant species producing even more recalcitrant litter therefore promoting a slower nutrient cycling.

In contrast, the "diet model" focuses on another aspect of herbivore-plant-soil interactions and distinguishes two types of herbivores, those consuming exploitative plants, and those eating conservative plants. Indeed, although most vertebrate herbivores either feed non-selectively or prefer high-quality plants (Hofmann 1989; Clauss et al. 2003), many insects prefer tougher plants (Ibanez et al. 2013) and perform better on seemingly low-quality diets (Cease et al. 2012; Talal et al. 2020). Herbivores feeding on exploitative plants favor the growth and survival of conservative plants and therefore the accumulation of more recalcitrant litter. This slows down organic matter cycling, as predicted by the productivity model according to which herbivores preferentially feed on higher quality plants in the least productive ecosystems (Bardgett and Wardle 2010). Instead, herbivores feeding on conservative plants transfer low quality litter into the fast cycle, which accelerates decomposition and nutrient cycling, and promotes exploitative plant species (Belovsky and Slade 2000). During the last 20 years, several experiments controlling for herbivory either by ungulates or

insects provided evidence in favor of the diet model. When herbivores fed on the most conservative
plants, they increased primary production and/or nutrient availability (McNaughton et al. 1997;
Belovsky and Slade 2000, 2018; Garibaldi et al. 2007; Schmitz 2008; Nitschke et al. 2014). In
contrast, when herbivores fed on the most exploitative plants they had an opposite effect (Pastor et
al. 1993; Ritchie et al. 1998; van Wijnen et al. 1999; Harrison and Bardgett 2004; Schmitz 2008;
Belovsky and Slade 2018). The two experiments showing such combination of both effects used
polyphagous grasshopper species for which the diet changes either according to the type of
predators present (Schmitz 2008) or depending on intraspecific variation of leaf water content
(Belovsky and Slade 2018). However, these experiments as well as others (e.g. Deraison et al.
2015) manipulated either the presence or the diets of herbivores in a single type of ecosystem, which
hampers the articulation of the diet model with the productivity model.
104
Although the productivity and diet models focus on different aspects (ecosystem
productivity and herbivore diet), they can be combined. In some cases, the predictions of both
models are aligned (Table 1). However, in less productive ecosystems inhabited by herbivores
selecting low quality plants, nutrient cycling may be affected in both directions, depending on the
relative strength of the various processes at play in the productivity and diet models. Similarly,
herbivores feeding on high quality plants in fertile ecosystems may have contrasting effects on
matter turn-over (Table 1). For instance, under fertile conditions herbivores feeding on high quality
plants can favor their dominance (Buckland and Grime 2000), presumably because high resource
availability allows compensatory growth for these plant species. This may in turn promote nutrient
cycling and soil fertility (Bardgett and Wardle 2010). However, if high-quality plants invest in
defensive secondary compounds instead of displaying compensatory growth, and if the herbivores
can cope with them, slow-growing species may become dominant and nutrient cycling may slow
down.
118
Furthermore, while the productivity and diet models explicitly consider plant and herbivore resource
acquisition strategies, they do not consider the resource acquisition traits of soil microbes. Yet,
microbial communities are key regulators of nutrient recycling in the plant-soil system, and recent
developments suggest that including microbial communities in a multitrophic functional trait
framework could strengthen our mechanistic understanding of ecosystem functioning (Malik et al.
2020; Piton et al. 2020b). Resource acquisition strategies of microbial communities can be partly
inferred from the characterization of enzymatic activities involved in C, N and P acquisition through
the extracellular depolymerization of organic compounds (Piton et al. 2020b, a). A higher investment
in resource acquisition traits (i.e. increased extracellular enzyme activities) is expected along a
gradient of decreasing soil resource availability, paralleling a shift in plant resource acquisition
strategy along the exploitative-conservative continuum (Piton et al. 2020b). Increased quantity (i.e.
soil organic matter, N and P amounts) and quality (i.e. lower soil C/N, higher nitrate, ammonium
contents) of available resources from directly assimilable fecal material enriched in nutrients (e.g.
Fielding et al. 2013) or increased labile carbon inputs from root exudates (Hamilton and Frank 2001;
Paterson et al. 2003; Hamilton et al. 2008) are expected to promote microbial communities (Wardle
et al. 2004; Grigulis et al. 2013) with a high yield strategy (Y strategy, Malik et al. 2020) characterized
by a low investment in extracellular enzyme production (Piton et al. 2020b). On the contrary, a
reduction of available resource quality should promote microbial communities with a resource

acquisition strategy characterized by a high investment in extracellular enzymes targeting complex
polymeric organic matter (A strategy, Malik et al. 2020). Higher nitrogen availability in the form of
labile compounds (amino acids, NH_4^+ or NO_3^-) resulting from herbivory should reduce the microbial
demand for this element, resulting in a lower investment in N-acquiring enzymes. Yet, increased
availability of labile C from root exudates following herbivory might also increase nutrient limitation of
microbial communities, especially if herbivory stimulates plant productivity and associated nutrient
acquisition. This would result in higher nutrient immobilization, which is more likely under less fertile
conditions that offer low resource availability for microbes. Moreover, the productivity model focuses
on the contrasting effects of herbivory along a gradient of plant productivity, but soil nutrient cycling is
ultimately controlled by microbial communities, whose responses to herbivory may rather depend on
a gradient of soil resources, on which they rely more directly than on plant productivity. Both
gradients are often correlated, productive ecosystems being associated with low carbon content
(Wardle et al. 2004; Grigulis et al. 2013), but decoupling can also be observed (Piton et al. 2020b).
150
We therefore identified two important gaps in previous works regarding the effects of herbivory on
ecosystem functioning: first, the two existing frameworks should be combined (Table 1), and second,
the functional characteristics of soil microbial communities should explicitly be included within the
frameworks. This calls for experimental evidence controlling for herbivore diet in ecosystems with
contrasting plant productivity and soil resource levels, in order to test the effect of herbivory on
microbial communities. In the present contribution, we bridge these gaps with a semi-controlled
experiment using insect-proof cages, where four types of grasslands (managed or natural,
dominated by grasses or by forbs) in 12 montane sites were subjected to herbivory by three
grasshopper species which either preferred grasses, forbs or a mixture of the two. Since within each
site grasses had a higher C/N ratio than forbs (Ibanez et al. 2021), grass-feeders were expected to
accelerate matter cycling, while an opposite effect was predicted for forb-feeders. Following two
years of herbivory, activities of extracellular soil enzymes specialized in carbon, nitrogen and
phosphorus acquisition were measured, as well as nitrogen mineralization potential and microbial
biomass. We hypothesized that the effect of herbivory on these soil microbial characteristics
depends on herbivore diet and plant productivity. In cases where the two frameworks make opposite
predictions (Table 1), the experiment was designed to compare the relative importance of diet and
productivity effects.

Material & methods

Study sites and experimental design

The experiment was conducted in the French Pre-Alps, in Northern Vercors on the calcareous plateau of Autrans-Méaudre (45°10'N, 5°32'E). Four grassland types were chosen on the base of their botanical composition: extensively managed and fertilized grasslands dominated by the grass *Festuca rubra*, intensively managed and fertilized grasslands dominated by the grass *Lolium perenne*, natural grasslands dominated by forbs characterized by *Heracleum sphondylium* and *Chaerophyllum hirsutum* and natural warm grasslands mainly formed by the grass *Bromopsis erecta*. The 6 managed grasslands are disseminated on the Autrans plateau, and the 6 natural grasslands form two distinct clusters containing either 2 forbs-dominated communities and 1 warm grassland, or 1 forbs-dominated community and 2 warm grasslands (Sup. Fig. 1, site coordinates in Sup. Tab. 1). Since the two clusters contain communities having contrasting plant productivity, botanical composition and soil characteristics, spatial autocorrelation is unlikely to affect the results. All grasslands were composed of a mixture of forbs and grasses, although the dominant functional group depended on the grassland type. Their productivity ranged from 156 to 840 g dw yr-1 m-2, managed grasslands being more productive than natural grasslands (Sup. Fig. 2). Three replicates per grassland type were randomly chosen in the study area, for a total of twelve sites ranging from 1000 to 1300 m.a.s.l. In each site, five 1m² plots (60 plots in total) were selected randomly within a 5m-radius and assigned to one of the five following grasshopper herbivory treatments: *Miramella alpina*, *Pseudochorthippus parallelus*, *Stauroderus scalaris*, an equal mixture of the three species, no grasshoppers (control). If single species treatments have contrasted effects on soil microbes, the mixture treatment should discriminate between additive and non-additive effects. The grasshopper species were chosen according to their diet, as *M. alpina* prefers to feed on forbs, *P. parallelus* on grasses and *S. scalaris* mainly on grasses but also on forbs (Ibanez et al. 2013). Plots were covered by a 1m^3 metallic cage covered by insect-proof mesh (PE 22.30, 920 x 920 µm; DIATEX, Saint Genis Laval, France), into which the adult grasshoppers were introduced. The number of grasshoppers per cage was adjusted to fit 1-g of grasshopper for 300-g of plant aboveground biomass (dry mass). Preliminary field observations revealed that this ratio corresponds to the maximal natural insect densities locally observed in the study area. This standardization is necessary to avoid confounding correlations between herbivory load and herbivore diet, or between herbivory load and plant productivity. For the standardization, the dry mass of each species and each sex was also considered, after preliminary measurements. Grasshoppers were present inside the cages during 65 days from the 2nd week of July until the 3rd week of September, during two consecutive years 2016 and 2017. During these periods, the number of grasshoppers per cage was checked every two weeks, including the control cages from which a few grasshoppers were occasionally removed. Several ecosystem characteristics were measured, as detailed in the following sections. In short, soil abiotic and microbial characteristics as well as plant biomass were measured at the end of the experiment, in mid-September 2017. Herbivory marks on standing biomass was visually estimated in 2016, while plant specific abundance was estimated before the experiment in 2016 and after one year in 2017, in order to calculate the effect of herbivory.

Soil characteristics

In mid-September 2017, four 10-g soil samples were collected in each plot at 5-cm depth, and then bulked. The samples were sieved at 5-mm and stored at 4°C before processing within 48h. Subsamples of 5-g of soil were dried at 70°C for 1 week to determine soil moisture, followed by 4h at 550°C to determine soil organic matter (SOM) content. Soil subsamples were air dried and ground to powder to measure total C and N contents using a FlashEA 1112 elemental analyzer (Fisher Scientific Inc., Waltham, MA, USA), and to determine soil pH in a 1:2.5 (soil : distilled water) solution. Solution of 0.5M K_2SO_4 was used to extract soil nitrate (NO_3^-), ammonium (NH_4^+), total dissolved nitrogen (TDN) (Jones and Willett 2006), and phosphate (PO_4^-) on 10-g of fresh soil. N and P concentrations were measured on an automated photometric analyzer using standard colorimetric methods (Gallery Plus: Thermo Fisher Scientific, Waltham, Massachusetts, USA). Dissolved organic nitrogen (DON) was calculated as the difference between TDN and the mineral N (NO_3^- + NH_4^+).

Microbial biomass, activities and traits

The same soil samples were used for the analysis of microbial characteristics. Microbial biomass nitrogen (MBN) content was based on the difference of soil N content before and after chloroform-fumigation extraction of 10 g of fresh soil (Vance et al. 1987). We then calculated MBN in µgN/g using a correction factor of 0.45 (Jenkinson et al. 2004). Potential nitrogen mineralization rates (PNM) were estimated after incubation under anaerobic conditions of 10 g of fresh soil for 7 days at 40°C in the dark (Wienhold 2007). Mineralized organic N was accumulated as NH_4^+ during this incubation and PNM rates were calculated based on the difference between NH_4^+ content before and after incubation and expressed as µgN/g dry soil/day. Microbial resource acquisition strategies were characterized using different microbial community-weighted mean traits (Piton et al. 2020b). We measured the potential activity of seven extracellular enzymes contributing to the degradation of C-rich substrates (α-glucosidase, β-1,4-glucosidase, β-d-cellobiosidase and β-xylosidase), N-rich substrates (β-1,4-N acetylglucosaminidase and leucine aminopeptidase), and P-rich substrates (phosphatase) using standardized fluorimetric techniques (Bell et al. 2013). We homogenized 2.75 g of fresh soil (1 minute in a Waring blender) in 200 ml of sodium acetate buffer solution that was adjusted at the mean soil pH observed in the present study (i.e. 6.2). The resulting soil slurries were added in duplicates to 96-deep-well microplates. We then added a substrate solution for each enzyme. Duplicated standard curves (0–100-µM concentration) were prepared for each soil sample by mixing 800 ml of soil slurry with 200 ml of 4-methylumbelliferone (MUB) or 7-amino-4-methylcoumarin (MUC) in 96-deep-well microplates. Microplates were incubated during 3hrs (dark, 175 rpm, 20°C), and centrifuged at 2,900 g for 3 min. Soil slurries (250 µl) were then transferred into black Greiner flat-bottomed microplate and scanned on a Varioskan Flash (Thermo Scientific) reader using excitation at 365 nm and emission at 450 nm (Bell et al. 2013). For each soil sample, the four enzyme activities degrading C-rich substrates, the two enzymes activities degrading N-rich substrates and all the seven enzymes were summed to obtain extracellular enzyme activity for C-rich substrates (EEC), N-rich substrates (EEN) respectively. Phosphatase activity was used to represent extracellular enzyme activity for P-rich substrates (EEP). EEC, EEN and EEP were calculated per gram of dry soil per hour (global activities, nmol activity g-1 dry soil h-1). Microbial resource limitation

248 and associated trade-offs between C and N acquisition was assessed through the calculation of the
249 ecoenzymatic ratio EEC:EEN (Sinsabaugh et al. 2009).

Vegetation surveys and plant biomass

251 In order to quantify the diet of each grasshopper species, a visual estimation of herbivory intensity
252 was conducted in August 2016. Ten individuals belonging to each of the 6 most dominant plant
253 species in each site were inspected for herbivory marks by grasshoppers and the percentage of leaf
254 area eaten was visually estimated.
255
A botanical survey was conducted at the beginning of the experiment (June 2016) and after one year
of controlled herbivory (June 2017). The point quadrat method was used (Levy and Madden 1933;
Vittoz and Guisan 2007; Lavorel et al. 2008), with 50 regularly spaced points in each 1m² cage. This
method allows to determine the relative abundance of the most common species in each plot, but
does not constitute a complete inventory of the specific richness. To assess the effect of herbivory,
we computed the proportion of forbs in 2017 minus the proportion of forbs in 2016 in each cage. At
the end of the experiment in September 2017 the aboveground biomass was harvested in each plot,
dried 48h at 40°C, sorted into forbs and grasses and weighed. In order to get a community-level
measure of leaf C/N ratio, a representative sample of the harvested biomass was ground to powder,
homogenized, and 5 mg of the leaf powder were then analyzed for carbon and nitrogen
concentration using a CHN analyser (FlashEA 1112, ThermoFisher Scientific, MA, USA). These
measures also provided the amount of nitrogen in aboveground plant biomass per unit area (gN/m²,
N_B).

Statistical analysis

270 Linear mixed models in which the random effects corresponded to the 12 sites (with 5 pseudo-
271 replicas per site) were used to predict the logit of the total percentage of leaf biomass eaten in
272 function of grassland type, herbivore identity, plant functional group (forbs *vs* grasses), and the
273 interaction between these three factors.
274
Coinertia analysis (Dray et al. 2003) was used to quantify the coefficient of correlation (RV
coefficient, ranging between 0 and 1) between vegetation characteristics (total aboveground
biomass, forbs biomass, N_B, community-level plant C/N ratio, plant Shannon index), soil abiotic
characteristics (water content, pH, SOM, TDN, phosphorus content, C/N ratio) and soil microbial
characteristics (EEC, EEN, EEP, EEC:ENN ratio, PNM and microbial biomass). Redundancy
analysis (RDA, Borcard et al. 1992) was used to quantify the amount of variation of either microbial,
soil abiotic and vegetation characteristics explained by either grassland type or site.
282
283 In order to quantify the effect of herbivory on the six microbial characteristics, the standardized
284 response to herbivory was calculated as the difference between grasshopper treatments and the
285 control treatment (no grasshoppers) of the same site, divided by the standard deviation of the
286 corresponding site. Coinertia analysis was used to test if the standardized responses depended on
287 soil abiotic and/or on vegetation characteristics.

288
289 To test if these effects of herbivory depended on grassland characteristics (either SOM or plant
290 productivity N_B), and on herbivore identity, the standardized responses of microbial characteristics
291 were then used as a response variable in separate linear mixed models including sites as random
292 effects. Since the warm grassland #1 had a particularly high SOM, the same models as above were
293 performed excluding this site to check if it had a disproportionate effect on the results.
294
295 In all mixed models, the denominator degrees of freedom were calculated using Satterthwaite's
296 approximation (Satterthwaite 1946). Type III sums of squares were used for models including
297 interactions between factors. For each model, the normality and homoscedasticity hypothesis were
298 visually checked. All statistical analyses were performed with the R software (R Core Team 2019),
299 using the packages lmerTest (Kuznetsova et al. 2017) and ade4 (Dray and Dufour 2007).

Results

Leaf quality and consumption

The percentage of leaf area eaten depended to a large extent on the interaction between plant functional group and herbivore identity (p<0.001), indicating that the selected grasshoppers had contrasted diets as expected. *M. alpina* preferred forbs over grasses (7% of leaf area consumed *vs* 3.2%), *S. scalaris* preferred grasses over forbs (7.2% *vs* 3%), while *P. parallelus* ate almost exclusively grasses (11.3% *vs* 0.9%, Sup. Fig. 3A). When these three species were combined, grasses were more impacted than forbs (7.4% *vs* 4.8%). The overall effect of grassland type on the leaf area eaten was not significant (p=0.50), which reflects the fact that the number of introduced grasshoppers in each cage depended on the estimated plant standing biomass (1-g of grasshopper for 300-g of plant aboveground biomass). However, there was a significant interaction between grassland type and plant functional group (p<0.001), because in forb dominated communities grass species were much more heavily impacted than forb species (15.3% *vs* 2.7%), whereas in the other grassland types both functional groups were consumed in similar proportions (Sup. Fig. 3B). The p-value of the three-way interaction between plant functional group, grassland type and grasshopper identity equaled 0.58, which indicates that the diet of each grasshopper species did not depend on grassland type. The elementary analysis of fresh plant material showed that the leaf C/N ratio was higher for grasses than for forbs, whatever the grassland type (p<0.001 in all cases). However, the magnitude of the difference depended on grassland type (interaction between plant functional group and grassland type, p<0.001). The largest difference was in warm grasslands (forbs: 25, grasses: 38) and the smallest in communities dominated by forbs (forbs: 22, grasses: 27, Sup. Fig. 4). The botanical survey conducted after one year of herbivory revealed that the proportion of forbs did not vary in most treatments (p>0.4), except in the cages containing the forb-feeding *M. alpina* where forbs declined by 7.2% (Sup. Fig. 5, p=0.013).

Relations between vegetation, soil microbial and abiotic characteristics

The coinertia analysis between six microbial soil characteristics (nitrogen mineralization potential, EEN, EEC, EEP, microbial biomass, EEC/EEN ratio) and six abiotic soil characteristics (water content, pH, SOM, TDN, phosphorus content, C/N ratio) had a RV coefficient of 0.59 (p<0.001). One axis of covariation between the two types of variables corresponds to an anticorrelation between soil pH and soil phosphorus content on the one hand, and EEP on the other hand, as well as EEC/EEN ratio to a lesser extent (Figure 1A). Therefore, in acidic soils having low phosphorus content, microbes invested more in phosphorus acquisition enzymes and slightly more on carbon than on nitrogen acquisition enzymes. Another axis of covariation was between the abiotic characteristics TDN, SOM and water content, and the microbial characteristics EEC, EEN and PNM. The RDAs indicated that grassland types were heterogeneous with respect to soil characteristics, grassland type explaining 28% of the variation for the six soil abiotic characteristics but only 9% for the six

microbial characteristics. In contrast, the soil characteristics were homogeneous at the site level
(Sup. Fig. 6).

Turning to vegetation, the coinertia analysis between the six abiotic soil characteristics and the five
vegetation characteristics (total aboveground biomass, forbs biomass, N_B, community-level plant C/N
ratio, plant Shannon index) had a RV coefficient of 0.22 (p<0.001). Ecosystem productivity was
negatively associated with SOM, pH and soil phosphorus content, and was not linked to TDN (Figure
1B). The RV coefficient of the coinertia between five vegetation characteristics and the six microbial
characteristics was equal to 0.29 (p<0.001). This much lower RV coefficient than that between soil
microbial and soil abiotic characteristics (0.59) suggests that the microbial soil characteristics were
more related to soil abiotic than to vegetation parameters. The EEC/EEN ratio covaried with plant
productivity (aboveground biomass and N_B) while EEC, EEN and PNM were associated with the leaf
C/N ratio and the proportion of grasses (Figure 1C). The RDA indicated that grassland types were
homogeneous with respect to vegetation characteristics with 53% of variation explained by grassland
type (Sup. Fig. 6). Indeed, warm grasslands were characterized by high leaf C/N ratio, forb-
dominated grassland by a high proportion of forbs and a high Shannon diversity, intensive
grasslands by a high biomass and nitrogen content, while extensive grasslands had intermediate
vegetation characteristics. Means ± sd of each vegetation, soil microbial and abiotic characteristics
are given for each of the 12 sites in Sup. Tab. 1.

Effect of herbivory on microbial characteristics

Another coinertia analysis, this time involving the standardized responses of microbial communities
to herbivory, revealed that they were related to soil abiotic characteristics (RV=0.22, p<0.001). The
coinertia analysis (Figure 2) and the linear mixed models (Figure 3) both indicated that the responses
of PNM and EEN to herbivory increased with SOM, as well as EEC and EEP although to a lesser
extent (see Figure 3 for the p-values of the linear mixed models). More specifically, below 10% of
SOM herbivory decreased PNM and soil enzymatic activities while above 15% of SOM herbivory had
a positive effect on these variables (Figure 3). The response of microbial biomass to herbivory did
not depend on SOM (p=0.75) but varied according to TDN. In sites with TDN below 30 µgN/g,
herbivory increased microbial biomass, while in sites with TDN above 50 µgN/g herbivory decreased
microbial biomass (Figure 3E, p=0.014). Finally, the response of EEP to herbivory was positively
related to pH (p=0.007). In contrast, the effect of herbivory on microbial characteristics did not
depend on vegetation traits (RV=0.09, p=0.17). In particular, neither plant biomass nor productivity
(N_B) were related to changes in the standardized responses of microbial characteristics (details not
shown, p>0.05). In all cases, the effects of herbivory on microbial characteristics depended neither
on herbivore diet nor on the interaction between herbivore diet and ecosystem characteristics such
as SOM, TDN or productivity (details not shown, p>0.05).

Discussion

Covariations between ecosystem properties

We first explore the covariations between vegetation, soil biotic and soil abiotic characteristics, in order to better identify the ecological factors which differentiate the studied grasslands. In the next sections, we discuss how these differences condition various types of responses to herbivory. The comparison of the 12 study sites indicated that, in line with the productivity model, soil C content measured by SOM was low in sites having the higher plant biomass (Figure 1B). However, there was a positive relationship between SOM and TDN (Figure 1A), and no relationship between TDN and plant biomass (Figure 1B). The decoupling between TDN and plant biomass might be explained by management, since extensive and intensive grasslands are fertilized with manure each year; intensive grasslands being the most fertilized and productive ones in the study area (Piton et al. 2020b & Sup. Fig. 2). Nutrients available after fertilization are likely to be rapidly assimilated by the fast growing and productive plant species, resulting in both high productivity and low measured nutrient availability. After plant growth, mowing exports nutrients away from these grasslands, while fertilization reintroduces the lacking nutrients. Therefore, in managed grasslands a high proportion of annual primary production is consumed and indirectly returned to the soil through subsequent fertilization, as are natural fertile ecosystems in the productivity model (Bardgett and Wardle 2010).

Turning to microbes, those living in soils with high SOM and TDN accelerated the nutrient cycle (high PNM, EEC and EEN, Figure 1A). However, microbial biomass did not increase with SOM, which implies that high SOM comes along with higher mass-specific extracellular activities for carbon uptake and higher mass-specific nitrogen mineralization, which corresponds to a resource acquisition (A) strategy at the expense of growth yield or carbon use efficiency (sensu Malik et al. 2020). Microbial biomass did not covary with any soil parameters (Figure 1 A & C), as in Farrell et al. (2011, see their Table 1) and Chu & Grogan (2010, see their Tables 1 & 2). This may reflect the fact that microbial biomass results from the complex interplay between potentially opposing factors such as microbial resource use efficiency, resource acquisition and resource availability, itself depending on SOM and plant input quantity and quality (e.g. root exudates). Moreover, microbial biomass depends on the whole soil trophic network (Calderón-Sanou et al. 2021). Furthermore, as found in alpine ecosystems in Cordillera Darwin, Tierra del Fuego (Chile) (Thébault et al. 2014), we suggest that competition between soil microbes and plants for nutrient resources in SOM and TDN-rich soils may contribute to the higher mass-specific microbial investment in resource acquisition in these soils. This result stresses the importance to consider the resource acquisition strategy of soil microbes and not only their biomass to understand their response to environmental gradients and their effect on nutrient recycling in soil (Piton et al. 2020a). Recent literature has proposed that microbes with an A strategy could be favored either when resources are scarce (Malik et al. 2019, 2020; Piton et al. 2020b), or when resources are abundant (Wood et al. 2018). Since microbes rely on two main carbon sources, not only soil organic matter but also those exuded by plants in the rhizosphere, it is necessary to explore the relationships between the traits of the microbial and plant communities. We found that microbes increased their investment in extracellular enzymatic activities in plant communities having higher leaf C/N ratios (Figure 1C), which corresponds to conservative plant

resource use strategy. Since exploitative plants have been shown to produce more root exudates
(Williams et al. 2022), an extra labile carbon source would be available to microbes in exploitative
plant communities, favoring microbes with a high yield (Y) strategy characterized by low enzyme
production, according to Malik et al. (2020). This might also be reinforced by organic inputs in the
most productive intensive grasslands. In contrast, in conservative plant communities, microbes would
lack labile carbon and nutrients sources and rely more on the biodegradation of polymerized soil
organic matter for resource acquisition. This implies that A strategists characterized by the
production of extracellular enzymes would be favored, which ultimately releases nitrogen from soil
organic matter (Figure 1C).

In summary, the pattern of covariations between ecosystem properties suggests a distinction
between two kinds of grasslands. Although both microbial strategies are likely to coexist in each soil
sample, both grassland kinds favor either strategy. More specifically, on the one hand the most
productive sites (>400 g dw/m²) are characterized by exploitative plants, by soils having relatively low
SOM (<10%) and TDN (<40 µgN/g) contents, and by Y-strategist microbes producing fewer
extracellular enzymes, possibly because of higher C supply in the rhizosphere (Table 2, left column).
On the other hand, in the least productive sites dominated by conservative plants, the SOM and TDN
contents are higher and the microbial community is characterized by an A strategy since microbes
need to produce extracellular enzymes to get access to SOM resources (Table 2, right column).
431

Herbivory effects on enzymatic activities

The present section explores if the effects of herbivory on microbial communities depend on the two
grassland kinds described above (Table 2). We found a covariation between SOM and the
responses of soil enzymatic activities and PNM to herbivory (Figure 2). Herbivory had a negative
effect on soil enzymatic activities and PNM when SOM was below 10%, and a positive effect when
SOM was above 15% (Figure 3), which means that herbivory can decelerate or accelerate nutrient
cycling depending on soil conditions. This occurs in relation to microbial strategies; in this section we
interpret the variations of soil enzymatic activities following herbivory as responses of the microbial
community, favoring either A or Y strategy. In soils rich in SOM already characterized by high mass-
specific extracellular enzyme activities, microbes invested even more in extracellular enzymes in
response to herbivory indicating that A strategists were favored under these conditions. Herbivory
provides fresh organic matter (FOM such as green fall, feces, cadavers) to the soil, which contains
nutrients that are more easily available than SOM but which nevertheless needs to be depolymerized
before assimilation (e.g. chitin). Since in rich-SOM-sites, A strategists possess the enzymatic traits
that give them access to this resource, this would explain why herbivory promoted the investment of
A strategists in extracellular enzymes. Consequently, the enhancement of extracellular enzymatic
activities might have cascaded on PNM, as expected according to the key limiting role of organic
matter depolymerization before microbial assimilation and subsequent N mineralization (Schimel and
Bennett 2004). In contrast, in low SOM soils, Y strategists decreased even more their investment in
extracellular enzymes. A possible explanation is that litter inputs are not the only consequences of
herbivory on ecosystem functioning (Hunter et al. 2012). In particular, herbivory stimulates root
exudation (Holland 1995; Holland et al. 1996), which is known to increase microbial activity in the

rhizosphere and subsequently to favor compensatory plant growth (Hamilton and Frank 2001;
Hamilton et al. 2008). Williams et al. (2022) found that root exudation is more intense in exploitative
than in conservative plant communities, but it is not known whether the effect of herbivory on root
exudation depends on plant functional groups. If the stimulation of root exudation by herbivory is
more intense for exploitative than for conservative plants, this would explain the finding that herbivory
decreased the production of extracellular enzymes in sites having low SOM, since they were
dominated by exploitative plants. However, although the effect of herbivory on soil enzymatic
activities and PNM depended on SOM content (Figure 3), it did not directly depend on the mean leaf
C/N ratio (details not shown, $p > 0.05$). Root exudation following herbivory has been found to enhance
PNM for the benefit of plants (Hamilton and Frank 2001), while our results suggest the opposite.
However, inputs of available C may also result in N immobilization in the microbial biomass (Lovett
and Ruesink 1995), which is this time in line with our results. This calls for both qualitative and
quantitative characterization of root exudation in future works to further elucidate how soil and plant
properties jointly control ecosystem response to herbivory. Herbivory may also decelerate nitrogen
mineralization through the induction of recalcitrant plant secondary compounds (Schultz and Baldwin
1982; Rhoades 1985; Agrawal 1999), which subsequently slows down litter decomposition (Findlay
et al. 1996; Hattenschwiler and Vitousek 2000). However, we did not find any effect of herbivory on
litter decomposition in a parallel study (Ibanez et al. 2021).
472
Turning to microbial biomass, herbivory had a positive effect in sites having low TDN, and a negative
effect when TDN was high (Figure 3E), but contrary to the enzymatic activities the effect of herbivory
on microbial biomass did not depend on the SOM content (Figure 2A). In previous studies, herbivory
increased microbial biomass through an input of labile C contained in the feces (Lovett and Ruesink
1995; Van Der Wal et al. 2004). Yet, herbivory was also found to decrease microbial biomass when
plants' C inputs to the soil are instead reduced due to herbivores respiration (Sankaran and
Augustine 2004). Since TDN was positively linked with mean leaf C/N ratio (Figure 1B, Robson et al.
2007; Legay et al. 2013), itself being negatively correlated to root exudates (Williams et al. 2022), we
hypothesize that in low TDN sites, herbivory triggered an input of labile C into the rhizosphere,
thereby stimulating the growth of microbial biomass, whereas in sites with high TDN, herbivory
reduced the plants C inputs, with a negative impact on microbial biomass.

No effect of herbivore diet

485
The proportion of forbs in the diet of *M. alpina* was the highest, followed by the mixed feeder *S.
scalaris* and then by the grass feeder *P. parallelus* (Sup. Fig. 2A), in line with previous studies
(Ibanez et al. 2013). Since the grasshoppers' diet did not depend on the grasslands they were
introduced in, it was possible to test if the effects of herbivory on soil enzymatic activities depended
on their diet. The diet model predicts that herbivores feeding on high quality plants should favor the
accumulation of poor-quality litter and thereby slow down nutrient cycling, and that herbivores
feeding on low quality plants should favor the accumulation of high-quality litter, with an accelerating
effect on nutrient cycling (Belovsky and Slade 2000).
494

In the present study, high quality plants characterized by low leaf C/N ratio (Sup. Fig. 4) were
consumed preferentially by the forb feeder *M. alpina* (Sup. Fig. 2A), which decreased their relative
abundance by about 7% (Sup. Fig. 5). In warm and forb-dominated prairies, litter decomposition
during winter has been found to be faster for forbs (28% of mass loss) than for grasses (17%),
although in extensive and intensive grasslands winter decomposition was similar for both plant
functional groups, due to the intermittent presence of snow cover (Ibanez et al. 2021). In any case,
this suggests that the year-round decomposition rate is higher for forbs than for grasses, as generally
described (Tilman 1988). Given that *M. alpina* modifies the balance between forbs and grasses, this
should affect the overall decomposition rate, with potential impacts on soil microbial communities.
However, *M. alpina* did not have any contrasted effect on soil microbial characteristics, in
comparison to the other grasshopper species. Perhaps this would have required a heavier impact on
the relative abundance of forbs, rather than only 7%.
507
Lower quality plants (high leaf C/N ratio) were consumed preferentially by the mixed feeder *S.*
scalaris and almost exclusively by the grass feeder *P. parallelus* (Sup. Fig. 2A), but without impact
on the relative abundance of forbs and grasses (Sup. Fig. 5). Also, grass feeders did not have any
distinguishable effect on soil microbes, relatively to the forb feeding *M. alpina*. Previous short-term
experiments using similar insect loads have reported that the effects of grasshoppers on soil
processes depended on their diet type (e.g. Schmitz 2008; Belovsky and Slade 2018). Our findings
question the generality of these results, since we found that the effect of herbivory on enzymatic
activities did not depend on grasshopper species identity.

Conclusion

517
Previous works showed that herbivory has contrasted impacts on ecosystem functioning in general
(Brown and Gange 1992; Bardgett 2005; Schmitz 2008), and on soil microbes in particular (Denton
et al. 1998; Stark and Grellmann 2002), in function of the environmental conditions and of the diet of
herbivores. The motivation of this work was to observe such contrasted impacts in a single
experiment across different types of grasslands with contrasting plant productivity, using herbivores
having different diets, in order to better understand the pathways which create this apparently
idiosyncratic pattern. We did not find any effect of herbivore diet on soil microbial characteristics,
contrary to our expectation. In contrast, the effects of herbivory on soil microbes depended on
several properties of the 12 study sites. Table 2 summarizes our main findings and can provide some
lines of interpretation, although its dichotomic schematization does not fully represent the
multidimensionality of the results (Figure 1-3). On the one hand, the most productive sites were
characterized by a higher biomass of exploitative plant species which depleted N resources in the
soil, and by yield-strategists microbes having a smaller investment in extracellular enzymes. Since
exploitative plant species tend to produce more root exudates (Williams et al. 2022), we postulate
that herbivory increased C supply in the rhizosphere (e.g. Holland et al. 1996; Hamilton and Frank
2001), which would explain the observed increase in microbial biomass and the lesser production of
extracellular enzymes. On the other hand, the least productive sites were characterized by a lower
biomass of conservative plants, which led to high soil C content, and by acquisition-strategists
microbes having a larger investment in extracellular enzymes. We hypothesize that, in those sites,

the consumption of plants results in lower soil C inputs due to herbivore respiration (Sankaran and

Augustine 2004), which would explain why herbivory eventually decreased microbial biomass and

increased the investment in extracellular enzymes in these less productive sites. Although the

framework presented in Table 2 has some similarities with the productivity model (Bardgett and

Wardle 2010), our findings point towards an acceleration of N cycling in less productive sites and a

deceleration in more productive sites, in opposition with the productivity model. However, both

frameworks do not consider the same time scales. The current experiment was conducted over two

years and focuses on physiological time scales (root exudation, microbial enzyme production), while

the productivity model encompasses plant community dynamics, which occur on longer time scales.

In any case, we are convinced that none of these frameworks fully grasp the complex relationships

between plants, soil, microbes and the effects of herbivory, which are most likely multidimensional.

At the very least, these frameworks have some heuristic value and can be used for the design of

future experiments.

Figures & Tables

Figure 1

Coinertia of (A) soil microbial and abiotic characteristics, (B) soil abiotic and vegetation characteristics and (C) microbial and vegetation characteristics of the 60 cages. In brown, soil abiotic characteristics (tdn=total dissolved nitrogen, hum=soil water content, som=soil organic matter, CN=soil C/N ratio, pho=soil phosphorus content). In blue, soil microbial characteristics (PNM=potential nitrogen mineralization, mbn=microbial biomass, EEN, EEC, EEP= extracellular enzyme activities related to either nitrogen, carbon or phosphorus, ratio=EEC/EEN). In green, vegetation characteristics (plantCN=community-level plant C/N ratio, biom=total aboveground plant biomass, forbs=forbs biomass (g), N_B=nitrogen content in total aboveground plant biomass (N$_B$), shan=Shannon diversity index of plants). The % indicate the projected inertia.

Figure 2

Coinertia of soil abiotic characteristics and the standardized response to herbivory of microbial characteristics. (A) Canonical weights of soil abiotic characteristics (in brown) and standardized response to herbivory of microbial characteristics (in purple). (B) The 48 arrows (60 cages - 12 controls) correspond to all the plots containing grasshoppers. The soil abiotic characteristics are at the beginning of the arrows, the standardized response to herbivory of microbial characteristics are at their end. tdn=total dissolved nitrogen, hum=soil water content, som=soil organic matter, CN=soil C/N ratio, pho=soil phosphorus content, PNM=potential nitrogen mineralization, mbn=microbial biomass, EEN, EEC, EEP= extracellular enzyme activities related to either nitrogen, carbon or phosphorus, ratio=EEC/EEN. The % indicate the projected inertia.

Figure 3

Relationship between soil organic matter and the standardized response to herbivory, calculated as the difference between each grasshopper treatment and the control treatment (no grasshoppers) of the same site, divided by the standard deviation of this site (n=48, 60 cages - 12 controls). p is the p-value of the linear mixed model including all sites (continuous regression line), while p' excludes the warm grassland #1 having SOM values >30 (dashed regression line).

Table 1

Expected effects of herbivory on matter cycling (either accelerating or decelerating), according to the productivity and diet model. In each of the four combinations, the prediction of the diet model is reported on the top (in orange), and that of the productivity model on the bottom (in green). In two cases out of four, the predictions are opposite.

| | | Diet model | |
		High quality diet	Low quality diet
Productivity model	Productive ecosystems (high plant biomass, low soil organic matter)	decelerate accelerate	accelerate accelerate
	Less productive ecosystems (low plant biomass, high soil organic matter)	decelerate decelerate	accelerate decelerate

590 # Table 2

592 Schematic representation of the main results, and their articulation with previous works.

	Most productive sites	Least productive sites	Source
Plants	High biomass	Low biomass	Figure 1
	Low leaf C/N	High leaf C/N	Figure 1
	Exploitative plants	Conservative plants	Figure 1
	High C supply in the rhizosphere	Low C supply in the rhizosphere	Williams et al. 2020
Soil	Low soil organic matter	High soil organic matter	Figure 1
	Low total dissolved nitrogen	High total dissolved nitrogen	Figure 1
Microbes	Low extracellular enzymes	High extracellular enzymes	Figure 1
	High yield strategy Y	Resource acquisition strategy A	Malik et al. 2020
Effects of herbivory on microbes	Decreases extracellular enzymes	Increases extracellular enzymes	Figures 2 & 3
	Decreases mineralization	Increases mineralization	Figures 2 & 3
	Increases microbial biomass	Decreases microbial biomass	Figures 2 & 3
	Increases C supply is the rhizosphere (root exudates)[1]	Herbivore respiration decreases C inputs to the soil[2]	[1]: e.g. Hamilton & Frank 2001 [2]: e.g. Sankaran & Augustine 2004

Acknowledgements

This work was funded by the ECO-SERVE project through the 2013–2014 BiodivERsA/FACCE-JPI joint call for research proposals, with the national funders ANR, NWO, FCT (BiodivERsA/001/2014), MINECO, FORMAS and SNF. This work was also funded by the Alpine Ecology Lab. We thank the municipality of Autrans-Méaudre, the local farmers and the Refuge de Gève for their authorization to access the study sites, Jonathan Crison for his hospitality. We are grateful to Jean-Noël Avrillier, Annie Millery, Hugo Girard, Matteo Tolosano, Louise Maris, Pablo Raguet, Océane Guillot, Jessica Barbe & Alison Dillien for their assistance during field and lab work.

References

Agrawal AA (1999) Induced responses to herbivory in wild radish: effects on several herbivores and plant fitness. Ecology 80:1713–1723

Bakker ES, Olff H, Boekhoff M, et al (2004) Impact of herbivores on nitrogen cycling: contrasting effects of small and large species. Oecologia 138:91–101. https://doi.org/10.1007/s00442-003-1402-5

Bardgett R (2005) The biology of soil: a community and ecosystem approach. Oxford university press

Bardgett RD, Wardle DA (2010) Aboveground-belowground linkages: biotic interactions, ecosystem processes, and global change. Oxford University Press Oxford

Bardgett RD, Wardle DA (2003) Herbivore-mediated linkages between aboveground and belowground communities. Ecology 84:2258–2268

Bell CW, Fricks BE, Rocca JD, et al (2013) High-throughput fluorometric measurement of potential soil extracellular enzyme activities. JoVE J Vis Exp e50961

Belovsky GE, Slade JB (2000) Insect herbivory accelerates nutrient cycling and increases plant production. Proc Natl Acad Sci 97:14412–14417. https://doi.org/10.1073/pnas.250483797

Belovsky GE, Slade JB (2018) Grasshoppers affect grassland ecosystem functioning: Spatial and temporal variation. Basic Appl Ecol 26:24–34

Borcard D, Legendre P, Drapeau P (1992) Partialling out the spatial component of ecological variation. Ecology 73:1045–1055

Brown VK, Gange AC (1992) Secondary plant succession: how is it modified by insect herbivory? Vegetatio 101:3–13

Buckland SM, Grime JP (2000) The effects of trophic structure and soil fertility on the assembly of plant communities: a microcosm experiment. Oikos 91:336–352. https://doi.org/10.1034/j.1600-0706.2000.910214.x

Calderón-Sanou I, Münkemüller T, Zinger L, et al (2021) Cascading effects of moth outbreaks on subarctic soil food webs. Sci Rep 11:15054. https://doi.org/10.1038/s41598-021-94227-z

Cease AJ, Elser JJ, Ford CF, et al (2012) Heavy Livestock Grazing Promotes Locust Outbreaks by Lowering Plant Nitrogen Content. Science 335:467–469. https://doi.org/10.1126/science.1214433

Chu H, Grogan P (2010) Soil microbial biomass, nutrient availability and nitrogen mineralization potential among vegetation-types in a low arctic tundra landscape. Plant Soil 329:411–420. https://doi.org/10.1007/s11104-009-0167-y

Clauss M, Lechner-Doll M, Streich WJ (2003) Ruminant diversification as an adaptation to the physicomechanical characteristics of forage. A reevaluation of an old debate and a new hypothesis. Oikos 102:253–262

640   Denton CS, Bardgett RD, Cook R, Hobbs PJ (1998) Low amounts of root herbivory positively
641   influence the rhizosphere microbial community in a temperate grassland soil. Soil Biol
642   Biochem 31:155–165

643   Deraison H, Badenhausser I, Loeuille N, et al (2015) Functional trait diversity across trophic levels
644   determines herbivore impact on plant community biomass. Ecol Lett 18:1346–1355

645   Dray S, Chessel D, Thioulouse J (2003) Co-inertia analysis and the linking of ecological data tables.
646   Ecology 84:3078–3089

647   Dray S, Dufour A-B (2007) The ade4 package: implementing the duality diagram for ecologists. J
648   Stat Softw 22:1–20

649   Farrell M, Hill PW, Farrar J, et al (2011) Seasonal variation in soluble soil carbon and nitrogen across
650   a grassland productivity gradient. Soil Biol Biochem 43:835–844.
651   https://doi.org/10.1016/j.soilbio.2010.12.022

652   Fielding DJ, Trainor E, Zhang M (2013) Diet influences rates of carbon and nitrogen mineralization
653   from decomposing grasshopper frass and cadavers. Biol Fertil Soils 49:537–544

654   Findlay S, Carreiro M, Krischik V, Jones CG (1996) Effects of damage to living plants on leaf litter
655   quality. Ecol Appl 6:269–275

656   Frank DA, Groffman PM, Evans RD, Tracy BF (2000) Ungulate stimulation of nitrogen cycling and
657   retention in Yellowstone Park grasslands. Oecologia 123:116–121

658   Garibaldi LA, Semmartin M, Chaneton EJ (2007) Grazing-induced changes in plant composition
659   affect litter quality and nutrient cycling in flooding Pampa grasslands. Oecologia 151:650–662

660   Grigulis K, Lavorel S, Krainer U, et al (2013) Relative contributions of plant traits and soil microbial
661   properties to mountain grassland ecosystem services. J Ecol 101:47–57.
662   https://doi.org/10.1111/1365-2745.12014

663   Hamilton EW, Frank DA (2001) Can plants stimulate soil microbes and their own nutrient supply?
664   Evidence from a grazing tolerant grass. Ecology 82:2397–2402

665   Hamilton EW, Frank DA, Hinchey PM, Murray TR (2008) Defoliation induces root exudation and
666   triggers positive rhizospheric feedbacks in a temperate grassland. Soil Biol Biochem
667   40:2865–2873

668   Harrison KA, Bardgett RD (2004) Browsing by red deer negatively impacts on soil nitrogen
669   availability in regenerating native forest. Soil Biol Biochem 36:115–126.
670   https://doi.org/10.1016/j.soilbio.2003.08.022

671   Hattenschwiler S, Vitousek PM (2000) The role of polyphenols in terrestrial ecosystem nutrient
672   cycling. Trends Ecol Evol 15:238–243

673   Hofmann RR (1989) Evolutionary steps of ecophysiological adaptation and diversification of
674   ruminants: a comparative view of their digestive system. Oecologia 78:443–457

675   Holland JN (1995) Effects of above-ground herbivory on soil microbial biomass in conventional and
676   no-tillage agroecosystems. Appl Soil Ecol 2:275–279

Holland JN, Cheng W, Crossley D (1996) Herbivore-induced changes in plant carbon allocation: assessment of below-ground C fluxes using carbon-14. Oecologia 107:87–94

Hunter MD (2001) Insect population dynamics meets ecosystem ecology: effects of herbivory on soil nutrient dynamics. Agric For Entomol 3:77–84

Hunter MD, Reynolds BC, Hall MC, Frost CJ (2012) Effects of herbivores on terrestrial ecosystem processes: the role of trait-mediated indirect effects. In: Trait-mediated indirect interactions: ecological and evolutionary perspectives. Takayuki Ohgushi, Oswald Schmitz, Robert D. Holt, Cambridge University Press, pp 339–370

Ibanez S, Brun C, Millery A, et al (2021) Litter and soil characteristics mediate the buffering effect of snow cover on litter decomposition. Plant Soil 460:511–525. https://doi.org/10.1007/s11104-020-04803-x

Ibanez S, Lavorel S, Puijalon S, Moretti M (2013) Herbivory mediated by coupling between biomechanical traits of plants and grasshoppers. Funct Ecol 27:479–489. https://doi.org/10.1111/1365-2435.12058

Jenkinson DS, Brookes PC, Powlson DS (2004) Measuring soil microbial biomass. Soil Biol Biochem 36:5–7. https://doi.org/10.1016/j.soilbio.2003.10.002

Jones DL, Willett VB (2006) Experimental evaluation of methods to quantify dissolved organic nitrogen (DON) and dissolved organic carbon (DOC) in soil. Soil Biol Biochem 38:991–999

Kristensen JÅ, Rousk J, Metcalfe DB (2020) Below-ground responses to insect herbivory in ecosystems with woody plant canopies: A meta-analysis. J Ecol 108:917–930

Kuznetsova A, Brockhoff PB, Christensen RHB (2017) lmerTest Package: Tests in Linear Mixed Effects Models. J Stat Softw 82:1–26. https://doi.org/10.18637/jss.v082.i13

Lavorel S, Grigulis K, McIntyre S, et al (2008) Assessing functional diversity in the field – methodology matters! Funct Ecol 22:134–147. https://doi.org/10.1111/j.1365-2435.2007.01339.x

Legay N, Grassein F, Robson TM, et al (2013) Comparison of inorganic nitrogen uptake dynamics following snowmelt and at peak biomass in subalpine grasslands. Biogeosciences Discuss 10:8887–8917

Levy EG, Madden EA (1933) The point method of vegetation analysis. N Z J Agric 46:267–279

Lovett GM, Ruesink AE (1995) Carbon and nitrogen mineralization from decomposing gypsy moth frass. Oecologia 104:133–138. https://doi.org/10.1007/BF00328577

Malik AA, Martiny JBH, Brodie EL, et al (2020) Defining trait-based microbial strategies with consequences for soil carbon cycling under climate change. ISME J 14:1–9. https://doi.org/10.1038/s41396-019-0510-0

Malik AA, Puissant J, Goodall T, et al (2019) Soil microbial communities with greater investment in resource acquisition have lower growth yield. Soil Biol Biochem 132:36–39. https://doi.org/10.1016/j.soilbio.2019.01.025

McNaughton SJ (1983) Compensatory plant growth as a response to herbivory. Oikos 329–336

McNaughton SJ, Banyikwa FF, McNaughton MM (1997) Promotion of the cycling of diet-enhancing
nutrients by African grazers. Science 278:1798–1800

McNaughton SJ, Ruess RW, Seagle SW (1988) Large Mammals and Process Dynamics in African
Ecosystems. BioScience 38:794–800. https://doi.org/10.2307/1310789

Nitschke N, Wiesner K, Hilke I, et al (2014) Increase of fast nutrient cycling in grassland microcosms
through insect herbivory depends on plant functional composition and species diversity.
Oikos

Parker TC, Sadowsky J, Dunleavy H, et al (2017) Slowed biogeochemical cycling in sub-arctic birch
forest linked to reduced mycorrhizal growth and community change after a defoliation event.
Ecosystems 20:316–330

Pastor J, Dewey B, Naiman RJ, et al (1993) Moose browsing and soil fertility in the boreal forests of
Isle Royale National Park. Ecology 74:467–480

Paterson E, Thornton B, Sim A, Pratt S (2003) Effects of defoliation and atmospheric CO_2 depletion
on nitrate acquisition, and exudation of organic compounds by roots of Festuca rubra. Plant
Soil 250:293–305

Piton G, Foulquier A, Martínez-García LB, et al (2020a) Disentangling drivers of soil microbial
potential enzyme activity across rain regimes: An approach based on the functional trait
framework. Soil Biol Biochem 148:107881. https://doi.org/10.1016/j.soilbio.2020.107881

Piton G, Legay N, Arnoldi C, et al (2020b) Using proxies of microbial community-weighted means
traits to explain the cascading effect of management intensity, soil and plant traits on
ecosystem resilience in mountain grasslands. J Ecol 108:876–893.
https://doi.org/10.1111/1365-2745.13327

Potthast K, Meyer S, Tischer A, et al (2021) Grasshopper herbivory immediately affects element
cycling but not export rates in an N-limited grassland system. Ecosphere 12:e03449.
https://doi.org/10.1002/ecs2.3449

R Core Team (2019) R: A language and environment for statistical computing.

Reich PB (2014) The world-wide 'fast–slow' plant economics spectrum: a traits manifesto. J Ecol
102:275–301

Rhoades DF (1985) Offensive-defensive interactions between herbivores and plants: their relevance
in herbivore population dynamics and ecological theory. Am Nat 125:205–238

Ritchie ME, Tilman D, Knops JMH (1998) Herbivore effects on plant and nitrogen dynamics in oak
savanna. Ecology 79:13

Robson TM, Lavorel S, Clement J-C, Roux XL (2007) Neglect of mowing and manuring leads to
slower nitrogen cycling in subalpine grasslands. Soil Biol Biochem 39:930–941.
https://doi.org/10.1016/j.soilbio.2006.11.004

Sandén H, Mayer M, Stark S, et al (2020) Moth outbreaks reduce decomposition in subarctic forest
soils. Ecosystems 23:151–163

Sankaran M, Augustine DJ (2004) Large herbivores suppress decomposer abundance in a semiarid
grazing ecosystem. Ecology 85:1052–1061

Satterthwaite FE (1946) An approximate distribution of estimates of variance components. Biom Bull
2:110–114

Schimel JP, Bennett J (2004) Nitrogen mineralization: challenges of a changing paradigm. Ecology
85:591–602

Schmitz OJ (2008) Effects of predator hunting mode on grassland ecosystem function. Science
319:952–954

Schultz JC, Baldwin IT (1982) Oak Leaf Quality Declines in Response to Defoliation by Gypsy Moth
Larvae. Science 217:149–151. https://doi.org/10.1126/science.217.4555.149

Singer FJ, Schoenecker KA (2003) Do ungulates accelerate or decelerate nitrogen cycling? For Ecol
Manag 181:189–204

Sinsabaugh RL, Hill BH, Follstad Shah JJ (2009) Ecoenzymatic stoichiometry of microbial organic
nutrient acquisition in soil and sediment. Nature 462:795–798.
https://doi.org/10.1038/nature08632

Stark S, Grellmann D (2002) Soil microbial responses to herbivory in an arctic tundra heath at two
levels of nutrient availability. Ecology 83:2736–2744

Talal S, Cease AJ, Youngblood JP, et al (2020) Plant carbohydrate content limits performance and
lipid accumulation of an outbreaking herbivore. Proc R Soc B 287:20202500

Thébault A, Clément J-C, Ibanez S, et al (2014) Nitrogen limitation and microbial diversity at the
treeline. Oikos 123:729–740. https://doi.org/10.1111/j.1600-0706.2013.00860.x

Tuomi M, Stark S, Hoset KS, et al (2019) Herbivore Effects on Ecosystem Process Rates in a Low-
Productive System. Ecosystems 22:827–843. https://doi.org/10.1007/s10021-018-0307-4

Van Der Wal R, Bardgett RD, Harrison KA, Stien A (2004) Vertebrate herbivores and ecosystem
control: cascading effects of faeces on tundra ecosystems. Ecography 27:242–252

van Wijnen HJ, van der Wal R, Bakker JP (1999) The impact of herbivores on nitrogen mineralization
rate: consequences for salt-marsh succession. Oecologia 118:225–231

Vance ED, Brookes PC, Jenkinson DS (1987) An extraction method for measuring soil microbial
biomass C. Soil Biol Biochem 19:703–707

Vittoz P, Guisan A (2007) How reliable is the monitoring of permanent vegetation plots? A test with
multiple observers. J Veg Sci 18:413–422

Wardle D A, Bardgett Richard D, Klironomos John N, et al (2004) Ecological Linkages Between
Aboveground and Belowground Biota. Sci Wash DC 304:1629–1633

Wienhold BJ (2007) Comparison of laboratory methods and an in situ method for estimating nitrogen
mineralization in an irrigated silt-loam soil. Commun Soil Sci Plant Anal 38:1721–1732

Williams A, Langridge H, Straathof AL, et al (2022) Root functional traits explain root exudation rate
and composition across a range of grassland species. J Ecol 110:21–33

Wood JL, Tang C, Franks AE (2018) Competitive traits are more important than stress-tolerance
traits in a cadmium-contaminated rhizosphere: a role for trait theory in microbial ecology.
Front Microbiol 9:121

Supplementary Figures & Tables

Sup. Fig. 1

Geographical distribution of the 12 study sites (F = Forbs communities, W = Warm grasslands, IN = Intensive grasslands, EX = Extensive grasslands). F1, W1 and F3 are close to each other, as well as F2, W2 and W3. Grenoble corresponds to the large city at the east.

Sup. Fig. 2

Aboveground plant biomass for grasses and forbs, in function of grassland type. There are 12 sites, 4 grassland types and 3 replicates per type. Each site is composed of 5 1m² cage.

806 # Sup. Fig. 3

807
808 Mean percentage of leaf area eaten for each plant functional group, in function of (A) grasshopper
809 treatment, and (B) grassland type. The percentage of leaf area eaten was estimated from
810 observations (mid July 2017) of the dominant plant species in each plot.
811

813 **Sup. Fig. 4**

814 Leaf C/N ratio of forbs and grasses in the four grassland types.

Sup. Fig. 5

819   Variation of the percentage of forbs estimated by the point quadrat method after one year of
820   herbivory, in function of the herbivory treatment.

Sup. Fig. 6

Principal component analysis of A. microbial, B. soil abiotic and C. vegetation characteristics. The percentage of variation explained by either the 4 grassland types or the 12 sites are given (redundancy analysis).

Sup. Tab. 1

Mean±sd of each of the 12 sites (with 5 plots per site). SWC: Soil water content (%), SOM: Soil organic matter (%), TDN: Total dissolved nitrogen (µg/g), PNM: Potential nitrogen mineralization (µgN/g dry soil/day), MBN: Microbial biomass nitrogen (µg/g), EEN: Nitrogen-related enzymes (nmol activity/ g dry soil/h), EEC: Carbon-related enzymes, EEP: Phosphorous-related enzymes, Biomass: Plant biomass (g/m²), N_B: Nitrogen in aboveground plant biomass (g/m²), Forbs: Forb biomass (g/m²). Geographical coordinates are also given.

	Extensive prairies			Intensive prairies			Forb-dominated grasslands			Warm grasslands		
	EX1	EX2	EX3	IN1	IN2	IN3	MG1	MG2	MG3	BR1	BR2	BR3
SWC	24.6 ± 2.3	27.3 ± 2.5	24.1 ± 1.6	21.2 ± 2.2	19.9 ± 3.4	33.7 ± 1.8	30.5 ± 0.6	29.1 ± 1.3	35.6 ± 1.2	45.9 ± 2.3	27.3 ± 2.2	26.6 ± 4.9
SOM	9.09 ± 1.11	9.55 ± 0.58	8.07 ± 0.39	8.45 ± 0.63	6.35 ± 0.61	12.7 ± 0.6	10.8 ± 0.9	9.53 ± 0.57	15.4 ± 1.3	32.6 ± 4	15.9 ± 1.8	14.8 ± 2
pH	5.21 ± 0.33	5.14 ± 0.1	5.04 ± 0.04	6.24 ± 0.31	4.98 ± 0.03	7.07 ± 0.1	4.15 ± 0.07	7.83 ± 0.1	6.45 ± 0.64	6.51 ± 0.16	7.69 ± 0.29	7.89 ± 0.09
TDN	31.6 ± 3.5	44.8 ± 3.5	42.2 ± 2.7	19 ± 1.1	48.5 ± 8.2	38.3 ± 3.2	36.5 ± 3	26 ± 1.9	46.5 ± 18.6	98.3 ± 23.1	46.3 ± 8	43.3 ± 8.5
Soil C/N	18.2 ± 2.3	18.2 ± 1.3	18.2 ± 2	23 ± 6.4	21.5 ± 5.6	15.4 ± 1.3	25 ± 10	23.3 ± 3.1	19 ± 3.4	16.7 ± 3.8	30.5 ± 13.3	27 ± 9.8
PNM	8.84 ± 5.41	9.76 ± 2.24	11.4 ± 4.9	9 ± 2.27	8.16 ± 1.51	8.14 ± 3.68	6.43 ± 1.45	4.72 ± 1.4	10.1 ± 1.6	28.2 ± 8.8	15 ± 3.8	16.6 ± 4.9
MBN	42.8 ± 5.6	47.7 ± 5	39.9 ± 4	49.8 ± 2.6	28.8 ± 6.9	68.3 ± 3.9	41.6 ± 18.8	54.8 ± 3.8	60.3 ± 4.7	43.9 ± 22.7	29.7 ± 14.7	50.2 ± 9.5
EEN	673 ± 68	466 ± 48	337 ± 65	451 ± 147	336 ± 50	728 ± 382	228 ± 44	1356 ± 136	637 ± 100	1155 ± 132	1155 ± 459	1429 ± 254
EEC/EEN	447 ± 52	351 ± 88	262 ± 79	511 ± 196	352 ± 98	742 ± 333	279 ± 78	483 ± 92	390 ± 130	1038 ± 705	593 ± 369	481 ± 128
EEP	1165 ± 229	1248 ± 236	1131 ± 190	838 ± 254	742 ± 125	480 ± 116	1505 ± 167	105 ± 40	651 ± 186	1827 ± 210	164 ± 127	135 ± 32
EEC/EEN	0.67 ± 0.1	0.75 ± 0.17	0.77 ± 0.1	1.12 ± 0.08	1.03 ± 0.17	1.04 ± 0.14	1.25 ± 0.34	0.35 ± 0.04	0.6 ± 0.12	0.86 ± 0.47	0.7 ± 0.81	0.34 ± 0.06
Biomass	388 ± 66	464 ± 26	465 ± 84	491 ± 89	645 ± 129	702 ± 110	440 ± 31	225 ± 47	412 ± 48	351 ± 98	391 ± 82	274 ± 55
Leaf C/N	28.1 ± 2.4	27.3 ± 2.4	28.7 ± 3.4	23.1 ± 2.2	27.8 ± 5.2	31.3 ± 2.3	23.8 ± 1.8	25.8 ± 1.9	20.5 ± 1.1	32.9 ± 3.8	34.4 ± 2.2	40.6 ± 3.8
N_B	7.35 ± 1.31	9.33 ± 0.74	8.5 ± 1.88	11.4 ± 2.6	12 ± 2.2	11.3 ± 2	9.78 ± 0.92	4.19 ± 1.03	9.94 ± 1.58	6.2 ± 2.44	6.34 ± 1.97	3.47 ± 0.62
Shannon	1.8 ± 0.16	1.69 ± 0.19	1.66 ± 0.2	2.37 ± 0.09	1.46 ± 0.2	1.98 ± 0.13	1.96 ± 0.21	2.28 ± 0.27	2.68 ± 0.03	2.07 ± 0.53	1.63 ± 0.2	0.91 ± 0.41
Forbs	161 ± 44	167 ± 58	126 ± 51	123 ± 70	36.8 ± 37.6	4.69 ± 3.31	369 ± 47	203 ± 47	376 ± 47	64.5 ± 47.9	118 ± 42	37.7 ± 33.3
Lat	45.1589114	45.1665751	45.1590965	45.1509678	45.15276	45.1630496	45.2170777	45.1687254	45.2172083	45.2185605	45.1690667	45.1693922
Long	5.52397504	5.53694229	5.53318494	5.55091331	5.52717685	5.52653451	5.55924632	5.56912743	5.55883236	5.56011453	5.56860468	5.56848548

Data accessibility

841   Data used for the analysis:

842   https://doi.org/10.5061/dryad.5tb2rbp6j

844   R script used for the analysis:

845   https://doi.org/10.5281/zenodo.6831330