Gyne production is regulated by the brood in a social bee (*Bombus impatiens*)

Etya Amsalem[*#], Priscila K. F. Santos[*], Ella Messner, Cameron S. Murray

Department of Entomology, Pennsylvania State University, University Park, PA 16802 U.S.A.

[#]Corresponding author

[*]Equal contribution

Address: 215 Huck Life Sciences building, University Park 16802

Email: eua6@psu.edu

ORCID: 0000-0001-7935-049X

Abstract

Sexual production in social insects marks the peak of colony development, yet the mechanisms regulating it remain unclear. We investigated the role of brood in colony development, worker reproduction, and sexual production in *Bombus impatiens*. While larvae are known to reduce worker egg-laying and enhance the queen's reproductive inhibition, these effects were previously tested only in small groups. We manipulated brood size in full-sized, young colonies by doubling or removing brood and monitored development. Colonies with doubled brood produced significantly more gynes, independent of the number of workers, while reduced-brood colonies exhibited a non-significant increase in male production that was driven by colony size. Worker ovary activation was lower in double-brood colonies, with no change in aggression. A follow-up experiment directly testing the effect of colony size showed that higher worker density led to higher ovary activation in workers but did not affect sexual production. These results suggest that brood strongly influences colony development and sexual production, possibly reflecting an extended phenotype of the queen, whereas worker ovary activation appears to be a more flexible process influenced by either brood presence or colony size. Understanding brood dynamics may be key to understanding the evolution of female castes in social insects.

Keywords: brood, sociality, density, caste determination, gynes, reproduction, aggression

Introduction

Social insects can form giant colonies; however, most individuals are sterile helpers produced to support the generation of a limited number of sexuals. Annual social species like bumble bees (1) and social wasps (2) slowly build up the worker force to support sexual production in a single event via a strategy called "bang-bang" theory (an abrupt shift from producing workers to sexuals) (3), whereas perennial species like honey bees and many species of ants alternate between cycles of worker and sexual production. In hymenopteran social insects, gynes (new queens) are morphologically, physiologically, and behaviourally different from female workers (4) and their life trajectory is often determined during larval development (5, 6). The determination of female caste early in development sets the foundation for the reproductive division of labour among females, and therefore, for the social organization (6). Timing the production of sexuals, and particularly of gynes, is critical for synchronizing sexual emergence with mating and floral availability and is regulated carefully to prevent loss of fitness. However, the factors triggering this event remain mostly unknown (7), and studying them is of significant importance to understanding social behaviour, resource allocation, and the evolution of female castes.

Research about the transition to sexual production in social insects has focused on a limited number of species, mostly bumble bees, honeybees, social wasps, and some ants. It is best understood through the lens of optimal resource allocation theory (8, 9), which predicts that colonies should initiate investment in sexuals once sufficient resources and workforce are secured to support reproductive success. Empirical studies support this framework but highlight the dominant role of internal sociobiological cues over environmental factors (7, 10). For instance, mathematical models and experimental data in bumble bees suggest that colony size and worker accumulation influence the optimal timing of reproduction (8, 11), yet actual triggers are often mediated by queen signalling. Queen presence (12) and age (13), pheromonal cues, and the type of brood in the colony (14, 15) regulate the onset of gyne production in *Bombus terrestris*. In ants such as *Linepithema humile* and *Aphaenogaster senilis*, queen control and pheromones, colony size and environmental seasonality jointly modulate sexual production (16-20), while in other species it might be seasonal (*Apis mellifera*) (21) or even random (*Formica exsecta*) (10). Across taxa, the lack of influence from worker age, mortality, or density (13, 22,

23) emphasize that sexual production is less a product of environmental feedback and more a

colony-driven process shaped by the need to maximize inclusive fitness.

Recent work has begun to highlight the role of brood in shaping colony-level reproductive

dynamics, particularly in Hymenoptera (24). Brood at various developmental stages, including

larvae and pupae in *Apis mellifera* (25-27), larvae in *Novomessor cockerelli* (28) and *Ooceraea*

biroi (29), eggs in *Camponotus floridanus* (30), and larvae and pupae in *Bombus impatiens* (31)

has been shown to influence female reproduction and behaviour. Brood signals not only suppress

worker reproduction but also regulate foraging behaviour, shifting workers from nursing to

foraging roles and increasing foraging effort (27, 32-34). In several species of ants, different

stages of brood were shown to stimulate egg production in fertile females. For example, late

instar larvae in fire ants (35), and the presence of larvae in pharaoh's ants drive queen

reproduction (36). It was further suggested that the brood-to-worker ratio may influence gyne

production in bumble bees (37), that workers can recognize and respond differently to sexual

versus worker brood in Argentine ants (20), and that late-stage larvae in *Monomorium pharaonis*

ants affect gyne production (38); however, the broader potential role of brood as a key trigger for

sexual production has largely been overlooked.

Here we examined how the amount of brood affects colony development and demography using

full size colonies of bumble bees (*Bombus impatiens)*. Bumble bees are an excellent system to

examine these questions since they are annual and semelparous. The life cycle of the colony

starts with a single, mated queen that lays eggs following a winter diapause. Initially, the queen

performs all the tasks in the colony but she switches exclusively to egg laying once the first

worker emerges (1). The queen monopolizes reproduction for approximately 4-5 weeks from the

first worker emergence but loses dominance as the colony grows and transitions into the

competition phase. During this phase, which highly correlates with the timing of gyne production

(39), workers compete with the queen and among themselves over male production by exhibiting

aggressive behaviour, oophagy and egg laying (40, 41). Gynes and males are typically produced

towards the end of the season, though males can be produced earlier (42). Colonies differ

substantially in the number and type of sexuals they produce, with some colonies specializing in

producing gynes and others in producing males (43). This split sex ratio is partially explained by

the diapause regime the queen experiences prior to funding a colony (43).

Recent studies in *Bombus impatiens* show that (a) young larvae decrease while pupae increase

worker egg laying (31); (b) the impact of the queen on worker ovary activation is stronger when

combined with the brood (44); and (c) the queen pheromonal secretion is only effective when

combined with brood (45). Altogether, these studies demonstrate how significant is the brood to

the social organization. To test the effects of brood on worker reproduction and sexual

production, we manipulated brood amount in 16 queenright, young colonies prior to the onset of

gyne production by transferring all brood from one colony to another. This resulted in five

colonies with no brood, five colonies with double brood, and six unmanipulated colonies at the

start of the experiment. We examined colony growth, aggressive behaviour toward and by the

queen, worker ovary activation, and the production of egg batches, brood, workers, and sexuals

over 26 days. From day 27 onward, we continued monitoring all adults emerging in the colonies

until the emergence of the last bee. In a follow-up experiment, we monitored an additional 15

colonies, each restricted to either 30 or 90 workers, for 54 days using the same protocol. We

hypothesized that an increased amount of brood would decrease worker ovarian activation and

aggression and support the production of more gynes. We further hypothesized that, with the

same amount of brood, colony size would affect worker ovary activation but not gyne

production.

Material and Methods

Bumble bee rearing. A total of 31 *Bombus impatiens* colonies were used in this study. All

colonies were obtained from Koppert Biological Systems (Howell, MI, USA) and arrived a few

weeks after the emergence of the first worker, containing a queen, several dozen workers, and

brood at various stages. Colonies were maintained in the laboratory under constant darkness,

60% relative humidity, and a temperature of 28–30 °C. They were provided with unlimited 60%

sucrose solution and fresh pollen collected by honeybees, purchased from Koppert. All handling

was performed under red light.

Experimental design. In the first experiment (Figures 1-4, S1-2), 16 colonies of approximately
equal wet mass and similar numbers of workers and brood were assigned to three treatments: in
five colonies, all brood was transferred to another five colonies, creating 'no brood' and 'double
brood' groups, respectively. The remaining six colonies were left unmanipulated and served as
controls. Because precise brood counts were not possible, 'double' refers to receiving brood
from two colonies. All colonies retained their original queen and workers. The experiment was
conducted in two consecutive replicates ('rounds'), each including eight colonies (three double
brood, three no brood, and two controls in the first round; the remainder in the second). For 26
days post-manipulation, we monitored colony growth and mass (see below), observed aggression
involving the queen, sampled workers for ovary activation at five timepoints, and recorded the
number of new egg batches daily. On day 26, we removed all bees and brood, excluding pupae.
We then counted the numbers of eggs, larvae, and pupae, and measured larval body mass. Pupae
were left in the colonies until adult emergence, at which point emerging individuals were sexed
and counted and added to the total number of individuals in each group.

In the second experiment (Figures 5-8, S3), we used 15 colonies. For each colony, all individuals
were removed, the brood wet mass was measured, and then the queen was returned along with
either 30 or 90 workers, resulting in two treatments: 30 workers (n = 8) and 90 workers (n = 7).
For 54 days following the manipulation, we repeated the same measurements as in the first
experiment, excluding observations of aggressive behaviour. Previous studies have shown that
even small colonies with as few as 20 workers are capable of producing gynes (13).

Control for colony growth: To account for differences in colony growth during the first
experiment (i.e., colonies without brood could not immediately produce new workers), newly
emerged workers (<24 hours old) were collected and redistributed daily. These workers were
equally divided among all colonies to ensure comparable growth regardless of initial brood
quantity. In each round, a minimum of 8 and a maximum of 67 workers were collected daily and
evenly distributed across the colonies. When the number of workers did not divide evenly by
eight, the remaining individuals were allocated in the following order: no brood, control, and
then double brood colonies, with balancing adjustments made on the consecutive day. In the
second experiment, colony size was maintained at 30 or 90 workers by removing workers daily.

Although these methods were generally effective, perfect standardization was not possible due to

the large number of workers per colony. Therefore, we also examined and controlled for the

effects of the initial, final, and total number of workers on the measured variables (see Results).

Colony mass: In the first experiment, colonies were weighed twice at the start, before and after

the social manipulation, and then every other day throughout the experiment using an electronic

scale. In the second experiment, colonies were first emptied of adults and food to obtain the

actual brood mass, then weighed with the adults, and subsequently weighed daily. Weighing was

performed by placing the entire colony on a scale, a method that is non-intrusive and not

stressful to the bees. The recorded mass includes the bees, brood, cells, and the plastic box

housing the colony, but excludes the sugar reservoir. All colonies were kept in their original

identical Koppert plastic boxes.

Aggressive behaviour: Colonies of the first experiment were video recorded every other day to

assess aggression towards and by the queen. Each colony was recorded for 20 minutes between 9

am and 1 pm. Approximately 70 hours of video footage were analysed by an observer blind to

the experimental design and hypotheses. Three behaviours were identified and summed: (1)

attack: overt aggression including pulling, climbing, dragging, or attempted stinging; (2) darting:

a sudden movement toward another bee without physical contact; and (3) humming: a series of

short wing vibrations directed at another bee while in motion (46-48). Behaviours performed by

the queen toward workers ("queen aggression") and by workers toward the queen ("worker

aggression") were analysed separately and reported as the average number of behaviours

observed per 20-minute recording.

Worker ovarian activation: In the first experiment, 10–15% of the workers in each colony were

collected at five timepoints (days 1, 7, 13, 19, and 26), and a subset (5–10 workers per colony

per timepoint) was dissected to assess ovary activation. The number of workers removed was

negligible relative to the total colony size and did not meaningfully affect colony dynamics.

Similar sampling approaches have been used in previous studies and are considered

representative of the workers' overall reproductive state (49-51). Since workers were not

individually tagged and newly emerged callows were redistributed daily, some dissected

individuals may have originated from other colonies. However, as these workers were added
upon emergence, their reproductive behaviour is assumed to align with that of the host colony
(48). In the second experiment, a subset of the workers removed daily was dissected at 9
timepoints (approximately every 5–10 days). To measure terminal oocyte size, the abdomen was
dissected under a stereomicroscope, and both ovaries were placed in a drop of water. The three
largest oocytes were measured using a stereomicroscope ruler, and their lengths were averaged
(52). Results are presented in millimetres.

Larval body mass. At the end of both experiments, larvae were weighed individually using an
electronic scale. Mass distribution within each colony is presented for all larvae with a body
mass greater than 0.1 g. This cutoff was selected because caste-related differences in body mass
(queen vs. worker) become detectable only at the third instar, which corresponds to
approximately 0.1 g (53). Although the precise mechanism of caste determination in *B. impatiens*
is unknown, it is unlikely to occur earlier than in *B. terrestris* (54). Our goal was to examine
whether larvae were distributed bimodally, reflecting the production of both workers/males and
gynes, or unimodally, reflecting the production of workers/males only.

Statistical analyses. Statistical analyses and visualization of the main text figures were
performed using JMP 18 Pro. Colony wet mass (Figures 1 and 5) was examined using a linear
mixed model (LMM) with repeated measures. The model included treatment (control, double
brood, no brood), day and their interaction as fixed effects, 'round' as a random factor, with
colony ID as the repeated subject and a AR(1) covariance structure to account for correlations
across time. A Generalized Linear Mixed Model (GLMM) was conducted to examine the effects
of treatment, day and their interaction on the number of egg batches laid per day in Figures 2 and
6 and the count of aggression in Figure S1. The model specified a negative binominal
distribution for the response variable and included 'round' and colony ID as random effects. The
effect of treatment, day and interaction on ovarian activation (Figures 3 and 7) was examined
using LMM, with 'round' and 'colony ID' included as random factors. The number of eggs,
larvae, pupae, males, gynes and workers at the end of the experiment, as well as the initial and
total number of workers and the initial mass of brood (Figures 4 and 8) were tested using
GLMM. We included 'round' and colony ID as random factors, and the initial and total number

of workers as covariates, whenever possible (see results for details). We ran the models with the

following distributions: normal for eggs, larvae and pupae, lognormal for the initial, end and

total number of worker and negative binomial for gynes and males. The initial mass of brood

was fitted with a normal distribution. Kaplan–Meier survival analysis was used to compare the

timing of gyne production across treatments, with differences between groups assessed using the

Log-Rank test. To test the effect of treatment on the bimodality of larval body mass distribution,

we use the ACR method. Statistical significance was accepted at α=0.05. All tests were followed

up by multiple comparisons (Tukey post hoc for all pairwise comparisons). Data are presented as

box plots excluding Figures 2 and 7 which show individual data points along with a non-

parametric regression fit.

Results

Colony wet mass (Figure 1). The analysis revealed a significant effect of treatment ($F_{2,12.9}$=22.4,

p<0.001 followed up by Tukey post-hoc test p<0.001 for no brood vs. the other treatments), day

($F_{1,148.9}$=557.3, p<0.001) and the interaction between them ($F_{2,148.9}$=67.6, p<0.001). All colonies

gained mass over time, reflecting the increase in worker populations and brood. However,

colonies in the double brood group exhibited significantly greater mass gain over time compared

to the control group, while colonies in the no brood group gained significantly less mass over the

same period.

Production of new egg batches (Figure 2). Neither treatment ($F_{2,\ 5.3}$=0.34, p=0.32) nor day ($F_{1,\ 265.5}$=0.98, p=0.32) had a significant main effect; however, their interaction was significant ($F_{2,\ 262.5}$=12.6, p<0.001), indicating that colonies in the no brood treatment exhibited a modest

increase in egg laying shortly after the onset of the experiment, peaking around day 7. This

pattern likely reflects an immediate compensatory response to brood removal. Modest increases

in egg laying were observed in the other two treatments at later time points, likely indicating the

onset of worker reproduction.

Workers' ovary activation (Figure 3). The model showed a significant main effect of treatment

($F_{2,12.2}$= 4.26, p=0.039 followed by post hoc Tukey p=0.03 for no brood vs. double brood), day

($F_{4,731.2}$= 22.79, p<0.0001 followed by Tukey test p<0.001 for d1 vs. all days and p=0.002 for

d13 vs d25) and the interaction between them ($F_{8,731.3}= 2.6$, p=0.008) indicating that treatment

effects varied across time. The interaction estimates further highlighted no differences across

treatments on day 1 and a pronounced increase in the no brood group by day 7. The no brood

reached the highest observed value on day 13 (estimate = 2.16, 95% CI: 1.64 to 2.67), exceeding

both control (1.62, 95% CI: 1.11 to 2.13) and double brood (1.17, 95% CI: 0.66 to 1.68).

Although responses declined slightly on days 19 and 25, the no brood group consistently

maintained higher values than the other treatments.

The total number of brood and adults (Figure 4). The number of workers at the end (Figure 4B)

did not vary between treatments ($F_{2,1}=40.2$, p=0.11), but both the initial number of workers

($F_{2,12.1}=5.66$, p=0.01 followed by Tukey post-hoc test p=0.02 for double brood vs. control and

p=0.04 for double brood vs. no brood, Figure 4A) and the total number of workers ($F_{2,9.4}=7.17$,

p=0.01 followed by Tukey post-hoc test p=0.01 for double brood vs. no brood, Figure 4C) were

affected by the treatment. Therefore, to test the effect of treatment on the number of brood,

males, and gynes, we included in the subsequent models both the random effects (round, colony

ID) and the initial and total number of workers as covariates.

Treatment had no effect on the number of eggs present at the end of the experiment ($F_{2,10.4}=0.5$,

p=0.62, Figure 4D). The number of larvae at the end of the experiment was affected by the

treatment ($F_{2,11}=4.79$, 0.03, Figure 4E), with more larvae found in the no brood compared to the

double brood (p=0.02). This difference was significantly associated with the total number of

workers ($F_{1,11}=8.47$, p=0.01) but not with the initial number of workers ($F_{1,11}=1.43$, p=0.25).

Running the model without the covariates did not converge. The number of pupae at the end of

the experiment (Figure 4F) was not affected by the treatment ($F_{2,10.6}=0.74$, p=0.49), but was

significantly associated with the total number of workers ($F_{1,10.6}=6.6$. p=0.02).

A treatment-only model excluding covariates showed no effect of treatment on the number of

males ($F_{2,12.5}=2.72$, p=0.1, Figure 4G). Adding the total number of workers to the model did not

converge, likely due to model complexity combined with the limited sample size. Including only

the initial number of workers maintained the non-significance of the model ($F_{2,12}=0.44$, p=0.65)

and showed no association with the covariate ($F_{1,12}=0.001$, p=0.97). Including both covariates in

the model also failed to converge.

A treatment-only model excluding covariates showed a strong effect of treatment on the number

of gynes ($F_{2,12.2}=8.21$, p =0.005, Figure 4H) with double brood colonies producing more gynes

than both the no brood (p = 0.007) and control colonies (p = 0.01). Adding the total number of

workers as a covariate resulted in a stronger treatment effect ($F_{2,10.1}=11.1$, p=0.002) and no

association with the covariate ($F_{1,10}=2.38$, p=0.15). Adding also the initial number of workers to

the model did not converge. Including the initial number of workers in the model caused it to fail

to converge. Overall, the treatments did not affect the number of eggs, pupae, or males. The

number of larvae was affected by treatment but mediated by colony size, whereas the number of

gynes was affected by treatment and could not be explained by colony size. We further

conducted a survival analysis to compare the timing of gyne production across the three

treatments. The Kaplan–Meier survival curves did not differ significantly but approached

significance (Log-Rank test: $\chi^2 = 4.94$, $df = 2$, p = 0.08), suggesting earlier gyne production in

the double brood group.

Aggression by and towards the queen (Figure S1). There were no differences in the total queen

aggression ($F_{2,12.8}=1.57$, p=0.24; Figure S1) or the total worker aggression ($F_{2,11.9}=1.6$, p=0.24).

One double brood colony was a clear outlier in the number of the aggressive behaviours

performed by and towards the queen on day 4. Thus, we also analysed the data without this

colony on day 4 (Figures S1) but this did not change the results.

Distribution of larva body mass (Figure S2). The differences between the treatments in gyne

production are reflected also in the body mass distribution of larvae that were collected and

weighed on the last day of the experiment (day 26). The larvae in double brood colonies showed

a bimodal distribution of larva mass, corresponding to larvae that will develop into

workers/males and gynes (ACR mode test, p<0.001), while control and no brood colonies

showed a unimodal distribution, indicating the production of workers or males (ACR mode test,

p>0.114).

Colony wet mass in colonies restricted to 30/90 workers (Figure 5). The analysis revealed a

significant effect of treatment ($F_{1,13.9}$=11.5, p=0.004), day ($F_{1,316.7}$=382.1, p<0.001) and the

interaction between them ($F_{1,316.7}$=20.7, p<0.001). All colonies gained mass over time, reflecting

the increase in worker populations and brood. However, colonies with 90 workers gained

significantly greater mass over time compared to colonies with 30 workers over the same period.

Production of new egg batches in colonies restricted to 30/90 workers (Figure 6). Neither the

treatment ($F_{1, 13}$=0.86, p=0.37) nor the interaction between day and treatment ($F_{1, 686.8}$=0.52,

p=0.46) had a significant main effect; however the day alone differed significantly ($F_{1, 686.8}$=71.3,

p<0.001), indicating an increase in egg laying over time in both treatments.

Workers' ovary activation in colonies restricted to 30/90 workers (Figure 7). The model showed

a significant main effect of treatment ($F_{1,16.9}$= 7.01, p=0.01), but not of day ($F_{8,639.9}$= 1.41,

p=0.18) or of the interaction ($F_{8,639.9}$= 1.35, p=0.21), indicating a higher and consistent ovarian

activation in the 90-worker treatment.

The total number of brood and adults in colonies restricted to 30/90 workers (Figure 8). The

initial mass of brood differed significantly and, despite randomization, was slightly higher in the

90-workers treatment ($F_{1,13}$=6.6, p=0.02, Figure 8A). The number of workers at the end of the

experiment was also higher in the 90-worker treatment, in line with our manipulation ($F_{1,13}$=58.3,

p<0.001, Figure 8C). While all colonies had more than 30 and 90 workers respectively, the 90-

worker colonies had, on average, three times more workers than the 30-worker colonies at the

end of the experiment. Neither the initial number of workers prior to the manipulation

($F_{1,12.9}$=1.99, p=0.18, Figure 8B) nor the total number of workers ($F_{1,13}$=0.41, p=0.53, Figure 8D)

differed significantly. Therefore, to test the effect of treatment on the number of brood, males,

and gynes, we included in the subsequent models both the random effect (colony ID) and the

initial brood mass and the end number of workers as covariates.

The number of eggs present at the end of the experiment was not affected by the treatment

($F_{1,11}$=1.31, p=0.27, Figure 8E) or associated with the initial brood mass ($F_{1,11}$=1.26, p=0.28) or

the end number of workers ($F_{1,11}$=3.47, p=0.09). The number of larvae at the end of the

experiment was not affected by the treatment ($F_{1,11}=0.93$, p=0.35, Figure 8F) or associated with

the initial brood mass ($F_{1,11}=1.14$, p=0.3) or the end number of workers ($F_{1,11}=1.81$, p=0.2).

Similar results were obtained for pupae (Treatment: $F_{1,11}=0.15$, p=0.7; Initial mass: $F_{1,11}=0.001$,

p=0.97; End workers: $F_{1,11}=0.03$, p=0.85, Figure 8G). The number of males produced by the

colonies did not differ between treatments ($F_{1,10.4}= 2.77$, p=0.12, Figure 8H) and was not

associated with any of the covariates (Initial mass: $F_{1,11}=2.79$, p=0.12; End workers: $F_{1,10.4}=1.14$,

p=0.3). The number of gynes in the two treatments is not presented since only one colony out of

15 produced any gynes during the experiment. This colony was in the 30-worker treatment and

produced 3 gynes in total. All in all, the manipulation of the number of workers did not affect the

number of brood or sexuals in the colonies.

Distribution of larva body mass in colonies restricted to 30/90 workers (Figure S3). The larvae

collected from the colonies at the end of the experiment showed a unimodal distribution

indicating the production of workers and/or males only (ACR mode test, p>0.4).

Discussion

In this study, we demonstrate that manipulating brood quantity in a social colony significantly

impacts colony development, gyne production, and worker ovary activation, while showing no

significant effect on aggressive behaviour or male production. Differences in gyne numbers

between treatments could not be explained by the number of workers, suggesting that brood may

function as an extended phenotype of the queen, influencing not only individual reproductive

decisions of workers but also colony-level strategies for investing in future queens. Our second

experiment, in which colonies were restricted to 30 or 90 workers, unexpectedly resulted in only

one colony producing gynes and no significant differences in male production. However, clear

differences emerged in worker ovarian activation, with larger colonies showing higher ovary

activation of workers. Based on data from both experiments, we suggest that worker ovary

activation is a flexible process influenced by colony size, density, or brood amount, whereas

sexual production, particularly of gynes, is primarily determined by the amount of brood

produced by the colony. Below, we discuss evolutionary explanations for the reliance on

sociobiological cues in regulating sexual production, within the framework of optimal resource

allocation theory. We also address the evolution of female castes in social insects, along with key

caveats and directions for future research.

Optimal resource allocation theories predict that insects secure sufficient resources before

switching to sexual production. Evidence across insect species indicates that this transition is

influenced not only by external ecological factors but also by internal sociobiological cues (8,

55). These internal cues include queen signalling, queen age, queen presence, and, as

demonstrated in this study, brood presence. Such cues are likely more reliable than ecological

factors like weather, which may fluctuate, or the number of workers, which reflects past

conditions and cannot be reliably extrapolated to predict future ones. Brood may offer a more

accurate estimate of ecological resources available for brood rearing, the queen's fecundity, and

the workers' capacity to support gyne production. Our second experiment demonstrates that

colony size alone is not a sufficient cue for gyne production and suggests that a threshold

amount of brood is required to trigger its onset. This finding aligns with previous models

predicting that bumblebee colonies optimally initiate sexual production approximately one brood

development period before colony termination to maximize growth and ensure the successful

maturation of sexual offspring (11).

In contrast to gyne production, bumblebee workers appear to rely on multiple cues to initiate

male production. These include brood amount and the number of workers as found in this study

and also in (31, 56), colony and/or queen age (57), and queen fecundity and health (58). When

brood levels are high, workers may delay reproduction to allow for gyne rearing, from which

they gain greater indirect fitness benefits compared to males (59). In contrast, the absence of

brood suggests that gynes will not be reared, making delayed reproduction unnecessary.

Additionally, the number of workers may signal the approaching end of the season, prompting

workers to seize their final opportunity for direct reproduction. This supports the idea that

worker reproduction is more flexible and less risky than gyne production, likely because males

can be produced by both queen and workers (60) and are less costly to produce .

Previous studies have shown that gyne production and the onset of worker reproduction are

highly correlated events (39). Ovarian activation by workers during the competition phase occurs

in most colonies, but only some colonies produce gynes. Therefore, these correlations apply only

to colonies where both events occur. If the onset of worker reproduction is triggered by various

cues - only one of which also triggers gyne production - this could explain both the observed

variation among colonies and the correlation between the two events in colonies that exhibit

both. Data from other species support this pattern; for example, in *Bombus perplexus*, an increase

in worker number is positively correlated with male production, but not necessarily with queen

production (61).

A few more points are worth mentioning. First, relatedness could potentially affect the results.

All colonies contained non-related workers (as newly emerged workers were distributed daily

across all the colonies, including the controls), but only the double brood colonies contained

partially unrelated brood, raising the question whether gynes were more likely to develop in

colonies with mixed brood. In a previous study we did not find any evidence of the impact of

relatedness on brood care (31) and therefore consider it unlikely to have an impact here. We also

did not observe any differential behaviour towards the brood in our study in any of the colonies.

Whether unrelated brood is more likely to develop into gynes compared to related brood is a

question worth investigating. Second, despite our limited number of colonies, we remain highly

confident in the effect brood has on the number of sexuals for several reasons: the substantial

effect size, the relatively low variation in the number of gynes and males in both the control and

no brood colonies compared to the double brood colonies, and our study design which

effectively controlled for many confounding factors inherent in the variable nature of a colony.

Conclusions

Overall, our results show that brood regulates not only egg laying in small groups of workers, as

previously demonstrated (31), but also influences colony-level events such as the onset of

reproductive competition and the timing of gyne production. We also found that worker ovarian

activation is associated with the total number of workers or colony density, even among colonies

with similar brood amounts. These findings suggest that brood serves as an indicator of queen

fecundity, promoting gyne production when abundant, and possibly triggering the production of

cheaper males when absent. In contrast, density alone, though sufficient to induce worker ovary

activation, is not a reliable cue for gyne production. Our findings in bumble bees are consistent

with results from annual wasps, where partial brood removal increased queen egg laying but did not suppress worker reproduction (62). This may point to a general mechanism in annual social species, in which brood irreversibly influences the level of cooperation between queens and workers and affects sexual production. Based on our second experiment directly testing the effect of density, gyne production likely requires a threshold amount of brood; without it, colonies may instead invest in male production. Indeed, the only colony that produced gynes had 312 larvae at the end of the experiment, while the average in all other colonies was 193.2 ± 19 larvae (mean $\pm$ SE). Altogether, our findings highlight the central role of brood in maintaining and shaping social organization in social insects and underscore the need to explore its diverse functions across taxa.

Declarations

Funding. This work was funded by the National Science Foundation IOS-1942127 to EA.

Conflict of interests. The authors declare no conflicts of interests

Availability of data and material. The data used and/or analyzed during the current study are available as supplementary material.

Acknowledgements. We would like to thank John M. Craig, Andrea Medero and Katie Barie for assisting in the experiments. We also thank the two anonymous reviewers for their thoughtful and constructive comments, which significantly improved the manuscript.

Author's contributions. EA, PKFS and EM designed the experiments, PKFS and EM collected the data, CSM performed the behavioral observations, EA analyzed the data, generated the figures and wrote the manuscript. PKFS and EM generated the figures in the supplementary materials. All authors edited the manuscript and gave final approval for publication.

References

1.   Amsalem E, Grozinger CM, Padilla M, Hefetz A. The physiological and genomic bases of
bumble bee social behaviour. In: Amro Z, Clement FK, editors. Genomics, Physiology and
Behaviour of Social Insects. 48. Adv In Insect Phys: Academic Press; 2015. p. 37-93.
2.   Jandt JM, Tibbetts EA, Toth AL. Polistes paper wasps: a model genus for the study of
social dominance hierarchies. Insectes Sociaux. 2013;61(1):11-27.
3.   Macevicz S, Oster G. Modeling social insect populations II: Optimal reproductive
strategies in annual eusocial insect colonies. Behavioral Ecology and Sociobiology.
1976;1(3):265-82.
4.   Treanore ED, Derstine N, Amsalem E. What can mechanisms underlying derived traits
tell us about the evolution of social behavior? Annals of the Entomological Society of America.
2020;114(5):547-61.
5.   Trible W, Kronauer DJ. Caste development and evolution in ants: it's all about size. J Exp
Biol. 2017;220(Pt 1):53-62.
6.   Wilson EO. The evolution of caste systems in social insects. Proceedings of the American
Philosophical Society. 1979;123(4):204-10.
7.   Hovestadt T, Degen T, Mitesser O. Suitable triggers for timing the transition from worker
to sexual production in annual eusocial insects. Insectes Sociaux. 2018;65(4):609-17.
8.   Hovestadt T, Mitesser O, Poethke HJ. The evolution of optimal timing strategies in
annual eusocial insects: colony size, worker mortality and the timing of reproduction. Oikos.
2018;127:964-73.
9.   Oster GF, Wilson EO. Caste and ecology in social insects. Princeton University Press.
1979.
10.   Liautard C, Brown WD, Helms KR, Keller L. Temporal and spatial variations of gyne
production in the ant Formica exsecta. Oecologia. 2003;136(4):558-64.
11.   Beekman M, Lingeman R, Kleijne FM, Sabelis MW. Optimal timing of the production of
sexuals in bumblebee colonies. Entomologia Experimentalis et Applicata. 1998;88:147-54.
12.   Lopez-Vaamonde C, Brown RM, Lucas ER, Pereboom JJM, Jordan WC, Bourke AFG. Effect
of the queen on worker reproduction and new queen production in the bumble beeBombus
terrestris. Apidologie. 2007;38(2):171-80.
13.   Alaux C, Jaisson P, Hefetz A. Reproductive decision-making in semelparous colonies of
the bumblebee bombus terrestris. Behavioral Ecology and Sociobiology. 2005;59(2):270-7.
14.   Alaux C, Jaisson P, Hefetz A. Regulation of worker reproduction in bumblebees (*Bombus
terrestris*): workers eavesdrop on a queen signal. Behavioral Ecology and Sociobiology.
2006;60(3):439-46.
15.   Bloch G. Regulation of queen-worker conflict in bumble-bee (*Bombus terrestris*)
colonies. Proc Biol Sci. 1999;266(1437):2465-9.
16.   Vargo EL, Passera L. Pheromonal and behavioral queen control over the production of
gynes in the Argentine ant Iridomyrmex humilis (Mayr). Behavioral Ecology and Sociobiology.
1993;28:161-9.
17.   Boulay R, Hefetz A, Cerdá X, Devers S, Francke W, Twele R, et al. Production of sexuals in
a fission-performing ant: dual effects of queen pheromones and colony size. Behavioral Ecology
and Sociobiology. 2007;61(10):1531-41.

18.   Vargo EL, Passera L. Gyne development in the Argentine ant *Iridomyrmex humilis*: role of
overwintering and queen control. Physiological Entomology. 1992;17:193-201.
19.   Passera L, Aron S, Bach D. Elimination of sexual brood in the Argentine ant Linepithema
humile: queen effect and brood recognition. Entomologia Experimentalis et Applicata.
1995;75:203-12.
20.   Aron S, Keller L, Passera L. Role of resource availability on sex, caste and reproductive
allocation ratios in the Argentine ant Linepithema humile. Journal of Animal Ecology.
2001;70:831-9.
21.   Seeley TD. Honeybee democracy. Princeton, New Jersey: Princeton University Press;
2010.
22.   Shykoff JA, Muller CB. Reproductive decisions in bumble-Bee colonies: The influence of
worker mortality in *Bombus terrestris* (Hymenoptera, Apidae). Functional Ecology.
1995;9(1):106-12.
23.   Muller C, Schmid-Hempel P. Variation in life-history pattern in relation to worker
mortality in the bumble-bee, Bombus lucorum. Functional Ecology. 1992:48-56.
24.   Schultner E, Oettler J, Helantera H. The role of brood in eusocial Hymenoptera. Q Rev
Biol. 2017;92(1):39-78.
25.   Mohammedi A, Paris A, Crauser D, Le Conte Y. Effect of aliphatic esters on ovary
development of queenless bees (*Apis mellifera L.*). Naturwissenschaften. 1998;85(9):455-8.
26.   Jay SC. Ovary development of worker honeybees when separated from worker brood by
various methods. Canadian Journal of Zoology. 1972;50(5):661-4.
27.   Maisonnasse A, Lenoir JC, Beslay D, Crauser D, Le Conte Y. E-beta-ocimene, a volatile
brood pheromone involved in social regulation in the honey bee colony (*Apis mellifera*). PLoS
One. 2010;5(10):e13531.
28.   Ebie JD, Holldobler B, Liebig J. Larval regulation of worker reproduction in the
polydomous ant *Novomessor cockerelli*. Naturwissenschaften. 2015;102(11-12):72.
29.   Ravary F, Jahyny B, Jaisson P. Brood stimulation controls the phasic reproductive cycle of
the parthenogenetic ant Cerapachys biroi. Insectes Sociaux. 2006;53(1):20-6.
30.   Endler A, Liebig J, Schmitt T, Parker JE, Jones GR, Schreier P, et al. Surface hydrocarbons
of queen eggs regulate worker reproduction in a social insect. Proc Natl Acad Sci U S A.
2004;101(9):2945-50.
31.   Starkey J, Brown A, Amsalem E. The road to sociality: brood regulation of worker
reproduction in the simple eusocial bee *Bombus impatiens*. Animal Behaviour. 2019;154:57-65.
32.   Pankiw T. Brood pheromone regulates foraging activity of honey bees (Hymenoptera:
Apidae). J Econ Entomol. 2004;97(3):748-51.
33.   Pankiw T, Page Jr RE, Kim Fondrk M. Brood pheromone stimulates pollen foraging in
honey bees (*Apis mellifera*). Behavioral Ecology and Sociobiology. 1998;44(3):193-8.
34.   Ulrich Y, Burns D, Libbrecht R, Kronauer DJ. Ant larvae regulate worker foraging behavior
and ovarian activity in a dose-dependent manner. Behav Ecol Sociobiol. 2016;70(7):1011-8.
35.   Tschinkel WR. Stimulation of fire ant queen fecundity by a highly specific brood stage.
Annals of the Entomological Society of America. 1995;88(6):876-82.

36.   Børgesen LW, Jensen PV. Influence of larvae and workers on egg production of queens
of the pharaoh's ant, Monomorium pharaonis (L.) (Hymenoptera: Formicidae). Insectes Sociaux.
1995;42(1):103-12.
37.   Free JB. Queen production in colonies of bumblebees. Proceedings of the Royal
Entomological Society of London Series A, General Entomology. 2009;30(1-3):19-25.
38.   Warner MR, Kovaka K, Linksvayer TA. Late-instar ant worker larvae play a prominent
role in colony-level caste regulation. Insectes Sociaux. 2016;63(4):575-83.
39.   Cnaani J, Robinson GE, Bloch G, Borst D, Hefetz A. The effect of queen-worker conflict
on caste determination in the bumblebee *Bombus terrestris*. Behavioral Ecology and
Sociobiology. 2000;47(5):346-52.
40.   Duchateau MJ, Velthuis HHW. Development and reproductive strategies in *Bombus*
terrestris colonies. Behavior. 1988;107:186-207.
41.   Amsalem E, Shamia D, Hefetz A. Aggression or ovarian development as determinants of
reproductive dominance in *Bombus terrestris*: interpretation using a simulation model. Insectes
Sociaux. 2013;60(2):213-22.
42.   Holland JG, Guidat FS, Bourke AF. Queen control of a key life-history event in a eusocial
insect. Biol Lett. 2013;9(3):20130056.
43.   Duchateau MJ. Sex ratio variation in the bumblebee *Bombus terrestris*. Behavioral
Ecology. 2004;15(1):71-82.
44.   Orlova M, Starkey J, Amsalem E. A small family business: synergistic and additive effects
of the queen and the brood on worker reproduction in a primitively eusocial bee. J Exp Biol.
2020;223(Pt 3):jeb217547.
45.   Orlova M, Amsalem E. Bumble bee queen pheromones are context-dependent. Sci Rep.
2021;11(1):16931.
46.   Duchateau MJ. Agonistic behaviours in colonies of the bumblebee *Bombus terrestris*.
Journal of Ethology. 1989;7(2):141-51.
47.   Amsalem E, Grozinger CM. Evaluating the molecular, physiological and behavioral
impacts of CO(2) narcosis in bumble bees (Bombus impatiens). J Insect Physiol. 2017;101:57-65.
48.   Amsalem E, Hefetz A. The appeasement effect of sterility signaling in dominance
contests among *Bombus terrestris* workers. Behavioral Ecology and Sociobiology.
2010;64(10):1685-94.
49.   Padilla M, Amsalem E, Altman N, Hefetz A, Grozinger CM. Chemical communication is
not sufficient to explain reproductive inhibition in the bumblebee *Bombus impatiens*. Royal
Society Open Science. 2016;3(10):160576.
50.   Treanore ED, Kiner JM, Kerner ME, Amsalem E. Shift in worker physiology and gene
expression pattern from reproductive to diapause-like with colony age in the bumble bee
Bombus impatiens. J Exp Biol. 2020;223(Pt 9):jeb218768.
51.   Cnaani J, Schmid-Hempel R, Schmidt JO. Colony development, larval development and
worker reproduction in *Bombus impatiens Cresson*. Insectes Sociaux. 2002;49(2):164-70.
52.   Amsalem E, Twele R, Francke W, Hefetz A. Reproductive competition in the bumble-bee
Bombus terrestris: do workers advertise sterility? Proc Biol Sci. 2009;276(1660):1295-304.

53.   Hefetz A, Robinson GE, Huang Z, Borst DW, Cnaani J. Caste Determination in Bombus
terrestris: Differences in Development and Rates of JH Biosynthesis between Queen and
Worker Larvae. J Insect Physiol. 1997;43(4):373-81.
54.   Barie K, Amsalem E. The influence of the social environment on larval development and
resulting caste in Bombus impatiens. bioRxiv. 2022:2022.09.09.507355.
55.   Mitesser O, Weissel N, Strohm E, Poethke HJ. Adaptive dynamic resource allocation in
annual eusocial insects: environmental variation will not necessarily promote graded control.
BMC Ecol. 2007;7:16.
56.   Amsalem E, Hefetz A. The effect of group size on the interplay between dominance and
reproduction in *Bombus terrestris*. PLoS One. 2011;6(3):e18238.
57.   Alaux C. Regulation of worker reproduction in bumblebees (Bombus terrestris): workers
eavesdrop on a queen signal. Behavioral Ecology and Sociobiology. 2006;60(3):439-46.
58.   Collins DH, Prince DC, Donelan JL, Chapman T, Bourke AFG. Costs of reproduction are
present but latent in eusocial bumblebee queens. BMC Biol. 2023;21(1):153.
59.   Bourke AF, Ratnieks FL. Kin-selected conflict in the bumble-bee *Bombus terrestris*
(Hymenoptera: Apidae). Proc Biol Sci. 2001;268(1465):347-55.
60.   Alaux C, Savarit F, Jaisson P, Hefetz A. Does the queen win it all? Queen-worker conflict
over male production in the bumblebee, *Bombus terrestris*. Naturwissenschaften.
2004;91(8):400-3.
61.   Pomeroy N, Plowright RC. The relation between worker numbers and the production of
males and queens in the bumble bee Bombus perplexus. Canadian Journal of Zoology.
1982;60:954-7.
62.   Liebig J, Monnin T, Turillazzi S. Direct assessment of queen quality and lack of worker
suppression in a paper wasp. Proc Biol Sci. 2005;272(1570):1339-44.
609
610

Legend to figures

Figure 1: The average colony wet mass throughout the experiment in the three treatment groups:

no brood, double brood, and control. Timepoints 0 and 1 refers to the colony wet mass before

and after brood manipulation, respectively.

Figure 2: The number of egg batches laid per day throughout the experiment in the three

treatment groups: no brood, double brood, and control.

Figure 3: The average terminal oocyte size in workers in the three treatment groups: no brood,

double brood, and control. Oocyte size was measured in five timepoints throughout the

experiment using a subset of workers from each colony.

Figure 4: The effect of treatment (no brood, double brood and control) on the initial number of

workers (A), the number of workers at the end of the experiment (B), the total number of

workers produced (C), the number of eggs (D), larvae (E), and pupae (F) found at the end of the

experiment, and the number of males (F) and gynes (H) produced in total. Different letters

denote statistical differences at α=0.05.

Figure 5: The average colony wet mass throughout the experiment in colonies limited in size to

30 or 90 workers.

Figure 6: The number of egg batches laid per day throughout the experiment in colonies limited

in size to 30 or 90 workers.

Figure 7: The average terminal oocyte size in workers in colonies restricted to 30 or 90 workers.

Oocyte size was measured in nine timepoints throughout the experiment using a subset of

workers from each treatment.

Figure 8: The effect of treatment (30/90 worker colonies) on the initial brood mass (A), The

initial number of worker before the manipulation (B), The number of workers found at the end of

the experiment (C), the total number of workers produced (D), the number of eggs (E), larvae

(F), and pupae (G) found at the end of the experiment, and the number of males (H) produced in

total. The total number of gynes produced are not shown since only one colony produced gynes.

Different letters denote statistical differences at α=0.05.

Figure 1

643
644

645 <div style="text-align:center">Figure 2</div>

646
647

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7

Figure 8