Gyne production is regulated by the brood in a social bee (*Bombus impatiens*)

Priscila K. F. Santos, Cameron S. Murray, Etya Amsalem*

Department of Entomology, Pennsylvania State University, University Park, PA 16802 U.S.A.

*Corresponding author

Address: 215 Huck Life Sciences building, University Park 16802

Email: eua6@psu.edu

ORCID: 0000-0001-7935-049X

Abstract

Sexual production in social insects marks the peak of colony development, yet the mechanisms regulating it remain unclear. We investigated the role of brood in colony development and sexual production in *Bombus impatiens*. While larvae are known to reduce worker egg-laying and enhance the queen's reproductive inhibition, these effects were previously tested only in small groups. We manipulated brood size in full-size, young colonies by doubling or removing brood and monitored development. Colonies with doubled brood produced significantly more gynes, independent of colony size, while reduced-brood colonies showed a slight, non-significant increase in male production. Worker reproduction was delayed in double-brood colonies, with no change in aggression. These findings suggest brood strongly influences colony development, possibly acting as an extended phenotype of the queen or shaping worker behaviour. Understanding brood dynamics may be key to unravelling female caste determination in social insects.

Keywords: brood, sociality, caste determination, gynes, reproduction, aggression

Introduction

Social insects can produce giant colonies, however most of the individuals are sterile helpers that are produced to support the production of a limited number of sexuals. Annual social species like bumble bees (1) and social wasps (2) slowly build up the worker force to support sexual production in a single event via a strategy called "bang-bang" theory (an abrupt shift from producing workers to sexuals) (3), whereas perennial species like honey bees and many species of ants alternate between cycles of worker and sexual production. Timing the production of sexuals is critical for synchronizing sexual emergence with mating and floral availability and is regulated carefully to prevent loss of fitness. However, the factors triggering this event remain mostly unknown (4).

In hymenopteran social insects, gynes (new queens) are morphologically, physiologically, and behaviourally different from female workers (5) and their life trajectory is often determined during larval development (6, 7). The determination of female caste early in development sets the foundation for the reproductive division of labour among females, and therefore, for the social organization (7). For example, in bumble bees, the timing of gyne production is highly correlated with the onset of worker reproduction (8). Thus, understanding the triggers for producing sexuals is of a significant importance to the studies of social behaviour and the evolution of female castes.

The triggers leading to sexual production in social insects are poorly understood and research has focused on a limited number of species, mostly bumble bees, honeybees, social wasps, and some ants. Sexual production, particularly of gynes, is only weakly correlated with environmental factors (4, 9), and is more strongly associated with sociobiological factors related to the queen. For example, preventing workers from contacting the queen (10), exposing workers to gyne larvae (11), manipulating the queen's age (12) and replacing the queen's eggs with male eggs (13) all contributed to early gyne production in bumble bees and/or wasps. Contrary, manipulating workers age, density, or mortality have little to no effect on the onset of gyne production (12, 14, 15). Gyne production is sometimes fixed in time, as in *Polistes* wasps (*P. gallicus L*) (16), stochastic as in *Formica exsecta* (9) or seasonal as in perennial species like the honeybee, where gyne production is controlled by the queen and gynes are produced as the

population increases throughout the season (17). Gyne production is controlled by the queen also

in Argentine ants, (18), and in *Aphaenogaster senilis*, where both the queen pheromone and

colony size play a role in regulating sexual production (19). Overall, data on the triggers leading

to sexual production vary and there is no single explanation shared across taxa.

In recent years, there was a significant growth in evidence demonstrating the role of the brood in

shaping the social environment in Hymenoptera (20). Studies showed that different stages of

brood regulate worker reproduction in several species, including larva and pupae *Apis mellifera*

(21-23), larvae in *Novomessor cockerelly*, eggs in *Camponotus floridanus* (24, 25), larvae in

Ooceraea biroi (26), and larvae and pupae in *Bombus impatiens* (27). The brood can also

regulate workers' task allocation like accelerating the transition from nursing to foraging tasks,

increasing the number of foraging trips and the size of pollen loads in *Apis mellifera* (23, 28, 29),

and increasing the foraging activity of *Oocerae biroi* (30). Furthermore, the ratio between the

brood and workers was suggested to regulate queen production in several bumble bee species in

an earlier study (31). However, the data are mostly descriptive and the role of brood in regulating

sexual production remained overlooked.

Here we examined directly how the amount of brood affects colony development and

demography using full size colonies of bumble bees (*Bombus impatiens*). Bumble bees are an

excellent system to examine these questions since they are annual and semelparous. The life

cycle of the colony starts with a single, mated queen that lays eggs following a winter diapause.

Initially, the queen performs all the tasks in the colony but switches exclusively to egg laying

once the first worker emerges (1). The queen monopolizes reproduction for approximately 4-5

weeks following the first emergence but loses dominance as the colony grows and transitions

into the competition phase. During this phase, that highly correlates with the timing of gyne

production (8), workers compete with the queen and among themselves on male production by

exhibiting aggressive behaviour, oophagy and egg laying (32, 33). Gynes are produced towards

the end of the season, and typically also males, though males can be produced earlier (34), and

colonies differ substantially in the number and type of sexuals they produce, with some colonies

specializing in producing gynes and others in producing males (35). This split sex ratio was

partially explained in the diapause regime queen experienced prior to funding a colony (35).

Recent studies in this species show that (a) young larvae decrease while pupae increase worker egg laying (27); (b) the impact of the queen on worker ovary activation is stronger when combined with the brood (36), and (c) the queen pheromonal secretion is only effective when combined with brood (37). Altogether, these studies demonstrate how significant is the brood to the social organization. To test the effects of brood on worker reproduction and sexual production, we manipulated the amount of brood in queenright, young colonies, prior to the stage of gyne production by transferring all the brood from one colony to another, resulting in colonies with no brood (nb, n=5), or with a double amount of brood (db, n=5) at the onset of the experiment. These colonies were compared to unmanipulated colonies that served as control (c, n=6). We examined colony growth in all 16 colonies, aggressive behaviour towards and by the queen, worker ovary activation, and the production of egg batches, brood, workers and sexuals for 26 days. We continued monitoring all the adults that emerged in the colonies from day 27 until the emergence of the last bee. We hypothesized that an increased amount of brood will decrease worker reproduction and aggression, and a larger amount of brood is expected to support an earlier production and a larger number of gynes. Since the brood serve as an extended phenotype of the queen, reduced amount of brood can be potentially used by workers to underestimate the quality of the queen and to lead to aggression and early worker reproduction of males.

Material and Methods

Bumble bee rearing. 16 *Bombus impatiens* colonies at the approximate age of 3-4 weeks from the emergence of the first worker were obtained from Koppert Biological Systems (Howell, MI, USA). All colonies contained a queen, workers (72 ± 5, mean ± SE, the number of workers per colony is provided in Table S1), and brood of different developmental stages. The colonies were kept in the laboratory under constant darkness, 60% relative humidity, and temperature of 28-30° C. They were provided with unlimited 60% sugar solution and fresh pollen collected by honeybees that was purchased from Koppert. Colonies were handled under red light.

Experimental design. Sixteen colonies of approximately the same wet mass (see Methods and Figure 1) and approximately the same number of workers and brood (Table S1) were assigned to

three treatments. Six colonies containing queen, workers and brood remained unmanipulated and served as controls (c). The remaining 10 colonies were randomly divided to two, and all the brood from five colonies was removed and transferred to the other five, resulting in colonies with no brood (nb) and in colonies with doubled amount of brood (db) in the first day of the experiment. Since we were not able to count the brood precisely, the 'double' refers to brood from two colonies. All colonies remained with their own original queen and workers. The experiment was conducted in two consecutive replicates, each containing eight colonies. Each replicate included all three treatments. Colonies were provided unlimited food and kept in the conditions above. During the first 26 days, we controlled for colony growth (see below), observed aggression towards and by the queen (208 observations of 20 min each, 65-78 per treatment, see below), sampled workers for ovary activation (a subset of workers were sampled at 5 timepoints from all colonies, n=799 workers, 250-299, see below), and measured the colony wet mass (mass was collected every other day, see below). The number of new egg batches was counted daily. On day 26, we removed all the bees and the brood excluding the pupae. The numbers of eggs and larvae were counted, and the larva body mass was measured (3432 larvae, 904-1459 per treatment). The pupae were also counted but were kept in the colonies until they all emerged. All individuals that emerged from the pupae were counted for the total number of adults (workers, gynes, males) produced in each colony. Larva body size distribution in each colony is presented using all larvae with a body mass larger than 0.1 g. This cutoff was chosen since differences in body mass between castes (queen/workers) are measurable only in the third instar, corresponding to approximately 0.1 g (38), and while the caste determination mechanism in *B. impatiens* is unknown, it is unlikely to be earlier than in *B. terrestris* (39).

Control for colony growth: To control for differences in the growth of colonies throughout the experiment (i.e., colonies with no brood could not produce new workers right away), newly emerged workers (< 24 h old) were collected daily from all worker-producing colonies and were equally redistributed across all colonies, creating a comparable growth in all colonies regardless of the amount of brood they had. A minimum of 8 and a maximum of 67 workers were collected daily in each replicate of the experiment and were equally distributed across the colonies. If the number of workers emerged did not divide by 8, extra workers were placed in colonies in the following order: nb, c and then db. While this method was effective in creating a comparable

growth across all colonies (Figure 1), it wasn't perfect since locating all newly emerged workers

in colonies containing hundreds of workers before they became indistinguishable was nearly

impossible. Therefore, we also examined the effect of the colony size (as measured by the

number of workers on the last day) on the variables tested (see results).

Colony mass: Colonies were weighted twice at the beginning of the experiment before and after

the social manipulation, and then every other day throughout the experiment using an electronic

scale. This measurement was done by placing the entire colony on a scale and is not intrusive nor

stressful. The mass reported includes the bees, brood, cells, and the plastic box in which the

colony resides but not its sugar reservoir. All colonies were placed in identical plastic boxes.

Aggressive behaviour: colonies were video recorded for aggression towards and by the queen

every other day. Video recording (20 minutes per colony) was performed between 9 a.m. to 1

p.m. Approximately 70 hours of videos were analysed by an observer blind to the experimental

design and hypotheses. Three different behaviours were counted and summed: (1) attack: this

behaviour includes overt aggression in the form of pulling, climbing, dragging or an attempt to

sting; (2) darting: a sudden movement of a bee towards another bee without a body contact, and

(3) humming: a series of short wing vibrations towards another bee that are conducted while the

bee is in movement (40-42). The behaviours performed by the queen towards workers ("queen

aggression") and by workers towards the queen ("worker aggression") were summed separately

and are presented as the average of the sum behaviours in 20 minutes.

Worker ovarian activation: 10-15% of the workers in each colony were collected at five

timepoints (on days 1, 7, 13, 19, and 26), and a subset of them (5-10 workers / colony / time

point) were dissected for ovary activation. The number of workers removed was negligible in

relation to the number of workers in the colonies and did not change the colony size in a

meaningful way. Similar samples were taken in previous studies and are representative of the

overall reproductive state of the colony (43-45). Since workers were not individually tagged,

some of the workers we dissected could be from another colony (since callows were daily

distributed), but since we added them to the colonies upon emergence, we assume their

reproductive behaviour is similar to the rest of the workers in the host colony (42). To measure

the terminal oocyte size, worker abdomen was dissected under a stereomicroscope. The two

ovaries were transferred to a drop of water and the three largest oocytes were measured using a

stereomicroscope ruler. The length of the three largest oocytes was averaged (46) and is

presented in mm.

Statistical analyses. Statistical analyses were performed using R Studio-v1.2.5033. To examine

the effect of treatment on colony wet mass, we used the plm-v2.6.1 R package to fit a time series

model and estimate the effect of treatment in colony mass while accounting for the correlation

over time. For new egg batches, worker oocyte size, aggression, the total number of eggs, larvae,

pupae, workers, males, and gynes, we used either a linear model (lm), or a generalized linear

model (glm) with a Poisson distribution, depending on whether the residuals of the model were

normally distributed or not (Shapiro-Wilk normality test, $p \geq 0.05$). A Poisson distribution was

replaced by a negative binomial distribution in case of overdispersion which was tested using

dispersiontest from MASS-7.3.55 package. lmm and glmm were used to compare the average

oocyte size and total aggression, respectively, with maternal colony as random effect. lm was

used to compare the total number of eggs, larvae, pupae, and workers across the treatments. glm

was used to compare the number of new egg batches by date, glm.nb was used for aggression by

date and the total number of males, and lmm was used for the total number of gynes. The linear

models were fitted using lmer or lm functions from lme4-v1.1.26 R package (47) and the

generalized linear models using glm or glm.nb functions from stats-v4.1.1 and MASS-7.3.55

packages included in R-v.3.6.3. All models were fitted using the treatment as a fixed effect.

Since the experiment was conducted in two consecutive replications, we also included the term

"repeat" as a random effect when it improved the model fit. The best model (with or without the

"repeat" variable) was determined using ANOVA. The effect of brood on the number of sexuals

was tested with and without the colony size (number of workers on the last day) as a fixed effect

in the model. Post-hoc pairwise comparisons across the treatments were performed using

estimated marginal means using emmeans-v1.5.4 R package with Tukey test as adjustment

method for multiple comparisons. To test the effect of treatment on the bimodality of larval body

mass distribution, we use the ACR method implemented in the multimode-v1.5 R package (48,

49). Figures were created using ggplot2-v3.3.5 and ggpubr-v0.4.0. Statistical significance was

accepted at $\alpha=0.05$. Data are presented as means $\pm$ S.E.M. The R markdown file and the data

used for the statistical analysis can be found at

https://github.com/pkfsantos/Brood_B_impatiens_colonies/tree/main.

Results

*Colony wet mass***.** The wet mass of the colonies at the beginning of the experiment, prior to the

manipulation ("day zero", Figure 1), was similar across all three treatments and was modified

("day 1") according to the desired manipulation with the db colonies weighing more compared to

the nb colonies and the control colonies being an intermediate group. Colony wet mass increased

throughout the experiment in all three treatments, reflecting the increase in worker populations

and brood and indicating a normal and healthy development of all colonies. The difference

among treatments remains throughout the experiment as evidenced by an average increase in

colony mass of 141.22 g in db colonies compared to control (plm; p=5.863e-15), and an average

decrease of 152.62 g in nb colonies compared to control (plm;p=2.2e-16). It should be noted that

the nb colonies, although had no brood at the onset of the experiment, started to produce new

brood right after the manipulation and continue to do so until the experiment was terminated on

day 26.

Production of new egg batches. The number of newly laid egg batches increased significantly in

the nb colonies, approximately one week after the onset of the experiment as compared to at least

one of the other two treatments (Figure 2). These differences were maintained for about two

weeks and were significantly higher in nb compared to the other treatments on days 6 (glm

followed by Tukey's post hoc test p=0.05 for nb vs. db), 7 (p=0.03 for nb vs. c; p=0.02 for nb vs.

db), 9 (p=0.04 for nb vs. db), and 12 (p=0.02 for nb vs. db). db colonies produced less egg

batches than the other treatments throughout the experiment, but a significant difference

compared to the control was observed only on day 16 (p=0.03 for db vs. c; Figure 2).

Aggression by and towards the queen. There were no differences in the average aggressive

behaviour performed by the queen or towards the queen between the treatments (glmm followed

by Tukey's post-hoc test, p>0.22; Figure S1). There were also no differences in the aggression

performed by/towards the queen between the treatments throughout the experiments (glm.nb

followed by Tukey's post-hoc test, p>0.065; Figure S2). One db colony was a clear outlier in the

number of the behaviours performed by the queen on day 4 (33 aggressive acts compared to 0-9

in other colonies) and towards the queen (18 compared to 0-12 in other colonies). Thus, we

analysed the data also without this colony on day 4 (Figures S1 and S2) but this did not change

the results.

Workers' ovary activation. On the first day of the experiment, all workers had inactivated ovaries

and there were no significant differences in the average terminal oocyte of workers, as we would

expect from young, queenright colonies (lmm followed by Tukey's post hoc test $p>0.64$; Figure

3). The control colonies exhibit normal development throughout the experiment, as evidence by

an increase in the average terminal oocyte of workers about two weeks after the experiment

onset. The manipulation was conducted on colonies that are estimated to be approximately 3-4

weeks old (counted from the emergence of the first worker) so an additional two weeks brought

these colonies to the competition phase where workers activate their ovaries (32, 45). The oocyte

size of nb workers was approximately double the size of workers in c (lmm followed by Tukey's

post-hoc test, $p=0.005$) and db ($p=0.038$) on day 7 to the manipulation. The oocyte size was

approximately doubled in nb compared to db also on day 13 ($p=0.004$) and day 25 ($p=0.04$). On

day 19, the differences were apparent but smaller and not significant ($p>0.05$).

Production of brood and adults. The total number of brood (eggs, larvae, and pupae) did not

vary across treatments on the last day of the experiment (day 26) (lm: Eggs or Larvae or Pupae ~

Treatment; Tukey's post hoc test $p>0.58$; Figure 4), meaning that the nb colonies have

compensated for the initial loss of brood throughout the experiment. However, the total number

of workers (including those that emerged from pupae after the last day of the experiment) was

significantly different between the treatments. All colonies had at least 200 workers, but nb

colonies (375±50 workers, means ± SE) were smaller than db (678±50) and control (643±46)

colonies (lm followed by Tukey's post hoc test $p=0.004$ (nb vs. c); $p=0.002$ (nb vs. db); Figure

5A). db colonies produced significantly more gynes than c and nb (lmer followed by Tukey's

post-hoc, $p=0.02$), with 132±51 gynes, on average, in db compared to 0.6±0.6 and 3.8±2 in nb

and the control, respectively. When we controlled for the number of workers by the end of the

experiment, db and c remain significantly different (lmer followed by Tukey's post-hoc,

$p=0.035$; Figure 5B), but db and nb were no longer significantly different. Many more males

were produced in the nb colonies (64±27) compared to db (40±29) and c (15±13), but the variation across colonies was high and the average number of males did not significantly vary between the treatments before or after controlling for the number of workers (glm.nb followed by Tukey's post-hoc, $p>0.15$; Figure 5C).

The number of gynes could not be explained by brood differentiating to gynes prior to the experimental manipulation since all the gyne-producing colonies, excluding two, produced gynes 25-43 days following the manipulation (Table S1). Thus, at the time of the brood manipulation, the diploid brood was not produced yet or was too young to commit to being a worker or a gyne. The critical period for female differentiation in *B. terrestris* is during the first or second instar, ~11 days after eggs are laid (50) and is unknown in *B. impatiens*. However, even according to a conservative estimate of an early critical period as in *B. terrestris*, the first gyne emerging in most colonies was unlikely to be committed to be a gyne at the onset of the manipulation. In two colonies, the first gyne emerged 19 days after the onset of the manipulation and could potentially be committed to be gynes at the manipulation onset, but both these colonies produced >200 gynes in total over an extended period, so the majority of gynes produced in these colonies differentiate after the manipulation onset. Moreover, although the number of colonies producing gynes in db and control was close (4/5 db and 3/6 c), the total number of gynes produced in the db colonies was 23 times higher compared to c colonies.

The workers:larvae ratio by the end of the experiment was biased in favour of workers in db colonies (mean 3.4:1), nearly balanced in c colonies (mean 1.3:1) and biased in favour of larvae in nb colonies (mean 0.6:1) (Supplementary Table S2). Although these ratios were not assessed at the beginning of the experiment, they can be inferred from the data, with the highest W:L ratio in nb colonies (~70 workers, no brood) and a much lower ratio in the db colonies compared to the control, as they maintained a similar number of workers but had brood from two colonies. Overall, the ratios by the beginning of the experiment were flipped towards the end of the experiment.

Distribution of larva body mass. The differences between the treatments in gyne production are reflected also in the body mass distribution of larvae that were collected and weighted on the last

day of the experiment (day 26). The larvae in db colonies showed a bimodal distribution of larva

mass, corresponding to larvae that will develop into workers/males and gynes (ACR mode test,

p<2.2e-16), while c and nb colonies showed a unimodal distribution, indicating the production of

workers/males only (ACR mode test, p>0.114; Figure 6). Larval weight distribution per colony

can be found in the Supplementary Figure S3.

Discussion

In this study, we demonstrate that manipulating brood quantity in a social colony significantly

impacts colony development, gyne production, and worker reproduction, while showing no

significant effect on aggressive behaviour or male production. Additionally, differences in gyne

numbers between treatments could not be attributed to colony size. The elevated production of

gynes in colonies with doubled brood suggests that brood may function as an extended

phenotype of the queen, influencing not only individual reproductive decisions of workers but

also colony-level strategies for investing in future queens.

In social insect species, gynes are typically larger and take longer to develop whereas males are

smaller and cheaper to produce (45, 51). Investment in either depends on the availability of

resources, and in social hymenopteran species, also reflects the outcomes of the conflict between

the queen and the workers (52). Maintaining the social structure depends on workers gaining

inclusive fitness by helping the queen rear her offspring. If the queen is no longer fecund, it is in

the interest of workers to start producing males, and since the brood is the ultimate testimony to

the queen's fecundity, its absence or a reduction in its amount are sufficient for the workers to

break down the social structure. The effect of brood on gyne production could be a result of a

direct impact of the brood or could be affected by the queen and/or the workers in response to the

amount of the brood. Such indirect effects could be due to a change in the feeding regime,

pheromones produced by females or predetermination of caste during egg laying.

Our data also suggest that reduction in brood may lead colonies to invest in males. While these

data were insignificant, the number of males produced in nb colonies was approximately x2

compared to db and x5 compared to the control. Males can be produced by the queen (when she

stitches to lay eggs earlier in the life cycle), or by the workers when they start competing over

male production but it is not clear who produced them in each of the treatments.

If the males in nb were produced by the queen, this may reflect a trade off in the queen's strategy

to invest in sexuals that are cheaper to produce; A previous study (35) showed that queens

following a longer winter diapause produced colonies with more males, whereas shorter diapause

resulted in colonies producing more gynes, potentially resulting from the physiological state of

the queen and the amount of brood she produced. Queens that switch early to male production

typically produce smaller colonies, fewer gynes and more males in line with the profile of the nb

colonies (32). Alternatively, the males emerging in the nb colonies could have been produced by

workers that competed with the queen over male production following the absence of brood and

the perception of the queen as not fecund or of lower quality. While we don't know for sure who

laid the male eggs in the nb colonies, it is likely that at least some of the eggs were laid by the

queen since workers emerged in nb more than 30 days after the onset of the experiment, meaning

that at least some eggs in these colonies were diploid and were laid by the queen. On the other

hand, the number of egg batches produced per day in these colonies is likely too many for one

individual to lay them all, thus at least some of the eggs were laid by workers in nb colonies. If

males were produced by workers, it is an indication that the queen was perceived as weak. If

both the queen and workers produced males simultaneously, this would represent an unusual

case of peaceful competition since aggression in the nb colonies was no higher compared to the

other treatments.

A few more points are worth mentioning. First, the manipulation we conducted to the amount of

brood, although extreme, did not affect colony survival or health. All manipulated colonies

recovered quickly, as evident by the increase in mass throughout the experiment (Figure 1) and

by no significant differences in the amount of brood by the end of the experiment (Figure 5).

This quick recovery was partially achieved by a temporary increase in egg laying in nb colonies

(Figure 2), indicative of the plasticity of the queen's behaviour. The second point is the potential

impact of relatedness on the results. All colonies contained non-related workers (as newly

emerged workers were distributed daily across all the colonies, including the controls), but only

the db colonies contained partially unrelated brood, raising the question whether gynes were

more likely to develop in colonies with mixed brood. In a previous study we didn't find any

evidence of the impact of relatedness on brood care (27) and therefore consider it unlikely to

have an impact here. We also didn't observe any differential behaviour towards the brood in our

study in any of the colonies. Whether unrelated brood is more likely to develop into gynes

compared to related brood is a question worth investigating. Finally, the distinct effects of colony

size and brood quantity on worker reproduction and gyne production remain to be disentangled.

Although these two processes are correlated (8) and often assumed to share common triggers,

emerging evidence suggests they are likely regulated independently. While colony size has been

shown to influence worker reproduction before (53), gyne production—despite occurring in

close proximity to the onset of worker reproduction—remains unaffected by colony size. The

worker-to-larvae ratio does not appear to be critical for determining the onset of gyne production

although it has never been tested directly and can only be indirectly inferred from other studies

(12, 13), including ours. Altogether, this may suggest that the shift toward gyne production may

depend more on the queen's perception of her age (12) or the absolute amount of brood present

(regardless of whether the brood is her own), rather than on the number of workers. However,

further research is needed to pinpoint the exact mechanism driving this transition.

Conclusions

Overall, the results of our study show that the brood regulates not only egg laying in small

groups of workers, as we found before (27), but also influences colony-level events such as the

beginning of the reproductive competition and the timing of gyne production. These data suggest

that the brood provides a testimony to the fecundity of the queen and leads to gyne production

when it is abundant, and possibly also to the production of cheap males in its absent. Our

findings in bumble bees align with data from wasps, also annual, in which partial brood removal

led to increased egg laying by the queens but did not prevent worker reproduction (54). This may

suggest a general mechanism in annual social species where the brood determines in an

irreversible way the level of cooperation between the workers and the queens and further

influences the production of sexuals. Altogether, our findings emphasize the importance of the

brood in maintaining and shaping the social organization in social insects and the need to

investigate its diverse roles across other taxa.

Declarations

Funding. This work was funded by the National Science Foundation IOS-1942127 to EA.

Conflict of interests. The authors declare no conflicts of interests

Availability of data and material. The data used and/or analyzed during the current study are available as a supplementary material.

Acknowledgements. We would like to thank John M. Craig, Andrea Medero and Katie Barie for assisting in the experiments.

Author's contributions. PKFS and EA designed the experiments, PKFS collected and analyzed the data, CSM performed the behavioral observations, PKFS and EA wrote the manuscript. All authors gave final approval for publication and agree to be held accountable for the work performed therein.

References

References

1.   Amsalem E, Grozinger CM, Padilla M, Hefetz A. The physiological and genomic bases of
bumble bee social behaviour. In: Amro Z, Clement FK, editors. Genomics, Physiology and
Behaviour of Social Insects. 48. Adv In Insect Phys: Academic Press; 2015. p. 37-93.
2.   Jandt JM, Tibbetts EA, Toth AL. Polistes paper wasps: a model genus for the study of
social dominance hierarchies. Insectes Sociaux. 2013;61(1):11-27.
3.   Macevicz S, Oster G. Modeling social insect populations II: Optimal reproductive
strategies in annual eusocial insect colonies. Behavioral Ecology and Sociobiology.
1976;1(3):265-82.
4.   Hovestadt T, Degen T, Mitesser O. Suitable triggers for timing the transition from worker
to sexual production in annual eusocial insects. Insectes Sociaux. 2018;65(4):609-17.
5.   Treanore ED, Derstine N, Amsalem E. What can mechanisms underlying derived traits
tell us about the evolution of social behavior? Annals of the Entomological Society of America.
2020;114(5):547-61.
6.   Trible W, Kronauer DJ. Caste development and evolution in ants: it's all about size. J Exp
Biol. 2017;220(Pt 1):53-62.
7.   Wilson EO. The Evolution of Caste Systems in Social Insects. Proceedings of the
American Philosophical Society. 1979;123(4):204-10.
8.   Cnaani J, Robinson GE, Bloch G, Borst D, Hefetz A. The effect of queen-worker conflict
on caste determination in the bumblebee *Bombus terrestris*. Behavioral Ecology and
Sociobiology. 2000;47(5):346-52.
9.   Liautard C, Brown WD, Helms KR, Keller L. Temporal and spatial variations of gyne
production in the ant Formica exsecta. Oecologia. 2003;136(4):558-64.
10.   Lopez-Vaamonde C, Brown RM, Lucas ER, Pereboom JJM, Jordan WC, Bourke AFG. Effect
of the queen on worker reproduction and new queen production in the bumble beeBombus
terrestris. Apidologie. 2007;38(2):171-80.
11.   Alaux C, Jaisson P, Hefetz A. Regulation of worker reproduction in bumblebees (*Bombus
terrestris*): workers eavesdrop on a queen signal. Behavioral Ecology and Sociobiology.
2006;60(3):439-46.
12.   Alaux C, Jaisson P, Hefetz A. Reproductive decision-making in semelparous colonies of
the bumblebee bombus terrestris. Behavioral Ecology and Sociobiology. 2005;59(2):270-7.
13.   Bloch G. Regulation of queen-worker conflict in bumble-bee (*Bombus terrestris*)
colonies. Proc Biol Sci. 1999;266(1437):2465-9.
14.   Shykoff JA, Muller CB. Reproductive Decisions in Bumble-Bee Colonies: The Influence of
Worker Mortality in Bombus terrestris (Hymenoptera, Apidae). Functional Ecology.
1995;9(1):106-12.
15.   Muller C, Schmid-Hempel P. Variation in life-history pattern in relation to worker
mortality in the bumble-bee, Bombus lucorum. Functional Ecology. 1992:48-56.
16.   Deleurance EP. Sur Le Mecanisme De La Monogynie Fonctionnelle Chez Les Polistes
(Hymenopteres-Vespides). Comptes Rendus Hebdomadaires Des Seances De L Academie Des
Sciences. 1950;230(8):782-4.
17.   Seeley TD. Honeybee democracy. Princeton, New Jersey: Princeton University Press;
2010.

18.   Vargo EL, Passera L. Pheromonal and behavioral queen control over the production of
gynes in the Argentine ant Iridomyrmex humilis (Mayr). Behavioral Ecology and Sociobiology.
1993;28:161-9.
19.   Boulay R, Hefetz A, Cerdá X, Devers S, Francke W, Twele R, et al. Production of sexuals in
a fission-performing ant: dual effects of queen pheromones and colony size. Behavioral Ecology
and Sociobiology. 2007;61(10):1531-41.
20.   Schultner E, Oettler J, Helantera H. The Role of Brood in Eusocial Hymenoptera. Q Rev
Biol. 2017;92(1):39-78.
21.   Mohammedi A, Paris A, Crauser D, Le Conte Y. Effect of Aliphatic Esters on Ovary
Development of Queenless Bees (Apis mellifera L.). Naturwissenschaften. 1998;85(9):455-8.
22.   Jay SC. Ovary development of worker honeybees when separated from worker brood by
various methods. Canadian Journal of Zoology. 1972;50(5):661-4.
23.   Maisonnasse A, Lenoir JC, Beslay D, Crauser D, Le Conte Y. E-beta-ocimene, a volatile
brood pheromone involved in social regulation in the honey bee colony (*Apis mellifera*). PLoS
One. 2010;5(10):e13531.
24.   Endler A, Liebig J, Schmitt T, Parker JE, Jones GR, Schreier P, et al. Surface hydrocarbons
of queen eggs regulate worker reproduction in a social insect. Proc Natl Acad Sci U S A.
2004;101(9):2945-50.
25.   Ebie JD, Holldobler B, Liebig J. Larval regulation of worker reproduction in the
polydomous ant *Novomessor cockerelli*. Naturwissenschaften. 2015;102(11-12):72.
26.   Ravary F, Jahyny B, Jaisson P. Brood stimulation controls the phasic reproductive cycle of
the parthenogenetic ant Cerapachys biroi. Insectes Sociaux. 2006;53(1):20-6.
27.   Starkey J, Brown A, Amsalem E. The road to sociality: brood regulation of worker
reproduction in the simple eusocial bee *Bombus impatiens*. Animal Behaviour. 2019;154:57-65.
28.   Pankiw T. Brood pheromone regulates foraging activity of honey bees (Hymenoptera:
Apidae). J Econ Entomol. 2004;97(3):748-51.
29.   Pankiw T, Page Jr RE, Kim Fondrk M. Brood pheromone stimulates pollen foraging in
honey bees (Apis mellifera). Behavioral Ecology and Sociobiology. 1998;44(3):193-8.
30.   Ulrich Y, Burns D, Libbrecht R, Kronauer DJ. Ant larvae regulate worker foraging behavior
and ovarian activity in a dose-dependent manner. Behav Ecol Sociobiol. 2016;70(7):1011-8.
31.   Free JB. Queen Production in Colonies of Bumblebees. Proceedings of the Royal
Entomological Society of London Series A, General Entomology. 2009;30(1-3):19-25.
32.   Duchateau MJ, Velthuis HHW. Development and reproductive strategies in *Bombus
terrestris* colonies. Behavior. 1988;107:186-207.
33.   Amsalem E, Shamia D, Hefetz A. Aggression or ovarian development as determinants of
reproductive dominance in *Bombus terrestris*: interpretation using a simulation model. Insectes
Sociaux. 2013;60(2):213-22.
34.   Holland JG, Guidat FS, Bourke AF. Queen control of a key life-history event in a eusocial
insect. Biol Lett. 2013;9(3):20130056.
35.   Duchateau MJ. Sex ratio variation in the bumblebee *Bombus terrestris*. Behavioral
Ecology. 2004;15(1):71-82.

36.   Orlova M, Starkey J, Amsalem E. A small family business: synergistic and additive effects
of the queen and the brood on worker reproduction in a primitively eusocial bee. J Exp Biol.
2020;223(Pt 3):jeb217547.
37.   Orlova M, Amsalem E. Bumble bee queen pheromones are context-dependent. Sci Rep.
2021;11(1):16931.
38.   Hefetz A, Robinson GE, Huang Z, Borst DW, Cnaani J. Caste Determination in Bombus
terrestris: Differences in Development and Rates of JH Biosynthesis between Queen and
Worker Larvae. J Insect Physiol. 1997;43(4):373-81.
39.   Barie K, Amsalem E. The influence of the social environment on larval development and
resulting caste in Bombus impatiens. bioRxiv. 2022:2022.09.09.507355.
40.   Duchateau MJ. Agonistic behaviours in colonies of the bumblebeeBombus terrestris.
Journal of Ethology. 1989;7(2):141-51.
41.   Amsalem E, Grozinger CM. Evaluating the molecular, physiological and behavioral
impacts of CO_2 narcosis in bumble bees (Bombus impatiens). J Insect Physiol. 2017;101:57-65.
42.   Amsalem E, Hefetz A. The appeasement effect of sterility signaling in dominance
contests among Bombus terrestris workers. Behavioral Ecology and Sociobiology.
2010;64(10):1685-94.
43.   Padilla M, Amsalem E, Altman N, Hefetz A, Grozinger CM. Chemical communication is
not sufficient to explain reproductive inhibition in the bumblebee Bombus impatiens. Royal
Society Open Science. 2016;3(10):160576.
44.   Treanore ED, Kiner JM, Kerner ME, Amsalem E. Shift in worker physiology and gene
expression pattern from reproductive to diapause-like with colony age in the bumble bee
Bombus impatiens. J Exp Biol. 2020;223(Pt 9):jeb218768.
45.   Cnaani J, Schmid-Hempel R, Schmidt JO. Colony development, larval development and
worker reproduction in Bombus impatiens Cresson. Insectes Sociaux. 2002;49(2):164-70.
46.   Amsalem E, Twele R, Francke W, Hefetz A. Reproductive competition in the bumble-bee
Bombus terrestris: do workers advertise sterility? Proc Biol Sci. 2009;276(1660):1295-304.
47.   Bates D, Mächler M, Bolker B, Walker S. Fitting Linear Mixed-Effects Models Usinglme4.
Journal of Statistical Software. 2015;67(1):1-48.
48.   Ameijeiras-Alonso J, Crujeiras RM, Rodríguez-Casal A. Mode testing, critical bandwidth
and excess mass. Test. 2018;28(3):900-19.
49.   Ameijeiras-Alonso J, Crujeiras RM, Rodriguez-Casal A. multimode: An R Package for
Mode Assessment. Journal of Statistical Software. 2021;97(9):1-32.
50.   Cnaani J, Robinson GE, Hefetz A. The critical period for caste determination in Bombus
terrestris and its juvenile hormone correlates. J Comp Physiol A. 2000;186(11):1089-94.
51.   Fisher RA, Bennett JH. The genetical theory of natural selection: a complete variorum
edition. (No Title). 1930.
52.   Bourke AF. Sex investment ratios in eusocial Hymenoptera support inclusive fitness
theory. J Evol Biol. 2015;28(11):2106-11.
53.   Amsalem E, Hefetz A. The effect of group size on the interplay between dominance and
reproduction in Bombus terrestris. PLoS One. 2011;6(3):e18238.
54.   Liebig J, Monnin T, Turillazzi S. Direct assessment of queen quality and lack of worker
suppression in a paper wasp. Proc Biol Sci. 2005;272(1570):1339-44.

Legend to figures

Figure 1: The average colony wet mass throughout the experiment in the three treatment groups:

colonies containing no brood (nb), colonies containing a double amount of brood (db), and

control colonies that remain unmanipulated (c). Timepoints 0 and 1 refers to the colony wet mass

before and after brood manipulation, respectively. Data are presented as means ± S.E.M.

Figure 2: The average number of newly laid egg batches per day throughout the experiment in

the three treatment groups: colonies containing no brood (nb), colonies containing a double

amount of brood (db) and control colonies that remain unmanipulated (c). Data are presented as

means ± S.E.M. Asterisks denote statistical difference at $p < 0.05$.

Figure 3: The average terminal oocyte size in workers in the three treatment groups: colonies

containing no brood (nb), colonies containing a double amount of brood (db), and control

colonies that remain unmanipulated (c). Oocyte size was measured in five timepoints throughout

the duration of the experiment using 5-10 workers from each colony. The Number above

columns represent the number of workers collected from all colonies. Different letters denote

significant differences at $p < 0.05$.

Figure 4: The effect of brood on the total number of eggs, larvae, and pupae on the last day of

experiment in the three treatment groups: colonies containing no brood (nb), colonies containing

a double amount of brood (db), and control colonies that remain unmanipulated (c). Different

letters denote significant differences at $p < 0.05$.

Figure 5: The effect of brood on the total number of workers, gynes and males in the three

treatment groups: colonies containing no brood (nb), colonies containing a double amount of

brood (db), and control colonies that remain unmanipulated (c). Different letters denote

significant differences at $p < 0.05$.

Figure 6: The effect of brood on the distribution of larval weight by the end of the experiment

grouped by treatment: colonies containing no brood (nb), colonies containing a double amount of

brood (db), and control colonies that remain unmanipulated (c).