Title: Global quantitative understanding of nonequilibrium cell fate decision making in response to pheromone

Authors: Sheng Li [1,2], Qiong Liu[2]*, Erkang Wang [1,2], Jin Wang[3]*

Affiliations:

[1]College of Chemistry, Jilin University, Changchun, Jilin, 130012 P. R. China.

[2]State Key Laboratory of Electroanalytical Chemistry, Changchun Institute of Applied Chemistry, Chinese Academy of Sciences, Changchun, Jilin, 130022, China.

[3]Department of Chemistry and of Physics and astronomy, State University of New York at Stony Brook, Stony Brook, New York, 11794-3400, USA.

*Corresponding author. Email: liuqiong@ciac.ac.cn; jin.wang.1@stonybrook.edu.

Abstract:

Cell cycle arrest and polarized cell growth are commonly used to qualitatively characterize the fate of yeast in response to pheromone. However, the quantitative decision-making process underlying the time-dependent changes in cell fate remains unclear. Here, by observing the multi-dimensional responses at the single-cell level experimentally, we find that yeast cells have various fates. Multiple states are revealed, along with the kinetic switching rates and pathways among them, giving rise to a quantitative landscape of mating response. We developed a theoretical framework using a nonequilibrium landscape and flux theory to account for the cell morphology observed experimentally and performed a stochastic simulation of biochemical reactions to explain the signal transduction and cell growth. Our experimental results established the first global quantitative demonstration of the real-time synchronization of intracellular signaling with their physiological growth and morphological functions which reveals the underlying physical mechanism. This study provides an emerging mechanistic approach for understanding the nonequilibrium global pheromone-regulated cell fate decision-making in growth and morphology.

Keywords:

Mating pathway; Cellular fate decision-making; Landscape; Flux; Nonequilibrium.

     Main Text:

INTRODUCTION

Facing a variety of external stimuli, cells integrate information from various sources and
initiate appropriate stress responses, such as survival or death, division or differentiation, and
shape change or migration (*1-7*). To better understand the cellular response to external stimuli,
we chose the pheromone pathway of *Saccharomyces cerevisiae*, which is a classical model of the
mitogen-activated protein kinase (MAPK) signaling pathway (*8-11*). *Saccharomyces cerevisiae*
has two modes of reproduction, sexual mating reproduction and asexual budding reproduction
(*12-16*). These two reproductive modes can be converted either with or without the assistance of
external pheromone stimulation from a partner (*17*). For sexual reproduction, mating to form a
diploid cell is the natural behavior of heterosexual haploid *Saccharomyces cerevisiae* to cope
with an unfavorable environment and to improve their survival rate for generations (*18-20*).
Pheromones, a type of sex hormone secreted by *Saccharomyces cerevisiae*, serve as the yeast's
mating signal, informing its partner to prepare for the beginning of cell fusion. (*18-26*). What
will happen to the yeast cell if its partner does not arrive and this normal stimulation continues?
From microscopic gene network **(Fig. 1)**, external signals are transmitted to Fus3 through a
series of internal proteins such as prototype heterotrimeric GTP binding protein and the MAPK
kinase cascade (*27-29*). Then, Fus3 shuttles back and forth across the nuclear membrane, directly
or indirectly activating genes for cell cycle arrest (Far1) and polar growth (Bni1) (*30-38*). Thus,
the current understanding of the pheromone-induced fate of yeast cells can be qualitatively
summarized as the occurrence of cell cycle arrest and polarized cell growth (*39-41*).
During qualitative studies of cell fate, researchers have recognized that cellular decision-
making plays a pivotal role in determining the ultimate fate of cells in the presence of various
external stimuli (*42-54*). From a biological perspective, cell fate decisions can be conceptually
characterized as the cascading transmission of signals along a static causal pathway (*55-59*).
However, the quantitative dynamics of cell fate decision-making over time during global
responses at both the mesoscopic and microscopic levels remain poorly understood. Recent
advances in live cell fluorescence imaging platforms have led to increasing focus on studying the
dynamics of biological processes and the underlying molecular mechanisms in living cells (*54,*
60-73). Nevertheless, disruptions in the external environment can perturb the delicate balance of
cellular activities and exacerbate internal imbalances in yeast cells, as living cells are complex
non-equilibrium microsystems (*74-83*). Therefore, quantifying these underlying cell fate
decisions in response to external stimuli remains a significant challenge.
Indeed, from a microscopic perspective, the cellular decision-making of yeast in response to
mating information is accomplished through multiple feedback loops, not only including the
positive feedback regulation mediated by Fus3, but also through certain negative feedback
regulation that can effectively reduce the transcriptional output of this pathway (*84-89*). After
pheromone stimulation, there are two pathways that can lead to polar growth of the cells **(Fig. 1)**.
The first path (P_1) is "Fus3 → Far1/Cdc24 → Cdc42 → Bni1" (*40, 90-93*). This is realized by
Fus3 entering the nucleus to activate Far1, which can escape from the nucleus to activate Bni1
indirectly. The second path (P_2) is direct activation by Fus3 of Bni1 in the cytoplasm, "Fus3 →
Bni1" (*31, 94*). In the process of these two polar growth pathways acting alone or in concert, the
pheromone-induced self-activation of Fus3 ("Fus3 → Fus3") facilitates the rapid transmission of
the signals (*95-98*). Activation of the pheromone pathway also induces multiple negative
feedback loops, such as I_1 (Sst2 and Gpa1) (*87-89, 99-101*), I_2 (Msg5 and Bni1) (*31, 85-89, 102-*

104) in the pathway. Only at high-dose pheromone levels is the negative feedback of the
downstream Msg5 upregulated (*85*) and only at high concentrations of Bni1 is activity of Fus3
indirectly reversed by Bni1 (*31, 94, 105-107*).
Here, by observing the multi-dimensional response at the single-cell level, we discovered
the non-equilibrium steady states, which demonstrate the cell fates quantitatively, in response to
different concentrations of pheromone. These steady states or cellular destinies include two gene
expression levels, four growth rates, and four morphological fates. We quantified these
responsive fates of the yeast cells in real time from various dimensions, including expression
levels of Fus3 inside and outside of the nucleus, cell morphology, cell growth rate, and the
stimulation concentrations. Multiple states, as well as switching kinetic rates and pathways
among them, were revealed, giving rise to a quantitative landscape of the mating response. The
applications of landscape and flux theory to this biological system enabled us to quantify the
non-equilibrium dynamics of the yeast cell mating responses. Our results established a global
and physical framework for understanding cell fate decision making and mating dynamics.
Furthermore, we experimentally validated the proposed molecular mechanism for the formation
of these states. These molecular mechanisms establish the links between molecular events to
cellular characteristics across scales. They allow for the real-time synchronization of intracellular
signaling with their physiological growth and morphological functions, bridging microscopic
mechanisms and mesoscopic functions. To further elucidate these microscopic mechanisms, we
conducted biochemical reaction simulations to demonstrate the emergence of these states. These
findings shed new light on the global signaling mechanisms that govern how yeast determines its
cellular fate in response to pheromone.

RESULTS

Quantifying the cellular decision-making threshold

Studies have shown that yeast, for which mating decisions are an all-or-none switch-like
response, can automatically filter out very low-dose pheromone signals and only respond near a
critical concentration or higher (*101, 108, 109*). This is due to the fact that very low doses of
pheromone induce insufficient accumulation of intracellular signals, such as Fus3, which are
incapable of initiating the mating-readiness project (*110-112*). To determine a pheromone
concentration capable of arresting the cell cycle, we measured the fluorescence intensity of
FUS3-GFP by flow cytometry in yeast cells incubated for 24 h with varying pheromone
concentrations **(Fig. 2A)**. According to the fitted statistics, the expression of Fus3 at a
pheromone dose of 0.01–0.4 μM showed unimodal distribution of the fluorescence intensities.
The bimodal distribution representing the bistable states gradually began to appear when the
dose was about 0.6 μM.
In order to comprehend the correlation between gene expression levels of Fus3 and the
threshold of cellular response, the fluorescence intensity trajectories of Fus3 expression in single
cells were associated with the microscopy-based cellular response under various pheromone
concentrations over time. A microfluidic device was used to culture yeast under constant
temperature conditions in real time. Using total internal reflection fluorescence microscopy, the
survival status and fluorescent signals of the yeast cells were captured in real time **(Movies S1–
S7)**. Compared to 0.7 μM pheromone, the expression of Fus3 in the yeast cells did not fluctuate
significantly at 0.2 μM, resulting in its inability to inhibit cell budding reproduction **(Fig. 2B)**.
However, the expression of Fus3 at a pheromone concentration of 0.7 μM exhibited four stages:

the initial stage (0–b_1), the adaptive stage (b_1–b_2), the stable stage (b_2–b_3), and the recovery stage
(b_3–b_4), corresponding to the biological behavior of yeast cells at different stages. Therefore, the
unimodal distribution of 0.01–0.4 μM pheromone occurred when only the initial cells (0-b_1) were
present in the flow cytometer sample. This very low dose of pheromone was unable to arrest the
cell cycle from the beginning in yeast. If the pheromone had arrested the cell cycle of yeast cells
from the beginning, but not for more than 24 h, some yeast cells would have resumed budding at
a later stage. As demonstrated by the bimodal distribution of 0.4–1.5 μM pheromone, the sample
contained yeast cells in two distinct states (b_2–b_3 and b_3–b_4).
The results measured by flow cytometry were only rough estimates of critical mating
decision thresholds, whereas microscopy of individual cells cultured in microfluidics revealed
more precise cellular decisions. During culturing with varying concentrations of pheromone, we
found that the time required for yeast cells to arrest their cell cycle was directly proportional to
the pheromone concentration **(Fig. 2C)**. By extending the duration of the pheromone stimulation,
the yeast cells gradually adapted to the external surroundings and resumed budding reproduction.

Two cell fate decision states reflected in the expressions of Fus3

To explore the yeast cell fate decision-making during the stable stage (b_2–b_3 in **Fig.2B**), the
fluorescence intensity trajectories of Fus3 inside and outside the nucleus were plotted over time
(Fig. 3A and Fig. S1–S2). The trajectories demonstrated that after approximately 600 min of
yeast cell cultivation, Fus3 entered a stable stage where its fluorescence fluctuated around a
certain value. In non-equilibrium statistical physics, the fluorescence intensity curve in this stable
stage can be interpreted as a non-equilibrium steady state, whereas the rising phase of the first
600 min can be interpreted as a non-equilibrium non-steady state, or the relaxation process of the
response. To visualize the cell fate of yeast at the level of gene expression, the three-dimensional
statistical distribution of Fus3 (cytoplasmic and nuclear fluorescent signals) was plotted during
the non-equilibrium steady-state phases **(Fig. 3B and Fig. S3–S5)**. Clearly, the landscape of
Fus3 displayed two peaks, which correspond to the two cell fates that existed both inside and
outside the nucleus.
To comprehend the logical relationship between the cell fate at expression levels of Fus3
and pheromone-influenced signaling in the transduction process, we explored the biological
mechanism underlying the emergence of these two Fus3 fates. We proposed that the two fates of
Fus3 expression observed in signaling could be explained by a logical sequence involving
functional depletion and feedback loops. First, functional depletion means that when the amount
of Fus3 in the nucleus was sufficient to arrest the cell cycle and activate the mating protein
("$\text{Outer\_P}_1 \rightarrow \text{Far}1 \ \& \ Ste12$"), the excess Fus3 was transported out of the nucleus for polar
growth ("$\text{Outer\_P}_1 \rightarrow \text{Inner\_P}_1$" and "$\text{Inner\_P}_2 \rightarrow \text{Outer\_P}_2$") **(Fig. 3C)**. Second, the feedback
loop means that yeast actively regulated Fus3 expression by activating or inhibiting signals.
Previously, experimentally validated feedback loops were categorized into two models: "A →
Fus3 ⊣ A" focuses primarily on the pathways involving Gpa1 and Sst2, whereas "Fus3 → B ⊣
Fus3" focuses primarily on the pathways involving Ste12 and Msg5 **(Fig. 1 and Fig. 3D)**.
This dynamic sequential mechanism results in two behaviors: the highly coordinated
expression levels of Fus3 inside and outside the nucleus, and the alternation of the two fates. As
experimental evidence for our proposed molecular mechanism, we observed the existence of two
behaviors. We revealed a correlation between intra- and extra-nuclear fluorescence intensities
using Pearson's coefficient. A linear correlation of approximately 0.80 between these two types
of fluorescence trajectories at various pheromone concentrations indicated a strong synergy

between them **(Table 1)**. In addition, using a hidden Markov chain model, we characterized the two cellular fates using gene expression levels by fitting the fluorescence trajectories into high states (H_1, H_2) and low states (L_1, L_2) **(Fig. 3E)**. The fitted red line in **Fig. 3E** demonstrates that the two trajectory fates (high and low states) inside and outside the nucleus could switch to each other, supporting our logical interpretation of the state switching above.

Uncovering the physical characteristics of cellular decision-making landscapes

To uncover the physical characteristics of the underlying bistable landscapes, we employed a Hidden Markov Chain model to compute the transition probability, transition rate, and residence time between the two gene expression fates. In the non-equilibrium steady state, the steady state probability can be used to quantify the population landscape P or the potential landscape U (*81, 113-115*), where U is defined as the negative logarithm of the steady state distribution P of gene expression, $U = -\ln P$. As can be seen, the population and potential landscapes (P and U) at various pheromone concentrations have two basins of attractions that are separated by barriers **(Fig. 3B and Fig. S3–S5).** The transition probabilities and residence times of the two expressions of *FUS3*, high state (H_1, H_2) and low state (L_1, L_2), can be determined through the statistical analysis of the experimental data fitted by the hidden Markov chain model. The transition rate can be derived from the transition probability, whereas the barrier height is determined by fitting two peaks into the landscape. The barrier height is determined by the depth of the basin on the potential energy landscape ($U = -\ln P$) obtained directly from the statistical histogram (P) of the fluorescence signals of the gene expressions **(Table 2)**.

At pheromone concentrations between 0.7 μM and 0.8 μM, the barrier heights of the low state (L_1, L_2) were less than those of high state (H_1, H_2), whereas the opposite was true between 1.0 μM and 3.0 μM **(Fig. 4A)**. In physics, the transition between steady states becomes harder as the barrier height increases, resulting in a longer residence time (*113, 116*). According to the statistical results of the experimental data, the distribution of high and low states of residence times in different pheromone concentrations is consistent with the distribution of residence times **(Fig. 4B)**. The correlation coefficient between barrier heights and residence times for all concentrations was 0.8806, indicating that they were significantly positively correlated **(Fig. 4C)**. This substantiates the statistical non-equilibrium physics claims regarding barriers and residence time.

To explore the biological implications of the landscape physical characteristics, we calculated the distribution of their ratios for two gene expression fates as a function of pheromone concentration. The opposite trend of the transfer rate ratio (k_{21}/k_{12}) relative to the barrier height ratio or the residence time ratio (high/low, H/L) in response to changes in pheromone concentration indicated that transition rates were lower the deeper the attraction basin **(Fig. 4D)**. To better comprehend this relationship, the potential landscape topographies under varying pheromone concentrations were illustrated **(Fig. 4E)**. Compared to a pheromone concentration of 0.7 μM, 0.8 μM had a deeper attraction basin for its high state than for its low state, making the transition from low to high states easier than that from high to low states. The yeast's cellular decision-making at the level of gene expression was consequently more likely to remain in the high state than in the low state. At 1.0–3.0 μM, the attraction basin of the low state was deeper than that of the high state, leading to the yeast's preference to remain in the low state. In general, as pheromone doses increase from low to high, the basin of the high state gradually becomes shallower, whereas the basin of the low state gradually becomes deeper. The increased height of the low-state potential barrier is due to the negative feedback regulation that is able to

212 inhibit the expression of Fus3, resulting in a greater number of yeast cells in a low-expression
213 state. This supports the biological notion that high-dose pheromone elicits a stronger negative
214 feedback response compared to the low-dose pheromone.

Quantifying the cellular deformation rate in polar growth

Upon observing the polar growth of yeast cells in real time via microscopy, we noticed that
the rate of cell deformation did not vary uniformly over time. Meanwhile, except for the top,
other parts of the cell changed slightly as the yeast stretched forward **(Movies S3–S7)**. To
accurately measure the spatiotemporal change rate of the cell morphology, we used a circular
filling pattern to segment the yeast cells **(Fig. 5A)**. To quantify the various cell deformations that
occur during cell growth, we considered a value (H_n) comparable to the harmonic mean to
characterize the cell morphology. H_n equals the sum of the reciprocal radii of the filled circles
multiplied by the number of filled circles, i.e., $H_n = n(\frac{1}{R_1} + \frac{1}{R_2} + \ldots + \frac{1}{R_n})$. The significant
advantage of this parameter is that it is particularly sensitive to small morphological changes at
various locations of the cell, allowing differential non-directional polar growth and directional
normal growth (i.e., lateral growth and longitudinal growth) to be characterized in real time.
From the real-time trajectory of the changes in the cell morphology, as measured by the H_n
of the cell-filled circles, we found that there were roughly four types of cell shape changes.
According to the definition of H_n describing the cell morphology, the rising phase ($t_1 - t_2$) of
the curve primarily represents the change in cell length, whereas the falling phase ($t_3 - t_4$)
primarily represents the change in cell width. At the ($t_2 - t_3$) and ($t_4 - t_5$) phases, the slopes of
the two curves are close to zero, indicating that cell morphology changes were minimal or
nonexistent **(Fig. 5B and Fig. S14)**. We collected the statistics on the distribution of the cell
growth rates or deformation rates **(Fig. 5C and Fig. S6)**. When the positive and negative values
were differentiated, it was evident that the cell growth rate could be divided into four states: the
lateral-fast rate (WF), the longitudinal-fast rate (LF), the lateral-slow rate (WS) and the
longitudinal-slow rate (LS). Among them, L and W represented the presence and absence of
direction in cell deformation, or the length and width, respectively. F and S represent the
magnitude of the "driving force" in cell deformation, or the fast and slow growth rates,
respectively.
To explore the molecular mechanisms underlying these four growth rates, we took both the
direction (L and W) and force (F and S) of cell growth into account. Many studies have shown
that Fus3 and Far1, which determine the direction of cell growth along its long axis, are essential
genes for bud formation during polar cell growth, and their absence results in mis-localization of
shmoo projections (*90, 91, 93, 94, 117*). Thus, the indirect pathway (P_1: "Fus3 → Far1/Cdc24 →
Cdc42 → Bni1"), which is more directional than the direct pathway (P_2: "Fus3 → Bni1"), is the
leader signaling pathway for longitudinal cell growth. Given that the P_1 pathway requires the
sequential activation of multiple proteins and that proteins such as Fus3, Far1, and Cdc24 must
be transported into or out of the nucleus, we proposed that the P_2 pathway had a quicker response
speed to stimulate cell growth than did the P_1 pathway **(Fig. 5D)**. Consequently, the temporal
delay in the actions of these two pathways resulted in P_2-dominated undirected growth (model W:
P_2), followed by the two growth pathways co-stimulating the cell's polar growth (model L: P_1+P_2)
once the P_1 pathway was fully functional. In addition, we proposed that F and S corresponded to
the high and low states in Fus3, respectively. When Fus3 was at a high state (H_1, H_2), there was
enough Fus3 to stimulate growth (model F). Following entry into the low state (L_1, L_2), only a

relatively small amount of Fus3 was available for polar growth (model S).
To confirm that there was indeed a time lag in multi-level protein signaling, we constructed
a dual fluorescence system (*CDC24*_GFP-*FUS3*_RFP strain), in which *CDC24* was linked to the
green fluorescent protein (S65T) and *FUS3* was linked to the red fluorescent protein (yomCherry)
(Fig. 5E and Movies S8–S9). Taking into account the order of signal transduction in yeast in
response to pheromone, the expression levels of Fus3 and Cdc24 would achieve the highest
degree of correlation or match after a certain time lag. Using a cross-correlation function, we
analyzed the in- and out-of-nucleus fluorescence trajectories of Fus3 and Cdc24 to compare the
lag times of various activation sequences **(Fig. 5F)**. The figure shows that the lag time between
Fus3 in the nucleus and Cdc24 (red curve) reaching a maximum correlation was approximately
120 min, while the lag time between Fus3 outside the nucleus and Cdc24 (blue curve) reaching a
maximum correlation was approximately 300 min. The fact that the lag time of Fus3_in and
Cdc24_in/out (red curve) in the figure was shorter than that of Fus3_out and Cdc24_in/out (blue
curve) confirms the existence of a temporal delay effect in multi-level signaling and provides
direct evidence for the temporal distinction in the functioning of the two growth pathways.
By taking the growth rate absolute value, we divided the four states into fast growth rate
and slow growth rate categories **(Fig. 5G and Fig. S7)**. To confirm that the two forces (F and S)
corresponded to the high and low states of Fus3, respectively, the cell growth rate states and
Fus3 gene expression states were measured as the cell polar growth changed over time.
According to the proportion of statistics in the figure, the High (expression) state was primarily
contained within the Fast (growth) state, whereas the Low (expression) state was primarily
contained within the Slow (growth) state **(Fig. 5H and Table 3)**. Consequently, this also
provided quantitative experimental evidence for the cellular growth rate molecular mechanism.

Quantifying the four morphological fates and the phase transition trend

The morphological trajectory of the cell shape over time as described by H_n fluctuated
continuously within a given range **(Fig. 5B)**. Using a hidden Markov chain model to fit the time-
varying trajectories of cell morphology, yeast cells at different pheromone concentrations
exhibited four distinct morphological fates. These four morphological fates corresponded
statistically to the four peaks in the distribution of all cell shapes (F_1–F_4) **(Fig. 6A and Fig. S8)**.
Due to the fact that H_n is a parameter that is extremely sensitive to changes in the size and
number of filled circles within the cell, minute variations in the length or width of a portion of
the cell observed through a microscope can result in the cell being classified as having a different
morphological fate **(Fig. 6B)**. We proposed, on the basis that cell morphology is an accumulation
in growth, that the molecular mechanisms underlying the formation of the four morphological
fates were dependent on the different capacities of cells to grow laterally and longitudinally. To
test the claim that four morphological fates were formed, the synergistic effect of growth ability
in both directions was calculated. The lateral growth capacity of cells was expressed by the rate
of change of the average radius of the filled circle within the cell, whereas the longitudinal
growth capacity was expressed by the rate of change of the sum of the radii of the filled circle.
The existence of four distinct states in the cooperative distribution of the two data, as depicted in
the figure, strongly suggested that the capacity to grow in different directions is a crucial factor
in determining cell morphology **(Fig. 6C and Fig. S9)**.
Notably, a phase transition trend from four states to a dominant state in the cell morphology
distribution between 0.7 μM and 3.0 μM was revealed **(Fig. 6A)**. Moreover, the morphology of
the cells observed under a microscope at a concentration of 3.0 μM was significantly smaller

than that observed at other concentrations **(Fig. 6B)**. By measuring the average cell length, it is possible to conclude that a high-dose pheromone concentration (3.0 μM) disrupted the monotonic linear trend observed at low doses (0.7 μM-2.0 μM) **(Fig. 6D and Table 4)**. The microscopic explanation of this phase transition trend is the enhancement of negative feedback regulation under high doses (I_2), suppressing the second polar growth pathway (P_2) **(Fig. 1 and Fig. 3C)**. Three-quarters of the cell fates (with the exception of F_2) showed a sudden reduction in morphology at a high pheromone dose (3.0 μM), most likely attributed to this impaired P_2 function **(Fig. 6A and Fig. S8)**.

To explain the physical mechanism of the phase transition trend that occurs at high dose (3.0 μM), we quantified the degree of nonequilibrium in the cellular morphological system using the net flux. From the cell morphological trajectories analyzed by the hidden Markov chain model, we determined the transition probability between various cell morphologies. The sum of the three net fluxes, which was obtained by decomposing the probability loops in the transition matrix, was used to represent the degree of the detailed balance collapsed in the cellular morphological system **(Fig. 6E)**. As the concentration of pheromone increased, the intensity of the net flux decreased first and then increased, exhibiting a significant phase transition trend at 3.0 μM. Since the net flux is rotational due to its steady state nature under local probability conservation, it tends to destabilize the point attractors. Therefore, the significant changes in net flux can lead to the instability of the cell attractor states, giving rise to possible phase transition trend. The quantification of this non-equilibrium dynamic explains the physical mechanism by which the phase transition trend in a morphological system was caused by the enhanced capability of the yeast cells negative feedback regulation at high doses.

Simulations for the signal transduction and the cell growth

Based on the functional and quantitative regulation obtained from databases, such as KEGG (Kyoto Encyclopedia of Genes and Genomes, https://www.kegg.jp/), SGD (the *Saccharomyces* Genome Database, https://www.yeastgenome.org/) and EVEX (http://evexdb.org/), we developed a simplified model of signal transduction in the context of global gene regulatory networks **(Fig. 1, Fig. 6F)**. This model simulated a series of biochemical reactions with the Gillespie algorithm (*118-120*) **(Table S6–8)**. By simulating the biochemical reactions associated with the pheromone pathway in yeast cells, the expression levels of Fus3 and the distribution of cell morphology were determined. Fus3 gene expression obtained from biochemical reaction results displayed a two-state distribution inside and outside of the nucleus, validating our understanding of the molecular mechanism underlying the bimodal fluorescence state within the context of the global response **(Fig. 6G)**. Meanwhile, the Bni1 produced by the reactions simulated the dynamic process of cell growth. Although both the growth pathways (P_1 and P_2) were involved in the process of cellular length and width growth, the relative weights of the pathways that grew in the two directions were significantly different. Therefore, we simply defined Bni1_in produced by the P_1 pathway that plays a major role as the longitudinal growth and Bni1_out produced by the P_2 pathway as the lateral growth.

In this simulation, a denotes the longitudinal length of the yeast cell, while b denotes the lateral length of the yeast cell **(Fig. 6F)**. Each iteration increased the longitudinal and lateral lengths of the yeast cells by Δa and Δb, respectively. Δa is proportional to "PBni1_in" and inversely proportional to b; Δb is proportional to "PBni1_out" and inversely proportional to a. That is, $\Delta a = \frac{r_a(\text{PBni1\_in})}{b}$, $\Delta b = \frac{r_b(\text{PBni1\_out})}{a}$. The value N corresponds to the rounding operation

345 of $\frac{a}{b}$, i.e., $\left[\frac{a}{b}\right]$, and c represents the amount remaining after rounding. The H_n is $(N +$
346 $1)(N/b + 1/c)$. As predicted by the simulation of the cell growth process, the distribution of
347 cell morphology revealed four distinct fates, which corresponded to the experimental
348 observations **(Fig. 6H)**.

DISCUSSION

In this study, we quantitatively uncovered and interpreted the yeast cell fate decision-
making in response to pheromone using biological and physical methods. Using flow cytometry,
we examined the induction of yeast cells by different concentrations of pheromone to determine
the critical threshold for eliciting a cellular response. The purpose of the fitting curve in **Fig. 2A**
was to estimate the pheromone dose by determining at what concentration the expression of Fus3
exhibited double peaks. Continuous microscopy observations of *FUS3*-GFP strain yeast cells
provided us with a microscopic and mesoscopic view of how the cells responded to pheromone.
The four stages of Fus3 expression levels at a 0.7 µM pheromone concentration accurately
depicted the mesoscopic cell behavior of cell cycle arrest and polar growth **(Fig. 2B)**. Among
them, the Fus3 expression levels in the stable stage (b2–b3) fluctuate around a fixed value, which
was considered a non-equilibrium steady state period. As our subsequent analysis centered on the
non-equilibrium steady-state phase, we chose five pheromone concentrations as stimuli that
allowed the yeast cells to maintain a stable-state phase for a sufficient amount of time.
To explore the fate of Fus3 gene expression in a non-equilibrium steady state, we chose data
that, after 600 min, brought all the cell fluorescence trajectories into stable-state phase **(Fig. 3A)**.
The two fates of Fus3 that resulted from cellular decision-making were separated by the Markov
fitting of the trajectories **(Fig. 3B)**. The criterion for data fitting is the probability that the
fluorescence intensity at a particular moment in the trajectory belonged to a high state (H_1, H_2)
or a low state (L_1, L_2) multiplied by their respective transition probabilities **(Fig. 3E)**.
Consequently, there was a chance that the value of the fluorescence intensity in the low state was
greater than the value of the fluorescence intensity in the high state. We proposed that the two
fates of Fus3 expressions observed in signaling were explained by a logical sequence of point-to-
point feedback regulation between proteins. Notably, even though the protein-to-protein
feedback regulation described above had been validated by previous biological experiments (*85-
89, 95-104*), our claims focused primarily on establishing a dynamical logical connection
between these isolated regulatory processes from a global perspective. For the purpose of
validating this dynamic molecular mechanism, one may wonder why the (L_1, H_2) and (H_1, L_2)
states did not appear in this scheme. We know from experimental evidence that the absence of
the (L_1, H_2) and (H_1, L_2) states was due to the strong correlation between intra- and extra-
nuclear fluorescence intensities. If the four fates emerge on the landscape of intra- and extra-
nuclear fluorescence intensities of Fus3, the Pearson correlation coefficient between these two
types of trajectories would be zero (*121*). Moreover, the feedback model of forward activation
and negative inhibition favored the formation of two states from a physical standpoint **(Fig. 3D)**.
Between steady states, the cellular decision-making is primarily reflected by the transition
rate, transition probability, residence time, and potential barrier height. Due to the relative nature
of the high and low states at each pheromone concentration, their physical characteristics can be
compared directly, but not at different concentrations. To compare the relative significance of
these physical characteristics at different concentrations, a similar normalized approach
(high/low) was used. At 0.7–0.8 µM, a slight increase in barrier height significantly lengthened
the residence time of the high state. At 1.0–3.0 µM, the residence time also decreased gradually

as the barrier height decreased **(Fig. 4D)**. In our biological system, the positive correlation
statement regarding barrier height and residence time was confirmed **(Fig. 4C)**. Certainly, such
physical characteristics that quantify cellular decision-making have significant biological
implications for the study of yeast response behavior. Positive feedback regulation in the gene
network ("Pheromone → Fus3" and "Fus3 → Fus3"), for example, increased the potential
barrier height on the gene expression landscape, whereas negative feedback regulation decreased
it (I_1, I_2) **(Fig. 1 and Fig. 3C)**. The residence time at various pheromone concentrations revealed
which gene expression fate yeast cells preferred in response to the pheromone.

Some studies believe that the establishment of polarity helps organisms to survive better in
nature (*122, 123*). Due to the inability of yeast's chemotactic response to swim as that of E. coli,
its polar growth could only grow in the direction of a high concentration of pheromone,
achieving mutual contact and fusion between heterosexual yeasts (*4, 124, 125*). To quantify the
mesoscopic behavior of yeast cells in polar growth, which grew and stopped intermittently, the
yeast cells were characterized as a combination of filled circles. The white edge of the yeast cell
image was the cell wall **(Fig. 5A).** Therefore, the cell filling circles were based on the outermost
edge of the cell wall as the cell boundary. Rather than using the cell area, the total length, or the
arithmetic mean of the length, we chose to describe the cell morphology using the reciprocal of
the radius in the cell-filled circle multiplied by the number of filled circles, i.e., $H_n = n(\frac{1}{R_1} +$
$\frac{1}{R_2} + \ldots + \frac{1}{R_n})$. This method has the following advantages: first, it could reflect the deformation of
different parts of the cell; and second, H_n is a statistic that can reflect both the process and
outcome of the cell polarity growth. It was important to note that when a cell grew laterally, its
length also increased, so the terms lateral growth and longitudinal growth were relative rather
than absolute descriptions of length and width.

The cell growth rate primarily characterized the process of cell polar growth, whereas the
cell morphology primarily characterized the outcome. Specifically, the cell growth rate is
actually the cell deformation rate. To verify the molecular mechanism of the growth rate for
directional (model L) and non-directional (model W) growth, we constructed a dual fluorescence
system (*CDC24*_GFP-*FUS3*_RFP) to demonstrate that there was a time lag in the multilevel
protein signaling behavior. Due to the different maturation times of the two fluorescent proteins
(GFP and RFP), the lag time between Fus3 and Cdc24 was only a relative quantity. Nonetheless,
the delay time of the dual-fluorescence system at different positions was used to quantify the
practical significance of multipolar signal transduction. Notably, the non-directional growth
(model W: P_2) did not consist solely of the polar growth pathway of P_2, but rather a relative
weight simplification to differentiate it from the directional growth (model L: $P_1 + P_2$). To confirm
that the fast growth rate (model F) and slow growth rate (model S) corresponded to the high and
low states of Fus3, we explored the association between growth rates and dual fluorescent gene
expression states. The high state (H_1, H_2) was primarily included in the fast growth rate, whereas
the low state (L_1, L_2) was primarily included in the slow growth rate, confirming the molecular
mechanism of F and S **(Fig. 5H and Table 3)**. In addition, there must be a lag between the
expression of Fus3p and the observed cell growth, resulting in a low fluorescence state during
fast cell growth and a high fluorescence state during slow cell growth.

For statistical analysis of different cell morphologies, we distinguished four cell
morphological fates roughly as F_1-F_4 to provide a more intuitive description **(Fig. 6A)**.
Nevertheless, the division of actual cell morphological fate was determined by fitting the
trajectories, and not arbitrarily by the size of H_n. Given that H_n is extremely sensitive to changes

in length and width in different parts of the cell, the area of cells that were morphologically
divided into the four categories did not differ significantly from their overall appearance **(Fig.**
6B). Our justification for proposing the molecular mechanisms of the four cell morphological
fates is that morphology is the accumulation of a growth process. Due to the fact that the cell
morphology described by H_n resembled the relative length and width of a cell, the capacity to
grow in both directions can explain the emergence of cell morphologies **(Fig. 6C)**. In addition, in
the statistical graph of cell morphology, other fates tended to disappear, with the exception of the
increase in the proportion of F_2. This indicated that the intracellular phase transition trend
occurred at 3.0 μM as a result of enhanced negative feedback regulation (I_2 in **Fig. 1**). The
coordinates of the cell morphology statistics chart at 3.0 μM were deviated from those of other
concentrations due to the phase transition trend in the system, which altered the measurement
scale of the four morphological fates. To explain the physics of the phase transitions in the
biological system, we quantified the degree to which the detailed balance collapsed by
employing the net flux characterizing the non-equilibrium statistical physics. The results
demonstrated that the non-equilibrium dynamics of the biological system was supported by the
significant increase in net flux of the morphological system at high pheromone dose (3.0 μM),
causing the instability and phase transition.
Finally, we developed a simplified model of the gene regulatory network for signal
transduction in order to confirm the global rationality of the dynamic sequential mechanism.
Thus far, we have established the logical links between the functions of regulated proteins in the
pheromone pathway for mating through the underlying signal transduction process. While our
validation method differs from the conventional approach of constructing yeast mutants, it has
been successful in observing realistic cellular response behavior. Furthermore, our future
research will focus on utilizing fluorescent labels to enable real-time visualization of the
dynamics of signaling pathways. Specifically, we plan to construct a multi-dimensional
fluorescent system to elucidate the temporal and spatial dynamics of cellular responses,
including potential crosstalk between different signaling pathways. This approach will provide a
more comprehensive and quantitative understanding of how cells respond to various stimuli, and
how these responses are integrated at the molecular level.
464

MATERIALS AND METHODS

Yeast strains and Growth media

The yeast strain used in this experiment is *Saccharomyces cerevisiae* S288C (ATCC 201388: *MATahis3*Δ1 *leu*2Δ0 *met*15Δ0 *ura*3Δ0) (*126*). In the yeast fluorescence library, we choose a special yeast as the research object, whose c-terminal of Fus3 is fused with GFP as the reporter protein. The yeasts are cultured by inoculating 5ml of YNB [Yeast Nitrogen Base Without Amino Acids (6.8 g/L), Dextrose (5 g/L), Uracil (76mg/L), 50×MEM Amino Acids (20ml/L)] medium with a colony from a YPD agar plate [Yeast extract (10 g/L), Peptone (20 g/L), Dextrose (20 g/L), Agar (10 g/L)]. Cells are grown overnight at 30□ shaking at 250 rpm (about 14h). Before starting the experiment, we take out 20ul of overnight cultures and dilute 100-fold in 2ml YNB medium to obtain an optical density of 0.1 at 600 nm (OD600nm).

Flow cytometry

To determine the critical concentration of pheromone, we chose a flow cytometer for preliminary screening. We set 8 different concentrations (0.01μM, 0.05μM, 0.1μM, 0.2 μM, 0.4μM, 0.6μM, 1.0μM, 1.5μM) for the pheromone medium, and culture the yeast cells in a shaker at 250 rpm and 30□. In order to avoid contamination of the yeast cells, we have set up 5 sets of parallel culture samples for each concentration of the medium. After culturing on a shaker for specific times (3h, 6h, 9h, 12h and 24h), 105,000 cells are taken from each sample for analysis using the flow cytometer.

Microfluidic system platform

We use the 4-chamber cell culture plate, CellASIC™ ONIX Y04C-02 Microfluidic Plate, for live cell culture at 30°C. The plate is designed for use with the CellASIC™ ONIX Microfluidic System, which provides a controlled dynamical microenvironment for cells. First, the yeast cells are cultured in a microfluidic plate containing conventional YNB medium for 1 hour to adapt to the new environment, and then switch to YNB medium supplemented with pheromone and culture at a flow rate of 1 psi for 30 hours.

Microscopy Measurements

The fluorescence values of the single cells were measured using an inverted fluorescence microscopy (Ti-E, Nikon) with automated stage and focus, equipped with a high NA oil-immersion objective (1.45NA, 100×). We applied 488nm laser and set the output power at 30mW (only 10% of the laser beam into the microscope objective), the fluorescence signals were collected by a cooled EM-CCD camera (897U, Andor). All images were acquired using both bright field imaging and fluorescent field imaging. These images were acquired by Nikon software. Data analysis were accomplished through a combination of manual and automated analysis using custom Matlab code. Many trajectories were taken from a time-lapse microscopy. The fluorescent images were periodically captured and recorded every 10 minutes. The fluorescent data of each cell at each time point were collected for the following discussion.

References and Notes:

1. H. C. Berg, D. A. Brown, Chemotaxis in Escherichia coli analysed by three-dimensional tracking. *Nature* **239**, 500-504 (1972).
2. R. M. Macnab, D. E. Koshland Jr, The gradient-sensing mechanism in bacterial chemotaxis. *Proceedings of the National Academy of Sciences* **69**, 2509-2512 (1972).
3. J. Yuan, K. A. Fahrner, L. Turner, H. C. Berg, Asymmetry in the clockwise and counterclockwise rotation of the bacterial flagellar motor. *Proceedings of the National Academy of Sciences* **107**, 12846-12849 (2010).
4. V. Sourjik, N. S. Wingreen, Responding to chemical gradients: bacterial chemotaxis. *Current opinion in cell biology* **24**, 262-268 (2012).
5. S. De Oliveira, E. E. Rosowski, A. Huttenlocher, Neutrophil migration in infection and wound repair: going forward in reverse. *Nature Reviews Immunology* **16**, 378-391 (2016).
6. P. Zhou *et al.*, Stochasticity triggers activation of the S-phase checkpoint pathway in budding yeast. *Physical Review X* **11**, 011004 (2021).
7. B. Fu *et al.*, Metal-induced sensor mobilization turns on affinity to activate regulator for metal detoxification in live bacteria. *Proceedings of the National Academy of Sciences* **117**, 13248-13255 (2020).
8. J. E. Slessareva, H. G. Dohlman, G protein signaling in yeast: new components, new connections, new compartments. *Science* **314**, 1412-1413 (2006).
9. R. E. Chen, J. Thorner, Function and regulation in MAPK signaling pathways: Lessons learned from the yeast Saccharomyces cerevisiae. *Bba-Mol Cell Res* **1773**, 1311-1340 (2007).
10. Y. Q. Wang, H. G. Dohlman, Pheromone signaling mechanisms in yeast: A prototypical sex machine. *Science* **306**, 1508-1509 (2004).
11. M. C. Gustin, J. Albertyn, M. Alexander, K. Davenport, MAP Kinase Pathways in the YeastSaccharomyces cerevisiae. *Microbiology and Molecular biology reviews* **62**, 1264-1300 (1998).
12. I. Herskowitz, Life cycle of the budding yeast Saccharomyces cerevisiae. *Microbiological reviews* **52**, 536 (1988).
13. C. M. Hull, R. M. Raisner, A. D. Johnson, Evidence for mating of the" asexual" yeast Candida albicans in a mammalian host. *Science* **289**, 307-310 (2000).
14. J. Warringer *et al.*, Trait variation in yeast is defined by population history. *PLoS Genet* **7**, e1002111 (2011).
15. E. Zörgö *et al.*, Ancient evolutionary trade-offs between yeast ploidy states. *PLoS Genet* **9**, e1003388 (2013).
16. D. Norris, M. Osley, The two gene pairs encoding H2A and H2B play different roles in the Saccharomyces cerevisiae life cycle. *Molecular and Cellular Biology* **7**, 3473-3481 (1987).
17. J. E. Haber, Mating-type genes and MAT switching in Saccharomyces cerevisiae. *Genetics* **191**, 33-64 (2012).
18. M. Z. Bao, M. A. Schwartz, G. T. Cantin, J. R. Yates III, H. D. Madhani, Pheromone-dependent destruction of the Tec1 transcription factor is required for MAP kinase signaling specificity in yeast. *Cell* **119**, 991-1000 (2004).
19. D. D. Jenness, A. C. Burkholder, L. H. Hartwell, Binding of α-factor pheromone to yeast a cells: chemical and genetic evidence for an α-factor receptor. *Cell* **35**, 521-529 (1983).
20. H. Youk, W. A. Lim, Secreting and sensing the same molecule allows cells to achieve

versatile social behaviors. *Science* **343**, 1242782 (2014).

21.   H. Youk, W. A. Lim, Secreting and Sensing the Same Molecule Allows Cells to Achieve
Versatile Social Behaviors. *Science* **343**, 628-+ (2014).

22.   M. D. Rose, Nuclear fusion in the yeast Saccharomyces cerevisiae. *Annu Rev Cell Dev Bi*
12, 663-695 (1996).

23.   L. Marsh, M. Rose, The pathway of cell and nuclear fusion during mating in S. cerevisiae.
COLD SPRING HARBOR MONOGRAPH SERIES **21**, 827-888 (1997).

24.   J. M. White, M. D. Rose, Yeast mating: getting close to membrane merger. *Current*
biology **11**, R16-R20 (2001).

25.   L. Bardwell, A walk-through of the yeast mating pheromone response pathway (vol 25,
pg 1465, 2004). *Peptides* **26**, 337-+ (2005).

26.   J. N. Molk, K. Bloom, Microtubule dynamics in the budding yeast mating pathway.
Journal of cell science **119**, 3485-3490 (2006).

27.   H. G. Dohlman, J. W. Thorner, Regulation of G protein-initiated signal transduction in
yeast: Paradigms and principles. *Annu Rev Biochem* **70**, 703-754 (2001).

28.   A. M. Neiman, I. Herskowitz, Reconstitution of a Yeast Protein-Kinase Cascade in-Vitro
- Activation of the Yeast Mek Homolog Ste7 by Ste11. *P Natl Acad Sci USA* **91**, 3398-
3402 (1994).

29.   B. Errede, A. Gartner, Z. Zhou, K. Nasmyth, G. Ammerer, MAP kinase-related FUS3
from S. cerevisiae is activated by STE7 in vitro. *Nature* **362**, 261-264 (1993).

30.   P. M. P. Anne-Christine Butty, Linda S. Huang, Ira Herskowitz, Matthias Peter*, The
Role of Far1p in Linking the Heterotrimeric G Protein to Polarity Establishment Proteins
During Yeast Mating. *SCIENCE*, (1998).

31.   D. Matheos, M. Metodiev, E. Muller, D. Stone, M. D. Rose, Pheromone-induced
polarization is dependent on the Fus3p MAPK acting through the formin Bni1p. *The*
Journal of cell biology **165**, 99-109 (2004).

32.   M. Peter, A. Gartner, J. Horecka, G. Ammerer, I. Herskowitz, FAR1 links the signal
transduction pathway to the cell cycle machinery in yeast. *Cell* **73**, 747-760 (1993).

33.   E. Elion, B. Satterberg, J. Kranz, FUS3 phosphorylates multiple components of the
mating signal transduction cascade: evidence for STE12 and FAR1. *Molecular biology of*
the cell **4**, 495-510 (1993).

34.   R. L. Roberts, G. R. Fink, Elements of a single MAP kinase cascade in Saccharomyces
cerevisiae mediate two developmental programs in the same cell type: mating and
invasive growth. *Genes & development* **8**, 2974-2985 (1994).

35.   I. Herskowitz, MAP kinase pathways in yeast: for mating and more. *Cell* **80**, 187-197
(1995).

36.   E. A. Elion, J. A. Brill, G. R. Fink, FUS3 represses CLN1 and CLN2 and in concert with
KSS1 promotes signal transduction. *Proceedings of the National Academy of Sciences* **88**,
9392-9396 (1991).

37.   E. Blackwell *et al.*, Effect of the pheromone-responsive Gα and phosphatase proteins of
Saccharomyces cerevisiae on the subcellular localization of the Fus3 mitogen-activated
protein kinase. *Molecular and cellular biology* **23**, 1135-1150 (2003).

38.   F. van Drogen, V. M. Stucke, G. Jorritsma, M. Peter, MAP kinase dynamics in response
to pheromones in budding yeast. *Nature cell biology* **3**, 1051-1059 (2001).

39.   F. Chang, I. Herskowitz, Identification of a gene necessary for cell cycle arrest by a
negative growth factor of yeast: FAR1 is an inhibitor of a G1 cyclin, CLN2. *Cell* **63**, 999-

595   1011 (1990).

596 40. Y. Shimada, M.-P. Gulli, M. Peter, Nuclear sequestration of the exchange factor Cdc24 by
597   Far1 regulates cell polarity during yeast mating. *Nature cell biology* **2**, 117-124 (2000).

598 41. J. E. Segall, Polarization of yeast cells in spatial gradients of alpha mating factor.
599   *Proceedings of the National Academy of Sciences* **90**, 8332-8336 (1993).

600 42. X. N. Fang *et al.*, Cell fate potentials and switching kinetics uncovered in a classic
601   bistable genetic switch. *Nature Communications* **9**, (2018).

602 43. Z. L. Jiang *et al.*, The emergence of the two cell fates and their associated switching for a
603   negative auto-regulating gene. *Bmc Biology* **17**, (2019).

604 44. P. A. Takizawa, A. Sil, J. R. Swedlow, I. Herskowitz, R. D. Vale, Actin-dependent
605   localization of an RNA encoding a cell-fate determinant in yeast. *Nature* **389**, 90-93
606   (1997).

607 45. A. Colman-Lerner *et al.*, Regulated cell-to-cell variation in a cell-fate decision system.
608   *Nature* **437**, 699-706 (2005).

609 46. R. M. Gordley *et al.*, Engineering dynamical control of cell fate switching using synthetic
610   phospho-regulons. *P Natl Acad Sci USA* **113**, 13528-13533 (2016).

611 47. Z. Jiang *et al.*, The emergence of the two cell fates and their associated switching for a
612   negative auto-regulating gene. *BMC biology* **17**, 1-14 (2019).

613 48. L. Zeng *et al.*, Decision making at a subcellular level determines the outcome of
614   bacteriophage infection. *Cell* **141**, 682-691 (2010).

615 49. A. Colman-Lerner *et al.*, Regulated cell-to-cell variation in a cell-fate decision system.
616   *Nature* **437**, 699-706 (2005).

617 50. X. Fang *et al.*, Cell fate potentials and switching kinetics uncovered in a classic bistable
618   genetic switch. *Nature communications* **9**, 1-9 (2018).

619 51. F. St-Pierre, D. Endy, Determination of cell fate selection during phage lambda infection.
620   *Proceedings of the National Academy of Sciences* **105**, 20705-20710 (2008).

621 52. P. J. Choi, L. Cai, K. Frieda, X. S. Xie, A stochastic single-molecule event triggers
622   phenotype switching of a bacterial cell. *Science* **322**, 442-446 (2008).

623 53. L. S. Weinberger, J. C. Burnett, J. E. Toettcher, A. P. Arkin, D. V. Schaffer, Stochastic
624   gene expression in a lentiviral positive-feedback loop: HIV-1 Tat fluctuations drive
625   phenotypic diversity. *Cell* **122**, 169-182 (2005).

626 54. W. Wang *et al.*, Live-cell imaging and analysis reveal cell phenotypic transition dynamics
627   inherently missing in snapshot data. *Sci Adv* **6**, eaba9319 (2020).

628 55. N. Yosef, A. Regev, Impulse Control: Temporal Dynamics in Gene Transcription. *Cell*
629   **144**, 886-896 (2011).

630 56. Z. Bar-Joseph, A. Gitter, I. Simon, Studying and modelling dynamic biological processes
631   using time-series gene expression data. *Nature Reviews Genetics* **13**, 552-564 (2012).

632 57. M. Behar, N. Hao, H. G. Dohlman, T. C. Elston, Dose-to-Duration Encoding and
633   Signaling beyond Saturation in Intracellular Signaling Networks. *Plos Computational*
634   *Biology* **4**, (2008).

635 58. A. Doncic *et al.*, Compartmentalization of a Bistable Switch Enables Memory to Cross a
636   Feedback-Driven Transition. *Cell* **160**, 1182-1195 (2015).

637 59. J. Hasty, D. McMillen, F. Isaacs, J. J. Collins, Computational studies of gene regulatory
638   networks: in numero molecular biology. *Nature Reviews Genetics* **2**, 268-279 (2001).

639 60. F. Caudron, Y. Barral, A super-assembly of Whi3 encodes memory of deceptive
640   encounters by single cells during yeast courtship. *Cell* **155**, 1244-1257 (2013).

61. J. S. Weinberg, D. G. Drubin, Regulation of clathrin-mediated endocytosis by dynamic ubiquitination and deubiquitination. *Curr Biol* **24**, 951-959 (2014).

62. M. D. Leonetti, S. Sekine, D. Kamiyama, J. S. Weissman, B. Huang, A scalable strategy for high-throughput GFP tagging of endogenous human proteins. *Proc Natl Acad Sci U S A* **113**, E3501-3508 (2016).

63. R. G.-L. Patrick Conlon, Ambhighainath Ganesan, Jin Zhang, and Andre Levchenko, Single-cell dynamics and variability of MAPK activity in a yeast differentiation pathway. *PNAS*, (2016).

64. S. Ozsezen *et al.*, Inference of the High-Level Interaction Topology between the Metabolic and Cell-Cycle Oscillators from Single-Cell Dynamics. *Cell Syst* **9**, 354-365 e356 (2019).

65. A. K. Chakravarty, T. Smejkal, A. K. Itakura, D. M. Garcia, D. F. Jarosz, A Non-amyloid Prion Particle that Activates a Heritable Gene Expression Program. *Mol Cell* **77**, 251-265 e259 (2020).

66. C. L. Klaips, M. H. M. Gropp, M. S. Hipp, F. U. Hartl, Sis1 potentiates the stress response to protein aggregation and elevated temperature. *Nature Communications* **11**, (2020).

67. S. N. Mouton *et al.*, A physicochemical perspective of aging from single-cell analysis of pH, macromolecular and organellar crowding in yeast. *Elife* **9**, (2020).

68. A. K. Pfitzner *et al.*, An ESCRT-III Polymerization Sequence Drives Membrane Deformation and Fission. *Cell* **182**, 1140-1155 e1118 (2020).

69. C. Ptak *et al.*, Phosphorylation-dependent mitotic SUMOylation drives nuclear envelope-chromatin interactions. *J Cell Biol* **220**, (2021).

70. W. Feng, O. Argüello-Miranda, S. Qian, F. Wang, Cdc14 spatiotemporally dephosphorylates Atg13 to activate autophagy during meiotic divisions. *Journal of Cell Biology* **221**, (2022).

71. S. Pinheiro, S. Pandey, S. Pelet, Cellular heterogeneity: yeast-side story. *Fungal Biology Reviews* **39**, 34-45 (2022).

72. S. Chaturvedi *et al.*, Disrupting autorepression circuitry generates "open-loop lethality" to yield escape-resistant antiviral agents. *Cell* **185**, 2086-2102 e2022 (2022).

73. W. Wang, G. W. Li, C. Chen, X. S. Xie, X. Zhuang, Chromosome organization by a nucleoid-associated protein in live bacteria. *Science* **333**, 1445-1449 (2011).

74. H. Feng, J. Wang, Potential and flux decomposition for dynamical systems and non-equilibrium thermodynamics: Curvature, gauge field, and generalized fluctuation-dissipation theorem. *The Journal of chemical physics* **135**, 234511 (2011).

75. C. Li, J. Wang, Landscape and flux reveal a new global view and physical quantification of mammalian cell cycle. *Proceedings of the National Academy of Sciences* **111**, 14130-14135 (2014).

76. G. Balázsi, A. van Oudenaarden, J. J. Collins, Cellular decision making and biological noise: from microbes to mammals. *Cell* **144**, 910-925 (2011).

77. J. Wang, K. Zhang, L. Xu, E. Wang, Quantifying the Waddington landscape and biological paths for development and differentiation. *Proceedings of the National Academy of Sciences* **108**, 8257-8262 (2011).

78. J. Wang, K. Zhang, E. Wang, Kinetic paths, time scale, and underlying landscapes: A path integral framework to study global natures of nonequilibrium systems and networks. *The Journal of chemical physics* **133**, 09B613 (2010).

687   79.   J. Wang, C. Li, E. Wang, Potential and flux landscapes quantify the stability and
688   robustness of budding yeast cell cycle network. *Proceedings of the National Academy of*
689   *Sciences* **107**, 8195-8200 (2010).
690   80.   J. Wang, K. Zhang, E. Wang, Robustness and dissipation of mitogen-activated protein
691   kinases signal transduction network: Underlying funneled landscape against stochastic
692   fluctuations. *The Journal of chemical physics* **129**, 10B602 (2008).
693   81.   J. Wang, L. Xu, E. Wang, Potential landscape and flux framework of nonequilibrium
694   networks: robustness, dissipation, and coherence of biochemical oscillations. *Proceedings*
695   *of the National Academy of Sciences* **105**, 12271-12276 (2008).
696   82.   E. H. Davidson, D. H. Erwin, Gene regulatory networks and the evolution of animal body
697   plans. *Science* **311**, 796-800 (2006).
698   83.   N. G. Van Kampen, *Stochastic processes in physics and chemistry*. (Elsevier, 1992), vol.
699   1.
700   84.   W. Zheng, H. Zhao, E. Mancera, L. M. Steinmetz, M. Snyder, Genetic analysis of
701   variation in transcription factor binding in yeast. *Nature* **464**, 1187-1191 (2010).
702   85.   A. Anders, B. Ghosh, T. Glatter, V. Sourjik, Design of a MAPK signalling cascade
703   balances energetic cost versus accuracy of information transmission. *Nature*
704   *communications* **11**, 1-10 (2020).
705   86.   H. G. Dohlman, J. Song, D. Ma, W. E. Courchesne, J. Thorner, Sst2, a negative regulator
706   of pheromone signaling in the yeast Saccharomyces cerevisiae: expression, localization,
707   and genetic interaction and physical association with Gpa1 (the G-protein alpha subunit).
708   *Molecular and cellular biology* **16**, 5194-5209 (1996).
709   87.   M. Sugiyama, Y. Kaneko, S. Harashima, Yeast protein phosphatases Ptp2p and Msg5p are
710   involved in G1–S transition, CLN2 transcription, and vacuole morphogenesis. *Archives of*
711   *microbiology* **191**, 721-733 (2009).
712   88.   W. A. Laviña, M. Sugiyama, Y. Kaneko, S. Harashima, Identification of protein kinase
713   disruptions as suppressors of the calcium sensitivity of S. cerevisiae Δ ptp2 Δ msg5
714   protein phosphatase double disruptant. *Archives of microbiology* **192**, 157-165 (2010).
715   89.   W. A. Laviña, M. Sugiyama, Y. Kaneko, S. Harashima, Functionally redundant protein
716   phosphatase genes PTP2 and MSG5 co-regulate the calcium signaling pathway in
717   Saccharomyces cerevisiae upon exposure to high extracellular calcium concentration.
718   *Journal of bioscience and bioengineering* **115**, 138-146 (2013).
719   90.   A.-C. Butty, P. M. Pryciak, L. S. Huang, I. Herskowitz, M. Peter, The role of Far1p in
720   linking the heterotrimeric G protein to polarity establishment proteins during yeast
721   mating. *Science* **282**, 1511-1516 (1998).
722   91.   A. Nern, R. A. Arkowitz, A Cdc24p-Far1p-Gβγ protein complex required for yeast
723   orientation during mating. *The Journal of cell biology* **144**, 1187-1202 (1999).
724   92.   A. Nern, R. A. Arkowitz, Nucleocytoplasmic shuttling of the Cdc42p exchange factor
725   Cdc24p. *The Journal of cell biology* **148**, 1115-1122 (2000).
726   93.   A. Nern, R. A. Arkowitz, A Cdc24p-Far1p-Gbetagamma protein complex required for
727   yeast orientation during mating. *J Cell Biol* **144**, 1187-1202 (1999).
728   94.   E. H. Chen, *Cell fusion: overviews and methods*. (Springer, 2008).
729   95.   E. Szczurek, I. Gat☐Viks, J. Tiuryn, M. Vingron, Elucidating regulatory mechanisms
730   downstream of a signaling pathway using informative experiments. *Molecular Systems*
731   *Biology* **5**, 287 (2009).
732   96.   C. I. Maeder *et al.*, Spatial regulation of Fus3 MAP kinase activity through a reaction-

diffusion mechanism in yeast pheromone signalling. *Nature cell biology* **9**, 1319-1326 (2007).

97. A. Serrano *et al.*, Spatio-temporal MAPK dynamics mediate cell behavior coordination during fungal somatic cell fusion. *Journal of cell science* **131**, (2018).

98. J. E. Kranz, B. Satterberg, E. A. Elion, The MAP kinase Fus3 associates with and phosphorylates the upstream signaling component Ste5. *Genes & Development* **8**, 313-327 (1994).

99. M. J. Winters, R. E. Lamson, H. Nakanishi, A. M. Neiman, P. M. Pryciak, A membrane binding domain in the Ste5 scaffold synergizes with Gβγ binding to control localization and signaling in pheromone response. *Molecular cell* **20**, 21-32 (2005).

100. R. P. Bhattacharyya *et al.*, The Ste5 scaffold allosterically modulates signaling output of the yeast mating pathway. *Science* **311**, 822-826 (2006).

101. M. K. Malleshaiah, V. Shahrezaei, P. S. Swain, S. W. Michnick, The scaffold protein Ste5 directly controls a switch-like mating decision in yeast. *Nature* **465**, 101-105 (2010).

102. T. Chen, J. Kurjan, Saccharomyces cerevisiae Mpt5p interacts with Sst2p and plays roles in pheromone sensitivity and recovery from pheromone arrest. *Molecular and Cellular Biology* **17**, 3429-3439 (1997).

103. T. R. Garrison *et al.*, Feedback phosphorylation of an RGS protein by MAP kinase in yeast. *Journal of Biological Chemistry* **274**, 36387-36391 (1999).

104. S. C. Parnell *et al.*, Phosphorylation of the RGS protein Sst2 by the MAP kinase Fus3 and use of Sst2 as a model to analyze determinants of substrate sequence specificity. *Biochemistry* **44**, 8159-8166 (2005).

105. R. E. Chen, J. Thorner, Function and regulation in MAPK signaling pathways: lessons learned from the yeast Saccharomyces cerevisiae. *Biochimica et Biophysica Acta (BBA)-Molecular Cell Research* **1773**, 1311-1340 (2007).

106. H.-O. Park, E. Bi, Central roles of small GTPases in the development of cell polarity in yeast and beyond. *Microbiology and molecular biology reviews* **71**, 48-96 (2007).

107. J. B. Moseley, B. L. Goode, The yeast actin cytoskeleton: from cellular function to biochemical mechanism. *Microbiology and Molecular Biology Reviews* **70**, 605-645 (2006).

108. A. Doncic, M. Falleur-Fettig, J. M. Skotheim, Distinct interactions select and maintain a specific cell fate. *Molecular cell* **43**, 528-539 (2011).

109. N. Barkai, M. D. Rose, N. S. Wingreen, Protease helps yeast find mating partners. *Nature* **396**, 422-423 (1998).

110. B. D. Slaughter, J. W. Schwartz, R. Li, Mapping dynamic protein interactions in MAP kinase signaling using live-cell fluorescence fluctuation spectroscopy and imaging. *Proceedings of the National Academy of Sciences* **104**, 20320-20325 (2007).

111. E. A. Elion, M. Qi, W. Chen, Signaling specificity in yeast. *Science* **307**, 687-688 (2005).

112. D. Fiedler *et al.*, Functional organization of the S. cerevisiae phosphorylation network. *Cell* **136**, 952-963 (2009).

113. J. Wang, Landscape and flux theory of non-equilibrium dynamical systems with application to biology. *Advances in Physics* **64**, 1-137 (2015).

114. X. Fang, K. Kruse, T. Lu, J. Wang, Nonequilibrium physics in biology. *Reviews of Modern Physics* **91**, 045004 (2019).

115. H. Feng, K. Zhang, J. Wang, Non-equilibrium transition state rate theory. *Chemical Science* **5**, 3761-3769 (2014).

116. X. N. Fang, K. Kruse, T. Lu, J. Wang, Nonequilibrium physics in biology. *Reviews of*
Modern Physics **91**, (2019).
117. A. Nern, R. A. Arkowitz, G proteins mediate changes in cell shape by stabilizing the axis
of polarity. *Molecular cell* **5**, 853-864 (2000).
118. D. T. Gillespie, The chemical Langevin equation. *The Journal of Chemical Physics* **113**,
297-306 (2000).
119. D. T. Gillespie, Approximate accelerated stochastic simulation of chemically reacting
systems. *The Journal of chemical physics* **115**, 1716-1733 (2001).
120. H. De Jong, Modeling and simulation of genetic regulatory systems: a literature review.
Journal of computational biology **9**, 67-103 (2002).
121. J. Lee Rodgers, W. A. Nicewander, Thirteen ways to look at the correlation coefficient.
The American Statistician **42**, 59-66 (1988).
122. A. Treuner-Lange, L. Sogaard-Andersen, Regulation of cell polarity in bacteria. *J Cell*
Biol **206**, 7-17 (2014).
123. J. Dworkin, Cellular Polarity in Prokaryotic Organisms. *Csh Perspect Biol* **1**, (2009).
124. R. M. Macnab, D. E. Koshland, The gradient-sensing mechanism in bacterial chemotaxis.
Proceedings of the National Academy of Sciences **69**, 2509-2512 (1972).
125. L. Bardwell, A walk-through of the yeast mating pheromone response pathway. *Peptides*
25, 1465-1476 (2004).
126. S. Ghaemmaghami *et al.*, Global analysis of protein expression in yeast. *Nature* **425**, 737-
741 (2003).

800
801

Acknowledgments:

SL, QL, and EW acknowledge the support of the National Natural Science Foundation of
China with Grant No. 21721003.

Competing interests: Authors declare that they have no competing interests.

Data and materials availability: All data are available in the main text or the supplementary
materials.

808
809

Figure Legends:

Fig. 1. Schematic diagram of the mating signal pathway of the yeast cell pheromone
pathway. The red horizontal line represents the inhibition of negative feedback; the green arrow
represents the activation of positive feedback; the dashed arrow represents a shift in localization;
I_1 and I_2 represent different negative feedback adjustment pathways; and P_1 and P_2 represent the
two polar growth signaling pathways that are connected by light green arrows.
817

818
Fig. 2. A non-equilibrium biological model for cellular responses. (A) The fluorescence
intensity of *FUS3*-GFP was measured using flow cytometry. The yeast cells were cultured in
YPD medium containing different concentrations of pheromone for 24 h; the black curve
represents the overall fluorescence intensity statistics; and the green and red curves represent the
two fitted statistical peaks. The sample size for each pheromone concentration test was 105,000
yeast cells. **(B)** Fus3 gene expression was observed microscopically in response to 0.2 μM and
0.7 μM pheromone. The x-axis represents the duration of exposure to the pheromone-containing
medium. The y-axis represents the fluorescence intensity of *FUS3*-GFP; and images a_1–a_3 and
b_1–b_3 depict the living states of yeast cells as observed through a microscope at their
corresponding times. **(C)** The microscopically captured living state of yeast at various
pheromone concentrations. The green fluorescence in cells represents the expression of *FUS3*-
GFP; 60 min on the x-axis represents the time when yeast cells were switched to a culture
medium containing pheromone.

Fig. 3. The two steady states of expression levels of Fus3. (**A**) Trajectories of fluorescence intensity of Fus3 at 0.7 μM inside and outside the nucleus (only a portion shown). The red vertical line at 600 min was used to approximate the time node at which all cell fluorescence trajectories had entered a non-equilibrium steady state. (**B**) Three-dimensional distribution graph of Fus3 fluorescence intensity inside and outside the nucleus of yeast cells in the stationary phase under different pheromone concentrations. On the left is the 3D distribution of fluorescence or the 3D population landscape, in the middle is the 2D histograms or the 2D underlying potential landscapes U in exponential scale (defined as $p \sim e^{-U}$), which is also the population landscape; on the right is the 2D underlying potential landscapes U ($U = -\ln P$). The sample sizes at steady state for each pheromone concentration are as follows: 0.7 μM was equivalent to 21,335 cells, 0.8 μM to 18,408 cells, 1.0 μM to 23,041 cells, 2.0 μM to 36,886 cells, and 3.0 μM to 18,276 cells. (**C**) Diagram illustrating the molecular mechanism by which yeast cells respond to pheromone. The Outer_P_1 represents the indirect pathway taken by Fus3 from the cytoplasm to the nucleus in order to inhibit the cell cycle; Inner_P_1 represents the indirect pathway by which Fus3 in the nucleus promoted polar growth; Outer_P_2 represents the direct pathway of Fus3 in the cytoplasm for polar growth; Inner_P_2 represents the direct pathway for the transfer of Fus3 from the nucleus to the cytoplasm for polar growth; I_1 and I_2 represent the inhibitory effects of the negative feedback regulation; and the two gray dashed lines represent the cell membrane and the

nuclear membrane. **(D)** Schematic representation of two negative feedback models for regulating Fus3 gene expression. A and B represent, respectively, two different types of proteins that interact with Fus3 in the signaling pathway; "α" stands for α-factor pheromone. **(E)** The fluorescence intensity trajectories of Fus3 inside and outside the nucleus are over time. The red line represents the fitting line of the high state and the low state.

Fig. 4. The physical characteristics of cellular decision-making landscapes. (A) Trends in the
statistical distribution of barrier heights for high and low states at varying pheromone
concentrations. **(B)** Trends in the statistical distribution of residence time for high and low states
at varying pheromone concentrations. **(C)** Correlation analysis between residence time and
barrier height. The red circles are the data points; the blue line is the fitted line of the data. **(D)**
The ratio of the physical characteristics of high and low states at various concentrations. The k_{12}
is the transition rate from a low state to a high state; k_{21} is the transition rate from a high state to
a low state. **(E)** Simple schematic diagram of the potential landscape topography under different
pheromone dosages. ① stands for the low state, ② stands for the high state; k_{12} and k_{21} are the
transition rates between the low state and the high state, respectively; h_1 and h_2 represent the
respective barrier heights of the low and high states, respectively.

870
Fig. 5. Quantification of the deformation of yeast cells during polar growth. (A) A simple
diagram of cell shapes with circle filling patterns. The image on the left depicts a yeast cell
cultured for 1,020 min in a medium containing 0.7 μM pheromone; the red line indicates the
contour of the cell; the image on the right is the filling model for the image on the left; $R_1 - R_4$
are the radii of the circle. **(B)** Real-time trajectory of the cell morphology (H_n) at 0.7 μM. **(C)**
The distribution statistics of the cell growth rate at 0.7 μM. The red dashed line serves as the
dividing line between positive and negative data. WF and LS represent the lateral-fast and
lateral-slow rates, respectively; LF and WS represent the longitudinal-fast and longitudinal-slow

rates, respectively. **(D)** Schematic illustration of the molecular mechanism underlying the formation of the four growth rates. Model W indicates that the polar growth pathway P_1 is not yet connected, so only the P_2 pathway operates; Model L depicts the cooperative operation of polar growth pathways P_1 and P_2. Models F and S describe the polar growth patterns of the two growth forces, which correspond to the high and low states of Fus3, respectively. **(E)** Dual fluorescent protein system in yeast. The three images on the left, from top to bottom, depict bright field cells, cells excited at 488 nm, and cells excited at 561 nm. The white circle within the cell represents the nucleus boundary; S65T and yomCherry are the fluorescent proteins that were linked to *CDC24* and *FUS3*, respectively. **(F)** Cross-correlation of two levels of gene expression at distinct positions. Gene_in and gene_out represent the gene expression levels inside and outside the nucleus, respectively. **(G)** Statistical graph of the distribution statistics of the absolute value of the cell growth rate at 0.7 μM. **(H)** The proportion of high-state and low-state data present in high and low states at various pheromone concentrations.

Fig. 6. The interpretation of the different cell morphological fates. (**A**) Statistical distribution of the cell morphology at 0.7 μM. The red dashed lines roughly correspond to the boundary between distinct cell morphological fates; F_1–F_4 represent the four cell morphological fates. (**B**) Photographs taken with a fluorescence microscope of cells exhibiting four distinct morphological fates in response to varying pheromone concentrations. (**C**) The synergistic effect of both lateral and longitudinal cell growth capabilities at a pheromone concentration of 0.7 μM. The x-axis represents the ability of the cells to grow longitudinally, as indicated by the rate of change in the sum of the radii of the filled circles within the cells, i.e., $(R_1 + R_2 + \cdots + R_n)'$; the y-axis represents the ability of cells to grow laterally, as indicated by the rate of change in the average radius of the filled circles within the cells, i.e., $(\frac{R_1 + R_2 + \cdots + R_n}{n})'$. (**D**) Changes in the average cell length as a function of pheromone concentration. (**E**) The sum of the net fluxes among the four cell morphologies at various pheromone concentrations. (**F**) The scheme for the simulated cell growth; a represents the length of the cell; Δa is the increased length of the cell; b represents the width of the cell; and Δb is the increased width of the cell; the scheme of the H_n calculation; c is the remainder of the cell length divided by b; N represents the number of b. (**G**) Distribution graph of the negative log of the value of the Fus3 fluorescence intensity inside and outside the nucleus of yeast cells in the stationary phase using simulation. (**H**) The distribution of the cell morphology (H_n) using simulation.

912 **Table 1. Correlation coefficients of fluorescence trajectories.** The "Pearson Correlation
913 Coefficient" between the intranuclear and extranuclear fluorescence intensity trajectories at
914 various pheromone concentrations.

Pheromone doses (μM)	0.7	0.8	1.0	2.0	3.0
Correlation (inside & outside)	0.8321	0.8647	0.8446	0.7752	0.8879

916 **Table 2. Physical characterization of gene expression landscapes.** The transition probability,
917 transition rate, residence time, and barrier height between the high and low states at various
918 pheromone concentrations.

Pheromone doses (μM)		0.7	0.8	1.0	2.0	3.0
Transition probability $P = \begin{pmatrix} P_{11} & P_{12} \\ P_{21} & P_{22} \end{pmatrix}$	P_{11}	0.69	0.74	0.5369	0.5502	0.6143
	P_{12}	0.31	0.26	0.4631	0.4498	0.3857
	P_{21}	0.3711	0.4251	0.5053	0.445	0.3686
	P_{22}	0.6289	0.6289	0.6289	0.6289	0.628
Transition rate	k_{12}	0.0623	0.0717	0.1803	0.112	0.0687
	k_{21}	0.052	0.0439	0.1652	0.1132	0.0718
Residence time (min)	low	357.43	270	691.5	947.98	750.74
	high	498.17	941.84	539.63	301.44	212.3
Barrier height	low	0.1131	0.0997	0.1323	0.2175	0.1716
	high	0.1371	0.1657	0.1125	0.0285	0.0451

Table 3. The proportional distribution of high state (H_1, H_2) and low state (L_1, L_2) in two growth rates. "FH" denotes the "High state" in the "Fast growth rate"; "FL" denotes the "Low state" in the "Fast growth rate"; "SH" denotes the "High state" in the "Slow growth rate"; "SL" denotes the "Low state" in the "Slow growth rate"; "$FH + FL$" and "$SH + SL$" represent fast and slow growth rate data volumes.

Pheromone doses (μM)	$\dfrac{FH}{(FH+FL)}$	$\dfrac{FL}{(FH+FL)}$	$(FH+FL)$	$\dfrac{SH}{(SH+SL)}$	$\dfrac{SL}{(SH+SL)}$	$(SH+SL)$
0.7	0.6847	0.3153	13888	0.1433	0.8567	7447
0.8	0.8500	0.1500	5027	0.2879	0.7121	13381
1.0	0.6782	0.3218	1157	0.2872	0.7128	21884
2.0	0.7266	0.2734	2249	0.1353	0.8647	34637
3.0	0.9530	0.0470	1679	0.0244	0.9756	16597

Table 4. The Average length of yeast cells at different pheromone doses.

Pheromone doses (μM)	0.7	0.8	1.0	2.0	3.0
Average length	31.98	32.27	32.99	33.99	25.01