Obox4 secures zygotic genome activation upon loss of *Dux*

Youjia Guo[#], Tomohiro Kitano[#], Kensaku Murano, Ten D. Li, Akihiko Sakashita, Hirotsugu Ishizu, Masayuki Sato, Haruhiko Siomi*

Department of Molecular Biology, Keio University School of Medicine, Tokyo 1608582, Japan.

[#]These authors contributed equally: Youjia Guo and Tomohiro Kitano.

*Corresponding author:
Haruhiko Siomi
awa403@keio.jp

Abstract

Once fertilized, mouse zygotes rapidly proceed to zygotic genome activation (ZGA), during which long terminal repeats (LTRs) of murine endogenous retroviruses with leucine tRNA primer (MERVL) are potently activated by a conserved homeodomain-containing pioneer factor, DUX. However, *Dux*-knockout embryos produce fertile mice, suggesting that ZGA is redundantly driven by unknown factor(s). In the present study, we report that the multicopy mouse homeobox gene, *Obox4*, encodes a transcription factor that redundantly drives ZGA. OBOX4 is highly expressed in mouse 2-cell embryos. Single knockdown of *Obox4* or *Dux* is tolerated by embryogenesis, whereas *Obox4/Dux* double knockdown completely compromises embryonic development. Genome-wide epigenetic profiling revealed that OBOX4 binds to MERVL LTRs as well as murine endogenous retroviruses with lysine tRNA primer (MERVK) LTRs, and mediates the deposition of active histone modifications. Our study identified OBOX4 as a pioneer factor that provides genetic redundancy to pre-implantation development.

Introduction

The mechanism by which zygotes acquire totipotency is a major question in Developmental Biology. Following fertilization, the zygote must build a *de novo* conceptus with a transcriptionally quiescent genome. The genome rapidly undergoes zygotic genome activation (ZGA), during which epigenetic reprogramming and expression of nascent transcripts enforce the replacement of parental infrastructures by their zygotic counterparts [1]. Fulfillment of metazoan ZGA results in blastomeres, a collection of cells that are totipotent enough to reflect their potential to individually produce both embryos and extraembryonic appendages [2,3]. Totipotency lasts until the morula stage, where the first cell fate decision takes place, following which it is remolded into pluripotency in cells destined for the embryonic lineage [4], to which embryonic stem cells (ESCs) and induced pluripotent stem cells correspond. In placental mammals, ZGA is characterized by the massive reactivation of transposable elements (TEs) and epigenome remodeling, predominantly implemented by a

collection of ZGA genes that have adapted long terminal repeats (LTRs) of endogenous retroviruses (ERVs) as stage-specific *cis*-regulatory elements [5].

Homeodomains are DNA-binding amino acid motifs encoded by a class of conserved genomic sequences termed homeoboxes [6-8]. Homeodomains are prevalent transcription factors that regulate development because of their ability to bind chromatin in a sequence-specific manner [9-11]. Among all homeobox-containing genes, Paired-like (PRD-like) homeobox class genes are particularly associated with ZGA [12]. Studies have shown that the PRD-like homeobox gene, double homeobox (*Dux*), plays a critical role in pre-implantation embryogenesis [13,14]. *Dux* is conserved throughout placentalia and is highly expressed during ZGA. In humans and mice, DUX specifically binds and activates endogenous retroviruses with leucine tRNA primer (ERVL) LTR-derived promoters, resulting in the expression of downstream ZGA genes [15-17]. *Dux* expression induces a 2-cell-embryo-like (2C-like) state in mouse embryonic stem cells (mESCs) [15-17], whereas *Dux*-knockout (KO) undermines mouse blastocyst formation [15]. These observations have led to the assumption that *Dux* resides at the top of the eutherian ZGA regulatory hierarchy [16]. However, subsequent studies have revealed that mouse embryogenesis is compatible with the loss of *Dux*, suggesting a redundant pathway of ZGA that is controlled by uncharacterized pioneer factor(s) [18-21].

In this study, we sought to identify the redundant pioneer factor that drives ZGA in the absence of *Dux*. We discovered that the multicopy homeobox gene, oocyte-specific homeobox 4 (*Obox4*), is essential for *Dux*-less ZGA. *Obox4* is abundantly expressed in mouse 2C-embryos, 2C-like mESCs, and totipotent blastomere-like cells (TBLCs). *Obox4* and *Dux* are collectively, but not respectively, essential for ZGA. Mechanistically, OBOX4 promotes ZGA by binding to the LTRs of murine endogenous retroviruses with leucine tRNA primer (MERVL) and murine endogenous retroviruses with lysine tRNA primer (MERVK), and thereby guiding the deposition of active epigenetic modifications. Taken together, our findings substantiate a pre-implantation development model, in which the initiation of ZGA is redundantly secured by *Dux* and *Obox4*.

Results

Expression profiling identifies pioneer factor candidates

Genetic redundancy is commonly observed among genes that share sequence homology [22]. Because *Dux* is a homeobox gene, we speculated that its redundant factors are homeobox genes. We examined the expression profiles of all mouse homeobox genes during pre-implantation embryogenesis using published single-cell RNA-seq (scRNA-seq) data [23]. Clustering analysis defined a collection of 45 homeobox genes, whose transcript levels reached a maximum before the late 2C stage (Figure 1A). We then sought to identify *bona fide* ZGA factors from this collection by looking for genes whose transcripts were of zygotic origin, instead of ones that were parentally inherited. TBLCs have recently been established *as in vitro* counterparts of 2C-embryos, which are derived from splicing-inhibited mESCs, and hence, are free of parentally inherited transcripts [24]. Analysis of published RNA sequencing (RNA-seq) data showed that the expression of four homeobox genes, *Duxf3*, *Emx2*, *Hoxd13*, and *Obox4*, was ten times higher in TBLCs than in ESCs (Figure 1B). We examined whether the expression of these genes was affected in the *Dux* knockout embryos. Using published RNA-seq data [18,20], we

found that the expression of all single-copy candidate genes and a subset of the multicopy gene *Obox4* was unaffected in *Dux* knockout embryos (Figure 1C).

Obox4 potently activates 2C-genes and TEs

Next, we examined whether these candidate genes promoted ZGA gene expression. Mouse ZGA is characterized by the surging activity of genes that have co-opted MERVL LTRs as stage-specific promoters [25]. We constructed a 2C::tdTomato reporter mESC line with transgenic tdTomato red fluorescence protein driven by MERVL 5'-LTR. [26]. Candidate genes were cloned and overexpressed in the reporter cell line (Figures 1D and S1A). Interestingly, ectopic expression of *Obox4* markedly induced tdTomato expression, similar to that induced by *Dux* (Figures 1E, 1F, and S1B). To characterize the impact of *Obox4* expression on the mESC transcriptome, we established an ESC stable line bearing an inducible *Obox4* transgene and performed RNA-seq analysis with or without induced *Obox4* expression (Figure S1C and S1D). Differentially expressed gene analysis revealed that *Obox4* expression is associated with drastic transcriptome changes in mESCs (Figure 1G). Specifically, *Obox4* expression resulted in the upregulation of genes that were specifically expressed in early-to-middle 2C-embryos (Figure 1H). Typical 2C-genes including *Zscan4*, *Zfp352*, *Dppa3*, and *Sp110*, as well as 2C-specific TEs, MERVL and MERVK, were highly activated upon *Obox4* expression (Figure 1I and 1J). These results indicated that *Obox4* is a potent inducer of the 2C-like state in mESCs.

Obox4/Dux double knockdown compromises pre-implantation development

We then examined whether *Obox4* was crucial for ZGA, particularly in the absence of *Dux*. First, we confirmed the presence of endogenous OBOX4 protein, as the *Obox4* loci are marked as pseudogenes in the current genome annotation GRCm39 [27]. We generated mouse monoclonal antibodies that recognized OBOX4 (Figure S2A). The sequence of OBOX4 was significantly divergent from those of other OBOX members (Figure S2B), and monoclonal antibodies were negative for cross-reactivity with OBOX2 (Figure S2C). Immunofluorescence staining with anti-OBOX4 monoclonal antibodies confirmed that endogenous OBOX4 was highly abundant in 2C-embryos (Figure 2A) and 2C-like mESCs (Figure 2B). To determine whether *Obox4* is essential for pre-implantation development, microinjection of antisense oligonucleotide (ASO) into male pronuclei of zygotes was performed to knockdown *Obox4* and *Dux* (Figure 2C). At the transcript level, both *Obox4* and *Dux* knockdown reduced *Obox4* expression in 2C-embryos (Figure 2D). At the protein level, ASO targets were depleted in knockdown 2C-embryos (Figure 2E and 2F), whereas OBOX4 was retained in *Dux* knockdown 2C-embryos (Figure 2G). Monitoring development until 4.5 days post cotium (dpc) showed that *Obox4* single knockdown resulted in moderate developmental retardation, with almost 50% of embryos reaching the blastocyst stage. Similarly, blastocyst formation was preserved in nearly 25% of *Dux* single knockdown embryos, which is consistent with previously reported *Dux* knockdown/knockout experiments [15,18-21]. Intriguingly, the *Obox4/Dux* double knockdown completely disrupted pre-implantation development, with most embryos arrested at the 2C stage (Figure 2H and 2I). These observations showed that the expression of *Obox4* and *Dux* is collectively required for pre-implantation development and strongly suggest that *Obox4* is capable of securing ZGA in a *Dux*-independent manner.

OBOX4 rescues *Obox4/Dux* double knockdown in pre-implantation embryos

We then sought to verify whether OBOX4 is essential for pre-implantation development in the absence of DUX, by determining whether the presence of OBOX4 can rescue the developmental arrest of *Obox4/Dux* double knockdown embryos. First, codon-optimized *Obox4* mRNA without the ASO target motif was produced using *in vitro* transcription. The rescue experiment was performed by means of co-microinjection of *Obox4* mRNA with *Obox4/Dux* double knockdown ASOs (Figure 3A). Restoration of OBOX4 in *Obox4* mRNA-microinjected embryos was confirmed using immunofluorescence staining at the 2C stage (Figure 3B). As expected, development monitoring showed that *Obox4/Dux* double knockdown embryos failed blastocyst formation at 4.5 dpc, whereas among *Obox4* mRNA-rescued embryos, blastocyst formation was retained at a similar level to that observed in the *Dux* single knockdown experiment (Figures 3C, 3D, 2H, and 2I). These results suggested that OBOX4 is responsible for securing development in the absence of DUX. As *Obox4* has been shown to activate 2C-genes and TEs in mESCs, we questioned whether embryonic depletion of OBOX4 impairs ZGA. Consistent with the observation of embryonic development, transcriptome analysis revealed that the ZGA pattern of *Dux/Obox4* double knockdown 2C-embryos was markedly different from that of control embryos, while single knockdown embryos were less affected (Figure 3E). The dysregulated transcriptomes were characterized by extensive hypoactivation of MERVL/MERVK elements (Figure 3F) and 2C-genes (Figure 3G), suggesting that the compromised pre-implantation development was due to ZGA failure.

ic*Obox4* and *Dux* are essential for ESC-to-TBLC transition

In 2021, it was reported that administration of the splicing inhibitor pladienolide B (PlaB) to ESC culture drives pluripotent-to-totipotent transition, which results in TBLCs that are functionally and molecularly comparable with totipotent blastomeres [24]. Since it remains unclear whether the ESC-to-TBLC transition relies on pioneer factor activity, we attempted to knockout *Obox4* and *Dux* in mESCs, to examine whether they contribute to *in vitro* totipotency acquisition in TBLCs. Following a previously published protocol [18], plasmids encoding Cas9 and single guide RNA (sgRNA) flanking the *Dux* cluster were introduced into wild-type (WT) mESCs, to knockout the *Dux* loci (Figure S3A). Plasmids encoding Cas9 and sgRNAs flanking the *Obox4* cluster, and sgRNAs targeting the *Obox4* internal sequence were then introduced into *Dux* knockout mESCs and WT mESCs, to knockout the *Obox4* loci (Figure S3B). Bi-allelic deletions of the *Dux* cluster and *Dux* internal sequences were confirmed in the monoclonized knockout cell lines (Figure S3C and S3D). Deletion of the *Obox4* cluster was confirmed in the monoclonized cell lines (Figure S3E). Knockout mESCs were treated with PlaB to enforce TBLC transition and then examined for the depletion of OBOX4 at the protein level. Immunostaining and western blotting revealed OBOX4 depletion in *Obox4/Dux* double knockout and *Obox4* single knockout cell lines (Figure S3F and S3G). The cell lines were subjected to transcriptome profiling for further validation (Figure 4A). RNA-seq analysis showed that both *Obox4* and *Dux* were highly expressed in WT TBLC, while a single knockout of either gene resulted in inactivation of the other (Figure 4B), which was not observed in embryo knockdown experiments (Figure 2D), suggesting that *Obox4* and *Dux* are individually required for TBLC transition. Indeed, transcriptome analysis showed that *Dux* and *Obox4* single knockout was sufficient to hinder the activation of TBLC-specific genes and TEs, while double knockout led to the dysregulation of the TBLC transcriptome to a greater extent (Figure 4C and 4D). Consistent with the

observation in embryos, the expression of critical 2C-genes was abolished in PlaB-treated *Dux/Obox4* double knockout mESCs (Figure 4E).

Obox4 binds to 2C-gene promoters and LTR elements

Obox4 contains a homeobox, activates 2C-genes and TEs, and is essential for *Dux*-less pre-implantation embryogenesis, which prompted us to examine whether OBOX4 is a pioneer factor that activates ZGA-associated loci through direct binding with *cis*-regulatory elements. Cleavage under targets and release using nuclease sequencing (CUT&RUN-seq) is an alternative method to chromatin immunoprecipitation sequencing (ChIP-seq) that profiles protein-DNA interactions. This technique combines antibody-targeted cleavage of proximal DNA with high-throughput sequencing, to identify the binding sites of DNA-associated proteins [28]. To profile genome-wide OBOX4 binding, we performed CUT&RUN-seq against triple FLAG-tagged OBOX4 (3×FLAG-OBOX4) expressed in mESCs, using a high-affinity anti-FLAG antibody [29] (Figure 5A). CUT&RUN-seq yielded ~11,000 peaks that were shared among biological triplicates over the control samples (q<0.05). Peak annotation revealed that OBOX4 was enriched in the promoter region and LTR elements (Figure 5B). As expected, we found that most genes (3304 of 5172, 63.9%) whose promoters were occupied by OBOX4 were highly expressed in 2C-embryos. Unlike DUX, which strictly binds to a conserved motif exclusively found in MERVL LTR [17], OBOX4 displayed a broad binding spectrum that was strongly biased towards both LTRs of MERVK and MERVL (Figure 5C). Consistently, *de novo* motif discovery failed to identify a consensus OBOX4 binding motif shared among the MERVK and MERVL elements, while clear patterns were discovered in each LTR element family (Figure 5D). Hence, we hypothesized that OBOX4 binds to the MERVL and MERVK elements, to promote downstream gene expression. To test this hypothesis, we constructed tdTomato reporters driven by LTRs of MERVL or MERVK and co-transfected them with *Dux* and *Obox4* into *Dux/Obox4* double knockout ESCs (Figure 5E). Fluorescence-activated cell sorting (FACS) analysis of the tdTomato-positive population showed that *Obox4* activated both MERVK and MERVL LTRs, whereas *Dux* only activated MERVL LTR (Figure 5F). As OBOX4 and DUX share the MERVL LTR as binding targets, we reasoned that OBOX4 and DUX are mutually redundant factors that activate core totipotency genes by activating their MERVL LTR promoters. Indeed, by comparing OBOX4 CUT&RUN-seq and previously published DUX ChIP-seq data [16], we found that OBOX4 targets more than half (1,089 out of 2,068) of the DUX target genes (Figure 5G), among which are critical totipotency genes, including *Dppa3*, *Sp110*, and *Zscan4d* (Figure 5H).

OBOX4 guides deposition of active histone marks

Pioneer factors have been proposed to modulate the epigenetic landscape at their binding loci [30]. We analyzed previously published ChIP-seq data [31,32] and found that OBOX4 target loci were enriched with the active histone marks H3K4me3 and H3K27ac (Figure 6A) in 2C-embryos. To examine whether OBOX4 mediates the deposition of active histone modifications, we performed CUT&RUN-seq in WT and *Obox4* knockout mESCs and TBLCs (Figure 6B). Differential analysis of histone modification revealed that upon PlaB treatment, *Obox4* knockout mESCs manifested extensive loss of H3K4me3, but not H3K27me3, at the OBOX4 target MERVL and MERVK loci (Figure 6C and 6D). We then investigated whether the MERVL and MERVK loci were discriminated in terms of histone modifications. Interestingly, most of the strongest OBOX4-associated

H3K27ac loci were deposited at the ERVK, but not ERVL loci (Figure 6E). As H3K27ac is highly associated with
enhancer activity, together with our finding that *Obox4* activates MERVK LTR and is associated with intense
H3K27ac deposition, we speculated that despite being functionally redundant to *Dux*, *Obox4* also acts as an
activator of ZGA-specific ERVK enhancers. This hypothesis is supported by the observation that in addition to
H3K4me3 enrichment at promoter regions, H3K27ac is also deposited at OBOX4 target ERVKs that are
proximal to critical ZGA-genes (Figure 6F).
197

Discussion

Starting with a transcriptionally inert genome, the eutherian ZGA engages in massive yet coordinated
expression of genes driven by LTR TEs. In mice, it has been reported that the pioneer factor DUX accesses and
opens condensed chromatin of the MERVL LTR loci, which results in the activation of downstream ZGA genes
[15-17]. However, the establishment of fertile *Dux* knockout mice suggests that ZGA is redundantly fulfilled by
unknown pioneer factor(s) [18-21]. In the present study, we found that *Obox4*, a cryptic multi-copy cluster gene,
encodes a homeodomain-containing protein that potently induces ZGA gene expression. Remarkably,
Dux/Obox4 double knockdown completely compromised pre-implantation development, which tolerated
single knockdown of either gene. Double knockdown embryos exhibited severely dysregulated transcriptomes,
characterized by MERVL and ZGA gene activation failure. We then focused on characterizing the molecular
mechanisms underlying the biological significance of *Obox4* as well as its relevance to *Dux*. Multi-omics
analysis revealed that OBOX4 directly binds to MERVL (the target of DUX) and MERVK loci, to mediate active
histone mark deposition. We demonstrated that *Obox4* activates MERVL and MERVK in a *Dux*-independent
manner and that the *Obox4*-targeted MERVK elements are enriched with the active enhancer mark H3K27Ac
in TBLCs. In summary, our results highlight that OBOX4 is a pioneer factor that is functionally redundant to
DUX during ZGA.
The human PRD-like homeobox gene, TPRX2, is highly expressed during ZGA [14]. Ectopic expression of
TPRX2 in human embryonic stem cells (hESCs) and fibroblasts alters the expression of genes related to
totipotency and pluripotency [14,33]. *Obox* is a divergent ortholog of *TPRX2* in mice [33]. Obox transcripts were first
discovered in gonads [34], and were later found to be highly abundant in mouse 2C-embryos [35]. The *Obox* family
has 67 members clustered in the sub-telomeric region of mouse chromosome 7 and is further divided into six
subfamilies: *Obox1*, *Obox2*, *Obox3*, *Obox4*, *Obox5*, and *Obox6*, with *Obox4* constituting nearly 60% (39 of 67)
of the family population [36,37]. Although *Obox4* dominates the *Obox* family population, members of the entire
subfamily have been annotated as pseudogenes and have largely been neglected from investigation. Despite
being abundantly transcribed in the gonads, the biological significance of *Obox* genes is poorly understood.
The role of *Obox4* is particularly mysterious, as it remains to be determined whether *Obox4* is a functional
protein-coding gene. Transcription of *Obox4* was originally reported to be testis-specific and contradicted by
the detection of *Obox4* transcripts in the ovaries and oocytes [38]. *Obox4* is involved in oocyte maturation and
ESC differentiation [39-43]. However, these observations are supported by limited evidence, with no evidence at
the protein level. By detecting OBOX4 in various totipotent cell types using high-quality monoclonal antibodies,
we provided decisive evidence that *Obox4* is a functional protein-coding gene. Our data demonstrated that

OBOX4 binds to genomic loci of MERVL-derived promoters and MERVK-derived enhancers and mediates their activation during ZGA. However, whether and how other *Obox* family members contribute to ZGA remains unclear.

Genetic redundancy is widespread in higher organisms and typically arises in signaling networks, in which multiple functionally overlapping factors commit to a shared teleological objective, to counteract sporadic mutations or defects [44,45]. This principle is of particular importance to embryogenesis, a process that gives rise to a reproducing organism. Indeed, genetic redundancy in pluripotency has been evidenced by the observation that naïve pluripotency can be elicited and maintained through various "independent inputs that operate through both unique and convergent targets" [46]. Although the molecular pathway that governs totipotency has been largely restricted to the scope that centers on *Dux*, it is reasonable to assume that the acquisition of totipotency also complies with the rule of redundancy. This assumption was validated by the generation of fertile *Dux* knockout mice; however, the panorama of ZGA redundancy remained hidden. Our work on *Obox4* sheds light on this topic by discovering that *Obox4* serves as a redundant gene to *Dux* during ZGA and is a bivalent activator of MERVL and MERVK elements. Although the function of MERVL has been extensively investigated, particularly during early embryogenesis, the role of MERVK remains unclear. Interestingly, recent studies have suggested that MERVK serves as a meiosis-specific enhancer of spermatogenesis [47]. Mammalian Y chromosomes have undergone rapid evolution and have been intensively inserted by TEs [48]. As *Obox4* is highly expressed in the testes, it is worth investigating whether *Obox4* plays a regulatory role during spermatogenesis, by activating MERVK-derived enhancers.

Unlike *Dux*, which is structurally and functionally conserved throughout placentalia [49], the ancestral locus of the *Obox* family appears to have undergone mouse-specific duplication and generated a gene cluster that is collectively syntenic to the *Tprx2* locus in other mammals [33]. Despite the distal homology between *Obox4* and *Tprx2*, it has been reported that human *TPRX2* is expressed in 8-cell embryos [14] and that ectopic expression of *TPRX2* in hESCs affects the expression of totipotency- and pluripotency-related genes [14,33]. Whether *TPRX2* plays a similar role in humans to *Obox4* in mice remains to be elucidated. In addition, *Obox4* and *Dux* are zygotically expressed, denying their role as ZGA initiators. How they are activated after fertilization and whether they share the same activation mechanism remain elusive. Both *Dux* and *Obox4* are tandem repeat genes that cluster in either sub-telomeric or centromeric regions and form heterochromatin in most cell lineages. These observations have led to the hypothesis that pioneer factors, such as *Obox4* and *Dux*, are passively activated by biased epigenetic remodeling towards their loci.

Figure Legends

Figure 1. *Obox4* promotes transcription of 2C-genes and transposable elements.

(A) Mouse homeobox genes are specifically expressed during ZGA. The genes were identified by means of statistically determined *k*-means clustering based on their expression in pre-implantation embryos. *Dux* is shown in a bold italic font.

(B) Dot plot showing per-gene normalized read counts in mouse totipotent TBLCs *versus* ESCs. Genes with more than 10-fold normalized read counts (FC=10) have been highlighted.

(C) Bar graph displaying gene expression in *Dux* knockout 2C-embryos *versus* WT 2C-embryos. Loss of *Dux*
partially deactivates the multicopy homeobox gene *Obox4*.
(D) Diagram of the 2C::tdTomato reporter assay. ESCs bearing the tdTomato expression cassette under the
control of the MERVL promoter showed red fluorescence upon entering the 2C-like state. The expression
of the pioneer factor increased the 2C-like population, as detected using FACS. An EGFP expression
plasmid was co-transfected with a gene of interest, to normalize the transfection efficiency.
(E) Boxplot showing normalized 2C-like cell percentage in 2C::tdTomato reporter ESCs overexpressing
candidate pioneer factors. *Dux* and *Obox4* potently induced a 2C-like state.
(F) Fluorescence microscope view of 2C::tdTomato ESCs transfected with empty plasmid, plasmid encoding
DUX, or plasmid encoding OBOX4. Plasmids encoding EGFP were co-transfected to normalize the
transfection efficiency in FACS analysis.
(G) Volcano plot of DEGs in *Obox4*-overexpressing mESCs. Representative ZGA genes are labeled with gene
symbols.
(H) Expression profile of genes up-regulated by *Obox4* during embryogenesis.
(I) Volcano plot of differentially expressed transposable elements in *Obox4*-overexpressing ESCs. ERVL and
ERVK were activated upon *Obox4* induction.
(J) Expression profile of ERVL and ERVK elements during pre-implantation embryogenesis.
284
Figure 2. OBOX4 is necessary for DUX-less preimplantation development.
(A) Immunofluorescence staining of histone H3, OBOX4, and MERVL GAG at different pre-implantation
embryo stages.
(B) Immunofluorescence staining of DUX, OBOX4, and OCT4 in 2C-like ESCs.
(C) Schematic representation of *Dux* and *Obox4* knockdown experiments in pre-implantation embryos. Male
pronuclei of zygotes were microinjected with ASO targeting *Dux* or *Obox4* transcripts. The embryos were
monitored for development until 4.5 dpc, or subjected to RNA-seq at the middle 2C stage.
(D) Transcript levels of *Dux* and *Obox4* in ASO-injected embryos, as measured using RNA-seq. CPM, counts per
million
(E) Immunofluorescence staining of OBOX4 in embryos microinjected with *Obox4* ASO.
(F) Immunofluorescence staining of DUX in embryos microinjected with *Dux* ASO.
(G) Immunofluorescence staining of OBOX4 in embryos microinjected with *Dux* ASO.
(H) Development monitoring of KD embryos at 1.5 dpc and 4.5 dpc.
(I) The percentages of embryonic stages observed at 1.5 dpc, 2.5 dpc, 3.5 dpc, and 4.5 dpc. The plot represents
the sum of three independent experiments, with 150–200 embryos per condition.
300
301 **Figure 3. *Obox4/Dux* double knockdown embryos manifest a dysregulated transcriptome.**
302 (A) Schematic of the double knockdown rescue experiment. Male pronuclei of zygotes were injected with ASO
303   targeting *Obox4* and *Dux* transcripts as well as *in vitro*-transcribed codon-optimized *Obox4* mRNA.
304   Embryos were monitored for development until 4.5 dpc.

(B) Immunofluorescence staining of OBOX4 in embryos microinjected with scrambled ASO, double ASO (*Obox4* and *Dux*), or double ASO with codon-optimized *Obox4* mRNA.

(C) Development monitoring of knockdown and rescue embryos at 1.5 dpc and 4.5 dpc.

(D) The percentages of embryonic stages observed at 1.5 dpc, 2.5 dpc, 3.5 dpc, and 4.5 dpc. The plot represents the sum of three independent experiments, with 80–100 embryos per condition.

(E) Heatmap of expression level of 2C-genes in ASO-injected embryos.

(F) Volcano plot showing the results of DEG analysis of 2C-genes and TEs in knockdown embryos, as compared to that in scramble ASO-injected embryos. Standard deviations of $\log_2$(fold-change) were used to represent the degree of transcriptome dysregulation. Stdev, standard deviation

(G) Bar chart showing the differential expression of representative 2C-genes in ASO-injected embryos; n=3 biological replicates.

Figure 4. *Obox4* is necessary for TBLC formation.

(A) Schematic diagram of *Obox4* and *Dux* knockout using CRISPR-Cas9 in mESCs. Previously reported sgRNAs flanking the *Dux* cluster were introduced into mESCs with SpCas9, to knockout *Dux*. sgRNAs flanking the *Obox4* cluster as well as the homeobox sequence of individual *Obox4* were introduced into mESCs with SpCas9, to knockout *Obox4*. WT and knockout mESCs were treated with PlaB, to induce TBLC formation. RNA-seq was performed on untreated and PlaB-treated mESCs, to profile transcriptome changes.

(B) Bar charts showing transcript levels of *Obox4* and *Dux* in mESCs and TBLCs are described in (A). WT, wild-type; mean $\pm$ SD (n=3 biological replicates).

(C) Heatmap of TBLC gene expression levels in normally cultured mESCs and PlaB-treated mESCs.

(D) Volcano plot showing differential expression of TBLC genes and TEs in PlaB-treated knockout mESCs, as compared to that in WT TBLCs. Standard deviations of all $\log_2$ values (fold change) were used to represent the degree of transcriptome dysregulation. Stdev, standard deviation

(E) Bar chart showing differential expression of representative 2C-genes (upper panel) and pluripotency genes (lower panel) in PlaB-treated knockout mESCs, as compared to that in WT TBLCs; n=3 biological replicates.

Figure 5. OBOX4 binds to LTR elements and activates 2C-genes.

(A) Schematic diagram of the CUT&RUN-seq workflow using ESCs with induced expression of FLAG-tagged OBOX4. ESCs bearing a doxycycline-inducible 3×FLAG-OBOX4 expression cassette were induced with doxycycline for 24 h. OBOX4-associated DNA was pulled down in the CUT&RUN assay, using a high-affinity anti-FLAG antibody.

(B) Pie-chart displaying proportions of annotated features in the OBOX4 binding sites.

(C) Heatmap showing the OBOX4 binding site distribution near 2C-gene promoters and LTR elements. OBOX4 binding is highly condensed in the 2C-gene TSS.

(D) The OBOX4 binding motif was enriched *de novo* with the MEME-ChIP algorithm, using the top 500 CUT&RUN-seq peaks discovered at the MERVK loci or all peaks discovered at the MERVL loci. The stacked letter suggested divergence of the motif, which matched 339 out of 500 peaks at the MERVK and 126 out of

138 peaks at the MERVL loci. The histogram given in the figure displays the distribution of conserved

motifs within the 500 nt windows flanking the peak summits.

(E) Schematic design of the reporter assay. Plasmids bearing a tdTomato ORF downstream of the ERVL or

ERVK LTR were co-transfected with *Dux* or *Obox4* expression plasmids. Activation of LTR elements

resulted in an increased red fluorescence-positive mESC population, which was measured using FACS.

EGFP expression plasmid was co-transfected into the culture, following which the green

fluorescence-positive population was measured, to normalize the transfection efficiency.

(F) Box plot showing the percentage of red fluorescence-positive mESCs upon expression of *Dux* or *Obox4*.

Both *Dux* and *Obox4* can activate ERVL, whereas only *Obox4* activates ERVK.

(G) Euler diagram showing the overlap of DUX and OBOX4 target genes. Selected 2C-genes are listed on the

right.

(H) Genomic track view showing DUX and OBOX4 binding sites at the *Sp110, Dppa3*, and *Zscan4d* loci and

their expression levels upon *Obox4* overexpression. Reads were converted to CPM for normalization of the

sequencing depth. The OBOX4 binding sites overlapped with those of DUX. *Sp110, Dppa3*, and *Zscan4d*

were highly expressed upon overexpression of *Obox4*.

Figure 6. OBOX4 guides deposition of active histone marks at its binding loci.

(A) Heatmap showing the distributions of H3K4me3, H3K27Ac, and H3K27me3 near the OBOX4 binding sites

in 2C-embryos.

(B) Schematic diagram showing the strategy for profiling histone modifications in PlaB-treated mESCs.

PlaB-treated or untreated WT and *Obox4* knockout mESCs were examined using CUT&RUN-seq, with

antibodies against H3K4me3, H3K27Ac, and H3K27me3. Loci subjected to TBLC and *Obox4*-associated

histone modifications were identified by comparing the signal intensities between PlaB-treated and

non-treated mESCs.

(C) Change in H3K4me3 modification intensity at OBOX4-binding ERV loci in PlaB-treated mESCs, as

compared to that in PlaB-untreated mESCs.

(D) Change in H3K27me3 modification intensity at OBOX4-binding ERV loci in PlaB-treated mESCs, as

compared to that in PlaB-untreated mESCs.

(E) Change in H3K27Ac modification intensity at OBOX4-binding ERV loci in PlaB-treated mESCs, as compared

to that in PlaB-untreated mESCs.

(F) Genomic track view of histone modification levels at *Dppa3, Zscan4c*, and selected ERVK loci in

PlaB-treated WT and *Obox4* knockout mESCs. Reads were converted to CPM, for normalization of the

sequencing depth. OBOX4 binding is associated with the deposition of active histone modifications

H3K4me3 and H3K27Ac, but not with the repressive histone modification H3K27me3.

Supplemental figure legends

Figure S1

(A) Western blot showing protein products of the candidate genes ectopically expressed in mESCs.

(B) FACS analysis of 2C::tdTomato cells co-transfected with candidate gene expression plasmids and EGFP.

(C) Schematic of *Obox4*-inducible cell line construction.

(D) Western blot showing 3×FLAG-OBOX4 in *Obox4*-inducible cell line, induced by different concentrations of

doxycycline. Expression of 3×FLAG-OBOX4 was carried out in a dose-dependent manner.

Figure S2

(A) Schematic of the production of monoclonal antibodies. Mice were immunized with the GST-OBOX4 fusion

protein. After fusing B-cells with myelomas, the antibodies produced by hybridomas were examined

against MBP-OBOX4 fusion protein, followed by MBP-OBOX2 fusion protein, for identification of

OBOX4-specific antibodies.

(B) Analysis of protein sequence similarity among the OBOX family members. The Smith–Waterman local

alignment method was used to align the amino acid sequences, and the divergence among different

sequences was represented in the form of a block substitution matrix.

(C) ELISA signal of anti-OBOX4 monoclonal antibodies against the MBP-OBOX4 and MBP-OBOX2 fusion

proteins. Anti-OBOX4 monoclonal antibodies were not reactive to MBP-OBOX2.

Figure S3

(A) Schematic of the strategy for knocking out the *Dux* cluster (not drawn to scale). *Dux* cluster was removed

by means of CRISPR-Cas9 targeting of the flanking regions. PCR-based genotyping, with primers targeting

the knockout allele and *Dux* internal sequence, was used to validate the genomic removal of *Dux*.

(B) Schematic of the strategy for knocking out the *Obox4* cluster (not drawn to scale). The *Obox4* cluster was

removed by means of CRISPR-Cas9 targeting of the flanking regions as well as the internal regions of

individual *Obox4* copies. PCR-based genotyping, with primers targeting the knockout allele, was used to

validate the genomic removal of the cluster.

(C) Agarose gel image of PCR-based genotyping of the *Dux* knockout allele in founder cell lines. The red arrow

indicates a cell line bearing a bi-allelic deletion of the *Dux* cluster.

(D) Agarose gel image of PCR-based genotyping of the *Dux* internal sequences in founder cell lines. The red

arrow indicates a cell line without the genomic *Dux* sequence.

(E) Agarose gel image of PCR-based genotyping of the *Obox4* knockout allele in founder cell lines. The red

arrow indicates a cell line bearing a bi-allelic deletion of the *Obox4* cluster.

(F) Immunofluorescence analysis of OBOX4 in *Obox4* knockout, *Dux* knockout, and double knockout cell lines.

(G) Western blot of OBOX4 and DUX in *Obox4* knockout, *Dux* knockout, and double knockout cell lines.

Methods

Mice

418   Male/female BDF1 and female BALB/c mice were purchased from Japan SLC Inc. Mice were fed regular

419   chow and housed in a controlled room under a 14 h/10 h light/dark cycle, at 22°C. All animal experiments

were approved by the Animal Care and Use Committee of Keio University and conducted in compliance with the Keio University Code of Research Ethics.

Cell culture

All mouse mESC lines were cultured in Dulbecco's Modified Eagle Medium (DMEM) (catalog no. 08488-55, Nacalai Tesque) supplemented with 10% fetal bovine serum (lot no. S10581S1820, Biowest), 1× GlutaMAX™ (catalog no. 35050061, Gibco), 1× penicillin-streptomycin (catalog no. 15140163, Gibco), 1× non-essential amino acids (catalog no. 11140050, Gibco), 1× sodium pyruvate (catalog no. S8636, Merck), 50 μM 2-mercaptoethanol (catalog no. 21985023, Gibco), 1 μM CHIR99021 (catalog no. 034-23103, Wako), 3 μM PD0325901 (catalog no. 168-25293, Wako), and mouse leukemia inhibitory factor produced in-house. The culture medium was supplemented with iMatrix-511 silk (recombinant human laminin-511 E8 fragment) (catalog no. 892021, Matrixome) at a concentration of 125 $ng.cm^{-2}$ culture vessel area, before seeding mESCs into it. The cells were cultured in a humidified atmosphere containing 5% CO_2 at 37°C, with the media changed every 48 h.

TBLCs were generated by culturing mESCs in an ordinary medium supplemented with 2.5 nM PLaB (catalog no. 16538, Cayman Chemicals). Cells were sub-cultured every 2–4 d, at a seeding density of $1×10^5$ cm^{-2} culture vessel area. After five rounds of sub-culture, the cells were deemed TBLCs.

SP2/O-Ag14 myeloma and primary clones of hybridomas were cultured in GIT medium (catalog no. 63725715, Wako) supplemented with recombinant human interleukin-6 (IL-6) (catalog no. 20006, PeproTech). Cells were cultured in a humidified atmosphere containing 5% CO_2 at 37°C and sub-cultured every day, at a seeding density of $3×10^5$ mL^{-1}.

For monoclonal antibody production, hybridomas were cultured in hybridoma serum-free medium supplemented with IL-6. The cells were cultured in a humidified atmosphere containing 5% CO_2 at 37°C, until they were over-confluent.

Cell transfection

Plasmid transfections were performed using jetOPTIMUS® DNA Transfection Reagent (catalog no. 101000025, Polyplus), according to the manufacturer's instructions. Briefly, mESCs were seeded at a cell density of $8×10^4$ cm^{-2} culture vessel area. After 20 min, the jetOPTIMUS® reagent and plasmids were diluted in jetOPTIMUS® buffer, incubated for 10 min at room temperature, and then applied to the cell culture. After 48 h, the cells were collected for downstream experiments.

cDNA synthesis and cloning

Total RNA was extracted from TBLCs using ISOGEN (catalog no. 311-02501, Nippon Gene), according to the manufacturer's instructions. Briefly, 500 μL ISOGEN was added to $1.5×10^6$ freshly harvested TBLCs. After 5 min of incubation at room temperature, 100 μL of chloroform (catalog no. 03802606, Wako) was added to the cells and mixed by means of vigorous shaking. After 2 min of incubation at room temperature, the samples were centrifuged at $12,000 × g$ for 15 min at 4°C. The upper aqueous phase was then transferred to a new tube and mixed with 240 μL isopropanol (catalog no. 15-2320, Merck), to precipitate the RNA. After 5 min of

incubation at room temperature, the samples were centrifuged at 12,000 × g for 15 min at 4°C. The
supernatant was then discarded, following which the RNA pellets were washed with 70% ethanol (catalog no.
057-00456, Wako). The RNA pellets were then air-dried and dissolved in RNase-free water. The RNA solution
was subjected to DNase treatment using TURBO DNA-*free*™ kit (catalog no. AM1907, Thermo Fisher),
according to the manufacturer's instructions, to remove genomic DNA carryover. RNA (1 µg) was reverse
transcribed to cDNA using a Transcriptor First Strand cDNA Synthesis Kit (catalog no. 04379012001, Roche),
according to the manufacturer's instructions. PrimeSTAR® Max DNA Polymerase (catalog no. R045A, TaKaRa)
and ProFlex™ PCR System (catalog no. 4484073, Thermo Fisher) were used to amplify the sequence of interest
from the cDNA. NEBuilder® HiFi DNA Assembly Kit (catalog no. E2621L, NEB) was used to clone the sequence
of interest into plasmid backbone.

Generation of transgenic mESC lines
The 2C::tdTomato reporter cell line was generated by transfection of EB3 mESCs (catalog no. AES0139,
RIKEN BRC Cell Bank) with a linearized 2C::tdTomato reporter plasmid (catalog no. 40281, Addgene). After 48
h, the cells were subjected to 500 µg.mL^{-1} hygromycin (catalog no. 08906151, Wako) for 7 d. The selected cells
were then seeded at a density of 2×10^2 cm^{-2} in culture medium containing 250 µg.mL^{-1} hygromycin. Single-cell
clones were picked and expanded after 7 d.
A tetracycline-controlled transcription activation system and a piggyBac transposon system were used
as previously described [50], to generate an *Obox4*-inducible cell line. Briefly, pPB-TRE-3×FLAG-Obox4,
pPB-CAG-rtTA3G-IRES-Hygro, and pCMV-HyPBase-PGK-Puro plasmids were co-transfected into EB3 mESCs.
After 48 h, the cells were subjected to selection with 500 µg.mL^{-1} hygromycin and 500 µg.mL^{-1} G418 (catalog
no. 07805961, Wako), for 7 d. The selected cells were then seeded at a density of 2×10^2 cm^{-2} in a culture
medium containing 250 µg.mL^{-1} hygromycin and 250 µg.mL^{-1} G418. Single-cell clones were picked and
expanded after 7 d.

Generation of knockout mESC lines
The *Dux* knockout cell line was generated by means of co-transfection of EB3 mESCs with
pX330-DuxKO-A and pX330-DuxKO-B plasmids. After 48 h, the cells were seeded at a very low density.
Single-cell clones were picked and expanded after 7 d and then subjected to genotyping.
The *Obox4* knockout cell line was generated by means of co-transfection of EB3 mESCs with the
pX330-Puro-Obox4KO-3, pX330-Puro-Obox4KO-4, pX330-Puro-Obox4KO-6, and pX330-Puro-Obox4KO-7
plasmids. After 18 h, the cells were subjected to selection with 1 µg.mL^{-1} puromycin (catalog no. 16023151,
Wako) for 54 h. The selected cells were then seeded at a very low density in culture medium without
antibiotics. Single-cell clones were picked and expanded after 7 d, and then subjected to genotyping.

Flow cytometry
The cells were digested with 0.25% trypsin-EDTA (catalog no. 25200072, Gibco), at 37°C for 5 min, and
then re-suspended in FluoroBrite™ DMEM (catalog no. A1896701, Gibco). The cell suspension was filtered

through a 35 μM cell strainer (catalog no. 352235, Falcon). The samples were analyzed using a cell sorter (catalog no. SH800Z, Sony). Data analysis was performed using the Sony SH800Z cell sorter software.

Western blot

Freshly harvested cells were re-suspended in phosphate-buffered saline (PBS) (catalog no. 14249-95, Nacalai Tesque) and lysed by means of sonication. Whole cell lysates were mixed with sample buffer containing reducing reagent (catalog no. 09499-14, Nacalai Tesque) and boiled at 95°C for 5 min. Protein samples were loaded on a 10% tris-glycine gel, run in AllView PAGE Buffer® (catalog no. DS520, BioDynamics Laboratory), and then transferred to nitrocellulose membranes (catalog no. 10600003, Cytiva) using a Power Blotter System (catalog no. PB0012, Thermo Fisher). The membranes were blocked with PBST containing 2% skim milk (catalog no. 4902720131292, Morinaga), at room temperature for 15 min. Membranes were incubated with primary antibodies in PBST containing 2% skim milk, at room temperature for 30 min. After three washes with PBST, the membrane was incubated with secondary antibodies in PBST containing 2% skim milk, at room temperature for 15 min, with shaking. After three washes with PBST, the membrane was incubated with ECL reagents (catalog no. RPN2232, Cytiva) and exposed to an X-ray film (catalog no. 28906839, Cytiva) or detected using a digital chemiluminescence imager (catalog no. 17001402JA, Bio-Rad).

Generation of monoclonal antibodies

The anti-OBOX4, anti-DUX, and anti-GAG monoclonal antibodies were produced as previously described [51]. Briefly, BALB/c mice were immunized by means of intraperitoneal injection of glutathione S-transferase-fused protein-of-interest (PoI). The mice were routinely immunized until their sera tested positive for the PoI-fused maltose binding protein (MPB), after which two boosting immunizations were performed. Following boosting, splenocytes from immunized mice were fused with SP2/O myeloma and cultured in hypoxanthine-aminopterin-thymidine medium for 10 d, to select for hybridomas. Hybridomas were subsequently screened using enzyme-linked immunosorbent assay against PoI-fused MBP. The positive polyclonal hybridomas were monoclonalized and expanded. For anti-OBOX4 hybridomas, additional screening using OBOX2-fused MBP was conducted. The culture supernatants of monoclonal hybridomas were sterilized by passing them through a 0.22 μM pore size filter (catalog no. 430769, Corning) and used directly as an antibody solution in other assays.

***In vitro* transcription**

The *Obox4* coding sequence was codon-optimized using GeneArt Instant Designer (Thermo Fisher), to remove the ASO target motif and improve translation efficiency. Codon-optimized DNA was synthesized using Prime Gene Synthesis Services (Thermo Fisher). Codon-optimized *Obox4* mRNA was transcribed *in vitro* using the mMESSAGE mMACHINE™ T7 Transcription Kit (catalog no. AM1344, Thermo Fisher) and then polyadenylated using a poly(A) tailing kit (catalog no. AM1350, Thermo Fisher), according to the manufacturer's instructions. The polyadenylated mRNA was purified using the RNeasy Mini Kit (catalog no. 74004, Qiagen), according to the manufacturer's instructions, and dissolved in RNase-free water, at a concentration of 100 ng.μL^{-1}.

Pronuclear injection of mouse embryos

Eight-week-old BDF1 female mice were injected with 150 µL of CARD HyperOva (catalog no. KYD-010-EX, Kyudo Co., Ltd.). After 48 hours, the female mice were injected with 5IU human chorionic gonadotropin (hCG) (catalog no. GONATROPIN, ASKA Animal Health Co. Ltd.) and mated with BDF1 male mice. After 22 h, the PN3 zygotes were collected. Cumulus cells were removed by briefly culturing the zygotes in potassium simplex optimization medium (KSOM) (catalog no. MR-101-D, Merck) supplemented with 0.3 µg.µL^{-1} hyaluronidase (catalog no. H4272, Merck). Embryos were cultured in KSOM medium drops covered with liquid paraffin (catalog no. 26137-85, Nacalai Tesque), in a humidified atmosphere with 5% CO_2, at 37°C. After 4 h, PBS containing either 40 µM scramble ASO, 20 µM anti-*Dux* ASO, 20 µM anti-*Obox4* ASO, 40 µM equimolar anti-*Dux*/anti-*Obox4* ASO mixture, or 40 µM equimolar anti-*Dux*/anti-*Obox4* ASO mixture with 100 ng.µL^{-1} codon-optimized *Obox4* mRNA was microinjected into the male pronuclei of zygotes using a microinjector (catalog no. 5252000013, Eppendorf). For developmental monitoring, embryos were cultured for 4 d after microinjection and assessed for developmental stage at 18 h, 42 h, 66 h, and 90 h after microinjection, which corresponded to 1.5 dpc, 2.5 dpc, 3.5 dpc, and 4.5 dpc, respectively.

Immunofluorescence

For embryos, middle-2C-embryos were collected at 1.5 dpc. The zona pellucida was removed by treating the embryos with acidic Tyrode's solution (catalog no. MR-0040-D, Merck). The embryos were fixed in 4% paraformaldehyde (catalog no. 09154-14, Nacalai Tesque) in PBS for 10 min. Following three washes in PBS, fixed embryos were permeabilized with 0.1% Triton™ X-100 (catalog no. 1610407, Bio-Rad) in PBS, for 20 min at room temperature. Following three washes in PBS, permeabilized embryos were blocked in 2% bovine serum albumin (BSA; catalog no. 011-27055, Wako) in PBS, for 20 min at room temperature. The blocked embryos were incubated with primary antibodies diluted in 2% BSA in PBS, at 4°C overnight. Following three washes in PBS, the embryos were incubated with 2% BSA in PBS containing 1:200 diluted 4',6-diamidino-2-phenylindole (DAPI) (catalog no. 19178-91, Nacalai Tesque) and 1:500 diluted Alexa Fluor™-conjugated anti-mouse IgG1 secondary antibody (catalog no. A-21127, Thermo Fisher) or Alexa Fluor™-conjugated anti-mouse IgG2a secondary antibody (catalog no. A-21131, Thermo Fisher) and, for 1 h at room temperature. Following three washes in PBS, the embryos were transferred into liquid paraffin-covered PBS drops on a glass-bottom dish (catalog no. D11130H, Matsunami).

For culturing, cells seeded on glass bottom chamber slides (catalog no. SCS-N02, Matsunami) were fixed with 4% paraformaldehyde in PBS, for 10 min. After three washes in PBS, the fixed cells were permeabilized with 0.1% Triton™ X-100 in PBS, for 15 min at room temperature. Following three washes in PBS, permeabilized cells were incubated with primary antibodies diluted with 2% skim milk in PBS, for 30 min at room temperature. Following three washes in PBS, the cells were incubated with 2% BSA in PBS containing 1:500 diluted Alexa Fluor™-conjugated secondary antibody and 1:200 diluted DAPI, for 1 h at room temperature. Following three washes in PBS, the chamber was removed and the slides were mounted with ProLong™ Glass Antifade Mountant (catalog no. P36982, Thermo Fisher). Fluorescence images were taken using a confocal laser scanning microscope (catalog no. FV3000, Olympus).

scRNA-seq analysis

Raw scRNA-seq data was obtained from the dataset of Deng et al. (GSE45719). Quality control and adapter trimming was done using fastp [52] v0.23.2. Quality-controlled reads were aligned to the GRCm38.p6 reference genome using STAR [53] v2.7.9a, with default arguments. Reads were counted against GRCm38.p6 comprehensive gene annotation [54] and mm10 repeats from the University of California, Santa Cruz (UCSC) RepeatMasker using Subread [55] v2.0.1 featureCounts function, and multi-mapping reads were discarded for non-TE features and counted fractionally for TEs. Seurat [56] v4.1.0 was used to process the read counts of scRNA-seq. Cells with greater than 7.5% mitochondrial reads or less than 14,000 annotated features were discarded. Expression levels were log-normalized. The expression profiles of homeobox genes listed in HomeoDB2 [57] were clustered into 11 k-means after z-score transformation.

RNA-seq

For embryos, 30 middle-2C-embryos were collected 20 h after microinjection in each independent biological replicate. The zona pellucida was removed by treating embryos with acidic Tyrode's solution. Libraries were constructed using the SMART-Seq® Stranded Kit (catalog no. 634442, TaKaRa) and indexed using the SMARTer® RNA Unique Dual Index Kit (catalog no. 634451, TaKaRa), according to the manufacturer's instructions.

For cell culture, total RNA was prepared using the RNeasy Kit, according to the manufacturer's instructions. The RNA solution was subjected to DNase treatment to remove genomic DNA carryover. Using 1 µg total RNA per sample, libraries were constructed with the NEBNext® Ultra™ II Directional RNA Library Prep Kit for Illumina (catalog no. E7760L, NEB) and indexed with NEBNext® Multiplex Oligos for Illumina (catalog no. E6440S, NEB), according to the manufacturer's instructions.

The libraries were quantified with a 2100 Bioanalyzer (catalog no. G2939BA, Agilent) using a High Sensitivity DNA Kit (catalog no. 5067-4626, Agilent). Quantified libraries were pooled and sequenced using an Illumina NovaSeq 6000 System in 150 bp paired-end mode (Illumina). Base calling and de-multiplexing were performed using the bcl2fastq2 (Illumina) v2.20. De-multiplexed reads were quality-controlled, aligned, and counted, as described above. DESeq2 [58] v1.32.0 was used to perform differential expression analysis. Raw RNA-seq reads for TBLCs were downloaded from the dataset of Shen et al. (GSE168728). Raw RNA-seq reads of *Dux* knockout embryos were downloaded from the datasets of Chen & Zhang and De Iaco et al. (GSE121746 and GSE141321 respectively). The published data were analyzed using the same method.

CUT&RUN-seq

Cultured cells (1×10^5) freshly prepared by trypsin-EDTA digestion were used to perform CUT&RUN with the CUT&RUN Assay Kit (catalog no. 86652, Cell Signaling Technology), according to the manufacturer's instructions. Enriched DNA was subjected to library construction using the NEBNext® Ultra™ II DNA Library Prep Kit for Illumina (catalog no. E7645L, NEB), with the adapter ligation step performed at 50℃ instead of 65℃, to prevent the denaturation of small DNA inserts. The libraries were indexed using NEBNext® Multiplex Oligos for Illumina, according to the manufacturer's instructions.

The libraries were quantified, pooled, sequenced, de-multiplexed, and quality controlled, as described above. Processed reads were aligned to the GRCm38.p6 reference genome using Bowtie2 [59] v2.4.1, with default settings. Peaks were called in each biological replicate using the MACS3 [60] v3.0.0a6 callpeak function (-f BAMPE). Alignment tracks were first generated using deepTools [61] v3.5.1 bamCoverage function (--binSize 10 --normalized using CPM --smoothLength 30), and then normalized by subtracting the signal from non-immune IgG using the deepTools bamCompare function (--scaleFactorssMethod None --operation subtract --binSize 10 --smoothLength 30). ChIPseeker [62] v1.28.3 was used to annotate the peaks. The MEME suite [63] v5.4.1 was used to identify the binding motifs. Quantitative histone modification analysis of replicated samples was performed using MAnorm2 [64] v1.2.0. Visualization of genomic tracks was performed using trackplot [65] v1.3.10. Raw ChIP-seq reads of DUXs in mESCs were downloaded from Hendrickson et al. (GSE85632). The published data were analyzed using the same method.

Quantification and statistical analysis

Descriptive and comparative statistics were employed in the manuscript as described in the figure legends with the number of replicates indicated. Significance is defined as a *p-value* less than 0.05 indicated with asterisk (* *p-value* < 0.05, ** *p-value* < 0.01, *** *p-value* < 0.001). Error bar represents the standard deviation (SD) of the mean of the replicates.

Data and code availability

The RNA-seq and CUT&RUN-seq data generated in this study have been deposited at NCBI Gene Expression Omnibus (GEO) database under the accession code GSE196671.

Acknowledgements

We thank all the members of the Siomi Laboratory for their discussions and comments on this work. We thank Takehiko Yokomizo (Department of Biochemistry, Juntendo University) for providing the anti-DYKDDDDK monoclonal antibody (clone 2H8). We also thank Daisuke Motooka (Research Institute for Microbial Diseases, Osaka University) for generating the sequencing data. We are grateful to Azusa Inoue (Center for Integrative Medical Sciences, RIKEN) and Katsuhiko Hayashi (Department of Genome Biology, Osaka University) for their comments on this manuscript. This work was supported by the MEXT Grant-in-Aid for Scientific Research in Innovative Areas (19H05753 to H.S.), AMED project to elucidate and control mechanisms of aging and longevity (1005442 to H.S.), JSPS Grant-in-Aid for Scientific Research KAKENHI (20K21507 to K.M.), Mochida Memorial Foundation Research Grant to K.M., Sumitomo Foundation Research Grant to K.M., Keio University Doctorate Student Grant-in-Aid Program to Y.G., and the JST Doctoral Program Student Support Fellowship to Y.G.

Author contributions

H.S. and Y.G. conceived and designed the project; Y.G., T.K., K.M., T.D.L., A.S., H.I., and M.S. performed the experiments; Y.G. and T.K. analyzed the data; H.I. developed the ASO-mediated knockdown experiments; H.S. and K.M. supervised and coordinated the experiments; Y.G. and H.S. wrote the paper with input from all authors. All authors reviewed and approved the final version of the manuscript.

Declaration of interests

The authors declare no competing interests.

References

1.   Lee, M.T., Bonneau, A.R. & Giraldez, A.J. Zygotic genome activation during the maternal-to-zygotic transition. *Annual review of cell and developmental biology* **30**, 581-613 (2014).

2.   Tarkowski, A.K. Experiments on the development of isolated blastomeres of mouse eggs. *Nature* **184**, 1286-1287 (1959).

3.   Tarkowski, A.K. Experimental studies on regulation in the development of isolated blastomeres of mouse eggs; Badania eksperymentalne nad rozwojem izolowanych blastomerów jaj myszy. *Acta Theriologica* **3**, 191-267 (1959).

4.   Suwińska, A., Czołowska, R., Ożdżeński, W. & Tarkowski, A.K. Blastomeres of the mouse embryo lose totipotency after the fifth cleavage division: expression of Cdx2 and Oct4 and developmental potential of inner and outer blastomeres of 16-and 32-cell embryos. *Developmental biology* **322**, 133-144 (2008).

5.   Svoboda, P. *et al.* RNAi and expression of retrotransposons MuERV-L and IAP in preimplantation mouse embryos. *Developmental biology* **269**, 276-285 (2004).

6.   McGinnis, W., Levine, M.S., Hafen, E., Kuroiwa, A. & Gehring, W.J. A conserved DNA sequence in homoeotic genes of the Drosophila Antennapedia and bithorax complexes. *Nature* **308**, 428-433 (1984).

7.   Carrasco, A.E., McGinnis, W., Gehring, W.J. & De Robertis, E.M. Cloning of an X. laevis gene expressed during early embryogenesis coding for a peptide region homologous to Drosophila homeotic genes. *Cell* **37**, 409-414 (1984).

8.   McGinnis, W., Hart, C.P., Gehring, W.J. & Ruddle, F.H. Molecular cloning and chromosome mapping of a mouse DNA sequence homologous to homeotic genes of drosophila. *Cell* **38**, 675-680 (1984).

9.   Scott, M.P. & Weiner, A.J. Structural relationships among genes that control development: sequence homology between the Antennapedia, Ultrabithorax, and fushi tarazu loci of Drosophila. *Proceedings of the National Academy of Sciences* **81**, 4115-4119 (1984).

10.   Gehring, W.J. & Hiromi, Y. Homeotic genes and the homeobox. *Annual review of genetics* **20**, 147-173 (1986).

11.   Gehring, W.J. Homeo boxes in the study of development. *Science* **236**, 1245-1252 (1987).

12.   Lewin, T.D., Royall, A.H. & Holland, P.W. Dynamic Molecular Evolution of Mammalian Homeobox Genes: Duplication, Loss, Divergence and Gene Conversion Sculpt PRD Class Repertoires. *Journal of Molecular Evolution*, 1-19 (2021).

13.   Töhönen, V. *et al.* Novel PRD-like homeodomain transcription factors and retrotransposon elements in early human development. *Nature communications* **6**, 8207 (2015).

14.   Madissoon, E. *et al.* Characterization and target genes of nine human PRD-like homeobox domain genes expressed exclusively in early embryos. *Scientific reports* **6**, 1-14 (2016).

15.   De Iaco, A. *et al.* DUX-family transcription factors regulate zygotic genome activation in placental mammals.
Nature genetics **49**, 941-945 (2017).

16.   Hendrickson, P.G. *et al.* Conserved roles of mouse DUX and human DUX4 in activating cleavage-stage genes
and MERVL/HERVL retrotransposons. *Nature genetics* **49**, 925-934 (2017).

17.   Whiddon, J.L., Langford, A.T., Wong, C.-J., Zhong, J.W. & Tapscott, S.J. Conservation and innovation in the
DUX4-family gene network. *Nature genetics* **49**, 935-940 (2017).

18.   Chen, Z. & Zhang, Y. Loss of DUX causes minor defects in zygotic genome activation and is compatible with
mouse development. *Nature genetics* **51**, 947-951 (2019).

19.   Guo, M. *et al.* Precise temporal regulation of Dux is important for embryo development. *Cell research* **29**,
956-959 (2019).

20.   De Iaco, A., Verp, S., Offner, S., Grun, D. & Trono, D. DUX is a non-essential synchronizer of zygotic genome
activation. *Development* **147**, dev177725 (2020).

21.   Bosnakovski, D., Gearhart, M.D., Ho Choi, S. & Kyba, M. Dux facilitates post-implantation development, but is
not essential for zygotic genome activation. *Biology of reproduction* **104**, 83-93 (2021).

22.   Wagner, A. Genetic redundancy caused by gene duplications and its evolution in networks of transcriptional
regulators. *Biological cybernetics* **74**, 557-567 (1996).

23.   Deng, Q., Ramsköld, D., Reinius, B. & Sandberg, R. Single-cell RNA-seq reveals dynamic, random monoallelic
gene expression in mammalian cells. *Science* **343**, 193-196 (2014).

24.   Shen, H. *et al.* Mouse totipotent stem cells captured and maintained through spliceosomal repression. *Cell*
184, 2843-2859. e20 (2021).

25.   Macfarlan, T.S. *et al.* Endogenous retroviruses and neighboring genes are coordinately repressed by
LSD1/KDM1A. *Genes & development* **25**, 594-607 (2011).

26.   Macfarlan, T.S. *et al.* Embryonic stem cell potency fluctuates with endogenous retrovirus activity. *Nature*
487, 57-63 (2012).

27.   Howe, K.L. *et al.* Ensembl 2021. *Nucleic Acids Research* **49**, D884-D891 (2020).

28.   Skene, P.J. & Henikoff, S. An efficient targeted nuclease strategy for high-resolution mapping of DNA binding
sites. *Elife* **6**, e21856 (2017).

29.   Sasaki, F. *et al.* A high-affinity monoclonal antibody against the FLAG tag useful for G-protein-coupled
receptor study. *Analytical biochemistry* **425**, 157-165 (2012).

30.   Mayran, A. & Drouin, J. Pioneer transcription factors shape the epigenetic landscape. *Journal of Biological*
Chemistry **293**, 13795-13804 (2018).

31.   Liu, X. *et al.* Distinct features of H3K4me3 and H3K27me3 chromatin domains in pre-implantation embryos.
Nature **537**, 558-562 (2016).

32.   Dahl, J.A. *et al.* Broad histone H3K4me3 domains in mouse oocytes modulate maternal-to-zygotic transition.
Nature **537**, 548-552 (2016).

33.   Maeso, I. *et al.* Evolutionary origin and functional divergence of totipotent cell homeobox genes in eutherian
mammals. *BMC biology* **14**, 1-14 (2016).

34.   Rajkovic, A., Yan, C., Yan, W., Klysik, M. & Matzuk, M.M. Obox, a family of homeobox genes preferentially
expressed in germ cells. *Genomics* **79**, 711-717 (2002).

35.   Ge, S.X. Exploratory bioinformatics investigation reveals importance of "junk" DNA in early embryo
development. *BMC genomics* **18**, 1-19 (2017).

36.   Zhong, Y.-f. & Holland, P.W. The dynamics of vertebrate homeobox gene evolution: gain and loss of genes in
mouse and human lineages. *BMC evolutionary biology* **11**, 1-13 (2011).

37.   Wilming, L.G., Boychenko, V. & Harrow, J.L. Comprehensive comparative homeobox gene annotation in
736   human and mouse. *Database* **2015**(2015).
737   38.   Lee, H.-S. & Lee, K.-A. Characterization and functional analysis of Obox4 during oocyte maturation by RNA
738   interference. *Clinical and Experimental Reproductive Medicine* **34**, 293-303 (2007).
739   39.   Lee, H.S. *et al.* Obox4 critically regulates cAMP-dependent meiotic arrest and MI-MII transition in oocytes.
740   *The FASEB Journal* **24**, 2314-2324 (2010).
741   40.   Lee, K.-A., Lee, H.-S., Kim, E.-Y., Kim, K.-H. & Kim, Y. Oocyte-specific homeobox 4 (Obox4) is a key regulatory
742   component of the cAMP-dependent meiotic arrest during in vitro oocyte maturation. *Fertility and Sterility*
743   **90**, S108 (2008).
744   41.   Lee, H.-S., Kim, E.-Y. & Lee, K.-A. Changes in gene expression associated with oocyte meiosis after Obox4
745   RNAi. *Clinical and experimental reproductive medicine* **38**, 68 (2011).
746   42.   Lee, H.-S. *et al.* Obox4-silencing-activated STAT3 and MPF/MAPK signaling accelerate nuclear membrane
747   breakdown in mouse oocytes. *Reproduction* **151**, 369-378 (2016).
748   43.   Kim, H.-M. *et al.* Obox4 regulates the expression of histone family genes and promotes differentiation of
749   mouse embryonic stem cells. *FEBS letters* **584**, 605-611 (2010).
750   44.   Nowak, M.A., Boerlijst, M.C., Cooke, J. & Smith, J.M. Evolution of genetic redundancy. *Nature* **388**, 167-171
751   (1997).
752   45.   Vavouri, T., Semple, J.I. & Lehner, B. Widespread conservation of genetic redundancy during a billion years
753   of eukaryotic evolution. *Trends in genetics* **24**, 485-488 (2008).
754   46.   Hackett, J.A. & Surani, M.A. Regulatory principles of pluripotency: from the ground state up. *Cell stem cell* **15**,
755   416-430 (2014).
756   47.   Sakashita, A. *et al.* Endogenous retroviruses drive species-specific germline transcriptomes in mammals.
757   *Nature Structural & Molecular Biology* **27**, 967-977 (2020).
758   48.   Bachtrog, D. Y-chromosome evolution: emerging insights into processes of Y-chromosome degeneration.
759   *Nature Reviews Genetics* **14**, 113-124 (2013).
760   49.   Leidenroth, A. & Hewitt, J.E. A family history of DUX4: phylogenetic analysis of DUXA, B, C and Duxbl reveals
761   the ancestral DUX gene. *BMC evolutionary biology* **10**, 1-13 (2010).
762   50.   Takeuchi, C., Murano, K., Ishikawa, M., Okano, H. & Iwasaki, W., Y. Generation of stable drosophila ovarian
763   somatic cell lines using piggyBac system. *Methods in Molecular Biology* **In press**(2022).
764   51.   Guo, Y. *et al.* Potent mouse monoclonal antibodies that block SARS-CoV-2 infection. *Journal of Biological*
765   *Chemistry* **296**(2021).
766   52.   Chen, S., Zhou, Y., Chen, Y. & Gu, J. fastp: an ultra-fast all-in-one FASTQ preprocessor. *Bioinformatics* **34**,
767   i884-i890 (2018).
768   53.   Dobin, A. *et al.* STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**, 15-21 (2013).
769   54.   Frankish, A. *et al.* GENCODE 2021. *Nucleic Acids Research* **49**, D916-D923 (2020).
770   55.   Liao, Y., Smyth, G.K. & Shi, W. featureCounts: an efficient general purpose program for assigning sequence
771   reads to genomic features. *Bioinformatics* **30**, 923-930 (2014).
772   56.   Hao, Y. *et al.* Integrated analysis of multimodal single-cell data. *Cell* (2021).
773   57.   Zhong, Y.f. & Holland, P.W. HomeoDB2: functional expansion of a comparative homeobox gene database for
774   evolutionary developmental biology. *Evolution & development* **13**, 567-568 (2011).
775   58.   Love, M.I., Huber, W. & Anders, S. Moderated estimation of fold change and dispersion for RNA-seq data
776   with DESeq2. *Genome biology* **15**, 1-21 (2014).
777   59.   Langmead, B. & Salzberg, S.L. Fast gapped-read alignment with Bowtie 2. *Nature methods* **9**, 357-359
778   (2012).

60.   Zhang, Y. *et al.* Model-based analysis of ChIP-Seq (MACS). *Genome biology* **9**, 1-9 (2008).

61.   Ramírez, F. *et al.* deepTools2: a next generation web server for deep-sequencing data analysis. *Nucleic acids*
research **44**, W160-W165 (2016).

62.   Yu, G., Wang, L.-G. & He, Q.-Y. ChIPseeker: an R/Bioconductor package for ChIP peak annotation, comparison
and visualization. *Bioinformatics* **31**, 2382-2383 (2015).

63.   Bailey, T.L. *et al.* MEME SUITE: tools for motif discovery and searching. *Nucleic acids research* **37**,
W202-W208 (2009).

64.   Tu, S. *et al.* MAnorm2 for quantitatively comparing groups of ChIP-seq samples. *Genome Research* **31**,
131-145 (2021).

65.   Pohl, A. & Beato, M. bwtool: a tool for bigWig files. *Bioinformatics* **30**, 1618-1619 (2014).

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure S1

Figure S2

Figure S3