Distinguishing between canonical and non-canonical tRNA genes reveals that Thermococcaceae adhere to the standard archaeal tRNA gene set

Peter T.S. van der Gulik[1]*, Martijn Egas[2], Ken Kraaijeveld[3], Nina Dombrowski[4], Astrid T. Groot[5], Anja Spang[4], Wouter D. Hoff[6]*, Jenna Gallie[7]*

[1] Centrum Wiskunde & Informatica, Amsterdam, 1090 GB, The Netherlands
[2] Department of Evolutionary and Population Biology, Institute for Biodiversity and Ecosystem Dynamics, University of Amsterdam, Amsterdam, 1090 GE, The Netherlands
[3] Leiden Centre for Applied Bioscience, University of Applied Sciences Leiden, Leiden, 2333CR, The Netherlands
[4] Department of Marine Microbiology and Biogeochemistry, NIOZ, Royal Netherlands Institute for Sea Research, Den Burg, 1790 AB, The Netherlands
[5] Institute for Biodiversity and Ecosystem Dynamics (IBED), University of Amsterdam, Amsterdam, 1090 GE, The Netherlands
[6] Department of Microbiology and Molecular Genetics, Oklahoma State University, Stillwater, Oklahoma, 74078, USA
[7] Department of Evolutionary Theory, Max Planck Institute for Evolutionary Biology, Plön, 24306, Germany

* To whom correspondence should be addressed. Tel: +31 20592 4232 Email: Peter.van.der.Gulik@cwi.nl
Correspondence may also be addressed to Jenna Gallie (gallie@evolbio.mpg.de) or Wouter Hoff (wouter.hoff@okstate.edu)

ABSTRACT

Automated genome annotation is an essential tool for extracting biological information from sequence data. The identification and annotation of tRNA genes is frequently performed by the software package tRNAscan-SE, the output of which is listed – for selected genomes – in the Genomic tRNA database (GtRNAdb). Given the central role of tRNA in molecular biology, the accuracy and proper application of tRNAscan-SE is important for both interpretation of the output, and continued improvement of the software. Here, we report a manual annotation of the predicted tRNA gene sets for 20 complete genomes from the archaeal taxon Thermococcaceae. According to GtRNAdb, these 20 genomes contain a number of putative deviations from the standard set of canonical tRNA genes in Archaea. However, manual annotation reveals that only one represents a true divergence; the other instances are either (i) non-canonical tRNA genes resulting from the integration of horizontally transferred genetic elements, or CRISPR-Cas activity, or (ii) attributable to errors in the input DNA sequence. To distinguish between canonical and non-canonical archaeal tRNA genes, we recommend using a combination of automated pseudogene detection by tRNAscan-SE and the tRNAscan-SE isotype score, greatly reducing manual annotation efforts and leading to improved predictions of tRNA gene sets in Archaea.

INTRODUCTION

Canonical transfer RNAs (tRNAs) implement the genetic code by coupling mRNA codons to their respective amino acids during protein synthesis. Of the 64 codons in the genetic code, three generally serve as stop codons. The remaining 61 ('sense') codons are decoded into 20 amino acids by tRNAs with various complementary anticodons. Given that some anticodons can recognize more than one codon – through wobble base pairing (1, 2) – organisms carry fewer than 61 types of tRNA (3). Together with knowledge of wobble base pairing rules, systematic studies of tRNA gene sets have yielded important generalities regarding tRNA requirements for efficient translation; standard tRNA gene sets have been identified for each of the three domains of life (4, 5). In Archaea – the focus of this work – a standard set of 46 tRNA types, each encoded by a single gene copy, has been proposed (examples depicted in Figure 1A). Standard tRNA gene sets are poised to play a pivotal role in understanding tRNA gene set function and evolution. However, a range of reported deviations from these sets (*e.g.*, (3, 6)) may be masking their role in tRNA biology. Specifically, two types of tRNA gene set deviations can occur: (i) evolutionary adaptations of the translationally active tRNA gene set in some organisms (see below), and (ii) artifactual deviations due to imperfections in genome sequencing/assembly, or downstream computational analyses of putative tRNA genes. This manuscript aims to distinguish between these two cases, using the tRNA gene sets carried by the archaeal Thermococcaceae family as an example.

As a result of remarkable progress in experimental methods for determining the genome sequences, automated annotation has become an essential tool for extracting biological information from sequence data. In the field of tRNA biology, several tRNA gene prediction tools are available (7–13), the most widely used of which is tRNAscan-SE (7, 14, 15). tRNAscan-SE effectively performs the challenging tasks of identifying putative tRNA genes from input DNA sequence, and annotating each predicted gene – including the definition of the anticodon and amino acid carried (*i.e.*, the tRNA isotype). Additionally, tRNAscan-SE distinguishes between three isotypes carrying the anticodon 5'-CAU-3': (i) initiator tRNA-Met(CAU), which serves in the initiation stage of translation, (ii) elongator tRNA-Met(CAU), which functions in peptide chain elongation, and (iii) tRNA-Ile(CAU), in which the anticodon is post-transcriptionally modified to recognize isoleucine codon 5'-AUA-3' (16).

The final output of tRNAscan-SE is a biochemically detailed, automatically annotated list of putative tRNA genes from the input DNA sequence (*e.g.*, genome assembly, plasmid). The tRNAscan-SE output for a selected set of complete (or near-complete) genomes is collected in the publicly accessible database, the Genomic tRNA database (GtRNAdb) (3, 17). The accumulation and comparison of complete tRNA gene set data paves the way for conclusions regarding the evolution of tRNA gene sets across the tree of life (5, 18–26). In addition, such information may have important medical implications (27–31). For example, both prokaryotic and eukaryotic tRNA gene sets have been shown to evolve rapidly in response to changing translational demands (32–34), and a correlation has been demonstrated between tRNA gene copy number and bacterial growth rate (35). However, conclusions regarding tRNA gene sets are dependent on the accuracy and proper use of the tRNA detection software.

Here, we evaluate the performance of tRNAscan-SE through detailed manual inspection of

the tRNA genes it identifies and annotates within a set of 20 genomes from the archaeal family

Thermococcaceae. This family is an archaeal clade within the phylum *Euryarchaeota* (according to

the National Center for Biotechnology Information (NCBI) taxonomy (36)), or the phylum

Methanobacteriota (according to the Genome Taxonomy Database, GTDB (37)). The

Thermococcaceae family consists of at least six genera (*i.e.*, *Pyrococcus, Thermococcus*,

Thermococcus A, Thermococcus B, Thermococcus C, and *Palaeococcus*) that are typically found in

hydrothermal marine environments (38, chapter 27). Members of the Thermococcaceae family

reportedly adhere very closely to the standard archaeal tRNA gene set (38, page 364). Given this

information, the proportion of Thermococcaceae genomes listed in GtRNAdb as encoding variant

tRNA gene sets is – as described below – unexpectedly high (seven deviant genomes out of the 20

listed genomes = 35 %). However, manual annotation of these predicted tRNA gene sets resolves

many of the apparent differences, leaving only a single probably truly divergent tRNA gene set.

MATERIALS AND METHODS

20 Thermococcaceae tRNA gene sets. All Thermococcaceae family tRNA gene sets available in

GtRNAdb (Data Release 19; June 2021) (3) were downloaded (see Supplementary Table S5). Details

of the 20 organisms – including members of the *Pyrococcus, Thermococcus, Thermococcus A,*

Thermococcus B, and *Palaeococcus* genera – are provided in Supplementary Table S1. In addition,

the tRNA gene sets of these organisms were predicted from the genome assemblies (downloaded

from NCBI) using locally run tRNAscan-SE (version 2.0.6, the same version that was reportedly used

for the current GtRNAdb listings for all 20 organisms), with standard settings for Archaea (option -A)

and pseudogene detection activated (see Supplementary Table S5). To display the output, options -H

and --detail were added.

Manual annotation of tRNA gene sequences. Putative tRNA genes of interest were manually

annotated using the following six steps: (i) Identify the 3' CCA terminus (nucleotides 74-76), the

discriminator nucleotide (nucleotide 73) and the acceptor stem (nucleotides 1-7, base pairing with

nucleotides 72-66) (39). (ii) Working 3' to 5', identify the T-arm prior to nucleotide 66

(>>>>>UUCRANN<<<<< , where R is base A or G, and each >< indicates a base pair). (iii) Identify

the section between the T-arm and the anticodon-arm (nucleotides 44-45-46-47-48 or 44-45-47-48, or

an extra arm). (iv) Identify the anticodon-arm (>>>>>NNNNNNN<<<<<). With very few exceptions, the

anticodon loop consists of precisely seven nucleotides (N), and very often begins with CU. (v) Identify

the D-arm (>>>xNNNNNNNNx<<<, where x denotes a degree of flexibility in the number of Ns). (vi)

Identify nucleotides 8-9 (between the acceptor stem and D-arm) and 26 (between the D-arm and

anticodon arm), which play a role in realization of the three dimensional structure of tRNA molecules.

Phylogenetic tree construction. The Thermococcaceae family phylogenetic tree in Supplementary Figure S1 was generated by first identifying suitable marker genes in genomes from 20 *Thermococcus* and three *Methanococcaceae* (outgroup) genomes, as described previously (40). Briefly, HMM profiles of an extended TIGRFAM database were queried against a protein database of all 23 archaeal genomes, using hmmsearch v3.1b2 and the following settings: hmmsearch --tblout sequence_results.txt -o results_all.txt --domtblout domain_results.txt --notextw extended_TIGR.hmm All_Genomes.faa. The extended TIGRFAM database can be found at https://zenodo.org/record/3839790#.YjByaVzMI3g. Only hits with an e-value <= 1e-3 were included, and the best hit per protein based on the bit score and e-value was selected. Markers with =>10% of duplicated proteins were excluded, leaving 45 markers for downstream analyses. Note that TIGR00335 and TIGR00483 were excluded from these analyses, because they comprised a large number of paralogues. The final set of 43 marker proteins (listed in Supplementary Table S4) was extracted from the larger protein database and individually aligned using MAFFT LINS-i v7.407 (settings: –reorder) (41) and trimmed using BMGE v1.12 (settings: -t AA -m BLOSUM30 -h 0.55) (42). Single-protein phylogenies were inferred with IQ-TREE v2.1.2 (settings: -m LG -T AUTO -keep-ident --threads-max 2 -bb 1000 -bnni) (43). Trees were manually inspected and all potential paralogues removed. The remaining proteins were realigned, trimmed (as above) and concatenated using a custom perl script; catfasta2phyml.pl (https://github.com/nylander/catfasta2phyml). The phylogenetic tree was inferred based on a final alignment of 12,403 positions using IQ-TREE (v2.1.2, -m LG+C60+F+R -T AUTO --threads-max 80 -bb 1000 -alrt 1000) (43), visualised using FigTree (v1.4.4) and annotated using Adobe Illustrator CC2018.

210 archaeal tRNA gene sets. The tRNA gene sets of 217 Archaea are currently listed in GtRNAdb (Data Release 19; June 2021). For seven of these organisms, the underlying genome assemblies were no longer available from NCBI (*Aciduliprofundum* sp. MAR08-339, *Methanococcus maripaludis* C6, *Methanococcus maripaludis* C7, *Methanococcus voltae* A3, *Methanolinea tarda* NOBI-1, *Methanosarcina barkeri* str. Fusaro, and *Saccharolobus solfataricus* 98/2 (formerly *Sulfolobus solfataricus* 98/2)). The remaining 210 organisms span five phyla and include 59 Crenarchaeota, 138 Euryarchaeota, 1 Korarchaeota, 1 Nanoarchaeota, and 11 Thaumarchaeota. The genomes of these organisms are available on Zenodo (https://zenodo.org/record/6782366#.YsQS5pDP01B). For each of these 210 organisms, the tRNA gene sets and associated scores listed in GtRNAdb were downloaded (resulting in the ~10,000 tRNA genes listed in Supplementary Table S6). In addition, the genome assemblies (including chromosomes and plasmids) used to generate the GtRNAdb entries were downloaded from NCBI, and tRNAscan-SE (version 2.0.6) was locally run on each genome, using standard settings for Archaea (options -A, -H, --detail) and with pseudogene detection activated. In accordance with expectations, ~10,000 tRNA genes were predicted (Supplementary Table S6).

RESULTS

GtRNAdb lists divergent tRNA gene sets for seven of 20 Thermococcaceae genomes

The tRNA gene sets of 20 genomes belonging to the Thermococcaceae family are listed in GtRNAdb (Data Release 19; June 2021) (Supplementary Table S1). Our phylogenetic tree of these 20 genomes confirmed the phylogeny-informed taxonomy of the Thermococcaceae family established by Rinke *et al* (44) and accurately resolved the relationships between the five genera to which these genomes belong, with *Palaeococcus* forming the earliest diverging branch (Supplementary Figure S1).

All 20 of the genomes of interest are complete genomes assembled into single circular contigs, with an average size of 1.9 Mb (ranging from 1.7 Mb to 2.2 Mb) and a mean genomic GC content of 47 % (ranging from 40 % to 56 %). Genome descriptions have been published for 18 of the 20 genomes (see Supplementary Table S1). Inspection of the associated GtRNAdb tRNA gene set entries revealed that, of the 20 genomes, only 13 (65 %) are predicted to carry the standard archaeal tRNA gene set of 46 canonical tRNA types encoded by single-copy genes (see Introduction), while the remaining seven (35 %) are predicted to differ. There are 15 distinct anomalies across the seven deviant genomes, consisting of twelve additional and three absent putative tRNA genes (Table 1; Figure 1A).

When the same set of genomes were analyzed by running tRNAscan-SE locally (version 2.0.6, see Materials and Methods), six of the genes causing tRNA gene set deviations were flagged as possible pseudogenes. Inspection of the documentation for the entries of these genomes on GtRNAdb indicated that automated pseudogene detection was not activated during the associated tRNAscan-SE run, resulting in the inclusion of the six potential pseudogenes within the canonical, standard tRNA gene sets for the organisms involved.

Manual annotation reveals only a single, actually divergent tRNA gene set

Given the unexpectedly high degree of deviation from the standard archaeal tRNA gene set, the 929 tRNA genes predicted across the 20 Thermococcaceae genomes were subjected to careful manual inspection (see Materials and Methods). This revealed that the 15 observed anomalies fall into three categories, each of which is described below (see also Table 1).

Category 1: Erroneously predicted tRNA genes arising from tRNA gene fragments

Eight of the 15 observed anomalies result from the erroneous listing of partial tRNA gene sequences as either standard tRNA genes (seven cases) or genes encoding tRNAs of undetermined isotypes (one case) in GtRNAdb (Table 1). As described further below, we found that the partial tRNA sequences are the result of two processes: (i) the acquisition of fragments of viral tRNA genes by CRISPR-Cas activity (accounting for one anomaly; category 1a in Table 1), and (ii) the partial duplication of tRNA genes during the integration of horizontally transferred genomic elements (accounting for seven anomalies; category 1b in Table 1).

Thermococcus nautili 30-1 (NCBI Reference Sequence NZ_CP007264.1) is predicted to carry two distinct genes encoding Leu-CAA (positions 10,012–10,099 and 706,451–706,668). The first of these has characteristics typical of canonical leucyl-tRNA genes (*e.g.*, 88-bp long, isotype score

134.8), while the second does not (*e.g.*, 218-bp long, very low isotype score of 2.3). Furthermore, the

RNA product of the second is predicted to form an unusual secondary structure, in which the T-arm is

the only readily-identifiable tRNA feature (see Supplementary Text S1). Closer inspection of the *T.*

nautili 30-1 genome reveals that the second, atypical Leu-CAA sequence occurs within a CRISPR

array of over 40 identical short repeats interspersed with unique spacer sequences that are derived

from foreign, invading DNA (Figure 2A) (for a recent review of CRISPR-Cas systems, see (45)). The

tenth spacer of this array appears to be derived from a viral tRNA gene, encompassing the T-arm and

one side of the acceptor stem (Figure 2B-C). A search for nucleotide sequence matches to the

spacer-10 sequence and viral DNA sequences (using the Nucleotide Basic Local Alignment Search

Tool, blastn) revealed a good match between the spacer and the Thr-TGT gene of several isolates

from the *Myoviridae* family, a family of bacteriophage known to infect Archaea and Bacteria (Figure

2D). Based on these results, we conclude that the tRNA gene fragment in *T. nautili* 30-1 is derived

from a viral tRNA gene. While tRNA genes have been documented in a range of bacteriophage

genomes (46, 47), CRISPR spacers do not commonly contain viral tRNA gene fragments (48, 49); to

the best of our knowledge, this is the first report of a viral tRNA gene fragment appearing in a

CRISPR-Cas array. Together, these results are consistent with the CRISPR-Cas-mediated

incorporation of a viral tRNA gene fragment into the *T. nautili* 30-1 genome, followed by the erroneous

listing of an additional, canonical tRNA gene in GtRNAdb.

Inspection of a further seven of the deviating tRNA genes revealed that they involve

fragments of tRNA genes resulting from insertion of mobile genetic elements into the genome. A

particularly clear example of such an element has been previously reported in *Pyrococcus yayanosii*

CH1 (NCBI Reference Sequence NC_015680.1; (50)), one of the 20 Thermococcaceae genomes: the

~21.4 kb mobile genetic element PYG1 lies between partial and complete Gln-TTG copies (Figure 3A-

C) (51). Both the partial and complete Gln-TTG sequences are listed in GtRNAdb as distinct, standard

tRNA genes (Figure 3D-E). Our dataset includes a further six examples of partial tRNA genes

resulting from similar integration events, with evidence for the integration of between ~9 kb and ~23

kb of viral DNA into various archaeal tRNA genes (see Supplementary Table S2). These findings are

in line with published results indicating that the incorporation of mobile genetic elements (*e.g.*,

temperate phages, integrative and conjugative elements (ICEs)) into prokaryotic genomes is a widely

occurring source of tRNA gene fragments; tRNA genes are common targets for the integration of such

elements, presumably as a result of tRNA gene stability and pervasiveness (52–54).

The ICE integration process typically results in one complete and one partial copy of the

target tRNA gene at opposing ends of the integrated element, providing a reliable method for their

delineation (55). Notably, the identification of these types of partial tRNA genes, and their full tRNA

gene counterparts, can also provide evidence of genome rearrangement events. For example, our

analysis indicates the integration of a ~20.7 kb prophage into the canonical Arg-TCT gene of the

common ancestor of *Thermococcus gammatolerans* EJ3 and *Thermococcus kodakarensis* KOD1,

followed by an inversion event in the KOD1 lineage that lead to the occurrence of the partial and full

tRNA gene copies on opposite strands of the chromosome, and a considerable distance apart (~179

kb; see Supplementary Table S2).

Category 2: Erroneously assigned tRNA isotypes resulting from ambiguous input sequence

The *Thermococcus litoralis* DSM 5473 genome (NCBI Reference Sequence NC_022084.1; (56)) contains seven of the 15 anomalies, including all three of the missing tRNA genes (Leu-CAG, Pro-GGG, Gln-CTG) and four additional tRNA genes (Leu-NAG, Pro-NGG, Und-NTG, Leu-CAA) (Figure 1A; Table 1). Careful analysis of the primary and secondary structures strongly indicates that three of the additional tRNA genes account for the three missing tRNA genes: additional gene Leu-NAG is the apparently missing gene Leu-CAG, Pro-NGG is Pro-GGG, and Und-NTG is Gln-CTG (see Supplementary Text S2 for tRNA structures). In each case, the presence of one or more N (*i.e.*, identity unknown) nucleotides in the tRNA gene sequence – and the *T. litoralis* DSM 5473 genome sequence – hampers the ability of tRNAscan-SE to predict the anticodon and/or isoacceptor family of the putative tRNA gene. Together, these errors account for a further six of the anomalies (Table 1). The seventh anomaly is described above in category 1b (a tRNA gene fragment resulting from the integration of incoming DNA). Hence, after manual annotation, the *T. litoralis* DSM 5473 genome is predicted to encode the standard archaeal tRNA gene set.

Category 3: Legitimate, additional tRNA genes

In the two categories above, manual annotation has accounted for 14 of the 15 observed tRNA gene set anomalies. The remaining anomaly is an additional copy of the Ala-TGC gene in *Palaeococcus pacificus* DY20341 (NCBI Reference Sequence NZ_CP006019.1; (57)). The two Ala-TGC gene copies reportedly occur within distinct, distally separated ribosomal RNA (*rrn*) operons, each consisting of a 16S gene, Ala-TGC, and a 23S gene. With the exception of an additional 32-bp stretch of DNA in one of the 23S genes, the two reported *rrn* loci are identical in primary sequence (including Ala-TGC). Notably, the other 19 Thermococcaceae genomes each encodes only a single *rrn* operon (and hence only one Ala-TGC gene; see Supplementary Table S3) (58). These results are consistent with intra-genomic duplication of the *rrn* operon in *Pa. pacificus* DY20341, leading to an actually divergent tRNA gene set in this taxonomically distinct organism (see Supplementary Figure S1). However, it should be noted that long stretches of nearly identical DNA sequence (such as *rrn* operons) pose considerable technical challenges for genome sequencing and assembly, and hence that *rrn* copy numbers (and any tRNA genes contained) may be inaccurate (59, 60).

The results so far show that, after manual annotation, the tRNA gene sets encoded in the 20 Thermococcaceae genomes adhere closely to the standard archaeal tRNA gene set of 46 single-copy tRNA genes; only a single probable deviation – duplication of Ala-TGC in *Pa. pacificus* DY20341 – remains (Figure 1B).

The tRNAscan-SE isotype score as an indicator of non-canonical tRNA genes

Manual annotation requires expert knowledge of tRNA structure, and detailed manual annotation may be too time consuming for very large data sets. It is therefore desirable to devise automated methods for distinguishing between canonical and non-canonical tRNA genes. As reported above, the automated detection of possible pseudogenes offered by tRNAscan-SE is valuable in this regard, as it

flags multiple tRNA gene fragments identified here as probable non-canonical tRNA genes. In
addition, during our manual annotation process, we noted lower than usual isotype scores for non-
canonical tRNA genes. Specifically, nine predicted tRNA genes contributing to the deviations of the
standard archaeal tRNA gene set exhibited relatively low isotype model scores (see Table 1).
Therefore, we performed a more systematic analysis of the value of both approaches (pseudogene
detection feature and isotype score) for detecting non-canonical tRNA genes. To this end, we
obtained the isotype scores of all predicted tRNA genes for the 20 Thermococcaceae genomes from
GtRNAdb and locally run tRNAscan-SE (version 2.0.6; see Materials and Methods), and extracted the
tRNA genes flagged as possible pseudogenes from the output of locally run tRNAscan-SE.
In each case, 929 tRNA genes were predicted, with isotype scores ranging between 2.3 and
138.2 (median=115.8; Figure 4A-B; Supplementary Table S5). While the isotype scores of most
predicted tRNA genes fell into a Gaussian-like large peak with a maximum of ~120, eight putative
tRNA genes – each corresponding to a partial tRNA gene sequence listed in Table 1 – received low
isotype scores (<85; Figure 4C-D). One of these low-scoring putative tRNA genes – *Thermococcus*
sp. ES1 Und-NNN (127-bp long; isotype score 16.8) – is called as a non-canonical tRNA of
undetermined or unknown isotype by tRNAscan-SE (and in GtRNAdb). A further six were identified as
pseudogenes by locally run tRNAscan-SE, yet listed as standard tRNA genes in GtRNAdb. The lack
of pseudogene definition in GtRNAdb appears to result from the active disablement of pseudogene
checking by tRNAscan-SE (as can be seen under the "Run Options/Stats" tab for individual
organisms listed in GtRNAdb). The decision to deactivate the possible pseudogene detection feature
when scanning archaeal (and bacterial) genomes for the current GtRNAdb listings was made in order
to avoid mis-classifying some low-scoring, but likely canonical, putative tRNA genes as pseudogenes
(Todd Lowe, personal communication).
The final relatively low-scoring putative tRNA gene – Val-CAC of *T. kodakarensis* KOD1
(isotype score 83.4) – is called as a standard tRNA gene by both tRNAscan-SE and GtRNAdb.
However, manual inspection of the primary sequence and the predicted secondary structure reveals
defects in the acceptor stem and D-arm that render the putative tRNA highly unlikely to be canonically
functional (see Supplementary Text S1). The tRNA with the next lowest isotype model score was
Pyrococcus furiosus DSM 3638 Arg-GCG (isotype score 88.1). Given that this single-copy tRNA gene
encodes an essential tRNA isotype, it is highly likely to give rise to a functional, canonical tRNA.
Indeed, its relatively low isotype score can be accounted for by the absence of a single nucleotide in
the acceptor stem (G70), which could arise from a point mutation or an error in the DSM 3638
reference genome sequence (see Supplementary Text S1). Overall, these results show that for the
tRNA gene dataset from the 20 Thermococcaceae genomes, a combination of (i) the automated
detection of probable pseudogenes/fragments, (ii) the tRNAscan-SE isotype score, and (iii) manual
inspection of tRNA genes with borderline isotype scores (~60-90) can be used as a valuable indicator
of the nature of putative tRNA genes (canonical versus non-canonical).
To investigate whether the above observations extend to a wider archaeal dataset, the
distribution of isotype scores for all putative tRNA genes in 210 complete (or nearly complete)
archaeal genomes listed in GtRNAdb were examined. For these 210 organisms, approximately

10,000 putative tRNA genes were listed in GtRNAdb and also predicted by locally run tRNAscan-SE

(the latter with pseudogene detection activated; see Materials and Methods). Isotype scores ranged

between -14.5 and 155.9, with a median of 105.9 (Figure 5; Supplementary Table S6). As observed

for the 20 Thermococcaceae genomes, the distribution of isotype scores for the tRNAs encoded in the

210 archaeal genomes showed a large Gaussian-like peak near 110 and a long tail of low-scoring

putative tRNA genes. tRNAscan-SE predicted 839 putative tRNA genes scoring below 85, accounting

for ~8 % of all predicted tRNA genes, and ~94 % of all predicted possible pseudogenes. Notably,

tRNAscan-SE predicts 742 standard tRNA genes with isotype scores below 85. On closer inspection,

many of these are unlikely to be canonically functional (*e.g.*, are predicted to encode truncated

tRNAs, or tRNAs with unusual secondary structures). Yet some, particularly those with isotype scores

between ~60 and ~90, are predicted to form tRNAs that may indeed be functional (for an example,

see *Thermoplasmatales archaeon* BRNA1 Ala-CGC in Supplementary Text S1). We also examined

the value of other types of scores generated by tRNAscan-SE (*e.g.*, HMM score, infernal score) in

identifying tRNA gene fragment but, compared to the isotype score, these resulted in a less clear

division of canonical and non-canonical tRNA genes (see Supplementary Figure S2). In summary,

while both the automated pseudogene detection and isotype scores are clearly informative, they

currently do not offer a watertight separation of canonical and non-canonical tRNA genes in Archaea.

DISCUSSION

In this work, we have performed a detailed manual analysis of the computationally predicted tRNA

gene sets listed in GtRNAdb for each of 20 high quality genome sequences from the archaeal

Thermococcaceae family. Foremost, we highlight that the tRNAscan-SE software has done a truly

impressive job of identifying putative tRNA genes in these organisms; with a degree of manual

curation, sensible and translationally complete canonical tRNA gene sets are recognizable in all 20

organisms (see Figure 1B). Additionally, and arguably equally usefully, tRNAscan-SE identifies a

number of non-canonical tRNA gene fragments resulting from biologically relevant and widespread

phenomena: CRISPR-Cas activity and the integration of incoming genetic elements (see Table 1).

However, the process of distinguishing between these two categories (canonical and non-canonical),

particularly on the widely-used, public database GtRNAdb, leaves some room for improvement.

345   Our analyses provide insights that allow progress towards three goals: (i) obtaining estimates

of the accuracy with which tRNAscan-SE identifies and subsequently annotates putative tRNA genes,

(ii) identifying specific sources of error, with the aim of highlighting specific avenues for possible

further improvement of tRNAscan-SE and/or its application by users, and (iii) gaining insight into the

properties and evolutionary dynamics of archaeal tRNA gene sets. Each of these is discussed below.

(i) The accuracy of automated tRNA gene prediction and annotation

In eleven independent cases (see Table 1, rows 1-11), our manual curation process yielded

conclusively different annotations of predicted tRNA genes than those obtained by the automated

process (listed in GtRNAdb). These included the complete removal of eight predicted tRNA genes

from the canonical tRNA gene sets, and the re-annotation of a further three to other – previously
absent – isotypes (Table 1, rows 9-11). Overall, fourteen apparent deviations were removed from a
set of 929 predicted tRNA genes, yielding an error rate of ~1.5 % for the 20 Thermococcaceae
GtRNAdb listings.
359

(*ii) Observed sources of error in automated tRNA gene annotations, and possible solutions*
When predicting the above tRNA gene sets, one source of error was the annotation of tRNA-like
sequences as standard tRNA genes (or, less commonly, tRNA genes of unknown or unidentified
isotypes) in GtRNAdb. Eight such instances were detected within our dataset of 20 Thermococcaceae
genomes (see Table 1). The existence of these sequences was attributed to two distinct mechanisms:
the integration of horizontally transferred genetic elements into host tRNA genes (seven instances),
and CRISPR-Cas activity (one instance).
The first mechanism – integration of genetic elements – is a general process in both Archaea
and Bacteria (52, 53) and, as such, similar tRNA gene-like remnants are expected to be a pervasive
feature of prokaryotic genomes. Indeed, in addition to the seven examples detected in our dataset,
inspection of GtRNAdb entries rapidly revealed instances of similar partial tRNA genes in Bacteria.
For example, GtRNAdb lists *Pseudomonas fluorescens* SBW25 as carrying two copies of the
Cys-GCA gene, with these occurring on either side of a putatively identified ~121 kb genomic island
(61). One copy is low-scoring (tRNA-Cys-GCA-2-1, isotype score 29.0), and the corresponding
putative tRNA was not detected in the mature tRNA pool of this bacterium (32). Notably, a third
possible source of genomic sequences with tRNA-like features in Bacteria is transfer-messenger RNA
(tmRNA) genes (62). Such tmRNA genes encode RNA products of a few hundred base pairs, and
which perform a role in the quality control of translation (as opposed to elongation). Since tmRNA
molecules carry some tRNA features, they have the potential to be identified and annotated by
tRNAscan-SE. An example is seen in the GtRNAdb listing for *Thiobacillus denitrificans* ATCC 25259:
tRNA-Und-NNN-3-1 (91-bp; isotype score 23.7) is actually the 3' end of a tmRNA gene.
The accurate detection and annotation of pervasive tRNA-like genomic sequences poses a
clear challenge to any tRNA identification software. To minimize the resulting overestimates of
canonical tRNA genes in Archaea, we recommend that users predict canonical tRNA gene sets – or
indeed, tRNA gene remnant sets – of interest using (i) locally run tRNAscan-SE, paying particular
attention to the pseudogene detection settings, and (ii) manual inspection, using the tRNAscan-SE
isotype score to focus on tRNA genes with isotype scores below ~85 (many of which are unlikely to
encode standard, canonical tRNA genes), and any detected possible pseudogenes (if applicable).
Further, we suggest that the addition of two new tRNA gene categories to the output of tRNAscan-SE
– *attR* (in line with established nomenclature denoting the ends of integrative elements; *e.g.*, (54)),
and tmRNA – would be both useful and biologically meaningful.
A second source of error in our dataset was the presence of N nucleotides in input genome
sequence leading to the mis-annotation of putative tRNA genes. Every such mis-annotation
generated two errors: a false positive tRNA gene (due to the erroneous presence of the mis-
annotated tRNA gene), and the corresponding false negative (due to the erroneous absence of the

correctly annotated tRNA gene). Three examples were detected in our dataset, resulting in six errors
(Table 1). This source of error could be minimized by a combination of users excluding (or, at a
minimum, noting) genomes containing significant numbers of Ns, and/or by the addition of a note to
the tRNAscan-SE output when Ns are detected within a putative tRNA gene sequence.
399
(iii) The properties and evolutionary dynamics of archaeal tRNA gene sets
The results reported here provide insight into the evolutionary dynamics of tRNA gene sets in
Archaea. Primarily, while the GC content of the 20 analyzed genomes varies widely (between 40 %
and 56 %; see Supplementary Table S1), the predicted tRNA gene sets exhibit remarkable stability;
with the likely exception of a single duplicate tRNA gene in *Pa. pacificus* DY20341, all 20 organisms
contain a tRNA gene set encoded by 46 single-copy tRNA genes (see Figure 1B). These
observations are consistent with previous suggestions of a 'standard' archaeal tRNA gene set (4, 5),
with occasional, small – and perhaps evolutionarily fleeting – instances of divergence.
408

AVAILABILITY

Raw data relating to this project is available at https://zenodo.org/record/6782366#.YsQSaZDP01A
The extended TIGRFAM database can be found at https://zenodo.org/record/3839790#.YjByaVzMI3g
The perl script used during phylogenetic tree construction, catfasta2phyml.pl, is available in the
GitHub repository (https://github.com/nylander/catfasta2phyml)
tRNAscan-SE is a freely available resource available online (http://lowelab.ucsc.edu/tRNAscan-SE/)
GtRNAdb is a publicly accessible resource available online (http://gtrnadb.ucsc.edu/)
NCBI is a publicly accessible resource available online (https://www.ncbi.nlm.nih.gov/)
rrnDB is a publicly accessible resource available online (https://rrndb.umms.med.umich.edu/)
BLAST is a publicly accessible resource available online (https://blast.ncbi.nlm.nih.gov/Blast.cgi)
419

SUPPLEMENTARY DATA

Supplementary Data are available online.

ACKNOWLEDGEMENT

None.

FUNDING

JG is supported by the Max Planck Society; AS is supported by the European Research Council
(ERC STG ASymbEL: 947317).

CONFLICT OF INTEREST

The authors declare no conflict of interest exists.

REFERENCES

1. Crick,F.H.C. (1966) Codon-anticodon pairing: The wobble hypothesis. *J. Mol. Biol.*, **19**, 548–555.

2. Agris,P.F., Eruysal,E.R., Narendran,A., Väre,V.Y.P., Vangaveti,S. and Ranganathan,S.V. (2018)
Celebrating wobble decoding: Half a century and still much is new. *RNA Biol.*, **15**, 537–553.

3. Chan,P.P. and Lowe,T.M. (2016) GtRNAdb 2.0: An expanded database of transfer RNA genes
identified in complete and draft genomes. *Nucl. Acids Res.,* **44**, D184-189.

4. Grosjean,H., de Crécy-Lagard,V. and Marck,C. (2010) Deciphering synonymous codons in the
three domains of life: Co-evolution with specific tRNA modification enzymes. *FEBS Lett.*, **584**, 252-
264.

5. Marck,C. and Grosjean,H. (2002) tRNomics: Analysis of tRNA genes from 50 genomes of Eukarya,
Archaea, and Bacteria reveals anticodon-sparing strategies and domain-specific features. *RNA*, **9**,
1189–1232.

6. Ehrlich,R., Davyt,M., López,I., Chalar,C. and Marín,M. (2021) On the track of the missing tRNA
genes: A source of non-canonical functions? *Front. Mol. Biosci.*, **8**, 643701.

7. Chan,P.P., Lin,B.Y., Mak,A.J. and Lowe,T.M. (2021) tRNAscan-SE 2.0: improved detection and
functional classification of transfer RNA genes. *Nucl. Acids Res.*, **49**, 9077–9096.

8. Laslett,D. (2004) ARAGORN, a program to detect tRNA genes and tmRNA genes in nucleotide
sequences. *Nucl. Acids Res.*, **32**, 11–16.

9. Wyman,S.K., Jansen,R.K. and Boore,J.L. (2004) Automatic annotation of organellar genomes with
DOGMA. *Bioinformatics*, **20**, 3252–3255.

10. Laslett,D. and Canback,B. (2008) ARWEN: a program to detect tRNA genes in metazoan
mitochondrial nucleotide sequences. *Bioinformatics*, **24**, 172–175.

11. Jühling,F., Pütz,J., Bernt,M., Donath,A., Middendorf,M., Florentz,C. and Stadler,P.F. (2012)
Improved systematic tRNA gene annotation allows new insights into the evolution of mitochondrial
tRNA structures and into the mechanisms of mitochondrial genome rearrangements. *Nucl. Acids
Res.*, **40**, 2833–2845.

12. Taquist,H., Cui,Y. and Ardell,D.H. (2007) TFAM 1.0: an online tRNA function classifier. *Nucl.
Acids Res.*, **35**, W350–W353.

13. Kinouchi,M. and Kurokawa,K. (2006) tRNAfinder: A software system to find all tRNA genes in the
DNA sequence based on the cloverleaf secondary structure. *J. Comput. Aided Chem.*, **7**, 116–124.

14. Chan,P.P. and Lowe,T.M. (2019) tRNAscan-SE: Searching for tRNA genes in genomic
sequences. *Methods Mol. Biol.*, **1962**, 1-14.

15. Lowe,T.M. and Chan,P.P. (2016) tRNAscan-SE on-line: integrating search and context for
analysis of transfer RNA genes. *Nucl. Acids Res.*, **44**, W54–W57.

16. Muramatsu,T., Nishikawa,K., Nemoto,F., Kuchino,Y., Nishimura,S., Miyazawa,T. and
Yokoyama,S. (1988) Codon and amino-acid specificities of a transfer RNA are both converted by a
single post-transcriptional modification. *Nature*, **336**, 179–181.

17. Chan,P.P. and Lowe,T.M. (2009) GtRNAdb: A database of transfer RNA genes detected in
genomic sequence. *Nucl. Acids Res.*, **37**, D93–D97.

18. Diwan,G.D. and Agashe,D. (2018) Wobbling forth and drifting back: The evolutionary history and
impact of bacterial tRNA modifications. *Mol. Biol. Evol.*, **35**, 2046–2059.

19. Withers,M., Wernisch,L. and Dos Reis,M. (2006) Archaeology and evolution of transfer RNA
genes in the *Escherichia coli* genome. *RNA*, **12**, 933–942.

20. Tremblay-Savard,O., Benzaid,B., Lang,B.F. and El-Mabrouk,N. (2015) Evolution of tRNA
repertoires in bacillus inferred with OrthoAlign. *Mol. Biol. Evol.*, **32**, 1643–1656.

21. Fujishima,K. and Kanai,A. (2014) tRNA gene diversity in the three domains of life. *Front. Genet.*,
5, 142.

22. Wald,N. and Margalit,H. (2014) Auxiliary tRNAs: Large-scale analysis of tRNA genes reveals
patterns of tRNA repertoire dynamics. *Nucl. Acids Res.*, **42**, 6552–6566.

23. McDonald,M.J., Chou,C.H., Swamy,K.B.S., Huang,H.D. and Leu,J.Y. (2015) The evolutionary
dynamics of tRNA-gene copy number and codon-use in *E. coli. BMC Evol. Biol.*, **15**, 163.

24. Novoa,E.M., Pavon-Eternod,M., Pan,T. and Ribas De Pouplana,L. (2012) A role for tRNA
modifications in genome structure and codon usage. *Cell*, **149**, 202–213.

25. Novoa,E.M., Jungreis,I., Jaillon,O. and Kellis,M. (2019) Elucidation of codon usage signatures
across the domains of life. *Mol. Biol. Evol.*, **36**, 2328–2339.

26. van der Gulik,P.T.S. and Hoff,W.D. (2016) Anticodon modifications in the tRNA set of LUCA and
488   the fundamental regularity in the standard genetic code. *PLoS ONE*, **11**, e0158342.

27. de la Torre,D. and Chin,J.W. (2021) Reprogramming the genetic code. *Nat. Rev. Genet.*, **22**, 169–
184.

28. Schimmel,P. (2018) The emerging complexity of the tRNA world: mammalian tRNAs beyond
protein synthesis. *Nat. Rev. Mol. Cell. Biol.*, **19**, 45–58.

29. Abbott,J.A., Francklyn,C.S. and Robey-Bond,S.M. (2014) Transfer RNA and human disease.
Front. Genet., **5**, 158.

30. Rak,R., Dahan,O. and Pilpel,Y. (2018) Repertoires of tRNAs: The couplers of genomics and
proteomics. *Annu. Rev. Cell Dev. Biol.*, **34**, 239–264.

31. Li,J., Zhu,L., Cheng,J. and Peng,Y. (2021) Transfer RNA-derived small RNA: A rising star in
oncology. *Semin. Cancer Biol.*, **75**, 29–37.

32. Ayan,G.B., Park,H.J. and Gallie,J. (2020) The birth of a bacterial tRNA gene by large-scale,
tandem duplication events. *eLife*, **9**, e57947.

33. Yona,A.H., Bloom-Ackermann,Z., Frumkin,I., Hanson-Smith,V., Charpak-Amikam,Y., Feng,Q.,
Boeke,J.D., Dahan,O. and Pilpel,Y. (2013) tRNA genes rapidly change in evolution to meet novel
translational demands. *eLife*, **2**, e01339.

34. Nilsson,A.I., Zorzet,A., Kanth,A., Dahlström,S., Berg,O.G. and Andersson,D.I. (2006) Reducing
the fitness cost of antibiotic resistance by amplification of initiator tRNA genes. *Proc. Nat. Acad.*
Sci. U.S.A, **103**, 6976–6981.

35. Rocha,E.P.C. (2004) Codon usage bias from tRNA's point of view: Redundancy, specialization,
and efficient decoding for translation optimization. *Genome Res.*, **14**, 2279–2286.

36. Schoch,C.L., Ciufo,S., Domrachev,M., Hotton,C.L., Kannan,S., Khovanskaya,R., Leipe,D.,
Mcveigh,R., O/Neill,K., Robbertse,B., *et al.* (2020) NCBI Taxonomy: a comprehensive update on
curation, resources and tools. *Database (Oxford)*, baaaa062.

37. Parks,D.H., Chuvochina,M., Rinke,C., Mussig,A.J., Chaumeil,P.-A. and Hugenholtz,P. (2022)
GTDB: an ongoing census of bacterial and archaeal diversity through a phylogenetically
consistent, rank normalized and complete genome-based taxonomy. *Nucl. Acids Res.*, **50**, D785-
D794.

38. Schut,G.J., Lipscomb,G.L., Han,Y., Notey,J.S., Kelly,R.M. and Adams,M.M.W. (2014) The order
Thermococcales and the family Thermococcaceae. In: Rosenberg,E., DeLong,E.F., Lory,S.,
Stackebrandt,E., Thompson,F. (eds), *The Prokaryotes*. Springer Berlin Heidelberg, Berlin,
Heidelberg, pp. 363–383.

39. Rich,A. and RajBhandary,U.L. (1976) Transfer RNA: Molecular structure, sequence, and
properties. *Annu. Rev. Biochem.*, **45**, 805–860.

40. Dombrowski,N., Williams,T.A., Sun,J., Woodcroft,B.J., Lee,J.-H., Minh,B.Q., Rinke,C. and
Spang,A. (2020) Undinarchaeota illuminate DPANN phylogeny and the impact of gene transfer on
archaeal evolution. *Nat. Commun.*, **11**, 3939.

41. Katoh,K. (2002) MAFFT: a novel method for rapid multiple sequence alignment based on fast
Fourier transform. *Nucl. Acids Res.*, **30**, 3059–3066.

42. Criscuolo,A. and Gribaldo,S. (2010) BMGE (Block Mapping and Gathering with Entropy): a new
software for selection of phylogenetic informative regions from multiple sequence alignments. *BMC*
Evol. Biol., **10**, 210.

43. Minh,B.Q., Schmidt,H.A., Chernomor,O., Schrempf,D., Woodhams,M.D., von Haeseler,A. and
Lanfear,R. (2020) IQ-TREE 2: New models and efficient methods for phylogenetic inference in the
genomic era. *Mol. Biol. Evol.*, **37**, 1530–1534.

44. Rinke,C., Chuvochina,M., Mussig, A.J., Chaumeil,P.-A., Davín,A.A., Waite,D.W., Whitman,W.B.,
Parks,D.H. and Hugenholtz,P. (2021) A standardized archaeal taxonomy for the Genome
Taxonomy Database. *Nat. Microbiol.*, **6**, 946-959.

45. Nidhi,S., Anand,U., Oleksak,P., Tripathi,P., Lal,J.A., Thomas,G., Kuca,K. and Tripathi,V. (2021)
Novel CRISPR–Cas systems: An updated review of the current achievements, applications, and
future research perspectives. *Int. J. Mol. Sci.*, **22**, 3327.

46. Kunisawa,T. (2000) Functional Role of Mycobacteriophage Transfer RNAs. *J. Theoret. Biol.*, **205**,
167–170.

47. Bailly-Bechet,M., Vergassola,M. and Rocha,E. (2007) Causes for the intriguing presence of
tRNAs in phages. *Genome Res.*, **17**, 1486–1495.

48. Nasko,D.J., Ferrell,B.D., Moore,R.M., Bhavsar,J.D., Polson,S.W. and Wommack,K.E. (2019)
CRISPR spacers indicate preferential matching of specific virioplankton genes. *mBio*, **10**, e02651-
18.

49. Paez-Espino,D., Morovic,W., Sun,C.L., Thomas,B.C., Ueda,K., Stahl,B., Barrangou,R. and
Banfield,J.F. (2013) Strong bias in the bacterial CRISPR elements that confer immunity to phage.
Nat. Commun., **4**, 1430.

50. Jun,X., Lupeng,L., Minjuan,X., Oger,P., Fengping,W., Jebbar,M. and Xiang,X. (2011) Complete
genome sequence of the obligate piezophilic hyperthermophilic archaeon *Pyrococcus yayanosii*
CH1. *J. Bacteriol.*, **193**, 4297–4298.

51. Li,Z., Li,X., Xiao,X. and Xu,J. (2016) An integrative genomic island affects the adaptations of the
piezophilic hyperthermophilic archaeon *Pyrococcus yayanosii* to high temperature and high
hydrostatic pressure. *Front. Microbiol.*, **7**, 1927.

52. Williams,K.P. (2002) Integration sites for genetic elements in prokaryotic tRNA and tmRNA genes:
sublocation preference of integrase subfamilies. *Nucl. Acids Res.*, **30**, 866–875.

53. Reiter,W.-D., Palm,P. and Yeats,S. (1989) Transfer RNA genes frequently serve as integration
sites for prokaryotic genetic elements. *Nucl. Acids Res.*, **17**, 1907–1914.

54. Johnson,C.M. and Grossman,A.D. (2015) Integrative and conjugative elements (ICEs): What they
do and how they work. *Annu. Rev. Genet.*, **49**, 577–601.

55. Hudson,C.M., Lau,B.Y. and Williams,K.P. (2015) Islander: a database of precisely mapped
genomic islands in tRNA and tmRNA genes. *Nucl. Acids Res.*, **43**, D48–D53.

56. Gardner,A.F., Kumar,S. and Perler,F.B. (2012) Genome sequence of the model hyperthermophilic
archaeon *Thermococcus litoralis* NS-C. *J. Bacteriol.*, **194**, 2375–2376.

57. Zeng,X., Jebbar,M. and Shao,Z. (2015) Complete genome sequence of hyperthermophilic
piezophilic archaeon *Palaeococcus pacificus* DY20341^{T}, isolated from deep-sea hydrothermal
sediments. *Genome Announc.*, **3**, e01080-15.

58. Stoddard,S.F., Smith,B.J., Hein,R., Roller,B.R.K. and Schmidt,T.M. (2015) rrnDB: improved tools
for interpreting rRNA gene abundance in Bacteria and Archaea and a new foundation for future
development. *Nucl. Acids Res.*, **43**, D593-598.

59. Rastogi,R., Wu,M., DasGupta,I. and Fox,G.E. (2009) Visualization of ribosomal RNA operon copy
number distribution. *BMC Microbiol.*, **9**, 208.

60. Acinas,S.G., Marcelino,L.A., Klepac-Ceraj,V. and Polz,M.F. (2004) Divergence and redundancy of
16S rRNA sequences in genomes with multiple *rrn* operons. *J. Bacteriol.*, **186**, 2629–2635.

61. Silby,M.W., Cerdeño-Tárraga,A.M., Vernikos,G.S., Giddens,S.R., Jackson,R.W., Preston,G.M.,
Zhang,X.X., Moon,C.D., Gehrig,S.M., Godfrey,S.A.C., *et al.* (2009) Genomic and genetic analyses
of diversity and plant interactions of *Pseudomonas fluorescens*. *Genome Biol.*, **10**, R51.

62. Keiler,K.C. and Ramadoss,N.S. (2011) Bifunctional transfer-messenger RNA. *Biochimie*, **93**,
1993–1997.

63. Tisza,M.J. and Buck,C.B. (2021) A catalog of tens of thousands of viruses from human
metagenomes reveals hidden associations with chronic diseases. *Proc. Natl. Acad. Sci. U.S.A.*,
118, e2023202118.

TABLES AND FIGURES

Table 1. Fifteen instances in which the 20 Thermococcaceae genomes listed in GtRNAdb
reportedly differ from the standard archaeal tRNA gene set

No.[a]	Affected isotype(s)[b]	Size (bp)[c]	Isotype Model Score[d]	Organism	Locus (strand)	tRNA scan-SE[e]	GtRNAdb[f]	Cat.[g]
1	(+)Ser-CGA	53	10.4	*T. gammatolerans* EJ3	621,672 - 621,724 (-)	Ps	St	1b
2	(+)Leu-CAA	218	2.3	*T. nautili* 30-1	706,451 - 706,668 (+)	Ps	St	1a
3	(+)Arg-TCT	83	21.4	*T. kodakarensis* KOD1	320,075 - 320,157 (-)	Ps	St	1b
4	(+)Val-CAC	72	83.4	*T. kodakarensis* KOD1	62,858 - 62,929 (-)	St	St	1b
5	(+)Gln-TTG	90	13.4	*P. yayanosii* CH1	1,321,660 - 1,321,749 (-)	Ps	St	1b
6	(+)Ser-CGA	69	30.9	*T.* sp. ES1	747,982 - 748,050 (+)	Ps	St	1b
7	(+)Und-NNN	127	16.8	*T.* sp. ES1	603,860 - 603,986 (-)	Und	Und	1b
8	(+)Leu-CAA	76	6.3	*T. litoralis* DSM 5473	326,949 - 327,024 (+)	Ps	St	1b
9-10	(+)Leu-NAG (-)Leu-CAG	88	135.1 (134.9)	*T. litoralis* DSM 5473	648,439 - 648,526 (+)	Ps	St	2
11-12	(+)Pro-NGG (-)Pro-GGG	78	120.3 (120.3)	*T. litoralis* DSM 5473	546,230 - 546,307 (-)	Ps	St	2
13-14	(+)Und-NTG (-)Gln-CTG	76	117.9 (117.9)	*T. litoralis* DSM 5473	825,204 - 825,279 (-)	Ps	St	2
15	(+)Ala-TGC	77	122.6	*Pa. pacificus* DY20341	1,527,238 - 1,527,314 (-)	Ps	St	3

Rows are ordered according to the phylogenetic tree in Supplementary Figure S1.
[a]Anomaly number (1-15)
[b](+) indicates additional tRNA genes, (-) indicates missing tRNA genes. Pairs of additional and
missing tRNA genes are placed together (see also Supplementary Text S1).
[c]Length in base pairs of the predicted tRNA gene
[d]Isotype score reported by GtRNAdb
[e,f]Classification by locally run tRNAscan-SE (version 2.0.6; pseudogene detection active), or
GtRNAdb (Ps=pseudogene, St=standard tRNA, Und=tRNA of undetermined or unknown isotype)
[g]Category that the anomaly falls into. 1=erroneously predicted tRNA gene resulting from (1a)
CRISPR-Cas activity or (1b) integration of a mobile genetic element (*i.e.*, *attR* sites), 2=isotype
misassignment, 3=legitimate additional tRNA gene).

601
Figure 1. Heatmap depicting putative tRNA gene sets for 20 Thermococcaceae family organisms. (**A**) tRNA gene sets as listed on GtRNAdb (predicted
via tRNAscan-SE 2.0.6, with pseudogene detection deactivated), and (**B**) following the manual annotation procedure described in the Materials and Methods.
Organisms with non-standard tRNA gene sets are marked by an asterisk (seven in panel A, one in panel B).

Figure 2. A partial tRNA gene resulting from CRISPR-Cas activity in _T. nautili_ 30-1 (category 1a in Table 1). (**A**) Cartoon depicting the _T. nautili_ 30-1 CRISPR-Cas locus of interest (genomic bases 705,938–708,770). The array consists of 41 identical 30-bp direct repeats (grey rectangles) interspersed with 41 unique spacers of ~36 bp (various shapes). The 218-bp Leu-CAA-2-1 tRNA identified by GtRNAdb contains sequence from spacers 7-10 and repeats 7-9 (bases 706,451–706,668). Of these, spacer 10 contains the actual tRNA gene remnant. (**B**) Enlargement of spacer 10 and flanking repeats (bases 706,604–706,699). Blue text indicates the 32-bp region matching to viral tRNA sequence (see panel D). (**C**) The partial tRNA gene sequence in spacer 10 is predicted by tRNAscan-SE to form a secondary structure in which the T-arm (turquoise) and part of the acceptor stem (magenta) are identifiable. (**D**) The T-arm and acceptor stem of tRNA-Thr(UGU) from _Myoviridae_ isolates (_Myoviridae_ sp. Isolate ctU5h13 (63) is shown here) are a good match for spacer 10; the sequence matches 29 of the 32 blue bases in panel B (matches indicated by thick grey lines).

620
Figure 3. A partial tRNA gene resulting from integration of a mobile genetic element in _P._
yayanosii CH1 (category 1b in Table 1). (**A**) _P. yayanosii_ CH1 contains a partial and a complete
copy of the Gln-TTG tRNA gene, separated by the ~21.4 kb genomic island, PYG1 (51). PYG1 is
flanked by two 44-bp direct repeats (1,321,660–1,321,703 in red; 1,342,940–1,342,983 in green), only
the second of which forms part of a standard, canonical Gln-TTG gene. (**B, C**) Primary sequences of
the 90-bp and 76-bp Gln-TTG genes listed as putative standard tRNA genes in GtRNAdb. The 44-bp
direct repeats are highlighted (red/green), with * indicating a 1-bp difference. (**D, E**) tRNAscan-SE-
derived secondary structures of the 90-bp and 76-bp Gln-TTG gene sequences, respectively. Thick
red and green lines indicate the 44-bp repeats.
630

Figure 4. tRNAscan-SE isotype score as a predictor of canonical versus non-canonical tRNA
genes in the 20 Thermococcaceae genomes. Histograms of the frequency of tRNA isotype scores
from all 929 tRNA genes listed in (**A**) GtRNAdb, and (**B**) predicted by locally run tRNAscan-SE, for the
20 Thermococcaceae genomes. (**C-D**) Closeup of isotype scores 0-100 in panels A and B. Numbers
1-8 relate to the anomalies listed in Table 1. tRNA gene categories are generated by tRNAscan-SE.
Note that in panels A and C (GtRNAdb) the pseudogene category ('pseudo') is absent, due to
disabling of pseudogene detection function (see 'Run Options/Stats' tab in GtRNAdb listings).

Figure 5. tRNAscan-SE isotype score as a predictor of canonical versus non-canonical tRNA genes in 210 archaeal genomes. (**A**) Histogram of the frequency of tRNA isotype scores from ~10,000 tRNA genes from 210 archaeal genomes listed in (**A**) GtRNAdb, and (**B**) predicted by locally run tRNAscan-SE. (**C-D**) Closeup of isotype scores 0-100 in panels A and B. Note that in panels A and C (GtRNAdb) the pseudogene category ('pseudo') is again absent, due to disabling of pseudogene detection function (see 'Run Options/Stats' tab in GtRNAdb listings). Category 'seleno' denotes putative selenocysteine tRNA genes. Dotted lines demonstrate the use of tRNAscan-SE isotype scores as a useful indicator of the nature of putative archaeal tRNA genes: <60=likely non-canonical, 60-90=uncertain (*i.e.*, may be either canonical or non-canonical), >90=likely canonical.

SUPPLEMENTARY MATERIALS LIST AND LEGENDS

Supplementary Figure S1. Phylogenetic tree highlighting the evolutionary relationships of the
20 Thermococcaceae genomes listed in GtRNAdb. The complete genome sequences of all
genomes were obtained (see Supplementary Tables S1, S5, and S6). Alignments were generated
using 12,403 amino acid positions across 43 shared marker proteins (see Supplementary Table S4).
The support shown next to the branches is based on 100 bootstrap replicates. Three genomes -
Methanococcus vannielii SB, *Methanococcus maripaludis*, and *Methanococcus aeolicus* – were used
to root the tree.

Supplementary Figure S2. tRNAscan-SE isotype score can be used as an indicator of whether
a putative archaeal tRNA gene is canonical or non-canonical. Frequencies of locally run
tRNAscan-SE (version 2.0.6)-derived scores for ~10,000 putative tRNA genes in 210 archaeal
genomes. (**A**) Isotype score (also presented in Figure 5B). (**B**) Infernal score, (**C**) HMM score, and (**D**)
Secondary structure-only score. See Chan and Lowe 2021 (doi:10.1093/nar/gkab688) and references
therein for a detailed description of score calculations.
669

670   **Supplementary Text S1. Primary and predicted secondary structures of putative tRNA genes of**
671   **interest.** Primary sequences were obtained from GtRNAdb. Secondary structures and listed scores
672   were predicted using the web based tRNAscan-SE 2.0 (sequence source: archaeal; search mode:
673   default).
674

675   **Supplementary Table S1: Genomic characteristics of the 20 Thermococcaceae organisms.**
676

677   **Supplementary Table S2: Evidence for partial tRNA sequences resulting from the integration**
678   **of genetic elements.** Genomic details of seven partial tRNA fragments resulting from the integration
679   of prophages (*i.e.*, erroneously identified as canonical tRNA genes; category 1b in Table 1).
680

681   **Supplementary Table S3: Details of ribosomal RNA (*rrn*) operons and associated tRNA genes**
682   **in the 20 *Thermococcaceae* genomes.** Only one genome differs from the norm of a single *rrn*
683   operon and Ala-TGC gene: *Pa. pacificus* DY20341 carries two *rrn* operons and thus two Ala-TGC
684   genes (*i.e.*, a real deviation; category 3).
685

686   **Supplementary Table S4: Details of markers for drawing of the Thermococcaceae phylogenetic**
687   **tree.**
688

689   **Supplementary Table S5: tRNA gene predictions, annotations, and scores for 20**
690   **Thermococcaceae organisms.** Tab 1: downloaded from GtRNAdb. Tab 2: predicted using locally
691   run tRNAscan-SE (version 2.0.6, as for the GtRNAdb entries in Tab 1).
692

693 **Supplementary Table S6: tRNA gene predictions, annotations, and scores for 210 archaea.** Tab

694 1: downloaded from GtRNAdb. Tab 2: predicted using locally run tRNAscan-SE (version 2.0.6, as for

695 the GtRNAdb entries in Tab 1).