A Buprenorphine depot formulation provides effective sustained post-surgical analgesia for 72h in mouse femoral fracture models

Authors: Angelique Wolter[1,2,3], Christian H. Bucher[4,5], Sebastian Kurmies[1,2], Viktoria Schreiner[6], Frank Konietschke[5,7], Katharina Hohlbaum[3,8], Robert Klopfleisch[9], Max Löhning[1,2], Christa Thöne-Reineke[3], Frank Buttgereit[1,2], Jörg Huwyler[6], Paulin Jirkof[10,11], Anna E. Rapp[1,2,12*#], Annemarie Lang[1,2,13*#]

*these authors contributed equally

Affiliations:

[1]Charité – Universitätsmedizin Berlin, corporate member of Freie Universität Berlin and Humboldt-Universität zu Berlin, Department of Rheumatology and Clinical Immunology, Berlin, Germany

[2]German Rheumatism Research Centre (DRFZ), a Leibniz Institute, Berlin, Germany

[3]Institute of Animal Welfare, Animal Behavior and Laboratory Animal Science, Department of Veterinary Medicine, Freie Universität Berlin, Berlin, Germany

[4]Julius Wolff Institute, Charité – Universitätsmedizin Berlin, Berlin, Germany

[5]Berlin Institute of Health Center for Regenerative Therapies (BCRT), Charité – Universitätsmedizin Berlin, Berlin, Germany

[6]Division of Pharmaceutical Technology, Department of Pharmaceutical Sciences, University of Basel, Basel, Switzerland

[7]Charité-Universitätsmedizin Berlin, Corporate Member of Freie Universität Berlin and Humboldt-Universität zu Berlin, Institute of Biometry and Clinical Epidemiology, Berlin, Germany

[8]German Federal Institute for Risk Assessment (BfR), German Centre for the Protection of Laboratory Animals (Bf3R), Berlin, Germany

[9]Institute of Veterinary Pathology, Department of Veterinary Medicine, Freie Universität Berlin, Berlin, Germany

[10]Center for Surgical Research, University Hospital Zurich, University Zurich, Zurich, Switzerland

[11]Office for Animal Welfare and 3Rs, University of Zurich, Zurich, Switzerland

[12]Dr. Rolf M. Schwiete Research Unit for Osteoarthritis, Department of Orthopedics (Friedrichsheim), University Hospital Frankfurt, Goethe University, Frankfurt, Germany

[13]Departments of Orthopaedic Surgery and Bioengineering, University of Pennsylvania, PA, United States

#Correspondences:

Anna E. Rapp, Dr. Rolf M. Schwiete Research Unit for Osteoarthritis, Department of Orthopedics (Friedrichsheim), University Hospital Frankfurt, Goethe University, Frankfurt, Germany

Rapp@med.uni-frankfurt.de

Annemarie Lang, Departments of Orthopaedic Surgery and Bioengineering, University of Pennsylvania, PA, United States

Annemarie.lang@pennmedicine.upenn.edu

Summary

Adequate pain management is essential for ethical and scientific reasons in animal experiments and should completely cover the period of expected pain without the need for frequent re-application. However, current depot formulations of Buprenorphine are only available in the USA and have limited duration of action. Recently, a new microparticulate Buprenorphine formulation for sustained release has been developed and safely used in first studies serving as an alternative product within Europe. Pharmacokinetics indicate a potential effectiveness for about 72h; the period that must be covered with analgesia in mouse fracture models. Here, we investigated whether the administration of the newly developed sustained-release Buprenorphine formulation ensures a continuous and sufficient analgesia in mouse femoral fracture models as a potent alternative to the application of Tramadol via drinking water. Both protocols were examined for analgesic effectiveness and side effects on experimental readout in two femoral fracture models using rigid and flexible external fixators in male and female C57BL/6N mice. We monitored general parameters of wellbeing, and model-specific pain parameters. Both, Tramadol in the drinking water and the sustained-release Buprenorphine provided effective analgesia. However, in male mice with flexible fixation, composite pain scores were significantly higher with Tramadol than in the other groups, and locomotion during gait analysis was more profoundly altered. Fracture healing outcome was not different between analgesic regimes. The availability of a sustained-release formulation of Buprenorphine in Europe would be beneficial for pain management in preclinical studies to increase animal welfare and actively support the refinement of painful animal experiments.

Keywords: pain management, refinement, bone repair, animal model, tramadol

1 Introduction

Despite the increasing progress in alternative methods, animals – especially mice – are still widely used and required in fundamental and translational research to study the complexity of biological and pathophysiological processes. Therefore, the active implementation of the 3R principle (Replace – Reduce – Refine), with a particular importance of *Refinement* forms the indispensable basis for a humane approach to conduct animal experiments. Thus, adequate pain assessment and medication in animals before, during and after the experimental procedure is crucial to decrease suffering and support data quality. Furthermore, handling-induced stress and insufficiently treated pain can affect animal behavior and physiological responses, especially in an immunological context, leading to potential falsification of the scientific outcomes and reduced reproducibility (Balcombe et al., 2004; Carstens and Moberg, 2000; Guo and Dipietro, 2010; Page, 2003; Peterson et al., 2017; Jirkof, 2017). However, evidence-based data on individual pain management efficiencies in surgical mouse models is still rare (Flecknell, 2018; Jirkof, 2017) and the reporting quality of the used analgesic protocols is often insufficient (Carbone and Austin, 2016; Wolter et al., 2021).

The potent opioid Buprenorphine is very often used in rodents after surgical intervention (Guarnieri et al., 2012; Roughan and Flecknell, 2002). Due to a short half-life of approximately 3 hours (Bullingham et al., 1980; Gades et al., 2000; Yu et al., 2006), frequent application intervals are required resulting in repeated handling of the animals. The commonly reported application intervals of subcutaneous or intraperitoneal injection every 8 – 12h can lead to pain peaks and insufficient analgesic coverage (Jirkof et al., 2015; Wolter et al., 2021). The parenteral application of other opioids such as Morphine, Tramadol and Fentanyl is most often not suitable for pain alleviation in rodents, as their half-life is even shorter (Gades et al., 2000; Evangelista-Vaz et al., 2018; Girón et al., 2002). It has also been suggested that the experience of pain and sensitivity to opioid analgesia can be strain and sex-dependent (Fillingim and Gear, 2004; Kest et al., 2000; Smith, 2019). Thus, the design of adequate pain management regimes needs to consider the duration and nature of the expected pain, animal species-specific sensitivity to opioid analgesia (including strain and sex) and the evaluation of options to reduce re-injection intervals and therefore, handling time.

To reduce handling-associated stress and to ensure continuous analgesic coverage, an alternative application route of Buprenorphine or Tramadol in the drinking water has been routinely used e.g., in orthopedics-related mouse models (Bucher et al., 2019; Jirkof et al., 2019; Rapp et al., 2018; Histing et al., 2016). As the uptake of analgesics with the drinking water is dependent on the drinking frequency and amount, the overall effectiveness of this treatment strategy might be highly influenced by e.g., reduced activity and water intake after anesthesia and surgery and circadian activity (Sauer et al., 2016). A drug formulation that extends the analgesic effects due to sustained parenteral drug release can overcome such challenges and serve as a powerful tool to refine today's analgesic regimens after surgical interventions in animal experiments. Current depot/sustained-release formulations of Buprenorphine for mice and rats are either not commercially available or only available in the United States of America (USA). Examples include Buprenorphine SR-LAB (ZooPharm) or Ethiqa XR (Fidelis Animal Health). Attempts to import these products to Europe have failed due to missing approval through the European Medical Evaluation Agency (EMEA).

To fill the gap, Schreiner et al. have successfully developed a poly-lactic-co-glycolic acid (PLGA) based

microparticulate drug formulation for sustained drug release of Buprenorphine (Schreiner et al., 2020;

Schreiner et al., 2021). In a proof-of-concept study in female C57BL/6J mice, they observed therapeutic-

relevant drug levels of the sustained-release Buprenorphine (BUP-Depot) in the brain for more than 24h,

an antinociceptive effect in the hot plate test and pain relief after a minor abdominal surgery for at least

72h. To test the BUP-Depot for the effectiveness in a wider range of surgical interventions, we here

performed a pain management study in two mouse femoral fracture models. The selection of appropriate

analgesics in experimental fracture healing studies is limited by potential adverse effects of nonsteroidal

anti-inflammatory drugs (NSAIDs) on inflammatory events during the initial and proinflammatory phase

of fracture healing (Bove et al., 2009; Cottrell and O'Connor, 2010; Gerstenfeld et al., 2003). Moreover,

the nature of pain divides into the pain sensitization during wound healing and the bone-related pain

due to the cutting of the periosteum during osteotomy and is therefore a useful representation of

potential pain conditions (Tappe-Theodor et al., 2022).

To this end, we tested the newly developed BUP-Depot as a potential alternative to the established

application of Tramadol via the drinking water. Both pain management protocols were examined for their

analgesic efficacy and adverse effects on experimental readouts in two femoral fracture models using

rigid and flexible external fixators. We monitored (i) general parameters of wellbeing e.g., body weight,

food and water intake, nestbuilding and explorative behavior and (ii) model-specific pain parameters

including walking behavior (limp score) and CatWalk analysis. In addition, fracture healing outcomes

were examined at the end of the study to exclude negative influences on the regeneration process. To

overcome potential sex-dependent differences in response to the analgesic protocol male and female

mice were included. A refined analgesic regime using a sustained-release Buprenorphine was

implemented for post-surgical pain management in a mouse bone fracture models to explore its potential

to improve animal welfare in a wider range of pre-clinical animal models.

2 Animals, Material and Methods

Animals and husbandry

The study was conducted according to the guidelines of the German Animal Welfare Act, National Animal Welfare Guidelines, and was approved by the local Berlin state authority (Landesamt für Gesundheit und Soziales – LAGeSo; permit number: G0044/20). Health monitoring in the animal facility was performed according to the FELASA guidelines (Supplementary Information). A total of 40 male and 40 female C57BL/6N mice aged 8 weeks were either provided by the Experimental Medicine Research Facilities (Charité-Universitätsmedizin Berlin, Berlin, Germany) or purchased from Charles River Laboratories (Sulzfeld, Germany). Mice underwent the first intervention (anesthesia/analgesia) at 10 weeks (body weight – males: 25.96 ± 2.1 g; females: 21.41 ± 1.3 g) and osteotomy with 12 weeks (body weight – males: 27.61 ± 1.9 g; females: 22.75 ± 1.3 g). Mice were housed in a semi-barrier facility with limited access, protective clothing, and individually ventilated cages (IVC). Housing conditions included a 12/12–h light/dark cycle (lights on from 6:00 a.m. to 6:00 p.m.), room temperature between 22 ± 2°C and a humidity of 55 ± 10%. Mice had access to food (Standard mouse diet, Ssniff Spezialdiäten, Soest, Germany) and tap water *ad libitum*.

Upon arrival, the mice were randomly divided into groups of two per cage. All mice were housed in Eurostandard Type II clear-transparent plastic cages. In some instances, two males had to be separated due to aggressive behavior. Those mice were housed in a separated pair housing system consisting of compact cages (Green Line IVC Sealsafe PLUS Rat GR 900, Tecniplast, Milan, Italy), which were divided in two equally sized compartments by a perforated transparent cage partition wall enabling olfactory, acoustic, and visual interaction (Hohlbaum et al., 2020). All cages contained wooden chips (SAFE FS 14, Safe Bedding, Rosenberg, Germany) as well as 20 g Envirodri (Shepherd Specialty Papers, USA) and one additional shredded paper towel as bedding and nesting material. Cages were also equipped with a clear handling tube (Datesand Group, Bredbury, UK) and a mouse double swing (Datesand Group, Bredbury, UK). No houses were provided to allow reliable scoring and to reduce the risk of injury after osteotomy. In addition, tunnels and double swings were removed from the cages to reduce the risk of injury due to the external fixator. Two single swings (Datesand Group, Bredbury, UK) per cage were reinstalled 5 days after osteotomy. Cages were changed weekly by the experimenters only. When nesting material was dry, it was transferred to the new cage and replenished to 20 g with new nesting material. In the week of anesthesia and of the osteotomy, the cages were renewed the day before the intervention, to provide a clean surrounding and still allow the animals to get accustomed to the new environment and to build proper nests beforehand. Animals were tunnel handled only and all experimenters performing behavioral analyses were female. Animal husbandry and care was in accordance with contemporary best practice and all individuals involved in the study had a high level of competency.

Humane endpoints were predefined and divided into clinical parameters (e.g., body weight reduction >20%, apathy) and model-specific parameters (e.g., open skin wound, no usage of operated limb). No humane endpoints were reached during the conduct of the study. Potential 3R applications were thoroughly researched and if possible, implemented within this study e.g. enrichment, positive reinforcement during training (white chocolate) and tunnel handling.

Study design and experimental timeline

The study included 8 groups with each n= 9–10 mice, comparing male and female mice, rigid and flexible

external fixators, and two different pain management protocols (Tramadol via drinking water vs.

sustained-release Buprenorphine (BUP-Depot) (**Fig. 1A**). Exclusion of mice and data are detailed in the

Supplementary Information. Cages and mice received individual random numbers which did not allow

any inferences for the analgesic regimen or fixation group. Experimenters performing the pre- and post-

surgical training and investigations were blinded for the analgesic regimen. Reporting of this study was

carried out in compliance with the ARRIVE 2.0 guidelines, including the "Arrive Essential 10" as well as

a majority of the "Arrive Recommended Set". The study was pre-registered in the Animal Study Registry

(Bf3R, Germany; DOI: 10.17590/asr.0000221).

After arrival, mice were allowed to acclimate for 5 days. Training was then performed by one female

experimenter, following a precise 2-week schedule to accustom the mice to the experimenter, tunnel

handling, the Noldus CatWalk XT (Noldus, Wageningen, Netherlands) and observation boxes (Ugo

Basile, Gemonio, Italy). Baseline measurements were also obtained during this period (**Fig. 1B**). To

correct for individual behavioral changes induced by anesthesia and analgesia alone, mice were first

anesthetized and received their assigned analgesic protocol without any further surgical procedure (first

intervention). This was followed by closely scheduled monitoring and assessment at 12h, 24h, 48h, and

72h. At day 14 after the first intervention, mice were then subjected to anesthesia, analgesia, and

osteotomy and again assessed at the same timepoints as well as at day 10 post-surgical. Mice were

euthanized at day 14 post-osteotomy to retrieve the osteotomized left femur.

Figure 1. Group assignment and study design. Overview on (**A**) the group assignment, as well as (**B**) time points and
measurements of the different parameters. To assure a rapid onset of the analgetic effect, each animal received a single s.c. dose
of Temgesic (1 mg/kg) at the beginning of each intervention. Tramadol (0.1 mg/ml) in drinking water was provided one day before
and for three consecutive days after interventions. A single dose of sustained-release BUP-Depot (1.2 mg/kg s.c) was
administered after interventions.

Analgesic regimes

Each mouse received a single s.c. injection of fast acting standard Buprenorphine (1 mg/kg Temgesic, RB Pharmaceuticals, Heidelberg, Germany) at the beginning of each intervention (anesthesia/analgesia alone and osteotomy) (**Fig. 1**). Depending on the randomly assigned group, mice either additionally received Tramadol administered with the drinking water (0.1 mg/ml, Drops, Grünenthal, Stolberg, Germany) or a s.c. injection of the sustained-release Buprenorphine (BUP-Depot; 1.2 mg/kg). Tramadol via the drinking water was administered one day before and for three consecutive days after both interventions. The BUP-Depot was injected once at the end of both interventions. The PLGA based BUP-Depot (RG 502 H-Big) was prepared at the University of Basel, Switzerland, as described previously by Schreiner et al. (Schreiner et al., 2020; Schreiner et al., 2021). The BUP-Depot was imported in accordance with national regulations for controlled substances (BtM import authorization No. 4679477). Four different batches of BUP-Depot were used during the study. Each batch was analyzed before shipment with respect to drug content, reconstitution time, and drug release kinetics as described previously (Schreiner et al., 2020; Schreiner et al., 2021). The-BUP-Depot was stored as a lyophilizate in crimped glass vials at 4°C. Each vial was reconstituted with physiological saline (0.9% NaCl) immediately before administration.

Anesthesia and osteotomy

Independent of the intervention, all mice were anesthetized with isoflurane (~2–3%; provided in 100% oxygen) (CP-Pharma, Burgdorf, Germany) in an induction chamber before being weighed, and moved onto a heating pad (37°C). Anesthesia was maintained at ~2% via a nose cone. Eye ointment, physiological saline (0.5 ml, 0.9% NaCl (B. Braun Melsungen, Melsungen, Germany)), Clindamycin (45 mg/kg, Ratiopharm, Ulm, Germany) and a single s.c. injection of Buprenorphine (1 mg/kg; Temgesic) were administered. Anesthesia was then upheld for 15 minutes for the first intervention. For osteotomy, the left femur was shaved and disinfected with Braunoderm (B. Braun Melsungen, Melsungen, Germany), the osteotomy was conducted as described previously (Jirkof et al., 2019; Lang et al., 2019; Lang et al., 2022; Rapp et al., 2018). In brief, a longitudinal skin incision was made between knee and hip, and the musculus vastus lateralis and musculus biceps femoris were bluntly separated to expose the femur. Two standardized external fixators (rigid: 18.1 N/mm; flexible: 3.2 N/mm, both RISystem, Davos, Switzerland) were used for stabilization. The external bar of the fixator was positioned parallel to the femur and all pins were positioned accordingly. Afterwards, an approximately 0.5 mm osteotomy gap was created between the second and third pin using a Gigli wire saw (0.44 mm; RISystem, Davos, Switzerland) and the gap was flushed with saline (NaCl (B. Braun Melsungen, Melsungen, Germany)). Muscle and skin were closed with two layers of sutures (muscle: 5-0 Vicryl, skin: Ethilon 5-0, both Ethicon, Raritan, USA). For recovery, the mice were returned to their home cages under an infrared lamp and were closely monitored.

Body weight and food/water intake

Animals were weighed right before the intervention (defined as 0h), and at 12h, 24h, 48h and 72h after each intervention and then every other day until osteotomy/euthanasia. Food/water intake were measured per cage (i.e., two mice) by weighing the food and water bottles every 24h, beginning 1 day

prior to each intervention and ending 3 days after. The difference to the previous value was calculated.

All measurements were normalized to the respective baseline values at time point 0h.

Explorative test and nest complexity score

Both scores were assessed before any other assessment or handling of the mice. To examine the

motivation of the mice to explore and interact (sniffing, holding with forepaws, or carrying) with a foreign

object in the home cage, we added a Nestlet (Ancare, Bellmore, USA) to the cages and observed the

mice for one minute. The explorative test was scored 1 (interaction) or 0 (no interaction) per cage while

an interaction of one animal of the cage was deemed as sufficient for a positive score. Nest complexity

scoring was performed following Hess et al. (Hess et al., 2008) in the home cage assigning scores

between 0 – 5.

Composite pain score

A composite pain score was used to combine the assessment of facial expression and clinical

appearance (Jirkof et al., 2019; Langford et al., 2010) (**Table 1**). The maximal score was 9. Mice were

transferred into a clear observation box (Ugo Basile, Gemonio, Italy) and were individually filmed for 3

min after an acclimatization period of 1 min (Basler Video Recording Software, Ahrensburg, Germany).

Video analysis was performed by one blinded observer at timepoints 12h, 24h, 48h, and 72h post-

intervention. As anesthesia and analgesia also impact the composite pain score, we calculated the delta

composite pain score for each mouse by subtracting the scores from each individual mouse post-

anesthesia from their respective scores post-osteotomy. This allows us to evaluate the effect of the

surgical procedure on an individual base without the interference of behavioral or clinical changes

induced by the anesthesia and analgesia alone (without surgical intervention) (Miller et al., 2015).

Table 1: Composite Pain Score

Parameter	Specification	Score (0 = not present)
Orbital tightening	Faint narrowing of the orbital area up to a tightly closed eyelid	Range: 0 – 2 0.5 = minimal 1 = moderately 1.5 = moderately severe 2 = severe
Ear position	Ears pulled back or rotated outwards and/or backwards	Range: 0 – 2 1 = moderately 2 = severe
Posture	Crouched posture, head and nose positioned towards the ground	Range: 0 – 1 0.5 = held for less than 10 s 1 = held for more than 10 s
Coat condition	Coat appeared disheveled or unkempt	Range: 0 – 1 0.5 = only certain body parts 1 = all over the body
	Erected fur, mice appear scruffy	Range: 0 – 1 0.5 = only on one body part 1 = on more than one body part, generalized
Movement	Apathetic, sedated, tipsy (0 – 1); Crawling (0 – 1)	Range: 0 – 1 0.5 = minimal 1 = moderately

Walking behavior – limp score

To assess the walking behavior of each individual mouse, limp scoring was performed as described by

Jirkof et al. (Jirkof et al., 2019). The mice were individually transferred to conventional type III cages that

contained the same type of wooden chips as the home cages. After an acclimation period of 5 min, a 3

min video was recorded. Walking behavior was examined at time points concurrent with the CatWalk

analysis at 24h, 48h, and 72h post-anesthesia and post-osteotomy as well as 10d post-osteotomy. Video

analysis was performed by two blinded observers and scores from 0 – 4 were assigned (**Table 2**). If

walking seemed to be impaired due to a mechanical problem (e.g., displacement of the patella) one

point was subtracted from the assigned score.

Table 2: Limp Score

Specification	Limp Score
Normal use, physiological gait	0
Complete ground contact and sporadic limping or alteration of the gait pattern	1
No complete ground contact or limping, constant alteration of the gait pattern	2
Partial non-use of the limb	3
Complete lack of use	4

CatWalk analysis

Specific gait analysis was performed using the CatWalk XT Gait Analysis system for rodents (Noldus,

Wageningen, Netherlands). Prior to data acquisition, mice were trained to the setup during three training

sessions. Data was then acquired at 24h, 48h, 72h and 10d post-anesthesia (not shown in results) and

post-osteotomy. To acquire at least three evaluable runs for analysis per timepoint, multiple runs per

animal were recorded. Post-acquisition, the runs were screened, and non-compliable runs as well as

interrupted runs (e.g., by sniffing, rearing) were deleted. Runs that noticeably differed from the rest of

the runs of these trials were also deleted, leaving an average of 4.4 runs per animal and timepoint for

analyses. All runs were classified automatically by the Noldus CatWalk XT software (version XT10.6).

Afterwards, every run was reviewed by one researcher and classification errors were corrected

manually.

From the obtained data, mean speed (cm/s) (velocity) and the following parameters were analyzed for

the osteotomized left hind leg: mean intensity, stand duration (s) and stride length (cm). The Noldus

CatWalk automatically calculated an average speed (velocity) of all runs per animal and timepoint. The

two mean velocity values from both baseline timepoints were used to create one baseline average

velocity. The velocity values at all time points were then normalized to the baseline values (timepoint

value divided by average baseline value). The same calculation was done for mean intensity (a measure

of load bearing), stand duration and stride length, resulting in relative mean intensity, relative stand

duration and relative stride length.

Euthanasia and sample collection

Euthanasia was carried out according to contemporary best practice. At 14 days post-osteotomy, mice

euthanized by cervical dislocation in deep anesthesia. The osteotomized femora were retrieved and

fixed in 4% paraformaldehyde (PFA; Electron Microscopy Sciences, Hatfield, USA) at 4°C for 6–8h. The

femora were then transferred into PBS until ex-vivo µCT was completed.

Ex-vivo µCT

To determine bone formation three-dimensionally, femurs were scanned in a SkyScan 1172 high-resolution µCT (Bruker, Kontich, Belgium). To ensure sufficient stability, the bones were placed within a narrow PBS filled serological pipette and the pins and the external fixator bar were carefully removed. Voxel size was set to 8 µm and the bones were scanned with a source energy of 70 kV, 142 µA, a rotation step of 0.2 degree and an 0.5 mm aluminum filter. Subsequently, scan images were reconstructed using NRecon (Bruker, Kontich, Belgium), applying ring artefact reduction and beam hardening corrections. The CT Analyser software (version 1.20.3.0; both Bruker, Kontich, Belgium) was used for 2D and 3D analyses. By excluding the original cortical bone within the callus, the total volume (TV, mm^3), the total bone volume (BV, mm^3) and the bone volume fraction (BV/TV) of the newly formed bone were analyzed in a manually defined VOI (Bucher et al., 2019).

Histology and immunofluorescence

Following *ex-vivo* µCT, bones were placed in an ascending sugar solution as a cryoprotectant (10%, 20%, 30%) at 4°C for 24h each, then cryo-embedded in SCEM medium (Sectionlab, Japan) and stored at -80°C. Consecutive sections of 7 µm were prepared from the cryo-embedded bones using a cryotome (Leica, Wetzlar, Germany) and cryotapes (Cryofilm 2C(9), Sectionlab, Japan). Sections were fixed onto glass slides, air-dried, and stored at -80 °C until staining.

Movat's pentachrome staining comprised the following steps: briefly, sections were air dried for 15 min, fixed with 4% PFA (Electron Microscopy Sciences, Hatfield, USA) for 30 min at room temperature and washed in distilled water (H_2O_{dest}) for 5 min. Then, sections were pretreated for 3 min in 3% acetic acid (Carl Roth, Karlsruhe, Germany), stained 30 min in 1% alcian blue (Sigma Aldrich, St. Louis, USA) pH 2.5, followed by 10 min washing in 3% acetic acid under light microscopic control. Sections were rinsed in H_2O_{dest} and immersed in alkaline ethanol (Carl Roth, Karlsruhe, Germany) for 60 min, washed in tap water for 10 min, rinsed with H_2O_{dest} and incubated in Weigert's hematoxylin (Waldeck, Münster, Germany) for 15 min. Then, the sections were washed in tap water for 10 min, shortly rinsed with H_2O_{dest} and stained in crocein scarlet-acid fuchsin (Waldeck, Münster, Germany) for 15 min. Afterwards, the slices were treated with 0.5% acetic acid for 1 min, followed by 20 min incubation in 5% phosphotungstic acid (Waldeck, Münster, Germany), and again 1 min in 0.5% acetic acid. The sections were washed three times for 2 min in 100% ethanol, followed by alcoholic Saffron du Gâtinais (Waldeck, Münster, Germany) for 60 min. Finally, the slides were dehydrated in 100% ethanol three times for 2 min and cleared shortly in xylene. Sections were covered with Vitro-Clud (R. Langenbrinck, Emmendingen, Germany) and a cover slip. Imaging was performed on a Leica light microscope using AxioVision software (Carl Zeiss Microscopy, Jena, Germany) at 10x magnification. Quantitative analyses of the Movat's pentachrome staining were evaluated using an ImageJ macro. All analyses were performed blinded to sex, fixation, and pain management protocol.

Immunofluorescence staining was performed as described previously (Lang et al., 2019; Lang et al., 2022) using the following antibody: Endomucin (Emcn) (V.7C7 unconjugated, rat monoclonal, sc-65495, 1:100; Santa Cruz Biotechnology, Dallas, USA), goat anti-rat A647 (1:500; A-21247, polyclonal, Invitrogen, Thermo Fisher Scientific, Waltham, USA) and DAPI (1:1,000; Thermo Fisher Scientific, Waltham, USA). Blocking was performed with 10% FCS/PBS and staining solution contained 5% FCS

with 0.1% Tween20 (Sigma Aldrich, St. Louis, USA). Images were acquired using a Keyence BZ9000
microscope (Keyence, Osaka, Japan). The images were processed and analyzed with ImageJ
(Rasband, 1997-2018; Schneider et al., 2012). An area of interest was established and managed via
the built-in ROI-Manager, while cell number and signal distribution within the area were determined
using the plug-ins Cell-counter and Calculator Plus. Data was processed with the ImageJ plugin
OriginPro.
347

Statistical analysis
Sample size calculation was performed based on own preliminary data (Jirkof et al., 2019) using a
nonparametric ranking procedure to analyze longitudinal data. In detail, the required number of animals
was modeled in R using the package nparLD (Noguchi et al., 2012). The test revealed a 20% difference
with a power of ~80% when the number of animals per group is n=10.
Statistical analysis was performed using RStudio and graphs were created in GraphPad Prism (V8).
The package "nparLD" was used in RStudio (Noguchi et al., 2012). To test whether the data from female
and male mice are homogenous and would allow for an integrated data analysis, we first exemplarily
compared the body weight data of both sexes using the F2-LD-F1 design. Since we found significant
interaction with regard to sex, we therefore analyzed both sexes separately for all statistical analyses.
Thus, nonparametric analysis of longitudinal data (F1-LD-F1 design; further on named as non-
parametric ANOVA-type tests), was further used to test for significant differences in the main effect of
time and main effect of group, separated by sex. When mean group differences *p≤0.05* were detected,
group comparison for each timepoint was performed using the Kruskal-Wallis test (Kruskal and Wallis,
1952). To determine group differences, Dunn's posthoc test with Bonferroni correction was performed
for each timepoint, specifying which means were significant from the others (Dinno, 2015; Dunn, 1961).
Non-parametric ANOVA-type tests as well as *p-values*, chi-squared and df for each analysis of the
following parameters are provided in the Supplementary Information: body weight, food and water
intake, composite pain score, delta composite pain score, limp score, CatWalk (relative velocity, relative
mean intensity, relative stand duration, relative stride length), *ex-vivo* μCT (BV/TV, BV, TV), histology
(relative bone fraction, relative cartilage fraction) and immunofluorescence (Endomucin, DAPI). Mice or
data that were excluded during the study and following analyses are explained in detail in the
Supplementary Information.
371

3 Results

Body weight, food and water intake are affected post-anesthesia and post-osteotomy independent of analgesic regime and fixation

An integrative study design was chosen to i) generate intra-individual controls for the behavioral assessments and ii) reduce animal numbers used in this study (**Fig. 1**). In brief, animals underwent a first intervention consisting of isoflurane anesthesia and administration of the assigned analgesics. Assessments were performed at 12h, 24h, 48h and 72h (referred to in the following as "post-anesthesia"). 14 days after the first intervention, the same animals underwent a second intervention including isoflurane anesthesia, administration of the respective analgesics and osteotomy on the left femur. Same assessments as post-anesthesia were performed at 12h, 24h, 48h and 72h (in the following referred to as "post-osteotomy") (**Fig 1B**).

To assess general indications for wellbeing, the body weight was monitored at 12h, 24h, 48h and 72h post-anesthesia (initial body weight at 0h – males: 25.96 ± 2.1 g; females: 21.41 ± 1.3 g) and post-osteotomy (initial body weight at 0h – males: 27.61 ± 1.9 g; females: 22.75 ± 1.3 g), showing a statistically significant reduction in the range of 5% in all groups at 12h and 24h regardless of sex, analgesic regime, and intervention (**Fig. 2A**). At 48h post-anesthesia, body weight normalized in all groups and even exceeded the pre-intervention weight (**Fig. 2A, B**). Of note, mice did not undergo osteotomy during the first intervention (= post-anesthesia). However, they were already assigned to the respective groups post-osteotomy. After osteotomy, we found that male mice showed prolonged body weight loss over 48h, when compared to post-anesthesia (**Fig. 2A**). In contrast, female mice had less body weight loss at 24h compared to post-anesthesia, but a similar recovery at 48h (**Fig. 2B**). The body weight increase compared to the initial value was higher post-anesthesia than post-osteotomy at 72h in all groups independent of sex, analgesia, and fixation. After both periods of close monitoring, body weight was measured every other day until osteotomy/euthanasia with comparable weight development at any time point (**Fig. S1**).

Food and water intake per cage were assessed at 24h, 48h and 72h post-anesthesia and post-osteotomy. Initial values at 0h covering the previous 24h per cage were as follows: post-anesthesia - males 8.1 ± 0.7 g (food) and 9.2 ± 1.5 ml (water); females 7.8 ± 1.2 g (food) and 9.2 ± 1.2 ml (water); post-osteotomy – males 8.7 ± 1.0 g (food) and 9.9 ± 1.5 ml (water); females 8.4 ± 0.9 g (food) and 9.5 ± 0.9 ml (water). The lowest food and water intake (approximately 50% reduction to initial values) across all groups and sexes was measured 24h after each intervention and reached the level of the initial values at 48h (post-anesthesia) or 72h (post-osteotomy) (**Fig. 2C–F**). No significant main effects were detected between treatment groups (**Table S2, S3**). To rule out any constipating adverse effects of the BUP-Depot, all groups were closely monitored for defecation during the assessments, as constipation is a known-side effect of chronic-opioid usage (Jirkof et al., 2015; Webster et al., 2016). However, a reduction in defecation was only noticeable at 12h after both interventions (**Fig. S2**). The reduction in defecation was considerably more pronounced in female than in male mice but showed no differences between the Tramadol and BUP-Depot groups (**Fig. S2**).

Figure 2. Reduction of body weight as well as food and water intake can be observed at 24h and 48h post-anesthesia and post-osteotomy. (**A, B**) Body weight was measured at 12h, 24h, 48h and 72h; (**C-F**) and food/water intake was measured at 12h, 24h, 48h and 72h post-anesthesia and post-osteotomy and normalized to the initial value (0h = 100%). Of note, mice did not undergo osteotomy during the first intervention (= post-anesthesia). However, they were already assigned to the respective groups post-osteotomy. All graphs show median with interquartile range for n= 9–10 (body weight) and n=4–5 (food/water intake). Non-parametric ANOVA-type test - main effects of time and of group are represented in the graphs; exact *p-values* are listed in Table S1–S3; *$p < 0.05$, ***$p < 0.001$. To determine group differences Kruskal-Wallis test and Dunn's posthoc test with Bonferroni correction was performed. *a* – significant difference Tramadol flexible vs. BUP-Depot flexible; *b* – significant difference Tramadol rigid vs. Tramadol flexible.

Nest building and explorative behaviors are not influenced by the analgesic regime

To detect model independent changes in spontaneous behavior, monitoring of nest complexity scores and explorative behaviors was performed at 24h, 48h and 72h post-anesthesia and post-osteotomy. Analyses showed no differences in the nest building performance between the Tramadol and BUP-Depot groups post-anesthesia and post-osteotomy as medians ranged between scores of 4.5 to 5 (**Fig. 3A**). Explorative behavior was present in all cages with male mice post-anesthesia while reduced in some cages at 24h post-osteotomy (8/20) independent of analgesic regime or fixation. In the female mice, 3 out of 20 cages (Tramadol flexible and BUP-Depot flexible) showed no exploration 24h after both interventions (**Fig. 3B**).

Figure 3. Nest building behavior remains unaffected post-anesthesia and post-osteotomy while exploration was negatively influenced post-osteotomy in male mice at 24h. (**A**) Nest building and (**B**) explorative behavior were monitored per cage at 24h, 48h and 72h post-anesthesia and post-osteotomy. (**A**) Shows median with interquartile range for n=4–5 (nest complexity score) based on cages (pair housing), (**B**) shows n = 5 (explorative behavior) based on cages (pair housing).

Delta composite pain score indicates limited analgesic capacity of Tramadol in male mice with flexible fixation

The composite pain score includes parameters of facial expression and overall appearance was assessed at 24h, 48h and 72h post-anesthesia and post-osteotomy. Since we observed a pronounced influence of the anesthesia and analgesic regime alone (**Fig. S3**; **Table S4–S4.3**), we corrected the

individual score post-osteotomy for the respective score post-anesthesia to obtain a delta composite
pain score, that depicts the isolated effect of the osteotomy without the interfering effects of anesthesia
and analgesia. The delta composite pain score was highest in all groups at 12h and 24h after osteotomy,
and declined after 48h and 72h, reaching lowest scores in all groups after 72h (significant main time
effect in both sexes, *p<0.001;* **Fig. 4A**; **Table S5**). We found a significant higher delta composite pain
score in male mice from the flexible fixation group treated with Tramadol at 24h (median score = 2) and
48h (median score = 1.5) after osteotomy (Kruskal-Wallis test of all groups at 24h: *p= 0.031* and at 48h:
p= <0.001; **Table S5.1**) when compared to the other groups (Dunn's posthoc test for Tramadol flexible
vs. BUP-Depot flexible, BUP-Depot rigid and Tramadol rigid, respectively: 24h *p= 0.077, 0.042, 0.032*;
48h *p= 0.004, <0.001, 0.018* ; **Fig 4A**; **Table S5.2–S5.3**). Female mice showed comparable score
developments between treatment and fixation groups with the highest median scores (1-1.5) at 12h.

453
Figure 4. Delta composite pain score suggests adequate pain alleviation in almost all groups while limping behavior
shows slight differences between groups post-osteotomy. (**A**) The delta composite pain score was evaluated at 12h, 24h,
48h and 72h post-anesthesia and post-osteotomy; to achieve the delta composite score, scores from each individual mouse post-
anesthesia were subtracted from their respective scores post-osteotomy. (**B**) Limp score was assessed at 24h, 48h, 72h and 10d
post-osteotomy. All graphs show median with interquartile range for n = 8–10 (delta composite pain score) and n= 9–10 (limp
score). Non-parametric ANOVA-type test - main effects of time and of group are represented in the graphs; exact *p-values* are
listed in Table S5-S5.3 and Table S6; **p < 0.05, ***p < 0.001.* To determine group differences Kruskal-Wallis test and Dunn's
posthoc test with Bonferroni correction were performed. *a* – significant difference Tramadol rigid vs. Tramadol flexible; *b* –
significant difference Tramadol flexible vs. BUP-Depot rigid; c – significant difference Tramadol flexible vs. BUP-Depot flexible.
463

Limping and gait analysis indicate model-related alterations in walking behavior with only slight differences between groups

Walking behavior was assessed by i) using a metric limping score applied to individual 3 min videos and ii) analyses of gait and locomotion using the Noldus CatWalk XT at 24h, 48h, 72h, and 10d post-osteotomy. The limping score was only considered post-osteotomy as mice did not show any alterations in walking post-anesthesia during routine monitoring. Individual baseline measurements for the CatWalk were taken before both interventions to correct the data post-osteotomy for variances in velocity.

As expected, limping was observed at all timepoints post-osteotomy until 10d with a general improving trend (significant main time effect in males *p<0.001* and females *p= 0.018*; **Fig. 4B**; **Table S6**). In general, median scores ranged between 0 – 1 at 24h and 48h independent of sex and analgesics, while higher variations were seen at 72h (male BUP-Depot rigid; female BUP-Depot flexible; **Fig. 4B**). Slight alterations in limping behavior were still visible after 10d in some individual mice in the female BUP-Depot groups or male Tramadol groups (all = score 1; sporadic limping; up to two animals, respectively). Further analyses of gait and locomotion (CatWalk XT) exhibited a clear reduction in velocity after osteotomy in all groups (significant main time effect post-osteotomy for both sexes *p<0.001;* **Fig. 5A, B**; **Table S7**) with an upward trend surpassing baseline values in nearly all groups at 10d (exception: male, Tramadol, flexible fixation). Significant differences between the male groups were found at 24h and 48h with male mice (flexible fixation group treated with Tramadol) showing a lower relative velocity (non-parametric ANOVA-type test for group *p= 0.034*; Kruskal-Wallis test 24h *p= 0.052* and 48h *p= 0.038*; Dunn's posthoc test – *p<0.05* when compared to BUP-Depot rigid; **Fig. 5A**; **Table S7.1–S7.3**). Moreover, female mice showed significant group differences at 24h (non-parametric ANOVA-type test for group *p= 0.035*; Kruskal-Wallis test 24h *p= 0.009*; **Fig. 5B**; **Table S7.4–S7.5***)* as mice with rigid fixation treated with Tramadol exhibited elevated velocity compared to the other groups which was comparable to their baseline values (median 1.015). With respect to the relative mean intensity (a measure for load bearing) and relative stand duration, we did not observe significant group effects independent of sex and analgesic (**Fig. 5C–F**), while time-dependent reductions at 24h, 48h and 72h improved over time (relative mean intensity: non-parametric ANOVA-type test *p< 0.001* in both sexes; relative stand duration *p< 0.001* in males and *p= 0.005* in females; **Table S8–S9**). As observed before, the relative mean intensity in male mice (flexible fixation group treated with Tramadol) was also numerically reduced (**Fig. 5C**) with no differences in the relative stand duration (**Fig. 5E**). Interestingly, the relative stand duration was only slightly increased in all female mice over 10 days (median range 10d: 0.68 – 0.75; **Fig. 5F**). In addition, the stride length was only markedly modulated over the first 72h and normalized to baseline values at 10d (**Fig. 5G, H**; **Table S10**). Only male mice with flexible fixation and Tramadol treatment showed shortened stride length which was specifically significant at 24h compared to BUP-Depot rigid, but not to Tramadol rigid (non-parametric ANOVA-type test *p= 0.024*; Kruskal-Wallis test 24h *p= 0.044*; Dunn's posthoc test – *p<0.05* when compared to BUP-Depot rigid; **Fig. 5G**; **Table S10–S10.2**). To evaluate the influence of the velocity on mean intensity, stand duration and stride length, we performed additional correlation analyses (spearman correlation) (**Fig. S4A–C**) which indicated that mean intensity (males r= 0.6, females r= 0.34; **Fig. S4A**) and stride length (males r= 0.8, females r= 0.52; **Fig. S4C**) correlate with the velocity (all *p< 0.001*) after osteotomy. Therefore, observations of those parameters need to be discussed in context of the overall velocity.

Figure 5. Results of limping and gait analyses yield model-related alterations in walking behavior which was specifically pronounced in male mice with flexible fixation and Tramadol treatment. CatWalk analysis was conducted at 24h, 48h, 72h and 10d post-osteotomy and normalized to the initial mean baseline value specifically focusing on (**A**) relative velocity, (**B**) relative mean intensity, (**C**) relative stand duration and (**D**) relative stride length. All graphs show median with interquartile range for n= 8–10. Non-parametric ANOVA-type test - main effects of time and of group are represented in the graphs; exact *p-values* are listed in Table S7–S10.2; *$p < 0.05$, ***$p < 0.001$. To determine group differences Kruskal-Wallis test and Dunn's posthoc test with Bonferroni correction were performed. *a* – significant difference Tramadol flexible vs. BUP-Depot rigid; *b* – significant difference Tramadol rigid vs. BUP-Depot flexible.

Fracture healing outcome is affected by fixation stability but not analgesic regime

To evaluate potential effects of the BUP-Depot on fracture healing outcomes, we performed *ex-vivo* μCT, histomorphometric analysis and vessel staining at day 14 post-osteotomy (**Fig. 6**). We observed a numerically lower BV/TV when comparing flexible fixation compared to rigid - except for males treated with Tramadol, however the differences did not reach statistical significance (BV/TV; **Fig. 6A, B**; **Fig. S5**; **Table S11**). The differences were slightly more pronounced in the BUP-Depot groups (male: median rigid = 25.87% vs. median flexible = 17.80%; female: median rigid = 26.12% vs. median flexible = 20.58%) than the Tramadol group (male: median rigid = 21.89% vs. median flexible = 21.52%; female: median rigid = 30.32% vs. median flexible = 23.31%).

Histomorphometric analysis revealed comparable differences between the rigid and flexible fixation in relative bone and cartilage fraction; as expected, bone was reduced while cartilage formation was elevated in the flexible groups compared to rigid fixation in both sexes and treatment groups (**Fig 6C–H**; **Table S12**). Analysis of the vessel formation (Endomucin/Emcn staining) within the callus area did not show significant or obvious differences in the male groups (**Fig. 6I**; **Fig. S5**; **Table S13**). Differences in vessel formation (Emcn) were conditionally statistically significant between fixation in female mice (Kruskal-Wallis test *p= 0.045*; Dunn's posthoc test *p> 0.05* between all groups; **Fig. 6J**; **Table S13.1**) with flexible fixation groups showing less relative Emcn+ areas when compared to the rigid fixation. Additionally, DAPI staining indicated lower cellularity in the female mice with flexible fixation independent of analgesic treatment (**Fig. S5**).

**Figure 6. Analgesic regimes did not negatively affect fracture healing outcome at day 14, while the different fixations
lead to differences in new bone formation.** (**A, B**) Relative bone volume (BV/TV) (%), (**C, D**) relative bone fraction (%) and (**E,
F**) relative cartilage fraction (%). (**G, H**) Exemplary images of the Movat's pentachrome staining: yellow = mineralized bone, green
= cartilage, magenta = bone marrow; scale bar 500 μm. (**I, J**) Immunofluorescence staining of vessel formation (Endomucin)
including quantification and exemplary images; scale bar 200 μm. All graphs show median with interquartile range for n = 8–10
(**A–H**) and n = 6–9 (**I–J**). To determine group differences, Kruskal-Wallis test and Dunn's posthoc test with Bonferroni correction
were performed in RStudio; exact *p-values* are listed in Table S11–S13.1; *p < 0.05*.

4 Discussion

We evaluated the analgesic efficacy and possible side-effects of a newly developed sustained-release Buprenorphine (BUP-Depot) in comparison to an already established protocol (Tramadol in the drinking water) after a pre-operative injection of Buprenorphine in two mouse-osteotomy models with different fixation stiffnesses (Schreiner et al., 2020; Schreiner et al., 2021; Jirkof et al., 2019; Rapp et al., 2015). The BUP-Depot delivered reliable pain relief over 72h post-surgical without side effects on fracture healing outcome, comparable to Tramadol applied with the drinking water. Unexpectedly, we found a possibly limited analgesic capacity of Tramadol in male mice with a more flexible fixation of the osteotomy based on an elevated composite pain score and more altered gait. Due to individual differences in behavioral changes and pain perception (Hohlbaum et al., 2018; Jirkof et al., 2019), we chose a consecutive study design to analyze the effect of anesthesia and analgesia alone and in combination with an osteotomy in the same animal.

General clinical parameters such as body weight, and food and water intake were noticeably reduced post-anesthesia and post-osteotomy. Reduction of food intake and therefore negatively influenced body weight development are known side-effects post-surgical but can also be related to anesthesia or Buprenorphine/Tramadol administration (Brennan et al., 2009; Clark et al., 2004; Goecke et al., 2005; Jirkof et al., 2015). Reduction of body size observed in the present study was very low (in the range of 5% and similar to the anesthesia only control), which is indicative of an excellent pain management and wellbeing in all animals. In none of the animals' values (i.e.20% weight loss) were reached, which would have necessitated immediate termination of the experiment. These findings are as well indicative of a high tolerability of the used drugs. Lowest values in body weight and food intake were reached after 24h in all groups, with an incline over 48h and 72h post-anesthesia and post-osteotomy. Mice were 10 weeks of age during the first intervention (anesthesia and analgesia alone) and 12 weeks at osteotomy. It is known that mice show a rapid body weight development until skeletal maturity between approx. 10-12 weeks which also includes the formation of more body fat (Ferguson et al., 2003; Yang et al., 2014). Since interventions entailing anesthesia results in short-term starvation during the recovery period, it can be speculated that mice at 10 weeks loose body weight more rapidly due to limited body fat reserves when compared to more mature mice (12 weeks) with most probably more body fat. The BUP-Depot showed similar effects on the body weight development as well as food and water intake as the established Tramadol treatment. We did not find differences in water intake between groups, excluding a negative effect of the Tramadol containing water on the overall drinking amount and ensuring a continuous uptake of medication as shown previously (Jirkof et al., 2019; Evangelista Vaz et al., 2018). However, the measurement of the overall 24h water intake does not allow to verify sufficient water intake during the first hours after surgery which might still be reduced as also reported by Evangelista et al. (Evangelista-Vaz et al., 2018). To reduce the chance of insufficient water intake due to altered water taste after surgery and to provide a pre-emptive level of analgesic, we administered Tramadol already one day before surgery (Ehrnthaller et al., 2013; Rapp et al., 2015). To improve the food intake and therefore body weight development, the food can be alternatively provided on the cage floor to prevent the animals from having to stand on their hind legs (Lang et al., 2016). However, this does not allow for precise monitoring of food intake as we did here, as food pellets might be hidden in the nest or be buried.

The use of high-caloric dietary gels could be a valid alternative ensuring the consumption of food and
liquids and can be also used as route for oral analgesic administration (Nizar et al., 2018).

Changes in nest building and the willingness to explore foreign objects can indicate alterations in well-
being in laboratory mice (Gaskill et al., 2013; Jirkof et al., 2013; Jirkof, 2014; Jirkof et al., 2019; Rock et
al., 2014). In our study, nest building behavior was only scarcely influenced by anesthesia or osteotomy
in the BUP-Depot and Tramadol groups independent of sex and fixation stiffness. This is in line with
other studies suggesting that nest complexity scoring might not be perceptive enough to differentiate
between minor pain and other potential stressors or reduced wellbeing after surgery (Jirkof et al., 2013).
However, these findings are in contrary to our previous study where we have reported a reduction in the
nest building performance after osteotomy (Jirkof et al., 2019). Technical differences (more or less
nesting material) or individual differences in scoring might explain the variations in our findings and
underline the necessity for more objective approaches. Moreover, the explorative behavior in male mice
seemed to be negatively impacted by osteotomy, especially at 24h post-surgery, when compared to the
post-anesthesia controls and female mice. This is in accordance with observations reported by Gallo et
al. where they found that male CD1 mice more often failed to pass a Time to Incorporate into Nest Test
(TINT) 3h after surgery (carotid artery catheterization) as compared to their baseline measurements
after anesthesia alone (Gallo et al., 2020). In addition, Hohlbaum et al. showed that female C57BL/6JRj
mice exhibited shorter latency to explore than male mice 1 day after last anesthesia in a repeated
inhalation anesthesia trial indicating sex-specific difference that is in accordance with our findings with
respect to explorative behavior (Hohlbaum et al., 2017).

We used a composite pain score to combine the assessment of facial expression, i.e., the mouse
grimace scale, and clinical appearance post-anesthesia and post-osteotomy. Based on our consecutive
study design, we were able to calculate a delta for each individual mouse representing numerically the
"real" osteotomy effect. In line with other studies, the composite pain score was already slightly impacted
by anesthesia and analgesia alone, indicating that some components of the composite pain score might
not only be influenced by pain but rather also depict stress or discomfort (Jirkof et al., 2019; Miller et al.,
2015). The highest delta composite scores were reached 12h and 24h after osteotomy suggesting the
pain/discomfort peak due to the surgical procedure which constantly declined till 72h post-surgical.
Median delta scores varied around 1 indicating only limited residual pain and/or discomfort over the first
48h. The BUP-Depot provided comparable and sufficient alleviation of pain signs as the Tramadol
treatment and our data indicate that pain relief can even be achieved over 72h. This is an important
advantage as compared to the commercially available sustained-release formulation Buprenorphine
SR-LAB (ZooPharm – USA only), which yields effective serum concentrations over a maximum of 24h
to 48h only with a need for re-administration (Clark et al., 2014; Kendall et al., 2014).

As model-specific parameters, we assessed limping behavior and locomotion using gait analysis. After
osteotomy, limping behavior was observed over 72h post-osteotomy in all groups, irrespective of sex,
fixation, and analgesic, with no significant differences between the groups. More detailed gait analysis
using the CatWalk system revealed reduced velocity and altered gait patterns over 72h and up to 10

days. While velocity and mean intensity ameliorated over 10 days, stand duration and stride length
remained reduced in all groups. Due to the functional impairment, limping as well as changes in gait
parameters after osteotomy are expected as a mixture of residual pain due to local inflammatory
processes and periosteal pain as well as the restricted functionality of the femur. Magnusdottir et al.
reported mechanical hyperalgesia and cold sensitivity in male mice after sham surgery (no external
fixator and osteotomy) and femur fracture over 6 weeks while the sham group significantly improved
after 3 weeks (Magnusdottir et al., 2021). This indicates that the wound healing itself already plays an
important role in potential pain sensitization after osteotomy. A reduction in the stride length over 14
weeks was also observed by Histing et al. (Histing et al., 2010). In addition, limited mobility and changed
gait are also common in human patients with e.g., proximal femur fractures or the femoral diaphysis and
are not directly related to pain (Bizzoca et al., 2020; Wong et al., 2004).
Of note, the sensitive and in-depth analyses of gait pattern and velocity used in the present study (as
opposed to simple limp scores) is important since most gait parameters correlate to velocity (Batka et
al., 2014; Jacobs et al., 2014). Failing to address possible impacts of changes in velocity can affect the
outcome of gait related data (Batka et al., 2014; Cendelín et al., 2010).
664
Unexpectedly, at 24h and 48h, the delta composite pain scores of male mice receiving Tramadol after
osteotomy with a more flexible fixator showed statistically significant higher scores than all other groups.
Together with more pronouncedly altered changes in relative velocity, our data suggest a clinically
relevant degree of discomfort/pain which has not been described previously. One explanation could be
that the effective Tramadol dose was not sufficiently achieved due to the higher body weight of male
mice. Moreover, Evangelista et al. showed that male mice had lower serum concentrations of Tramadol
than female mice when applying Tramadol (0.2 mg/ml) in the drinking water for up to 30h, although the
analgesic effective M1 metabolite was similar between male and female mice (Evangelista Vaz et al.,
2018) which is in line with a previous study performed in rats (Matthiesen et al., 1998). In contrast, other
studies report lower sensitivity of female mice or rats to Tramadol (Dai et al., 2008; Liu et al., 2003).
With respect to the influence of the body weight on the local degree of interfragmentary movement,
Röntgen et al. characterized two configurations of the external fixator, the rigid one (18.1 N/mm) and a
more flexible one (0.82 N/mm) calculating an interfragmentary strain for a 25 g mouse with 2.8% and
61%, respectively (Röntgen et al., 2010). However, when applied to female C57BL/6 mice, they did not
find differences in body weight, ground reaction force and motion over 18 days when treated 3d post-
surgical with 0.025 mg/ml Tramadol in the drinking water (Röntgen et al., 2010). This additionally
highlights potential sex-specific differences in pain perception but also the response to analgesic
medication (Fillingim and Gear, 2004). Sex-specific adaption of pain management regimes might be
advisable in future studies.
684
In terms of fracture healing outcome analyzed by *ex-vivo* µCT and histomorphometry, we found no
differences between the analgesic regimens, indicating safe use of the newly developed BUP-Depot. A
more flexible fixation allows higher interfragmentary movements and, therefore, promotes cartilage
rather than bone formation, as seen in the histomorphometric analysis, as well as formation of a larger
periosteal callus (Röntgen et al., 2010). Staining for endomucin, representing vessels, revealed no

difference between fixations or analgesic in male mice However, female mice with more flexible
stabilized osteotomies showed a reduced Emcn-positive area as well as lower cellularity, regardless of
their analgesic regime. Besides mechanical hinderance of revascularization due to higher strains, the
higher proportion of cartilage observed in female mice with flexible stabilized osteotomies might prevent
revascularization due to the intrinsic anti-angiogenic nature of cartilage e.g., due to chondromodulin-1
(Patra and Sandell, 2012; Shukunami and Hiraki, 2001). This reduced vascularization is in accordance
with earlier observations in sheep (Lienau et al., 2005). Revascularization occurs in a two-waved fashion
during fracture healing (Bahney et al., 2015; Schmidt-Bleek et al., 2015). In the first wave, hypoxia, and
invading immune cells secreting pro-angiogenic factors such as vascular endothelial growth factor and
platelet-derived growth factor, induce angiogenesis, predominantly in the marrow space. The second
wave of revascularization goes along with the conversion of the avascular hypertrophic cartilage to bone.
The different patterns of Emcn staining in males and females might also hint on a more advanced callus
remodeling and therefore a more rapid healing progression in males than females (Haffner-Luntzer et
al., 2021).
704
An effective sustained-release Buprenorphine in Europe would not only be a conceivable alternative to
the application of Tramadol via the drinking water as investigated in this study, but also a potential
alternative to repeated injections of Buprenorphine, which are still more frequently used for pain
management in femur fracture models compared to Tramadol in the drinking water (Wolter et al., 2021).
Assessment of the analgesic effect of Buprenorphine is most often based measurement of plasma or
blood serum levels. Studies specified therapeutic effective concentrations of Buprenorphine in plasma
at a threshold of around 1 ng/ml or 1 ng/g in mice and rats (Yassen et al., 2005; Yun et al., 2010;
Guarnieri et al., 2012). However, Buprenorphine works through the μ-, κ- and δ-opioid receptors in the
brain (Villiger and Taylor, 1981; Lutfy and Cowan, 2004) and it can be hypothesized that reliable pain
alleviation is more reliant on specific binding concentration values in the brain than specific plasma
concentrations (Schreiner et al., 2020). Ohtani et al. also demonstrated a correlation between analgesic
effects and specific binding concentrations of Buprenorphine in the brain of rats (Ohtani et al., 1995).
Schreiner et al. contemplated that specific binding concentrations of 5 ng/g in the brain might be needed
for reliable pain relieve in mice, as observed 24h after injection of the BUP-Depot. Binding
concentrations of less than 3 ng/g at 48h injection (BUP-Depot) – which is comparable to levels observed
12h after Temgesic injection – still resulted in high, but not significantly increased withdrawal latencies
compared to a single injection with Temgesic or NaCl. Schreiner et al. therefore suggests that alleviation
of strong pain through the BUP-Depot might require the administration every 24h (Schreiner et al.,
2020). However, based on our assessment of the clinical, behavioral, and model-specific parameters,
we can postulate that the analgesic properties of the BUP-Depot are sufficient for 72h post-operative
analgesia in our specific mouse-osteotomy model of moderate severity. Nonetheless, the potential use
of the BUP-Depot in other, more painful models still needs to be critically assessed and evaluated.
727
Generally, the use of a sustained-release Buprenorphine results in less handling induced stress and
prevents potential gaps in analgesic coverage due to infrequent application intervals when injected
repeatedly. Drug administration via the drinking water always bears the risk of insufficient water intake

after anesthesia/surgery and due to circadian rhythm (Sauer et al., 2016). Therefore, adequate pain
management requires a constant reevaluation of the chosen protocol as well as the consideration of
strain, sex, procedure, interindividual differences in animals, and more, and the use of the most suitable
analgesic protocol must be separately evaluated and regularly reevaluated for each animal experiment.
As not only sex, but also genetic heritability influences experienced pain and response to analgesia in
mice (Smith, 2019), further studies using different strains are needed in the evaluation of commonly
used analgesic regimens in laboratory rodents. However, it needs to be considered that animals were
kept in IVC cages, leading to a naïve immune system that could not maturate accordingly, potentially
leading to different bone healing properties under different housing conditions that could not be
assessed within this study (Bucher et al., 2019).
741
Taken together, our assessment of clinical, behavioral, and model-specific parameters suggest that the
analgesic properties of the BUP-Depot were sufficient for 72h post-operative analgesia in male and
female C57BL/6J mice after femoral osteotomy. Therefore, the BUP-depot provides an excellent
alternative for extended pain relief in preclinical studies. There is no question that the availability of such
a sustained-release formulation of Buprenorphine in Europe would be beneficial for pain management
in preclinical studies.
748

Conflict of interest statement

The authors declare that they have no conflict of interests.

Acknowledgement

We like to thank Kristina Ullmann, Bianka Verrett and Sylvia Gundelach for their support during the
planning and performance of the animal experiments. Special thanks to Darryl Borland and PD Dr.
Maxim Puchkov for expert technical support and the preparation of BUP-Depot. This study was funded
by Charite3R, Charité-Universitätsmedizin Berlin (RefineMOMo 2.0; RefineLab) and a grant from
Interpharma Basel.

749 **Data availability statement**

The authors declare that all data supporting the findings of this study are available within the paper and
its Supplementary Information file. Further information is made available by the authors upon request.

References
Bahney, C. S., Hu, D. P., Miclau, T. et al. (2015). The multifaceted role of the vasculature in
endochondral fracture repair. *Frontiers in Endocrinology 6*, doi:10.3389/fendo.2015.00004
Balcombe, J. P., Barnard, N. D. and Sandusky, C. (2004). Laboratory routines cause animal stress.
Contemp Top Lab Anim Sci 43, 42-51.
Batka, R. J., Brown, T. J., McMillan, K. P. et al. (2014). The need for speed in rodent locomotion
analyses. *Anat Rec (Hoboken) 297*, 1839-1864. doi:10.1002/ar.22955
Bizzoca, D., Bortone, I., Vicenti, G. et al. (2020). Gait analysis in the postoperative assessment of
intertrochanteric femur fractures. *J Biol Regul Homeost Agents 34*, 345-351. Congress of the
Italian Orthopaedic Research Society.
Bove, S. E., Flatters, S. J., Inglis, J. J. et al. (2009). New advances in musculoskeletal pain. *Brain Res
Rev 60*, 187-201. doi:10.1016/j.brainresrev.2008.12.012

Brennan, M. P., Sinusas, A. J., Horvath, T. L. et al. (2009). Correlation between body weight changes
and postoperative pain in rats treated with meloxicam or buprenorphine. *Lab Anim (NY) 38*, 87-
93. doi:10.1038/laban0309-87
Bucher, C. H., Schlundt, C., Wulsten, D. et al. (2019). Experience in the adaptive immunity impacts bone
homeostasis, remodeling, and healing. *Front Immunol 10*, 797. doi:10.3389/fimmu.2019.00797
Bullingham, R. E., McQuay, H. J., Moore, A. et al. (1980). Buprenorphine kinetics. *Clin Pharmacol Ther*
28, 667-672. doi:10.1038/clpt.1980.219
Carbone, L. and Austin, J. (2016). Pain and laboratory animals: Publication practices for better data
reproducibility and better animal welfare. *PLoS One 11*, e0155001.
doi:10.1371/journal.pone.0155001
Carstens, E. and Moberg, G. P. (2000). Recognizing pain and distress in laboratory animals. *Ilar j 41*,
62-71. doi:10.1093/ilar.41.2.62
Cendelín, J., Voller, J. and Vožeh, F. (2010). Ataxic gait analysis in a mouse model of the olivocerebellar
degeneration. *Behavioural Brain Research 210*, 8-15.
doi:https://doi.org/10.1016/j.bbr.2010.01.035
Clark, M. D., Krugner-Higby, L., Smith, L. J. et al. (2004). Evaluation of liposome-encapsulated
oxymorphone hydrochloride in mice after splenectomy. *Comp Med 54*, 558-563.
Clark, T. S., Clark, D. D. and Hoyt, R. F., Jr. (2014). Pharmacokinetic comparison of sustained-release
and standard buprenorphine in mice. *Journal of the American Association for Laboratory Animal*
Science : JAALAS 53, 387-391.
Cottrell, J. and O'Connor, J. P. (2010). Effect of non-steroidal anti-inflammatory drugs on bone healing.
Pharmaceuticals (Basel) 3, 1668-1693. doi:10.3390/ph3051668
Dai, X., Brunson, C. D., Rockhold, R. W. et al. (2008). Gender differences in the antinociceptive effect
of tramadol, alone or in combination with gabapentin, in mice. *J Biomed Sci 15*, 645-651.
doi:10.1007/s11373-008-9252-0
Dinno, A. (2015). Nonparametric pairwise multiple comparisons in independent groups using dunn's
test. *The Stata Journal 15*, 292-300. doi:10.1177/1536867x1501500117
Dunn, O. J. (1961). Multiple comparisons among means. *Journal of the American statistical association*
56, 52-64.
Ehrnthaller, C., Huber-Lang, M., Nilsson, P. et al. (2013). Complement c3 and c5 deficiency affects
fracture healing. *PLoS One 8*, e81341. doi:10.1371/journal.pone.0081341
Evangelista-Vaz, R., Bergadano, A., Arras, M. et al. (2018). Analgesic efficacy of subcutaneous-oral
dosage of tramadol after surgery in c57bl/6j mice. *J Am Assoc Lab Anim Sci 57*, 368-375.
doi:10.30802/AALAS-JAALAS-17-000118
Evangelista Vaz, R., Draganov, D. I., Rapp, C. et al. (2018). Preliminary pharmacokinetics of tramadol
hydrochloride after administration via different routes in male and female b6 mice. *Vet Anaesth*
Analg 45, 111-122. doi:10.1016/j.vaa.2016.09.007
Ferguson, V. L., Ayers, R. A., Bateman, T. A. et al. (2003). Bone development and age-related bone
loss in male c57bl/6j mice. *Bone 33*, 387-398. doi:https://doi.org/10.1016/S8756-
3282(03)00199-6
Fillingim, R. B. and Gear, R. W. (2004). Sex differences in opioid analgesia: Clinical and experimental
findings. *Eur J Pain 8*, 413-425. doi:10.1016/j.ejpain.2004.01.007
Flecknell, P. (2018). Rodent analgesia: Assessment and therapeutics. *Vet J 232*, 70-77.
doi:10.1016/j.tvjl.2017.12.017
Gades, N. M., Danneman, P. J., Wixson, S. K. et al. (2000). The magnitude and duration of the analgesic
effect of morphine, butorphanol, and buprenorphine in rats and mice. *Contemp Top Lab Anim*
Sci 39, 8-13.
Gallo, M. S., Karas, A. Z., Pritchett-Corning, K. et al. (2020). Tell-tale tint: Does the time to incorporate
into nest test evaluate postsurgical pain or welfare in mice? *Journal of the American Association*
for Laboratory Animal Science : JAALAS 59, 37-45. doi:10.30802/AALAS-JAALAS-19-000044
Gaskill, B. N., Karas, A. Z., Garner, J. P. et al. (2013). Nest building as an indicator of health and welfare
in laboratory mice. *J Vis Exp* 51012. doi:10.3791/51012
Gerstenfeld, L. C., Thiede, M., Seibert, K. et al. (2003). Differential inhibition of fracture healing by non-
selective and cyclooxygenase-2 selective non-steroidal anti-inflammatory drugs. *Journal of*
Orthopaedic Research 21, 670-675. doi:https://doi.org/10.1016/S0736-0266(03)00003-2
Girón, R., Abalo, R., Goicoechea, C. et al. (2002). Synthesis and opioid activity of new fentanyl analogs.
Life Sci 71, 1023-1034. doi:10.1016/s0024-3205(02)01798-8
Goecke, J. C., Awad, H., Lawson, J. C. et al. (2005). Evaluating postoperative analgesics in mice using
telemetry. *Comp Med 55*, 37-44.
Guarnieri, M., Brayton, C., DeTolla, L. et al. (2012). Safety and efficacy of buprenorphine for analgesia
in laboratory mice and rats. *Lab Anim (NY) 41*, 337-343. doi:10.1038/laban.152

Guo, S. and Dipietro, L. A. (2010). Factors affecting wound healing. *J Dent Res 89*, 219-229.
doi:10.1177/0022034509359125
Haffner-Luntzer, M., Fischer, V. and Ignatius, A. (2021). Differences in fracture healing between female
and male c57bl/6j mice. *Frontiers in Physiology 12*, doi:10.3389/fphys.2021.712494
Hess, S. E., Rohr, S., Dufour, B. D. et al. (2008). Home improvement: C57bl/6j mice given more
naturalistic nesting materials build better nests. *J Am Assoc Lab Anim Sci 47*, 25-31.
Histing, T., Kristen, A., Roth, C. et al. (2010). In vivo gait analysis in a mouse femur fracture model.
Journal of Biomechanics 43, 3240-3243. doi:https://doi.org/10.1016/j.jbiomech.2010.07.019
Histing, T., Andonyan, A., Klein, M. et al. (2016). Obesity does not affect the healing of femur fractures
in mice. *Injury 47*, 1435-1444. doi:10.1016/j.injury.2016.04.030
Hohlbaum, K., Bert, B., Dietze, S. et al. (2017). Severity classification of repeated isoflurane anesthesia
in c57bl/6jrj mice-assessing the degree of distress. *PloS one 12*, e0179588-e0179588.
doi:10.1371/journal.pone.0179588
Hohlbaum, K., Bert, B., Dietze, S. et al. (2018). Impact of repeated anesthesia with ketamine and
xylazine on the well-being of c57bl/6jrj mice. *PLoS One 13*, e0203559.
doi:10.1371/journal.pone.0203559
Hohlbaum, K., Frahm, S., Rex, A. et al. (2020). Social enrichment by separated pair housing of male
c57bl/6jrj mice. *Sci Rep 10*, 11165. doi:10.1038/s41598-020-67902-w
Jacobs, B. Y., Kloefkorn, H. E. and Allen, K. D. (2014). Gait analysis methods for rodent models of
osteoarthritis. *Curr Pain Headache Rep 18*, 456. doi:10.1007/s11916-014-0456-x
Jirkof, P., Fleischmann, T., Cesarovic, N. et al. (2013). Assessment of postsurgical distress and pain in
laboratory mice by nest complexity scoring. *Lab Anim 47*, 153-161.
doi:10.1177/0023677213475603
Jirkof, P. (2014). Burrowing and nest building behavior as indicators of well-being in mice. *J Neurosci
Methods 234*, 139-146. doi:10.1016/j.jneumeth.2014.02.001
Jirkof, P., Tourvieille, A., Cinelli, P. et al. (2015). Buprenorphine for pain relief in mice: Repeated
injections vs sustained-release depot formulation. *Lab Anim 49*, 177-187.
doi:10.1177/0023677214562849
Jirkof, P. (2017). Side effects of pain and analgesia in animal experimentation. *Lab Anim (NY) 46*, 123-
128. doi:10.1038/laban.1216
Jirkof, P., Durst, M., Klopfleisch, R. et al. (2019). Administration of tramadol or buprenorphine via the
drinking water for post-operative analgesia in a mouse-osteotomy model. *Sci Rep 9*, 10749.
doi:10.1038/s41598-019-47186-5
Kendall, L. V., Hansen, R. J., Dorsey, K. et al. (2014). Pharmacokinetics of sustained-release analgesics
in mice. *J Am Assoc Lab Anim Sci 53*, 478-484.
Kest, B., Sarton, E. and Dahan, A. (2000). Gender differences in opioid-mediated analgesia: Animal and
human studies. *Anesthesiology 93*, 539-547. doi:10.1097/00000542-200008000-00034
Kruskal, W. H. and Wallis, W. A. (1952). Use of ranks in one-criterion variance analysis. *Journal of the
American Statistical Association 47*, 583-621. doi:10.1080/01621459.1952.10483441
Lang, A., Schulz, A., Ellinghaus, A. et al. (2016). Osteotomy models - the current status on pain scoring
and management in small rodents. *Lab Anim 50*, 433-441. doi:10.1177/0023677216675007
Lang, A., Kirchner, M., Stefanowski, J. et al. (2019). Collagen i-based scaffolds negatively impact
fracture healing in a mouse-osteotomy-model although used routinely in research and clinical
application. *Acta Biomater 86*, 171-184. doi:10.1016/j.actbio.2018.12.043
Lang, A., Stefanowski, J., Pfeiffenberger, M. et al. (2022). Mif does only marginally enhance the pro-
regenerative capacities of dfo in a mouse-osteotomy-model of compromised bone healing
conditions. *Bone 154*, 116247. doi:10.1016/j.bone.2021.116247
Langford, D. J., Bailey, A. L., Chanda, M. L. et al. (2010). Coding of facial expressions of pain in the
laboratory mouse. *Nat Methods 7*, 447-449. doi:10.1038/nmeth.1455
Lienau, J., Schell, H., Duda, G. N. et al. (2005). Initial vascularization and tissue differentiation are
influenced by fixation stability. *Journal of Orthopaedic Research 23*, 639-645.
doi:https://doi.org/10.1016/j.orthres.2004.09.006
Liu, H. C., Jin, S. M. and Wang, Y. L. (2003). Gender-related differences in pharmacokinetics of
enantiomers of trans-tramadol and its active metabolite, trans-o-demethyltramadol, in rats. *Acta
Pharmacol Sin 24*, 1265-1269.
Lutfy, K. and Cowan, A. (2004). Buprenorphine: A unique drug with complex pharmacology. *Curr
Neuropharmacol 2*, 395-402. doi:10.2174/1570159043359477
Magnusdottir, R., Gohin, S., ter Heegde, F. et al. (2021). Fracture-induced pain-like behaviours in a
femoral fracture mouse model. *Osteoporosis International 32*, 2347-2359. doi:10.1007/s00198-
021-05991-7
Matthiesen, T., Wöhrmann, T., Coogan, T. P. et al. (1998). The experimental toxicology of tramadol: An
overview. *Toxicol Lett 95*, 63-71. doi:10.1016/s0378-4274(98)00023-x

Miller, A., Kitson, G., Skalkoyannis, B. et al. (2015). The effect of isoflurane anaesthesia and
buprenorphine on the mouse grimace scale and behaviour in cba and dba/2 mice. *Applied*
animal behaviour science 172, 58-62. doi:10.1016/j.applanim.2015.08.038
Nizar, J. M., Bouby, N., Bankir, L. et al. (2018). Improved protocols for the study of urinary electrolyte
excretion and blood pressure in rodents: Use of gel food and stepwise changes in diet
composition. *American Journal of Physiology-Renal Physiology 314*, F1129-F1137.
doi:10.1152/ajprenal.00474.2017
Noguchi, K., Gel, Y. R., Brunner, E. et al. (2012). Nparld: An r software package for the nonparametric
analysis of longitudinal data in factorial experiments. *Journal of Statistical Software 50*, 1 - 23.
doi:10.18637/jss.v050.i12
Ohtani, M., Kotaki, H., Sawada, Y. et al. (1995). Comparative analysis of buprenorphine- and
norbuprenorphine-induced analgesic effects based on pharmacokinetic-pharmacodynamic
modeling. *J Pharmacol Exp Ther 272*, 505-510.
Page, G. G. (2003). The immune-suppressive effects of pain. *Adv Exp Med Biol 521*, 117-125.
Patra, D. and Sandell, L. J. (2012). Antiangiogenic and anticancer molecules in cartilage. *Expert*
Reviews in Molecular Medicine 14, e10. doi:10.1017/erm.2012.3
Peterson, N. C., Nunamaker, E. A. and Turner, P. V. (2017). To treat or not to treat: The effects of pain
on experimental parameters. *Comp Med 67*, 469-482.
Rapp, A. E., Kroner, J., Baur, S. et al. (2015). Analgesia via blockade of ngf/trka signaling does not
influence fracture healing in mice. *J Orthop Res 33*, 1235-1241. doi:10.1002/jor.22892
Rapp, A. E., Hachemi, Y., Kemmler, J. et al. (2018). Induced global deletion of glucocorticoid receptor
impairs fracture healing. *Faseb j 32*, 2235-2245. doi:10.1096/fj.201700459RR
Rock, M. L., Karas, A. Z., Rodriguez, K. B. et al. (2014). The time-to-integrate-to-nest test as an indicator
of wellbeing in laboratory mice. *J Am Assoc Lab Anim Sci 53*, 24-28.
Röntgen, V., Blakytny, R., Matthys, R. et al. (2010). Fracture healing in mice under controlled rigid and
flexible conditions using an adjustable external fixator. *J Orthop Res 28*, 1456-1462.
doi:10.1002/jor.21148
Roughan, J. V. and Flecknell, P. A. (2002). Buprenorphine: A reappraisal of its antinociceptive effects
and therapeutic use in alleviating post-operative pain in animals. *Lab Anim 36*, 322-343.
doi:10.1258/002367702320162423
Sauer, M., Fleischmann, T., Lipiski, M. et al. (2016). Buprenorphine via drinking water and combined
oral-injection protocols for pain relief in mice. *Applied Animal Behaviour Science 185*, 103-112.
doi:10.1016/j.applanim.2016.09.009
Schmidt-Bleek, K., Kwee, B. J., Mooney, D. J. et al. (2015). Boon and bane of inflammation in bone
tissue regeneration and its link with angiogenesis. *Tissue Engineering Part B: Reviews 21*, 354-
364. doi:10.1089/ten.teb.2014.0677
Schneider, C. A., Rasband, W. S. and Eliceiri, K. W. (2012). Nih image to imagej: 25 years of image
analysis. *Nature Methods 9*, 671-675. doi:10.1038/nmeth.2089
Schreiner, V., Durst, M., Arras, M. et al. (2020). Design and in vivo evaluation of a microparticulate depot
formulation of buprenorphine for veterinary use. *Sci Rep 10*, 17295. doi:10.1038/s41598-020-
74230-6
Schreiner, V., Detampel, P., Jirkof, P. et al. (2021). Buprenorphine loaded plga microparticles:
Characterization of a sustained-release formulation. *Journal of Drug Delivery Science and*
Technology 63, doi:10.1016/j.jddst.2021.102558
Shukunami, C. and Hiraki, Y. (2001). Role of cartilage-derived anti-angiogenic factor, chondromodulin-
i, during endochondral bone formation. *Osteoarthritis Cartilage 9 Suppl A*, S91-101.
doi:10.1053/joca.2001.0450
Smith, J. C. (2019). A review of strain and sex differences in response to pain and analgesia in mice.
Comp Med 69, 490-500. doi:10.30802/AALAS-CM-19-000066
Tappe-Theodor, A., Pitzer, C., Lewejohann, L. et al. (2022). The "wwhow" concept for prospective
categorization of post-operative severity assessment in mice and rats. *Front Vet Sci 9*, 841431.
doi:10.3389/fvets.2022.841431
Villiger, J. W. and Taylor, K. M. (1981). Buprenorphine : Characteristics of binding sites in the rat central
nervous system. *Life Sci 29*, 2699-2708. doi:10.1016/0024-3205(81)90529-4
Webster, L. R., Camilleri, M. and Finn, A. (2016). Opioid-induced constipation: Rationale for the role of
norbuprenorphine in buprenorphine-treated individuals. *Subst Abuse Rehabil 7*, 81-86.
doi:10.2147/SAR.S100998
Wolter, A., Rapp, A. E., Durst, M. S. et al. (2021). Systematic review on the reporting accuracy of
experimental details in publications using mouse femoral fracture models. *Bone 152*, 116088.
doi:10.1016/j.bone.2021.116088

Wong, J., Boyd, R., Keenan, N. W. et al. (2004). Gait patterns after fracture of the femoral shaft in
957   children, managed by external fixation or early hip spica cast. *J Pediatr Orthop 24*, 463-471.
958   doi:10.1097/00004694-200409000-00003
959   Yang, Y., Smith, D. L., Jr., Keating, K. D. et al. (2014). Variations in body weight, food intake and body
960   composition after long-term high-fat diet feeding in c57bl/6j mice. *Obesity (Silver Spring, Md.)*
961   *22*, 2147-2155. doi:10.1002/oby.20811
962   Yassen, A., Olofsen, E., Dahan, A. et al. (2005). Pharmacokinetic-pharmacodynamic modeling of the
963   antinociceptive effect of buprenorphine and fentanyl in rats: Role of receptor equilibration
964   kinetics. *J Pharmacol Exp Ther 313*, 1136-1149. doi:10.1124/jpet.104.082560
965   Yu, S., Zhang, X., Sun, Y. et al. (2006). Pharmacokinetics of buprenorphine after intravenous
966   administration in the mouse. *J Am Assoc Lab Anim Sci 45*, 12-16.
967   Yun, M.-H., Jeong, S.-W., Pai, C.-M. et al. (2010). Pharmacokinetic-pharmacodynamic modeling of the
968   analgesic effect of bupredermtm, in mice. *Health 02*, 824-831. doi:10.4236/health.2010.28124
969