Cancer subclone detection based on DNA copy number in single cell and spatial omic sequencing data

Chi-Yun Wu[1,2,3], Anuja Sathe[4], Jiazhen Rong[1,2], Paul R. Hess[2,5], Billy T. Lau[4,6], Susan M. Grimes[6], Hanlee P. Ji[4,6], Nancy R. Zhang[1,2]

Institutions

[1]Graduate Group in Genomics and Computational Biology, University of Pennsylvania, Philadelphia, PA

[2]Department of Statistics, University of Pennsylvania, Philadelphia, PA

[3]Gladstone Institutes, San Francisco, CA

[4]Division of Oncology, Department of Medicine, Stanford University School of Medicine, Stanford, CA

[5]Department of Pathology and Laboratory Medicine, University of Pennsylvania, Philadelphia, PA

[6]Stanford Genome Technology Center, Stanford University, Palo Alto, CA

Corresponding authors

Nancy R. Zhang

Email: nzh@wharton.upenn.edu

Abstract

In cancer, somatic mutations such as copy number alterations (CNAs) accumulate during disease progression and lead to functional intra-tumor heterogeneity that can influence the efficacy of cancer therapy. Therefore, studying the functional characteristics and spatial distribution of genetically distinct subclones is crucial to the understanding of tumor evolution and the design of cancer treatment. Here, we present Clonalscope, a method for subclone detection using copy number profiles that can be applied to spatial transcriptomics (ST) data and data from single-cell sequencing platforms such as scRNA-seq and scATAC-seq. Clonalscope implements a nested Chinese restaurant process to identify *de novo* subclones within one or multiple samples from the same patient. Clonalscope incorporates prior information from paired whole-genome or whole-exome sequencing (WGS/WES) data to achieve more reliable subclone detection and malignant cell labeling. On scRNA-seq and scATAC-seq data from four gastrointestinal tumor samples, Clonalscope successfully labeled malignant cells and identified genetically different subclones, which were validated in detail using matched scDNA-seq data. On ST data from a squamous cell carcinoma and two invasive ductal carcinoma samples, Clonalscope successfully labelled malignant spots, traced subclones between associated datasets, and identified spatially segregated subclones expressing genes associated with drug resistance and survival.

Introduction

Cancer is a disease of intra-tissue clonal evolution, where DNA mutations combine with epigenetic alterations to confer selective advantages on neoplastic cells, allowing them to proliferate, invade, metastasize, and resist therapeutic treatment [1, 2]. Detection and characterization of genetically different subclones is crucial for the study of cancer and for developing effective treatments and prognostic strategies. Copy number alteration (CNA), one type of genomic variation, is a prevalent manifestation of genome instability and a key driver of tumor evolution [3-5]. Subclones distinguished by CNA have been successfully profiled using bulk and single-cell DNA sequencing data [6-14]. CNA-based subclone detection in other types of single-cell omic data, such as single-cell RNA sequencing (scRNA-seq) [15-19] and single-cell transposase-accessible chromatin-sequencing (scATAC-seq) [20, 21] further allows the joint analysis of genetic mutations with gene expression and chromatin state. Recently, the maturation of spatial transcriptomic (ST) technologies [22, 23] is heralding new opportunities for the study of spatial subclonal dynamics through the mapping of DNA alterations of cells in situ [24, 25].

There is a need for reliable methods for CNA estimation and subclone detection for scRNA-seq, scATAC-seq and ST data. Since these methods do not profile DNA, read coverage along the chromosome is, at best, a noisy and indirect proxy of the underlying DNA copy number. Existing methods for scRNA-seq data rely on smoothing across adjacent genes, e.g. inferCNV [16] and CopyKAT [15]. Similar strategies were also used in CNA analysis for scATAC-seq data [21]. Other methods such as HoneyBADGER [18] and CaSpER [17] consider allelic expression in their copy number detection model, but the sparsity of allele-specific reads limits the improvement achieved by these approaches.

For spatial transcriptomics data, there is yet no vetted published method for CNA-based

subclone detection. STARCH, developed for CNA detection for ST data, leverages the

signals from nearby spots to overcome sparsity [25]. Since scRNA-seq and ST both

measure gene expression, some emerging studies apply inferCNV, developed for

scRNA-seq data, to analyze CNA in tumor ST data [24].

The above methods partition cells/spots into clusters with distinct

transcriptomic/epigenetic profiles, but to what degree do the clusters represent subclones

with distinct DNA-level copy number profiles? Since broad megabase-level shifts in

accessibility/transcription have been observed between normal cells of different lineages,

and may reflect chromatin-level remodeling instead of copy number, RNA-based CNA

detection can be prone to false-positives. In contrast, CONICS [19] profiles CNAs in scRNA-

seq data with segmentation obtained from matched bulk DNA-seq data to reduce false

signals, but it was mainly developed for the full-length scRNA-seq data and does not infer

intratumor subclones. Like the above approaches, CONICS relies on hierarchical

clustering to cluster cells into subclones. Hierarchical clustering works when there are

clear, well-separated clusters in the data, and breaks down when the signal-to-noise ratio

is low as in the case of copy number estimates from scRNA-seq, scATAC-seq, and ST

data. Hierarchical clustering does not leverage the prior information in matched bulk DNA

sequencing data, which is often available, nor does it leverage the specific structure of

subclonal copy number changes that could be used to distinguish signal from noise.

We present Clonalscope, a method for CNA-based subclone detection that can be

applied to diverse single cell and spatial sequencing data types, such as scRNA-seq,

scATAC-seq, and spatial transcriptomics data. Clonalscope is based on nonparametric

Bayesian clustering via a novel nested Chinese restaurant process, which exploits the

sparse nature of subclonal CNA differences to identify *de novo* subclones with related,

but distinct, copy number profiles. Importantly, Clonalscope allows the leveraging of prior

information from paired whole-genome or whole-exome sequencing (WGS/WES) data to

achieve more reliable subclone detection and malignant cell labeling. When multiple

samples are sequenced for a patient, Clonalscope also allows the linking of subclones

across samples for lineage tracing.

We first consider CNA estimation, malignant cell labeling and cancer subclone

detection in scRNA-seq data, where tissue-matched bulk and single-cell DNA sequencing

data are used for benchmarking. We show that CNA estimates derived from scRNA-seq

data alone, without leveraging matched bulk DNA-seq data, can be inaccurate, and that

the errors propagate downstream to affect such common tasks as malignant cell

identification. Leveraging WGS/WES data, Clonalscope obtains precise copy number

estimation and significantly improves the accuracy of malignant cell labeling and subclone

detection, as compared to existing approaches. Next, we apply Clonalscope to scATAC-

seq data from a gastric cancer cell line with matched scDNA-seq data for validation, and

show that Clonalscope can identify subclones based on allele-specific copy number

profiles. Finally, on ST data from breast, colorectal, and skin cancer, with expert pathology

annotation as validation, we show that Clonalscope successfully distinguishes malignant

from non-malignant cells and achieves reproducible spatial subclone mapping across

adjacent tissue slices.

Results

Overview of the Clonalscope model and algorithm (Fig. 1)

Clonalscope starts by segmenting the genome into regions of homogeneous copy

number states in matched bulk whole-genome sequencing (WGS) or whole-exome

sequencing (WES) data (Step. 1). Next, in single-cell RNA/ATAC sequencing (scRNA-

seq/scATAC-seq) data, the relative coverage fold change for each cell i in each region r

is estimated under a Poisson model with the baseline coverage computed from a set of

pre-chosen diploid cells, such as immune cells (Step. 2; Methods). This reduces the cell-

by-gene matrix to a cell-by-region matrix, which serves as input to the non-parametric

Bayesian clustering model for subclone detection and subclonal copy number inference

(Step. 3). The non-parametric Bayesian clustering model is initialized to two clusters: a

cluster representing normal cells with fold change initialized to 1 across all regions, and

a cluster representing the major tumor clone with fold change in each region initialized to

its fold change in the matched bulk DNA sequencing (DNA-seq) data. Then, Clonalscope

applies a nested Chinese Restaurant Process to iteratively add/select copy number states

for each genome segment, seed new subclones, and assign each cell to a subclone. We

use the term "nested" since the algorithm involves two Chinese Restaurant Processes:

The first Chinese Restaurant Process samples subclone membership and seeds new

subclones, while for each new cancer subclone, a second Chinese Restaurant Process

samples the copy number of each region, seeding new copy number states as needed.

Clonalscope outputs the estimated numbers of subclones, the copy number profiles for

the subclones, and the subclone assignment of each cell.

For spatial transcriptomic (ST) data, the same algorithm as described above applies, with

"cell" replaced by "spot". We do not make use of the spatial coordinates in the clustering,

and we don't smooth the cluster memberships across space. This is because we do not

want to make a priori assumptions on the degree of intermixing of subclones in space.

Clonalscope can also make use of allele-specific coverage, if such data is available.

As above, we start with a segmentation of the genome derived from bulk DNA-seq data,

and then use Alleloscope [9] to estimate both the relative fold change and allelic ratio for

each cell i in each region r. The non-parametric Bayesian clustering then considers both

relative fold change and allelic ratio in computing the likelihood of a cell given a copy

number profile. See Methods for details.

Figure 1.

Figure 1: Overview of subclone detection with Clonalscope.

Step 1. The algorithm starts from a segmentation of the genome based on matched WGS or WES data. Step 2. For each segment, Clonalscope computes the coverage change of each cell in scRNA-seq data by normalizing to a set of normal cells. This step reduces the raw gene-by-cell matrix for the UMI counts to the segment-by-cell matrix. Step 3. Clonalscope implements a nested Chinese Restaurant Process to iteratively assign each cell into one of the existing clusters, add or select copy number states for each segment (1st Chinese Restaurant Process), and seeds new clusters (2nd Chinese Restaurant Process) (Methods). For ST data, the setting is similar but "cell" is substituted with "spot". Optionally, Clonalscope can also operate on both coverages and allelic ratios for subclone detection on scATAC-seq data. In addition to the segment-by-cell matrix for fold changes, Alleloscope can be used to retrieve the segment-by-cell matrix for allelic ratios. Clonalscope then detects subclones based on both types of information.

Subclone detection in scRNA-seq data validated by matched scDNA-seq data

Most existing copy number aberration (CNAs) estimation methods for scRNA-seq

data do not allow integration with bulk WGS data, and rely solely on a

smoothing/averaging of the RNA expression data for copy number estimation. Then,

these methods usually apply hierarchical clustering on the estimated single cell copy

number profiles for subclone detection or malignant cell labeling. There can be

considerable differences between RNA expression and DNA copy number [26], even when

the former is smoothed across large contiguous regions [27]. To examine how well scRNA-

seq data can recapitulate the underlying *DNA-level* heterogeneity, we consider scRNA-

seq data from three primary gastrointestinal cancer samples (P5847, P5931, and P8823)

and one colorectal metastasis (P6198). For P5847 and P5931, scDNA-seq data is

available for validation, and for P6198 and P8823, bulk DNA-seq is available for

assessment of concordance of mean copy number profiles. We benchmarked

Clonalscope against inferCNV and CopyKAT, two commonly used methods. First, we

compared the correlations between mean copy number estimates derived from epithelial

cells in the scRNA-seq data with copy number estimates derived from the matched DNA-

seq data (Fig. 2a). All epithelial cells in the metastasized tumor P6198 are malignant cells.

The three primary tumors (P5847, P5931, and P8823) have high purity and thus most of

the epithelial cells are expected to be malignant. Therefore, we expect high correlation

between the CNA profiles of epithelial cells in scRNA-seq data and that from the matched

DNA-seq data. Across the four samples (Supplementary Table 1), correlations of

Clonoscope estimates all exceed 0.7, while existing methods which rely solely on scRNA-

seq data achieve correlations around 0.5. The detailed copy number estimates are shown

in Supplementary Fig. 1. Across the four samples, the lack of concordance between copy

number estimates obtained by existing methods and gold standard values derived from

DNA-seq demonstrates the difficulty of copy number estimation solely from RNA

expression, and motivates the integration with matched bulk DNA-seq for copy number-

based subclone detection in scRNA-seq data.

In addition to providing the genome segmentation which serves as the starting

point of Clonoscope's single-cell subclone detection, the copy number profile from tissue-

matched bulk DNA-seq data is also informative for the assignment of scRNA-seq cells to

malignant versus non-malignant status, a common analysis task in single cell cancer

sequencing. Existing methods use hierarchical clustering for this task, without utilizing

the tissue-matched bulk DNA-seq data. Consider P6198, the metastasized tumor from

above where all epithelial cells are known to be of malignant status and thus can be used

to benchmark copy number-based malignant cell identification accuracy. The results of

copy number-based malignant cell labeling by Clonalscope (left), CopyKAT (middle) and

hierarchical clustering (right) are shown in Fig. 2 (inferCNV does not perform malignant

cell assignment). By utilizing the CNA profile from matched bulk DNA-seq data,

Clonalscope's accuracy in malignant cell labeling reaches ~0.9986 accuracy. In

comparison, CopyKAT's malignant cell labeling, which does not utilize matched bulk

DNA-seq data, has accuracy of ~0.4529 on this sample (Supplementary Fig. 2). While we

recognize that tissue-matched bulk DNA-seq data is not always available, this example

shows that when it is available, it can be harnessed by Clonalscope to substantially boost

accuracy. The accuracy boost is derived not just from the DNA-based segmentation, but

also from Clonalscope's clustering model which utilize the bulk DNA profile as prior. This

can be seen from the fact that even when the segmentation from matched bulk DNA-seq

data is used to delineate the CNA regions, hierarchical clustering is still not able to reliably

label malignant cells (Supplementary Fig. 3).

Next, we compared the accuracy of subclone identification by Clonalscope,

CopyKAT and inferCNV for scRNA-seq data. Existing methods for copy number-based

subclone detection for scRNA-seq data have not explicitly evaluated the accuracy of

intratumor subclones using scDNA-seq gold standard. Instead, benchmarking was only

done at the bulk level. The comparison here was performed on the P5931 primary gastric

tumor sample since it has a clear subclonal structure from scDNA-seq data (Fig. 2c left

and [9]), and furthermore, it has matched normal scRNA-seq data generated for

benchmarking the performance of subclone identification (Supplementary Fig. 4).

Focusing on the malignant cells in the P5931 scRNA-seq dataset, the subclones identified

from Clonalscope (top), CopyKAT (middle), and inferCNV (bottom) were compared to the

three main subclones (D1 to D3) observed in matched scDNA-seq data (Fig. 2c-e;

Methods). Proportions of each subclone detected by each method are shown in

Supplementary Fig. 5. Clonalscope successfully identified all three major subclones with

proportions comparable to those in scDNA-seq data (Supplementary Fig. 5).

Comparatively, subclone D3 was apparently missed in the results of CopyKAT and

inferCNV (Fig. 2d, e). Also, compared to existing methods, Clonalscope does not require

stringent filters and thus retains more low coverage cells in the analysis (Supplementary

Fig. 5). Thus, with validation using scDNA-seq data, we have shown that Clonalscope is

able to improve the accuracy of malignant cell labeling and intratumor subclones

detection. The improvements are achieved by harnessing tissue-matched bulk DNA-seq

data and by a nonparametric Bayesian model tailored for clustering of clonal copy number

profiles.

Figure 2.

Figure 2: Subclone detection and malignant cell labeling for scRNA-seq data validated by matched scDNA-seq data.

a. Correlations between the copy number profiles from matched DNA-seq data and those from different segmentation methods on four gastrointestinal tumor samples. b. Malignant cell labeling from Clonalscope (left), CopyKAT (middle), and Hierarchical clustering with 2 major groups (right) on the

P6198 scRNA-seq data. T and N represent the estimated tumor and normal cells respectively. G1 and G2 stand for the first and second major groups from hierarchical clustering. c. Mapping the detected subclones from Clonalscope (top), CopyKAT (middle), and inferCNV(bottom) in scRNA-seq data (right) to those observed in the matched scDNA-seq data (left) for the P5931 malignant cells (Methods).

Subclone detection based on coverage and allelic ratio

Clonalscope can also utilize allele-specific information for subclone detection,

when enough allele-specific reads can be extracted from single-cell sequencing data. We

demonstrate this application on scATAC-seq data for SNU601, a gastric cancer cell line

with multiple subclones delineated by scDNA-seq data. Some subclones in SNU601 are

differentiated mostly by their allele-specific copy number changes [9]. In our previous study

[9],we applied a supervised classification approach to assign each cell into one of the six

major subclones detected in the matched scDNA-seq data. If we were, instead, to perform

unsupervised hierarchical clustering on the scATAC-seq data, we would not be able to

recapitulate the subclones due to the low signal-to-noise ratio (Supplementary Fig. 6).

The results for Clonoscope are shown in Fig. 3. Based on the segmentation of the

pseudo-bulk (Fig. 3a), cells in the SNU601 scDNA-seq data can be clustered into six

major subclones with 10 marker regions shown in Fig. 3b, which can be viewed as the

ground truth for evaluating the subclone detection results in the much shallower scATAC-

seq data (Fig. 3c).

We applied Clonalscope to the scATAC-seq data in the 10 regions in Fig. 3a , and

identified 13 subclones in an unsupervised manner (Fig. 3c). The leftmost column shows

the subclone identity of each cell assigned by Clonalscope (C). For comparison, we also

conducted supervised classification—simply assigning each cell in scATAC-seq to one of

the six subclones identified by scDNA-seq, utilizing the subclonal mean copy number

profiles from scDNA-seq data. The clusters from supervised classification (S) are also

shown in Fig. 3c. A detailed comparison of this supervised subclone assignment versus

Clonalscope unsupervised clustering are shown in the contingency table in

Supplementary Fig. 7. We see that the six major subclones (right) all have counterparts

in Clonalscope's clusters that exhibited high enrichment. Specifically, the cells assigned

to S1 to S6 are mainly enriched in C10, C1, C5, C4, C12, and C3 respectively. The

consensus plot also shows comparable allele-specific copy number profiles between the

mapped clones from Clonalscope and those derived from the scDNA-seq data

(Supplementary Fig. 8). Additionally, we computed the UMAP coordinates for cells in the

scATAC-seq data based on their peak profiles and colored the cells based on their clonal

assignment from Clonalscope and from supervised classification (Supplementary Fig. 9;

[9]). Clusters from the two methods show similar local enrichment patterns on the UMAP

plots. The example here indicates that Clonalscope can detect subclones not only based

on total coverage but also based on integrated modeling of total coverage and allele-

specific copy number profiles.

Figure 3.

Figure 3: Subclone detection based on coverage and allelic ratio in the scATAC-seq data from the SNU601 gastric cancer cell line.

a. Segmentation plot of the pseudo-bulk scDNA-seq dataset from the SNU601 sample. b. The six major subclones detected on the matched scDNA-seq data from hierarchical clustering with ten regions shown in plot c. The result of subclone detection from Clonalscope on the scATAC dataset. The clusters from

Clonalscope (C) and supervised classification (S) are shown parallelly in the left. In the color legend, M and m represent the major haplotype and minor haplotype, respectively.

Malignant spot labeling and subclone tracing on spatial transcriptomic data

We next demonstrate the application of Clonalscope on ST data. First, consider

an ST data set consisting of two adjacent slices from a squamous cell carcinoma (SCC)

sample, with matched bulk WES data. Leveraging the segmentation and bulk copy

number profile from the matched WES data (Fig. 4a), Clonalscope identified malignant

cells in both slices, which are spatially concentrated despite the fact that Clonalscope did

not make use of spatial coordinates in subclone detection (Fig. 4b). Clonalscope's

malignant spot labeling are highly concordant with expert pathology annotation (Fig. 4c).

Further, across the two spatially adjacent slices, Clonalscope's copy number estimates,

for the spots labeled as malignant, are highly correlated (correlation ~0.84 Fig. 4d,e).

Since the two spatially adjacent slices serve as approximate technical replicates of each

other, this analysis demonstrates the accuracy and reproducibility of Clonoscope in

malignant cell labelling for spatial transcriptomic data. Detailed heatmaps are shown in

Supplementary Fig. 10.

Currently, there is no method for reliable subclone tracing between samples

collected at multiple timepoints, anatomic positions, or spatial slices for the same patient,

which is common in today's study designs. Since Clonalscope allows the flexible

integration of prior information in subclone detection, it can be applied to link subclones

between samples for the same patient (Fig. 4f). Subclone tracing by Clonalscope is

demonstrated on an invasive ductal carcinoma (IDC) data set with samples from two

tissue   sections   (https://www.10xgenomics.com/resources/datasets/human-breast-

cancer-block-a-section-1-1-standard-1-0-0). Since there are no bulk DNA sequencing

data available for these samples, we simply segmented the genome by chromosome arm.

First, we applied Clonalscope on the ST data from one section, yielding a total of 15

clusters (Fig. 4g; left). As before, even though the clustering algorithm does not conduct

any smoothing across adjacent spots, the detected subclones are mostly spatially

contiguous, which is evidence that the subclones are truly spatially localized. Next, these

clusters were used as priors for the subclone detection for the second tissue section,

while allowing new clusters to be seeded and prior clusters to be modified/removed (Fig.

4f). In the second tissue section, 11 out of the 15 clusters detected by Clonoscope were

preserved with one novel cluster detected (Fig. 4g; right). Comparing the spatial

distributions of the detected clusters in the two adjacent sections, we see that the sites of

the corresponding clusters match well between the two slides. In both tissue sections, the

shapes and positions of subclones delineated by Clonalscope are concordant with the

outlines of the tumor in the pathology annotation (Fig. 4h).

Figure 4.

a HMM segmentation across all chroms

Figure 4: Malignant spot labeling and subclone tracing on spatial transcriptomics (ST) data.

a. Segmentation plot of the whole-exome sequencing (WES) data from the squamous cell carcinoma (SCC) Patient 6 (P6) sample. b. Malignant cell labeling by Clonalscope on the ST datasets from SCC P6 replicate 1 (left) and replicate 2 (right). c. The H&E image of the SCC P6 replicate 1 sample with pathology annotation. d. Normalized genome-wide copy number profiles between the identified malignant cells in the two replicates. e. The scatter plot of the mean read counts for each region across the identified malignant cells for SCC P6 replicate 1 against those for replicate 2. f. Illustration of the scheme for subclone detection using Clonalscope. Clonalscope can leverage the prior subclones from the first dataset to detect subclones in another related dataset. g. Subclones detected from Clonalscope in the ST datasets from section 2 (left) and section 1 (right) of an invasive ductal carcinoma (IDC) sample. h. The H&E image of the IDC section 1 sample with pathology annotation.

Clonalscope identified spatially segregated subclones expressing genes associated with drug resistance and poor survival

As a second demonstration of the application of Clonalscope to subclone detection in ST data, we analyzed VISIUM 10x data from an invasive ductal carcinoma (IDC sample) (https://www.10xgenomics.com/resources/datasets/human-breast-cancer-ductal-carcinoma-in-situ-invasive-carcinoma-ffpe-1-standard-1-3-0). Since there are no matched bulk WGS data for this sample, copy numbers were estimated at the chromosome arm level. In this sample, Clonalscope identified nine clusters based on the copy number profiles (Supplementary Fig. 11). The nine clusters are locally enriched in the tissue (Fig. 5a) although no spatial information was used in the estimation. Among the clusters, cluster 2 is the major clone; cluster 3 and cluster 4 are the minor subclones spatially segregated in the tissue. Most spots containing malignant cells are contained in cluster 2. Comparatively, spots in cluster 3 are only limited to the connected region labeled in blue, while spots in cluster 4 are limited to three separate ducts in the right bottom region. The three clusters (cluster 2, 3, and 4) were separated due to difference in their copy number states for regions such as chr2q and chr8q (Supplementary Fig. 11). To illustrate the difference in the copy number profiles between the three clones, the mean fold changes across the spots for each of the three subclones are shown in Fig. 5b. Four

regions with significant shifts in the values of at least one of the three subclones were

highlighted in the density plots (Fig. 5b; top).

Next, we performed differential expression analysis on the three subclones

detected by Clonalscope. Gene expression of the top 6 highly expressed genes for

subclone 3 and subclone 4 are shown in Fig. 5c and Fig. 5d respectively. Expression of

these genes corresponds well to spatial distributions of the two subclones. These genes

were also discovered to be associated with outcomes of therapies and different

adaptability of tumors. For example, *FGG* is highly expressed in subclone 4. Previous

study discovered that breast cancer patients with expression of this gene responded

poorly   to   chemotherapy   and   had   poor   survivals

(https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7136924/). Another example is the highly

expressed *IGFBP5* gene in subclone 3. This gene was found to be associated with

metastatic   phenotype   in   breast   tumors

(https://bmccancer.biomedcentral.com/articles/10.1186/1471-2407-9-103).   Higher

expression of other genes in the two subclones such as *NNMT, TGM2*, and *CAPN13*

were all shown to be associated with poor survival and drug resistance in breast cancer

patients [28-30]. The results suggest that the subclones detected by Clonalscope can

potentially be the sources of resistance to drug and therapy.

Figure 5.

Figure 5: Subclone detection and differential gene analysis on an invasive ductal carcinoma sample.

a. Subclones detected from Clonalscope on the ST dataset from an invasive ductal carcinoma sample. b. The mean copy number profiles across the cells assigned to the three spatially segregated subclones (bottom), and the density plots of four regions with significant differences in the copy number states for at least one of the three subclones (top). c. and d. show the top 6 highly expressed genes for cluster 3 and cluster 4. The color scale indicates the numbers of UMIs for each gene.

Discussion

With the development of single-cell sequencing technologies, there is increased

interest in subclone detection based on chromosomal copy numbers estimated at the

single cell level. Existing copy number-based subclone detection methods are mostly

based on hierarchical clustering, and do not leverage information from matched DNA-seq

data. Here we presented Clonalscope, a subclone detection method for single-cell and

ST data that leverages prior information from matched bulk DNA-seq data for malignant

cell/spot labeling and subclone tracing between related datasets. We demonstrated

Clonalscope on diverse single-cell data types including scRNA-seq, scATAC-seq, and

spatial transcriptomic data. Subclone detection in scRNA-seq and scATAC-seq data were

benchmarked with matched bulk and single cell DNA-seq data. In spatial transcriptomic

data, expert pathology annotation was used to assess the accuracy of malignant spot

labeling, and comparisons between adjacent tissue slices was used to assess the

reproducibility of spatial subclone mapping.

Clonalscope estimates copy number states by normalizing to a set of diploid

cells/spots at the start. It is important to understand that this does not require the

classification of all cells into normal versus malignant status, as we only seek to identify

a small set of highly confident normal cells/spots to initialize the normalization. The diploid

cells can be a user-selected set of well-known immune or stromal cell types that are easily

identifiable with marker genes in standard single-cell data analysis pipelines. Alternatively,

Clonalscope can first normalize each cell by the median profile across all cells in the

dataset, and then use the malignancy labelling in each round of estimation to iteratively

update the diploid cell selection. For ST datasets, diploid spots can be selected based on

marker genes, similar to that for scRNA-seq data. Since H&E images are also provided

for the ST data, expert pathology annotation can also be useful in verifying/selecting spots

from normal regions.

Subclone detection in Clonalscope is based on a nested Chinese Restaurant

Process that captures the special, nested structure of subclonal copy number alterations.

Within a tumor, CNAs occurring at the early developmental stage are present in most

malignant cells, while some subclonal CNAs are only observed in a subset of malignant

cells. The nested Chinese Restaurant Process not only samples subclonal membership,

but also samples copy number states for each region to create a new copy number profile

for the membership assignment. In sampling copy number states, most regions retain the

pre-existing state, while there is a rare chance that a region moves to a new state,

mimicking the occurrence of a random mutation. The raw outputs from the nested

Chinese Restaurant Process depend on user parameters. For more stable estimation, we

include a post-processing step that constrains the detected clusters to at least n_{min} cells.

We also demonstrated the application of Clonalscope to haplotype-informed

subclone detection on scATAC-seq data. There is currently a need for reliable haplotype-

based subclone detection method for sparse single-cell sequencing data such as

scATAC-seq. Improving upon [9], Clonalscope does not rely on the known subclones

detected from the matched scDNA-seq data. Comparatively, most scRNA-seq protocols

only sequence the 3' or 5' end of transcripts, limiting allele-specific CNA detection in

scRNA-seq data. Recent technologies enable multi-modal sequencing such as RNA and

ATAC at the single-cell level [31], providing opportunities for broader applications of

Clonalscope in analysis of allele-specific CNA profiles.

On the ST dataset from an IDC sample, our detailed analysis identified minor subclones

(spots with the same copy number configuration) that might be resistant to therapy. Since

this data is not single-cell resolution (each spot contains 1 to 10 cells), Clonalscope also

identified clusters consisting of spots with mixtures of tumor and stromal cells based on

the gradients of the copy number values. These spots can be the regions of interest for

characterizing the interactions between tumor and stromal cells with the help of recently

developed cell type deconvolution methods [32-35].

Another unique feature of Clonalscope is its ability to trace subclones, a common analysis

step in studies where multiple tissue samples are sequenced for the same patient. In this

paper, we demonstrated subclone tracing in a data set involving adjacent tissue slices

from the same patient. Clonalscope can also be used to link subclones between samples

from different time points or different anatomic positions (*unpublished*). Tracing subclones

between sequential time points enables the study of temporal dynamics of subclones over

the treatment course of cancer, possibly identifying subclones that are resistant to therapy.

Materials and Methods

Datasets and pre-processing

The four 10x scRNA-seq datasets, five Visium spatial transcriptomics (ST) datasets, and the SNU601 scATAC-seq dataset analyzed in this study are summarized in Supplementary Table 1. P5847, P6198 and P8823 scRNA-seq data were generated using the protocol described in the previous study (https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7269843). The Cell Ranger v 3.0 (10x Genomics) was used to generate gene-by-cell matrices of unique molecular identifier (UMI) counts for scRNA-seq data by aligning to the GRCh38 reference genome. The ST datasets of the two replicates from squamous cell carcinoma (SCC) were provided by Ji et al [23]. The three ST datasets from invasive ductal carcinoma (IDC) were retrieved from the 10x website (https://www.10xgenomics.com/resources/datasets). The SNU601 scATAC-seq dataset was retrieved from our earlier Alleloscope paper [9].

Single-cell and bulk DNA sequencing and data preprocessing

Results of segmentation and subclone detection of the four 10x scRNA-seq datasets were validated with matched scDNA-seq and WGS data (Supplementary Table 3.2). The scDNA-seq data of P5847, P5931 were retrieved from [76] and P6198 were retrieved from [137]. WGS was performed on the P8823 sample for this study.

For WGS sequencing, DNA was first extracted using PureLink™ Genomic DNA Mini Kit as per the manufacturer's protocols. Genomic DNA was quantified using the Qubit 2.0 Fluorometer (ThermoFisher Scientific, Waltham, MA, USA). For library preparation,

NEBNext® Ultra™ DNA Library Prep Kit for Illumina, clustering, and sequencing reagents

was used throughout the process following the manufacturer's recommendations. Briefly,

the genomic DNA was fragmented by acoustic shearing with a Covaris S220 instrument.

Fragmented DNA was cleaned up and end repaired. Adapters were ligated after

adenylation of the 3'ends followed by enrichment by limited cycle PCR. DNA libraries

were validated using a High Sensitivity D1000 ScreenTape on the Agilent TapeStation

(Agilent Technologies, Palo Alto, CA, USA), and were quantified using Qubit 2.0

Fluorometer. The DNA libraries were also quantified by real time PCR (Applied

Biosystems, Carlsbad, CA, USA). Then the sequencing library was clustered onto a

flowcell. After clustering, the flowcell was loaded onto the Illumina HiSeq or equivalent

instrument according to manufacturer's instructions. The samples were sequenced using

a 2x150bp Paired End (PE) configuration. Image analysis and base calling were

conducted by the HiSeq Control Software (HCS). Raw sequence data (.bcl files)

generated from the Illumina machine was converted into fastq files and de-multiplexed

using Illumina bcl2fastq 2.17 software. One mis-match was allowed for index sequence

identification.

Segmentation on matched bulk DNA-seq data

The first step of Clonalscope is segmenting the genome into regions with different CNA

profiles. In this step, matched bulk DNA sequencing data (WGS/WES) or pseudo-bulk

data from scDNA-seq data can be segmented by an HMM-based segmentation method.

The HMM method operates on the binned counts of pooled cells, and assumes a Markov

transition matrix on four hidden states representing deletion, copy-neutral state, single-

copy amplification and double-copy amplification: $\begin{pmatrix} 1-3t & t & t & t \\ t & 1-3t & t & t \\ t & t & 1-3t & t \\ t & t & t & 1-3t \end{pmatrix}$,

where $t = 1 \times 10^{-6}$ by default. Emission probabilities follow a normal distribution with 0.2

standard deviation with means depend on individual datasets. The segmentation was

performed on pseudo-bulk scDNA-seq data for P5847, P5931, and P6198; WGS data for

P8823; and WES for SCC P6. Without matched bulk DNA-seq data, the segmentation

was performed on chromosome arms.

Selection of diploid cells

To identify diploid cells in the scRNA-seq data, the raw scRNA-seq UMI count matrices

were first filtered to include cells with at least 500 UMIs, at least 300 genes expressed,

and at most 20% of the reads mapped to mitochondria transcriptome. Next, the matrices

were normalized with size factors (http://dx.doi.org/10.1186/s13059-014-0550-8) and log-

transformed. Then PCA was performed, and 30 PCs were selected for clustering and

UMAP analysis (https://doi.org/10.1126/science.aax1971). For separate clusters on the

UMAP, we annotate the cell types in the gastrointestinal tumor tissues using the marker

genes from the previous study (https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7269843).

To confidently differentiate malignant cells from normal epithelial cells in the P5931

scRNA-seq dataset, we processed both tumor and normal scRNA-seq datasets together

using the same pipeline described above, and successfully labeled normal epithelial cells

in the tumor dataset in a separate cluster mainly containing cells in the normal dataset

(Supplementary Fig. 4).

To identify reference cells in the Visium ST datasets, we first ran the Seurat (4.0.1) [36]

pipeline like that for scRNA-seq data to retrieve some preliminary clusters at the spot

level. Since the resolution of the Visium datasets are 1-10 cells, marker gene expression

is not locally enriched but gradient. However, we could still select a cluster with least

expression of the genes in cancer cells (eg. ERBB2 for breast cancer cells) and high

expression of other marker genes for diploid cells. With the help of H&E images, we could

also more confidently select the clusters located in the regions mainly consisting of diploid

cells.

Computing coverage change

With each region r identified from matched DNA-seq data, we applied a Poisson model

to reduce a gene-by-cell matrix to a region-by-cell matrix for scRNA-seq data. For cell i

and gene g, we defined the raw UMI count matrix as N_{ig} with

456
$$N_{ig} \sim Poisson(\lambda_{ig} = s_i b_g X_{ir});$$

457 , where s_i is the normalizing factor of cell i; b_g is the baseline expression of gene g; and

458 X_{ir} is the relative fold change of region r for cell i. The normalizing factor s_i is computed

459 as follows:

460
$$s_i = \frac{\sum_g N_{ig}}{median_i(\sum_g N_{ig})}$$

461 With a set of control cells being defined, $\delta_{ir}'s$ for these cells are equal to 1. Therefore, b_g

462 can be estimated by solving

463
$$\hat{b}_g = argmax_{b_g} \left(-\sum_{c=1}^{n_c} (s_c b_g - N_{cg} \log(s_c b_g)) \right)$$

464 By derivation, $\hat{b}_g$ can be solved with direct computation.

465
$$\hat{b}_g = \left(\frac{\sum_c N_{cg}}{\sum_c s_c} \right)$$

466 Then for each cell i and region r,

467
$$\hat{X}_{ir} = argmax_{x_{ir}} \left(-\sum_{g \in r(g)} (s_i \hat{b}_g X_{ir} - N_{ig} \log(s_i \hat{b}_g X_{ir})) \right)$$

468 , where $r(g)$ is region r which gene g belong to.

469 By derivation, $\hat{X}_{ir}$ can be solved with direct computation.

470
$$\hat{X}_{ir} = \left(\frac{\sum_g N_{ig}}{s_i \sum_g \hat{b}_g} \right)$$

For ST datasets, the model is the same, but i is for spot instead of cell.

In the real estimation, we excluded the cycle genes, HLA genes and the mitochondria

genes learning from CopyKAT [15]. We also filtered out the extremely variable genes and

regions overlapping <150 genes.

Subclone detection with Nested Chinese Restaurant Process

After the relative fold changes for each cell and each region ($\hat{X}_{ir}$) are retrieved, A Nested

Chinese Restaurant Process was applied to detect subclones in scRNA-seq data in a

non-parametric manner. First, we consider the coverage-only mode. The model for X_{ir} is

$$X_{ir} \sim N(\theta_{ir}, \sigma_r^2)$$

$$\theta_{ir} = \mu_{Z_i,r}$$

, with $z_i \in \{1, \dots, K\}$ is subclone k which cell i is assigned to. μ_{kr} is the relative fold

change of subclone k in region r, and σ_r is the standard deviation of region r. The

targeted outputs of the Nested Chinese Restaurant Process are Z_i, the estimated

subclonal identity of cell i; and U_{kr}, the copy number states of region r for subclone k.

To include the prior information from the matched bulk DNA-seq data, we first set the

prior U_{prior} as two clusters

$$U_{prior} = \begin{bmatrix} \mu_{11} & \cdots & \mu_{1R} \\ \mu_{21} & \cdots & \mu_{2R} \end{bmatrix} \mid \mu_{1r} = 1; \ \mu_{2r} = \begin{cases} 0.5; deletion \\ 1.5; amplification \end{cases}$$

, with μ_{2r} known from the DNA-seq data. Then for each cell $i = 1 \sim N$,

$Z_{prior} = [z_{prior,1}, \dots, z_{prior,i}, \dots, z_{prior,N}] \ ; \ \hat{z}_{prior,i} = \begin{cases} 1; if \ P(X_i|\mu_1.) > P(X_i|\mu_2.) \\ 2; otherwise \end{cases}$

In the t^{th} iteration of the Nested Chinese Restaurant Process, we first iterate over all cell

i. To assign cell i into one of the existing clusters, the probability of

492
$$\tilde{\theta}_i \mid \tilde{\theta}_{-i}, X_i = \begin{cases} \tilde{\mu}_{k_t}, \; p_{i,k_t} = P(X_i|\tilde{\mu}_{k_t}) \times \dfrac{n_{k_t}^{\setminus i}}{N-1+\beta}; k_t = 1, \dots, K_t \\[2ex] \tilde{\mu}_{it}^{new}, \; p_{it}^{new} = P(X_i|\tilde{\mu}_{it}^{new}) \times \dfrac{\beta}{N-1+\beta} \end{cases}$$

, where $\tilde{\theta}_i \mid \tilde{\theta}_{-i}, X_i$ is the variable for the estimated mean of relative fold change for cell i

across all the regions, conditioning on all other cells and the observation. $n_{k_t}^{\setminus i}$ is the

number of cells assigned to cluster k from the t^{th} iteration excluding cell i; and N is the

total number of cells. β is the scalar parameter controlling the chance of accepting a new

cluster in the process. β was by default set as 2. $\tilde{\mu}_{k_t}$ is the relative fold changes across

all regions for subclone k form the t^{th} iteration, and p_{i,k_t} is the probability of cell i being

assigned to the k_t cluster. $\tilde{\mu}_{it}^{new}$ is the relative fold changes across all regions for a newly

seeded cluster for cell i in iteration t with the probability being p_{it}^{new}. p_{i,k_t} and p_{it}^{new} can be

calculated as follows

502
$$\log p_{i,k_t} \sim \sum_r \log P(X_{ir}|\tilde{\mu}_{k_{tr}}) + \log(n_{k_t}^{\setminus i})$$

503
$$\log p_{it}^{new} \sim \sum_r \log P(X_{ir}|\tilde{\mu}_{itr}^{new}) + \log(\beta)$$

504 To create $\tilde{\mu}_{it}^{new}$, the second Chinese Restaurant Process was executed to select one of

505 the existing copy number states or seed a new copy number state for each region. We

506 skip the subscript "it" for the following notation. For each region r,

507
$$\tilde{\mu}_r^{new} = \begin{cases} \mu_{rk}; with\ probability\ \dfrac{n_{k_t}}{N-1+\alpha} \\[2ex] \mu_r^{new} l; with\ probability\ \dfrac{\alpha}{N-1+\alpha} \end{cases}$$

508
$$\mu_r^{new} \sim \text{Unif}(0,3)$$

509 , where μ_r^{new} is a newly seeded copy number state for region r. n_{k_t} is the number of cells

510 assigned to cluster k from the t^{th} iteration. α is the scalar parameter controlling the

511 chance of accepting a new cluster in the process. By default, α is set as 2. With $p'_{i,k_t}s$ for

512 all k_t clusters and p_{it}^{new} being calculated, $\tilde{z}_{it}$ is sampled based on the probability.

513 After $\tilde{z}_{it}$'s are updated for all cell i, μ_{kr} is sampled based on the following distribution

514
$$\mu_{krt}|e.e \sim N\left(\bar{X}_{kr}, \frac{\sigma_{r(t-1)}^2}{n_{k_t}}\right)$$

515 , where $\sigma_{r(t-1)}$ is the standard deviation of region r for the $(t-1)^{th}$ iteration. Then σ_{rt}

516 can be updated by computing

517
$$\sigma_{rt} = \sqrt{\frac{1}{N}\sum_i \left(X_{ir} - \mu_{z_i rt}\right)^2}$$

518 For subclone detection with both coverage and allelic ratio, the model for X_{ir} is

519
$$X_{ir} = [P_{ir}, \Theta_{ir}]$$

520
$$P_{ir} \sim Normal\left(\mu_{z_i r\rho}, \sigma_{r\rho}\right)$$

521
$$\Theta_{ir} \sim Normal(\mu_{z_i r\theta}, \sigma_{r\theta})$$

In the formula, P_{ir} and Θ_{ir} are the estimated coverage and major haplotype proportion [9]

for region r in cell i. $z_i \in \{1, \dots, K\}$ is subclone k which cell i is assigned to, and $\mu_{kr\rho}$ and

$\mu_{kr\theta}$ are the relative fold change and major haplotype proportion of subclone k in region

r respectively. $\sigma_{r\rho}$ and $\sigma_{r\theta}$ are the standard deviations of relative fold change and major

haplotype proportion for region r. The Nested Chinese Restaurant Process is similar to

that for the coverage-only mode. The outputs from the estimation are Z_i, the estimated

subclonal identity of cell i; and $[U_{kr\rho}, U_{kr\theta}]$ the allele-specific copy number states of

subclone k for region r.

The two parameters α and β control the numbers of subclones detected from the Nested

Chinese Restaurant Process. To retrieve more stable estimation, the first few iterations

of the raw outputs from the Nested Chinese Restaurant Process are trimmed. By default,

200 iterations are run and the estimates from the first 100 iterations are trimmed. After

the trimming step, each cell is assigned into the most possible subclone k using the

majority vote strategy. Next, only the subclones with $> n_{min}$ cells assigned are kept, and

the smaller subclones are removed. By default, n_{min} is set as 1% of cells. Then the cells

originally assigned to the removed subclones are re-assigned to the existing subclones.

Then U_{kr} the copy number states of region r for subclone k is updated based on the re-

assignment.

Assessing segmentation on scRNA-seq data and on matched DNA-seq data

To evaluate different segmentation methods on scRNA-seq data, we calculated the

correlations between the mean copy number profiles in scRNA-seq data to those from

(pseudo)bulk DNA-seq data. Three segmentation methods for scRNA-seq data were

compared—segmentation from matched bulk DNA-seq data; segmentation from

CopyKAT and segmentation from inferCNV. CopyKAT and inferCNV were ran using the

default parameters. CopyKAT and inferCNV perform segmentation mainly at the gene

level, and they output the estimated copy number profiles for each gene in each cell.

Therefore, we computed the mean copy number profiles across all epithelial cells for each

gene. Since the output values for CopyKAT are at the log-scale, the values were

exponentially transformed to original values. Comparatively, we used Clonalscope to

perform segmentation on matched DNA-seq data and compute the mean fold changes

for each segment in scRNA-seq data. For easier comparison with CopyKAT and inferCNV,

the mean fold changes for each segment were transformed to the gene level.

All scDNA-seq data were pre-processed with Cell Ranger DNA pipeline

(https://support.10xgenomics.com/single-cell-dna/software/overview/welcome, beta

v.6002.16.0). For P5847 and P5931, we applied Alleloscope [9] on the scDNA-seq data,

and computed the mean coverage changes across all identified malignant cells for each

segment to retrieve the mean copy number profiles for the pseudo-bulk DNA-seq data.

For P6198, we generated the pseudo-bulk sample from the scDNA-seq data, and

computed the mean coverage changes for each segment. For P8823, we first quantified

the reads in each window with a 100000 bp size for both matched tumor and normal WGS

data by running the "intersect" command in the bedtools [37]. The profile of the mean

coverage changes for P8823 bulk DNA-seq data were then computed for each segment.

Similarly, we also transformed the vector of the mean coverage changes for all DNA-seq

data from the segment scale to the gene scale.

Malignant cell/spot labeling on scRNA-seq and spatial transcriptomic data

Copy number profiles from matched bulk DNA-seq data can also be used to reliably detect

malignant cells/spots in scRNA-seq and ST datasets. After the subclones are detected in

the scRNA-seq and ST datasets, $\hat{\mu}_k$, the estimated copy number profiles of subclone k

can be compared to the copy number profiles from matched DNA-seq data (μ_1) across all

regions. To determine subclone k is malignant or not, the revised cosine similarity (S_c)

was computed as

573
$$S_c = \frac{\sum_r[(\hat{\mu}_{kr} - 1) \cdot (\mu_{1r} - 1)]}{\sqrt{\sum_r(\hat{\mu}_{kr} - 1)^2} \sqrt{\sum_r(\mu_{1r} - 1)^2}}$$

If $S_c > S_{min}$ (0.5 by default), the subclone k is labeled as malignant.

Accuracy of malignant cell labeling was compared between Clonalscope, CopyKAT, and

hierarchical clustering on an scRNA-seq dataset. Clonalscope and CopyKAT were run

using default parameters. For hierarchical clustering, we first retrieved segments from

matched DNA-seq data, and estimated relative coverage fold changes for each cell in

each segment using the function of Clonalscope. Then hierarchical clustering was applied

to cluster the cells based on the estimated copy number profiles instead of using the

Bayesian non-parametric clustering in Clonalscope.

Comparison of subclones between scRNA-seq data and matched scDNA-seq data

On the malignant cells identified in the P5931 scRNA-seq data, we compared the

subclones detected by Clonalscope, inferCNV, and copyKAT to those in the matched

scDNA-seq data. In the scDNA-seq data of P5931, 3 major subclones (D1~D3) were

observed, which can be used as the ground truth of the sample. Since we knew for P5931,

CNAs are mainly on the four chromosomes from the matched pseudo-bulk DNA-seq data,

Clonalscope was run on the four chromosomes to detect subclones with default

parameters. For CopyKAT and inferCNV, hierarchical clustering requires specifying the

number of subclones. Since 6 and 5 subclones were used in the segmentation for

CopyKAT and inferCNV, we set the output numbers of subclones as 6 and 5 respectively.

After the subclones in scRNA-seq data were detected from the three methods, we

mapped the subclones to the corresponding "true" subclones observed in the matched

scDNA-seq data. For subclone mapping, the revised cosine similarity (S_c) was computed:

595
$$S_c = \frac{\sum_g [(\hat{\mu}_{cg} - 1) \cdot (\hat{\mu}_{dg} - 1)]}{\sqrt{\sum_g (\hat{\mu}_{cg} - 1)^2} \sqrt{\sum_g (\hat{\mu}_{dg} - 1)^2}}$$

596   , where $\hat{\mu}_{cg}$ and $\hat{\mu}_{dg}$ are the mean fold change for gene g across the cells assigned to

597   subclone c in scRNA-seq data and subclone $d \in [D1, D2, D3]$ in scDNA-seq data.

598   Subclone c in scRNA-seq data was mapped to the subclone d with the highest similarity

599   score. For Clonalscope, we summed over all the region r instead of gene g.

Subclone detection in scATAC-seq data

Following the pipeline described in our previous study[9], we processed the SNU601

scATAC-seq data and generated the region-by-cell matrices for relative fold changes and

allelic ratios. Next, we constrained the values for the relative fold changes to be < 3, and

the values for the allelic ratios > 0.05 and < 0.95 for more stable estimation. In our

previous study, six major subclones were detected in the SNU601 sample with difference

in their allele-specific copy number profiles mainly on 10 regions. Therefore, we selected

the 10 regions for the downstream subclone detection. Cells with no reads detected in

more than five out of the ten regions were filtered out. With the two processed matrices

as the input, Clonalscope was ran to detect the subclones based on both coverage

changes and allelic ratios using the default parameters. The final clusters were requested

to include > 50 cells. The subclones detected from Clonalscope were compared to the

supervised clustering used in the previous study, which requires a known number of

subclones and their corresponding copy number profiles from scDNA-seq data. The two

methods were shown parallelly for easier comparison.

WES data processing

The bam file of the WES data from SCC P6 was retrieved from GSE144240. On the bam

files for matched tumor and normal samples, we first quantified the reads in each window

with a 100,000 bp size by running the "intersect" command in the bedtools [37]. Then we

segmented the genome into regions with the same copy number states using the HMM

segmentation function in Clonalscope. Since the dataset was originally aligned to hg19,

we used the LiftOver tool (https://genome.ucsc.edu/cgi-bin/hgLiftOver) to convert the

segments to GRCh38 to be compatible with the matched ST dataset.

Spatial transcriptomic (ST) data analysis

Clonalscope was run to detect subclones without considering spatial loci in ST datasets.

The estimated clonal identity of each spot can then be visualized spatially, enabling

integration of the two types of signals. Seurat was used to align each spot to the

corresponding loci in the H&E image.

For SCC P6 sample, the copy number profiles of the identified malignant spots in the two

replicates were compared to each other by computing the correlation. The spatial

distributions of the malignant spots were also shown. On the IDC sample with 2 sections,

to trace subclones from one ST dataset to another ST dataset, we used the subclones

detected on the first section (section 2) as the prior clusters for Clonalscope's subclone

detection on another section (section 1). The prior clusters can be easily provided to

Clonalscope, and the direct information from bulk DNA-seq data is ignored in the mode.

On the IDC sample with three subclones detected (Fig. 5), we assessed the difference in

the mean copy number profiles of the three clones. Differential gene expression analysis

on the three clones was performed using the "FindMarkers" function in Seurat [36]. Each of

the three subclones were compared against the other two subclones in the comparison.

Data availability

The patient scRNA-seq and WGS data generated for this study are available under

dbGAP identifier phs001818. Data is available with dbGaP authorized access for

Health/Medical/Biomedical purposes. The other patient scRNA-seq data were obtained

from dbGAP under accession phs001818 [38, 39]. The patient scDNA-seq data were from

dbGAP under accession phs001711 [9] and phs001818 [39]. For the SNU601 sample,

scDNA-seq data and scATAC-seq data were retrieved from the SRA under accession

PRJNA598203 [40] and PRJNA674903 [9] respectively.

Code availability

Clonalscope is available on GitHub at https://github.com/seasoncloud/Clonalscope.

<u>Reference</u>

1. Magrangeas, F. et al. Minor clone provides a reservoir for relapse in multiple
myeloma. *Leukemia* **27**, 473-481 (2013).
2. Ding, L. et al. Clonal evolution in relapsed acute myeloid leukaemia revealed by
whole-genome sequencing. *Nature* **481**, 506-510 (2012).
3. Wang, K. et al. Genomic landscape of copy number aberrations enables the
identification of oncogenic drivers in hepatocellular carcinoma. *Hepatology* **58**,
706-717 (2013).
4. Beroukhim, R. et al. The landscape of somatic copy-number alteration across
human cancers. *Nature* **463**, 899-905 (2010).
5. Futreal, P.A. et al. A census of human cancer genes. *Nat Rev Cancer* **4**, 177-183
(2004).
6. Kim, C. et al. Chemoresistance Evolution in Triple-Negative Breast Cancer
Delineated by Single-Cell Sequencing. *Cell* **173**, 879-893 e813 (2018).
7. Laks, E. et al. Clonal Decomposition and DNA Replication States Defined by
Scaled Single-Cell Genome Sequencing. *Cell* **179**, 1207-1221 e1222 (2019).
8. Navin, N. et al. Tumour evolution inferred by single-cell sequencing. *Nature* **472**,
90-94 (2011).
9. Wu, C.Y. et al. Integrative single-cell analysis of allele-specific copy number
alterations and chromatin accessibility in cancer. *Nat Biotechnol* **39**, 1259-1269
(2021).
10. Niknafs, N., Beleva-Guthrie, V., Naiman, D.Q. & Karchin, R. SubClonal Hierarchy
Inference from Somatic Mutations: Automatic Reconstruction of Cancer
Evolutionary Trees from Multi-region Next Generation Sequencing. *PLoS Comput*
Biol **11**, e1004416 (2015).
11. Yuan, K., Sakoparnig, T., Markowetz, F. & Beerenwinkel, N. BitPhylogeny: a
probabilistic framework for reconstructing intra-tumor phylogenies. *Genome Biol*
16, 36 (2015).
12. Deshwar, A.G. et al. PhyloWGS: reconstructing subclonal composition and
evolution from whole-genome sequencing of tumors. *Genome Biol* **16**, 35 (2015).
13. El-Kebir, M., Satas, G., Oesper, L. & Raphael, B.J. Inferring the Mutational History
of a Tumor Using Multi-state Perfect Phylogeny Mixtures. *Cell Syst* **3**, 43-53 (2016).
14. Jiang, Y., Qiu, Y., Minn, A.J. & Zhang, N.R. Assessing intratumor heterogeneity
and tracking longitudinal and spatial clonal evolutionary history by next-generation
sequencing. *Proc Natl Acad Sci U S A* **113**, E5528-5537 (2016).
15. Gao, R. et al. Delineating copy number and clonal substructure in human tumors
from single-cell transcriptomes. *Nat Biotechnol* **39**, 599-608 (2021).
16. Patel, A.P. et al. Single-cell RNA-seq highlights intratumoral heterogeneity in
primary glioblastoma. *Science* **344**, 1396-1401 (2014).
17. Serin Harmanci, A., Harmanci, A.O. & Zhou, X. CaSpER identifies and visualizes
CNV events by integrative analysis of single-cell or bulk RNA-sequencing data.
Nat Commun **11**, 89 (2020).
18. Fan, J. et al. Linking transcriptional and genetic tumor heterogeneity through allele
analysis of single-cell RNA-seq data. *Genome Res* **28**, 1217-1227 (2018).

19.   Muller, S., Cho, A., Liu, S.J., Lim, D.A. & Diaz, A. CONICS integrates scRNA-seq
694   with DNA sequencing to map gene expression to tumor sub-clones. *Bioinformatics*
695   **34**, 3217-3219 (2018).
696   20.   Granja, J.M. et al. Single-cell multiomic analysis identifies regulatory programs in
697   mixed-phenotype acute leukemia. *Nat Biotechnol* **37**, 1458-1465 (2019).
698   21.   Satpathy, A.T. et al. Massively parallel single-cell chromatin landscapes of human
699   immune cell development and intratumoral T cell exhaustion. *Nat Biotechnol* **37**,
700   925-936 (2019).
701   22.   Stahl, P.L. et al. Visualization and analysis of gene expression in tissue sections
702   by spatial transcriptomics. *Science* **353**, 78-82 (2016).
703   23.   Ji, A.L. et al. Multimodal Analysis of Composition and Spatial Architecture in
704   Human Squamous Cell Carcinoma. *Cell* **182**, 1661-1662 (2020).
705   24.   Erickson, A. et al. The spatial landscape of clonal somatic mutations in benign and
706   malignant tissue. *bioRxiv* (2021).
707   25.   Elyanow, R., Zeira, R., Land, M. & Raphael, B.J. STARCH: copy number and clone
708   inference from spatial transcriptomics data. *Phys Biol* **18**, 035001 (2021).
709   26.   Myhre, S. et al. Influence of DNA copy number and mRNA levels on the expression
710   of breast cancer related proteins. *Mol Oncol* **7**, 704-718 (2013).
711   27.   Campbell, K.R. et al. clonealign: statistical integration of independent single-cell
712   RNA and DNA sequencing data from human cancers. *Genome Biol* **20**, 54 (2019).
713   28.   Ai, L. et al. The transglutaminase 2 gene (TGM2), a potential molecular marker for
714   chemotherapeutic drug sensitivity, is epigenetically silenced in breast cancer.
715   *Carcinogenesis* **29**, 510-518 (2008).
716   29.   Wang, Y. et al. Nicotinamide N-methyltransferase enhances chemoresistance in
717   breast cancer through SIRT1 protein stabilization. *Breast Cancer Res* **21**, 64
718   (2019).
719   30.   Bhattacharya, A. et al. A framework for transcriptome-wide association studies in
720   breast cancer in diverse study populations. *Genome Biol* **21**, 42 (2020).
721   31.   Zhu, C., Preissl, S. & Ren, B. Single-cell multimodal omics: the power of many.
722   *Nat Methods* **17**, 11-14 (2020).
723   32.   Cable, D.M. et al. Robust decomposition of cell type mixtures in spatial
724   transcriptomics. *Nat Biotechnol* **40**, 517-526 (2022).
725   33.   Elosua-Bayes, M., Nieto, P., Mereu, E., Gut, I. & Heyn, H. SPOTlight: seeded NMF
726   regression to deconvolute spatial transcriptomics spots with single-cell
727   transcriptomes. *Nucleic Acids Res* **49**, e50 (2021).
728   34.   Dong, R. & Yuan, G.C. SpatialDWLS: accurate deconvolution of spatial
729   transcriptomic data. *Genome Biol* **22**, 145 (2021).
730   35.   Kleshchevnikov, V. et al. Cell2location maps fine-grained cell types in spatial
731   transcriptomics. *Nat Biotechnol* **40**, 661-671 (2022).
732   36.   Hao, Y. et al. Integrated analysis of multimodal single-cell data. *Cell* **184**, 3573-
733   3587 e3529 (2021).
734   37.   Quinlan, A.R. & Hall, I.M. BEDTools: a flexible suite of utilities for comparing
735   genomic features. *Bioinformatics* **26**, 841-842 (2010).
736   38.   Sathe, A. et al. Single-Cell Genomic Characterization Reveals the Cellular
737   Reprogramming of the Gastric Tumor Microenvironment. *Clin Cancer Res* **26**,
738   2640-2653 (2020).

39.   Sathe, A. et al. The cellular genomic diversity, regulatory states and networking of
the metastatic colorectal cancer microenvironment. *bioRxiv* (2020).
40.   Andor, N. et al. Joint single cell DNA-seq and RNA-seq of gastric cancer cell lines
reveals rules of in vitro evolution. *NAR Genom Bioinform* **2**, lqaa016 (2020).

Acknowledgements

The work is supported by the National Institutes of Health [P01HG00205ESH to B.T.L.,

S.M.G. AND H.P.J., 5R01-HG006137-07 and 1U2CCA233285-01 to C-Y.W. and to

N.R.Z.]. Additional support to HPJ came from the Research Scholar Grant, RSG-13-297-

01-TBG from the American Cancer Society, Clayville Foundation and the Gastric Cancer

Foundation.

Author contributions

C.-Y.W. and N.R.Z. conceived the computational methods and designed the study with

help from H.P.J. C.-Y.W. developed and implemented the computational methods and

conducted all data analyses. A.S. performed all related sample preparation and

sequencing. J.R. helped with benchmarking and improvement of the method. P.R.H.

helped with pathology annotation. B.T.L. helped in data interpretation. S.M.G. performed

data pre-processing and coordinated data transfer. H.P.J. advised all experiments and

data collection. C.-Y.W. and N.R.Z wrote the manuscript. All authors read and approved

the final manuscript.

Competing interests

The authors declare no competing interests.