Beyond sequence similarity: cross-phyla protein annotation by structural prediction and alignment

Fabian Ruperti[1, 2⊘], Nikolaos Papadopoulos[1⊘], Jacob Musser[1], Detlev Arendt[1,3*]

1 Developmental Biology Unit, European Molecular Biology Laboratory, 69117 Heidelberg, Germany

2 Collaboration for joint Ph.D. degree between EMBL and Heidelberg University, Faculty of Biosciences 69117 Heidelberg, Germany.

3 Centre for Organismal Studies (COS), University of Heidelberg, 69120 Heidelberg, Germany

⊘: These authors contributed equally to this work. *: arendt@embl.de

Abstract

Background: Annotating protein function is a major goal in molecular biology, yet experimentally determined knowledge is often limited to a few model organisms. In non-model species, the sequence-based prediction of gene orthology can be used to infer function, however this approach loses predictive power with longer evolutionary distances. Here we propose a pipeline for the functional annotation of proteins using structural similarity, exploiting the fact that protein structures are directly linked to function and can be more conserved than protein sequences.

Results: We propose a pipeline for the functional annotation of proteins via structural similarity (CoFFE: **Co**lab**F**old-**F**oldSeek-**E**ggNOG) and use this to annotate the complete proteome of a sponge. Sponges are uniquely positioned for inferring the early history of animals, yet their proteomes remain sparsely annotated. CoFFE accurately identifies proteins with known homology in $> 90\%$ cases, annotating an additional 50% of the proteome beyond sequence-based methods. Using this, we uncover new functions for sponge cell types, including extensive FGF, TGF and Ephrin signalling in sponge epithelia, and redox metabolism and

control in myopeptidocytes. Notably, we also annotate genes that arose via horizontal gene transfer, which are specific to the enigmatic sponge mesocytes and likely participate in digesting algal and plant cell walls.

Conclusions: Our work demonstrates that structural similarity is a powerful approach that complements and extends sequence-based annotations to bridge long evolutionary distances. We anticipate this approach will expand functional annotations across the tree of life and boost discovery in numerous -omics datasets.

Keywords

Functional annotation, proteins, *Spongilla*, scRNA-seq, structural similarity, structure conservation, ortholog conjecture

Background

Knowledge of protein function is crucial for interpreting many types of high-throughput molecular datasets. Since protein functional studies are limited to a few model species, amino acid sequences have been used to infer the function of protein homologs. However, homology detection over longer evolutionary distances remains challenging owing to the decay of protein similarity that abolishes evidence of historical continuity. This presents a severe bottleneck for inferring protein function across a wide expanse of the tree of life, particularly in distant organisms where many proteins fall in the "twilight zone", only sharing a sequence identity between $10 - 20\%$ with proteins in characterised models [1, 2].

A way to venture more deeply into the twilight zone is to use structural similarity for functional annotation, as structure is more conserved in evolution [3]. Until recently, this was not feasible since predicting protein structures from amino acid sequence required prior inference of sequence homology [4]. This has changed with the advent of AlphaFold [5], a deep learning AI system that can predict *de novo* protein structures with atomic resolution, together with novel approaches for identifying similar structures in large databases [6]. Protein structures can now be predicted by AlphaFold for entire proteomes, and then aligned to structures from model systems with characterised functions.

In this work, we predicted structures for the sparsely annotated proteome of the freshwater sponge *Spongilla lacustris*, an early-branching animal, and aligned them against available structural databases. We show a 50% expansion of previous functional annotations. Revisiting recent single-cell RNA-sequencing data [7] we discovered additional aspects of sponge cell biology, such as extended cell signalling in pinacocytes, redox metabolism and control in myopeptidocytes, and polysaccharide digestion as a key function of the previously uncharacterized mesocytes.

Results

A protein structure-based pipeline enriches functional annotation for *Spongilla lacustris*

We created a structure-based pipeline for functional annotation transfer, which we refer to as CoFFE (**Co**lab**F**old-**F**oldSeek-**E**ggNOG). Instead of using amino acid sequences directly, we predict protein structures, align them against structural databases, and transfer the functional annotation of the best hits to the queries (for an overview, see Suppl. Fig. 5; details in Methods). As a test case, we chose to annotate proteins in the freshwater demosponge *Spongilla lacustris*, an early-branching animal. With only about 20 cell types, organised into 4 cell families, *Spongilla* is a key model for understanding the origins of specialised animal cells [7].

We used the ColabFold [8] pipeline to predict three-dimensional structures of all 41943 predicted *Spongilla* proteins, including isoforms (all structures and metadata deposited to ModelArchive, see Methods). We succeeded, except for 11 proteins, which were too long (>2900 amino acids) to be predicted by the available hardware. Confidence of predicted protein structures was assessed by calculating average pLDDT (predicted local distance difference) values. Notably, average pLDDT values for Spongilla protein structures were similar to those of well-characterised animal models (Fig. 1A).

Next, we used the predicted protein structures as queries to search with FoldSeek [6] against AlphaFoldDB [9], SwissProt [10], and PDB [11]. In general, bit scores for the best FoldSeek hits were positively correlated with mean pLDDT. Neither parameter correlates with predicted physico-chemical properties of the proteins such as hydrophobicity, isoelectric points or their instability index (Supplemental Note B, Supplemental Figs. 6-8). After removing lower-quality matches we transferred the functional annotation of the best hits to the sponge proteins, producing annotations for slightly more than 60% of the proteome (25232 proteins). This represents an approximately 50% increase in annotated proteins compared to the state-of-the-art orthology pipeline EggNOG-mapper [12]. This moves Spongilla into a comparable level of functional annotation as *C. elegans* (70% at least described), one of the most intensively studied model species (Fig. 1B).

Fig 1. Structural prediction and alignment of the *Spongilla* proteome A) Distribution of average pLDDT for predicted proteomes from common model species in comparison to *Spongilla lacustris*. B) Proportion of EggNOG or CoFFE protein annotations in *S. lacustris* and other eukaryotes. Highlighted organisms appear in A. C) Overlap between EggNOG and CoFFE annotations. *Ortholog*: proteins identified as belonging to the same orthology group in the most recent common ancestor in the EggNOG database. *Protein fam.*: proteins identified as belonging to the same eukaryote orthology group in the EggNOG database, indicating annotations represent homologs in the same gene family. 50% *PFAM*: half of the sequence-based PFAM domains are shared. *No agreement*: CoFFE and EggNOG annotations identify non-homologous proteins. Subcategories with "same name" denote the fractions where EggNOG and CoFFE returned the same preferred name for a protein. D) Annotated proportions of different categories of *S. lacustris* proteins.

CoFFE annotations agree with sequence-based annotation transfer

As a first step towards validating our pipeline, we compared CoFFE annotations to sequence-based annotations generated by EggNOG-mapper [12]. This method annotates proteins by searching precomputed gene trees in the eggNOG orthology database and assigns query proteins to fine-grained orthology groups. A total of 16589 proteins were annotated by both pipelines. For 90.6% of these proteins, best hits from EggNOG and CoFFE were homologs (Fig. 1C), including 56.7% orthologs and an additional 33.9% representing proteins in the same gene family. In the remaining cases, most proteins (∼5%) shared PFAM domains in

common [13], bringing the percentage of proteins with at least minimal agreement between CoFFE and EggNOG-mapper to >95%. These results indicate that CoFFE can successfully reproduce sequence based homology inferences.

We also compared our results to "legacy" annotations from the recently published Spongilla cell type atlas, which used BLASTp to supplement EggNOG-mapper annotations [7]. Compared to this combined sequence-based approach, CoFFE annotates more proteins proteome-wide ($\sim$60% to $\sim$40%). More importantly, CoFFE markedly improved the proportion of annotated cell type- and cell family-specific markers ($\sim$70%) compared with sequence-based approaches ($\sim$56%; Fig. 1D, Suppl. Fig. 9). This indicates CoFFE will be valuable for exploring cell type-specific functions in non-model organisms.

CoFFE expands annotation of signalling pathways in Sponge pinacocytes

Pinacocytes are contractile epithelial cells that line the sponge canal system, playing important roles in morphogenesis, barrier formation, and sponge whole-body contractions [14,15]. Musser *et al.* showed pinacocytes upregulate genes involved in TGF/BMP and Wnt signalling, key pathways regulating development and morphogenesis in multicellular animals [16,17]. Supporting this, CoFFE identified additional pinacocyte marker genes in each of these pathways.

In the FGF pathway, previous sequence-based annotations identified Fgf receptors and the FGF regulators Frs and Grb2, which are expressed in different pinacocyte cell types (Figure 2A). Extending this, CoFFE identified GAB1 and GAB2 (GBR2-associated-binding protein 1/2) as well as PTPN11/SHP2, which are necessary for signal transmission into the cell [18,19]. Notably, CoFFE also detected the FGF ligand in *Spongilla*, which was not found using sequence-based approaches. Structural superposition of the protein with its best FoldSeek hit (UniProtID: P48804 , *Gallus gallus* FGF4) revealed an extensive alignment of large parts of the protein, with a RMSD of 0.89 over 543 atoms despite a sequence similarity of only 11.8% (Fig. 2B). Newly annotated players in the FGF pathway also exhibited enrichment in pinacocytes, consistent with its previously known cellular role.

Musser *et al.* also described multiple genes involved in TGF-β signalling in the pinacocyte family, including Tgfbr1, Acvr, Smad and Smurf [20]. CoFFE extends the list of known actors in pinacocyte TGF signalling, adding important ligands such as INHBE, CHRDL1 or KCP.

Lastly, Ephrin and the ephrin receptor (Eph) are membrane-anchored signalling molecules that mediate communication between adjacent cells [21, 22]. Whereas multiple ephrin receptors were detected via sequence similarity, ephrin itself was only found in the *Spongilla* proteome using highly sensitive HMM profile searches [23, 24]. CoFFE annotates a *Spongilla* gene with differential expression in various pinacocytes as a homolog of *Caenorhabditis elegans* Efn-3 (UniProtID: Q19475). Although these proteins share only 22% sequence identity, the superimposed structures achieve an RMSD of 1.59 over 580 atoms (Fig. 2C). A separate Hmmer search [25] using the ephrin Pfam profile (PF00812) picked up the same gene, suggesting CoFFE is at least as sensitive as curated HMM profile searches.

FGF, TGF-beta as well as Ephrin interestingly exhibit converging downstream signalling pathways, including PI3K/Akt, ERK/MAPK1, JNK or RhoA/ROCK, responsible for cell growth, differentiation, migration and cytoskeletal organisation [21, 26, 27]. Together, CoFFE and sequence-based methods identified the principal proteins involved in the downstream pathways. The genes encoding these proteins are broadly expressed across most *Spongilla* cell types, consistent with their diverse functional roles. This example highlights the power of CoFFE to vastly extend annotations and further elaborate cell type specific functions.

Redox metabolism and control in myopeptidocytes

The mesohyl of sponges is a collagenous, dynamic tissue forming large parts of the body between pinacocytes and the feeding choanocyte chambers [28]. Musser *et al.* identified five novel mesenchymal cell types [7] in *Spongilla*. Among them, myopeptidocytes are an abundant uncharacterised cell type, forming long projections that contact other cells. Sequence-based annotations suggested myopeptidocytes function to generate and degrade hydrogen peroxide by expressing dual oxidase (Duox1), its maturation factor (DuoxA), and Catalase (Cat) [29]. Myopeptidocytes also express transporters of copper ions which in their reduced state react with H_2O_2 (Fenton reaction) [30] leading to the generation of hydroxyl radicals in cells. The existence of these prominent reactive oxygen species (ROS) (Fig. 3) is further supported by expression of Cyba [31]. However, further roles of ROS metabolism and function are unclear.

CoFFE identified key additional members of ROS generation, metabolism and response that are expressed in myopeptidocytes. CoFFE annotated Ferric-chelate reductase (FRRS1 and FRRS1L), which recycles Fe^{3+} to its reduced state, and suggests iron-based generation of H_2O_2.

At the same time, proteins newly annotated by CoFFE suggest downstream metabolism and function of ROSs: Disulfide oxidoreductase (DsbB) as well as Flavin carrier protein (FLC) both play a role in oxidative protein folding [32, 33]. NmrA-like acts as a redox sensor in the cell [34]. Consistent with a possible redox regulation role, myopeptidocytes express a range of additional ROS responsive proteins identified by CoFFE: Sulfiredoxin 1 (SRXN1) promotes resistance against oxidative stress damage [35], whereas AP endonuclease 1 (APEX1) protects against ROS induced DNA damage [36, 37]. We also identified a glutathione S-transferase member (HPGDS) as well as an additional member of methionine sulfoxide reductases (msrA), which are important enzymes involved in the repair of proteins damaged by oxidative stress [38, 39]. Finally, myopeptidocytes express the LDL-receptor LRP2 and its binding partner apolipoprotein M (ApoM) [40], suggesting a role in lipid metabolism and consistent with the observation myopeptidocytes exhibit round inclusions that may represent lipid droplets. Interestingly, lipid metabolism and redox control is tightly coupled in peroxisomes which are responsible for beta-oxidation of long-chain fatty acid [41].

Evolution of cell type function by horizontal gene transfer

Sponges feed by initial endocytosis of prey by pinacocytes and choanocytes before passing food particles on to digestive and phagocytic cells in the mesohyl [42]. These food particles often include bacteria and algae, which are protected by polysaccharide and glycoprotein cell walls [43] that require specialised enzymes to break down. Yet, those enzymes are notably absent from the metazoan digestive repertoire [44].

Mesocytes are newly discovered sponge cell types and functionally uncharacterised medium-sized cells, whose name refers to their location in the mesenchymal mesohyl [7]. Our new CoFFE annotations of previously unannotated mesocyte markers include genes such as expansin (yoaJ), glucan endo-1,3-beta-glucosidase (BG3), and spore cortex-lytic enzyme (sleB), all hydrolases that specifically degrade cell walls, cellulase, chitin, other polysaccharides [44–47] and proteins (Table 1). It is thus tempting to speculate that mesocytes represent cells specialised to digest polysaccharides that are otherwise difficult to hydrolyse.

CoFFE annotations of these mesocyte marker genes reveal that many have highest structural similarity to bacteria and plant/algal proteins, suggesting they may have arisen via horizontal gene transfer (HGT). A previous study analysing origins of genes in the marine

Annotation	Annotation origin	Taxonomy	Function
Aminohydrolase	legacy EggNOG	Bacteria	hydrolase acting on aminogroups
GMHA	legacy EggNOG	Bacteria	xanthan biosynthesis
Metallopeptidase M20	legacy EggNOG	Bacteria	metallopeptidase
yoaJ	refined by CoFFE	Bacteria	cell wall degradation
sleB	refined by CoFFE	Bacteria	hydrolase activity, cell wall organization
T5orf172	new annotation	Bacteria	hydrolase activity
BG3	new annotation	Streptophyta	cellulase activity
cellulase A family member	new annotation	Streptophyta	cellulase activity
Endochitinase	new annotation	Streptophyta	endochitinase
Chitinase class I	new annotation	Streptophyta	endochitinase

Table 1. Hydrolytic enzymes in *Spongilla* mesocytes.

demosponge *Amphimedon queenslandica* identified a high proportion of polysaccharide   151
hydrolases and metallopeptidases originating by HGT [48]. Analysis of the GC-content and   152
codon usage of genes shown in Table 1 compared to the entire Spongilla transcriptome revealed   153
no significant deviation for putative HGT genes (Figure 3A), consistent with the hypothesis   154
that they are present in the *Spongilla* genome. To confirm this and explore the relative timing   155
of HGT events, we searched for homologs of all proteins shown in Table 1 in openly available   156
genomes of sponge species (Figure 3B). We detected every protein in at least one other sponge.   157
The only protein present in all analysed sponges is a metallopeptidase, similar to that detected   158
in ctenophore comb jellies [49], indicating this transfer may have occurred prior to the last   159
common ancestor of all animals. Strikingly, we found that three hydrolases were shared by only   160
freshwater sponges, suggesting mesocyte specific functions may have expanded via HGT events   161
after colonisation of new freshwater environments. Finally, the presence of predicted chitinases   162
is intriguing, as chitin has been shown to be an important structural component of *Spongilla*   163
lacustris [50]. This finding suggests another possible role for mesocytes as structural remodelers   164
of the sponge skeleton made of polysaccharides.   165

Identifying novel well-folded proteins in *Spongilla lacustris*

Using the CoFFE pipeline with a stringent bit score cut-off and augmenting it with sequence-based annotations we annotated a total of 26633 out of 41943 predicted proteins. The remaining 15312 proteins may represent incomplete fragments, untranslated sequences, or sponge lineage-specific genes. Notably, we found 3875 unannotated proteins with a pLDDT score greater than 70, indicating well-folded structures. Although many of these had poor FoldSeek alignments falling below our accepted bit score threshold, 316 had no FoldSeek hit whatsoever. Manual inspection revealed that the overwhelming majority of these are predicted to be long helices, except for 35 non-helical structures. To ensure these hadn't somehow eluded sequence similarity searches, we used NCBI BLASTp to identify potential homologs in the NR database, even very remote ones (Suppl. Table 2). Seven of the sequences find no matches at any theshold nor get significant PFAM domain hits. Notably, several of them are broadly expressed in *Spongilla* cell types, presenting prime candidates for truly novel structures.

Fig 2. Signalling pathways in *Spongilla* pinacocytes A) Dotplot of pinacocyte signalling and effector genes. Cell types of the pinacocyte family are encased by a red square. Genes on blue background are annotated by sequence-based methods ("legacy"). Genes on yellow background are annotated by CoFFE. B) Superposition of *Spongilla* FGF (blue, 61 - 230 aa) and *Gallus gallus* FGF4 (UniProtID: P48804) (yellow, 54 - 194 aa) (RMSD = 0.89 over 543 atoms). C) Superposition of *Spongilla* ephrin (blue, 1 - 153 aa) and *C. elegans* efn-3 (UniProtID: Q19475) (yellow, 29 - 179 aa) (RMSD = 1.59 over 580 atoms). In both cases the structural similarity is apparent despite their low sequence identity of 11.8% and 22% respectively. Superpositions were created using the `super` command in PyMOL (v2.3.5). Cell type abbreviations: incPin - incurrent pinacocytes; apnPn - apendopinacocytes; lph - lophocytes; basPin - basopinacocytes; scp - sclerophorocytes; met - metabolocytes; chb - choanoblasts; cho - choanocytes; apo - apopylar cells; myp - myopeptidocytes; amb - amoebocytes; grl - granulocytes; nrd - neuroid cells; mes - mesocytes; arc - archaeocytes; scl - sclerocytes.

Fig 3. ROS metabolism and redox-control in Myopeptidocytes Myopeptidocytes differentially express multiple genes involved in redox control and ROS defence. Genes in black and green have been annotated using sequence based methods. Green proteins are known to interact. Genes in red have been identified using CoFFE.

Fig 4. Horizontal gene transfer of mesocyte marker genes A) Distribution of GC-content and codon usage of mesocyte marker genes of non-metazoan origin (colored dots) compared to the entire *Spongilla* transcriptome (blue background). B) Heatmap showing scores of best search hit of *Spongilla* mesocyte marker genes in various sponge species. The green dot denotes the last common freshwater sponge ancestor.

Discussion

We predicted the protein structures of the entire proteome of the freshwater sponge *Spongilla* *lacustris* and aligned them against all currently available databases. This approach increased the annotation of the proteome from 40% to 60%, an approximately 50% increase. For proteins with annotation from both sources, we found that in more than 90% of cases, sequence-based and structure-based annotation agreed at least to the protein superfamily level, a finding supported by recent work with well-annotated model species [51].

In ∼5% of cases CoFFE and eggnog-mapper annotations do not agree. This may result from inaccurate or incomplete orthology groups in the eggNOG database, or because the proteins evolved similar structures independently in evolution. Although protein structure may be more directly related to function, structural annotations depend on the quality of the predicted structure. Interestingly, pLDDT values for this category are on average only marginally lower than in cases were sequence and structure annotations agree only marginally lower than in cases were sequence and structure annotations agree, suggesting poor structural predictions are unlikely to be driving the incongruence. Considering that careful evaluation of sequence-based phylogenetic trees can identify a protein's evolutionary history [52], it will be interesting to explore the extent to which annotations based on structure provide better predictors of function than evolutionary homology.

To demonstrate the usefulness of structural similarity for functional annotation we revisited the annotation of marker genes previously identified from *Spongilla* scRNA-seq data [7]. In the epithelial pinacocyte family, this significantly improved sequence-based annotation via the detection of key players in FGF, TGF-β, as well as ephrin signalling. For the digestive peptidocyte family, we were able to infer complex modes of redox regulation as a possible function of the myopeptidocytes. Finally, we detected polysaccharide and protein hydrolyzing enzymes of bacterial and plant/algal origin in the enigmatic mesocytes which strongly suggests that these digest cell walls of endocytosed prey. This example reveals cell types whose functional role is largely defined by genes acquired via HGT from other pro- and eukaryotic organisms. Recent evidence from nematodes showed an expansion of dietary possibilities after HGT of a eukaryotic cellulase gene [53]. This is remarkable both from a cell type evolutionary perspective [54] and because such innovation had previously escaped our notice due to extensive

sequence divergence following the HGT event. By re-evaluating marker genes from the *Spongilla* cell type atlas it became obvious that structure-based annotations not only expand the molecular context of known cell type functions but also allow substantiated hypotheses about previously unexplored cell types. These hypotheses can serve as a next step for further investigation and discoveries.

Well-folded *Spongilla* proteins without any annotation constitute intriguing candidates for novel protein folds and functions specific to sponges. Many of these proteins are predicted to be long alpha helices. This could either be an artifact of spuriously translated proteins (as described in [55]) or present sponge specific proteins for structural integrity of the sponge body, similar to the "constant force spring" character of naturally occurring single α-helices [56]. In addition, unannotated proteins with globular structures, expressed in the scRNA-seq data, are most probably functional and offer a great resource to study the evolutionary emergence of protein folds.

Conclusion

The lack of reliable functional annotation has so far been a major bottleneck in the analysis of non-model species, in particular those separated from traditional models by large phylogenetic distances. Here, we demonstrate that by exploiting the evolutionary conservation of protein structure it is possible to dramatically improve protein functional annotations in non-model species. Although protein structural predictions for an entire proteome might be outside the technical capabilities of many labs, the pipeline described here can be used to query individual highly informative candidate genes from proteomics or single-cell -omics experiments. Additionally, plans to continously expand the AlphaFold database [9] have been announced. It is reasonable to expect that in the future protein sequences deposited in public databases will automatically receive predicted structures, paving the way for unique insights into biological functions across the tree of life.

Materials and methods

In all boxplots, the box extends from the lower to upper quartile values of the data, with a line at the median. The whiskers represent the $5 - 95\%$ percentiles.

Sequence-based annotation of the *Spongilla lacustris* proteome

Juvenile freshwater sponges (*Spongilla lacustris*), grown from gemmules, were used for bulk RNA isolation and sequencing. *De novo* transcriptome assembly with Trinity, returned 62180 putative isoforms, covering 95.2% of Metazoan BUSCOs [57]. To identify putative proteins, Transdecoder [58] (version 3.0.1) was used with a minimum open reading frame length of 70 amino acids, resulting in 41945 putative proteins. The longest putative protein per gene ID was kept. The resulting predicted proteome was annotated by EggNOG mapper [12, 59] (v2.1.7, default settings) via the website.

Legacy annotation

Musser *et al.* [7] used the putative proteins to create a phylome by constructing gene/protein trees for each protein (available online). The phylome information was used to refine the assignment of transcripts to genes. In some cases, 3' and 5' fragments of a gene were assigned to two different transcripts. These fragments were merged into the same merged gene name using the gene tree information. Functional annotations were supplemented by EggNOG mapper (v1) and `blastp` searches against human RefSeq (default parameters). This annotation was used in the original *Spongilla* scRNA sequencing publication and is present in the single-cell data. The legacy annotation was used for the single-cell data analysis, but not for the comparison between the CoFFE pipeline and the sequence-based annotation transfer.

Structure-based annotation of the *Spongilla lacustris* proteome

CoFFE and constituent tools

We designed a simple pipeline that goes from sequence to predicted structure to functional annotation (symbolically, **Co**lab**F**old-**F**oldSeek-**E**ggNOG, or CoFFE), and used it to annotate the *Spongilla lacustris* predicted proteome. We used ColabFold to predict structures and subsequently aligned the predicted structures against all currently available (solved and

predicted) protein structures using FoldSeek. Finally, we used structural similarity to transfer annotations from the best structural targets to their corresponding *Spongilla* protein queries. In the following we show an overview of the tools in use and a more detailed description of the CoFFE pipeline.

MMseqs2 [60] is a software suite for sequence-sequence, sequence-profile, and profile-profile search and clustering. It is orders of magnitude faster than BLAST at the same sensitivity and is widely adopted [12].

AlphaFold2 [5] is a neural network-based model that predicts protein three-dimensional structures from sequence, regularly achieving atomic accuracy even in cases where no similar structure is known. AlphaFold is widely considered to have revolutionised the field of structural bioinformatics, greatly outperforming the state of the art in the most recent iteration of the CASP challenge [61].

AlphaFold quantifies prediction confidence by pLDDT, the predicted local distance difference test on the $C\alpha$ atoms. Regions with pLDDT > 90 are modelled to high accuracy; regions with $70 <$ pLDDT < 90 should have a generally good backbone prediction; regions with $50 <$ pLDDT < 70 are low confidence and should be treated with caution; regions with pLDDT < 50 should not be interpreted and probably are disordered.

ColabFold [8] is a pipeline that combines fast homology searches via MMseqs2 [60] with AlphaFold2 [5] to predict protein structures 40 to 60 times faster than the original AlphaFold2.

FoldSeek [6] enables fast and sensitive comparison of large structure databases. FoldSeek's key innovation lies in the appropriate translation of structure states to a small alphabet, thus gaining access to all the heuristics of sequence search algorithms.

The CoFFE pipeline

A visual representation of the CoFFE pipeline can be found in supplement figure XX. To predict structures, we adapted the ColabFold pipeline as outlined in
`https://colabfold.mmseqs.com/`

Multiple sequence alignment generation: We downloaded reference sequence databases (UniRef30, ColabFold DB) (databases.sh, adapted from setup_databases.sh) and calculated indices locally (create_index.sh). We were interested in homology detection at the limit of the twilight zone, so UniRef30, a 30% sequence identity clustered database based on UniRef100 [62],

was the adequate choice. Similarly, we indexed the *Spongilla* predicted proteome ₂₉₁

(spongilla_create_index.sh). We calculated MSAs for each *Spongilla* predicted protein using ₂₉₂

MMseqs2 [60] (align.sh, adapted from colabfold_search.sh). ₂₉₃

Structure prediction: We predicted structures for all *Spongilla* predicted proteins using ₂₉₄

ColabFold [8] as a wrapper around AlphaFold2. ₂₉₅

We split the MSAs in 32 batches and submitted each one to the EMBL cluster system ₂₉₆

(managed by `slurm` [63]); we used default arguments but added `--stop-at-score 85`. ₂₉₇

(milot.sh, M. Mirdita, personal communication). The calculations were done on NVIDIA A100 ₂₉₈

GPUs, on computers running CentOS Linux 7. We used GCC [64] version 10.2.0 and CUDA ₂₉₉

version 11.1.1-GCC-10.2.0 [65]. We processed the resulting PDB-formatted model files with ₃₀₀

Biopython's PDB module [66]. ₃₀₁

Structure search and annotation transfer: Structural search was conducted using ₃₀₂

FoldSeek. FoldSeek allows fast comparison of large structural databases. We downloaded PDB, ₃₀₃

SwissProt, and AlphaFold DB (fs_pdb.sh, fs_sp.sh, and fs_afdb.sh, respectively). For each ₃₀₄

Spongilla protein we kept the best-scoring AlphaFold2 model, and used them to construct a ₃₀₅

FoldSeek database fs_query.sh. These models were then used to search against the three ₃₀₆

structural databases. For each search we kept the FoldSeek hit with the highest corrected bit ₃₀₇

score in each database and aggregated the three result tables (AlphaFoldDB, PDB, SwissProt) ₃₀₈

into one. We imposed a bit score cutoff (1xe-05) on FoldSeek hits based on their bimodal ₃₀₉

distribution. Annotations of the best hits were gathered from either UniProt via its API [67] or ₃₁₀

through EggNOG mapper (v2.1.7, default settings) [12] by using the sequences of the best ₃₁₁

structural hits (pulled by UPIMAPI [68]). To facilitate downstream analysis, we extracted ₃₁₂

summary tables from each resource type. This procedure can be found in the corresponding ₃₁₃

notebook. A total of 1401 proteins received sequence annotation but their FoldSeek best hits ₃₁₄

were below the bit score cutoff. ₃₁₅

Differential gene expression in single-cell transcriptomics data ₃₁₆

We obtained the processed Seurat file from [7], and downloaded the lists of differentially ₃₁₇

expressed genes of clusters, cell types, and cell type clades from the supplemental material of ₃₁₈

the same publication (Supplemental Data S1 to S3; file science.abj2949_data_s1.xlsx; tabs "Diff. ₃₁₉

exp. 42 clusters", "Diff. Exp. cell types", "Cell type clade genes (OU tests)"). The single-cell ₃₂₀

data operates on the level of genes, so we transformed the sequence-derived and CoFFE 321
annotations by merging isoform entries and keeping the entry with the best bit score. 322

We used the legacy annotation included in the file (phylome, EggNOG-mapper, and 323
BLASTp-based), the sequence-derived annotation, and the CoFFE output to propose names for 324
Spongilla proteins. We prioritized sequence-derived annotations (legacy annotation, EggNOG 325
preferred name, EggNOG description) and fell back to CoFFE (CoFFE preferred name, CoFFE 326
description) when there were none. We produced dotplots for the top 200 differentially 327
expressed genes in each cell type and manually inspected them. We focused on the terminal 328
(named) cell types; owing to the presence of continuously differentiating stem cells, the 329
single-cell data contains many clusters of maturing or differentiating cells whose expression 330
patterns do not distinguish them from their mature counterparts. Code and detailed 331
explanations are available in the corresponding notebook. 332

Protein structure visualisation and superposition 333

In order to visualize predicted structures from the *Spongilla* proteome, we used PyMOL Version 334
2.3.5 [69]. Superposition with their respective best Foldseek hit was carried out using the `super` 335
command. `super` creates sequence- independent superpositions and is more reliable for protein 336
pairs with low sequence similarity. 337

Detection of ephrin orthologs in *Spognilla* using hmmer 338

To validate the ephrin ortholog detected by CoFFE in *Spongilla*, we recapitulated a previous 339
effort to detect ephrins in different species using extensive hmmer protein profile search [24]. 340
For this, we performed a hmmer search (v3.3.2) using the ephrin Pfam profile (PF00812) 341
against the *Spongilla* proteome using default settings. 342

Detecting HGT in Spongilla 343

To detect HGT we followed the proposal by Degnan [70] and sought to get multiple indications 344
for HGT. In particular: phylogenetic evidence that the candidate gene is more closely related to 345
foreign than to animal genes; genome data showing the candidate gene assembles into a 346
contiguous stretch of DNA with neighboring genes unambiguously of animal origin (this 347
requires, of course, the availability of a sequenced and assembled animal genome; the more 348

complete the assembly, the more confident the HGT identification); and gene sequence revealing ₃₄₉ metazoan-like compositional traits, including presence of introns, GC content and codon usage. ₃₅₀ Where possible, gene expression data showing active transcription of candidate genes in animal ₃₅₁ cell nuclei can enormously strengthen a case, and also addresses the issue of whether or not the ₃₅₂ HGT-acquired gene is active in its new genomic context. ₃₅₃

We used `https://spongebase.net` to obtain transcriptomes and genomes for 13 sponge ₃₅₄ species, and obtained a 14th one directly from its repository ₃₅₅ (`https://bitbucket.org/molpalmuc/tethya_wilhelma-genome/`). We built sequence ₃₅₆ databases with MMseqs2 version 12-113e3 and searched against them (mmseqs easy-search) ₃₅₇ with the protein sequences of all isoforms of the HGT candidates. The resulting alignments ₃₅₈ were filtered to keep the best-scoring alignment per species per gene. ₃₅₉

The table of best hits per genome was visualised in Python, and a sponge phylogeny was ₃₆₀ added manually based on [71] (also A. Riesgo, personal communication). ₃₆₁

Acknowledgments

The authors thank Milot Mirdita and Martin Steinegger for discussions about ColabFold usage
and technical support; Alexandros Pittis for his technical feedback; the Arendt lab for
discussions; the ModelArchive team for their support in publishing the predicted structures;
Ana Riesgo for sharing demosponge phylogenies. NP was funded by MSCA individual
fellowship 101031984/DeCoDe Platy. FR has received funding from the European Union's
Horizon 2020 research and innovation programme under the Marie Sklosowska-Curie grant
agreement No. 764840/IGNITE as well as EMBL International PhD Program.

A CoFFE overview

Fig 5. Overview of CoFFE pipeline ColabFold is called with a proteome as input. It produces multiple sequence alignments for each protein using MMseqs2 and predicts structures for each structure using AlphaFold2. The best model per protein is kept and used as query in FoldSeek to search against PDB, AlphaFoldDB, and SwissProt. This creates an m8-formatted table for each database, where the significant hits for each query protein are kept. The tables are merged by keeping the target with the highest bit score. The sequence of the best hit is then retrieved from UniProt and used as a query for EggNOG-mapper. This finds the target sequence itself and retrieves the EggNOG entry, including preferred name and description.

B Extracting quality indicators from the CoFFE pipeline and comparing against physico-chemical parameters of proteins

To get a better understanding of the relationship between the various parameters of the CoFFE pipeline as well as physico-chemical parameters of the proteins, we assessed the correlation between them (Fig. 6). We hoped to find a combination of measures that together would be a robust indicator of prediction quality for CoFFE. Additionally we sought to explore the relationship between structural prediction quality and various intrinsic protein parameters.

Most obviously, query length, (uncorrected) structural bit score, sequence bit score (EggNOG-mapper) as well as molecular weight correlate very strongly. Longer proteins (= heavier proteins) will have more or longer aligned regions with their targets, and as the bit score is additive, will accumulate higher bit scores, regardless of whether the alignment happens in structure or sequence space. It seems these measures can be used interchangeably; out of principle, we would prefer to use query length, as it is the least derived.

A way to overcome the dependence on length is to normalize alignment scores by the length of the aligned region (*corrected bit score (FS)*). This measure correlates with the percent structure state identity. This is equivalent to percent sequence identity correlating strongly with the bit score in a sequence similarity search, and is not surprising at all. More interestingly, these measures, together with the MSA size, correlate with the average pLDDT as a measure for protein prediction quality. Nevertheless, this correlation is less pronounced, suggesting to look at pLDDT independently.

Relative alignment length (FS) (percentage of the query (*Spongilla*) structure aligned with the best target structure) does not correlate too strongly with other parameters.

Three things become apparent: first, higher pLDDT scores generally correlate with higher FoldSeek bit scores. This means that FoldSeek does a better job of identifying structural similarity for well-folded proteins. This seems intuitive: the atomic coordinates of well-folded proteins will be more deterministic than disordered proteins, so well-folded orthologs will overlap structurally over their entire length, while heavily disordered orthologs might only overlap structurally over very short patches. Additionally, high pLDDTs are reached when sufficiently large MSAs are available. Large MSAs contribute more information to AlphaFold's Evoformer module, leading to a higher predictive power.

Second, the relationship between pLDDT and (corrected) FoldSeek bit score is not dependent on the length/weight of the protein. Independent of their size, proteins structures are predicted equally good or bad.

Third, pLDDT as well as FoldSeek bit score do not correlate with various physico-chemical parameters of the proteins. Protein structure prediction seems to be independent from gravy score [75], aromaticity [74], isoelectric point, instability index [72] as well as the average of flexibility index [73] of proteins. Consequently, FoldSeek bit score is not dependent on these parameters.

Next, we were interested in the distribution of these parameters across different annotation categories. Proteins either got an annotation by CoFFE and/or a sequence based EggNOG-mapper search or not. (*'Annotation'* can entail a name or a description of the protein). After applying the FoldSeek bit score cut-off, we gain a total of 16285 proteins with annotations from CoFFE and EggNOG-mapper, 15599 proteins without annotations, 8914 proteins with CoFFE annotations only and 1147 proteins with sequence based annotations only (Fig. 7A). Comparing different CoFFE parameters, it becomes clear again that large MSAs are helpful for gaining CoFFE based annotations (Fig. 7B). Nevertheless, MSAs are not necessary for getting CoFFE based annotations, showing that AlphaFold is able to work well even without evolutionary information available. Comparing pLDDTs it becomes obvious that proteins without any annotation usually have worse structural prediction quality (Fig. 7C) and vice versa. Interestingly, Proteins with only sequence based annotation have on average relatively high pLDDT values. As expected, proteins with CoFFE annotations have significantly higher FoldSeek bit scores, reflecting the usefulness of the chosen bit score cut-off (Fig. 7D).

Comparing different annotation categories based on the distribution of physico-chemical parameters supports the statement above that they are not correlating with CoFFE parameters. Apart from molecular weight (Fig. 8B), none of the parameters such as gravy score, aromaticity index, isoelectric point, instability index and average flexibility score (Fig. 8B-F) differ significantly between the annotation categories.

isoform	q len	#MSA	%seq. id.	bit score	plddt	BLASTp
c88419_g1_i1	154	462	0.337	580.0	85.01	CRN5
c104256_g1_i1	375	163	0.272	785.0	82.42	CRN5
c104256_g1_i2	299	184	0.267	781.0	90.35	CRN5
c104256_g1_i3	336	167	0.262	777.0	88.39	CRN5
c104256_g1_i4	332	158	0.263	796.0	90.26	CRN5
c104256_g1_i6	253	196	0.275	774.0	92.81	CRN5
c94352_g2_i1	127	1725	0.500	299.0	86.06	DD3-3
c89711_g1_i1	144	2424	0.441	380.0	86.73	DD3-3
c89711_g2_i1	75	2340	0.611	210.0	77.48	DD3-3
c112476_g1_i1	179	1450	0.352	434.0	83.19	DD3-3
c103626_g1_i1	670	3392	0.324	2224.0	91.33	DD3-3
c103626_g1_i2	670	3424	0.329	2258.0	91.18	DD3-3
c103626_g2_i1	667	3427	0.326	2215.0	90.01	DD3-3
c102584_g1_i1	578	3328	0.355	2091.0	81.55	DD3-3
c102584_g1_i2	187	1994	0.416	522.0	89.04	DD3-3
c101972_g1_i1	778	3481	0.378	2536.0	85.58	DD3-3
c101972_g1_i2	778	3481	0.378	2534.0	85.58	DD3-3
c101972_g1_i3	778	3481	0.378	2537.0	85.54	DD3-3
c104453_g2_i1	359	368	0.177	662.0	75.80	foll. epith. yolk prot. subunit
c104453_g2_i2	280	356	0.172	658.0	89.90	foll. epith. yolk prot. subunit
c104453_g2_i3	340	378	0.185	634.0	78.61	foll. epith. yolk prot. subunit
c104453_g2_i4	236	374	0.183	594.0	83.73	foll. epith. yolk prot. subunit
c104453_g3_i1	182	341	0.117	298.0	70.54	foll. epith. yolk prot. subunit
c98985_g1_i2	73	1852	0.138	45.0	70.12	XPO5 (Exportin-5)
c88005_g3_i1	130	3213	0.274	363.0	94.66	hemopexin repeat-containing protein
c70965_g1_i1	169	299	0.209	354.0	77.62	pinocchio
c103612_g1_i1	89	1253	0.152	46.0	73.88	kinesin-like protein KIF23
c102197_g1_i3	312	257	0.179	649.0	86.28	bamB
c100331_g2_i1	281	43509	0.219	821.0	82.62	hydralysin-2-like
c93555_g2_i2	77	439	0.109	80.0	87.77	?
c86044_g1_i1	87	7052	0.077	90.0	88.87	?
c83472_g1_i1	98	553	0.177	90.0	84.77	?
c108675_g1_i1	73	3174	0.224	51.0	78.17	?
c100701_g1_i2	69	1	0.123	61.0	81.77	?
c100429_g1_i1	79	43996	0.152	85.0	71.74	?

Table 2. High confidence structures with non-helical appearance. The last column contains a summary of the significant BLASTp results in the `nr` database. Question marks denote peptides who did not find any named sequence homologs or significant PFAM domain hits.

Fig 6. Pairwise correlation of metrics from the CoFFE pipeline as well as physico-chemical protein parameters.

instability index: measure for proteins stability based on Guruprasad et al. [72]. *flexibility average*: average of structural flexibility scores across all amino acids of a protein based on Vihinen et al. [73]. *rel. ali. length (FS)*: percentage of the query (*Spongilla*) structure aligned with the best target structure. *isoelectric points*: prediction of the isoelectric point of a protein. *aromaticity*: measure for protein aromaticity calculated as the relative frequency of Phe/Trp/Tyr [74]. *gravy*: measure for the hydropathic caracter of a protein based on [75]. *bit score (FS)*: bit score of the FoldSeek alignment between the query *Spongilla* structure and the structure of the best FoldSeek target. *bit score (EggNOG-seq)*: bit score of the EggNOG alignment between the query *Spongilla* peptide sequence and the best target sequence. *Spongilla query length (AF)*: the query length of the input peptide. *molecular weight*: molecular weight of the protein. *plDDT (AF)*: average pLDDT of the predicted structure. *Spongilla MSA size (AF)*: size of the input MSA used for AlphaFold. *best match seq. id. (FS)*: structure state identity in the alignment between the query. *corrected bit score (FS)*: the FoldSeek bit score divided by alignment length.

Fig 7. Comparison of CoFFE parameters in different annotation categories.

A) Protein count in different annotation categories. Annotation includes either protein name or description. Protein counts are calculated *after* applying the FoldSeek bit score cut-off. *Structure Sequence*: Annotation by EggNOG-mapper sequence search as well as CoFFE *No structure No sequence*: No annotation by EggNOG-mapper sequence search as well as CoFFE. *Structure No sequence*: Annotation by CoFFE but no sequence based annotation by EggNOG-mapper. *No structure Sequence*: Annotation by EggNOG-mapper sequence search but not by CoFFE. B) Distribution of AlphaFold MSA sizes across annotation categories. C) Distribution of AlphaFold pLDDTs across annotation categories. D) Distribution of FoldSeek (uncorrected) bit scores across annotation categories.

Fig 8. Comparison of physico-chemical protein parameters in different annotation categories.

A) Distribution of molecular weights across annotation categories. B) Distribution of gravy scores across annotation categories. C) Distribution of aromaticity scores across annotation categories. D) Distribution of isoelectric points across annotation categories. E) Distribution of instability indices across annotation categories. F) Distribution of flexibility score averages across annotation categories.

C Differentially expressed genes per cell type

Fig 9. The percentage of marker genes per cluster as per Musser *et al.* that are annotated by different methods.

References

1. Rost B. Twilight zone of protein sequence alignments. Protein engineering. 1999;12(2):85–94.

2. Kabir MN, Wong L. EnsembleFam: towards more accurate protein family prediction in the twilight zone. BMC bioinformatics. 2022;23(1):1–20.

3. Illergård K, Ardell DH, Elofsson A. Structure is three to ten times more conserved than sequence—a study of structural response in protein cores. Proteins: Structure, Function, and Bioinformatics. 2009;77(3):499–508.

4. Waterhouse A, Bertoni M, Bienert S, Studer G, Tauriello G, Gumienny R, et al. SWISS-MODEL: homology modelling of protein structures and complexes. Nucleic acids research. 2018;46(W1):W296–W303.

5. Jumper J, Evans R, Pritzel A, Green T, Figurnov M, Ronneberger O, et al. Highly accurate protein structure prediction with AlphaFold. Nature. 2021;596(7873):583–589.

6. van Kempen M, Kim S, Tumescheit C, Mirdita M, Söding J, Steinegger M. Foldseek: fast and accurate protein structure search. bioRxiv. 2022;.

7. Musser JM, Schippers KJ, Nickel M, Mizzon G, Kohn AB, Pape C, et al. Profiling cellular diversity in sponges informs animal cell type and nervous system evolution. Science. 2021;374(6568):717–723.

8. Mirdita M, Schütze K, Moriwaki Y, Heo L, Ovchinnikov S, Steinegger M. ColabFold - Making protein folding accessible to all. bioRxiv. 2022;doi:10.1101/2021.08.15.456425.

9. Varadi M, Anyango S, Deshpande M, Nair S, Natassia C, Yordanova G, et al. AlphaFold Protein Structure Database: Massively expanding the structural coverage of protein-sequence space with high-accuracy models. Nucleic acids research. 2022;50(D1):D439–D444.

10. Boutet E, Lieberherr D, Tognolli M, Schneider M, Bansal P, Bridge AJ, et al. UniProtKB/Swiss-Prot, the manually annotated section of the UniProt KnowledgeBase: how to use the entry view. In: Plant Bioinformatics. Springer; 2016. p. 23–54.

11. Burley SK, Bhikadiya C, Bi C, Bittrich S, Chen L, Crichlow GV, et al. RCSB Protein Data Bank: powerful new tools for exploring 3D structures of biological macromolecules for basic and applied research and education in fundamental biology, biomedicine, biotechnology, bioengineering and energy sciences. Nucleic acids research. 2021;49(D1):D437–D451.

12. Cantalapiedra CP, Hernández-Plaza A, Letunic I, Bork P, Huerta-Cepas J. eggNOG-mapper v2: functional annotation, orthology assignments, and domain prediction at the metagenomic scale. Molecular biology and evolution. 2021;38(12):5825–5829.

13. Mistry J, Chuguransky S, Williams L, Qureshi M, Salazar GA, Sonnhammer EL, et al. Pfam: The protein families database in 2021. Nucleic acids research. 2021;49(D1):D412–D419.

14. Elliott GR, Leys SP. Coordinated contractions effectively expel water from the aquiferous system of a freshwater sponge. Journal of Experimental Biology. 2007;210(21):3736–3748.

15. Nickel M, Scheer C, Hammel JU, Herzen J, Beckmann F. The contractile sponge epithelium sensu lato–body contraction of the demosponge Tethya wilhelma is mediated by the pinacoderm. Journal of Experimental Biology. 2011;214(10):1692–1698.

16. Adamska M, Degnan SM, Green KM, Adamski M, Craigie A, Larroux C, et al. Wnt and TGF-β expression in the sponge Amphimedon queenslandica and the origin of metazoan embryonic patterning. PloS one. 2007;2(10):e1031.

17. Schippers KJ, Nichols SA. Evidence of signaling and adhesion roles for β-catenin in the sponge Ephydatia muelleri. Molecular Biology and Evolution. 2018;35(6):1407–1421.

18. Li H, Tao C, Cai Z, Hertzler-Schaefer K, Collins TN, Wang F, et al. Frs2α and Shp2 signal independently of Gab to mediate FGF signaling in lens development. Journal of cell science. 2014;127(3):571–582.

19. Mossahebi-Mohammadi M, Quan M, Zhang JS, Li X. FGF signaling pathway: a key regulator of stem cell pluripotency. Frontiers in cell and developmental biology. 2020;8:79.

20. Massagué J. TGFβ signalling in context. Nature reviews Molecular cell biology. 2012;13(10):616–630.

21. Darling TK, Lamb TJ. Emerging roles for Eph receptors and ephrin ligands in immunity. Frontiers in immunology. 2019;10:1473.

22. Kania A, Klein R. Mechanisms of ephrin–Eph signalling in development, physiology and disease. Nature reviews Molecular cell biology. 2016;17(4):240–256.

23. Arcas A, Wilkinson DG, Nieto MÁ. The Evolutionary History of Ephs and Ephrins: Toward Multicellular Organisms. Molecular biology and evolution. 2020;37(2):379–394.

24. Krishnan A, Degnan BM, Degnan SM. The first identification of complete Eph-ephrin signalling in ctenophores and sponges reveals a role for neofunctionalization in the emergence of signalling domains. BMC evolutionary biology. 2019;19(1):1–17.

25. Eddy SR. Accelerated profile HMM searches. PLoS computational biology. 2011;7(10):e1002195.

26. Horbelt D, Denkis A, Knaus P. A portrait of Transforming Growth Factor β superfamily signalling: Background matters. The international journal of biochemistry & cell biology. 2012;44(3):469–474.

27. Turner N, Grose R. Fibroblast growth factor signalling: from development to cancer. Nature Reviews Cancer. 2010;10(2):116–129.

28. Bonasoro F, Wilkie IC, Bavestrello G, Cerrano C, Carnevali M. Dynamic structure of the mesohyl in the sponge Chondrosia reniformis (Porifera, Demospongiae). Zoomorphology. 2001;121(2):109–121.

29. Ameziane-El-Hassani R, Morand S, Boucher JL, Frapart YM, Apostolou D, Agnandji D, et al. Dual oxidase-2 has an intrinsic Ca2+-dependent H2O2-generating activity. Journal of Biological Chemistry. 2005;280(34):30046–30054.

30. Miller CJ, Rose AL, Waite TD. Importance of iron complexation for Fenton-mediated hydroxyl radical production at circumneutral pH. Frontiers in Marine Science. 2016;3:134.

31. Ueno N, Takeya R, Miyano K, Kikuchi H, Sumimoto H. The NADPH oxidase Nox3 constitutively produces superoxide in a p22phox-dependent manner: its regulation by oxidase organizers and activators. Journal of Biological Chemistry. 2005;280(24):23328–23339.

32. Protchenko O, Rodriguez-Suarez R, Androphy R, Bussey H, Philpott CC. A screen for genes of heme uptake identifies the FLC family required for import of FAD into the endoplasmic reticulum. Journal of Biological Chemistry. 2006;281(30):21445–21457.

33. Missiakas D, Georgopoulos C, Raina S. Identification and characterization of the Escherichia coli gene dsbB, whose product is involved in the formation of disulfide bonds in vivo. Proceedings of the National Academy of Sciences. 1993;90(15):7084–7088.

34. Zhao Y, Zhang J, Li H, Li Y, Ren J, Luo M, et al. An NADPH sensor protein (HSCARG) down-regulates nitric oxide synthesis by association with argininosuccinate synthetase and is essential for epithelial cell viability. Journal of Biological Chemistry. 2008;283(16):11004–11013.

35. Chang TS, Jeong W, Woo HA, Lee SM, Park S, Rhee SG. Characterization of mammalian sulfiredoxin and its reactivation of hyperoxidized peroxiredoxin through reduction of cysteine sulfinic acid in the active site to cysteine. Journal of Biological Chemistry. 2004;279(49):50994–51001.

36. Pei DS, Jia PP, Luo JJ, Liu W, Strauss PR. AP endonuclease 1 (Apex1) influences brain development linking oxidative stress and DNA repair. Cell death & disease. 2019;10(5):1–14.

37. Hu Z, Ding X, Ji Y, Liu X, Ding Z. APEX1 protects against oxidative damage-induced cardiomyocyte apoptosis. Biocell. 2021;45(3):745.

38. Korbecki J, Baranowska-Bosiacka I, Gutowska I, Chlubek D. The effect of reactive oxygen species on the synthesis of prostanoids from arachidonic acid. J Physiol Pharmacol. 2013;64(4):409–421.

39. Boschi-Muller S, Azza S, Sanglier-Cianferani S, Talfournier F, Van Dorsselear A, Branlant G. A sulfenic acid enzyme intermediate is involved in the catalytic mechanism

of peptide methionine sulfoxide reductase fromEscherichia coli. Journal of Biological Chemistry. 2000;275(46):35908–35913.

40. Faber K, Hvidberg V, Moestrup SK, Dahläck B, Nielsen LB. Megalin is a receptor for apolipoprotein M, and kidney-specific megalin-deficiency confers urinary excretion of apolipoprotein M. Molecular endocrinology. 2006;20(1):212–218.

41. Ayala A, Muñoz MF, Argüelles S. Lipid peroxidation: production, metabolism, and signaling mechanisms of malondialdehyde and 4-hydroxy-2-nonenal. Oxidative medicine and cellular longevity. 2014;2014.

42. Imsiecke G. Ingestion, digestion, and egestion in Spongilla lacustris (Porifera, Spongillidae) after pulse feeding with Chlamydomonas reinhardtii (Volvocales). Zoomorphology. 1993;113(4):233–244.

43. Imam SH, Buchanan MJ, Shin HC, Snell WJ. The Chlamydomonas cell wall: characterization of the wall framework. The Journal of cell biology. 1985;101(4):1599–1607.

44. Davison A, Blaxter M. Ancient origin of glycosyl hydrolase family 9 cellulase genes. Molecular Biology and Evolution. 2005;22(5):1273–1284.

45. Kerff F, Amoroso A, Herman R, Sauvage E, Petrella S, Filée P, et al. Crystal structure and activity of Bacillus subtilis YoaJ (EXLX1), a bacterial expansin that promotes root colonization. Proceedings of the National Academy of Sciences. 2008;105(44):16876–16881.

46. Zavaliev R, Levy A, Gera A, Epel BL. Subcellular dynamics and role of Arabidopsis β-1, 3-glucanases in cell-to-cell movement of tobamoviruses. Molecular Plant-Microbe Interactions. 2013;26(9):1016–1030.

47. Makino S, Ito N, Inoue T, Miyata S, Moriyama R. A spore-lytic enzyme released from Bacillus cereus spores during germination. Microbiology. 1994;140(6):1403–1410.

48. Conaco C, Tsoulfas P, Sakarya O, Dolan A, Werren J, Kosik KS. Detection of prokaryotic genes in the Amphimedon queenslandica genome. PLoS One. 2016;11(3):e0151092.

49. Hernandez AM, Ryan JF. Horizontally transferred genes in the ctenophore Mnemiopsis leidyi. PeerJ. 2018;6:e5067.

50. Ehrlich H, Kaluzhnaya OV, Brunner E, Tsurkan MV, Ereskovsky A, Ilan M, et al. Identification and first insights into the structure and biosynthesis of chitin from the freshwater sponge Spongilla lacustris. Journal of Structural Biology. 2013;183(3):474–483.

51. Monzon VA, Paysan-Lafosse T, Wood V, Bateman A. Reciprocal Best Structure Hits: Using AlphaFold models to discover distant homologues. bioRxiv. 2022;.

52. Kapli P, Yang Z, Telford MJ. Phylogenetic tree building in the genomic age. Nature Reviews Genetics. 2020;21(7):428–444.

53. Han Z, Sieriebriennikov B, Susoy V, Lo WS, Igreja C, Dong C, et al. Horizontally acquired cellulases assist the expansion of dietary range in Pristionchus nematodes. Molecular biology and evolution. 2022;.

54. Arendt D, Musser JM, Baker CV, Bergman A, Cepko C, Erwin DH, et al. The origin and evolution of cell types. Nature Reviews Genetics. 2016;17(12):744–757.

55. Monzon V, Haft DH, Bateman A. Folding the unfoldable: using AlphaFold to explore spurious proteins. Bioinformatics Advances. 2022;2(1):vbab043.

56. Wolny M, Batchelor M, Bartlett GJ, Baker EG, Kurzawa M, Knight PJ, et al. Characterization of long and stable de novo single alpha-helix domains provides novel insight into their stability. Scientific reports. 2017;7(1):1–14.

57. Simão FA, Waterhouse RM, Ioannidis P, Kriventseva EV, Zdobnov EM. BUSCO: assessing genome assembly and annotation completeness with single-copy orthologs. Bioinformatics. 2015;31(19):3210–3212.

58. Haas BJ, Papanicolaou A, Yassour M, Grabherr M, Blood PD, Bowden J, et al. De novo transcript sequence reconstruction from RNA-seq using the Trinity platform for reference generation and analysis. Nature protocols. 2013;8(8):1494–1512.

59. Huerta-Cepas J, Szklarczyk D, Heller D, Hernández-Plaza A, Forslund SK, Cook H, et al. eggNOG 5.0: a hierarchical, functionally and phylogenetically annotated orthology

resource based on 5090 organisms and 2502 viruses. Nucleic acids research. 2019;47(D1):D309–D314.

60. Steinegger M, Söding J. MMseqs2 enables sensitive protein sequence searching for the analysis of massive data sets. Nature biotechnology. 2017;35(11):1026–1028.

61. Pereira J, Simpkin AJ, Hartmann MD, Rigden DJ, Keegan RM, Lupas AN. High-accuracy protein structure prediction in CASP14. Proteins: Structure, Function, and Bioinformatics. 2021;89(12):1687–1699.

62. Suzek BE, Huang H, McGarvey P, Mazumder R, Wu CH. UniRef: comprehensive and non-redundant UniProt reference clusters. Bioinformatics. 2007;23(10):1282–1288.

63. Yoo AB, Jette MA, Grondona M. Slurm: Simple linux utility for resource management. In: Workshop on job scheduling strategies for parallel processing. Springer; 2003. p. 44–60.

64. GCC, the GNU Compiler Collection; 2020. https://web.archive.org/web/20220528080831/https://gcc.gnu.org/.

65. NVIDIA, Vingelmann P, Fitzek FHP. CUDA, release: 11.1.1; 2020. Available from: https://docs.nvidia.com/cuda/archive/11.1.1/.

66. Hamelryck T, Manderick B. PDB file parser and structure class implemented in Python. Bioinformatics. 2003;19(17):2308–2310.

67. consortium TU. UniProt: the universal protein knowledgebase in 2021. Nucleic acids research. 2021;49(D1):D480–D489.

68. Sequeira JC, Rocha M, Alves MM, Salvador AF. UPIMAPI, reCOGnizer and KEGGCharter: Bioinformatics tools for functional annotation and visualization of (meta)-omics datasets. Computational and Structural Biotechnology Journal. 2022;20:1798–1810.

69. The PyMOL Molecular Graphics System; 2022. http://web.archive.org/web/20220624180204/https://pymol.org/.

70. Degnan SM. Think laterally: horizontal gene transfer from symbiotic microbes may extend the phenotype of marine sessile hosts. Frontiers in microbiology. 2014;5:638.

71. Plese B, Kenny NJ, Rossi ME, Cárdenas P, Schuster A, Taboada S, et al. Mitochondrial evolution in the Demospongiae (Porifera): Phylogeny, divergence time, and genome biology. Molecular Phylogenetics and Evolution. 2021;155:107011.

72. Guruprasad K, Reddy BB, Pandit MW. Correlation between stability of a protein and its dipeptide composition: a novel approach for predicting in vivo stability of a protein from its primary sequence. Protein Engineering, Design and Selection. 1990;4(2):155–161.

73. Vihinen M, Torkkila E, Riikonen P. Accuracy of protein flexibility predictions. Proteins: Structure, Function, and Bioinformatics. 1994;19(2):141–149.

74. Lobry J, Gautier C. Hydrophobicity, expressivity and aromaticity are the major trends of amino-acid usage in 999 Escherichia coli chromosome-encoded genes. Nucleic acids research. 1994;22(15):3174–3180.

75. Kyte J, Doolittle RF. A simple method for displaying the hydropathic character of a protein. Journal of molecular biology. 1982;157(1):105–132.