Co-opted Genes of Algal Origin Protect *C. elegans* against Cyanogenic Toxins

Bingying Wang[1], Taruna Pandey[1], Yong Long[2], Sofia E. Delgado-Rodriguez[3], Matthew

D. Daugherty[3, *], Dengke K. Ma[1, *]

[1]Cardiovascular Research Institute and Department of Physiology, University of

California San Francisco, San Francisco, USA

[2]State Key Laboratory of Freshwater Ecology and Biotechnology, Institute of

Hydrobiology, Chinese Academy of Sciences, Wuhan, China.

[3]Department of Molecular Biology, University of California San Diego, San Diego,

United States.

* Correspondence should be addressed to Dengke K. Ma, Ph.D.

(dengke.ma@ucsf.edu) and Matthew D. Daugherty, Ph.D. (MDDaugherty@UCSD.edu)

Abstract

Amygdalin is a cyanogenic glycoside widely used by many plants in herbivore defense.
Poisonous to most animals, amygdalin-derived cyanide is detoxified by potent enzymes
commonly found in bacteria and plants but not most animals. Here we show that the
nematode *C. elegans* can detoxify amygdalin by a genetic pathway comprising *cysl-1*,
egl-9, *hif-1* and *cysl-2*. Essential for amygdalin resistance, *cysl-1* encodes a protein
similar to cysteine synthetic enzymes in bacteria and plants, but functionally co-opted in
C. elegans. We identify exclusively HIF-activating *egl-9* mutations in a *cysl-1* suppressor
screen and show that *cysl-1* confers amygdalin resistance by regulating HIF-1-
dependent *cysl-2* transcription to protect against amygdalin toxicity. Phylogenetic
analysis suggests *cysl-1* and *cysl-2* were acquired from green algae by horizontal gene
transfer (HGT) and functionally co-opted by *C. elegans* in protection against amygdalin.
Unlike functional HGTs from bacteria to eukaryotes that have been widely observed and
characterized, our studies reveal previously unknown HGT from algae to nematodes,
supporting the emerging theme that HGT-mediated evolutionary changes can facilitate
host survival and adaptation to adverse environment stresses and biogenic toxins.

Introduction

In nature, many species of plants have evolved the ability to biosynthesize a group of
chemically related molecules called cyanogenic glycosides as chemical toxins used in
herbivore defense[1–3]. Amygdalin is a natural cyanogenic glycoside enriched in tissues of
many edible plants, including seeds of stone fruits such as cherry (*Prunus avium*),

peach (*Prunus persica*), apricot (*Prunus armeniaca*), and apples (*Malus domestica*).

These plant species synthesize amygdalin in defense against herbivore animals as

amygdalin generates poisonous cyanide upon plant tissue destruction[1–4]. After

consumption by herbivores, amygdalin is hydrolyzed by β-glucosidases and α-

hydroxynitrilases, releasing hydrogen cyanide that can be highly toxic and even lethal to

herbivore animals. To tolerate amygdalin, plants also evolved a family of cysteine

synthase-like enzyme, β-cyanoalanine synthases, to directly detoxify cyanides[4–6].

Caenorhabditis elegans (C. elegans) is a nematode that lives with rotten fruits and

bacteria in their natural ecological niches[7–9], thus may frequently encounter amygdalin.

Although this commonly used genetic model organism rarely encounters amygdalin

under laboratory conditions, loss-of-function mutations in the gene *egl-9* have been

shown to confer resistance to high doses of hydrogen cyanides generated by a

pathogenic bacterial strain of *Pseudomonas aeruginosa*[10]. EGL-9 is a prolyl hydroxylase

and negative regulator of HIF-1 (hypoxia inducible factor), a transcription factor that

directly activates expression of many genes involved in hypoxic adaptation, metabolic

reprogramming and stress responses[11–13]. As a transcriptional target of HIF-1, CYSL-2

is a protein strongly up-regulated in *egl-9* loss-of-function mutants and can detoxify

cyanides *in vitro* as a cyanoalanine synthase[14]. CYSL-1 is a paralogue of CYSL-2, both

of which share sequence similarities with a family of β-cyanoalanine synthases found in

bacteria and plants. We previously found that CYSL-1 regulates *cysl-2* transcription by

binding to EGL-9 in regulation of *cysl-2* expression[15]. Whether this pathway is involved

in response to cyanogenic amygdalin has not been characterized. While EGL-9 and

HIF-1 are evolutionarily conserved in metazoans, most animals do not have proteins

homologous to CYSL-1 and CYSL-2, except certain herbivorous arthropods[16]. In

addition, co-opted detoxification genes of plant origin in whiteflies can neutralize plant-

derived phenolic glycoside toxins[17,18]. However, the evolutionary origin, history of

domestication and co-option of *cysl* genes in *C. elegans* remain hitherto unknown.

By screening a natural product library, we identified amygdalin as a potent activator of

C. elegans cysl-2. We showed that the CYSL-1/EGL-9/HIF-1/CYSL-2 pathway plays a

crucial role in conferring resistance to amygdalin. The intriguing acquisition of amygdalin

resistance in *C. elegans* through two genes (*cysl-1* and *cysl-2*) not commonly found in

other animals into the conserved EGL-9/HIF-1 genetic pathway led us to further explore

the evolutionary origin and path of *cysl-1* and *cysl-2* based on phylogenetic analysis.

Our combined mechanistic and phylogenetic studies indicate that CYSL-1 and CYSL-2

may have originated from green algae by a horizontal gene transfer event in ancient

nematodes and evolved by gene co-option into separate roles in a cellular circuit to

respond to and detoxify cyanogenic cyanides, including amygdalin from plants today.

Results

We previously discovered several components of the HIF-1 pathway acting upstream of

the oxygen-sensing HIF hydroxylase EGL-9 in *C. elegans*[15]. To identify small-molecule

natural products that may modulate HIF-1 activity independently of oxygen-sensing in

C. elegans, we conducted a natural product library screen for compounds that can

activate a HIF-1-dependent transgenic reporter strain *cysl-2*p::GFP in liquid cultures

under normoxia (Fig. 1a). Among the identified compounds, a glycoside called

amygdalin emerged as a potent activator of *cysl-2*p::GFP. We validated the finding

using amygdalin from new preparations of different sources (vendor VWR or Sigma-

Aldrich) to treat *C. elegans* on solid NGM media and confirmed that amygdalin can

robustly activate *cysl-2*p::GFP under normoxic conditions, without affecting the

transgenic co-injection marker *myo-2*p::mCherry (Fig. 1b). Activation of *cysl-2*p::GFP by

amygdalin is dose-dependent and reaches peak level at approximately 48 hrs post

amygdalin treatment (Fig. 1c, d).

We performed RNAseq to identify genome-wide transcriptomic changes induced by

amygdalin. We treated wild-type mixed-stage *C. elegans* hermaphrodites cultivated at

20 °C with 2 mg/ml amygdalin for 48 hrs. After differential expression analyses of

triplicate samples, we identified many genes that are significantly (adjusted p value <

0.05) up- or down-regulated by amygdalin treatment (Fig. 1e). As expected, *cysl-2* is

among the most highly up-regulated genes ($\log_2$fold change = 2.62, adjusted p value =

3.11E-06), while expression of its upstream regulators including *hif-1*, *vhl-1* or *egl-9*

remains largely unchanged (Supplementary Fig. 1 and table S1). Gene Ontology

analysis (Wormbase) indicates that amygdalin also up-regulated several genes

encoding glutathione S-transferases involved in the xenobiotic detoxification pathway

(Supplementary Fig. 1 and table S1).

We next examined how amygdalin activates *cysl-2*p::GFP. HIF-1 and CYSL-1 are two

previously established positive regulators of *cysl-2p*::GFP[15,19–21]. We found that loss-of-

function mutations of *hif-1* or *cysl-1* abolished *cysl-2*p::GFP induction by low-dose

amygdalin (Fig. 2a). Amygdalin is a glycoside that can be metabolized sequentially to

prunasin, glucose, benzaldehyde and hydrogen cyanide (Fig. 2b)[1,4,22]. We treated *cysl-*

*2*p::GFP animals with each of these amygdalin-derived molecules and found prunasin

and cyanide (as in potassium cyanide) activated *cysl-2*p::GFP as amygdalin did, but

others had no effect (Fig. 2c). CYSL-1 and CYSL-2 are cysteine synthase-like proteins

and have been implicated in hydrogen sulfide and cyanide detoxification[14,15,20,23,24].

While CYSL-2 can enzymatically convert cyanide to nontoxic β-cyanoalanine[14], the

essential role of CYSL-1 in *cysl-2*p::GFP activation by amygdalin suggests CYSL-1 acts

as a transcriptional regulator rather than a cysteine synthase or an enzyme to detoxify

cyanide[15]. Taking advantage of the *cysl-2*p::GFP phenotype and lethal effects of

amygdalin in *cysl-1* mutants (see below), we performed a *cysl-1* suppressor screen and

identified exclusively three loss-of-function mutations in *egl-9* that led to constitutive

*cysl-2*p::GFP expression without amygdalin treatment (Fig. 2d). In further genetic

epistasis analysis, loss of *hif-1* suppressed not only *cysl-2*p::GFP induction by

amygdalin but also constitutive *cysl-2*p::GFP expression in *egl-9*; *cysl-1* mutants (Fig.

2E). These data suggest that amygdalin activates *cysl-2*p::GFP through a dis-inhibitory

regulatory pathway from cyanide, CYSL-1, EGL-9 to HIF-1 (Fig. 2f).

To assess the functional importance of the CYSL-1/EGL-9/HIF-1/CYSL-2 pathway, we

examined the time-dependent survival response to high-dose amygdalin (10 mg/ml on

NGM; effective concentrations *in vivo* are likely much lower due to cuticle and intestinal

absorption barriers) in animals with individual or combinatorial pathway gene mutations.

Wild-type animals exhibited no apparent decrease in survival upon amygdalin over

three days of treatment (Fig. 3a). By contrast, loss-of-function *cysl-1* mutants displayed

striking sensitivity to amygdalin, with nearly complete population death at day 3 of

amygdalin treatment (Fig. 3b). Loss-of-function of *cysl-2* also caused similar time-

dependent sensitivity to amygdalin (Fig. 3c). While loss of *egl-9* fully suppressed

vulnerability of *cysl-1* mutants to amygdalin (Fig. 3d), loss of either *hif-1* or *cysl-2*

rendered *egl-9* mutants sensitive to amygdalin (Fig. 3e and f). These data support

central roles of the CYSL-1/EGL-9/HIF-1/CYSL-2 pathway in the transcriptional

response and conferring resistance to amygdalin in *C. elegans*.

To investigate the evolutionary origin CYSL-1/2 in *C. elegans*, we searched for

homologs of CYSL-1 and CYSL-2 proteins across all domains of life. We identified

related proteins with >50% sequence identity in nematodes, green algae, land plants,

and bacteria. Outside of nematode proteins, the most closely related sequences, and

the only ones with sequence identities >60% were from the *Chlorophyta* clade of green

algae (Supplementary Fig. 2a and Fig. 4a).

To determine the phylogenetic origin of CYSL proteins in nematodes, we performed

maximum likelihood (ML) based phylogenetic analyses on CYSL homologs from a

broad range of eukaryotic and prokaryotic species (Supplementary Data Files 1, 2 and

Fig. 4A). Consistent with the high sequence similarity to proteins in green algae, we

found strong support for nematode proteins nesting within the chlorophyte clade of

green algae (Fig. 4b). Proteins from streptophytes (*Streptophyta*), which include some

green algae and all land plants, are found in a distinct region of the phylogenetic tree,

as are proteins from single-cell eukaryotes (*e.g.* groups including *SAR*, *Excavata*, and

Rhodophyta) and bacteria. These inferences were robust to the ML software we used

for analysis, as well as the substitution models we employed (Fig. 4b). Importantly,

cysteine synthase enzymes found in some arthropods, such as mites, which are the

result of a horizontal gene transfer (HGT) event from bacteria show less than 40%

sequence identity, consistent with their separate evolutionary origin[16] (Supplementary

Fig. 2b). Taken together, these data support a single HGT event from a green algal

species in the *Chlorophyta* that gave rise to nematode CYSL.

We further wished to analyze the distribution of CYSL proteins in nematodes to

determine at what point CYSL was acquired by nematodes and at what point gene

duplication occurred (Fig. 4c). We found CYSL proteins only in *Chromadoria* clade of

nematodes and not in the well sampled organisms in the *Trichinellida* (*e.g. Trichinella*

species). Although we cannot rule out that it was lost in the *Trichinellida*, the most

parsimonious explanation for this distribution is that CYSL was acquired by an ancestral

Chromodoria nematode after the divergence from other clades of nematodes. Within

Chromodoria, we only detect a single CYSL homolog in the most basally branching

Spirurina nematodes[25] (*e.g. Toxocara canis*) whereas we see two or more copies of

CYSL homologs in *Tylenchina* (*e.g. Bursaphelenchus* species) and *Rhabditina* (*e.g.*

Caenorhabditis species) lineages of nematodes. These data, coupled with the

branching of CYSL-1 and CYSL-2 paralogs at the base of the nematode CYSL protein

phylogeny (Fig. 4c), suggest that CYSL duplicated soon after acquisition in nematodes

to CYSL-1 like proteins and CYSL-2 like proteins. Based on our findings of the non-

overlapping functional importance of CYSL-1 and CYSL-2 in *C. elegans* for cyanide

detoxification, these phylogenetic findings suggest that this network of CYSL-1 and

CYSL-2 evolved soon after the acquisition of these enzymes by nematodes.

Discussion

In this study, we establish novel physiological roles of the CYSL-1/EGL-9/HIF-1/CYSL-2

pathway in detoxification of amygdalin in *C. elegans*. Our data indicate that CYSL-1 acts

upstream of the evolutionarily conserved HIF pathway as a key regulator of *cysl-2*

induction upon amygdalin exposure, resulting in detoxification of cyanide that derives

from the natural diet of *C. elegans* and other nematodes. How amygdalin or cyanide is

sensed by cells in *C. elegans* and regulates CYSL-1 and EGL-9 awaits further studies.

Plausible mechanisms include mitochondrial inhibition, posttranslational modification by

S-cyanylation, and allosteric modulation of CYSL-1 and EGL-9 interaction[15,26,27].

By phylogenetic analysis, we obtained evidence pointing to an intriguing evolutionary

origin of *cysl-1/cysl-2* in *C. elegans*. Maximum likelihood estimation based on protein

sequence similarity supports a single HGT event from a green algal species in the

Chlorophyta that gave rise to nematode *cysl* genes. Our analysis further suggests that a

cysl gene orthologue was acquired by ancestral *Chromodoria* nematodes after the

divergence from other clades of nematodes, and duplicated soon after their acquisition

to give rise to *cysl* gene paralogs. In *C. elegans*, *cysl-1* and *cysl-2* have further evolved

separate functions from ancestral cysteine synthase-like activity into division of labor.

While CYSL-2 retains cyanoalanine synthase enzymatic activity to detoxify cyanide,

CYSL-1 is co-opted to regulate *cysl-2* expression upon amygdalin/cyanide exposure in

a cellular circuit that detects and defends against amygdalin/cyanide. *Chromodoria*

nematodes diverged from other clades of nematodes more than 250 million year ago,

around which drastic geological and ecological changes (e.g. those leading to Permian-

Triassic mass extinction) may have exerted selection pressures on nematodes surviving

in environments rife with sulfide/cyanide toxicity and ancient algal species[28–32].

Following acquisition of *cysl* genes, nematodes may have retained and co-opted *cysl*

gene functions in adaptation to new environments, including living as plant parasites or

in rotten fruits associated with cyanide presence. While precise mechanisms of these

evolutionary changes particularly HGT remain unclear, the evolutionary pathway leading

to distinct cellular roles of *cysl-1* and *cysl-2* in *C. elegans* today exemplifies how

organisms can acquire novel biological traits to survive and reproduce by HGT and

gene co-option[32–37]. Unlike functional HGTs from bacteria to eukaryotes that have been

widely observed and characterized[38–40], our mechanistic and phylogenetic analyses

reveal previously unknown HGT from algae to nematodes, supporting the emerging

principle that HGTs can lead to evolutionary changes and novel traits to facilitate host

protection and adaptation to abiotic stress, chemical toxins and extreme environments.

Figures and Figure legends

Fig. 1. Screen of natural product library identifies amygdalin as a potent activator of *cysl-2*p::GFP. a, Schematic of compound screens that led to identification of amygdalin. **b**, Representative bright field and epifluorescence images showing dose-dependent up-regulation of *cysl-2*p::GFP by amygdalin. Scale bar: 50 μm. **c**, Quantification of percentages of animals with strong *cysl-2*p::GFP expression after 48 hrs of amygdalin treatment with increasing doses. Values are means ± S.D with ***P < 0.001 (N = 5 independent experiments, n > 50 animals per experiment). **d**, Quantification of percentages of animals with strong *cysl-2*p::GFP expression after

216   increasing durations of amygdalin treatment at the fixed dose of 2 mg/ml. Values are

217   means ± S.D with ***$P < 0.001$ (N = 5 independent experiments, n > 50 animals per

218   experiment). **e**, Volcano plot showing significantly (adjusted p value < 0.05) up- (green)

219   or down- (blue) regulated genes in amygdalin-treated animals, with *cysl-2* noted (arrow).

Fig. 2. Amygdalin activates *cysl-2*p::GFP through cyanide, CYSL-1, EGL-9 and

HIF-1. a, Representative bright field and epifluorescence images showing up-regulation

of *cysl-2*p::GFP by amygdalin (1 mg/ml) in wild-type but not *hif-1*(*ia4*), *cysl-1*(*ok762*)

loss-of-function mutant animals. Scale bar: 50 µm. **b**, Schematic illustrating the

metabolic pathway of amygdalin, leading to generation of prunasin, glucose, hydrogen

cyanide and benzaldehyde. **c**, Quantification of percentages of animals with strong *cysl-*

*2*p::GFP expression after 48 hrs of treatment with amygdalin (2 mg/ml), prunacin (2

mg/ml), glucose (2 mg/ml), potassium cyanide (0.2 mg/ml) and benzaldehyde (1 mg/ml).

Values are means ± S.D with ***P < 0.001 (N = 5 independent experiments, n > 50

animals per experiment). **d**, Schematic showing *cysl-1* suppressor screens that led to

232 identification of three mutants, all of which are allelic to *egl-9* based on whole-genome

233 sequencing and complementation tests. **e**, Table summary of *cysl-2*p::GFP activation

234 phenotypes in animals with indicated genotypes and treatment conditions. Penetrance

235 and numbers of animals examined are noted. **f**, Model illustrating the regulatory

236 pathway that mediates transcriptional response to amygdalin.

Fig. 3. Amygdalin resistance requires CYSL-1, HIF-1 and CYSL-2 acting in a regulatory pathway. a, Quantification of percentages of post-L4 stage wild-type animals survived after 24, 48 or 72 hrs of treatment with amygdalin (10 mg/ml). **b**, Quantification of percentages of post-L4 stage *cysl-1(ok762)* loss-of-function mutant animals survived after 24, 48 or 72 hrs of treatment with amygdalin (10 mg/ml). **c**, Quantification of percentages of post-L4 stage *cysl-2(ok3516)* loss-of-function mutant animals survived after 24, 48 or 72 hrs of treatment with amygdalin (10 mg/ml). **d**, Quantification of percentages of post-L4 stage *cysl-1(ok762); egl-9(sa307)* double loss-of-function mutant animals survived after 24, 48 or 72 hrs of treatment with amygdalin (10 mg/ml). **e**, Quantification of percentages of post-L4 stage *hif-1(ia04); egl-9(sa307)*

249   double loss-of-function mutant animals survived after 24, 48 or 72 hrs of treatment with

250   amygdalin (10 mg/ml). **f**, Quantification of percentages of post-L4 stage *cysl-2*(*ok3516*);

251   *egl-9(sa307)* double loss-of-function mutant animals survived after 24, 48 or 72 hrs of

252   treatment with amygdalin (10 mg/ml). Values are means ± S.D with ***$P < 0.001$ ($N = 5$

253   independent experiments, $n > 50$ animals per experiment).

Figure 4

a

Bacteria

Unicellular eukaryotes

Green algae

Nematodes

CYSL-1

100

CYSL-2

Land plants

0.2 substitutions/residue

b

Software Substitution model	Phylogenetic clade	Branch support Bootstrap	SH-aLRT
IQ-TREE *LG + I + G4(*)*	Nematode only	100	100
	Nematode + green algae	93	88
IQ-TREE *LG*	Nematode only	100	100
	Nematode + green algae	87	89
IQ-TREE *WAG + I + G4*	Nematode only	100	100
	Nematode + green algae	97	89
IQ-TREE *WAG*	Nematode only	100	100
	Nematode + green algae	73	76
IQ-TREE *JTT + I + G*	Nematode only	100	100
	Nematode + green algae	76	71
IQ-TREE *JTT*	Nematode only	100	100
	Nematode + green algae	47	73
RAxML *LG + Gamma(*)*	Nematode only	99	
	Nematode + green algae	32	

c

Fig. 4. Nematode CYSL genes were acquired from green algae by horizontal gene transfer. a, IQ-TREE generated maximum likelihood (ML) phylogenetic tree of homologs of nematode CYSL across a broad range of species. Species groups are shaded. Clades of similar proteins are collapsed and shown as triangles. Bootstrap support values for relevant branches are shown. Bolded values indicate the support for a clade that only contains CYSL-like proteins from nematodes (100% bootstrap support) and a well-supported clade that contains only CYSL-like proteins from nematodes and a group of green algae in the *Chlorophyta* lineage (93% bootstrap support). Within the nematodes, two clear, well-supported clades of CYSL proteins exist, one that contains *C. elegans* CYSL-1 and one that contains *C. elegans* CYSL-2. **b**, ML phylogenetic analyses were performed with multiple programs and multiple amino acid substitution models as indicated. Support values (bootstrap and SH-aLRT when available) are shown for a clade containing only nematode proteins and a clade containing nematode and chlorophyte green algae proteins as are bolded in part A. Asterisks next to model names indicate that best fitting model as described in Methods. **c**, Expanded phylogenetic tree of CYSL proteins from nematodes and their nearest homologs in green algae. Font colors denote major groups of nematode species as found in [25]. Bootstrap support values are shown at relevant nodes. For panels A-C, a complete list of all sequences used is found in Supplementary Data file 1 and all trees, including support values, are found in Supplementary Data file 2.

Supplementary Fig. 1. Amygdalin-independent and regulated genes identified by

RNAseq. a, Normalized fold induction by amygdalin (2 mg/ml on NGM for 48 hrs) of

genes acting upstream of *cysl-2* (*egl-9*, *hif-1*, *vhl-1*) and GST-encoding genes involved

in xenobiotic detoxification. **b**, Heat map showing hierarchical clustering of genes

differentially up-regulated by amygdalin (2 mg/ml on NGM for 48 hrs) from RNAseq

experiments.

a

C. elegans CYSL-1 alignment to top hit green algae (*Raphidocelis subcapitata*)

```
Blast score: 392 bit    Identities: 63%   Similarity: 79%

C. elegans CYSL-1 [NP_001369978.1]           4    RNSIGENAAALIGNTPMVYINKLTKGLPGTVAVKIEYMNPAGSVKDRIGAAMLAAAEKDG   63
                                                   + +I ++    LIGNTPMVY+ ++ +GLPG +A K+E M P  SVKDRIG +M++AAE DG
R. subcapitata (greenalgae) [GBF88368.1]     36   KMNIAKDVTELIGNTPMVYLKRVNEGLPGKIACKLELMEPCCSVKDRIGYSMISAAEADG   95

C. elegans CYSL-1 [NP_001369978.1]           64   TVIPGVTTLIEPTSGNTGIALAFVAAAKGYRCIVTMPASMSGERRTLLKAYGSEVVLTDP   123
                                                   +  G TTL+EPTSGNTGI LAF+AAAKGY+ ++TMPASMS ERR LL+A+G+E+VLTDP
R. subcapitata (greenalgae) [GBF88368.1]     96   KITAGKTTLVEPTSGNTGIGLAFIAAAKGYKLVLTMPASMSLERRILLRAFGAELVLTDP   155

C. elegans CYSL-1 [NP_001369978.1]           124  AKGMKGAIDMANQLKENIPGSIILAQFDNPNNPLVHYQTTGPEIWRQTKGTVDAVVFGVG   183
                                                   AKGMKGA+  A ++  N P S IL QF+NPNNP VHY+TTGPEIW+ T G +D +V GVG
R. subcapitata (greenalgae) [GBF88368.1]     156  AKGMKGAVLKAQEIAANTPDSYILQQFENPNNPKVHYETTGPEIWKATDGKIDFLVSGVG   215

C. elegans CYSL-1 [NP_001369978.1]           184  TGGTITGVGRYLQEQNPGVRVFAVEPEESAILSGRPAGPHKIQGIGAGFAPAVLDTKIYE   243
                                                   TGGTITG GRYL+EQNP V++ AVEP ES +LSG   GPHKIQGIGAGF P VLDT + +
R. subcapitata (greenalgae) [GBF88368.1]     216  TGGTITGAGRYLKEQNPNVKLVAVEPSESPVLSGGKPGPHKIQGIGAGFVPGVLDTHLID   275

C. elegans CYSL-1 [NP_001369978.1]           244  DVIRIHSDEAIVMAQRLSYEEGLLGGISSGANVAAALQLAARPEMAGKLIVTCLPSCGER   303
                                                   +VI++ SD+A+ MA+RL+ EEGLL GISSGA   AA+Q+AARPE AGKLI   LPS GER
R. subcapitata (greenalgae) [GBF88368.1]     276  EVIQVSSDDAVSMARRLATEEGLLTGISSGAATVAAIQVAARPENAGKLIAVILPSFGER   335

C. elegans CYSL-1 [NP_001369978.1]           304  YMTSPLYTDIRESAMALA   321
                                                   Y++S L+ +IR+ A A++
R. subcapitata (greenalgae) [GBF88368.1]     336  YLSSVLFQNIRDEAEAMS   353
```

b

C. elegans CYSL-1 alignment to *Tetranychus urticae* (mite) cysteine synthase

```
Blast score: 211 bit    Identities: 39%   Similarity: 56%

C. elegans CYSL-1 [NP_001369978.1]           3    DR-NSIGENAAALIGNTPMVYINKLTKGLPGTVAVKIEYMNPAGSVKDRIGAAMLAAAEK   61
                                                   DR N I +A  LIGNTP++ +++L  G PG + K E++NP  S+KDR    M+A A++
T. urticae (mite) [XP_015786551.1]           7    DRINGITPSALDLIGNTPLIALDRLWPG-PGRLLAKCEFLNPTASLKDRSSYYMIAKAKE   65

C. elegans CYSL-1 [NP_001369978.1]           62   DGTVIPGVTTLIEPTSGNTGIALAFVAAAKGYRCIVTMPASMSGERRTLLKAYGSEVVLT   121
                                                    G +  G  ++IE TSGN G  +A V A  G+   VTM    S +R  ++ A G+ V+L
T. urticae (mite) [XP_015786551.1]           66   SGQLKDG-ESVIEVTSGNQGGGIACVTAVMGHPFTVTMSKGNSPQRAIMMNALGANVILV   124

C. elegans CYSL-1 [NP_001369978.1]           122  DPAKGMKGAI---------DMANQLKENIPGSIILAQFDNPNNPLVHYQTTGPEIWRQTK   172
                                                   D   G  G +         + A +++E   +  + QF+NP N L HY+TTGPEIWRQT
T. urticae (mite) [XP_015786551.1]           125  DQVTGKPGNVTADDVAAAEETAMKIREET-NAYYVDQFNNPTNCLAHYETTGPEIWRQTN   183

C. elegans CYSL-1 [NP_001369978.1]           173  GTVDAVVFGVGTGGTITGVGRYLQEQNPGVRVFAVEPEESAILSG----RPAGPHKIQGI   228
                                                   G +DA + G GTGG   G  ++L+E+NP VR F VEPE     ++G      +P  H +QG
T. urticae (mite) [XP_015786551.1]           184  GRIDAFLVGCGTGGCFVGTSKFLKEKNPNVRCFVVEPEGCQPIAGCTITKPL--HLLQGS   241

C. elegans CYSL-1 [NP_001369978.1]           229  GAGFAPAVLDTKIYEDVIRIHSDEAIVMAQRLSYEEGLLGGISSGANVAAALQLAARPEM   288
                                                   G G  P + D K+Y D I +  +EAI   + L  +EGL  G ++G N+AAA++L    ++
T. urticae (mite) [XP_015786551.1]           242  GYGCVPTLFDKKVYNDSISVSDEEAIEYRKLLGQKEGLFCGFTTGGNIAAAIKLLKSGQL   301

C. elegans CYSL-1 [NP_001369978.1]           289  -AGKLIVTCLPSCGERY   304
                                                    +VT L   G +Y
T. urticae (mite) [XP_015786551.1]           302  PKDAWVVTILCDSGLKY   318
```

284 **Supplementary Fig. 2. Protein sequence alignments of CYSL-1 with orthologs**

285 **from green algae (a) and mite (b).** Blast scores, a.a. identities and similarities are

286 shown.

 Methods and Materials

 C. elegans

C. elegans strains were maintained under standard laboratory conditions unless

otherwise specified. The N2 Bristol strain was used as the reference wild type, and the

polymorphic Hawaiian strain CB4856 was used for genetic linkage mapping and SNP

analysis [41,42]. Forward genetic screens for *cysl-2*p::GFP; *cysl-1* suppressing mutants

after ethyl methanesulfonate (EMS)-induced random mutagenesis were performed as

described previously [15,43]. Identification of mutations by whole-genome sequencing and

complementation tests by crossing EMS mutants with *egl-9(sa307)* heterozygous males

were used to determine *dma430, dma432* and *dma437* as new loss-of-function alleles

of *egl-9*. Transgenic strains were generated by germline transformation as described [44].

Transgenic constructs were co-injected (at 10 - 50 ng/μl) with dominant *unc-*

54p::mCherry, and stable extrachromosomal lines of fluorescent animals were

established. Genotypes of strains used are as follows: Chr. II: *cysl-2*(*ok3516*)*;* Chr. IV:

nls470 [cysl-2p::GFP; unc-54p::mCherry], Chr. V: *egl-9(dma430, 432, 437, sa307), hif-*

1(ia4), Chr. X: *cysl-1(ok762).*

Sample preparation for RNA sequencing and data analysis

N2 control (M9 treated) and amygdalin-treated animals (2 mg/ml on NGM for 48 hrs)

were maintained at 20 °C and washed down from NGM plates using M9 solution and

subjected to RNA extraction using the RNeasy Mini Kit from Qiagen. RNA-seq library

preparation and data analysis were performed as previously described [45]. Three

biological replicates were included for each treatment. The cleaned RNAseq reads were

mapped to the genome sequence of *C. elegans* using hisat2 [46]. Abundance of genes

was expressed as FPKM (Reads per kilobase per million mapped reads). Identification

of differentially expressed genes was performed using the DESeq2 package [47].

C. elegans amygdalin sensitivity and survival assays

Animals were cultured under non-starved conditions for at least 4 generations before

amygdalin sensitivity assays. NGM plates spread with equal agar thickness seeded with

equal amounts of OP50 were used for M9 or amygdalin (dissolved in M9, TCA0443-

010G, VWR or equivalent A6005, Sigma-Aldrich) supplementation. Fresh preparations

of fully dissolved amygdalin in M9 were added (250 µl/ 60 cm plate) and spread evenly

on NGM plates with pre-seeded OP50. Once the plates with amygdalin were briefly air

dried, synchronized-stage (L4) animals were placed and monitored for survival over 24,

48 and 72 hrs post treatment. Animals were scored as dead if they showed no pumping

and movement upon light touch with the body necrosis subsequently confirmed.

Phylogenetic analysis

C. elegans CYSL-1 (accession NP_001369978.1) was used as a BLASTp [48] search

query against the nonredundant (NR) and Reference Sequence (RefSeq) protein

databases. Resulting sequences were aligned using Clustal Omega [49] using two

iterations of refinement and duplicate sequences were removed. Non-aligning termini

and alignment positions in which >95% of all sequences had a gap were manually

removed from subsequent analyses. To generate phylogenetic trees encompassing a

broad range of CYSL homologs, sequences with >80% sequence identity were reduced

to a single unique sequence using CD-HIT [50] with a 0.8 sequence identity cutoff. The

resulting 253 protein sequences (accession numbers and species names found in

Supplemental file 1) were realigned as above and used as input for maximum likelihood

phylogenetic programs. IQ-TREE [51] phylogenies were generated using the "-bb 1000 -

alrt 1000" commands for generation of 1000 ultrafast bootstrap [52] and SH-aLRT support

values. The best substitution model was determined by ModelFinder [53] using the "-m

AUTO" command or the substitution model was specified as shown in Fig. 4B using the

"-m" command. RAxML phylogenies were generated using "-f a" with 500 bootstrap

replicates and "-m PROTGAMMAAUTO" to determine the best substitution model with

gamma distribution. All phylogenetic trees generated from IQ-TREE and RAxML, with

support values, can be found in Supplemental file 2.

To generate a phylogeny of nematode CYSLs, the relevant subtree containing

nematodes and the nearest green algae was extracted for the larger phylogeny

generated above. Additional nematode sequences that had been removed using the

CD-HIT filtering described above were added back in. Sequences were realigned and

degapped as described above with a CDHIT similarity cutoff of 98% followed by

additional manual removal of poorly aligning sequences. An IQ-TREE phylogenetic tree

using ModelFinder (LG+I+G) was determined as described above. Accession numbers

and species names are found in Supplemental file 1, and complete IQ-TREE with

support values is found in Supplemental file 2.

Confocal and epifluorescence microscopic imaging

SPE epifluorescence compound microscopes (Leica) were used to capture

fluorescence images. Animals of different genotypes were randomly picked at the same

young adult stage (24 hrs post L4) and treated with 1 mM Levamisole sodium Azide in

M9 solution (31,742-250MG, Sigma-Aldrich), aligned on an 4% agar pad on slides for

imaging. Identical setting and conditions were used to compare genotypes, treatment

experimental groups with control.

Statistical analysis (wet lab data)

Data were analyzed using GraphPad Prism 9.2.0 Software (Graphpad, San Diego, CA)

and presented as means ± S.D. unless otherwise specified, with P values calculated by

unpaired two-tailed t-tests (comparisons between two groups), one-way ANOVA

(comparisons across more than two groups) and adjusted with Bonferroni's corrections.

Data availability

The RNAseq read datasets were deposited in NCBI SRA (Sequence Read Archive)

under the BioProject accession PRJNA843348. All other data generated for this study

are included in this article.

Acknowledgments

Some strains were provided by the CGC, which is funded by NIH Office of Research

Infrastructure Programs (P40 OD010440). The work was supported by NIH grant

1R35GM139618 (D.K.M.), Packard Fellowships for Science and Engineering (D.K.M.)

and the Pew Scholars Program in the Biomedical Sciences (D.K.M. and M.D.D.).

Author contributions

B.W., T.P. and D.K.M. conceptualized, performed and analyzed the wet experiments

and wrote the manuscript. Y.L. performed RNAseq bioinformatic analysis. S.D.R. and

M.D.D. performed phylogenetic analysis. M.D.D and D.K.M. supervised the project.

Competing interests

The authors declare no competing interests.

Correspondence

Correspondence should be addressed to Dengke K. Ma, Ph.D. (dengke.ma@ucsf.edu)

and Matthew D. Daugherty, Ph.D. (MDDaugherty@UCSD.edu)

References

1. Gleadow, R. M. & Møller, B. L. Cyanogenic glycosides: synthesis, physiology, and phenotypic plasticity. *Annu. Rev. Plant Biol.* **65**, 155–185 (2014).

2. Mithöfer, A. & Boland, W. Plant defense against herbivores: chemical aspects. *Annu. Rev. Plant Biol.* **63**, 431–450 (2012).

3. Møller, B. L. Functional diversifications of cyanogenic glucosides. *Curr. Opin. Plant Biol.* **13**, 338–347 (2010).

4. Cressey, P. & Reeve, J. Metabolism of cyanogenic glycosides: A review. *Food Chem. Toxicol. Int. J. Publ. Br. Ind. Biol. Res. Assoc.* **125**, 225–232 (2019).

5. Ohlen, M. van, Herfurth, A.-M. & Wittstock, U. *Herbivore Adaptations to Plant Cyanide Defenses*. *Herbivores* (IntechOpen, 2017). doi:10.5772/66277.

6. Machingura, M., Salomon, E., Jez, J. M. & Ebbs, S. D. The β-cyanoalanine synthase pathway: beyond cyanide detoxification. *Plant Cell Environ.* **39**, 2329–2341 (2016).

7. Schulenburg, H. & Félix, M.-A. The Natural Biotic Environment of Caenorhabditis elegans. *Genetics* **206**, 55–86 (2017).

8. Frézal, L. & Félix, M.-A. C. elegans outside the Petri dish. *eLife* **4**, (2015).

9. Kiontke, K. & Sudhaus, W. Ecology of Caenorhabditis species. *WormBook Online Rev. C Elegans Biol.*

1–14 (2006) doi:10.1895/wormbook.1.37.1.

10. Gallagher, L. A. & Manoil, C. Pseudomonas aeruginosa PAO1 kills Caenorhabditis elegans by cyanide

poisoning. *J. Bacteriol.* **183**, 6207–6214 (2001).

11. Semenza, G. L. Hypoxia-inducible factors in physiology and medicine. *Cell* **148**, 399–408 (2012).

12. Kaelin, W. G. & Ratcliffe, P. J. Oxygen sensing by metazoans: the central role of the HIF hydroxylase

pathway. *Mol. Cell* **30**, 393–402 (2008).

13. Powell-Coffman, J. A. Hypoxia signaling and resistance in C. elegans. *Trends Endocrinol. Metab. TEM*

21, 435–440 (2010).

14. Vozdek, R., Hnízda, A., Krijt, J., Será, L. & Kožich, V. Biochemical properties of nematode O-

acetylserine(thiol)lyase paralogs imply their distinct roles in hydrogen sulfide homeostasis. *Biochim.*

Biophys. Acta **1834**, 2691–2701 (2013).

15. Ma, D. K., Vozdek, R., Bhatla, N. & Horvitz, H. R. CYSL-1 interacts with the O2-sensing hydroxylase

EGL-9 to promote H2S-modulated hypoxia-induced behavioral plasticity in C. elegans. *Neuron* **73**,

925–940 (2012).

16. Wybouw, N. *et al.* A gene horizontally transferred from bacteria protects arthropods from host plant

cyanide poisoning. *eLife* **3**, e02365 (2014).

17. Xia, J. *et al.* Whitefly hijacks a plant detoxification gene that neutralizes plant toxins. *Cell* **184**, 3588

(2021).

18. Lapadula, W. J., Mascotti, M. L. & Juri Ayub, M. Whitefly genomes contain ribotoxin coding genes

acquired from plants. *Sci. Rep.* **10**, 15503 (2020).

19. Shen, C. & Powell-Coffman, J. A. Genetic analysis of hypoxia signaling and response in C elegans.

Ann. N. Y. Acad. Sci. **995**, 191–199 (2003).

20. Budde, M. W. & Roth, M. B. The response of Caenorhabditis elegans to hydrogen sulfide and

hydrogen cyanide. *Genetics* **189**, 521–532 (2011).

21. Shao, Z., Zhang, Y., Ye, Q., Saldanha, J. N. & Powell-Coffman, J. A. C. elegans SWAN-1 Binds to EGL-9

and regulates HIF-1-mediated resistance to the bacterial pathogen Pseudomonas aeruginosa PAO1.

PLoS Pathog. **6**, e1001075 (2010).

22. He, X.-Y. *et al.* Amygdalin - A pharmacological and toxicological review. *J. Ethnopharmacol.* **254**,

112717 (2020).

23. Topalidou, I. & Miller, D. L. Caenorhabditis elegans HIF-1 Is Broadly Required for Survival in

Hydrogen Sulfide. *G3 Bethesda Md* **7**, 3699–3704 (2017).

24. Burton, N. O. *et al.* Intergenerational adaptations to stress are evolutionarily conserved, stress-

specific, and have deleterious trade-offs. *eLife* **10**, e73425 (2021).

25. Smythe, A. B., Holovachov, O. & Kocot, K. M. Improved phylogenomic sampling of free-living

nematodes enhances resolution of higher-level nematode phylogeny. *BMC Evol. Biol.* **19**, 121

(2019).

26. García, I., Arenas-Alfonseca, L., Moreno, I., Gotor, C. & Romero, L. C. HCN Regulates Cellular

Processes through Posttranslational Modification of Proteins by S-cyanylation. *Plant Physiol.* **179**,

107–123 (2019).

27. Cooper, C. E. & Brown, G. C. The inhibition of mitochondrial cytochrome oxidase by the gases

carbon monoxide, nitric oxide, hydrogen cyanide and hydrogen sulfide: chemical mechanism and

physiological significance. *J. Bioenerg. Biomembr.* **40**, 533–539 (2008).

28. Grice, K. *et al.* Photic zone euxinia during the Permian-triassic superanoxic event. *Science* **307**, 706–

709 (2005).

29. Gill, B. C. *et al.* Geochemical evidence for widespread euxinia in the later Cambrian ocean. *Nature*

469, 80–83 (2011).

30. Meyer, K. M., Yu, M., Jost, A. B., Kelley, B. M. & Payne, J. L. δ13C evidence that high primary

productivity delayed recovery from end-Permian mass extinction. *Earth Planet. Sci. Lett.* **302**, 378–

384 (2011).

31. Quist, C. W., Smant, G. & Helder, J. Evolution of plant parasitism in the phylum Nematoda. *Annu.*

Rev. Phytopathol. **53**, 289–310 (2015).

32. Coghlan, A. Nematode genome evolution. *WormBook Online Rev. C Elegans Biol.* 1–15 (2005)

doi:10.1895/wormbook.1.15.1.

33. Delaux, P.-M. & Schornack, S. Plant evolution driven by interactions with symbiotic and pathogenic

microbes. *Science* **371**, eaba6605 (2021).

34. Oakley, T. H. Furcation and fusion: The phylogenetics of evolutionary novelty. *Dev. Biol.* **431**, 69–76

(2017).

35. Zhou, H., Beltrán, J. F. & Brito, I. L. Functions predict horizontal gene transfer and the emergence of

antibiotic resistance. *Sci. Adv.* **7**, eabj5056.

36. Hittinger, C. T. & Carroll, S. B. Gene duplication and the adaptive evolution of a classic genetic

switch. *Nature* **449**, 677–681 (2007).

37. True, J. R. & Carroll, S. B. Gene co-option in physiological and morphological evolution. *Annu. Rev.*

Cell Dev. Biol. **18**, 53–80 (2002).

38. Husnik, F. & McCutcheon, J. P. Functional horizontal gene transfer from bacteria to eukaryotes. *Nat.*

Rev. Microbiol. **16**, 67–79 (2018).

39. Keeling, P. J. Functional and ecological impacts of horizontal gene transfer in eukaryotes. *Curr. Opin.*

Genet. Dev. **19**, 613–619 (2009).

40. Crisp, A., Boschetti, C., Perry, M., Tunnacliffe, A. & Micklem, G. Expression of multiple horizontally

acquired genes is a hallmark of both vertebrate and invertebrate genomes. *Genome Biol.* **16**, 50

(2015).

41. Brenner, S. The genetics of Caenorhabditis elegans. *Genetics* **77**, 71–94 (1974).

42. Davis, M. W. *et al.* Rapid single nucleotide polymorphism mapping in C. elegans. *BMC Genomics* **6**,

118 (2005).

43. Ma, D. K. *et al.* Acyl-CoA Dehydrogenase Drives Heat Adaptation by Sequestering Fatty Acids. *Cell*

161, 1152–1163 (2015).

44. Mello, C. C., Kramer, J. M., Stinchcomb, D. & Ambros, V. Efficient gene transfer in C.elegans:

extrachromosomal maintenance and integration of transforming sequences. *EMBO J.* **10**, 3959–

3970 (1991).

45. Jiang, W. *et al.* A genetic program mediates cold-warming response and promotes stress-induced

phenoptosis in C. elegans. *eLife* **7**, (2018).

46. Kim, D., Langmead, B. & Salzberg, S. L. HISAT: a fast spliced aligner with low memory requirements.

Nat. Methods **12**, 357–360 (2015).

47. Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and dispersion for RNA-seq

data with DESeq2. *Genome Biol.* **15**, 550 (2014).

48. Altschul, S. F., Gish, W., Miller, W., Myers, E. W. & Lipman, D. J. Basic local alignment search tool. *J.*

Mol. Biol. **215**, 403–410 (1990).

49. Sievers, F. *et al.* Fast, scalable generation of high-quality protein multiple sequence alignments using

Clustal Omega. *Mol. Syst. Biol.* **7**, 539 (2011).

50. Fu, L., Niu, B., Zhu, Z., Wu, S. & Li, W. CD-HIT: accelerated for clustering the next-generation

sequencing data. *Bioinforma. Oxf. Engl.* **28**, 3150–3152 (2012).

51. Nguyen, L.-T., Schmidt, H. A., von Haeseler, A. & Minh, B. Q. IQ-TREE: a fast and effective stochastic

algorithm for estimating maximum-likelihood phylogenies. *Mol. Biol. Evol.* **32**, 268–274 (2015).

52. Hoang, D. T., Chernomor, O., von Haeseler, A., Minh, B. Q. & Vinh, L. S. UFBoot2: Improving the

Ultrafast Bootstrap Approximation. *Mol. Biol. Evol.* **35**, 518–522 (2018).

53. Kalyaanamoorthy, S., Minh, B. Q., Wong, T. K. F., von Haeseler, A. & Jermiin, L. S. ModelFinder: fast

model selection for accurate phylogenetic estimates. *Nat. Methods* **14**, 587–589 (2017).

495