An Early Miocene skeleton of *Brachydiceratherium* Lavocat, 1951 (Mammalia,

Perissodactyla) from the Baikal area, Russia, and a revised phylogeny of Eurasian

teleoceratines

Alexander Sizov[a, b], Alexey Klementiev[b] and Pierre-Olivier Antoine[c]

a. Geological Institute of Russian Academy of Science, Pyzhevskii pereulok, 7, bld., 1,

Moscow 119017, Russia

b. Institute of the Earth's Crust of the Russian Academy of Sciences, Siberian Branch,

Lermontova St., 128, Irkutsk, 664033, Russia.

c. Institut des Sciences de l'Evolution, UMR 5554 Univ Montpellier, CNRS, IRD, Place

Eugène Bataillon, 34095 Montpellier cedex 5, France

Abstract. Hippo-like rhinocerotids, or teleoceratines, were a conspicuous component of

Holarctic Miocene mammalian faunas, but their phylogenetic relationships remain poorly

known. Excavations in lower Miocene deposits of the Olkhon Island (Tagay locality, Eastern

Siberia; 16–18 Ma) have opened a unique window on the poorly-known early history of the

Lake Baikal ecosystems, notably by unearthing a skeleton of the teleoceratine

Brachydiceratherium shanwangense (Wang, 1965). The remains provide new insights into the

skull and postcranial morphology of this elusive species. The new material is compared with

other Eurasian teleoceratines and the relationships within Teleoceratina are investigated

through a phylogenetic analysis. *Diaceratherium* Dietrich, 1931 (earliest Miocene, Western

Europe) is found to be monotypic and is retrieved as the earliest teleoceratine offshoot. Other

genera have more than one species and are also found to be monophyletic, with

Prosantorhinus Heissig, 1974 (early Miocene, Eurasia) + *Teleoceras* Hatcher, 1894

(Miocene, North America) forming the sister clade of *Brachypotherium* Roger, 1904

(Miocene, Old World) + *Brachydiceratherium* Lavocat, 1951). *Brachydiceratherium* includes

eight species spanning the late Oligocene to Late Miocene in Europe and Asia. All

teleoceratine genera except *Diaceratherium* span considerable geographical and

stratigraphical ranges, likely related to their ultra-generalist ecological preferences.

Keywords: Rhinocerotidae, *Brachydiceratherium shanwangense*, Tagay, Early Miocene,

Siberia, Lake Baikal, phylogeny, biogeographical history.

INTRODUCTION

Although they are nearly extinct today, rhinocerotids were one of the most widespread

and successful groups of large mammals on all the northern continents for over 40 million

years. They have occurred across Eurasia and North America since middle Eocene times, and

are known from Africa since the Early Miocene (e.g., Prothero et al., 1989; Antoine et al.,

2003, accepted; Geraads, 2010). They have also occupied many different locomotory modes,

ranging from slender- and long-legged savannah roamers (e.g., elasmotheriines) to hippo-like

forms that apparently lived along rivers and lakes (teleoceratines; Prothero et al., 1989;

Antoine, 2002). Most hippo-like rhinocerotids are gathered within teleoceratines, a clade at

the tribal to sub-tribal level, the phylogenetic relationships of which have never been fully

elucidated (Antoine, 2002; Lu et al., 2021). Most teleoceratines had skulls that were either

hornless or with a small nasal horn, barrel-shaped bodies, and shortened limb bones.

Teleoceratines span the late Oligocene–latest Miocene in Eurasia (Antoine, in press), the

Miocene in Africa (Geraads & Miller, 2013), and the early Miocene–early Pliocene in North

and Central America (Prothero, 2005). Most of them are interpreted to have been browsers

(based on both dental morphology and isotopic studies; MacFadden, 1998; Hullot et al.,

2021).

In this study, we describe a skeleton of a teleoceratine from Lower Miocene deposits of

Olkhon Island, Lake Baikal area, Siberia. We identify its species assignment and compare it

to most teleoceratine species described from Eurasia. This in-depth comparison forms the

basis for performing a parsimony analysis of phylogenetic relationships among Eurasian

Teleoceratina, and for discussing key events in the paleobiogeography of teleoceratine

rhinocerotids.

LOCALITY AND GEOLOGICAL SETTINGS

Lake Baikal, located in the Baikal Rift System, is morphologically characterised by

three basins (Southern, Central and Northern). The Southern and Central basins are thought to

have existed permanently since the Paleogene, whereas the Northern Basin did not develop

before the Late Miocene (Mats et al., 2010, 2011). Olkhon Island (Russian: Ольхон) is

located in the transitional zone between the Central and the Northern basins of Lake Baikal. It

is separated from the mainland in the west by a shallow Maloe More strait (Russian: Малое

Море; in English literally the Small Sea) of the Northern Basin that extends far to the south.

In the south, Maloe More strait is connected through the narrow Olkhonskie Vorota strait

(Russian: Ольхонские Ворота; in English literally the Olkhon Gate) to the central part of

Lake Baikal. From the northwestern part of Olkhon Island, one locality known as Tagay or

Tagai (Russian: Тагай or Тогай) has yielded numerous terrestrial fossils of the Neogene (Fig.

1). The Neogene sediments in Tagay Bay belong to the Tagay Formation (Logachev et al.,

1964; Mats et al., 2001; Mats, 2013, 2015). Sediments are exposed in the northeastern part of

the bay in a steep erosional cliff up to 15 m high. Elsewhere along the shores of the bay, this

cliff is levelled by landslides. The cliff borders a large landslide cirque and a sandy beach

below.

The Tagay locality was discovered in the 1950s (Kitainik & Ivaniev, 1958). The first

paleontological studies of the large mammals were performed in 1958 under the direction of

N.A. Logachev (Logachev et al., 1964). Studies of small mammals have been carried out

since the 1970s (Pokatilov, 2004; Daxner-Höck et al., 2022). Tagay preserves an abundant

fossil fauna that includes molluscs and vertebrates such as fish, amphibians, reptiles, birds and

mammals. However, a significant part of the paleontological material was determined for the

longest time only tentatively: Mustelidae and Felidae among carnivorans, *Anchitherium* sp,

Metaschizotherium(?) sp., and *Dicerorhinus*(?) sp. among perissodactyls, *Palaeomeryx* sp.

and Bovidae indet. among artiodactyls (Logachev et al., 1964). The reexamination of

artiodactyl remains led to the identification of Cervidae (*Amphitragulus boulangeri*,

Lagomeryx parvulus, *Stephanocemas* sp.), Palaeomerycidae (*Orygotherium tagaiense*,

Palaeomeryx cf. *kaupi*) and Anthracotheriidae (*Brachyodus intermedius*) (Vislobokova, 1990,

1994, 2004). Chelonians were studied by Khosatzky and Chkhikvadze (1993) and the

ichthyofauna by Filippov & Sytchevskaya (2000).

Small mammals from the Tagay-1 section were recently revised, with a list of 21 taxa

documenting erinaceids, talpids, plesiosoricids, and soricids among eulipotyphlans,

palaeolagid lagomorphs, and sciurids, aplodontids, mylagaulids, glirids, castorids, eomyids,

and cricetodontine muroids among rodents (Daxner-Höck et al., 2022).

The sedimentological, stratigraphical, and palaeontological aspects of the sediments were

described by Kossler (2003). A new phase of the study of Tagay locality started in 2008, with

numerous publications since that time (Rage & Danilov, 2008; Klementiev, 2009; Danilov et

al., 2012; Syromyatnikova, 2014, 2015; Tesakov & Lopatin, 2015; Klementiev & Sizov,

2015; Zelenkov, 2016; Sotnikova et al, 2021).

The Tagay Formation consists of alternating beds of clays and clayey sands containing

interlayers and lenses of carbonate concretions of diagenetic origin. Deposits rest upon the

crystalline basement, and are submerged below water to the south. Clay beds are mostly green

and brown, sometimes black. There are also lenses and interlayers of brick red and red clay

and loam. Bone beds, deposited in successive sedimentary cycles, were given letters from top

to bottom (i.e., downsection: A–H; Fig. 2 A, B). Most clay beds have predominant

ferruginous-magnesian montmorillonites composition. A remarkable feature of these clay

sediments is the high (up to 8%) content of silt-psammite-psephite admixtures. Moreover,

most psammite-psephitic fragments are not rounded and have angular and indented outlines,

which indicates a lack of transportation. The lithology of the sections and further details on

the bone beds were described in other studies (Logachev et al., 1964; Sizov & Klementiev,

2015).

Neogene continental deposits in the late early Miocene Tagay locality have yielded a

diverse vertebrate fauna. Following most works, the age of the Tagay Fauna correlates to the

European Mammal Zones MN3 through MN5 (20–15 Ma; Rössner & Mörs, 2001;

Vislobokova, 2004; Klementiev & Sizov, 2015; Sotnikova et al., 2021). Other researchers

have correlated the Tagay fauna or to MN 7+8 and Chinese Mammal Unit NMU7 (13–11 Ma;

Daxner-Höck et al., 2013). More recently, Daxner-Höck et al. (2022) proposed a more precise

age of ~16.5-16.3 Ma based on micromammalian bicostratigraphy and the magnetic polarity

pattern of the Tagay-1 section (Fig. 1 B), which is in agreement with our preferred interval.

MATERIAL AND METHODS

All the remains described here belong to a single adult individual (IZK79-1-08C-1/),

stored in the collection of the Institute of the Earth's Crust (Irkutsk, Russia). Alexey

Klementiev and Gennady Turkin discovered this skeleton in 2008 at Tagay (Fig. 2 B, C)

(Klementiev, 2009).

Capital letters are used for upper teeth (I, C, D, P, M), and lower-case letters for lower teeth

(i, c, d, p, m). Rhinocerotid dental terminology follows Heissig (1969, 1972a: pl. 13) and

Antoine (2002), while dental and skeletal measurements were taken according to Guérin

(1980). Anatomical descriptions follow basically the same sequence as in Antoine (2002), and

Antoine et al. (2010). Dimensions are given in mm.

The stratigraphical framework is based on the Neogene geological time scale and

European Land Mammal Ages (Hilgen, Lourense & Van Dam, 2012; Raffi et al, 2020).

3D-rendering

All bones of the rhinoceros were scanned with a resolution of 0.25 mm using a

RangeVision Smart 3D scanner. RangeVision Smart has three areas of scanning and is

equipped with colour cameras 1.3 megapixels. We used the associated RangeVision 2020.2

software for visualization, segmentation and 3D rendering.

Phylogenetic analysis

The phylogenetic analysis was based on 282 cranio-mandibular, dental, and postcranial

characters primarily derived from the dataset of Antoine (2002, 2003) which was scored on 31

ceratomorph species (one tapirid and 30 rhinocerotoids). All multistate characters were

treated as additive, except for the characters 72, 94, 102, 140, 187, and 269 (non-additive; as

in Antoine, 2002).

The living Brazilian tapir *Tapirus terrestris* (Linnaeus, 1758), the Eocene non-

rhinocerotid rhinocerotoid *Hyrachyus eximius* Leidy, 1871 and the Paleogene stem

rhinocerotids *Trigonias osborni* Lucas, 1900 (Eocene of North America) and *Ronzotherium*

filholi (Osborn, 1900) (Oligocene of Western Europe) were selected as outgroups. We also

included a branching group (Antoine, 2002, 2003; Orliac et al., 2010; Boivin et al., 2019),

consisting of non-teleoceratine taxa, with the aim of testing the monophyly and relationships

of the Teleoceratina among the Rhinocerotinae. These consist of 12 species, including an early-diverging representative of Rhinocerotinae (*Plesiaceratherium mirallesi* (Crusafont, Villalta & Truyols, 1955)), three species of Aceratheriini (*Aceratherium incisivum* Kaup, 1832, *Acerorhinus zernowi* (Borissiak, 1914), and *Alicornops simorrense* (Lartet, 1851)), and eight members of the Rhinocerotina, encompassing all five living rhinoceroses, namely the Indian rhino (*Rhinoceros unicornis* Linnaeus, 1758), the Javan rhino (*Rhinoceros sondaicus* Desmarest, 1822), the Sumatran rhino (*Dicerorhinus sumatrensis* (Fischer, 1814)), the white rhino (*Ceratotherium simum* (Burchell, 1817)), and the black rhino (*Diceros bicornis* (Linnaeus, 1758)), in addition to three fossil species: *Lartetotherium sansaniense* (Lartet in Laurillard, 1848) (Miocene of Europe; Heissig, 2012), *Gaindatherium browni* Colbert, 1934 (Miocene of South Asia; Heissig, 1972a; Antoine, in press), and *Nesorhinus philippinensis* (Von Koenigswald, 1956) (early Middle Pleistocene of the Philippines; Antoine et al., 2022 and references therein).

The ingroup sensu stricto (Teleoceratina) comprises 15 taxa, with *Teleoceras fossiger* Cope, 1878 (late Miocene to earliest Pliocene, North America), *Brachypotherium brachypus* (Lartet in Laurillard, 1848) (late early and middle Miocene, Eurasia), *Brachypotherium perimense* (Falconer & Cautley, 1847) (Miocene, South Asia), *Prosantorhinus germanicus* (Wang, 1929) (late early and middle Miocene, Europe), *Prosantorhinus douvillei* (Osborn, 1900) (late early and early middle Miocene, Europe), *Prosantorhinus laubei* Heissig & Fejfar, 2007 (early Miocene, central Europe), and a comprehensive sample of taxa either classically or more recently assigned to *Diaceratherium* Dietrich, 1931. These consist of the type species *D. tomerdingense* Dietrich, 1931 from the earliest Miocene of Tomerdingen (Germany), *D. lemanense* (Pomel, 1853) from the latest Oligocene-early Miocene of Western Europe (also described under the *Diceratherium* (*Brachydiceratherium*) *lemanense* combination by Lavocat, 1951), *D. aurelianense* (Nouel, 1866) from the early Miocene of Western Europe.

The taxonomic sample also includes *D. asphaltense* (Depéret & Douxami, 1902) from the

earliest Miocene of Western Europe, *D. fatehjangense* (Pilgrim, 1910), from the Miocene of

Pakistan and early Miocene of Kazakhstan (previously described as "*Brachypotherium*

aurelianense Nouel, var. nov. *Gailiti*" by Borissiak, 1927), and *D. aginense* (Répelin, 1917)

from the earliest Miocene of Western Europe. Lastly, we have considered *D. shanwangense*

(Wang, 1965) from the late early Miocene of eastern China (Shanwang; Lu et al., 2021),

Japan, and eastern Siberia (Tagay; this work), and *D. lamilloquense* Michel, in Brunet et al.,

1987 from the late Oligocene of France. We also included *Aceratherium gajense intermedium*

Lydekker, 1884, which has disputed taxonomic affinities. It was previously assigned to the

aceratheriine genera *Subchilotherium* (e.g., Heissig, 1972a) or *Chilotherium* (e.g., Khan et al.,

2011), although based on a parsimony analysis taking into account the holotype and original

hypodigm, Antoine et al. (2003) considered that it might be a teleoceratine instead, of

uncertain generic assignment. The recognition of associated dental and postcranial remains

from the Potwar Plateau (late early to early late Miocene, Pakistan) allowed for defining the

new combination *Diaceratherium intermedium* (Lydekker, 1884) for this taxon, as recently

proposed by Antoine (in press).

Three representatives of Teleoceratina, *Diaceratherium* cf. *lamilloquense* from the late

Oligocene of Thailand (Marivaux et al., 2004), *Brachypotherium gajense* (Pilgrim, 1910),

from the late Oligocene–earliest Miocene of Pakistan, and *Prosantorhinus shahbazi* (Pilgrim,

1910), from the early Miocene of Pakistan (combinations proposed by Antoine et al., 2010

and Antoine, in press) were not included in the analysis, due to their poorly-known

hypodigms, restricted to a few elements.

Moreover, *Diaceratherium askazansorense* Kordikova, 2001 from the early Miocene of

Kazakhstan was not included, as dental and postcranial elements assigned to this taxon

closely resemble those of *Pleuroceros blanfordi*, a stem member of Rhinocerotinae (early

Miocene of South Asia; Antoine et al., 2010; Prieto et al., 2018) and, to a lesser extent, of

Pleuroceros pleuroceros (earliest Miocene of western Europe; Antoine et al., 2010; Antoine

& Becker, 2013). Similarly, the early late Oligocene species *Diaceratherium massiliae*

Ménouret & Guérin, 2009 was recently shown to be a junior synonym of the short-limbed and

early-diverging rhinocerotid *Ronzotherium romani* Kretzoi, 1940, through a re-examination of

most available material and the recognition of new associated dental and postcranial

specimens in Switzerland (Tissier et al., 2021).

Details on the specimens, collections, direct observation and/or literature used for

scoring taxa, with references used) are provided as taxon notes in the morphological data

matrix (supplementary file S3).

The parsimony analyses were performed through the heuristic search of PAUP 4 3.99.169.0

(Swofford, 2002), with tree-bisection-reconnection (reconnection limit = 8), 1000 replications

with random addition sequence (10 trees held at each step), gaps treated as missing, and no

differential weighting or topological constraints. Branch support was estimated through

Bremer indices (Bremer, 1994), also calculated in PAUP 4 3.99.169.0.

Systematics

Generic and suprageneric systematics follow the present parsimony analysis (see

below).

SYSTEMATIC PALAEONTOLOGY

Order Perissodactyla OWEN, 1848

Family Rhinocerotidae GRAY, 1821

Subfamily Rhinocerotinae GRAY, 1821

Tribe Rhinocerotini GRAY, 1821

Subtribe Teleoceratina HAY, 1902

Genus *Brachydiceratherium* LAVOCAT, 1951

Syn. *Diaceratherium* DIETRICH, 1931 (partim)

Type species: *Acerotherium lemanense* Pomel, 1853 by subsequent designation

(Lavocat, 1951)

Included species: *Rhinoceros aurelianensis* Nouel, 1866 from the early Miocene of

Western Europe; *Aceratherium intermedium* Lydekker, 1884, from the early–late Miocene of

the Indian Subcontinent and China (Deng and Gao, 2006; Antoine et al., 2013; Antoine, in

press); *Diceratherium asphaltense* Depéret & Douxami, 1902 from the earliest Miocene of

Western Europe; *Teleoceras fatehjangense* Pilgrim, 1910, from the Miocene of Pakistan and

early Miocene of Kazakhstan (senior synonym of "*Brachypotherium aurelianense* Nouel, var.

nov. *Gailiti*" by Borissiak, 1927); *Teleoceras aginense* Répelin, 1917 from the earliest

Miocene of Western Europe; *Plesiaceratherium shanwangense* Wang, 1965 from the late

early Miocene of eastern China (Shanwang; Lu et al., 2021), Japan, and eastern Siberia

(Tagay; this work); *Diaceratherium lamilloquense* Michel, in Brunet et al., 1987 from the late

Oligocene of France.

Diagnosis: Teleoceratines with a small nuchal tubercle, articular tubercle smooth on the

squamosal, with cement present on cheek teeth, protocone always constricted on P3-4, labial

cingulum usually absent on lower premolars and always present on lower molars, foramen

vertebrale lateralis present and axis-facets transversally concave on the atlas, a postero-distal

apophysis low on the tibia, and a latero-distal gutter located posteriorly on the fibula.

Distinct from *Diaceratherium tomerdingense* in possessing a long metaloph on M1-2,

no mesostyle on M2, a distal gutter on the humeral epicondyle, an anterior side of the

semilunate with a sharp distal border, no posterior expansion on the pyramidal-facet of the

unciform, and a trapezium-facet present on the McII.

Differs from representatives of *Brachypotherium* in having close parietal crests, a

median ridge on the occipital condyles, a mandibular symphysis less massive, a labial

cingulum usually or always absent on upper premolars, an external groove developed on the

ectolophid of lower cheek teeth, a V-shaped lingual opening of the posterior valley of lower

premolars (in lingual view), a paraconid developed on p2, no second distal radius-ulna facet, a

symmetric semilunate-pyramidal distal facet, a posterior McIII-facet present on the McII, and

a fibula-facet subvertical on the astragalus.

Distinct from species referred to as *Prosantorhinus* in showing no latero-ventral

apophysis on the nasals, close fronto-parietal crests, ad no posterior groove on the processus

zygomaticus of the squamosal.

Geographical and stratigraphical range: Late Oligocene and Miocene of Eurasia,

with an Early Miocene climax.

Brachydiceratherium shanwangense (Wang, 1965)

See synonymy list in Lu et al. (2021)

Holotype: IVPP V 3026, left maxilla with upper cheek tooth series (P2-M3), stored at

the Institute of Vertebrate Paleontology and Paleoanthropology, Chinese Academy of

Sciences (IVPP).

Stratum typicum and locus typicus: Early Miocene (Shanwang Formation,

Shanwangian Age/Stage of China, middle Burdigalian); Xiejiahe locality, Shanwang Basin,

Shandong, China (see Lu et al., 2021).

Diagnosis: Representative of *Brachydiceratherium* with a lateral apophysis present on

the nasals, a median nasal horn present on the nasals, premolar series short with respect to the

molar series, roots distinct on the cheek teeth, crochet always simple and lingual cingulum

usually absent and always reduced on P2-4, crista always present on P3, protocone strongly

constricted on M1-2, lingual cingulum usually absent on lower premolars and always absent

on lower molars, d1/p1 absent in adults, glenoid fossa with a medial border straight on the

scapula, distal gutter absent on the lateral epicondyle of the humerus, proximal radius-ulna

facets always fused, and trochanter major low on the femur.

Distinguished from *Bd. lamilloquense*, *Bd. lemanense*, *Bd. asphaltense*, and/or *Bd.*

aurelianense in having I1s oval in cross section, no labial cingulum on upper cheek teeth, a

strong paracone fold on M1-2 and a constricted hypocone on M1, M3s with a triangular

occlusal outline, a radius with a high posterior expansion of the scaphoid-facet, a femoral

head hemispheric, an astragalus with a laterodistal expansion, very low-and-smooth

intermediate reliefs on metapodials, and a long insertion of m. interossei on lateral

metapodials.

Differs from *Bd. aginense* in having a processus postorbitalis on the frontal bone and a

median ridge on the occipital condyle, but no posterior groove on the processus zygomaticus

of the squamosal, molariform P2s (protocone and hypocone lingually separate), a long

metaloph on M1-2, a posterior groove on M3, a shallow gutter for the m. extensor carpi on the

radius, a posterior MtII- MtIII facet developed, but no cuboid-MtIII contact.

Distinct from *D. intermedium* in showing usually a lingual cingulum on upper molars, a

strong paracone fold on M1-2, a lingual cingulum usually absent on lower premolars, and a

right angle between the cuboid-facet and the base of the tuber calcanei on the calcaneus

Differs from *Bd. lemanense* in possessing a low zygomatic arch with a processus postorbitalis,

a small processus posttympanicus and a well-developed processus paraoccipitalis.

Distinct from *Bd. asphaltense* in having closer fronto-parietal crests and a brachycephalic

shape.

Differs from *D. lamilloquense* in showing a protoloph joined to the ectoloph on P2 and

molariform P3-4s (protocone and hypocone lingually separate).

Differs from *D. aurelianense* in having no metaloph constriction on P2-4 and a protocone

weakly developed on P2.

Geographical and stratigraphical range: Late early Miocene of the Shanwang Basin,

Shandong Province, China (see Lu et al., 2021) and of Irkutsk Region, Russia (Tagay locality,

Olkhon Island, Lake Baikal).

Material studied: IZK79-1-08C-1, almost complete skeleton, including the skull

(occipital, parietal, frontal, the right zygomatic and lacrimal, both nasals, and temporals with

processes and also premaxillae), the jaws, most vertebrae and ribs, both humeri, radii and

ulnae, both femora, tibiae, right fibula, most metacarpals, and several metatarsals and

phalanges. The skeleton described herein was found disarticulated at the junction of layers of

sand and clay (Fig. 2 B, C). In general, the right side of the individual is much better

preserved than the left one.

DESCRIPTION

Skull

The skull (Fig. 3, Table 1) was found disarticulated, but there is no doubt that the separate bones belong to the same individual, because they were found in close proximity to one another with no extraneous elements, and they fit together well. The temporal, zygomatic and lacrimal, nasal, frontal, parietal and occipital fit each other perfectly. The remaining bones are matching in size, colour and texture. The skull is short and relatively wide (Length from condyles to nasals = 540 mm, Width at the frontals ≈ 190 mm), belonging to a large-sized adult rhinocerotid. The separated nasal bones are long and longer than the preserved part of the premaxilla, relatively thin and bear a lateral apophysis. Roughness for a small nasal horn is preserved at the tip of the nasals. In lateral view, the foramen infraorbitalis and the posterior border of the U-shaped nasal notch are both located above the P3, while the anterior border of the orbit is above the M1. The minimum distance between the posterior edge of the nasal notch and the anterior border of the orbit is 67.2 mm.

Cranial features. The skull was partly destroyed and some elements were reconstructed in anatomical position by one of us (AS). It is short, broad, and elevated. The dorsal profile of the skull is concave, with a small protuberance for a short nasal horn and an upraised parietal bone (50°). In lateral view, the nasals have a small ventrolateral prominence (lateral apophysis, sensu Antoine, 2002). The maxilla is badly damaged and the area of the foramen infraorbitalis is restored on both sides. Nevertheless, based on the preserved part of the maxilla, a position above P4 can be hypothesised. The posterior end of the nasal notch is located above the anterior part of P3. The nasal septum is not ossified at all. The premaxillae are broken rostrally. They form a short and elevated strip, slightly dipping frontward, with a deep ventral sulcus. Relations between nasal and lacrimal bones are not observable, and neither are the lacrimal processi. The anterior border of the orbit is situated above the middle of M1. On the frontal, a pair of smooth tubercles lay on the dorsal and posterodorsal edges of

the orbit (processus postorbitalis). The anterior base of the processus zygomaticus maxillari is

low, ~1 cm above the neckline of molars. The zygomatic arch forms a straight, low, and

oblique strip, with parallel dorsal and ventral borders. It is parallel to the dorsal outline of the

skull, with a rounded and rugose posterodorsal tip. A marked processus postorbitalis deforms

the dorsal edge of the zygomatic process, at the junction between the jugal and the squamosal.

Its tip, located on the latter bone, has a rugose aspect. Most of the temporal fossa elements are

not preserved and it is therefore impossible to consider the shape and relations of the foramina

sphenorbitale and rotundum. The area between the temporal and nuchal crests is depressed,

forming a deep gutter. The external auditory pseudo-meatus is partly closed ventrally. The

posterior side of the processus zygomaticus is flat in lateral view (no posterior groove). The

occipital side is inclined up- and forward, with a very salient nuchal tubercle (although small,

i.e., not extended on a wide area), determining a diamond-shape to the skull in dorsal view.

The occipital condyles are oriented in the same axis as the skull in lateral view. The posterior

tip of the tooth row reaches the posterior third of the skull. The pterygoids are not preserved,

as most of the basicranium, vomer, and basal foramina. The skull is brachycephalic

(interzygomatic width/total length ~0.57; Table 1). As observable in dorsal view, the nasals

have a sharp tip. They are long and unfused, fully separate by a deep groove from tip to tip.

There were no lateral nasal horns, but a small median nasal horn, as unambiguously shown by

the presence of axial vascularised rugosities in the anterior quarter of the nasal bones. In

contrast, the frontal bones have a smooth aspect, thus indicating the absence of a frontal horn.

The orbits were not projected laterally. The zygomatic arches are 1.51 times wider than the

frontals. From this frontal ambitus, run posteriorly two straight and smooth frontal crests,

getting closer by the parietals (minimum distance = 30 mm; Table 1), and then abruptly

diverging and forming an occipital crest that is concave posteriorly. The transition from the

maxilla to the processus zygomaticus maxillary is progressive, with no brutal inflection. The

articular tubercle of the squamosal is smooth (in lateral view) and straight (in sagittal view).

The right processus postglenoidalis forms a rounded right dihedron in ventral view. The

foramen postglenoideum is remote from the latter. The left one is not preserved. The occipital

side is wide and, accordingly, the processus posttympanicus and the processus paraoccipitalis

are distant. The former is poorly developed, while the latter is very long, slender, and vertical.

The foramen magnum is not preserved well enough to allow any observation. The occipital

condyle has a median ridge but no medial truncation.

Mandible

In lateral view, the symphysis is upraised, with an angular ventral profile determined by

two successive inflections. The foramen mentale is widely open and located below p2 (left)

and p3 (right). The corpus mandibulae is low, with a straight ventral border (Table 2). It is

getting gradually higher to the mandibular angle, smooth, rounded and hugely developed.

There is a shallow vascular incisure. The ramus is low, with a posterior border that is oblique

up- and forward and a vertical anterior border. The processus coronoideus is high, tapering

dorsally, and somewhat concave posteriorly. The condyloid process is high and sharp-edged,

separate from the latter by a deep mandibular notch. In dorsal view, the symphysis is massive,

well developed anteroposteriorly and narrow, with i2s and lateral edges parallel and two

circular alveoli for small i1s. The posterior border of the symphysis is located between the

trigonids of p3. The tooth rows are more parallel than the bodies (Fig. 4), which widely

diverge posteriorly. The spatium retromolare is wide on both sides. The mylohyoid sulci are

present but very shallow. The foramen mandibulare opens below the teeth-neck line.

Dental material

The dental formula is 1-0-4-3/2-0-3-3. No decidual dentition is known.

Upper dentition (Fig. 4, Table 3). The first upper incisors are not preserved, but

straight and sagittally-elongated alveoli point to an oval cross section for them (as usual in

teleoceratines). There are no I2, I3, or C. The premolar series is short with respect to the

molar series (LP3-4/LM1-3*100=48.7; Lp3-4/Lm1-3*100=45.8), which is further highlighted

by the small size of P2 and p2. The enamel is thick, wrinkled and corrugated, and partly

covered with a thin layer of cement. Teeth are low crowned, with roots partly joined. The

labial cingulum is absent on the upper cheek teeth. A thick paracone fold is present on P2-M3,

vanishing with wear on P2-M1 and marked until the neck on M2-3. There is no metacone fold

or mesostyle on the upper cheek teeth. Short and wide crochet is present on P3-4 (always

simple), but absent on P2. There is no metaloph constriction on P2-4. The lingual cingulum is

absent on all upper cheek teeth, except for a small tubercle on the anterolingual base of the

hypocone on both P4. The postfossette forms a small and deep isometric pit. The antecrochet

is getting stronger backward, from absent on P2 and short on P3-4 to very elongate on M1-3.

The first upper cheek tooth is most likely a persistent D1: it is much more worn than other

teeth and the enamel is also much thinner. It is preserved on the right side and its presence is

further attested by two alveoli on the left side (heart-shaped anterior root; peanut-shaped

posterior root). It has a sharp anterolingual cingulum, a straight lingual edge, and rounded

posterior and labial edges. P2-4 are fully molariform (bilophodont, with an open lingual

valley). On P2, the metaloph is transverse labially, but curved posterolingually due to the

position of the hypocone. The latter is much more developed than the protocone. The

protoloph is thin but continuous and transversely oriented. There is no medifossette on P3-4,

but a short crista on P4 and on P3 (mostly wiped out by wear on P3). The protocone is

constricted anteriorly on P3-4. The metaloph forms a dihedron on P3-P4, with the crochet as a

tip and the hypocone located posterior to the metacone. The protoloph is complete and

continuous and there is no pseudometaloph on P3. The metacone is not constricted or isolated

on P3-4. The crochet is long and sagittal on M1-M3, with a rounded tip on M1, and a sharp

tip on M2-3. There is no crista, medifossette, or cristella on upper molars. The lingual

cingulum is restricted to a small pair of tubercles on M2s and a smooth ridge on the hypocone

of M3. The protocone is strongly constricted on M1-3, and trefoil shaped on M3. The

parastyle is short and sagittal on M1-3; the paracone fold is very salient on M2 and especially

on M3. The metastyle is very long on M1-2. The metaloph is almost as long as the protoloph

on M1-2. In lingual view, the protocone is increasingly developed sagittally from M1 to M3.

A deep groove carves the anterolingual side of the hypocone on M2, and a shallower one is

observed on M1. The ectoloph is straight on M1 and concave on M2. The antecrochet and the

hypocone are close but separate on M1-2. There is no lingual groove on the lingual side of

M2. The posterior cingulum is complete on M1-2 and the postfossette is still narrower and

deeper than on premolars. The right M3 has a triangular outline in occlusal view, with a

straight ectometaloph (the left M3 is not preserved). The protoloph is transversely developed.

There is no posterior groove on the ectometaloph and the labial cingulum is restricted to a low

and smooth spur covering the lingual third of the former.

Lower dentition (Fig. 4, Table 3). There are small circular alveoli for both i1s, between

the i2s, in the symphyseal part of the dentary but the shape of the concerned teeth is unknown.

The presence of a short p2 is attested by three closely-appressed alveoli on the right side (area

unpreserved on the left side), but no d1 or p1 was present, as attested by the sharp ridge

running anterior to p2's alveoli. There are no vertical rugosities on the ectolophid of p3. On

the lower cheek teeth, ectolophid grooves are developed (U-shaped) and vanishing before the

neck, trigonids are rounded and forming a right angle in occlusal view, metaconids and

entoconids are unconstricted. The bottom of the lingual valleys is V-shaped in lingual view on

lower premolars. On lower premolars, the lingual cingulum is restricted to a low ridge

continuing the anterior cingulum on the trigonid of p3s, and the labial cingulum consists of a

small edge obtruding the ectolophid groove on p4s. Lower molars lack a lingual cingulum but

a small cingular ridge partly obtrudes the ectolophid groove. The hypolophid is oblique in

occlusal view on m1-3. There is no lingual groove on the entoconid of m2-3. The posterior

cingulum of m3 forms a low, horizontal, and transversely-elongated ridge.

Poscranial skeleton (Tables 4, 5)

Atlas. The atlas is wide and short sagittally. In dorsal view, the transverse processes

(partly broken) and the alar notches are developed and the axis-facets are concave. In anterior

view, the rachidian canal has a bulb-like outline. The occipital condylar facets are kidney-like.

The foramen vertebralis cuts across the anterior third of the dorsal surface on both sides and it

is continued by a shallow groove laterally (for the vertebral artery). In posterior view, the

foramen transversarium is present, wide and partly hidden by the lateral expansion of each

axis-facet (Fig. 5, A).

Axis. The axis is stocky, with thick and cylindric dens and tear-shaped atlas-facets

(convex transversely) on the prezygapophyses. The spinous process is thick and carinated.

The foramen vertebrale is large and subtriangular. The postzygapophyses have wide and

circular facets for the first thoracic vertebra, forming a ~45° angle with the horizontal line.

The centrum is very long anteroposteriorly, with a pentagonal outline in posterior view (Fig.

5, B). Most thoracic vertebrae are preserved. They are massive, with heart-shaped centrums,

and stocky transverse processes. The dorsal spines are slender and oblique (45° with the

vertical line), with a length reaching up to 250% of the centrum height for the T4-6.

Scapula. The scapulae are partly preserved. They are elongated dorsoventrally, notably

due to their anteroposterior narrowness (H/APD = ~0.50). The scapular spine is straight,

much developed and with an extremely salient tuberculum bent caudally. There is no pseudo-

acromion. The tuberculum supraglenoidale is well distinct from the cavitas glenoidalis. The

medial border of the cavitas glenoidalis is straight, determining a semi-circular outline in

ventral view.

Humerus. Both humeri are almost complete (Fig. 6, A-E). The humerus is a slender

bone, with a straight diaphysis. The trochiter is high, with a smooth and rounded outline. The

caput humeri is wide and rounded, with a rotation axis forming a 40° angle with the vertical

line. The deltoid crest is elongated, almost reaching the mid-bone. The deltoid tuberosity is

not much salient. The fossa radii is wide and shallow. The fossa olecrani is higher than wide.

The distal articulation is egg-cup shaped (sensu Antoine, 2002, 2003), without marked median

constriction. The trochlea is half-conical and the capitulum humeri is half-cylindrical. There is

no synovial fossa ("trochlear scar") on the anterodorsal edge of the trochlea. The lateral

epicondyle is elongated dorsoventrally and its ventral border ends dorsal to the capitulum

humeri, lacking a distal gutter.

Radius. The two bones are complete and undistorted (Fig. 06, F-J). The anterior border

of the proximal articulation is straight in dorsal view but convex in anterior view. The radius

is slender, with a distal extremity larger than the proximal one in anterior view. The diaphysis

is quite slender, especially in its proximal half. It has a straight medial border in anterior view,

but it is posterolaterally concave, which determines a wide spatium interosseum brachii when

the ulna is in anatomical connection. The proximal ulnar facets are fused on both sides. The

insertion of the m. biceps brachii is wide but shallow, with two small pits. Ulna and radius are

independent, apart from the proximal and distal articular areas. On the anterodistal part of the

diaphysis, the gutter for the m. extensor carpi is not marked at all. There is only one distal

facet for the ulna on the lateral side of the bone. The posterior expansion of the scaphoid-facet

is high, forming a right-angled rectangle. There is a wide pyramidal-facet on the distal

articulation.

Ulna. The bone is sturdy, with a long and heavy olecranon, the tip of which is wide and

diamond shaped (Fig. 6, K-O). The diaphysis is straight, triangular in cross- section and as

robust as the radius shaft. It forms a ~135° angle with the olecranon in lateral view. The

humeral facet is saddle-shaped. The proximal radio-ulna facets form a continuous pad, with a

wide medial strip and a high triangular lateral facet. A smooth but salient anterior tubercle is

located above the distal end of the bone. There is neither a second distal radius-facet on the

medial side of the diaphysis nor semilunate-facet on the distal side. The almond-shaped distal

radius-facet is separate proximally from a salient horizontal ridge by a deep and rugose

depression. The pyramidal-facet is concavo-convex, with a quarter-circle outline in distal

view.

CARPUS. The carpus is very low and massive, especially with respect to slender

stylopodial and zeugopodial elements (Fig. 7). All carpals have salient tubercles on the

anterior aspect of the bones. The right hand is more complete than the left one.

Scaphoid. The scaphoid is low and massive, with equal anterior and posterior heights

(Fig. 7, A-C). The proximal radial facet is diamond shaped in proximal view. The

posteroproximal semilunate facet is strongly distinct. It is oval, wide, and separated from all

other facets. A deep depression hollows the lateral side between the semilunate-facets. The

anterodistal semilunate-facet is nearly flat and crescent shaped. The magnum-facet is concave

in lateral view. The trapezium-facet is smaller than other distal facets, but it forms a wide

triangle, separated from the trapezoid-facet by a smooth ridge.

Semilunate. The bone is compact. In proximal view, the anterior facet only contacts the

radius, whereas the wide posteromedial facet is for the scaphoid (Fig. 7, F-I). The anterior

side is smooth (not keeled or carinated), with a sharp distal tip. On the lateral side, both

pyramidal-facets are closely appressed. The proximal one is almond shaped and the distal one

is comma like. The posterior tuberosity is short. Most of the distal side is articulated, medially

with the magnum and laterally with the unciform.

Pyramidal. The bone is almost cubic. The proximal side is square shaped, with a

saddle-shaped ulna-facet (Fig. 7, J-M). The semilunate-facets are sagittally elongated, with a

half-oval outline for the proximal one and a crescent-like shape for the distal one. The

pisiform-facet is comma shaped, with a concave sagittal profile and it overhangs a strong

lateral tuberosity. The distal facet, for the unciform, forms a right isosceles triangle with

rounded angles. There is no magnum-facet.

Pisiform. The right pisiform is short, high, and spatulate, with large and triangular

radius- and pyramidal-facets (Fig. 7, N-P). Both facets are separated by a sharp ridge and

form a right angle. There is no strong constriction separating the thick body and the

articulated part. The medial edge of the body is straight and vertical.

Trapezium. The right trapezium is preserved. It is a small proximo-distally flattened

bone with a circular outline in proximal view. The proximal side is almost entirely occupied

by a wide pentagonal scaphoid-facet (compatible with the large-sized trapezium-facet on the

scaphoids). The latero-distal side bears a trapezoid-facet with a right-triangled shape,

overhanging a deep pit. All other sides have a rugose aspect and they are devoid of articular

facets.

Trapezoid. Only the right trapezoid is documented (Fig. 7, D-E). It is wider than high,

almost cubic. Only the anterior and posterior sides (oval and pentagonal in shape,

respectively) are free of articular surfaces. The proximal side, saddle shaped and tapering

posteriorly, responds to the scaphoid. In medial view, the trapezium-facet is restricted to the

posterior half, with a deep insertion pit located close to the anterior edge. The lateral facet is

al low rectangle for the magnum. The distal side is weakly concavo-convex, and it consists of

a pentagonal McII-facet.

Magnum. The magnum has a very low anterior aspect, with a subrectangular outline

and a salient horizontally-elongated median pad (Fig. 7, Q-T). The proximal border is straight

in anterior view. In medial view, the anteromedial facets are in contact throughout their whole

length (no anterior groove). In lateral view, the dorsal pulley for the semilunate forms a low-

diameter half circle, further determining a question mark proximal profile. The distal facet is wide and tapering posteriorly. The posterior tuberosity is broken on the left magnum, and it is very short on the right specimen.

Unciform. The bone is compact, with a posterior tuberosity wide and much developed sagittally (Fig. 7, Y-W). The anterior side is wide and low, with a pentagonal outline and a maximum height on its lateral tip. The proximal side has two anterior facets flat transversally and convex sagittally, separated by a sharp sagittal edge. The medial one, triangular, is for the semilunate while the lateral one, diamond shaped, is for the pyramidal. The latter has a wide posterolateral expansion joining the lateral edge and the McV-facet (located on the distal side) on the right unciform. This part is broken on the left one. From the medial tip, the distal and distolateral sides have three contiguous facets, responding to the McIII (small and quadrangular), McIV (bulb-shaped), and McV (oval and deeply concave sagittally), respectively. They are only separated by smooth sagittal grooves. The McV-facet is oblique, which could suggest the presence of a functional McV (see Antoine, 2002, 2003; Boada-Saña et al., 2008).

METACARPUS. The hand and pes have a mesaxonian Bauplan. Although no McV is preserved, the hand was probably tetradactyl, as hypothesised by the vertical facet on the McIV (see above). The metapodials have salient insertions for the m. extensor carpalis. Their shafts are robust (wide transversally and flattened sagittally), with neither distal widening nor clear shortening (no brachypody; see Antoine, 2002). The insertions for the m. interossei are long, reaching the mid-shaft on all available metapodials. The intermediate reliefs do not reach the anterior aspect of the distal articulation on metapodials. The intermediate relief is moderately high and quite sharp on the McIII, but low and smooth on medial and lateral metapodials.

McII. In proximal view, the proximal side consists of a large tear-shaped trapezoid-facet medial to a narrow sagittally-elongated and strip-like magnum-facet. In medial view, the trapezium-facet is large and comma shaped, higher in its posterior tip. In lateral view, the magnum-facet is a straight and low strip, separated from the McIII-facets over their length. The McIII-facets are fused into a curved strip with a shallow disto-median constriction. The distal articulation, for the phalanx 1, has a sub-square outline in distal view, with rounded anterior angles. Above it, is a wide and salient medial tuberosity (Fig. 8, A-E).

McIII. The bone has a straight shaft. The proximal side is dominated by a wide and pentagonal magnum-facet, contiguous to two narrow sagittal strip-shaped facets (medially for the McII and laterally for the McIV). In anterior view, the proximal side consists of a subvertical medial edge (McII-facet), a very wide magnum-facet, weakly-concave medially, and a much narrower, oblique and straight McIV-facet. The magnum-facet is almost invisible in anterior view. Indeed, its dorsal outline is not much convex in medial view. The McII-facets are broadly connected, forming a thick strip with a shallow constriction in its disto-median part. In lateral view, the anterior McIV-facet is low, elongated sagittally, and tear-shaped. It is disconnected from the oval posterior McIV-facet by a narrow but deep oblique groove. This articulated surface overhangs a deep circular depression. There is no postero-distal tubercle on the diaphysis. In distal view, the distal articulation is wide and subrectangular, with straight medial and lateral edges, rounded antero-medial and -lateral angles, and a m-like posterior edge, due to a low but sharp intermediate relief (Fig. 8, F-J).

McIV. The McIV is the shortest and most robust metapodial preserved. The shaft is concave laterally in anterior view. The proximal aspect is trapezoid, deeper than wide, with a narrow medial strip (the sagittally-elongated 'anterior' McIII-facet) and a wide unciform-facet. In proximal view, there is no postero-lateral pad, but a small anterolateral tubercle in front of the McV-facet. In medial view, the proximal McIII-facets are connected (right

specimen) and form a right dihedron (L-shape), with a high posterior facet. The McV-facet is

vertical, suggesting a functional McV, in good agreement with the orientation of the McV-

facet on the unciform. In distal view, the distal articulation forms a quarter circle, with a

posteromedial right angle. There is almost no intermediate relief on the McIV (Fig. 8, K-O).

Phalanges. Only three phalanges are preserved for the manus (left/right first phalanges

and left second phalanx for the McII). They have strong interphalangeal insertions and

tubercles. The phalanx 1 is low and massive, with a kidney-like proximal side (McII-facet,

lacking a groove responding to the intermediate relief). The distal facet is oval and

transversely transversally elongated. The phalanx 2 is still lower, with a proximal facet

perfectly matching in shape the distal facet on the phalanx 1. The distal facet is slightly

concave transversely and convex sagittally. Both facets have similar width and depth.

Coxal. The pubic bones and ischia are lacking on both sides but the ilia are well

preserved. Dorsally, the iliac crest is regularly convex. The wing of the ilium is spatulated.

The sacral tuberosity has a rounded triangular shape, with a rugose aspect. The coxal

tuberosity, partly broken, was probably thick and high, also with a rugose aspect. The caudal

gluteal line is smooth, with a concave outline (forming a semi-circular curve). The

acetabulum has a subcircular outline.

Femur. The bone is quite slender, with a shaft straight in anterior view, concave

anteriorly in lateral view, and compressed sagittally (Fig. 9, A-F). The anterior part of the

trochanter major is high, but the caudal part is very low, i.e., much lower than the wide and

hemispheric head. The fovea capitis is deep, low, and wide, with a triangular outline. The

trochanter minor is elongated dorsoventrally. Its distal end reaches the mid-height of the third

trochanter. The latter is developed, wider distally and with smooth lateral borders. The

anteroproximal border of the patellar condyle is curved, with a medial lip much more

developed and salient than its lateral counterpart. In lateral view, the medial lip of the trochlea

and the diaphysis determine a broken angle (130°). In distal view, the anterior border of the patellar trochlea is convex medially and straight and transverse laterally. The tibial condyles are separate from the patellar trochlea by a narrow groove. The intercondylar fossa is deep and narrow. The medial condyle, with a diamond-shaped outline, is much more developed than the lateral one. The medial epicondyle is also more salient than the lateral epicondyle.

Patella. The patella is massive, wider than high, and with a triangular and rugose anterior aspect. The medial border is straight and vertical. The posterior side, almost fully articulated, contacts the femoral cochlea, with a wide medial lip, triangular (wider distally), and a narrower trapezoid lateral lip. In vertical view, the latter lip is almost straight while the former is more concave. The most striking feature is the weak anteroposterior development of the bone with respect to other dimensions.

Tibia. The tibia is high and relatively slender, with heavy extremities (Fig. 9, K-O). The medial border of the diaphysis is strikingly straight in anterior view, which widely contrasts with the concave lateral border of the shaft. This impression is highlighted by the median half of the proximal articulation being much higher than the lateral one. The patellar ridge is thick and bulbous, with a rough surface. The patellar groove is deep, short dorso-ventrally, and regularly concave. The proximal peroneal articulation is located low on the tibia (no contact with the lateral femoral condyle). There is neither an anterodistal groove nor medio-distal gutter (for the tendon m. tibialis posterior). Tibiae and fibulae are independent, apart from articulated areas, thus determining a wide spatium interosseum cruris. The distal fibula-facet is low, elongated, and crescent shaped, overhung by a rugose triangular area. The posterior apophysis is low and rounded. In distal view, the outline is a trapezoid, wider than deep. The astragalar cochlea has two lips, the medial one being narrower and deeper and the lateral one wider and shallower.

Fibula. The diaphysis is straight and particularly slender, in sharp contrast with two thick ends (and the robustness of the tibia) (Fig. 9, G-J). The proximal end is nevertheless flattened sagittally, with a smooth proximal tibia-facet. The distal end is robust, with a deep laterodistal gutter for the tendon m. peronaeus, located posteriorly, immediately posterior to a huge tubercle. The distal fibula-facet is low, elongated sagittally, and crescent shaped. It is contiguous to a flat and rectangular astragalus-facet, oriented at ~25° with respect to the vertical line.

PES. The pes is not completely known (Fig. 10). The naviculars, cuneiforms, MtIIIs, and most phalanges are not preserved. The metatarsals are shorter than the metacarpals.

Astragalus. The astragalus is thick (APD/H = 0.76), wide and low (TD/H = 1.29). The fibula-facet is subvertical, wide and flat dorsoventrally (Fig. 10, A-C). The medial trochlear ridge is rounded, whereas the lateral one is sharper. The collum tali is very high (up to ¼ of the height), especially with respect to the general proportion of the bone. The caudal border of the trochlea is sinuous in dorsal view (with a falciform shape). There is no anterodistal trochlear notch, but a wide foramen for an insertion located distally to the concerned area, in the mid-collum tali. In anterior view, the distal border is deeply concave medially (navicular-facet) and straight and oblique laterally (cuboid-facet). The medial tubercle is low and rounded, but much projected medially. The distal articulation is not twisted with respect to the axis of the trochlea (<15°), in distal view. The calcaneus-facet 1 has a wide and very low, triangular laterodistal expansion. This facet is nearly flat in lateral view. The calcaneus-facets 2 (low oval) and 3 (tear shaped and low) are distinct and separate by a wide groove. In distal view, the distal articulation is much wider than deep, with a cuboid-facet particularly wide transversely. The posterior stop on that cuboid-facet is abrupt and prolongated medially by a similar transversely-elongated inflection on the navicular-facet.

Calcaneus. The calcaneus is robust, with a tuber calcanei massive and oval in

posteroproximal view (Fig. 10, D-F). This tuber calcanei is strongly vascularised and rugged

with salient muscle/tendon insertion areas, The tibia-facet is low, wide, and almond shaped,

while the fibula-facet is round and oblique with respect to the vertical and sagittal lines. The

astragalus-facet 1 is lozenge shaped in anterior view and almost flat. The facet 2 is oval, wider

than high and flat. It is separate from the smaller and semi-oval facet 3. The sustentaculum

tali is low and very wide. In lateral view, the cuboid-facet and the posterior border of the tuber

form a right angle and the processus, at the level of the sustentaculum tali, is deeper (in terms

of APD) than the tuber calcanei. The insertion for the m. fibularis longus forms a salient and

rugose pad, but without sharp ridges. On the distal side, the cuboid-facet forms a transversely-

elongated hexagon. It is mostly flat but concave in its mediodistal quarter.

Cuboid. The cuboid is compact, wide, and low (Fig. 10, G-I). In proximal view, the

large articular surface is oval, slightly tapering posteriorly, and split into two equally-

developed and sagittally-elongated facets. The astragalus-facet (medial) is separated from the

calcaneus-facet (lateral) by a narrow and shallow groove. The anterior side is low and

pentagonal in anterior view, with a sharp proximal tip. In medial view, there are four facets.

The anteroproximal one is very low and crescent like (navicular-facet). Distally to it is a much

larger semi-circular ectocuneiform-facet. The posteroproximal navicular-facet, broadly

joining the proximal facet for the astragalus, has an 8-shaped outline. Contiguous to it, but

distally, is a semi-circular posterodistal ectocuneiform-facet. The posterior tuberosity is short

sagittally and narrow, but quite elevated: its acuminated distal tip is positioned much more

distally than the distal articulation (MtIV-facet). The latter facet is flat and trapezoid, with

larger anterior, medial, and posterior sides and a shorter lateral border. There is no MtIII-

facet.

MtII. The bone is short and robust (Fig. 11, A-E). The proximal side, with a triangular

outline (widening posteriorly), responds to the entocuneiform (posteromedial facet,

pentagonal, and oblique), the mesocuneiform (proximal-most facet, wide and trapezoid), and

to the ectocuneiform (wide strip-like facet oblique and tapering anteriorly). In lateral view, the

MtIII-facets are vertical, with a large triangular anterior facet and a much lower, oval

posterior facet. Both are widely connected. The shaft is straight and subcircular in cross

section. The distal end is stocky and square in distal view. The distal articulation has almost

no intermediate relief, even in its posterior aspect.

MtIV. The bone is short and massive, with a heavy proximal end (Fig. 11, F-J). The

proximal side is entirely occupied by a flat and sub-square cuboid-facet. There are two

distinct proximal tubercles at the anterolateral and posterolateral angles, but no continuous

pad. In medial view, there are two equally-wide MtIII-facets. The anterior one is located more

dorsally, elevated and with a half-oval outline, connecting the proximal side. The posterior

one is oval, isolated, and anteroventrally-posterodorsally elongated. The shaft is slightly

concave laterally but straight in lateral view, with a strong laterodistal tubercle. The distal side

is entirely articulated, deeper than wide (APD>TD), and lacking an intermediate relief. Only

the lateral lip is slightly concave transversely in its posterior aspect.

Phalanges. Only the first phalanges for the MtII and MtIV are known. They have strong

interphalangeal insertions and tubercles. There is no groove responding to the intermediate

relief. The MtII phalanx 1 is almost cubic, with a circular and slightly biconcave proximal

side (MtII-facet). The distal facet (phalanx 2) is kidney shaped. The MtIV phalanx 1 is as

wide as but lower than the former phalanx. The proximal facet is kidney shaped and almost

flat. The distal facet is oval and elongated transversely. In both phalanges, the distal facet is

smaller than the proximal facet, but also slightly convex sagittally and concave transversely.

COMPARISON

In the next pararaphs, the comparison will be organised following anatomical regions (skull, mandible, teeth, and postcranial skeleton) and we will use the generic assignments as supported by the phylogenetic analysis.

The rhino from Tagay cannot be assigned to *Diaceratherium* (*sensu stricto*, see below) as it does not have teeth with wrinkled enamel, M1-2 with a short metaloph, M2 with a mesostyle, semilunates with a distal border of the anterior side rounded, or unciforms with a posterior expansion of the pyramidal-facet always present (all these features are diagnosing its type and only species, *D. tomerdingense*). It cannot belong either to *Brachypotherium*, the representatives of which have an occipital condyle without a median ridge, a mandibular symphysis very massive, a labial cingulum usually present on upper premolars and always present on upper molars, lower cheek teeth with a flat ectolophid, lower premolars with a lingual opening of the posterior valley U-shaped, p2 with a paraconid reduced, radius-ulnae with a second distal articulation, pyramidals with a distal semilunate-facet asymmetric, posterior facet always absent on the McII-McIII, and a fibula-facet oblique on the astragali. Contrary to representatives of *Prosantorhinus*, it has no sagittal fronto-parietal crest, posterior groove on the processus zygomaticus of the squamosal, or metacone fold on M1-2, but a constricted metaloph on M1 (diagnostic features of the genus). At last, it is differing from *Teleoceras fossiger* in a wide array of cranio-mandibular, dental, and postcranial features (e.g., foramen infraorbitalis above premolars, processus postorbitalis present on the zygomatic arch, occipital side inclined up-and frontward, low-crowned cheek teeth, crista always present on P3, atlas with a bulb-like rachidian canal cavity, scapula elongated, or navicular with a lozengic outline in vertical view).

On the other hand, in its general shape, proportions and anatomical features, the skull from Tagay closely matches that of *Brachydiceratherium shanwangense* (Wang, 1965) as recently described by Lu et al. (2021). It also resembles that of late Oligocene–early Miocene

teleoceratines from Western Europe classically assigned to *Diaceratherium*, except for the

type species (*D. tomerdingense*, for which only an isolated nasal bone is preserved). Within

Brachydiceratherium, the arched dorsal profile, the short and slender premaxillae, the

zygomatic arch (straight, oblique, with a marked posterodorsal angle, and an anterior tip

starting progressively), and the processus paraoccipitalis long and narrow make it have the

closest affinities with *Bd. shanwangense*, *Bd. aginense* (earliest Miocene; Répelin, 1917), and

Bd. asphaltense (Becker et al., 2018). The only differences with the former do concern the tip

of the nasal bones, pointing upward, having a small median bump (suggesting the presence of

a terminal nasal horn) and a distolateral apophysis, and the stockier zygomatic arch as

observed in the Tagay specimen. More specifically, the Tagay skull differs from that of *Bd.*

aginense in having a processus postorbitalis on the frontal bone and a median ridge on the

occipital condyle, but no posterior groove on the processus zygomaticus of the squamosal

(Répelin, 1917). It is distinct from *Bd. lemanense* in possessing a low zygomatic arch, with a

processus postorbitalis, a small processus posttympanicus and a well-developed processus

paraoccipitalis (Lavocat, 1951), and from *Bd. asphaltense* in having closer fronto-parietal

crests and a brachycephalic shape (Becker et al., 2018; Jame et al., 2019). In contrast, the

shape of the processus zygomaticus, but also the presence of a small median nasal horn and of

a concave occipital crest make is somewhat resemble *Bd. asphaltense*.

As for mandibular features, the Tagay jaw is also particularly resembling those of *Bd.*

shanwangense and *Bd. aginense* among representatives of *Diaceratherium*, with an upraised

symphysis (distinct from that of *Bd. lamilloquense* and of *Bd. aurelianense*), low corpus,

vertical ramus and a deep and laterally-salient mandibular angle, with a shallow vascular

incisure.

With respect to all other representatives of *Diaceratherium* now referable to

Brachydiceratherium (see phylogenetic discussion), the most distinctive dental features of the

Tagay rhinocerotid are a short premolar series (also observed in *Bd. shanwangense*), an

enamel wrinkled and corrugated at the same time, crochets simple and lingual cingula usually

absent and always reduced on P2-4, a protocone strongly constricted on M1, a lingual

cingulum usually absent on lower premolars and always absent on lower molars, and the

absence of d1/p1 at an adult stage (also observed in *Bd. shanwangense*). It can be further

distinguished from *Bd. lamilloquense*, *Bd. lemanense*, *Bd. asphaltense*, and/or *Bd.*

aurelianense by its I1s oval in occlusal view, the absence of labial cingulum on upper

premolars and molars, the presence of a strong paracone fold on M1-2 and of a constricted

hypocone on M1, and M3s with a triangular occlusal outline. Contrary to *Bd. lamilloquense*,

the rhino from Tagay has a protoloph joined to the ectoloph on P2, but also a protocone and a

hypocone lingually separate on P3-4 (molariform). With respect to *Bd. aurelianense*, it has no

metaloph constriction on P2-4 and a protocone weakly developed on P2. In other words,

dental remains from Tagay are strictly similar to those of *Bd. shanwangense* (Lu et al., 2021).

They further have very close affinities with those of *Bd. aginense* and *Bd. intermedium* among

Brachydiceratherium representatives. Nevertheless, the Tagay rhinocerotid differs from *Bd.*

aginense in bearing a protocone and a hypocone lingually separate on P2 (molariform) and

from both species in having a long metaloph on M1-2 and a posterior groove on M3.

Even though postcranial elements are not known in all *Brachydiceratherium* species, the

cervical vertebrae and/or limb bones from Tagay are perfectly matching those of *Bd.*

shanwangense (Lu et al., 2021). They are also very similar to those of *Bd. aginense* and of *Bd.*

intermedium, notably in terms of size and proportions. They differ, however, from all

representatives of the genus (including the latter species), in having a scapular glenoid fossa

with a straight medial border and a tibia-facet on the calcaneus, but no distal gutter on the

humeral lateral epicondyle. It can be further distinguished from *Bd. lamilloquense*, *Bd.*

lemanense, *Bd. asphaltense*, and/or *Bd. aurelianense* by a radius with a high posterior

expansion of the scaphoid-facet, a femoral head hemispheric, an astragalus with a laterodistal

expansion, the presence of very low-and-smooth intermediate reliefs on metapodials but also

a long insertion of m. interossei on lateral metapodials. The Tagay rhinocerotid differs from

Bd. aginense in bearing a shallow gutter for the m. extensor carpi on the radius, a posterior

MtII-facet developed on the Mt3, but no contact between the cuboid and the MtIII, from *Bd.*

intermedium in showing a right angle between the cuboid-facet and the base of the tuber

calcanei on the calcaneus, and from both species in having a scaphoid with equal anterior and

posterior heights, a short posterior tuberosity on the magnum, a wider astragalus (TD/H>1.2),

and a fibula-facet on the calcaneus.

In fact, the Tagay rhinocerotid individual is identical in all aspects to corresponding

specimens of the complete skeleton from the early Miocene of eastern China recently

assigned to as *Diaceratherium shanwangense* by Lu et al. (2021). The only differences lie in

the occipital crest being more concave in the Tagay skull than in the Shanwang one, following

the description by Lu et al. (2021). This feature is likely to document either sexual

dimorphism, ontogenetic variation, or interindividual variability. Accordingly, we consider

unambiguously the Tagay rhinocerotid as documenting *Brachydiceratherium shanwangense*.

PHYLOGENETIC ANALYSIS

We have first run a preliminary analysis (see supplementary materials: files S1 and S2),

with 32 taxa i.e., the Tagay individual and *B. shanwangense* scored as two distinct terminals.

In that analysis, these terminals differ in a single and only feature (char. 36: occipital crest

concave in the former). Accordingly, we have merged them into a single terminal, for running

the final analysis (see supplementary materials: files S3 and S4). A single most parsimonious

tree is retrieved (length = 1316 steps; consistency index = 0.2698; retention index = 0.4918;

Fig. 13; see supplementary files S3 and S4). Twenty-four characters are constant, due to their

original definition for solving phylogenetic relationships within Elasmotheriina (Antoine,

2002), a rhinocerotid subtribe the representatives of which are not included here. Character

distribution at each node and corresponding indices are detailed in the supplementary

materials (file S4). Suprageneric relationships within Rhinocerotinae (i.e., the clade including

Rhinocerotini + Aceratheriini) are consistent with those proposed by Antoine (2002, 2003),

Antoine et al. (2010, 2022), Becker et al. (2013), Tissier et al. (2021), and Pandolfi et al.

(2021): *Plesiaceratherium mirallesi* is the earliest offshoot among Rhinocerotinae (node 1; 26

unambiguous synapomorphies; Bremer Support [BS] > 5). Aceratheriini (node 3; nine

unambiguous synapomorphies; BS = 2) and Rhinocerotini (node 5; eight unambiguous

synapomorphies; BS = 2) are sister clades (node 2; 13 unambiguous synapomorphies; BS =

4). Rhinocerotina (node 6; 18 unambiguous synapomorphies; BS > 5) and Teleoceratina

(node 13; five dental and postcranial unambiguous synapomorphies; BS = 1) are sister clades

within Rhinocerotini (Fig. 13). Aceratheriini comprise *Alicornops simorrense* as a sister

species to the (*Aceratherium incisivum*, *Acerorhinus zernowi*) clade (node 4). Rhinocerotina

include the (*Lartetotherium sansaniense*, *Gaindatherium browni*) clade (node 7; seven

unambiguous synapomorphies; BS = 5) as the first offshoot, then *Nesorhinus philippinensis*

(node 8; seven unambiguous synapomorphies; BS = 3), and the living rhino species (node 9;

nine unambiguous synapomorphies; BS = 2), with the *Rhinoceros* clade (node 10; four

unambiguous synapomorphies; BS = 1) being sister group to the (*Dicerorhinus sumatrensis*

plus African rhinos) clade (node 11; 13 unambiguous synapomorphies; BS = 3). The clade of

living African rhinos is the most supported node of the tree (node 12; 38 unambiguous

synapomorphies; BS > 5).

In the next paragraphs, we will focus on the topology, node support (Bremer Support:

BS), and apomorphy distribution regarding the Teleoceratina. The monophyly of the subtribe

is weakly supported by five dental and postcranial unambiguous synapomorphies (BS = 1): I1

with an almond-shaped cross section, hypocone isolated by an anterior constriction on M2,

ulna with the olecranon and the diaphysis forming a closed angle, robust limbs, and lateral

metapodials with insertions of the m. interossei short. The earliest-diverging teleoceratine is

Diaceratherium tomerdingense. This species is defined by ten dental and postcranial

autapomorphies (teeth with enamel wrinkled and roots separate, P2-3 with an antecrochet

usually absent, M1-2 with a metaloph short, M2 with a mesostyle, humerus without a distal

gutter on the lateral epicondyle, semilunate with a distal border of the anterior side rounded,

trapezoid with a proximal border asymmetric in anterior view, unciform with a posterior

expansion of the pyramidal-facet always present, and trapezium-facet always absent on the

McII; Table 6). Next node (node 14) segregates the *Brachypotherium* clade (node 15) from all

other teleoceratines scored here (node 16). Node 14 (BS = 2) is weakly supported by three

postcranial unambiguous synapomorphies (proximal ulna-radius facets usually fused, gutter

for the m. extensor carpi weakly developed on the radius, and McII with anterior and posterior

McIII-facets fused). Eleven cranio-mandibular, dental, and postcranial synapomorphies define

Brachypotherium (node 15; BS = 2): occipital condyle without a median ridge, mandibular

symphysis very massive, labial cingulum usually present on upper premolars and always

present on upper molars, lower cheek teeth with a flat ectolophid, lower premolars with a

lingual opening of the posterior valley U-shaped, p2 with a paraconid reduced, radius-ulna

with a second distal articulation, pyramidal with a distal semilunate-facet asymmetric,

posterior facet always absent on the McII-McIII, and fibula-facet oblique on the astragalus.

The Bremer Support is low, due to an alternative topology with *B. perimense* being sister

taxon to the (*B. brachypus*, node 16) clade appearing at 1317 steps. *Brachypotherium*

brachypus is particularly well differentiated, with 27 unambiguous cranio-mandibular, dental,

and postcranial autapomorphies each (see Table 6). From node 16 diverge two clades, with

(*Teleoceras* plus *Prosantorhinus*) on the one hand (node 17), and all species classically

assigned to *Diaceratherium* except the type species (node 20). Node 16 (BS = 2) is supported

by eight cranio-dental and postcranial unambiguous synapomorphies: vomer rounded,

protocone constriction usually absent on P3-4, antecrochet always present on P4, lingual

cingulum always present on lower premolars, pyramidal- and McV-facets always separate on

the unciform, McIV with a trapezoid outline in proximal view, calcaneus-facets 2 and 3

always independent on the astragalus, and fibula-facet always present on the calcaneus. Node

17 (BS = 4) places the highly-divergent *Teleoceras fossiger* (39 cranio-mandibular, dental,

and postcranial unambiguous autapomorphies; Table 6) as sister species to *Prosantorhinus*,

through 14 cranio-mandibular, dental, and postcranial synapomorphies: base of the processus

zygomaticus maxillary low on the maxilla, zygomatic arch high, articular tubercle of the

squamosal concave, lingual groove (sulcus mylohyoideus) absent on the corpus mandibulae,

metaloph transverse and protoloph sometimes interrupted on P2, mesostyle present on M2, d2

with a posterior valley usually open, scapula spatulated and with a medial border straight on

the glenoid fossa, a trochanter major low on the femur, MtII-facet always absent and cuboid-

facet present on the MtIII, and metapodials with high and acute intermediate reliefs.

Prosantorhinus (node 18; BS = 4) is monophyletic, with *P. germanicus* (thirteen cranio-dental

unambiguous autapomorphies; Table 6) as the first offshoot (node 18) and *P. laubei* and *P.*

douvillei being sister species (node 19). The monophyly of *Prosantorhinus* is supported by

seven cranio-dental unambiguous synapomorphies, some being optimised in *P. laubei* (no

cranial remains available; Heissig & Fejfar, 2007): lateral apophysis present on the nasals,

median nasal horn present (probably in males), presence of a sagittal fronto-parietal crest, of a

posterior groove on the processus zygomaticus of the squamosal, of a metacone fold on M1-2,

of an unconstricted metaloph on M1, and of an ectolophid fold on d2-3. *Prosantorhinus*

douvillei (nine unambiguous dental autapomorphies; Table 6) and *P. laubei* (six unambiguous

dental autapomorphies; Table 6) share six dental and postcranial unambiguous

synapomorphies (node 19; BS = 4): protocone unconstricted on P3-4 and M3, metaloph

unconstricted on M2, labial cingulum always present on lower molars, lingual groove always

present on d3, and expansion of the calcaneus-facet 1 always high and narrow on the

astragalus.

Node 20 (BS = 3) gathers eight terminal taxa (Fig. 13). It is supported by ten cranio-

dental and postcranial synapomorphies: nuchal tubercle small, articular tubercle smooth on

the squamosal, cement present on cheek teeth, protocone always constricted on P3-4, labial

cingulum usually absent on lower premolars and always present on lower molars, foramen

vertebrale lateralis present and axis-facets transversally concave on the atlas, postero-distal

apophysis low on the tibia, and latero-distal gutter located posteriorly on the fibula. Two

clades diverge from node 20. The first one (node 21, BS = 3) gathers *Brachydiceratherium*

shanwangense, *Bd*. *aginense*, and *Bd. intermedium*, based on eight dental and postcranial

synapomorphies: I1 with an oval occlusal outline, labial cingulum always absent on upper

premolars, crista usually present on P3, scapula elongated, fossa olecrani high on the

humerus, fovea capitis low and wide on the femur, latero-distal gutter deep on the fibula,

limbs slender, and insertions for the m. interossei long on lateral metapodials. Most of them

are optimised in *Bd. intermedium. Brachydiceratherium shanwangense* is well diagnosed,

with sixteen cranio-dental and postcranial unambiguous synapomorphies: lateral apophysis

present on the nasals, median nasal horn present on the nasals, premolar series short with

respect to the molar series, roots distinct on the cheek teeth, crochet always simple and lingual

cingulum usually absent and always reduced on P2-4, crista always present on P3, protocone

strongly constricted on M1-2, lingual cingulum usually absent on lower premolars and always

absent on lower molars, d1/p1 absent in adults, glenoid fossa with a medial border straight on

the scapula, distal gutter absent on the lateral epicondyle of the humerus, proximal radius-ulna

facets always fused, and trochanter major low on the femur. Node 22 (BS = 3) is supported by

five dental and postcranial unambiguous synapomorphies: metaloph short on M1-2, posterior

height exceeding the anterior height on the scaphoid, astragalus almost as high as wide (TD/H

ratio between 1 and 1.2), and tibia- and fibula-facets absent on the calcaneus.

Brachydiceratherium intermedium (five dental and postcranial unambiguous autapomorphies;

Table 6) is less derived than *Bd. aginense* (16 dental and postcranial unambiguous

autapomorphies; Table 6), which probably reflects the strong contrast in the completeness of

their hypodigms (e.g., no indisputable cranial remains are documented for *Bd. intermedium*).

The second clade diverging from the node 20 (i.e., node 23) places *Bd. fatehjangense* as

a sister taxon to (*Bd. aurelianense*, (*Bd. lamilloquense*, (*Bd. lemanense*, *Bd. asphaltense*))).

All the corresponding nodes are weakly supported ($1 \leq BS \leq 3$), with low numbers of

unambiguous synapomorphies (ranging from three to six). Node 23 is the least-supported one

(BS = 1), with five dental and postcranial synapomorphies (metacone fold present on M1-2,

second distal radius-ulna articulation present, posterior expansion of the scaphoid-facet low

on the radius, postero-proximal semilunate-facet usually absent on the scaphoid, and

expansion of the calcaneus-facet 1 usually wide and low on the astragalus).

DISCUSSION

Ontogenetic age and sex

Both the complete dental eruption and the wear stages of upper and lower teeth concur

to consider this individual as an adult, most likely ~7-15 years old (with reference to recent

rhinos; e.g., Hillman-Smith et al., 1986; Hullot et al., 2020). In the absence of I1s (usually

highly dimorphic in teleoceratines; see Antoine, 2002 regarding *Prosantorhinus douvillei*),

and due to the fragmentary state of i2s, it is not possible to determine its sex.

Taxonomic inferences

Surprisingly, *Diaceratherium tomerdingense* Dietrich, 1931 is retrieved as the first

offshoot among Teleoceratina (Fig. 13). Moreover, the assignment of this hornless and robust-

limbed rhinocerotine to the subtribe is not well supported at all (BS = 1): in other words, this

species could be closely related to Rhinocerotina instead among Rhinocerotini, as suggested

by some of its peculiar features, retrieved as autapomorphies in the current analysis (metaloph

short on M1-2; distal gutter on the lateral epicondyle absent on the humerus, distal border of

the anterior side of the semilunate rounded, and trapezium-facet absent on the McII).

Accordingly, and taking into account both the topology of the most parsimonious tree and the

character distribution along its branches, we propose that *Diaceratherium* Dietrich, 1931 shall

be restricted to the type species.

Indeed, all other species previously assigned to *Diaceratherium* in the last decades form

a well-supported clade remote from the type species (Fig. 13). This clade is split into two

sister clades encompassing three and five species, respectively (*D. shanwangense*, *D.*

aginense, and *D. intermedium*; *D. fatehjangense*, *D. aurelianense*, *D. lamilloquense*, *D.*

asphaltense, and *D. lemanense*). Except for *D. lamilloquense* Michel, 1987, these species

were originally or subsequently assigned to pre-existing genera, i) either unambiguously non-

related to Teleoceratina, such as *Aceratherium* (*D. lemanense*), *Diceratherium* (*D.*

asphaltense, *D. lemanense*), *Aprotodon* (*D. fatehjangense*), *Chilotherium* or *Subchilotherium*

(*D. intermedium*), and *Plesiaceratherium* (*D. shanwangense*), or ii) among Teleoceratina,

with *Teleoceras* and/or *Brachypotherium* (*D. aginense*, *D. aurelianense*, *D. shanwangense*,

and *D. fatehjangense*). Finally, and to our knowledge, the only species belonging to this clade

for which a genus-group name has been unambiguously proposed is *D. lemanense*. Indeed,

Lavocat (1951) has erected the subgenus *Brachydiceratherium* for "*Acerotherium lemanense*

Pomel, 1853". Interestingly, Lavocat did assign these species and subgenus to *Diceratherium*

Marsh, 1875, a genus consistently assigned to Elasmotheriinae in the last decades (e.g.,

Antoine, 2002). We propose that all these eight species be assigned to *Brachydiceratherium*

Lavocat, 1951, especially as the five-species clade, with *D. fatehjangense*, *D. aurelianense*,

D. lamilloquense, *D. asphaltense*, and *D. lemanense*, is not well supported (BS = 1; 5

unambiguous synapomorphies). Noteworthily, *D. asphaltense* and *D. lemanense* are sister

species in the most parsimonious tree, with a low number of morpho-anatomical

discrepancies. It should be noted that Jame et al. (2019) consider both species as being

distinct, based on a wide array of cranio-dental and postcranial features.

Other teleoceratine genera are monophyletic in the present analysis. *Brachypotherium*

Roger, 1904 includes *B. brachypus* and *B. perimense* and this genus is a sister group to a

clade gathering *Teleoceras* Hatcher, 1894 plus *Prosantorhinus* Heissig, 1974 on one branch

and *Brachydiceratherium* on the other one (see above).

Historical biogeography of Eurasian teleoceratines

During early Miocene times, Teleoceratina were particularly species-rich in Eurasia,

with 5–8 coeval species in any time slices (Fig. 14). A common thread between

Brachypotherium, *Brachydiceratherium*, and *Prosantorhinus* is their huge geographical range

at the generic level, encompassing most of the Eurasian landmasses for the latter two genera

(e.g., Heissig, 1999; Antoine et al., 2010, 2013), plus Afro-Arabia for *Brachypotherium* (e.g.,

Hooijer, 1963, Geraads & Miller, 2013; Pandolfi & Rook, 2019). An early representative of

Brachydiceratherium has been recognised in Thailand (*Bd.* cf. *lamilloquense*; Marivaux et al.,

2004). It has the closest affinities with *Bd. lamilloquense*, from the late Oligocene of Western

Europe (Fig. 15). To our knowledge, no occurrence has been reported between both areas for

this species. *Prosantorhinus* has a similar geographical range, extending from Western

Europe (*P. germanicus* and *P. douvillei*; Heissig, 1972b; Antoine et al., 2000; Heissig, 2017)

and Central Europe (*P. laubei*; Heissig & Fejfar, 2007) to Southern Pakistan (*P. shahbazi*;

Antoine et al., 2010, 2013). If confirmed, the recognition of *Bd. fatehjangense* in lower

Miocene beds of the Turgai region in Kazakhstan, previously described as a representative of

Bd. aurelianense by Borissiak (1927) and Lu et al. (2021), would considerably expand

latitudinally the range of this species, previously restricted to the Indian Subcontinent. It

would then be documented on both sides of the Himalayas (Fig. 15). The ubiquitous

distributions of most teleoceratine taxa likely underline ultra-generalist ecological preferences

(Hullot et al., 2021). Moreover, such ranges seemingly support the absence of efficient

ecological and geographical barriers at the Eurasian scale for the concerned teleoceratines, at

least by early Miocene times (Fig. 15).

Moreover, ghost lineages within *Brachypotherium* and *Prosantorhinus* (Fig. 14) are

likely to be bridged by *B. gajense* and *P. shahbazi*, from the latest Oligocene–earliest

Miocene and the early Miocene of Pakistan, respectively (for further discussion, see Antoine

et al., 2013 and Antoine, in press).

Brachydiceratherium shanwangense was previously only documented at Shanwang,

eastern China (N32°, E116.5°). The well-supported specific assignment of the Tagay

rhinoceros (N53°, E107.5°) points to an unsuspectedly wide geographical range for this

species, further pleading for both a low climatic and environmental gradient in the concerned

area at that time and very broad ecological preferences for this species (Fig. 15). Moreover, it

can be suspected that the smallest teleoceratine remains described over the early Miocene

interval in Japan (Kani and Mizunami formations) and referred to the *Brachypotherium*

pugnator (Matsumoto, 1921), otherwise of gigantic dimensions (Fukuchi & Kawai, 2011;

Tomida et al., 2013; Handa, 2020), may have particularly close affinities with those of *Bd.*

shanwangense. More generally, the concerned Japanese assemblages are very similar to the

Tagay and Shanwang ones (e.g., with the equid *Anchitherium* cf. *gobiense*, the proboscidean

Gomphotherium annectens, and the beaver *Youngofiber sinensis*; Qiu & Qiu, 2013), thus

strengthening the existence of a single eastern Asian biogeographical province at mid

latitudes at that time (Fig. 15). Indeed, closed forest environments under a subtropical climate,

with precipitation averaging ca. 1500 mm per year, are reported for the Shanwang Basin

based on early Miocene floras and vertebrates (Lu et al., 2021). The same proxies allow for

considering the Tagay area as a lake, also surrounded by dense forests under subtropical

conditions, with precipitation averaging ca. 1000-1500 mm per year (Logachev et al, 1964;

Belova, 1985; Sizov & Klementiev, 2015).

CONCLUSIONS

The numerous associated features documented and scored in the Tagay rhinocerotid

skeleton have allowed for assigning it to the same teleoceratine species (*Brachydiceratherium*

shanwangense) as in Shanwang, eastern China. These remains further contribute to a refined

depiction of phylogenetic relationships and to a revision of generic assignments among

Eurasian Teleoceratina.

The genus *Diaceratherium* Dietrich, 1931 should be restricted to the type species,

Diaceratherium tomerdingense Dietrich, 1931. This monotypic genus is the first offshoot

within Teleoceratina. Our results support the reappraisal of *Brachydiceratherium* Lavocat,

1951, with eight assigned species: *Brachydiceratherium lemanense* (Pomel, 1853),

Brachydiceratherium aurelianense (Nouel, 1866), *Brachydiceratherium intermedium*

(Lydekker, 1884), *Brachydiceratherium asphaltense* (Depéret & Douxami, 1902),

Brachydiceratherium fatehjangense (Pilgrim, 1910), *Brachydiceratherium aginense* (Répelin,

1917), *Brachydiceratherium shanwangense* (Wang, 1965) and *Brachydiceratherium*

lamilloquense Michel, 1983. *Brachydiceratherium* is a sister group to a clade encompassing

Prosantorhinus and the North American genus *Teleoceras*. *Brachypotherium* is more closely

related to the latter three genera than to *Diaceratherium*.

All Old World teleoceratines have extended geographical distributions at the genus level, which is also true for some species, such as the late Oligocene *Brachydiceratherium lamilloquense* and the early Miocene *Brachydiceratherium shanwangense*. The latter range supports the existence of a single eastern Asian biogeographical province at mid latitudes at that time for such megaherbivores.

ACKNOWLEDGMENTS

We are grateful to our colleagues for having participated in the 2008-2021 excavations. Special thanks to Gennady Turkin for logistical support and for assistance in the field. Valeria Burova and Ekaterina Nikulina are acknowledged for working on the skeleton reconstruction. Géraldine Véron and Christine Argot (Muséum National d'Histoire Naturelle (Paris, France) kindly provided access to zoological and paleontological collections under their care. We deeply acknowledge the Recommender, Faysal Bibi, and all three reviewers (Jérémy Tissier, Deng Tao, and Panagiotis Kampouridis), for their thorough revisions and constructive remarks on a widely-improvable previous version of the manuscript.

FUNDING

The work related to the study of the geological structure of the section was supported by the Russian Foundation for Basic Research, project numbers 22-17-00049 «Neotectonics and active tectonics of the northern part of Central Asia».

REFERENCES

Antoine P-O. 2002. Phylogénie et évolution des Elasmotheriina (Mammalia, Rhinocerotidae). *Mémoires du Muséum National d'Histoire Naturelle* **188**: 1–359.

Antoine P-O. in press. Rhinocerotids from the Siwalik faunal sequence. In: Badgley, C.,

Pilbeam, D. & Morgan, M. (Eds.), *At the Foot of the Himalayas: Paleontology and*

Ecosystem Dynamics of the Siwalik Record of Pakistan. Johns Hopkins University Press.

Antoine P-O, Welcomme J-L. 2000. A new rhinoceros from the Bugti Hills, Baluchistan,

Pakistan: the earliest elasmotheriine. *Palaeontology* **43**: 795–816.

Antoine P-O, Fleury G, Duranthon F. 2002. Le rhinocérotidé *Prosantorhinus douvillei*

(Osborn, 1900) de l'Orléanien supérieur de Captieux (Gironde). *Bulletin de la Société*

d'Histoire Naturelle de Toulouse **137**: 87–91.

Antoine P-O, Duranthon F, Welcomme J-L. 2003. *Alicornops* (Mammalia, Rhinocerotidae)

dans le Miocène supérieur des Collines Bugti (Balouchistan, Pakistan): implications

phylogénétiques. *Geodiversitas* **25**: 575–603.

Antoine P-O, Ducrocq S, Marivaux L, Chaimanee Y, Crochet J-Y, Jaeger J-J,

Welcomme J-L. 2003. Early rhinocerotids (Mammalia: Perissodactyla) from South Asia

and a review of the Holartic Paleogene rhinocerotid record. *Canadian Journal of Earth*

Sciences **40**: 365–374.

Antoine P-O, Downing KF, Crochet J-Y, Duranthon F, Flynn LJ, Marivaux L, Métais

G, Rajpar AR, Roohi G. 2010. A revision of *Aceratherium blanfordi* Lydekker, 1884

(Mammalia: Rhinocerotidae) from the Early Miocene of Pakistan: postcranials as a key.

Zoological Journal of the Linnean Society **160**: 139–194.

Antoine P-O, Becker D. 2013. A brief review of Agenian rhinocerotids in Western Europe.

Swiss Journal of Geosciences **106**: 135–146.

Antoine P-O, Becker D, Laurent Y, Duranthon F. 2018. The Early Miocene teleoceratine

Prosantorhinus aff. *douvillei* (Mammalia, Perissodactyla, Rhinocerotidae) from Béon 2,

Southwestern France. *Revue de Paléobiologie* **37**: 367–377.

Antoine P-O, Becker D, Pandolfi L, Geraads D. accepted. Evolution and fossil record of

Old World Rhinocerotidae. In: Melletti M, Balfour D, Talukdar B. (Eds.), *Rhinos of the*

World: Ecology, Conservation and Management. Fascinating Life Sciences, Springer

Nature.

Antoine P-O, Métais G, Orliac MJ, Crochet J-Y, Flynn LJ, Marivaux L, Rajpar AR,

Roohi G., Welcomme J-L. 2013. Mammalian Neogene biostratigraphy of the Sulaiman

Province, Pakistan. In: Wang X-m, Flynn LJ, Fortelius M (eds.), *Fossil mammals of Asia:*

Neogene Biostratigraphy and Chronology, Columbia University Press 400–422.

Antoine P-O, Reyes MC, Amano N, Bautista AP, Chang CH, Claude J, Vos JD, Ingicco

T. 2022. A new rhinoceros clade from the Pleistocene of Asia sheds light on mammal

dispersals to the Philippines. *Zoological Journal of the Linnean Society* **194**: 416–430.

DOI: 10.1093/zoolinnean/zlab009

Becker D, Antoine P-O, Maridet O. 2013. A new genus of Rhinocerotidae (Mammalia,

Perissodactyla) from the Oligocene of Europe. *Journal of Systematic Palaeontolo*gy **11**:

947-972.

Belova VA. 1985. *Vegetation and climate of the Late Cenozoic of the south of Eastern*

Siberia. Novosibirsk: Nauka [in Russian].

Boada-Saña A. 2008. *Phylogénie du rhinocérotidé* Diaceratherium Dietrich, 1931

(Mammalia, Perissodactyla). Master thesis dissertation. University of Montpellier,

France, 2.

Boivin M, Marivaux L, Antoine P-O. 2019. L'apport du registre paléogène d'Amazonie sur

la diversification initiale des Caviomorpha (Hystricognathi, Rodentia) : implications

phylogénétiques, macroévolutives et paléobiogéographiques. *Geodiversitas* **41**: 143–245.

Borissiak A. 1914. On the dental apparatus of *Elasmotherium caucasicum* n. sp. *Bulletin de*

l'Académie Impériale des Sciences de St-Pétersbourg **6**: 555–584.

Borissiak A. 1927. *Brachypotherium aurelianense* Nouel, var. nov. Gailiti, from the Miocene

deposits of the Turgai region. *Bulletin de l'Académie des Sciences de l'URSS* **21**: 273–

286.

Bremer KR. 1994. Branch support and tree stability. *Cladistics* **10**: 295–304.

Burchell WJ. 1817. Note sur une nouvelle espéce de Rhinoceros. *Bulletin de la Société*

Philomathique de Paris (June).

Cerdeño E. 1993. Étude sur *Diaceratherium aurelianense* et *Brachypotherium brachypus*

(Rhinocerotidae, Mammalia) du Miocène moyen de France. *Bulletin du Muséum national*

d'Histoire naturelle **15**: 25–77.

Cerdeño E. 1995. Cladistic analysis of the Family Rhinocerotidae (Perissodactyla). *American*

Museum Novitates **3143**: 1–25.

Crusafont M, Villalta JF, Truyols J. 1955. El Burdigaliense continental de la Cuenca del

Vallés-Penedés. *Memorias y Comunicaciones del Instituto Geológico, Barcelona* **12**: 1–

272.

Danilov IG, Syromyatnikova EV, Klementiev AM, Sizov AV, Martynovich NV,

Zelenkov NV, Sychevskaya EK, Tesakov AS. 2012. New data on Miocene vertebrates

of Tagay locality (Olkhon, Lake Baikal). In: Lopatin AV, Parkhaev PYu, Rozanov AYu

eds. *Modern Paleontology: classical and new methods. The ninth all-Russian scientific*

school for young scientists in Paleontology. Moscow: Paleontological Institute 19–20 [in

Russian].

Daxner-Höck G, Böhme M, Kossler A. 2013. New Data on Miocene Biostratigraphy and

Paleoclimatology of Olkhon Island (Lake Baikal, Siberia). In: Wang X, Flynn LJ,

Fortelius M, eds., *Fossil mammals of Asia: Neogene biostratigraphy and chronology*,

New York: Columbia University Press, 508-517.

Daxner-Höck G, Mörs T, Kazansky AY, Matasova GG, Ivanova VV, Shchetnikov AA, Filinov IA, Voyta L, Erbajeva MA, 2022. A synthesis of fauna, palaeoenvironments and stratigraphy of the Miocene Tagay locality (Olkhon Island, Lake Baikal, Eastern Siberia). *Palaeobio Palaeoenv* **102**, 969–983.

Deng T, Gao F. 2006. Perissodactyla. In: Qi GQ, Dong W (Eds). Lufengpithecus hudienensis *site*. Beijing: Science Press, 188–195 (in Chinese), 334–335 (in English).

Depéret C, Douxami H. 1902. Les Vertébrés oligocènes de Pyrimont-Challonges (Savoie). *Mémoires suisses de Paléontologie* **29**: 1–92.

Dietrich WO. 1931. Neue Nashornreste aus Schwaben (*Diaceratherium tomerdingensis* n. g. n. sp.). *Zeitschrift für Säugetierkunde* **6**: 201–223.

Falconer H, Cautley PT. 1847. *Fauna antiqua Sivalensis, being the fossil zoology of the Sewalik Hills, in the North of India (Sus, Rhinoceros, Chalicotherium).* London: Smith, Elder and Co..

Filippov AG, Sytchevskaya EK. 2000. Remains of snakeheaded fishes (Channidae) near Lake Baikal. *The Third Vereshchagin Baikal Conference, Aug 22-27, Irkutsk*, 251.

Fischer von Waldheim GF. 1814. *Zoögnosia tabulis synopticis illustrate, in usum Paeselectionum Academiae Imperialis Medicochirurgae*. Moscow: Nicolai Sergeidis Vsevolozsky.

Fukuchi A, Kawai K. 2011. Revision of fossil rhinoceroses from the Miocene Mizunami Group, Japan. *Palaeontological Research* **15**: 247–257.

Geraads D. 2010. Rhinocerotidae. In: Werdelin L, Sanders WJ (Eds.), *Cenozoic Mammals of Africa*. University of California Press, Berkeley, 675–689.

Geraads D, Miller E. 2013. *Brachypotherium minor* n. sp., and other Rhinocerotidae from the Early Miocene of Buluk. Northern Kenya. *Geodiversitas* **35**: 359–375. DOI: 10.5252/g2013n2a5.

Guérin C. 1980. Les rhinocéros (Mammalia, Perissodactyla) du Miocène terminal au

Pléistocène supérieur en Europe occidentale. Comparaison avec les espèces actuelles.

Documents des Laboratoires de Géologie de Lyon **79**: 1–1185.

Handa N. 2020. Reappraisal of a rhinocerotid (Mammalia, Perissodactyla) from the lower

Miocene Yotsuyaku Formation, Northeast Japan, with an overview of the early Miocene

Japanese rhinocerotids. *Paleontological Research* **24**: 183–191.

Heissig K. 1969. Die Rhinocerotidae (Mammalia) aus der oberoligozänen Spaltenfüllung von

Gaimersheim bei Ingolstadt in Bayern und ihre phylogenetische Stellung. *Verlag der*

Bayerischen Akademie der Wissenschaften **138**: 1–133.

Heissig K. 1972a. Paläontologische und geologische Untersuchungen im Tertiär von

Pakistan. 5 – Rhinocerotidae (Mamm.) aus den unteren und mittleren Siwalik-schichten.

Abhandlungen der Bayerischen Akademie der Wissenschaften, München, mathematische-

naturwissenschaftliche Klasse **152**: 1–112.

Heissig K. 1972b. Die obermiozäne Fossil-Lagerstätte Sandelzhausen. 5. Rhinocerotidae

(Mammalia), Systematik und Ökologie. *Mitteilungen der Bayerischen Staatssammlung*

für Paläontologie und historische Geologie **12**: 57–81.

Heissig K. 1999. Family Rhinocerotidae. In: Röessner GE, Heissig K. eds. *The Miocene Land*

Mammals of Europe. Dr Pfeil, Munich 175–188.

Heissig K, Fejfar O. 2007. Die fossilen Nashörner (Mammalia, Rhinocerotidae) aus dem

Untermiozän von Tuchořice in Nordwestboehmen. *Acta Musei Nationalis Pragae B* **63**:

19-64.

Heissig K. 2012. Les Rhinocerotidae (Perissodactyla) de Sansan. *Mémoires du Muséum*

national d'Histoire naturelle de Paris **203**: 317–485.

Heissig K. 2017. Revision of the European species of *Prosantorhinus* Heissig, 1974

(Mammalia, Perissodactyla, Rhinocerotidae). *Fossil Imprint* **122**: 265–294. DOI :

10.1515/if-2017-0014.

Hilgen FJ, Lourense LJ, Van Dam JA. 2012. The Neogene period. In: Gradstein FM, Ogg

JG, Schmitz MD, Ogg GM, eds. *The Geologic Time Scale 2012*. Oxford: Elsevier 923–

978.

Hooijer DA. 1963. Miocene Mammalia of the Congo. *Musée royal de l'Afrique Centrale.*

Annales - sciences géologiques **46**: 1–77.

Hullot M, Laurent Y, Merceron G, Antoine P-O. 2021. Paleoecology of the

Rhinocerotidae (Mammalia, Perissodactyla) from Béon 1, Montréal-du-Gers (late early

Miocene, SW France): Insights from dental microwear texture analysis, mesowear, and

enamel hypoplasia. *Palaeontologia Electronica* **24**: a27. DOI: 10.26879/1163

Jame C, Tissier J, Maridet O, Becker D 2019. Early Agenian rhinocerotids from Wischberg

(Canton Bern, Switzerland) and clarification of the systematics of the genus

Diaceratherium. *PeerJ* **7**: e7517**.**

Kitainik AF, Ivaniev LN. 1958. A note on the Tertiary deposits of Olkhon Island on Lake

Baikal. *Notes of the Irkutsk Regional Museum of Local History* 55–60.

Klementiev AM. 2009. Finding a Miocene rhinoceros on Olkhon Island (Lake Baikal). In:

Barskov IS & Nazarova VM eds. *200 years of national paleontology. Materials of the*

All-Russian meeting. Moscow: Paleontological Institute, 56–57 [in Russian].

Klementiev AM, Sizov AV. 2015. New record of anchithere (*Anchitherium aurelianense*) in

the Miocene of Eastern Siberia, Russia. *Russian Journal of Theriology* **14**: 133-143.

Kossler A. 2003. Neogene sediments of Olkhon and Svyatoy Nos (Baikal Rift System, East

Siberia): Suggestions about the development of Lake Baikal. *Berliner Paläobiologische*

Abhandlungen **4**: 55–63.

Khan MA, Akhtar M, Khan AM, Ghaffar A, Iqbal M, Samiullah K. 2011. New fossil

locality in the Middle Miocene of Lava from the Chinji Formation of the Lower Siwaliks,

Pakistan. *Pakistan Journal of Zoology* **43**: 61–72

Khosatzky LI, Chkhikvadze VM. 1993. New data about Miocene turtles of the genus

Baicalemys. *Bulletin of the Academy of Science of Georgia* **148**: 155–160 [in Russian].

von Koenigswald GHR. 1956. Fossil Mammals from the Philippines. *National Research

Council of the Philippines, University of the Philippines Diliman. Special reprint. Full

text illustrations of paper 22 Proceedings of the Fourth Far-Eastern Prehistory

Congress*: 1–14.

Kordikova EG. 2001. Remarks on the Oligocene-Miocene mammal paleontology and

sequence stratigraphy of South-Western Betpakdala Steppe, South Kazakhstan. *Neues

Jahrbuch für Geologie und Paläontologie Abhandlungen* **221**: 35–79.

Lartet E. 1851. *Notice sur la colline de Sansan*. Auch: Portes.

Laurillard F. 1848. Rhinocéros fossiles, in d'Orbigny CD (ed.), *Dictionnaire universel

d'Histoire naturelle*, volume 11. Renard, Martinet & Cie, Paris: 99–102.

Linnaeus C. 1758. *Systema Naturae per regna tria naturae, secundum classes, ordines,

genera, species, cum characteribus, differentiis, synonymis, locis.* Vol. 1. Tenth Edition.

Stockholm: Regnum animale.

Logachev NA, Lomonosova TK, Klimanova VM. 1964. *The Cenozoic deposits of the

Irkutsk amphitheatre*. Moscow: Nauka [in Russian].

Lavocat R. 1951. *Révision de la faune des mammifères oligocènes d'Auvergne et du Velay*.

Sciences et Avenir, Paris.

Lu X, Cerdeño E, Zheng X, Wang S, Deng T. 2021. The first Asian skeleton of

Diaceratherium from the early Miocene Shanwang Basin (Shandong, China), and

implications for its migration route. *Journal of Asian Earth Sciences: X* **6**: 100074

Lydekker R. 1884. Additional Siwalik Perissodactyla and Proboscidea. *Memoirs of the*

Geologcal Survey of India – Palaeontologia Indica **3**: 1–34.

MacFadden BJ. 1998. Equidae, In: Janis CM, Scott KM, Louis LL eds. *Evolution of Tertiary*

Mammals North America. vol. 1. Cambridge University Press 537–559.

Marivaux L, Chaimanee Y, Yamee C, Srisuk P, Jaeger J-J. 2004. Discovery of *Fallomus*

ladakhensis Nanda & Sahni, 1998 (Mammalia, Rodentia, Diatomyidae) in the lignites of

Nong Ya Plong (Phetchaburi Province, Thailand): systematic, biochronological and

paleoenvironmental implications. *Geodiversitas*, **26**: 493-507.

Mats VD, Ufimtsev GF, Mandelbaum MM., Alakshin AM, Pospeev AV, Shimaraev MN,

Khlystov OM. 2001. *Cenozoic of the Baikal rift zone*. Novosibirsk: GEO SO RAS Press

[in Russian].

Mats VD, Lomonosova TK, Vorobyeva GA, Vologina EG. 2010. Late Cretaceous-

Cenozoic sediments of the Baikal rift basin and changing natural conditions.

Geodynamics & Tectonophysics **1**: 75–86 [in Russian].

Mats VD, Yefimova IM. 2011. Paleogeographic scenario of the Late Cretaceous – Cenozoic

for the central part of the Baikal region. *Geodynamics & Tectonophysics* **2**: 175–193 [in

Russian].

Mats VD. 2013. Late Cretaceous and Cenozoic Stratigraphy of the Baikal Rift Sediments.

Stratigraphy and Geological Correlation **21**: 637–651. DOI:

10.1134/S0869593813060075

Mats VD. 2015. The Baikal rift: Pliocene (Miocene) – Quaternary episode or product of

extended development since the Late Cretaceous under various tectonic factors. A

review. *Geodynamics & Tectonophysics* **6**: 467–489 [in Russian]. DOI: 10.5800/GT-

2015-6-4-0190

Matsumoto H. 1921. Descriptions of some new fossil mammals from Kani District, Province of Mino, with revisions of some Asiatic fossil rhinocerotids. *The Science Reports of the Tohoku Imperial University. Second Series (Geology)* **5**: 75–91.

Ménouret B, Guérin C. 2009. *Diaceratherium massiliae* nov. sp. from the Oligocene clays of Saint-André and Saint-Henri in Marseille and Les Milles near Aix-en-Provence (South Eastern France), the first European large brachypod Rhinocerotidae. *Geobios* **42**: 293–327.

Nouel E. 1866. Mémoire sur un nouveau rhinocéros fossile. Mémoires de la Société d'Agriculture, Sciences, *Belle-Lettres et Art d'Orléans* **8**: 241–251.

Orliac MJ, Antoine P-O, Ducrocq S. 2010. Phylogenetic relationships of the Suidae (Mammalia, Cetartiodactyla): new insights on the relations within Suoidea. *Zoologica Scripta* **39**: 315–330.

Osborn HF. 1900. Phylogeny of the rhinoceroses of Europe. *Memoirs of the American Museum of Natural History* **13**: 229–267.

Pandolfi L, Rook L. 2019. The latest Miocene Rhinocerotidae from Sahabi (Libya). *Comptes Rendus Palevol* **18**: 442-448. DOI: 10.1016/j.crpv.2019.03.002.

Pandolfi L, Antoine P-O, Bukhsianidze M, Lordkipanidze D, Rook L. 2021. Northern Eurasian rhinocerotines (Mammalia, Perissodactyla) by the Pliocene–Pleistocene transition: phylogeny and historical biogeography. *Journal of Systematic Palaeontology.* DOI: 10.1080/14772019.2021.1995907

Pilgrim GE. 1910. Notice on new mammal genera and species from the Tertiaries of India. *Records of the Geological Survey of India* **15**: 63–71.

Pokatilov AG. 2004. *Paleontology and stratigraphy of the Cenozoic of the south of Eastern Siberia and adjacent territories*. Irkutsk: Irkutsk State Technical University Press [in Russian].

Pomel M. 1853. *Catalogue méthodologique et descriptif des vertébrés fossiles découverts*

dans le bassin hydrographique supérieur de la Loire, et surtout dans la vallée de son

affluent principal, l'Allier. Paris: Baillière Ed.

Prieto J., Antoine P-O, Böhme M, van der Made J, Métais G, Laq The Phuc, Quý

Trương Quan, Schneider S, Dang Ngoc Tran, Vasilyan D, Luong The Viet, 2018.

Biochronological and paleobiogeographical significance of the earliest Miocene mammal

fauna from Northern Vietnam. *Palaeobiodiversity and Palaeoenvironments* **98**: 287–313.

doi 10.1007/s12549-017-0295-y

Prothero DR, Guérin C, Manning E. 1989. The History of the Rhinocerotoidea. In:

Prothero DR & Schoch RM eds. *The Evolution of Perissodactyls*, New York: Oxford

University Press, 322–340.

Prothero DR. 2005. *The evolution of North American Rhinoceroses*. Cambridge; New York;

Melbourne: Cambridge University Press.

Qiu ZD, Qiu ZX. 2013. Early Miocene Xiejiahe and Sihong fossil localities and their faunas,

eastern China. In: Wang X-m, Flynn LJ, Fortelius M (eds.), *Fossil mammals of Asia:*

Neogene Biostratigraphy and Chronology, Columbia University Press 142–154.

Raffi I, Wade BS, Pälike H, Beu AG, Cooper R, Crundwell MP, Krijgsman W, Moore T,

Raine I, Sardella R, Vernyhorova YV. 2020. Chapter 29 - The Neogene Period. In:

Gradstein FM, Ogg JG, Schmitz MD, Ogg GM ads. *Geologic Time Scale 2020*. Elsevier

1141–1215. DOI: /10.1016/B978-0-12-824360-2.00029-2.

Rage JC, Danilov IG. 2008. A new Miocene fauna of snakes from eastern Siberia, Russia.:

Was the snake fauna largely homogenous in Eurasia during the Miocene? *Comptes*

Rendus Palevol **7**: 383–390. DOI: 10.1016/j.crpv.2008.05.004.

Répelin J. 1917. Études paléontologiques dans le sud-ouest de la France (Mammifères). Les

rhinocérotidés de l'Aquitanien supérieur de l'Agenais (Laugnac). *Annales du Muséum*

d'Histoire naturelle de Marseille **16**: 1–47.

Rössner G.E., Mörs T. 2001. A New Record of the Enigmatic Eurasian Miocene Ruminant

Artiodactyl *Orygotherium*. *Journal of Vertebrate Paleontology* **21**: 591-595.

Sizov AV, Klementiev AM. 2015. Geology and taphonomy of Tagay locality of early

Miocene vertebrate fauna. In: Lipnina EA & Berdnikov IM, eds. *Eurasia in the Cenozoic.*

Stratigraphy, paleoecology, cultures. Irkutsk: Irkutsk State University Press, 206–218 [in

Russian].

Sotnikova MV, Klementiev AM, Sizov AV, Tesakov AS. 2021. New species of *Ballusia*

Ginsburg and Morales, 1998 (Ursidae, Carnivora) from Miocene of Eastern Siberia,

Russia. *Historical Biology* **33**: 486-497, DOI: 10.1080/08912963.2019.1637864

Swofford, D.L. 2002. *PAUP*: Phylogenetic analysis using parsimony (*and other methods).*

Version 4.0b10, Sunderland, Sinauer Associates, Inc., Publishers, Sunderland.

Syromyatnikova EV. 2014. The first record of *Salamandrella* (Caudata: Hynobiidae) from

the Neogene of Russia. *Russian Journal of Herpetology* **21**: 217–220.

Syromyatnikova EV. 2015. A New Species of *Bufo* (Amphibia, Anura) from the Miocene of

Russia. *Russian Journal of Herpetology* **22**: 281–288.

Tesakov AS, Lopatin AS. 2015. First record of Mylagaulid rodents (Rodentia, Mammalia)

from the Miocene of Eastern Siberia (Olkhon Island, Baikal Lake, Irkutsk Region,

Russia). *Doklady Biological Sciences* **460**: 23–26.

Tissier J, Antoine P-O, Becker D. 2021. New species, revision, and phylogeny of

Ronzotherium Aymard, 1854 (Perissodactyla, Rhinocerotidae). *European Journal of*

*Taxonom*y **753**: 1–80. DOI: 10.5852/ejt.2021.753.1389

Tissier J, Geiger-Schütz P, Flückiger PF, Becker D. 2021. Neue Erkenntnisse über die

Nashorn-Funde von Rickenbach (SO) (Oberes Oligozän, Kanton Solthurn, Schweiz) aus

der Sammlung des Naturmuseums Olten. *Naturforschende Gesellschaft des Kantons*

Solothurn **44**: 25–50

Tomida Y, Nakaya H, Saegusa H, Miyata K, Fukuchi A. 2013. Miocene Land Mammals

and Stratigraphy of Japan. In: Wang X, Flynn LJ, Fortelius M, eds., *Fossil mammals of*

Asia: Neogene biostratigraphy and chronology, New York: Columbia University Press,

314–333.

Vislobokova IA. 1990. About artiodactyls from the Lower Miocene of the Tagay bay,

Olkhon island (Baikal). *Paleontological Journal* **2**: 134–138 [in Russian].

Vislobokova IA. 1994. The Lower Miocene artiodactyls of Tagay Bay, Olkhon Island, Lake

Baikal (Russia). *Palaeovertebrata*. **23**: 177–197.

Vislobokova I. 2004. New species of *Orygotherium* (Palaeomerycidae, Ruminantia) from the

Early and Late Miocene of Eurasia. *Annalen des Naturhistorischen Museums in Wien*

106: 371–385.

Wang KM. 1929. Die obermiozänen Rhinocerotiden von Bayern. *Paläontologische*

Zeitschrift, **10**: 184-212.

Wang BY. 1965. A new Miocene aceratheriine rhinoceros of Shanwang, Shandong.

Vertebrata PalAsiatica **9**: 109–112.

Zelenkov NV. 2016. The first fossil parrot (Aves, Psittaciformes) from Siberia and its

implications for the historical biogeography of Psittaciformes. *Biology Letters* **12**:

20160717. DOI: 10.1098/rsbl.2016.0717

1340

SUPPORTING INFORMATION

Additional supporting information may be found online in the supporting information

tab for this article. Supplementary Files:

• **S1**. Character matrix for the preliminary phylogenetic analysis, including 282 cranial,

dental, and postcranial characters controlled on 32 terminal taxa (one tapirid,

rhinocerotoids, and rhinocerotids), with Tagay rhinoceros and "Diaceratherium

shanwangense" as separate terminals.

• **S2.** Output log text of the preliminary phylogenetic analysis (282 characters and 32

taxa).

• **S3**. Character matrix for the final phylogenetic analysis, including 282 cranial, dental,

and postcranial characters controlled on 31 terminal taxa (one tapirid, rhinocerotoids,

and rhinocerotids).

• **S4.** Output log text of the final phylogenetic analysis, with Bremer Support

• **S5** Measurements for *Brachydiceratherium shanwangense* from Tagay site.

1355

1356
1357

1358   FIGURES

1359

1360

1361   **Fig. 1.** Geographic position of Tagay locality on Olkhon Island, Baikal Region, Russia (A)

1362   and age of Tagay Formation according to various authors (B). [Full width suggested]

1363

1364   **Fig. 2.** Geological structure of the Tagay section (A), photo (B) and plan (C) of the

1365   excavations of the Miocene rhinocerotid at Tagay site in 2008 (Olkhon Island, Baikal Region,

1366   Russia). [Full width suggested]

1367

1368

1369 **Fig. 3.** *Brachydiceratherium shanwangense* (Wang, 1965) from Tagay, Baikal Region,

1370 Russia, late early Miocene. Photos in dorsal (A) and lateral views (B) of the skull and

1371 mandible IZK79-1-08C-1. C - Scientific drawing of the right lateral view of the skull (based

1372 on B). Striped areas are reconstructed. D – Tentative reconstruction of the head in lateral

1373 view, by one of us (AS). [One column width suggested]

1374

Fig. 4. *Brachydiceratherium shanwangense* (Wang, 1965) from Tagay, Baikal Region,

Russia, late early Miocene. Mandible and dental material. A, B – Mandible in right lateral (A)

and occlusal views (B); C, D – Right upper cheek teeth (D1–M3) in occlusal view:

photograph (C) and interpretative sketch (D); E, F – Right lower cheek teeth (p3–m3) in

occlusal view: photograph (E) and interpretative sketch (F); G, H – Dental terminology used

for rhinocerotid upper tooth (G) and lower tooth (H). [Full width suggested]

1381

1382

Fig. 5. *Brachydiceratherium shanwangense* (Wang, 1965) from Tagay, Baikal Region,

Russia, late early Miocene. A-D – Atlas in dorsal (A), ventral (B), cranial (C) and posterior

views (D); E-I – Axis in anterior (E), left lateral (F), posterior (G), dorsal (H), and ventral

views (I). [Full width suggested]

1387

Fig. 6. *Brachydiceratherium shanwangense* (Wang, 1965) from Tagay, Baikal Region, Russia, late early Miocene. Long bones of the right forelimb. A-E –humerus in posterior (A), medial (B), anterior (C), lateral (D), and proximal views (E); F-J – radius in anterior (F), lateral (G), posterior (H), medial (I), and proximal views (J); K-O – ulna in proximal (K), medial (L), anterior (M), lateral (N), and posterior views (O). [Full width suggested]

1393

Fig. 7. *Brachydiceratherium shanwangense* (Wang, 1965) from Tagay, Baikal Region,

Russia, late early Miocene. Carpal bones. A-C – left scaphoid in posterior (A), proximal (B),

and distal views (C); D-E – right trapezoid in anterior (D), and distal views (E); F-I – left

semilunate in distal (F), proximal (G), medial (H), and lateral views (I); J-M – left pyramidal

in anterior (J), lateral (K), posterior (L), and medial views (M); N-P – right pisiform in

anterior (N), lateral (O), and proximal views (P); Q-T – right magnum in lateral (Q), medial

(R), distal (S), and anterior views (T); Y-W – right unciform in anterior (Y), proximal (V),

and distal (W). [Full width suggested]

Fig. 8. *Brachydiceratherium shanwangense* (Wang, 1965) from Tagay, Baikal Region, Russia, late early Miocene. Left metacarpal bones. A-E – second metacarpal in lateral (A), posterior (B), medial (C), anterior (D), and proximal views (E); F-J – third metacarpal in proximal (F), lateral (G), posterior (H), medial (I), and anterior views (J); K-O – fourth metacarpal in proximal (K), lateral (L), posterior (M), medial (N), and anterior views (O).

[Full width suggested]

1411

1412   **Fig. 9.** *Brachydiceratherium shanwangense* (Wang, 1965) from Tagay, Baikal Region,

1413   Russia, late early Miocene. Long bones of the left hind limb. A-F – femur in anterior (A),

1414   medial (B), posterior (C), lateral (D), proximal (E), and distal views (F); G-J – fibula in lateral

1415   (G), posterior (H), medial (I), and anterior views (J); K-O – tibia in anterior (K), medial (L),

1416   posterior (M), lateral (N), and proximal views (O). [Full width suggested]

1417

Fig. 10. *Brachydiceratherium shanwangense* (Wang, 1965) from Tagay, Baikal Region,

Russia, late early Miocene. Tarsal bones. A-C – left astragalus in anterior (A), distal (B), and

posterior views (C); D-F – left calcaneus in proximal (D), medial (E), and anterion views (F);

G-I – left cuboid in distal (G), lateral (H), and proximal views (I). [Full width suggested]

1422

1423

1424

1425

Fig. 11. *Brachydiceratherium shanwangense* (Wang, 1965) from Tagay, Baikal Region,

Russia, late early Miocene. Metatarsal bones. A-E – left second metatarsal in lateral (A),

posterior (B), medial (C), anterior (D), and proximal views (E); F-J – right fourth metatarsal

in proximal (F), medial (G), posterior (H), lateral (I), and anterior views (J). [Full width

suggested]

1431

1432

1433

1434

1435

1436

Fig. 12. Skulls of different species of *Brachydiceratherium* in right lateral view. **A -**

Brachydiceratherium shanwangense from Tagay (Baikal Region, Russia, late early Miocene)

№IZK79-1-08C-1/1; **B** - *Brachydiceratherium shanwangense* from Jijiazhuang locality STM

44–98 (deformed, mirrored) (MN4 - early Miocene, Shanwang Basin, Shandong Province,

China) №MHNT.PAL.2013.0.1001; **C** - *Brachydiceratherium aginense* (Répelin, 1917) from

Laugnac (MN2 - early Miocene, Lot-et-Garonne, France); **D** - *Brachydiceratherium*

lemanense from Gannat (MN1 - early Miocene, France) №MNHN-AC-2375, holotype; **E** -

Brachydiceratherium asphaltense (Depéret et Douxami, 1902) from Saulcet (MN1 - earliest

Miocene, Allier, France). №NMB–Sau1662; **F** - *Brachydiceratherium aurelianense* from

Neuville-aux-Bois (MN3 - early Miocene, France) №MHNT.PAL.2013.0.1001, cast of the

holotype; [Full width suggested]

1452 — 10 changes

Fig. 13. Phylogram of Rhinocerotinae, with a focus on Teleoceratina. Most parsimonious tree

(1316 steps; consistency index = 0.2698; retention index = 0.4918), retrieved from 282

unweighted cranio-mandibular, dental, and postcranial characters scored in 31 tapirid and

rhinocerotoid species (see S3 and S4). Node numbers appear in empty circles. Number of

unambiguous synapomorphies/Bremer Support are indicated left to nodes. [Full width

suggested]

1459

Fig. 14. Phylogenetic relationships of Teleoceratina versus time (see Fig. 13), with new

combinations. Although they were not included in the current parsimony analysis, the

temporal distributions of *Brachypotherium gajense* and *Prosantorhinus shahbazi* are provided

here, as these species might bridge a stratigraphic gap for the concerned genera. Red dotted

line for *B. shanwangense* stands for the age uncertainty of Tagay locality (MN3-5) [Full width

suggested]

1466

Fig. 15. Paleomap of Eurasia by early Miocene times (~20 Ma), showing the main

occurrences of representatives of the teleoceratine rhinocerotid *Brachydiceratherium*, at the

basin scale (apart from Tagay, Shanwang, and Nong Ya Plong localities). The Green area

depicts the interpolated geographical range of *B. shanwangense* (with possible occurrences on

Honshu Island, Japan). Based on data from Borissiak (1927), Cerdeño (1993), Antoine et al.

(2000, 2013), Becker et al. (2009), Antoine & Becker (2013), Tomida et al. (2013), Jame et

al. (2019), Handa (2020), Lu et al. (2021), Antoine (in press), and the present work.

PalaeoAtlas by Scotese (2016, under cc 4.0 license) with added paleomap for the Baikal area

(Mats et al., 2011). [Full width suggested]

Table 1. Cranial measurements of *Brachydiceratherium shanwangense*, from Tagay, early
Miocene of Eastern Siberia, in mm. 1, Length (occipital-premaxilla distance); 2, Length
(occipital-nasal distance); 3, Upper length (nasal-occipital crest distance); 4, Nasal incisure
length; 5, Minimal width; 6, Occipital crest-postorbital process distance; 8, Occipital crest-
lacrimal process distance; 9, Nasal incisure-orbit distance; 13, Post-M3-condyle distance; 14,
Nasal-orbit distance; 15, Occipital crest width; 16, Mastoid apophyses width; 17, Inter
frontoparietal crest minimal distance; 18, Postorbital process width; 19, Lacrimal process
width; 21, Zygomatic width; 22, Nasal incisure width; 23, Occipital height; 25, P2-level
height; 26, P4-M1-level height; 27, M3-level height; 31, Foramen magnum width; 32, Inter-
occipital condyle width. Numbers coincide with measurements as defined and illustrated by
Guérin (1980, fig. 1, table 1).

1493

1	2	3	4	5	6	8	9	13	14	15	16
505	540	455.8	(174)	125	249.6	293.6	(72.6)	(254)	219	(160)	224.8

17	18	19	21	22	23	25	26	27	31	32
26.7	179	189	308.2	69.7	144.6	(140)	(155)	(156)	43	112.4

1500
1501

Table 2. Right mandibular measurements of *Brachydiceratherium shanwangense*, from
Tagay, early Miocene of Eastern Siberia, in mm. 1, Maximal length; 2, Length without the
symphysis; 3–8, Heights of the corpus mandibulae, between p2-p3, p3-p4, p4-m1, m1-m2,
m2-m3, and behind m3, respectively; 9 and 10, Transverse diameters of the corpus
mandibulae, between p4-m1 and behind m3, respectively; 11, Antero-posterior length of the
symphysis; 13, Antero-posterior diameter of the ramus (at the level of the occlusal line); 14,
Transverse diameter of the articular condyle; 15, Height of the articular condyle; 16, Height
of the coronoid process. Numbers coincide with measurements as defined and illustrated by
Guérin (1980, fig. 1, table 3).

1511

1	2	3	4	5	6	7	8	9	10	11	13	14	15	16
467.4	136.8	68	68	70.7	75	76.4	77.7	38.1	41.3	118.2	136.8	96.2	203.3	229.2

1512
1513

Table 3. Dental measurements of *Brachydiceratherium shanwangense*, from Tagay, early

Miocene of Eastern Siberia, in mm. Abbreviations: H, crown height; L, length; W, width.

		P1	P2	P3	P4	M1	M2	M3	p2	p3	p4	m1	m2	m3
left	L		c25.1	31.5	35.3	47.0	50.5			28.0	31.8	38.6	43.8	43.9
	W		31.5	41.1	47.7	51.7	54.2			21.8	25.2	27.6	29.8	30.5
	H		17.1	21.7	28.3	30.5	37.5			18.2	23.2	28.8	30.2	29.8
right	L	19.3	27.2	29.0	35.5	47.1	50.6	42.1		29.8	32.0	37.7	44.1	45.4
	W	14.5	31.9	40.0	47.6	51.4	56.2	46.1		21.9	27.2	28.1	30.0	30.3
	H	8.8	18.0	21.1	27.0	29.7	37.6	37.8		19.8	23.0	24.8	28.7	27.0

Table 4. Postcranial measurements of *Brachydiceratherium shanwangense*, from Tagay, early

Miocene of Eastern Siberia, in mm. Forelimb bones. Abbreviations: ant, anterior; APD,

antero-posterior diameter; art, articulation; artic, articular; D, distance; del, deltoid; dist,

distal; ext, extremity; H, height; L, length; lat, lateral; mag, magnum; max, maximal; med,

medial; mid, middle; post, posterior; pyr, pyramidal; rad, radius; sml, semilunate; TD,

transverse diameter; tpz, trapezium; tpzd, trapezoid; troch, trochlea; tub, tuberosity; unc,

unciform.

1525

Bone										
Scapula	-	-	-	-	-	glenoid cavity TD　APD 65　86.8	-	-	-	-
Humerus	L 400.3	proximal ext TD　APD 120.7　126.8		diaphysis TD　APD 54.1　55.1		dist TD　APD 127.6　99	TD del tub 104	TD troch 92.5	APD trochlea med　mid　lat 75　42　57.2	
Radius	max L 324.1	proximal ext TD　APD 88.1　58		proximal art TD　max APD 87　47.7	diaphysis TD　APD 39　36.5		distal extremity TD　APD 95.5　54.4		distal articulation TD　APD 84.6　40	-
Ulna	artic L 353.9	olecranon TD　H 67.5　79.8		humeral cochlea TD　H 81.3　66.5	diaphysis TD　APD　46.3 31.4		pyramidal-facet TD　APD 32.1　46.5		dist radius-facet APD　H 16.5　7.2	-
Scaphoid	ant H 49	post H 46.4	TD 52.7	APD 73.6	rad-fac APD 39	tpz-fac TD　APD 18.6　12.4		trap-fac TD　APD 29.1　39.7	mag-fac TD　APD 30.4　31.3	-
Semilunate	TD 36.1	APD 63.6	H 47.1	post TD 28.1	mag-facet TD　APD 22.8　47.7		unc-facet TD　APD 17.8　33	scaph- fac D 10.7	pyram- fac D 6.3	-
Pyramidal	TD 45.3	APD 44	ant H 56.5	ul-fac APD 32.6	unciform-facet TD　APD 38.9　28.5		semilunate-facet D　prox H　dist H 6.8　12.7　9.5		-	-
Pisiform	TD 57.7	APD 26.4	tuberosity H　APD 40.9　15.1		ulna-fac TD 18.6	pyr-fac TD 18.9	-	-	-	-
Trapezoid	TD 34.4	APD 35	ant H　mid H　post H 27.6　19.6　29.4		-	-	-	-	-	-
Magnum	TD 49	ant H 26.1	H 47.7	APD 74.1	sml-fac APD 40.2	McIII-facet TD　APD 43.5　42.2		post tuberosity TD　H 17.6　25.9	-	-
Unciform	TD 52.8	H 40.1	APD max　min 71.7　56.9		post tuberosity TD　H 31　20.7	semilunate-fac TD　APD 22.2　36.8		pyramidal-facet TD　APD 33.8　40.7	McV-fac TD　APD 23.1　29.6	
McII	L 130.9	TD 41	proximal art TD　APD 41　37.3	trapezoid-facet TD APD 30.6　36.4		diaphysis TD　APD 37.3　17.1	dist articulation TD　APD 40.4　41.9	-		-
McIII	L 143.2	med L 133.2	proximal art TD　APD 55.9　40.7	McIV- facets D 12.3	magnum-facet TD　APD 43.5　41.5	diaphysis TD　APD 44　19.4		dist ext TD 54.4	distal art TD　APD 44.5　41	
McIV	L 114.7	proximal art TD　APD 38.1　40.2	unciform-facet TD　APD 29.1　39.6		McV-facet H　APD 12.7　15.3	diaphysis TD　APD 32.8　16.8		dist ext TD 43.2	distal art TD　APD 35.1　36.3	
McV	L 20	TD 27.2	APD 25.7	McIV-facet H　APD 12.9　19	unc-fac APD 22.9	-	-	-	-	-

1526
1527
1528
1529

Table 5. Postcranial measurements of *Brachydiceratherium shanwangense*, from Tagay, early Miocene of Eastern Siberia, in mm. Hind limb bones. Abbreviations: ant, anterior; APD, antero-posterior diameter; art, articulation; artic, articular; As3, astragalus-3 (sensu Heissig, 1972); astr, astragalus; Cc1, calcaneus-1 (sensu Heissig, 1972); Cc2, calcaneus-2 (sensu Heissig, 1972); D, distance; dist, distal; ext, extremity; H, height; L, length; lat, lateral; maj, major; max, maximal; med, medial; mesocun, mesocuneiform; mid, middle; min, minimum; post, posterior; sust, sustentaculum; TD, transverse diameter; tr, trochanter; troch, trochlea; tuber, tuberosity.

Femur	L 486.9	proximal ext		maj tr	third tr		diaphysis		dist ext		distal condyles	
		TD	APD	TD	TD	H	TD	APD	TD	APD	TD	D
		164.2	72.4	97	54.5	51.6	63.5	43	129.4	144.8	114.3	10.7

Patella	TD	APD	H	articulation		med lip	lateral lip		-	-	-	-
	80.1	34	84	TD	H	TD	TD	H				
				68.3	82.5	53.5	29.2	69.5				

Tibia	L	proximal ext		diaphysis		dist ext		fibula-facet		astr cochlea		-
	333	TD	APD	TD	APD	TD	APD	APD	H	TD	APD	
		115.3	100.5	47	40.5	89.5	60.6	33.6	9.7	70.6	50.1	

Fibula	L	proximal ext		diaphysis		dist ext		tibia-facet		astr facet	
	267.7	TD	APD	TD	APD	TD	APD	APD	H	APD	H
		25	39.3	47	40.5	11.8	14.4	37.1	9.6	37.1	18.7

Astragalus	max	troch	max	H			Cc1-facet		Cc2-facet		cuboid-facet	
	TD	TD	APD	medial	mid	lateral	TD	H	TD	H	L	W
	90.9	75.1	53.4	67.3	52.3	70.5	40.5	37.6	39.5	22.5	45	23.2

Calcaneus	H	artic	tuberosity		beak	sust TD	tuber min		As3-facet		cuboid-facet	
	128.9	H 67.3	TD	APD	APD	73.7	TD	APD	TD	H	TD	H
			46.8	61.8	62.9		28.8	51	30.1	11.4	43.9	20.4

Cuboid	TD		max	H		proximal art		distal art		-	-	-
	ant	post	APD	ant	post	TD	APD	TD	APD			
	43	31	61.9	29.9	49.4	39.6	43.8	37.3	37.1			

MtII	L	proximal art		mesocun-fac		lat facet H		diaphysis		distal art		-
	160	TD	APD	TD	APD	ant	post	TD	APD	TD	APD	
		31.3	39.9	25.4	28.4	5.4	12.9	28.7	19.4	33.8	39.5	

MtIV	L 101.7	proximal art		medial facets				diaphysis		dist ext	distal art	
		TD	APD	ant TD	ant H	post TD	post H	TD	APD	max TD	TD	APD
		31.7	33.3	11.3	16.9	18.7	14.8	30.5	18.1	38.3	30.4	40.2

 Table 6. Distribution of unambiguous apomorphic characters (synapomorphies and

autapomorphies, including reversals) among teleoceratine rhinocerotids, as retrieved in the

current phylogenetic analysis. Node numbers match those of Fig. 13. Binominal combinations

are as detailed in the Discussion.

Node 13 (Teleoceratina): -72^0, 129^1, 205^1, 279^1, 282^1

Diaceratherium tomerdingense (type and only species): -70^0, 90^1, 121^1, 130^1, -196^0, 212^1,

216^1, 223^3, 228^2, -251^0

Node 14: 199^2, 202^1, 227^1

Brachypotherium: -50^0, 54^2, -83^1, -114^0, 140^1, -146^0, 155^1, 203^1, 214^1, -226^0, 254^1

Brachypotherium brachypus: 3^2, -34^0, -39^0, 57^1, 62^1, 99^2, -109^1, -115^2, 118^1, -119^0, -135^0,

144^1, -149^0, 157^3, -159^0, -160^0, 170^1, 179^1, 180^1, 191^1, -193^0, -204^0, 209^1, 210^1, 239^1, 275^2,

-280^0

Brachypotherium perimense: 25^1, 41^1, 68^1, 76^1, -85^0, 121^1, -125^0, 128^1, -129^0, 151^3, 172^1,

173^1, 175^1, 181^1, 199^3, 200^1, -205^0, 228^2, 246^1, 248^1, 255^1, 263^3, -271^0, 272^1, 274^1, 277^1, -

282^0

Node 16: 38^1, 101^1, 107^3, -147^0, -222^0, -230^0, -263^0, 264^3

Node 17 (*Teleoceras* + *Prosantorhinus*): 10^1, 11^1, 40^1, 57^1, 95^1, 99^1, 130^1, 180^1, 190^2, 191^1,

234^1, 272^2, 275^2, -280^0

Teleoceras fossiger (type species): 2^1, 19^1, 25^1, 48^1, -56^0, 59^1, 60^0, 63^1, 68^1, 80^1, 87^2, 88^2, 91^2,

99^2, 102^1, 114^3, 116^1, 117^1, 121^1, 126^1, 128^1, 144^1, 151^3, 153^1, 157^3, 159^3, 165^1, 173^1, 175^1,

180^3, -196^0, 212^1, 220^1, 248^1, -249^0, 254^1, 268^1, 269^1, 276^1

Prosantorhinus: 1^1, 27^1, -35^0, 45^1, -119^0, -125^0, -177^0

Prosantorhinus germanicus (type species): 37^1, 85^3, 105^1, -109^1, -110^0, -111^2, 115^2, 147^2,

148^1, -149^1, -151^1, 166^1, -182^0

Node 19: -101^0, -129^0, -135^0, -159^1, 182^2, 261^3

Prosantorhinus laubei: -85^1, 90^1, -99^0, -114^0, -134^0, -180^0

Prosantorhinus douvillei: -88^0, 99^2, 114^2, 144^1, 156^2, -157^0, -159^0, 162^1, 180^3

Brachydiceratherium: -20^0, -39^0, 65^1, 101^3, -149^1, -159^0, 185^1, 187^2, 246^1, -251^0

Node 21: 72^1, 105^2, -190^0, -193^0, 238^1, 250^1, -279^0, -282^0

Brachydiceratherium shanwangense: 1^1, 27^1, 63^1, -70^0, -85^0, 87^2, 88^2, 105^3, 116^1, 147^2, 151^3,

157^3, 191^1, -196^0, 199^3, 234^1

Node 22: 121^1, 210^1, -252^1, -264^0, -265^0

Brachydiceratherium aginense: 112^1, -114^0, 128^2, 130^1, -149^0, 162^1, 214^1, 216^1, 219^1, -221^0, -

226^0, 256^1, -262^0, 263^1, 272^1, -280^0

Brachydiceratherium intermedium: -65^0, 114^3, 118^1, -129^0, 220^1

Node 23: -119^0, 203^1, -204^0, 207^1, 261^1

Brachydiceratherium fatehjangense: 10^1, 18^2, 48^1, -49^0, 53^1, 54^2, 57^1, -70^0, -94^1, -107^0, 140^1, -

149^0, 151^3, 155^1, 174^1, -202^0, -205^0, 209^1, -224^0, -227^0, -247^0, 261^2

Node 24: 118^1, -125^0, -134^0, 230^1

Brachydiceratherium aurelianense: -53^0, 86^1, 90^3, 114^3, 124^1, 130^1, 147^1, -193^0, 198^1, -199^0,

214^1, 220^1

Node 25: 105^1, 210^1, -253^0, -264^0, -279^0, -280^0

Brachydiceratherium lamilloquense: 53^2, -84^1, 88^1, 99^1, -102^1, 105^3, -111^2, 112^3, -135^0, -146^0,

207^2, 228^2, -243^0, -246^0, -247^0, 248^1, -259^0

Node 26: -109^0, -138^0, -221^0

Brachydiceratherium asphaltense: -23^0, 27^1, 83^3, -227^0

Brachydiceratherium lemanense (type species): 11^1, 40^1, 45^1, -47^0, 48^1, -50^0, -70^0, 147^1, -226^1

1588
1589
1590