**Conjunctive and complementary CA1 hippocampal cell populations relate sensory events to immobility and locomotion**

Samsoon Inayat[1*], Brendan B. McAllister[1], Ian Q. Whishaw[1], and Majid H. Mohajerani[1*]

1. Canadian Centre for Behavioural Neuroscience, University of Lethbridge, Lethbridge, Alberta, Canada

Corresponding authors: samsoon.inayat@gmail.com, mohajerani@uleth.ca

## Abstract

This study investigated the dynamics of recruitment of cells in the CA1 region of the hippocampus in response to sensory stimuli given during immobility, movement, and their transitions. Two-photon calcium imaging of somal activity in hippocampal CA1 neuron populations was done in head fixed mice given a light flash or an air puff when at rest, when moving, and when induced to run for a fixed distance on the conveyor belt when a brake was released. Overall, 99% of 2083 cells were active across 20 sensorimotor events. A larger proportion of cells were active during locomotion. Nevertheless, for any sensorimotor event, only about 17% cells were active. These consisted of conjunctive (C ∈ A and B) cells that were shared among events and complementary (C ∈ A not B or C ∈ B not A) cells that were active only during individual events. Whereas conjunctive cells characterised stable representations of repeated sensorimotor events, complementary cells characterised recruitment of new cells for encoding novel sensorimotor events. The moment-to-moment recruitment of conjunctive and complementary cells with changing sensorimotor events signifies the involvement of hippocampus in functional networks integrating sensory information with ongoing movement. This role of hippocampus is well suited for movement guidance that might include spatial behavior, episodic learning and memory, context representation, and scene construction.

## Introduction

The idea that the brain features behavior-specific electrographic events during the behavioral states of immobility and locomotion originated with the finding that electrical activity in the hippocampus is different during these two states [1,2]. The electrical activity measured as local field potentials (LFPs) features large amplitude irregular activity (LIA) during immobility and two kinds of rhythmical slow activity (RSA or theta activity, 6-12 Hz) that are sensory or movement related [1,3-9]. Sensory-induced

RSA is cholinergic-dependent whereas movement-related RSA can be cholinergic-independent [10-12]. The behavior-specific differentiation of electrical activity is also found at the cellular level. Whereas excitatory cells in both CA1 and CA2 modulate their firing in relation to location during immobility [13,14] and during locomotion [15,16], anatomically distinct types of inhibitory cells fire preferentially during locomotion versus immobility [17]. Location specific firing during locomotion has also been found in other subregions of the hippocampal formation such as DG, CA3, Subiculum, and medial entorhinal cortex as well as cortical regions including retrosplenial, visual, and sensorimotor cortices. There are reports of behavior-dependent cellular responses in the hippocampus when animals encounter sensory stimuli including odors [18-22], tones [23,24], tastes [25], textures [26], visual stimuli [27,28], somatosensory stimuli [29,30], and vestibular events [31].

Behavior-specific sensory processing is also reported in sensory and non-sensory cortices, and some subcortical structures display behavior-dependent spontaneous and evoked firing activity - for a review, see [32]. Compared to immobility, locomotion increases firing of sensory neurons in the visual [33,34] and somatosensory [35,36] cortices but reduces firing in the auditory cortex [37]. Locomotion also modulates higher order functions; for example, it improves sensory perception [38], guides navigation through the visual cortex [39], and alters the functional connectivity of visual and retrosplenial cortices with other cortical regions [40]. Movements other than locomotion such as whisking, facial and body movements, and arousal-related changes in pupil diameter also influence neuronal firing in the sensory cortices [41,42]. Furthermore, movement-related inputs to the sensory cortices reflect expectations of sensory information anticipated by movements [43,44]. Such studies have contributed to the idea that such activity mediates spatial navigation and context processing [see reviews, [45-48]].

Stimulated by the many finding of the relationship of electrical events to ongoing behavior, the present study examined such sensory behavior-specific hippocampal cellular responses to sensory

stimuli during immobility, locomotion, and their transitions. Calcium imaging was used in head-fixed mice to record hippocampal CA1 cell responses to sensory stimulation by a pressurized air stream or light flash while the mice were at rest or locomotin. A population level analysis investigated how cells were recruited during and between sensory stimuli applied during immobility versus locomotion. Subgroups of cells responding to different sensorimotor events were active in different stimulus-locomotion configurations. When two (or more) events were considered at a time, cells could be classified as complementary, responsive to only one event and conjunctive, responsive to both of events. From a large pool of responsive cells, complementary cells identified novel sensorimotor events and conjunctive cells encoded familiar sensorimotor events suggesting a recruitment with replacement organization.

## Results

### Experimental design, behavioral paradigm, and its characterization

Hippocampal CA1 cell activity was recorded from head-fixed mice resting or moving on a non-motorized conveyor belt (Fig. 1A). The application of a brake restricted movement and its release enabled movement and running. An air stimulus, delivered on the back, and whole-field illumination with an LED in front of the mouse served as sensory stimuli. The behavioral paradigm featured 7 configurations of motor activity and sensory stimulation (Fig. 1B) divided into two sets based on two brake conditions: Brake (B) and No-Brake (NB). In Configurations 1, 2, 6, and 7, the B-condition prevented walk/run behaviors whereas in Configurations 3, 4, and 5, the NB-condition allowed walk/run behaviors. In Configurations 1 and 2, ten pulses each of light (L; 200 ms with 5 s interval) and air (A; 5 s with 10 s interval) were applied, respectively. In Configurations 3, 4, and 5, the air stream was applied to cause stimulus-induced running for a fixed distance of 150cm, one lap of the belt (before the air was

stopped). Ten trials were given in each condition with an intertrial interval of 15 s. In Condition 4, a 200 ms light flash was also applied once the mouse had traveled 110 cm. Configurations 3 and 5, 1 and 6, and 2 and 7 were identical.

**Figure 1.** The experimental paradigm for studying hippocampal CA1 neuronal responses to sensory stimulation during immobility and locomotion. **(A)** Experimental setup. A head-fixed mouse on a linear conveyor belt receives a mild air stream on the back to motivate running. **(B)** Experiment timeline and list of 7 configurations that mice experienced while calcium imaging was performed. Inset shows schematic depicting air onset (AOn), air offset (AOff), and arbitrary (Arb) events. **(C)** Average speed over all trials by animals around stimulus and arbitrary (Arb) events during Configurations 3, 4, and 5. Magenta lines indicate occurrences of events. **(D)** Distribution of speeds at events mentioned in C. The numbers indicate overall mean±SEM speeds. **(E)** Average image (left) of the calcium imaging time

series from a representative animal, and arbitrarily color-coded regions of interest indicating identified cell bodies (right) in a ~400 × 400 μm imaging window. **(F)** Normalized calcium traces ($\Delta F/F_0$), from 100 representative cells from a representative animal for a whole recording of ~25 min, overlayed with indicators of light and air stimuli (vertical-colored lines). Cells are arranged by the time of peak firing. The speed signal is shown at the bottom (cyan color). Notice differential firing of cells during immobility vs. locomotion (cells in blue vs red ovals) and cell population changes from sensorimotor event to sensorimotor event. Thick black lines in (C) and (D) indicate mean over animals while shaded regions represent SEM.

Neuronal activity 2 s before and 2 s after each stimulus event was analyzed and compared across B and NB conditions. For air stimuli, activity around both air onset and offset events (AOn and AOff respectively) was characterized whereas for light stimuli, activity at only light onset (L) was analyzed because of the brevity of stimulus duration i.e., 200 ms. Because the hippocampus is an associative brain region, cellular activity may have resulted from intrinsic network activity representing task parameters or internal states. Therefore, the activity of cells was also analyzed around arbitrary (Arb) time points in which no explicit sensory stimulation occurred in both B and NB conditions. The time chosen for Arb events (Fig. 1B, inset), was the mid-point of the intertrial intervals (in Configurations 2, 3, 4, 5, and 7). For B and NB conditions, the Arb point was 5 s and 7.5s after air offset respectively. Finally, during NB-condition, activity of neurons was analyzed when animals voluntarily started or stopped their locomotion in the absence of explicit sensory stimuli.

The animals were immobile during the B-condition. During the NB-condition, their average speed was always greater than zero around stimulus and Arb events (Fig. 1C). For AOff and L events, the speed in all trials (from Configurations 3-5) of individual animals was greater than zero (Fig. 1D, middle panels). For AOn, however, $50.00 \pm 12.78$ % of the trials on average (over 5 animals) had speeds greater than zero (Fig. 1D, left panel). There was animal-to-animal variability with greater than zero speeds in 3, 26, 18, 17, 11 trials (corresponding to 5 animals) of the cumulative 30 air application trials. In trials, in which the animals were immobile, they initiated locomotion after air onset with an average

movement latency of $0.94 \pm 0.06$ sec (range: 0.05, 3.66, median: 0.62) and hence these trials were pooled with trials in which the mice were already moving at air onset. For the Arb events, there was animal-to-animal variability with greater than zero speeds in 2, 20, 21, 8, 6 trials (corresponding to 5 animals) of the cumulative 30 trials (Fig. 1D). Overall, however, the average speed around the Arb events was greater than zero (Fig. 1C).

**Calcium imaging of the activity of hippocampal CA1 pyramidal neurons**

Calcium imaging of CA1 pyramidal neurons was performed as the mice experienced the 7 behavioral configurations. Fig. 1E presents an average image of an imaging stack from a representative animal showing its densely packed dorsal CA1 neurons. Using Suite2p, a Python based analysis software, 603, 568, 315, 338, and 259 individual cells were identified from 5 mice (total: 2083 cells). An image of identified cells from a representative animal is displayed in Fig. 1E. The figure also shows that Suite2P did not recognize all visible cells in the average image, perhaps because some cells were seldom active (fired only occasionally or not at all). The raw calcium traces ($\Delta F/F_0$) of 100 cells from a representative animal shown in Fig. 1F demonstrate that cellular responses reflected brake conditions and/or sensorimotor events. In one example, the cell groups colored red were active in the B-condition and those colored blue were active in the NB-condition. For further quantitative analysis of cellular responses to sensory events and behavior, as well as study of population level characteristics, the calcium signals were deconvolved to estimate firing rates for each cell. In the subsequent text, neural activity refers to the firing rates estimated from the calcium signals.

To examine the changes in cellular firing properties across B and NB conditions, mean and maximum firing rates were determined for both conditions and for each cell. The mean and maximum firing rates were then calculated to obtain the respective values for each animal. These values were then

subjected to statistical comparison across conditions with a repeated measures analysis of variance (RM-ANOVA). The mean firing rate across B and NB conditions was not significantly different [$0.054 \pm 0.016$ vs. $0.073 \pm 0.022$; $F(1,4) = 7.05$, $p = .057$, $\eta^2 = .64$], but note a higher mean for NB-condition and a p value close to 0.05. The comparison of the maximum firing rates of cells for B and NB conditions gave a significant difference [$26.47 \pm 5.05$ vs. $48.51 \pm 10.14$; $F(1,4) = 13.89$, $p = .020$, $\eta^2 = .78$] showing that locomotion contributed to increased transient cell firing.

## Larger population of responsive cells for No-Brake condition vs Brake condition for AOn, AOff, and Arb events

Cell activity during B and NB conditions for AOn, AOff, and Arb events was investigated by finding responsive cells and their response characteristics. Cells active around these events, were examined by generating raster plots of neuronal activity (trials versus time) for each cell and event. Peri-event time histograms (PETHs) were then calculated from raster plots by finding mean neuronal activity over trials. Responsive cells were identified from PETHs by statistically comparing average pre- and post- event firing rates using a Student's t-test (alpha < 0.05). This method allowed an unbiased approach for identifying both excited (Exc) and inhibited (Inh) cells in which post-event firing rates were either significantly larger or smaller from their pre-event firing rates. Figure 2A displays raster plots for a representative AOn-Exc cell (left) and AOn-Inh cell (right) from the Configuration 2, with firing rates across time and across trials overlayed with the PETH. Population level response dynamics for responsive cells were observed with rate vector maps (Fig. 2B, top row). These show the normalized mean response of all cells organized by the time of each cell's peak response. The rate vector maps for both the B and NB conditions feature the binary nature of Exc and Inh cells (i.e., their responses of ON or OFF type) in relation to AOn. These are illustrated in population correlation plots (Fig. 2B, middle

row) in which each pixel represents a Pearson correlation between two columns of the corresponding

rate vector map. Population correlations displayed dark square-like regions for both the pre-event (-2 to

0 s) and post-event (0 to 2 s) periods corresponding to responses of Exc and Inh cells, respectively. Plots

of the average population correlation (Fig. 2B, bottom row) confirmed that the majority of Exc and Inh

cells were binary in nature. Plots generated for AOff and Arb events exhibited similar characteristics

with both Exc and Inh cells.

**Figure 2.** More cells active during the No-Brake compared to the Brake condition for AOn, AOff, and Arb events. **(A)** Raster plots and peri-event histograms of representative excited (Exc) and inhibited (Inh) cells showing increased and decreased firing, respectively, in response to AOn (t = 0, magenta line). The blue lines show the means across trials. **(B)** Rate vector maps (top row) and corresponding population vector correlation plots (middle row) of responsive cells to AOn from a representative animal for Brake vs. No-Brake conditions. Average population correlation plots (bottom row) from 5 animals. **(C-E)** Results of two-way RM-ANOVA for assessing responsivity (Resp), response fidelity (RF), and mutual information z-score (zMI) respectively, across brake conditions (B versus NB) and events (Aon, AOff, and Arb) **(F-H)** Results of three-way RM-ANOVA for assessing Resp, RF, and zMI respectively, across brake conditions, events, and cell types (Exc vs Inh). For all statistical tests, $n = 5$ mice. Error bars = SEM. *$p < .05$, **$p < .01$, ***$p < .001$.

To assess the responsivity (Res) of cell activation across B and NB conditions, the percent of responsive cells (including both Exc and Inh cells) around AOn, AOff, and Arb events was found. Resp was averaged over Configurations 2 and 7, and Configurations 3, 4, and 5 for B and NB conditions, respectively. The data were subject to a two within-subjects factors (Brake-State and Event-Type) RM-ANOVA. There were significant main effects of both Brake-State [$F$ (1,4) = 130.12, $p < .001$, $\eta^2 = .97$] and Event-Type [$F$ (2,8) = 19.22, $p = .009$, $\eta^2 = .83$] with no interaction. The Resp in the NB-condition was larger than in the B-condition and *post hoc* comparisons revealed that the percent of cells activated around AOn was larger than for AOff and Arb events (Fig. 2C).

Responsive cells found from PETHs might have had different response characteristics across trials during B and NB conditions in response to AOn, AOff, and Arb events. Therefore, two measures, response fidelity (RF) and mutual information (MI) were extracted from raster plots to quantitatively assess the strength and robustness of responses across trials. RF was defined as percent of trials in which the cell's firing rate was greater than zero at any point during a trial. MI between firing rate and time was determined from a raster plot [49] and then z-scored (zMI) using a distribution of 500 MI values calculated from randomly shuffled instances of the same raster plot. A larger zMI value indicates better/tighter temporal repeatability of response across trials. RF and zMI values were averaged across Configurations 2, 7, and Configurations 3, 4, and 5, for B and NB conditions, respectively. The results were subject to a two within-subjects factors (Brake-State and Event-Type) RM-ANOVA individually for RF and zMI. For RF, there were significant main effects of both Brake-State [$F$ (1,4) = 41.32, $p = .003$, $\eta^2 = .91$] and Event-Type [$F$ (2,8) = 30.62, $p = .000$, $\eta^2 = .88$] with no interaction. RF in the B-condition was larger compared to the NB-condition and post hoc comparisons revealed that the RF for AOn and AOff was larger compared to Arb events (Fig. 2D). For zMI, there was a main effect of Event-

Type [$F$ (2,8) = 42.50, $p$ = .001, $\eta^2$ = .91] with post hoc comparisons showing that zMI for AOn and AOff events was larger compared to Arb events and that for AOn was larger compared to AOff (Fig. 2E).

Taken together the results suggest that more cells are active during NB-condition compared to B-condition. Nevertheless, cells that were active during B-condition had more robust responses than occurred for cells active during the NB-condition. Additionally, cells that were activated by sensory air events had a more robust response compared to those activated around Arb events, especially at AOn.

**Characteristics of excited and inhibited cells is similar across Brake and No-Brake conditions but different for AOn and AOff events**

The analysis of cell populations activated across B and NB conditions around AOn, AOff, and Arb events was further extended to ask how Brake-State and Event-Type might have invoked different cell types (Exc and Inh cells). A three within-subjects factors (Brake-State, Event-Type, and Cell-Type) RM-ANOVA was used on the output variables: Resp, RF, and zMI.

For Resp, there was a significant three-way interaction, [$F$ (2,8) = 6.73, $p$ = .028, $\eta^2$ = .63]. Therefore, simple two-way interactions of two factors at each level of the third factor were assessed with a Bonferroni adjusted alpha value (see methods). Significant simple two-way interactions were found between Event-Type and Cell-Type for both Brake [$F$ (2,8) = 51.37, $p$ < .001, $\eta^2$ = .93] and NB-condition [$F$ (2,8) = 9.14, $p$ = .010, $\eta^2$ = .70]. In the next level of analysis, simple simple main effects were assessed with a Bonferroni adjusted alpha value. Only simple simple main effects of Event-Type were significant: B-condition, Exc cells [$F$ (2,8) = 56.15, $p$ < .001, $\eta^2$ = .93], Inh cells [$F$ (2,8) = 32.19, $p$ < .001, $\eta^2$ = .89], and NB-condition, Exc cells [$F$ (2,8) = 12.98, $p$ = .003, $\eta^2$ = .76]. Simple simple comparisons with Bonferroni correction then assessed differences in means for Event-Types (Fig. 2F).

The effect of AOn was stronger in generating Exc responses for both B and NB conditions, but AOff was more potent in generating Inh responses only for B-condition although the trend was similar for NB-condition.

For RF, there was a main effect of Brake-State [$F$ (1,4) = 32.21, $p$ = .005, $\eta^2$ = .89] and a two-way interaction between Event-Type and Cell-Type [$F$ (2,8) = 25.50, $p$ = .006, $\eta^2$ = .86]. Posthoc comparisons (Fig. 2G) showed that Exc and Inh cells activated around AOn and AOff respectively had the largest RF in both B and NB conditions. For zMI, there was a significant two-way interaction between Event-Type and Cell-Type [$F$ (2,8) = 21.26, $p$ = .004, $\eta^2$ = .84]. Post hoc comparisons revealed larger zMI values for AOn-Exc cells and AOff-Inh cells (Fig. 2H).

Taken together, the results suggest that Exc and Inh cells were similar in the B and NB conditions, but AOn was associated with more Exc cells and AOff was associated with more Inh cells. As found before, sensory air events generated more robust responses compared to Arb events for both Exc and Inh cells.

**Separate groups of cells in Brake and No-Brake conditions for AOn, AOff, and Arb events**

The population level activity of cells was classified across brake conditions and different events as conjunctive and complementary. Considering two cellular populations active around two sensorimotor events e.g., A and B, conjunctive (Conj) cells were defined as those active for both events (Conj ∈ A and B) and two types of complementary (Comp) cells were defined as those active for individual events (Comp1 ∈ A not B; and Comp2 ∈ B not A). The percent of these three cell populations (of the total number of cells) for different sensorimotor events was compared statistically or with a secondary level of analysis to observe relations among sensorimotor events and configurations.

**Figure 3.** Segregated cell populations in the Brake and the No-Brake conditions for AOn, AOff, and Arb events. **(A)** Results of a two-way RM-ANOVA for assessing the percent of cells across population types (Comp1, Comp2, and conjunctive) and events (AOn, AOff, and Arb). **(B)** Results of a three-way RM-ANOVA assessing the percent of cells across cell types (Exc vs Inh) in addition to factors in (A). **(C)** Overall responsivity, conjunction, and complementation across Brake and No-Brake conditions. **(D-E)** Average and SEM (insets) heatmaps (over 5 animals) of conjunction and complementation of Exc and Inh cell groups active in Brake and No-Brake conditions around AOn, AOff, and Arb events. **(F-H)** Dendrograms resulting from agglomerative hierarchical clustering of the average conjunction and complementation heatmaps/matrices. CC indicates cophenetic correlation. For all statistical tests, $n = 5$ mice. Error bars = SEM. $*p < .05$, $**p < .01$, $***p < .001$.

Comp1, Conj, and Comp2 cell populations were found for AOn, AOff, and Arb events where Comp1 and Comp2 were selectively active during B and NB conditions respectively and Conj were active in both. For this analysis, cells were pooled across Configurations 2, 7 and Configurations 3, 4, 5 for B and NB conditions, respectively. The population percent values were then subjected to a two-way RM-ANOVA with Event-Type and Population-Type as within-subjects factors. There was a significant main effect of both factors, Event-Type [$F$ (2,8) = 10.53, $p$ = .030, $\eta^2$ = .72] and Population-Type [$F$ (2,8) = 715.73, $p < .001$, $\eta^2$ = .99] but no significant interaction. The former result was expected, as described above for differences in Resp across sensory air and Arb events (Fig. 2C and Fig. 3A). The later result however showed differences in percent across population types with Conj cells forming a significantly smaller population compared to Comp1 and Comp2, as identified with post hoc comparisons (Fig. 3A).

The complementation and conjunction of Exc and Inh cells was then examined by extending the analysis by adding a third factor (Cell-Type) and performing a three-way RM-ANOVA. There was a significant three-way interaction [$F$ (4,16) = 7.19, $p$ = .002, $\eta^2$ = .64]. Further tests showed that the only simple two-way interaction was between Cell-Type and Event-Type for Comp1 type of cells [$F$ (2,8) = 20.68, $p$ = .003, $\eta^2$ = .84]. Simple simple main effects of both Cell-Type and Event-Type were significant: [$F$ (2,8) = 12.88, $p$ = .003, $\eta^2$ = .76] and [$F$ (2,8) = 11.31, $p$ = .005, $\eta^2$ = .74], respectively. Post hoc comparisons showed that for Exc cells, AOn formed a larger percentage compared to AOff . The opposite relation was obtained for Inh cells (Fig. 3B). For Conj and Comp2 cells, there was no difference across Exc and Inh cell types (Fig. 3B, pooled data).

When pooled across event and cell types, 65.5 ± 2.3 % of cells were responsive comprising three populations. One population was active during the B-condition (Comp 1, 9.7 ± 0.7 %), a second population was active during the NB-condition (Comp 2, 40.9 ± 0.6 %) and the third population was

active in both conditions (Conj, $14.9 \pm 2.2$ %) (see Fig. 3C, Venn Diagram). After pooling however, the

average percent of Comp1 cells was not significantly different from the average percentage of Conj cells

(Fig. 3C, RM-ANOVA and post hoc comparison). The latter relationship suggests that some cells might

have shared responsiveness across sensory air and Arb events. To further investigate this possibility, the

conjunction and complementation of all different groups of cells was analyzed. The conjunction and

complementation of different groups of cells categorized by brake conditions (B vs NB), event types

(AOn, AOff, and Arb) as well as cell types (Exc vs Inh) were plotted as heat maps. Figures 3D and 3E

show average conjunction and complementation heatmaps respectively (over 5 animals) and insets show

SEM heatmaps. In the average heatmaps, diagonal pixels are shown as white with no color coding and

numbers therein represents the percentage of responsive cells for the selected category. Off-Diagonal

pixels in these heatmaps show percentages of conjunctive and complementary cells. For the

complementation heatmap, pixels above and below the diagonal represent Comp1 and Comp2 cells,

respectively when considering row-column pairs.

Conjunctive cell populations were small, with ~3 % conjunction between Exc cells responsive to

AOn and Inh cells responsive to AOff during the Brake conditions (Fig. 3D, B-AOn-Exc vs B-AOff-Inh,

row 1 column 4). To find an overall classification of cell populations and their relationships with each

other, a secondary analysis with agglomerative hierarchical clustering (see methods) of the average

heatmap matrix was used. Clustering revealed four distinct clusters (Fig. 3F). The cyan cluster grouped

Exc cells responding to AOn with Inh cells responding to AOff during the B-condition indicating cells

that responded during air ON periods i.e., when air was on. A similar relation existed for the same types

of cells and events during the NB-condition (red cluster). The third cluster constituted cells from the B-

condition (purple cluster), which were active during air OFF periods i.e., when air was not on. The

fourth cluster (green) represented air OFF cells for the NB-condition. These results show that

conjunctive cells across sensory air and Arb events could be segregated into two groups for the B and NB conditions, and further subdivided into cells that responded to air ON and OFF periods.

The complementation heatmap showed segregation between B and NB conditions with a largest complementation between NB-AOn-Exc and B-AOff-Exc cell groups (~15.3 %). The complementation heatmap was split into two heatmaps, symmetric across the diagonal, by using the upper and lower triangular matrices corresponding to Comp1 and Comp2 cells. The clustering of Comp1 heatmap (Fig. 3G) showed separate grouping of cells across B (green cluster) and NB (blue and red clusters) conditions. The clustering of Comp2 (Fig. 3H) showed similar cluster segregation of B and NB conditions (cyan and red clusters, respectively). However, within the dendrograms of event-cell-type combinations, grouping mainly occurred based on event-type e.g., NB-AOff-Exc closer to NB-AOff-Inh and NB-AOn-Exc closer to NB-AOn-Inh. Taken together, the results show segregation of cell groups across B and NB conditions when cells were divided into subgroups based on AOn, AOff, and Arb events and cell types.

**Separate groups of cells for Light-onset in Brake and No-Brake conditions**

Light-onset cell populations were analyzed to investigate whether same or different cellular networks might be involved across B and NB conditions. The above-mentioned analysis (for AOn, AOff, and Arb events) was repeated for cell populations active around light stimulus events in Configurations 1 and 4 for B and NB conditions, respectively. Both Exc and Inh cells were found in a similar way as is explained above. Fig. 4A shows raster plots and peri-event histograms (PETHs) for representative Exc and Inh cells from Configuration 1. Rate vector maps (Fig. 4B, top row) and population vector correlation plots (Fig. 4B, middle and bottom rows) showed the binary nature of cellular responses for both representative populations in the B and NB conditions.

**Figure 4.** Segregated cell populations across the Brake and No-Brake conditions to a light stimulus. **(A)** Raster plots of representative Exc and Inh cells. The magenta line at t = 0 indicates light onset. The blue lines show means over trials. **(B)** Rate vector maps (top row) and corresponding population vector correlation plots (middle row) of responsive cells from a representative animal for Brake and No-Brake conditions. Average population correlation plots (bottom row) from 5 mice. **(C-E)** Results of one-way RM-ANOVA assessing responsiveness, response fidelity, and mutual information z-score of Exc and Inh cells for light stimulus in Brake condition (B-L). **(F-H)** Results of two-way RM-ANOVA assessing response variables for Exc and Inh cells in the No-Brake conditions for light stimulus combined with air (NB-AL) and identical Arb event where only air was present (NB-A). **(I)** Overall responsivity, conjunction, and complementation for B-L, NB-AL, and NB-A cell populations. **(J)** Results of three-way RM-ANOVA for assessing the percent of cells across population types (Comp1, Comp2, and Conj), cell types (Exc vs Inh) and event pairs e.g., B-L::NB-AL. For all statistical tests, $n$ = 5 mice. Error bars = SEM. *$p < .05$, **$p < .01$, ***$p < .001$.

The Resp, RF, and zMI were separately assessed for B and NB conditions because only the light stimulus was applied alone in B-condition (B-L), whereas for the NB-condition, light was applied while the air was ON (NB-AL, Air-Light).  For B-condition, values were averaged from Configurations 1 and 6. There was no significant difference between Exc and Inh cells for Resp and zMI values, but the RF of

Exc cells was larger than that of Inh Cells [$F$ (1,4) = 9.85, $p$ = .035, $\eta^2$ = .71] (Fig. 4C-E). For the NB-condition, NB-AL cell population from Condition 4 was compared with cells active for identical Arb events in Configurations 3 and 5 where only air was being applied with no light stimulation (NB-A, Air). A two-way RM-ANOVA with Cell-Type (Exc and Inh) and Event-Type (A and AL) as within-subjects factors was used and none of the main or interaction effects were significant for any of the variables (Fig. 4F-H).

Cell populations corresponding to the three sensorimotor events, B-L, NB-A, and NB-AL were then compared for complementation when taken collectively and pooled across Exc and Inh cell types (Fig. 4J). There were separate groups of cells active around the three events with NB-A having the largest percentage of complementary cells (tested with RM-ANOVA with Event-Type as within-subjects factor) [$F$ (2,8) = 8.99, $p$ = .009, $\eta^2$ = .69]. When taken two at a time, a three-way RM-ANOVA was conducted with Event-Type-Pair e.g., B-L::NB-AL, Population-Type (Comp1, Conj, and Comp2), and Cell-Type (Exc and Inh) as within-subjects factors. There was a significant three-way interaction [$F$ (4,16) = 5.83, $p$ = .004, $\eta^2$ = .59]. Simple two-way interactions were then examined, and the only difference was between Event-Type-Pair and Population-Type for Inh cells [$F$ (4,16) = 16.71, $p$ = .005, $\eta^2$ = .81]. Simple simple main effects of Population-Type were then individually determined for all Event-Type-Pairs and all were significant: B-L::NB-A [$F$ (2,8) = 54.13, $p$ < .001, $\eta^2$ = .93], B-L::NB-AL [$F$ (2,8) = 41.41, $p$ < .001, $\eta^2$ = .91], and NB-A::NB-AL [$F$ (2,8) = 12.12, $p$ = .004, $\eta^2$ = .75]. Post hoc comparisons revealed that for the first two Event-Type-Pairs involving B and NB conditions, there were more complementary cells than conjunctive cells, but this was not the case for the third Event-Type-Pair involving events in only NB-condition (Fig. 4J). The percent of complementary Exc cells was significantly larger than the percent of conjunctive cells (Fig. 4J) for all Event-Type-Pairs. These results

suggest a segregation of cell populations across B and NB conditions for both Exc and Inh cells active around light stimuli that are different from the cells active around only-air-Arb (A) event.

**Conjunction and complementation of cell populations characterize stable and dynamic cellular representations of sensorimotor events**

The dynamics of the recruitment of cells across different sensorimotor events in all behavioral configurations was analyzed by finding the conjunction and complementation of cell populations active around all sensory and Arb events after pooling across Exc and Inh cell types (Fig. 5A-B). Agglomerative hierarchical clustering of the average heatmaps was then used to identify relationships among conditions (Fig. 5C-E). The clustering analysis showed a segregation of cellular populations across B and NB conditions either with clear clustering or branch separation in dendrograms within the same clusters. However, there was some mixing of populations across B and NB conditions for Arb events.

The clustering from the conjunction heatmap (Fig. 5C) revealed that distinct subgroups of cells were active for distinct sensorimotor events. For example, the blue cluster shows the closeness of cells active around AOn in the NB-condition (Configurations 3, 4, and 5). Similar relationship exists for the B-condition (Configurations 2 and 7), see dendrogram branch in red cluster enclosed with dotted blue line. The green cluster represents cells active around AOff in the NB-condition and the branch in red cluster enclosed with dotted green line represents similar cells for the B-condition. The branch indicated with magenta arrowhead shows the closeness of populations active around light in the B-condition (Configurations 1 and 6). The cell populations active around Arb events in the NB-condition (3-Arb, 4-Arb, and 5-Arb) however, were closer to cell populations active in the B-condition. When the Arb, A,

and A-L events were excluded from clustering, more distinct grouping of cell populations occurred

which were active for distinct sensorimotor events (Fig. 5F) and there was clear separation between the

B and the NB conditions.

**Figure 5.** Separate subgroups of cells are active for distinct sensorimotor events. **(A-B)** Average and SEM (insets) heatmaps (over 5 mice) of conjunction and complementation of cell groups active in the Brake and No-Brake conditions around sensory air, light, and Arb events. **(C-E)** Dendrograms resulting from agglomerative hierarchical clustering of the average conjunction and complementation heatmaps/matrices. **(F-H)** Dendrograms resulting from agglomerative hierarchical clustering of the average conjunction and complementation heatmaps/matrices after excluding arbitrary events. CC indicates cophenetic correlation.

The clustering of both complementation heatmaps (Fig. 5D-E) revealed grouping of sensorimotor events that either belonged to the same Configuration or those that happened close in time. For example, in the purple and cyan clusters in Figures 5D and 5E respectively, 1-L was grouped with 2-AOff and 2-Arb events (for B-condition). Similarly, the green cluster in Figure 5D shows grouping of sensorimotor events in the NB-condition with branch proximity of sensorimotor events belonging to the same Configuration e.g., 5-AOff, 5-A, 5-Aon enclosed with dotted brown line. The cyan cluster in Fig. 5D shows grouping of events in Configurations 6 and 7 with 5-Arb event from the NB-condition. In both heatmaps for complementation, with Arb, A, and A-L events were excluded, a clear separation between B and NB conditions was seen. Furthermore, within branches, grouping of sensorimotor events occurred based on the Configuration (or Configurations) that occurred closer in time (Fig. 5G-H) e.g., 1-L closer to 2-AOn and 2-AOff in Fig. 5G.

The conjunction and complementation of the above-mentioned populations was also studied with cells that responded around voluntary motion onset and offset events when there was no sensory stimulation. During the no stimulus intervals of the NB-condition, animals sometimes walked or ran spontaneously. These events were identified and subdivided into components in which an animal started locomoting (motion onset, MOn) or stopped locomoting (motion offset, MOff). For 5 animals, the number of MOn events detected was 8, 21, 19, 20, and 19, and the number of MOff events was 35, 24, 31, 37, and 39. Cellular activity around these events (1.5 s pre- and post-event) was assessed by generating raster plots and PETHs to identify voluntary motion-related cellular responses. Both Exc and Inh cell responses (Fig. 6A) were identified. Rate vector maps (Fig. 6B, top row) and population correlation plots (Fig. 6B, middle and bottom rows) showed the binary nature of the responses. The effect of Event-Type (MOn vs. MOff) and Cell-Type (Exc vs. Inh) on the percentages of responsive cells was examined with a RM-ANOVA. There was a significant interaction effect [$F_{(1,4)} = 45.73$, $p =$

.002, $\eta^2$ = .92], with a greater number of cells showing an Exc than an Inh response to MOn ($p$ = .025), and vice versa for MOff ($p$ = .002) (Fig. 6C). The same analysis for response fidelity revealed a significant effect of Event-Type (Fig. 6D) [$F$ (1,4) = 8.12, $p$ = .046, $\eta^2$ = .67]. The fidelity of the cells responding to MOn was significantly greater than that of the cells responding to MOff. There was no significant main effect or interaction between factors for zMI values (Fig. 6E). These results suggest more cells responded during locomotion and these cells have a more robust response to MOn.

**Figure 6.** Separate cell groups at movement onset (MOn) and offset (MOff) events compared to other sensorimotor events. (**A**) Raster plots of representative Exc and Inh cells to MOn. The magenta line at t = 0 indicates motion onset. The blue lines show means over trials. (**B**) Rate vector maps (top row) and corresponding population vector correlation plots (middle row) of responsive cells from a representative animal for MOn and MOff events. Average population correlation plots (bottom row) for 5 animals. (**C-E**) Results of two-way RM-ANOVA for assessing responsivity, response fidelity, and mutual information z-score for MOn vs. MOff and Exc vs. Inh cells. (**F**) Overall responsivity, conjunction, and complementation for Brake, No-Brake, and Motion conditions. (**G**) Results of two-way RM-ANOVA for assessing the percent of cells across population types (Comp1, Conj, and Comp2) and condition pairs e.g., Brake and Motion (B-M). (**H-I**) Dendrograms resulting from agglomerative hierarchical clustering of the average conjunction and complementation heatmaps/matrices when MOn and MOff events were included. CC indicates cophenetic correlation. For all statistical tests, $n$ = 5 mice. Error bars = SEM. *$p$ < .05, **$p$ < .01.

To examine whether cells exhibiting responses to MOn and Moff (M-condition) responded to events during the B-condition and non-voluntary motion events in the NB-condition, cells were pooled across events (AOn, AOff, Arb, L, A, and A-L), response types (Exc vs. Inh), and stimulus types (air and light). Overall, there were $78.61 \pm 1.69$ % responsive cells with $8.8 \pm 0.4$ %, $30.8 \pm 1.4$ %, and $4.2 \pm 0.3$ % active in B-, NB-, and M- conditions, respectively and $6.9 \pm 0.9$% responding during all three conditions (Fig. 6F). When taken collectively, the percentage of complementary cells in the NB-condition was significantly larger than in the B and M conditions ($p < .001$ for both comparisons) whereas the percentage in the B-condition was larger than that in M-condition ($p = .003$): (Fig. 6F) [RM-ANOVA, $F (2,8) = 210.72$, $p < .001$, $\eta^2 = .98$]. When taken two at a time, the percent of complementary and conjunctive cells were compared by using a two-way RM-ANOVA with Condition-Pair e.g., B-M, NB-M, and Population-Type (Conj, Comp1, and Comp2) as within-subjects factors. There was a significant interaction between the two factors [$F (4,16) = 210.78$, $p < .001$, $\eta^2 = .98$]. Post hoc comparisons revealed that for the B-M comparison, the percentage of conjunctive cells was significantly smaller than the percentage of complementary cells (Fig. 6G, $p = .012$ and $p < .001$). However, there was a larger overlap of cells between NB and M conditions with a larger percent of complementary cells in the NB-condition (Fig. 6G, $p = .001$, $.0003$, and $.0047$).

The conjunction and complementation of cells responding to MOn and MOff events was also analyzed with all other sensorimotor events in B and NB conditions. Cell populations for MOn and MOff were closer to AOn and AOff events respectively in NB-condition revealed from clustering of conjunction heatmap (Fig. 6H). The clustering of complementation heatmaps showed that MOn and MOff formed a separate cluster which suggested that the complementary cells related to voluntary motion were distinct compared to other cell populations.

Taken together, these results suggest that there are separate cell populations which are active for distinct sensorimotor events in different behavioral configurations. Where the conjunctive cell populations indicated stable cellular representations, complementary cell populations featured the dynamics of recruitment across different sensorimotor events. This possibility was analyzed next by doing a trial-by-trial analyses on the activity of cells.

**Trial-wise turnover of cellular recruitment reveals larger a percent of complementary cells for No-Brake condition in early trials**

To examine how cells were recruited during individual trials during the presentation of air, light, and arbitrary events in relation to the B and NB conditions, responsive cells (firing rate > 0) were identified for each trial. The percent of responsive cells in each trial were found for the 20 different sensorimotor events (see Fig. 7A). Across all trials for all events, 99.7 ± 0.1% of cells were responsive. Nevertheless, for any given event (Fig. 7A, magenta line), average of 17.3 ± 0.1% (range: 13.07, 23.02, median: 17.34) of cells were responsive in any given event. To statistically compare the percentage of responsive cells across the 20 events and 10 trials, a two-way RM-ANOVA was used. There was a significant effect of Event-Type on the percent of responsive cells (Fig. 7C), [$F$ (19,76) = 2.10, $p$ = .012, $\eta^2$ = .34]. Post hoc comparisons revealed differences in means for a few events (Fig. 7C). Consistent with previous observations, a larger percent of cells responded to sensory stimuli during the NB-condition and there were abrupt increases in the percent of responsive cells at Brake to No-Brake transition, the condition permitting running onset in Configuration 3.

The dynamics of the percent of complementary and conjunctive cells was also assessed by considering adjacent pairs of trials (trials 1 and 2, trials 2 and 3, etc.). For the 9 pairs of trials for each of the 20 events, Conj, Comp1, and Comp2 cells were identified (Fig. 7B). Where Conj cells were active in

both trials in a trial-pair, Comp1 cells were active in the former trial but became silent in the later trial, and Comp2 cells started silent but became active in the later trial.

**Figure 7.** Adjacent trial-wise dynamics of cellular responses and conjunction/complementation in cellular populations. **(A)** Percent of responsive cells (firing rate > 0) over trials for 20 types of cellular responses to different sensorimotor events exhibited across the 7 behavioral configurations. **(B)** Percent of complementary and conjunctive cells for pairs of adjacent trials. Pairs 1-9 correspond to Trials 1-2, 2-3, 3-4, etc. up to 9-10. **(C)** Result of two-way RM-ANOVA assessing responsivity across 20 events and 10 trials. **(D-F)** Results of two-way RM-ANOVA assessing the percent of conjunction and complementation across 20 events and 9 trial-pairs. For all statistical tests, $n = 5$ mice. Error bars = SEM. *$p < .05$, **$p < .01$.

A two-way RM-ANOVA with Event-Type and Trial-Pairs as within-subjects factors of the percent of Conj cells gave a significant effect of Event-Type [$F_{(19,76)} = 5.87$, $p < 0.001$, $\eta^2 = .59$]. Post

hoc tests revealed significantly smaller percentages during Arb events (Fig. 7D, see significant post hoc comparisons). Similar analysis on the percent of complementary cells showed that percent of both Comp1 and Comp2 cells were significantly different across Event-Types: [$F$ (19,76) = 2.81, $p < 0.001$, $\eta^2 = .41$] and [$F$ (19,76) = 2.82, $p < 0.001$, $\eta^2 = .41$. Post hoc comparisons revealed a larger percent of both types during NB-condition compared to B-condition (Fig. 7E-F, see significant post hoc comparisons). The percent of Comp2 cells were also different across trial-pairs [$F$ (8,32) = 2.41, $p = .037$, $\eta^2 = .38$]. Post hoc comparisons did not show any differences in means across trial-pairs. However, if only air and light stimuli were considered, the percent of Comp2 cells was significantly different across events [$F$ (12,48) = 3.79, $p = .0005$, $\eta^2 = .49$] as well as trial-pairs [$F$ (8,32) = 2.76, $p = .019$, $\eta^2 = .41$] with trial-pair 1-2 having a larger percentage of cells compared to trial-pair 9-10 (post hoc comparison, $p = .021$). These results suggest locomotion contributed to more new cells being recruited immediately following a new event (experience).

The conjunction and complementation of cells for all trial-pairs (200 × 200) corresponding to 20 events and 10 trials were qualitatively assessed with heatmaps. The average heat maps for conjunctive and complementary cells are shown in Fig. 8. The heatmap of conjunctive cells (Fig. 8A) shows that there are separate subgroups of cells that are selectively active during different events. For example, see the checkerboard-like pattern in the middle of the map outlined by the thick brown box, which shows that separate subgroups of cells respond to AOn versus AOff (outlined with red boxes). Furthermore, some of the cells that participated during the first B-condition also did so selectively during the second B-condition (outlined with purple boxes). Cells that responded to light during the NB-condition had a high overlap with cells that responded to AOff during the NB-condition. To perform clustering on the conjunction heatmap, it was binned to average across trial-pairs i.e., all 100 pixels corresponding to 10 x 10 trial-pairs within each big black box were averaged. This way an average heatmap was obtained and

subjected to agglomerative hierarchical clustering. It showed that there was a clear segregation within

the dendrograms for the B versus NB conditions (Fig. 8C, dotted green box versus other branches).

Further segregation on branch level can be seen for grouping of sensory and arbitrary events (blue and

purple dotted boxes and cyan cluster). The segregation between B and NB conditions was clearer when

Arb and A events were excluded (Fig. 8F).

**Figure 8.** Trial-pair-wise dynamics of conjunction and complementation of cellular populations for all 20 sensorimotor events. (**A**) Heatmap of the average percentage of conjunctive cells (n = 5 mice) between trials from all 20 events. The 200 × 200 pixels (off diagonal) belong to trial-pairs from all 20 events, as shown by the x and y labels. Each big box enclosed by thin black lines shows 10 × 10 trial pairs of the given event. For example, the color value of the pixel in row 1, column 2 indicates the

percentage of conjunctive cells for trial 1 and 2 of presentation of the light stimulus. **(B)** Heatmap of the average percentages of complementary cells (n = 5 mice) between trial pairs (like A). Here, for example, the color value of the pixel in row 1, column 2 indicates the percentage of cells present in trial 1 and not present in trial 2. **(C-E)** Dendrograms resulting from agglomerative hierarchical clustering of the average conjunction and complementation heatmaps/matrices. **(F-H)** Dendrograms resulting from agglomerative hierarchical clustering of the average conjunction and complementation heatmaps/matrices after excluding arbitrary events. CC indicates cophenetic correlation.

The heatmap of complementary cells (Fig. 8B) revealed a larger complementation of the NB vs. B conditions (outlined with purple boxes). Furthermore, the checkerboard-like pattern in the middle region of the heatmap suggests selective activation during AOn versus AOff during the NB-condition. Clustering of the complementary heatmaps (Comp1 and Comp2) was done in a similar manner as described above after averaging over trial-pairs and separating the upper and lower triangular regions. Here, the clustering revealed segregation of B and NB conditions but with some mixing owing to the ongoing recruitment of complementary cells that were similar across events or Configurations which happened close in time. The exclusion of Arb and A events before clustering revealed similar results (Fig. 8G-H).

In summary, the results from the trial-by-trial analysis were consistent with findings on the responsiveness of cells averaging over trials. More cells were recruited during transition from the B to NB-condition with overall larger complementation for the NB-condition. Furthermore, complementation was larger in the initial trials (novel experience) and became smaller as the experience was repeated. Additionally, distinct smaller groups of cells encoded distinct sensorimotor events.

## Discussion

Calcium imaging documented the activity of hippocampal CA1 cells and revealed different individual and collective cell population responses to sensory stimulation during immobility, locomotion and the transitions between these behaviors. Head fixed mice stood on a conveyor belt and immobility

was encouraged by activating a conveyor belt brake, whereas locomotion could occur when the brake was released. Light and tactile air stimuli were systematically applied during B and NB conditions. Neuronal excitation or inhibition featured a salt and pepper arrangement suggesting that responsivity was not topographically organized and did not reflect a general function such as immobility vs locomotion or arousal. Overall, ~99% of 2083 cells were active within the imaging session, with ~78% responding in relation to individualized experimental sensory/movement events. On a given trial, combinations of sensory and locomotor events featured subpopulations of active neurons comprising about ~17% of all detected neurons organized in complementary and conjunctive patterns of activation. From a large population of cells, complementary and conjunctive cells encode relatively novel and familiar experiences respectively. These results suggest that hippocampal CA1 cells participate in functional networks that distinguish sensory events in relation to ongoing moment-to-moment behavior using a recruitment with replacement strategy.

A central finding of the present study was the individuation of cell responses to events featuring configurations of light, air, locomotion, and immobility. By observing cellular responses to different sensory stimuli during multiple behavioral states in a single paradigm, this study describes the sensorimotor relations of an unprecedented 99% responsive cells. This responsivity could be attributed to the bias of Suite2P software in identifying only active cells. However, overall ~78% of cells (determined from average responses) were tuned to the experimentally determined stimuli and motion events, with a larger percentage of cells active during locomotion vs immobility. Previous studies have reported locomotion-related responses including sensory-spatial [18,19,24,25,27], sensory-temporal [21,22], or spatial-temporal [50-52] populations with smaller percentages or limited numbers of active cells. For example, [21] reported sensory-temporal representations in up to 30% active cells, [53] reported responses related to 11 different rooms of 30 active cells per animal (freely behaving), and [54] used a 3D

environment and described 756 place cells from 13 rats. That the percentage of cells responsive in all conditions in the present study was negligible argues that to fully sample hippocampal cell diversity, it is necessary to study the activity of cells during an array of sensorimotor conditions.

Although the percent (~7%) of conjunctive cells; that is cells that were active to specific sensory-motor configurations such that C ∈ A and B, was less variable across conditions of immobility and locomotion, the percentages of complementary cells; that is cells that were active to a specific stimulus or condition such that C ∈ A not B or C ∈ B not A, were larger during locomotion. Moreover, complementary cells varied with experience (in later compared to former trials), with a larger percentage occurring in the first few trials. This observation supports the idea that ongoing cellular recruitment might depend on experience as well as moment-to-moment behavior, as novel experiences become familiar experiences. The existence of conjunctive populations across similar Configurations (e.g., 2 and 7 OR 3, 4, and 5), additionally argues that familiarity is also encoded in ongoing cell activity.

For both immobility and locomotor related cellular responses, the tactile air stimulus was more potent in eliciting sensory responses than was the visual light stimulus. There were also more cells, with higher response fidelity and stability that respond to air compared to light stimulus. This is consistent with a finding in rabbits reporting that repetitive tactile stimulation is more potent than repetitive visual stimulation in modulating hippocampal field potentials [12]. Here, cell responsivity could have been influenced by many factors including the shorter duration of the light stimulus (200 ms) compared to the air stimulus (5 s), differences in behavioral responses of "freezing" to the stimulus, differences in autonomic responses to the stimuli, including respiratory and pupil response changes or slight body movements by the animals. Future experiments using videography combined with calcium imaging can address these possibilities. For locomotion related cell responses, we cannot rule out higher order responses to sensory stimulation or behavior e.g., cellular responses to an AOff event might have some

contributions from a previous air or light onset event or vice versa. It should be noted that although we did not record behavioral concurrent field potentials a substantial literature reports that they change with transitions between immobility and movement (see introduction).

The cell/behavior relationship described in the present study provides insights into the sensorimotor functions of the hippocampus as a neural network. First, almost no cells were consistently active in association with all sensorimotor events, as might be expected in a system related to general arousal [55] or reflecting only ongoing movement [1]. Second, cell responses were not topographically organized as is found in primary sensory and motor regions of the neocortex, a finding consistent with the idea that the hippocampus is a network system [56]. Third, cells obtain specificity with respect to sensorimotor behaviors by featuring complementary and conjunctive subpopulations. For example, a subpopulation of cells might respond in relation to a tactile air stimulus when a mouse is relatively immobile, whereas a different, complementary subpopulation might respond to the same stimulus when the mouse is moving, and a smaller, conjunctive population might respond to the stimulus in both conditions. Fourth, complementary and conjunctive subpopulation cell numbers appeared relatively fixed around average numbers of 10% and 7%, respectively.

In summary, the complementary/conjunctive specificity of CA1 cells suggests that the hippocampus is a network that encodes ongoing sensorimotor events related to concurrent movement. As such cell-behavior relations are congruent with reports of the relationship between hippocampal field potentials and behavior. Given this complementary and conjunctive cell subpopulation organization it is not surprising that subpopulations of cells are related to features of spatial behavior [15,16], episodic learning and memory [57,58], context representation [47,59,60], and scene construction [61] as the hippocampus is downstream from the many cortical and subcortical systems that are also involved in these processes.

## Methods

### Animals and surgery

Animal procedures were performed in compliance with protocols approved by the Animal Welfare Committee of the University of Lethbridge and followed the guidelines for the ethical use of animals provided by the Canadian Council on Animal Care. In total 5 mice, 3 male and 2 female Thy1-GCaMP6s [C57BL/6J-Tg(Thy1-GCaMP6s)GP4.3Dkim/J] (JAX stock #024275) hemizygous transgenic mice were used. This mouse line exhibits stable and homogenous expression of GCaMP6s over a large population of cortical and subcortical excitatory neurons [62,63]. After cranial window surgery mice were housed in individually ventilated Optimice cages (Animal Care Systems) in temperature-controlled rooms (22 °C) maintained on a 12:12 light/dark cycle (lights on during the day). Food and water were provided *ad libitum*.

Mice were implanted with chronic cranial windows at ~5-7 months of age. Mice were injected subcutaneously with buprenorphine (0.075 mg/kg) 0.5 h before surgery. Immediately prior to surgery, animals were also injected subcutaneously with 0.5 ml of a 5% dextrose, 0.9% saline solution mixed with atropine (3 µg/ml). Animals were anesthetized with isoflurane (2-3% for induction, 1-1.5% for anesthesia) and their body temperature was maintained at 37 °C using a homoeothermic monitoring system and heating pad. Lidocaine (10 mg/kg) was injected under the scalp as a local anesthetic, and a section of skin was removed with scissors to expose the skull. A craniotomy, 3 mm in diameter, was made above the right hippocampus (center at -2.0 mm AP, 1.8 mm ML, relative to bregma). The dura underlying the craniotomy was resected, and a small region of neocortex was removed by aspiration, exposing the underlying white matter. A metal cylinder (3 mm outer diameter, 1.5 mm long) with a glass coverslip attached to the bottom end using optical adhesive (NOA71, Norland) was inserted into the cavity. The coverslip was gently pushed against the tissue to reduce motion, and the cylinder was

attached to the skull using cyanoacrylate tissue adhesive (Vetbond, 3M) and dental acrylic. To allow for

head-fixation of the mouse, a titanium head-plate was attached to the skull using adhesive cement (C&B

Metabond, Parkell). Two rubber rings were attached to the top of the head-plate, forming a well that

could be filled with distilled water for imaging with a water-immersion objective. Post-surgery,

subcutaneous injections of meloxicam (7.5 mg/kg) and enrofloxacin (10 mg/kg) were administered daily

for 2-3 days. Animals were allowed to recover for 3-8 weeks before behavioral training was started. The

age of the animals at the time of imaging was ~8 months.

**Animal training and behavioral experiment**

Animals were habituated to head-fixation over 3 days. Mice were head-fixed twice per day, with

sessions gradually increasing in length from 5 to 30 min. Following habituation, animals were trained to

run on a linear conveyor belt 150cm in length while head fixed. The conveyor belt has been previously

described [64,65]. The conveyor belt was not motorized; all movement of the belt were generated by the

mouse. The belt was made from Velcro material (Country Brook). Training was done in the dark, on a

blank conveyor belt devoid of sensory cues. Belts were washed after every training session. Mice were

trained for 20-30 min each day to run in response to a continuous, mild air stream applied to the back. A

solenoid valve based automated system was used to apply a continuous mild air stream from the

building's compressed air supply via a regulator. To escape this stimulus, the animal was required to run

a fixed distance on the belt, after which the air puff ceased. Following an inter-trial interval of 15 s, the

air stream would turn on again, initiating the next training trial. The distance the mouse was required to

run was increased gradually across training trials, starting each day with 30 cm, and increasing to a

maximum of 150 cm (i.e., one full turn of the belt). On a given training trial, if the mouse ran the

required distance with an average speed greater than 7 cm/s, the required distance on the next trial was

incremented by 7-15 cm. If the mouse failed to achieve the target speed (e.g., due to running too slowly, delayed starting, or repeated starting and stopping), the required distance did not change between trials. Mice were trained until they reached the maximum distance of 150 cm and achieve the target speed on >75% of trials across a training session. All mice reached criterion in 3 training sessions.

After training, mice were tested (re-trained) 10-18 days later and the following day they experienced a 7-configuration sensorimotor behavioral paradigm while activity of hippocampal CA1 neurons was recorded by two-photon calcium imaging. Behavioural data were simultaneously acquired and synchronized with the imaging data using a data acquisition system (Axon Digidata 1322A). These data included the distance traveled on the belt (measured by rotation encoders attached to the shafts of the treadmill wheels). A fixed reference point on the belt was also tracked, using reflective tape attached to the underside of the belt, which activated a photosensor once per belt lap. Air puff onset and offset, as well as brake and visual cue onset and offset, were controlled by a microprocessor board (Arduino Mega). The results of air training and analyses of distance and/or duration tuning from air onset and offset in Configurations 3, 4, and 5 as well as additional analyses will be presented elsewhere.

**Two-photon calcium imaging**

A multiphoton microscopy system (Bergamo II, Thorlabs) was used for two-photon calcium imaging. A Ti:Sapphire excitation laser (Coherent) was tuned to 920 nm, with a power of ~100 mW at the sample. Laser scanning was controlled by galvo and resonant scanners. A GaAsP photomultiplier tube was used to detect the green fluorescence signal from GCaMP6s. Samples were imaged through a 16×/0.8 NA water-immersion objective (Nikon). A black fabric cover was wrapped around the objective and head-plate to block contaminating light. For two mice, images of hippocampal CA1 pyramidal neurons were captured from two planes, with the dorsal plane ~125 µm beneath the tissue surface and

the second plane 50 µm deeper. Images were gathered at a frame rate of 29.16 Hz, but the effective

frame rate for each individual plane was 9.72 Hz (29.16 ÷ 3, because 1 extra fly-back frame was also

captured). For three mice images were gathered from only one plane with a frame rate of 29.16Hz. A

field of view of $418 \times 418$ µm was captured with a resolution of $512 \times 512$ pixels.

**Data analysis**

Suite2P [66] (http://mouseland.github.io/suite2p) was used to register images and to automatically

detect regions of interest (ROIs) assumed to represent individual cell bodies. Raw calcium traces were

determined for each ROI by averaging the fluorescence signal from all pixels within the ROI and

correcting for neuropil contamination. For each raw calcium fluorescence trace (F), baseline ($F_o$) was

estimated, and baseline subtracted signal ($\Delta F/F_o$) was then determined as ($F-F_o$)/$F_o$. From the $\Delta F/F_o$

calcium traces, deconvolved spike rates were determined using a first order auto-regressive model and

constrained nonnegative matrix factorization [67]. Further analysis of the deconvolved traces was

conducted using custom scripts written in MATLAB (MathWorks).

To make the raster plots, a time bin width of 0.11s was used (36 bins for 4 s) and the average

firing rate was calculated for each time bin. Response fidelity (RF) of a neuron for a raster was defined

as the percentage of trials in which the firing rate was greater than zero for any time bin. Mutual

information (MI) between firing rate and time was calculated for each raster using firing rate values

from all trials by binning them into four non-overlapping quantiles[49] using the following relation

$$MI = \sum_{i=1}^{Nb} \sum_{j=1}^{4} p_{ij} \log_2 \left( \frac{p_{ij}}{p_i \cdot p_j} \right)$$

where $p_i$ and $p_j$ are the probabilities of time bin $i$ and firing rate bin $j$, respectively; $p_{ij}$ is the joint

probability between time bin $i$ and firing rate bin $j$; $Nb$ is the number of time bins i.e., 36.

Peri-event time histogram (PETH) was obtained by averaging the response in the rasters over trials and was used to determine responsiveness of a cell using a two-tailed, two-sample, t-test comparing pre- and post-event firing rates (alpha value 0.05). Rate vector maps were obtained by plotting PETH of all cells after normalizing with the peak of response. The population vector correlation was obtained by finding the Pearson correlation between all pairs of the columns of the rate vector map i.e., each pixel shows correlation between two columns of rate vector map and hence the matrix is symmetric across the diagonal.

**Agglomerative Hierarchical Clustering**

This analysis was performed on the conjunction or complementation matrices (heatmaps) when multiple cell groups were considered. Complementation matrices were split into two heatmaps symmetric across the diagonal by using the upper and lower triangular matrices corresponding to Comp1 and Comp2 types of cells before clustering. Based on the conjunction and complementation percentages of each cell group with every other cell group, this analysis found sensorimotor events which were like each other i.e., close to each other with respect to their conjunction/complementation relationships with all other cell groups. The analysis was done by first calculating Euclidian distance for each pair of cell groups considering two rows of a matrix at a time (using Matlab function "pdist"). For example, if a matrix was n x n, there would be $\frac{n!}{2!(n-2)!}$ pairs which can then be arranged into a symmetric n x n matrix usually called the dissimilarity matrix owing to smaller Euclidian distance indicating smaller dissimilarity. Matlab function "linkage" was then used to operate on the dissimilarity matrix and define clusters based on the unweighted average distance between clusters. Clusters were then separated (color coded in figures using the Matlab function "dendrogram") by using a threshold of 70% of the maximum

linkage distance between any two clusters. The quality of clustering was assessed by determining cophenetic correlation (CC) reported with each dendrogram in figures.

**Statistical Analysis**

Statistical analysis was done in Matlab® 2018b (Ver 9.5.0.944444) and confirmed for accuracy with IBM® SPSS® (Ver 28.0.1.1). All means in the text are reported ± the standard error of the mean (SEM). Unless otherwise specified, a repeated measures analysis of variance (RM-ANOVA) test was used for statistically comparing differences in measured parameters from n = 5 animals. Shapiro-Wilk test was used for test of normality and was done using a file "swtest.m" downloaded from Matlab File Exchange Server [68]. The accuracy of the code was confirmed by comparing its output with similar results from SPSS. Minor violations of the normality test and the presence of outliers were ignored e.g., if some variables did not pass the normality test or had outliers, RM-ANOVA was still performed. The test for sphericity was conducted using Mauchly's Test. Greenhouse-Geisser corrected $p$-value is reported where sphericity was violated. All post hoc pairwise comparisons were conducted either using Tukey's Honestly Significant Difference (HSD) test or Bonferroni correction with an alpha = 0.05.

An alpha value of .05 was used to determine significance at the first level of analysis e.g., for performing a three-way RM-ANOVA. For the second level of analysis, it was adjusted with Bonferroni correction e.g., if there was a significant three-way interaction, in the second level of analysis to find simple two-way interactions, alpha was divided by the number of statistical tests performed. For example, if there were two simple two-way interactions to be tested, an alpha value of 0.05/2 = 0.025 was used. Similarly, at the next level of analysis, alpha was further divided by the number of statistical comparisons made e.g., if there were three simple simple main effects to be tested, alpha = 0.025/3 = 0.0083 was used.  *, **, and *** indicate $p$-values less than .05, .01, and .001, respectively. Actual $p$-

values are reported except for values less than .001 are reported as p < .001. Because of the small sample size (n = 5 animals), effect size is reported along with *p*-values as partial $\eta^2$ = (Sum of Squares $_{effect}$)/(Sum of Squares $_{effect}$ + Sum of Squares $_{error}$).

## Acknowledgements

We thank Dr. JianJun Sun for performing animal surgeries, Di Shao and the University of Lethbridge Animal Care Services staff for animal husbandry, and HaoRan Chang, Drs. Bruce McNaughton, Javad Karimi, Hardeep Ryait, and Žaneta Navrátilová for useful discussions regarding the preparation of this manuscript. We thank you Dr. Bryan Souza for providing Matlab Code for calculating mutual information.

## Author Contributions

All authors participated in the design of this study, with S.I. and I.Q.W. as the main contributors. S.I. and B.B.M. performed the experiments. S.I. performed the data analyses with participation of the other authors. S.I. and I.Q.W. wrote the manuscript, which all authors commented on and edited. M.H.M. supervised the study.

## Competing Interests

The authors declare no competing interests.

## References

Vanderwolf, C. H. Hippocampal electrical activity and voluntary movement in the rat. *Electroencephalogr Clin Neurophysiol* **26**, 407-418, doi:10.1016/0013-4694(69)90092-3 (1969).

Kay, K. & Frank, L. M. Three brain states in the hippocampus and cortex. *Hippocampus* **29**, 184-238, doi:10.1002/hipo.22956 (2019).

Whishaw, I. Q. & Vanderwolf, C. H. Hippocampal EEG and behavior: effects of variation in body temperature and relation of EEG to vibrissae movement, swimming and shivering. *Physiol Behav* **6**, 391-397, doi:10.1016/0031-9384(71)90172-7 (1971).

Whishaw, I. Q. & Vanderwolf, C. H. Hippocampal EEG and behavior: changes in amplitude and frequency of RSA (theta rhythm) associated with spontaneous and learned movement patterns in rats and cats. *Behav Biol* **8**, 461-484, doi:10.1016/s0091-6773(73)80041-0 (1973).

Buzsaki, G. & Watson, B. O. Brain rhythms and neural syntax: implications for efficient coding of cognitive content and neuropsychiatric disease. *Dialogues Clin Neurosci* **14**, 345-367 (2012).

Buzsáki, G. *Rhythms of the brain*.  (Oxford University Press, 2006).

Foster, T. C., Castro, C. A. & McNaughton, B. L. Spatial selectivity of rat hippocampal neurons: dependence on preparedness for movement. *Science* **244**, 1580-1582, doi:10.1126/science.2740902 (1989).

Whishaw, I. Q. The effects of alcohol and atropine on EEG and behavior in the rabbit. *Psychopharmacology (Berl)* **48**, 83-90, doi:10.1007/BF00423311 (1976).

Perentos, N., Krstulovic, M. & Morton, A. J. Deep brain electrophysiology in freely moving sheep. *Current biology : CB* **32**, 763-774 e764, doi:10.1016/j.cub.2021.12.035 (2022).

Kramis, R., Vanderwolf, C. H. & Bland, B. H. Two types of hippocampal rhythmical slow activity in both the rabbit and the rat: relations to behavior and effects of atropine, diethyl ether, urethane, and pentobarbital. *Exp Neurol* **49**, 58-85, doi:10.1016/0014-4886(75)90195-8 (1975).

Vanderwolf, C. H., Gutman, M. & Baker, G. B. Hypothalamic self-stimulation: the role of dopamine and possible relations to neocortical slow wave activity. *Behavioural brain research* **12**, 9-19, doi:10.1016/0166-4328(84)90198-0 (1984).

Whishaw, I. Q. & Dyck, R. Comparative potency of tactile, auditory, and visual stimulus repetition in eliciting activated forebrain EEG in the rabbit. *Behavioral neuroscience* **98**, 333-344, doi:10.1037//0735-7044.98.2.333 (1984).

Yu, J. Y. *et al.* Distinct hippocampal-cortical memory representations for experiences associated with movement versus immobility. *Elife* **6**, doi:10.7554/eLife.27621 (2017).

Kay, K. *et al.* A hippocampal network for spatial coding during immobility and sleep. *Nature* **531**, 185-190, doi:10.1038/nature17144 (2016).

O'Keefe, J. & Dostrovsky, J. The hippocampus as a spatial map. Preliminary evidence from unit activity in the freely-moving rat. *Brain research* **34**, 171-175, doi:10.1016/0006-8993(71)90358-1 (1971).

O'Keefe, J. & Nadel, L. *The Hippocampus as a Cognitive Map*.  (Oxford: Clarendon Press, 1978).

Arriaga, M. & Han, E. B. Dedicated Hippocampal Inhibitory Networks for Locomotion and Immobility. *The Journal of neuroscience : the official journal of the Society for Neuroscience* **37**, 9222-9238, doi:10.1523/JNEUROSCI.1076-17.2017 (2017).

Komorowski, R. W., Manns, J. R. & Eichenbaum, H. Robust conjunctive item-place coding by hippocampal neurons parallels learning what happens where. *The Journal of neuroscience : the official journal of the Society for Neuroscience* **29**, 9918-9929, doi:10.1523/JNEUROSCI.1378-09.2009 (2009).

Manns, J. R., Howard, M. W. & Eichenbaum, H. Gradual changes in hippocampal activity support remembering the order of events. *Neuron* **56**, 530-540, doi:10.1016/j.neuron.2007.08.017 (2007).

Igarashi, K. M., Lu, L., Colgin, L. L., Moser, M. B. & Moser, E. I. Coordination of entorhinal-hippocampal ensemble activity during associative learning. *Nature* **510**, 143-147, doi:10.1038/nature13162 (2014).

Taxidis, J. *et al.* Differential Emergence and Stability of Sensory and Temporal Representations in Context-Specific Hippocampal Sequences. *Neuron* **108**, 984-998 e989, doi:10.1016/j.neuron.2020.08.028 (2020).

MacDonald, C. J., Carrow, S., Place, R. & Eichenbaum, H. Distinct hippocampal time cell sequences represent odor memories in immobilized rats. *The Journal of neuroscience : the official journal of the Society for Neuroscience* **33**, 14607-14616, doi:10.1523/JNEUROSCI.1537-13.2013 (2013).

Sakurai, Y. Coding of auditory temporal and pitch information by hippocampal individual cells and cell assemblies in the rat. *Neuroscience* **115**, 1153-1163, doi:10.1016/s0306-4522(02)00509-2 (2002).

Moita, M. A., Rosis, S., Zhou, Y., LeDoux, J. E. & Blair, H. T. Hippocampal place cells acquire location-specific responses to the conditioned stimulus during auditory fear conditioning. *Neuron* **37**, 485-497, doi:10.1016/s0896-6273(03)00033-3 (2003).

Herzog, L. E. *et al.* Interaction of Taste and Place Coding in the Hippocampus. *The Journal of neuroscience : the official journal of the Society for Neuroscience* **39**, 3057-3069, doi:10.1523/JNEUROSCI.2478-18.2019 (2019).

Itskov, P. M., Vinnik, E. & Diamond, M. E. Hippocampal representation of touch-guided behavior in rats: persistent and independent traces of stimulus and reward location. *PloS one* **6**, e16462, doi:10.1371/journal.pone.0016462 (2011).

Zhao, X., Wang, Y., Spruston, N. & Magee, J. C. Membrane potential dynamics underlying context-dependent sensory responses in the hippocampus. *Nature neuroscience* **23**, 881-891, doi:10.1038/s41593-020-0646-2 (2020).

Chen, G., King, J. A., Burgess, N. & O'Keefe, J. How vision and movement combine in the hippocampal place code. *Proceedings of the National Academy of Sciences of the United States of America* **110**, 378-383, doi:10.1073/pnas.1215834110 (2013).

Wang, Y., Feng, Z., Wang, J. & Zheng, X. Somatosensory stimulation suppresses the excitability of pyramidal cells in the hippocampal CA1 region in rats. *Neural Regen Res* **9**, 1138-1144, doi:10.4103/1673-5374.135316 (2014).

Bellistri, E., Aguilar, J., Brotons-Mas, J. R., Foffani, G. & de la Prida, L. M. Basic properties of somatosensory-evoked responses in the dorsal hippocampus of the rat. *The Journal of physiology* **591**, 2667-2686, doi:10.1113/jphysiol.2013.251892 (2013).

Smith, P. F. Vestibular-hippocampal interactions. *Hippocampus* **7**, 465-471, doi:10.1002/(SICI)1098-1063(1997)7:5<465::AID-HIPO3>3.0.CO;2-G (1997).

Schneider, D. M. Reflections of action in sensory cortex. *Current opinion in neurobiology* **64**, 53-59, doi:10.1016/j.conb.2020.02.004 (2020).

Dipoppa, M. *et al.* Vision and Locomotion Shape the Interactions between Neuron Types in Mouse Visual Cortex. *Neuron* **98**, 602-615 e608, doi:10.1016/j.neuron.2018.03.037 (2018).

Niell, C. M. & Stryker, M. P. Modulation of visual responses by behavioral state in mouse visual cortex. *Neuron* **65**, 472-479, doi:10.1016/j.neuron.2010.01.033 (2010).

Ayaz, A. *et al.* Layer-specific integration of locomotion and sensory information in mouse barrel cortex. *Nature communications* **10**, 2585, doi:10.1038/s41467-019-10564-8 (2019).

Sofroniew, N. J., Vlasov, Y. A., Hires, S. A., Freeman, J. & Svoboda, K. Neural coding in barrel cortex during whisker-guided locomotion. *Elife* **4**, doi:10.7554/eLife.12559 (2015).

Schneider, D. M., Nelson, A. & Mooney, R. A synaptic and circuit basis for corollary discharge in the auditory cortex. *Nature* **513**, 189-194, doi:10.1038/nature13724 (2014).

Mineault, P. J., Tring, E., Trachtenberg, J. T. & Ringach, D. L. Enhanced Spatial Resolution During Locomotion and Heightened Attention in Mouse Primary Visual Cortex. *The Journal of neuroscience : the official journal of the Society for Neuroscience* **36**, 6382-6392, doi:10.1523/JNEUROSCI.0430-16.2016 (2016).

Saleem, A. B., Diamanti, E. M., Fournier, J., Harris, K. D. & Carandini, M. Coherent encoding of subjective spatial position in visual cortex and hippocampus. *Nature* **562**, 124-127, doi:10.1038/s41586-018-0516-1 (2018).

Clancy, K. B., Orsolic, I. & Mrsic-Flogel, T. D. Locomotion-dependent remapping of distributed cortical networks. *Nature neuroscience* **22**, 778-786, doi:10.1038/s41593-019-0357-8 (2019).

Stringer, C. *et al.* Spontaneous Behaviors Drive Multidimensional, Brain-wide Activity. *Science* **364**, 255, doi:10.1126/science.aav7893 (2019).

Musall, S., Kaufman, M. T., Juavinett, A. L., Gluf, S. & Churchland, A. K. Single-trial neural dynamics are dominated by richly varied movements. *Nature neuroscience* **22**, 1677-1686, doi:10.1038/s41593-019-0502-4 (2019).

Keller, G. B., Bonhoeffer, T. & Hubener, M. Sensorimotor mismatch signals in primary visual cortex of the behaving mouse. *Neuron* **74**, 809-815, doi:10.1016/j.neuron.2012.03.040 (2012).

Attinger, A., Wang, B. & Keller, G. B. Visuomotor Coupling Shapes the Functional Development of Mouse Visual Cortex. *Cell* **169**, 1291-1302 e1214, doi:10.1016/j.cell.2017.05.023 (2017).

Moser, E. I., Moser, M. B. & McNaughton, B. L. Spatial representation in the hippocampal formation: a history. *Nature neuroscience* **20**, 1448-1464, doi:10.1038/nn.4653 (2017).

Colgin, L. L. Five Decades of Hippocampal Place Cells and EEG Rhythms in Behaving Rats. *The Journal of neuroscience : the official journal of the Society for Neuroscience* **40**, 54-60, doi:10.1523/JNEUROSCI.0741-19.2019 (2020).

Smith, D. M. & Bulkin, D. A. The form and function of hippocampal context representations. *Neuroscience and biobehavioral reviews* **40**, 52-61, doi:10.1016/j.neubiorev.2014.01.005 (2014).

Eichenbaum, H. *et al.* Hippocampus at 25. *Hippocampus* **26**, 1238-1249, doi:10.1002/hipo.22616 (2016).

Souza, B. C., Pavao, R., Belchior, H. & Tort, A. B. L. On Information Metrics for Spatial Coding. *Neuroscience* **375**, 62-73, doi:10.1016/j.neuroscience.2018.01.066 (2018).

Kraus, B. J., Robinson, R. J., 2nd, White, J. A., Eichenbaum, H. & Hasselmo, M. E. Hippocampal "time cells": time versus path integration. *Neuron* **78**, 1090-1101, doi:10.1016/j.neuron.2013.04.015 (2013).

Haimerl, C. *et al.* Internal representation of hippocampal neuronal population spans a time-distance continuum. *Proceedings of the National Academy of Sciences of the United States of America* **116**, 7477-7482, doi:10.1073/pnas.1718518116 (2019).

Villette, V., Malvache, A., Tressard, T., Dupuy, N. & Cossart, R. Internally Recurring Hippocampal Sequences as a Population Template of Spatiotemporal Information. *Neuron* **88**, 357-366, doi:10.1016/j.neuron.2015.09.052 (2015).

Alme, C. B. *et al.* Place cells in the hippocampus: eleven maps for eleven rooms. *Proceedings of the National Academy of Sciences of the United States of America* **111**, 18428-18435, doi:10.1073/pnas.1421056111 (2014).

Grieves, R. M. *et al.* The place-cell representation of volumetric space in rats. *Nature communications* **11**, 789, doi:10.1038/s41467-020-14611-7 (2020).

Green, J. D. & Arduini, A. A. Hippocampal electrical activity in arousal. *Journal of neurophysiology* **17**, 533-557, doi:10.1152/jn.1954.17.6.533 (1954).

Buzsaki, G. The hippocampo-neocortical dialogue. *Cerebral cortex (New York, N.Y. : 1991)* **6**, 81-92, doi:10.1093/cercor/6.2.81 (1996).

Eichenbaum, H. Hippocampus: cognitive processes and neural representations that underlie declarative memory. *Neuron* **44**, 109-120, doi:10.1016/j.neuron.2004.08.028 (2004).

Eichenbaum, H. & Cohen, N. J. Can we reconcile the declarative memory and spatial navigation views on hippocampal function? *Neuron* **83**, 764-770, doi:10.1016/j.neuron.2014.07.032 (2014).

Rudy, J. W. Context representations, context functions, and the parahippocampal-hippocampal system. *Learning & memory (Cold Spring Harbor, N.Y.)* **16**, 573-585, doi:10.1101/lm.1494409 (2009).

Bulkin, D. A., Sinclair, D. G., Law, L. M. & Smith, D. M. Hippocampal state transitions at the boundaries between trial epochs. *Hippocampus* **30**, 582-595, doi:10.1002/hipo.23180 (2020).

McCormick, C., Dalton, M. A., Zeidman, P. & Maguire, E. A. Characterising the hippocampal response to perception, construction and complexity. *Cortex* **137**, 1-17, doi:10.1016/j.cortex.2020.12.018 (2021).

Chen, T. W. *et al.* Ultrasensitive fluorescent proteins for imaging neuronal activity. *Nature* **499**, 295-300, doi:10.1038/nature12354 (2013).

Dana, H. *et al.* Thy1-GCaMP6 transgenic mice for neuronal population imaging in vivo. *PloS one* **9**, e108697, doi:10.1371/journal.pone.0108697 (2014).

Mao, D., Kandler, S., McNaughton, B. L. & Bonin, V. in *Nature communications* Vol. 8    (2017).

Mao, D. *et al.* Hippocampus-dependent emergence of spatial sequence coding in retrosplenial cortex. *Proceedings of the National Academy of Sciences of the United States of America* **115**, 8015-8018, doi:10.1073/pnas.1803224115 (2018).

Pachitariu, M. *et al.* Suite2p: beyond 10,000 neurons with standard two-photon microscopy. *bioRxiv*, doi:10.1101/061507 (2017).

Friedrich, J., Zhou, P. & Paninski, L. in *PLoS computational biology* Vol. 13    (2017).

BenSaïda, A. in *MATLAB Central File Exchange*    (MathWorks, MATLAB Central File Exchange, 2022).