IBD-based estimation of X chromosome effective population size with application to sex-specific demographic history

Ruoyi Cai,[1*] Brian L. Browning,[1,2] Sharon R. Browning[1*]

[1] Department of Biostatistics, University of Washington

[2] Division of Medical Genetics, Department of Medicine, University of Washington

[*] Address for correspondence: RC: rcai2@uw.edu, SRB: sguy@uw.edu

Abstract

Effective population size (N_e) in the recent past can be estimated through analysis of identity-by-

descent (IBD) segments. Several methods have been developed for estimating N_e from autosomal IBD

segments, but no such effort has been made with X chromosome IBD segments. In this work, we

propose a method to estimate the X chromosome effective population size from X chromosome IBD

segments. We show how to use the estimated autosome N_e and X chromosome N_e to estimate female

and male effective population sizes. We demonstrate the accuracy of our autosome and X chromosome

N_e estimation with simulated data, and we show that pronounced differences in female and male

effective population size can be detected. In analysis of UK Biobank and the TOPMed project data, we

obtain results that are consistent with equal female and male effective population sizes in the UK White

British and UK Indian populations and in an African American population.

Introduction

The effective size of a population (N_e) is defined as the number of breeding individuals in an idealized

randomly mating population that has the same expected value of a parameter of interest as the actual

population under consideration.[1] The effective population size is a fundamental parameter in population

genetics because it determines the strength of genetic drift and the efficacy of evolutionary forces such

as mutation, selection, and migration.[2] Previous studies have demonstrated that estimates of recent

effective population size can reveal aspects of a population's demographic history, such as past

population growth or bottleneck events.[3,4]

Identity-by-descent (IBD) segments can be used to estimate effective population size in the recent past.

IBD segments are haplotypes which two or more individuals have inherited from a common ancestor.

IBD segments end at positions where crossovers have occurred in the meioses between the common

ancestor and the descendant individuals. IBD segments for which the common ancestor is in the distant

past tend to be shorter than IBD segments from a recent common ancestor because there are more

meioses since the common ancestor on which crossovers can occur. The autosomes and the X

chromosome are both subject to recombination, making them both amenable to IBD segment

analysis.[5,6] Previous studies have developed methods for estimating recent effective population size

from autosomal IBD segments.[3,4,7] However, no such effort has been made with X chromosome IBD

segments.

The autosomes are equally influenced by female and male demographic processes. In contrast, the X

chromosome is influenced more strongly by female demographic processes than by male demographic

processes. This is because females have two copies of the X chromosome, while males have only one.

Thus, comparison of statistics from the X chromosome and from autosome data can be used to estimate

sex-specific parameters such as female and male effective population sizes.[6,8-16]

The standard Wright-Fisher model used to define the effective population size assumes equal numbers

of breeding females and males. To define female and male effective population sizes, we consider an

idealized Wright-Fisher population modified to allow for different numbers of females and males. The

female and male effective sizes (N_e^f and N_e^m) of a non-idealized population are the values that would

give the same rates of IBD on autosomes and sex chromosomes as the Wright-Fisher population with

the corresponding numbers of females and males.[17] The effective sex ratio (ESR) is the ratio of the

effective number of females to the sum of the effective number of females and the effective number of

males, $N_e^f/(N_e^f + N_e^m)$.[18]

In this work, we develop an IBD-based method to estimate the X chromosome effective population size.

We also show that estimated X chromosome N_e can be combined with estimated autosome N_e to

estimate the female and male effective population sizes and the effective sex ratio. Through simulation

studies, we validate the theoretical relationship between the autosome and X chromosome N_e, and we

show that our method can accurately estimate the autosome and X chromosome N_e in simulated

populations. We examine the application of X chromosome N_e to estimate sex-specific N_e and show

that pronounced differences in female and male effective population sizes can be detected. We use our

method to infer autosome and X chromosome N_e, as well as sex-specific N_e, for several human

populations.

Methods

Probability modelling for the X chromosome

All meioses from mothers transmit an X chromosome, while approximately half of meioses from fathers

transmit an X chromosome. Over many generations, approximately two-thirds of meioses from parents

to offspring that transmit an X chromosome will be from females, as we show below. However, the

proportion of females in a sample of individuals affects the actual proportion of meioses transmitting an

X chromosome in recent generations that are from females. For example, if the sample is solely made up

of females (with two X chromosomes, one from each parent), half of the meioses in the most recent

generation that transmit an X chromosome are from females. If the sample is made up solely of males

(with only one X chromosome, inherited from the mother), all the meioses that directly contribute the X

chromosomes in the sample are from females. If half the sampled individuals are female and half are

male, then 2/3 of the meioses in the previous generation that contribute the X chromosomes in the

sample will be from females. We show in the next paragraph that when half of the samples are female

this 2/3 ratio will apply in all prior generations, regardless of sex-specific demographic forces.

Consider the lineage of a randomly selected haplotype at a point in the genome. Let p_g be the

probability that the ancestral haplotype at generation g before present is carried by a female, where

$g = 0$ corresponds to the generation of the sampled individuals, $g = 1$ to the generation of their

parents, and so on. A given haplotype carried by a female has a 50% probability that its parent

haplotype is carried by a female, while a given haplotype carried by a male always has its parent

haplotype carried by a female. Thus, for $g \geq 0$,

72
$$p_{g+1} = 0.5p_g + (1 - p_g) = 1 - 0.5p_g.$$

73   If $p_g = 2/3$, then $p_{g+1} = 2/3$. If the sampled haplotype is randomly chosen from a set of individuals

74   with equal numbers of females and males, then the sampled haplotype has probability $p_0 = 2/3$ of

75   being carried by a female, and as a result $p_g = 2/3$ for all g. More generally, it can be shown by

76   mathematical induction that

78
$$p_g = \frac{2}{3}\left(1 - \left(-\frac{1}{2}\right)^g\right) + \left(-\frac{1}{2}\right)^g p_0$$

which converges to 2/3 for large g. In what follows, we assume that $p_g = 2/3$ for all g.

We model the probability distribution of the length of an X chromosome IBD segment resulting from a

coalescence event occurring g generations ago as an exponential random variable with mean length

$50/g$ cM on the sex-averaged recombination map. This is the same model that we use for estimating

IBD-based effective population size in autosomal data.[3] The model is somewhat inaccurate for small g

on the X chromosome because of the large difference between the female and male recombination

maps (with the male-specific X chromosome map having a recombination rate of 0). For larger g,

averaging over multiple generations ensures that the exponential distribution in this model is a good fit.

We investigate the performance of the method with this model through simulations.

IBD-based estimation of X chromosome effective population size

Our method for IBD-based estimation of X chromosome effective size history is based on IBDNe which

was designed to estimate recent effective population size from autosomal IBD segments.[3] The IBDNe

method calculates the expected length distribution of IBD segments exceeding a given length threshold

(such as 2 cM) for a given effective population size history. It finds the effective size history that equates

the observed and expected IBD length distributions using an iterative scheme. IBDNe applies smoothing

over intervals of eight generations to avoid overfitting. Although IBDNe was designed for autosomal

data, we show that it can also be used with X chromosome data, with some adjustments to the analysis

procedure that we describe in the following paragraphs.

The first adjustment ensures proper inference of IBD segments on the X chromosome by encoding male

X chromosome genotypes as haploid. This coding conforms to the VCF specification.[19] Male X

chromosome genotypes are frequently coded as homozygous diploid genotypes rather than haploid

genotypes, which typically results in duplicate reported IBD segments when using IBD detection

methods designed for autosomal data. The hap-ibd[20] program can correctly analyze chromosome X data

with haploid male genotypes. We exclude the pseudo-autosomal regions from analysis.

The second adjustment is to use a sex-averaged genetic map, as we also do for the autosomes. X

chromosomes transmitted from females are subject to recombination, while X chromosomes

transmitted from males are not (except for the pseudo-autosomal regions which we exclude from all

analyses). Genetic maps, such as the HapMap map[21] and the deCODE map[22], typically report the female-

specific recombination map for the X chromosome. Since an average of 2/3 of meioses transmitting an X

chromosome are from females, the sex-averaged X chromosome recombination map can be obtained by

multiplying female-specific genetic distances by 2/3. For example, a region with length 3 cM on the

female-specific map has length 2 cM on the sex-averaged map. Equivalently, the sex-averaged

recombination rates can be obtained by multiplying female-specific recombination rates by 2/3. For

example, an X chromosome region with female-specific recombination rate of 3×10^{-8} per base pair

per generation has a sex-averaged recombination rate of 2×10^{-8} per base pair per generation.

The third adjustment ensures equal numbers of sampled females and males. If the sample is

unbalanced, we selectively remove some individuals to obtain equal numbers of females and males.

Consequently, p_0, the proportion of sampled X chromosome haplotypes carried by females, is 2/3, and

hence p_g, the probability that the ancestral haplotype of a sampled X chromosome haplotype g

generations before the present is carried by a female is always 2/3 (see the preceding "Probability

Modelling for the X chromosome" section).

The fourth adjustment modifies the IBDNe "npairs" parameter to be equal to the number of analyzed

haplotype pairs. By default, IBDNe assumes that each individual contributes two haplotypes to the

analysis, and that all cross-individual pairs are analyzed, resulting in $(2n)(2n-2)/2$ haplotype pairs

when there are n individuals. On the X chromosome, with n_f females and n_m males, the number of

haplotype pairs is

$$2n_f(2n_f - 2)/2 \; + \; n_m(n_m - 1)/2 \; + \; 2n_f n_m.$$

We set the IBDNe "npairs" parameter to the value in equation 1 when analyzing X chromosome data,

after adjusting it for removal of close relative pairs as described below.

The fifth adjustment is to manually remove detected IBD segments corresponding to close relatives

(parent-offspring and siblings). By default, IBDNe identifies the close relatives using the input IBD

segments, removes the IBD segments between them, and adjusts the "npairs" parameter to account for

the removed sample pairs. However, this strategy does not work for the X chromosome because one

cannot reliably detect close relatives using only X chromosome data. We thus turn off IBDNe's filtering

of close relatives by setting "filtersamples=false". We can identify close relatives based on autosomal

data or from a pedigree file if available, and then remove IBD segments for these pairs and update the

"npairs" parameter accordingly in the chromosome X analysis.

The sixth adjustment enables calculation of confidence intervals. IBDNe obtains confidence intervals for

the estimated effective sizes by bootstrapping over chromosomes. We thus divide the X chromosome

into six pieces of equal cM length and treat these as separate "chromosomes" in the analysis with

IBDNe.

From X chromosome effective population size to sex-specific effective population size

We next describe how the estimated X chromosome effective population size can be used in conjunction

with the estimated autosome effective population size to estimate female and male effective population

sizes. We will write N_g^X and N_g^A for the X chromosome and autosomal effective sizes at generation g.

And we will write N_g^f and N_g^m for the female and male effective population sizes at generation g, which

can be derived from the X chromosome and autosomal effective sizes as described below.

The IBD-based effective population size (for autosomes or X chromosome) is defined in terms of the

conditional coalescence probability for a Wright-Fisher population. We first consider autosomes. For a

randomly selected pair of haplotypes, let G be the number of generations before present that the

haplotypes coalesce. Conditional on the haplotypes not coalescing by generation $g - 1$ before present,

the ancestral haplotypes are distinct at that generation. For them to coalesce at generation g before

present, their two parental haplotypes at generation g must be the same haplotype. If the diploid

autosomal effective size is N_g^A at g generations before present, there are $2N_g^A$ autosomal haplotypes

available, and the probability that the two parental autosomal haplotypes are the same is thus

$1/(2N_g^A)$. That is, $P_A(G = g|G > g - 1) = 1/(2N_g^A)$. Thus, if we know the value of

$P_A(G = g|G > g - 1)$, then we can obtain the effective size g generations before present:

156
$$N_g^A = 1/\big(2P_A(G = g|G > g - 1)\big).$$

157   On the X chromosome, the conditional coalescence probability can be obtained by considering that 2/3

158   of meioses are from female parents, while 1/3 are from male parents. For coalescence to occur, both

159   haplotypes' parent haplotypes must be the same. This means both haplotypes must be inherited from

160   parents that have the same sex, and the second haplotype must have the same parental haplotype as

161   that of the first (within that sex). Thus,

162
$$P_X(G = g|G > g - 1) = \frac{(2/3)^2}{2N_g^f} + \frac{(1/3)^2}{N_g^m}.$$

163   And hence,

164
$$N_g^X = 1/\big(2P_X(G = g|G > g - 1)\big)$$

165
$$= \frac{9N_g^f N_g^m}{2N_g^f + 4N_g^m}$$

166   Equation 2

167   For comparison, on the autosomes, by the same reasoning but with half of the meioses from each sex

168   and with diploid males,

$$P_A(G = g | G > g - 1) = \frac{(1/2)^2}{2N_g^f} + \frac{(1/2)^2}{2N_g^m}.$$

170   And hence,

$$N_g^A = \frac{1}{2} \Big/ \left(\frac{(1/2)^2}{2N_g^f} + \frac{(1/2)^2}{2N_g^m} \right) = \frac{4N_g^f N_g^m}{N_g^f + N_g^m}.$$

172   Equation 3

173   From Equations 2 and 3, the ratio of X to autosomal effective size, which we denote as α, is

$$\alpha = \frac{N_g^X}{N_g^A}$$

$$= \frac{9(N_g^f + N_g^m)}{8(N_g^f + 2N_g^m)}$$

$$= \frac{9\left(1 + (N_g^m/N_g^f)\right)}{8\left(1 + 2(N_g^m/N_g^f)\right)}.$$

177   Equation 4

178   Thus $N_g^m/N_g^f \to 0$ as $\alpha \to 9/8$, and $N_g^m/N_g^f \to \infty$ as $\alpha \to 9/16$. Since α is a decreasing function of

179   N_g^m/N_g^f, the ratio of X to autosomal effective size satisfies $9/16 < \alpha < 9/8$.

180   With algebra, it can be shown that Equations 2 and 3 imply that

$$N_g^f = \frac{2N_g^X N_g^A}{9N_g^A - 8N_g^X}$$

182   Equation 5

183   and

184
$$N_g^m = \frac{2N_g^X N_g^A}{16N_g^X - 9N_g^A}.$$

185 Equation 6

186 Consequently, the effective sex ratio is

187
$$\text{ESR} = \frac{N_g^f}{N_g^f + N_g^m} = \frac{16N_g^X - 9N_g^A}{8N_g^X}$$

Equation 7

Thus, given estimates of N_g^X and N_g^A, one can use Equations 5, 6, and 7 to obtain estimates for N_g^f, N_g^m,

and the ESR. These are standard equations for estimating sex-specific effective population sizes based

on N_g^X and N_g^A,[23,24] although usually presented in the context of constant effective population sizes

across time.

According to Equation 4, the allowable range of X chromosome N_e is between $\frac{9}{16}$ and $\frac{9}{8}$ of the autosome

N_e at each generation. When the X chromosome N_e is overestimated or the autosome N_e is

underestimated, the estimated female effective size (Equation 5) can be negative. On the other hand,

underestimation of the X chromosome N_e or overestimation of the autosome N_e can result in a negative

estimate of the male effective size (Equation 6).

Analysis pipeline

We start with phased sequence data (true phase for the simulated data, and inferred phase for real

data), with males coded as haploid on the X chromosome. We use hap-ibd[20] to infer IBD segments. For

hap-ibd analysis on sequence data, we set the minimum seed length to 0.5 cM and the minimum

extension length to 0.2 cM. The relatively small minimum seed length and minimum extension length

increase power to detect short IBD segments. We exclude rare variants by setting the minimum minor

allele count filter to 100 because these lower frequency variants are less informative, have less accurate

phasing, and may be recent mutations. For SNP-array data, due to the lower marker density, we set the

minimum seed length to 1, the minimum extension length to 0.1, and the minimum number of markers

in a seed IBS segment to 50. Other parameters are left at their default values. The IBD analysis

parameters are the same for the autosomes and X chromosome.

We then run IBDNe on the detected IBD segments output, with one analysis for autosomes and a

separate analysis for the X chromosome. The genetic map file for the analysis is assumed to be a sex-

averaged map. For the X chromosome, this means multiplying cM positions in the female-specific map

by 2/3.

When applying IBDNe on the simulated data (autosomes or X), we set "filtersamples=false" and

"gmin=1" because the chromosomes are simulated independently so that a pair of individuals can share

ancestry one generation back (i.e. be siblings) on one chromosome without such sharing occurring on

other chromosomes. These settings tell IBDNe not to look for and remove close relatives, and to model

IBD from shared ancestry starting from one generation before present.

Real data often has an excess of close relatives due to the sampling scheme. Thus, in the analysis of real

autosomal data we allow IBDNe to detect and remove close relatives, which is the default behavior. The

X chromosome on its own is not sufficient to detect close relatives, so we use either available pedigree

information or the close relative pairs identified by IBDNe in the autosomal analysis to manually remove

IBD segments from close relatives in the X chromosome data. We then update the "npairs" parameter

accordingly (see below), and set "filtersamples=false" in the X chromosome IBDNe analysis.

In the IBDNe analysis of the X chromosome data, we set the "npairs" parameter equal to the number of

haplotype pairs for which IBD segments could be present (i.e., all pairs except for those for which we

have explicitly removed IBD segments). We first calculate the number of haplotype pairs based on the

numbers of females and males in the sample, using Equation 1. We then adjust the number of haplotype

pairs to account for the number of close relative pairs that were removed. Removing a male-male pair

decreases the count by 1. Removing a male-female pair decreases the count by 2. Removing a female-

female pair decreases the count by 4.

In order for IBDNe to obtain bootstrap confidence intervals for the X chromosome effective size

estimates, we divide the X chromosome into six pieces of equal cM length. We recode the chromosome

field of the IBD segment file using integer values between 1 and 6 according to the location of the IBD

segments. IBD segments that cross more than one of these "chromosomes" are split into subsegments

at the boundaries of the corresponding "chromosomes". We remove the centromere region from IBD

segments, so that each IBD segment that crosses the centromere region is replaced with two IBD

segments: a segment before and a segment after the centromere.

After obtaining the X chromosome and autosomal effective population sizes, we estimate the female

and male effective sizes and effective sex ratio using Equations 5, 6, and 7. We obtain bootstrap values

for these estimates by taking pairs of bootstrap values from the X and autosomal analyses. For example,

for the n-th bootstrap value of the female effective population size at generation g, we take the n-th

bootstrap value for the X chromosome effective size at generation g and the n-th bootstrap value for

the autosomal effective size at generation g, and apply Equation 5. After obtaining all the bootstrap

values, we use the 2.5[th] and 97.5[th] percentile to obtain an approximate 95% confidence interval for the

female (for example) effective population size at generation g.

Simulation study

We conducted a simulation study to evaluate the performance of our method. We used SLiM, a forward

simulator,[25] to simulate the demographic history for the most recent 5000 generations, and we used

msprime, a coalescent simulator, to complete the simulation back to full coalescence of the sample.[26,27]

For all scenarios, we simulated data for 30 autosomes of length 100 Mb and an X chromosome of length

180 Mb.

For the simulation in SLiM, we used a mutation rate of 10^{-8} per base pair per generation, and a

recombination rate of 10^{-8} per base pair per meiosis on the autosomes and for female meioses on the

X chromosome. We set the gene conversion initiation rate to 2×10^{-8} per base pair per meiosis on the

autosomes and per female meiosis on the X chromosome, with mean gene conversion tract length of

300 bp. We simulated populations with equal sex ratio, and we also simulated populations with 20%

females, 40% females, 60% females, and 80% females. We used the same total effective population size

$(N_g^f + N_g^m)$ in each simulation. We sampled 50,000 individuals comprising 25,000 females and 25,000

males for each analysis.

We simulated data under two different demographic models. The first model is a four-stage

exponentially growing population with increased growth rate over time, which we call the "UK-like"

scenario because it approximates the demographic history of the UK population.[28] The forward

simulation of this model in SLiM starts from 5000 generations ago with an initial size of 3000. At 300

generations ago, this population starts to grow at an exponential rate of 1.4% per generation. At 60

generations ago, the rate of the exponential growth increases to 6%. In the most recent 10 generations,

the growth rate further increases to 25%, and the population size reaches around 21 million at the time

of sampling.

The second demographic model was modified from the "UK-like" model by adding a bottleneck 10

generations before present, which approximates the immigration of Europeans to America. This

bottleneck reduces the population size to 100,000. The population then continues to grow at an

exponential rate of 25% and reaches a final size around 1.2 million. We call this simulated population

the "US-like" scenario.

When completing the simulations with msprime, we did not apply gene conversion or sex-specific

population size. In the msprime simulations, mutation occurred at a rate of 10^{-8} per base pair per

generation, and recombination at a rate of 10^{-8} per base pair per meiosis. There is no differentiation

between female and male meioses in the generations prior to the starting generation of the forward

simulation because msprime does not have sex-specific functionality. This lack of sex-specific treatment

in the period more than 5000 generations ago will affect the level of variation in the data but will not

affect the distribution of IBD segments of length > 2 cM (the segments analyzed by IBDNe) because such

segments typically have ancestry within the past three hundred generations.

Analysis of human populations

We applied the above analysis pipeline on SNP-array data from the UK Biobank[29] and whole genome

sequence data from the TOPMed project[30] to infer the sex-specific population history of several human

populations.

The UK Biobank is a large-scale biomedical database that contains in-depth genetic, physical and health

data collected between 2006 and 2010 on half a million UK participants aged between 40 and 69.[31]

Genome-wide genetic SNP-array data were collected on every participant using the UK Biobank Axiom

array that assays approximately 850,000 genetic variants across genome.[29] Previous IBDNe analyses of

SNP-array data have found that effective population size estimates are less precise when estimating the

effective population size more than 50 or so generations before the present due to uncertainty in the

IBD-segment endpoints.[3] A previous study on UK Biobank data also showed lower false positive rates in

estimated IBD segments that were at least 4 cM compared to estimated IBD segments with length

below 4 cM.[20] Therefore, when analyzing UK Biobank data, we used IBD segments that were at least 4

cM to estimate population history in the past 40 generations. We used the GRCh37 European

recombination map developed by Bherer et al. (2017) in the analyses.[32]

We first analyzed the population history of the White British participants, which is the largest ethnic

group in the UK Biobank cohort. The UK Biobank genotype data include 221,141 White British females

and 187,802 White British males. Since IBDNe has limits on the number of IBD segments that it can

process, we randomly selected 5,000 White British females and 5,000 White British males for estimation

of the autosome N_e. For analysis of the X chromosome N_e, we removed 33,339 randomly selected

females to ensure equal numbers of females and males in the sample. For the autosomes, IBDNe

removed close relatives using default settings. For the X chromosome, we used the UK Biobank's kinship

estimates to identify and remove IBD segments from sibling pairs and parent-offspring pairs.[29]

The UK Biobank cohort also includes participants from several ethnic minority groups including Black

British, Indian, Pakistani, Asian, and Bangladeshi. Among these, we chose to analyze the effective

population size of the Indian group, which is the largest minority group in the UK Biobank, although this

group itself contains considerable diversity. There are genotype data for 2885 males and 2775 females

with Indian ancestry in UK Biobank. We removed 110 randomly selected males to achieve equal

numbers of females and males for the subsequent analysis. By default, IBDNe automatically removes

IBD segments from pairs of related individuals and generates a list of these related pairs. We manually

removed X chromosome IBD segments for these pairs of related individuals prior to estimation of X

chromosome N_e using IBDNe.

The Hypertension Genetic Epidemiology Network Study (HyperGEN) study includes individuals with

hypertension from Birmingham, AL and Winston-Salem, NC. Whole genome sequencing was performed

as part of the TOPMed project (dbGaP: phs001293.v2.p1). Haplotype phasing was performed as part of

the phasing of a larger set of Freeze 8 TOPMed data.[33] We estimated the effective population size from

the 1,586 Black non-Hispanic participants from this study. Prior to analysis, we randomly removed 416

Black non-Hispanic female participants to ensure equal numbers of females and males in the sample.

IBD segments from closely related pairs of individuals were excluded in the IBDNe analysis on the

autosome data. X chromosome IBD segments from close relatives were manually removed using the list

of relative pairs identified by IBDNe in the autosomal analysis. We used 2 cM as the IBD length threshold

and estimated effective population size over the past 100 generations. We used the GRCh38 deCODE

map developed by Halldorsson et al. (2019) in the analyses.[22]

Results

Simulation study

We compared N_e estimated from the simulated autosome and X chromosome data to the actual N_e for

the UK-like and US-like demographic models. The actual autosome or X chromosome N_e can be

obtained from the sex-specific effective population sizes using Equations 2 and 3.

The estimated N_e generally matches the true N_e closely (Figure 1 and Figures S1-S2). However, some

discrepancies exist between the estimated and actual N_e because IBDNe cannot localize sharp changes

in the population size to one exact generation and it tends to over-smooth corners of the population-

size trajectory.[3] For the US-like simulations with a sharp bottleneck event that happened at a single

generation, the estimated duration of bottleneck is spread over multiple generations (Figure 1).

We next used the estimated X chromosome N_e and autosome N_e to estimate the sex-specific effective

population sizes and the effective sex ratio. We find that the formulas for the sex-specific N_e and ESR as

functions of the autosome and X chromosome N_e (Equation 5, 6, 7) are sensitive to errors in N_e

estimation. In UK-like scenarios, with more accurate estimates, the estimated sex-specific N_e and the

estimated ESR are similar to the actual values, except at around 20 generations ago, where there was a

large change in the population growth rate (Figures 1 and 2 and Figure S1). For the US-like scenarios,

around the time of the bottleneck event where there is greater inaccuracy in the estimated N_e, the

estimated sex-specific N_e differs significantly from the actual N_e (Figure 1). As a result, the ESR

estimated directly from the sex-specific N_e in the US-like simulation is also quite inaccurate (Figure 2 and

Figure S2), and the estimated confidence band generally fails to cover the true sex ratio.

UK Biobank data

The estimated autosome N_e for the UK Biobank's White British population (Figure 3) had a high rate of

growth in the most recent 20 generations and reached a current population size of 40 million (95%

confidence interval = 35-47 million). Before that, this population had no noticeable change in its

effective size. The autosome N_e estimated from the genotype data of the UK Biobank Indian participants

(Figure 4) shows slow growth from 40 generations ago until around 10 generations ago. The increased

growth beginning 10 generations ago results in an estimated current effective population size of 4.2

million (95% confidence interval = 3.7-5.0 million).

The inferred X chromosome N_e has the same shape as the inferred autosome N_e for both groups.

Notably, the estimated X chromosome N_e is close to 75% of the inferred autosomal effective size, which

is the expected effective size that would be obtained from the X chromosome data if the female and

male effective population sizes have been equal in these populations (Figures 3 and 4). For the UK Indian

population, the confidence band for the X chromosome N_e overlaps that of 75% of the autosome N_e

over the entire 40 generation period. For the UK White British population, the confidence band of 75%

of the autosome N_e is also covered by that of the X chromosome N_e in the most recent 10 generations.

This suggests that these two populations had a balanced sex ratio in their recent history.

To further investigate the sex composition of the UK white British population and the UK Indian

population in the recent past, we applied the X chromosome and autosome N_e to estimate the female

and male N_e and the effective sex ratio of these two populations. For both populations, the sex-specific

N_e has a similar overall trend to that of the entire population (Figures 3 and 4). The estimated effective

sex ratio has a high level of uncertainty in both populations (Figures 3 and 4).

 Trans-Omics for Precision Medicine (TOPMed) study data

The inferred effective population size history of the TOPMed HyperGEN African American cohort shows

a period of fast growth from 100 generations before present until around 40 generations ago. The

effective population size then stayed relatively stable for around 10 generations. A bottleneck event

from 32 to 11 generations ago reduced the N_e of this population from 1.4 million (95% CI: 1.1-1.7

million) to 112 thousand (95% CI: 105-119 thousand). A period of high growth since then has restored

the N_e to 3.6 million at the present time (95% CI: 2.7-5.4 million) (Figure 5). The recent bottleneck event

in the estimated effective-size trajectory of the HyperGEN Black non-Hispanic participants is consistent

with the immigration bottleneck resulting from the transatlantic slave trade in the history of the African

American population.

As in the analysis of UK Biobank data, the X chromosome effective size inferred from the HyperGEN

group follows the same trend as the autosome N_e of this group (Figure 5). However, we observed more

discrepancy between the estimated X chromosome N_e and the estimated autosomal N_e than was seen

in the UK Biobank data, matching the comparison of results found in the US-like and UK-like simulations.

Compared to the estimated autosome N_e, the X chromosome N_e estimate was high between 65 to 30

generations ago, which produced negative estimates of female-specific N_e during this period (Figure 5).

The estimated X chromosome N_e was also low compared to the estimated autosome N_e both during the

most recent 6 generations and 21 to 18 generations ago, which led to negative male-specific N_e (Figure

5). There is significant uncertainty in the ESR estimates.

Discussion

Previous studies have shown that the X chromosome can provide information about demographic

processes that cannot be revealed by the analysis of autosomes alone.[9,11,12,14,34,35] In this work, we

focused on utilizing IBD segments on the X chromosome to infer the X chromosome effective population

size. We developed a framework to model X chromosome N_e and derived the relationship between X

chromosome and autosome N_e by considering the different coalescence rates among X chromosomes

and autosomes. We also showed how to apply this information to calculate the female and male

effective population sizes and the effective sex ratio in a population as functions of the X chromosome

and autosome N_e.

We validated the performance of our method in simulated populations with similar histories to the UK

and the US populations. We found that pronounced differences between female and male effective

population sizes can be detected. Such differences are easier to detect in populations that have not

undergone recent bottlenecks, because recent bottlenecks create noise in the estimates of female and

male effective population sizes.

We applied our method to estimate the X chromosome effective size for the UK British and UK Indian

populations in the UK Biobank, as well as an African American population from the TOPMed HyperGEN

study. When using the X chromosome effective population size to estimate the sex-specific effective

population size, we obtained results consistent with gender-balanced demographic histories for each of

these populations.

Although X chromosome IBD information has been used by previous studies for the estimation of

genealogical relations between individuals, especially for kinship estimation in forensic settings[5,6,36],

there has been a lack of studies that use X chromosome IBD segments to estimate recent effective

population size. Our work thus fills a gap that existed in the application of X chromosome IBD

information in population genetic studies.

Previous methods for estimating sex-specific population history or sex bias in human populations have

relied on comparisons of genetic diversity between autosomes and the X chromosome using allele

frequency differentiation, patterns of neutral polymorphism, and the site frequency

spectrum.[13,15,16,18,35,37] Most of these methods considered only a single estimate of the effective sex ratio

over the entire history of a population, although this ratio can vary over time.[13,15,18,37] Some of these

methods also focused on a constant overall effective size across time, although changes in effective size

can distort these analyses.[35,37] Recently, Musharoff et al. developed a likelihood ratio test for population

sex bias that considered populations of non-constant size and changing sex ratios using site frequency

spectrum data.[16] However, this method requires demographic parameters to be constant within time

epochs. In comparison, our approach for estimating the X chromosome effective population size and the

female and male effective population sizes requires minimal assumptions and allows the effective

population sizes to vary independently over time. The ability of our IBD-based analyses to infer effective

population sizes in the past hundred generations distinguishes our approach from other methods.

There are several limitations to our approach. First, IBD-based estimation of effective population size

requires a large sample of individuals from the population, although this may be less of an issue given

the continuing increase in the size of genetic studies, particularly in humans. Second, our method for

estimating N_e is less accurate around generations where the population experienced a drastic change in

population size trajectory such as a bottleneck event. Joint estimation of autosomal and X chromosome

effective size could improve estimation of the effective sex ratio in such situations. In addition, the

performance of IBDNe is affected by the number and accuracy of detected IBD segments. For example,

we observe wider confidence bands for the estimated N_e in the most recent generations since there

tend to be fewer very long IBD segments in the sample. Similarly, we observe wider confidence bands

for the X chromosome N_e compared to the autosomal N_e due to the smaller amount of data in the X

chromosome. Moreover, the estimated autosome and X chromosome effective size tend to deviate

more from the simulated effective size in the more distant past due to the lower relative accuracy when

estimating the lengths of shorter IBD segments.

Given these limitations, we note that our method is not suitable for rigorously testing hypotheses on

sex-specific N_e and the effective sex ratio in a population, because small inaccuracies in estimation of

autosome and X chromosome effective population sizes are magnified when transforming these to

estimates of sex-specific effective population size and the effective sex ratio. However, our method can

still serve as an exploratory tool for detecting evidence of sex-specific past demographic events.

Data and code availability

441   UK Biobank data were downloaded from the European Genome-Phenome Archive (https://ega-

442   archive.org/datasets/EGAD00010001497). TOPMed freeze 8 data for study accession phs001293.v2.p1

443   (HyperGEN) were downloaded from dbGaP (https://www.ncbi.nlm.nih.gov/gap/). The IBDNe software is

444   available from https://faculty.washington.edu/browning/ibdne.html. The hap-ibd software is available

445   from https://github.com/browning-lab/hap-ibd.

Acknowledgements

Research reported in this publication was supported by the National Human Genome Research Institute

of the National Institutes of Health under award number HG005701. The content is solely the

responsibility of the authors and does not necessarily represent the official views of the National

Institutes of Health.

This research has been conducted using the UK Biobank Resource under Application Number 19934.

Molecular data for the Trans-Omics in Precision Medicine (TOPMed) program was supported by the

National Heart, Lung and Blood Institute (NHLBI). Core support including centralized genomic read

mapping and genotype calling, along with variant quality metrics and filtering were provided by the

TOPMed Informatics Research Center (3R01HL-117626-02S1; contract HHSN268201800002I). Core

support including phenotype harmonization, data management, sample-identity QC, and general

program coordination were provided by the TOPMed Data Coordinating Center (R01HL-120393; U01HL-

120393; contract HHSN268201800001I). We gratefully acknowledge the studies and participants who

provided biological samples and data for TOPMed. The Hypertension Genetic Epidemiology Network

Study is part of the NHLBI Family Blood Pressure Program; collection of the data represented here was

supported by grants U01 HL054472, U01 HL054473, U01 HL054495, and U01 HL054509; genome

sequencing was funded by R01HL055673.

Literature cited

1. Wright S. Evolution in Mendelian populations. *Genetics.* 1931;16(2):97.

2. Charlesworth B. Effective population size and patterns of molecular evolution and variation. *Nature Reviews Genetics.* 2009;10(3):195-205.

3. Browning SR, Browning BL. Accurate non-parametric estimation of recent effective population size from segments of identity by descent. *Am J Hum Genet.* 2015;97(3):404-418.

4. Browning SR, Browning BL, Daviglus ML, et al. Ancestry-specific recent effective population size in the Americas. *PLoS Genet.* 2018;14:e1007385.

5. Henden L, Wakeham D, Bahlo M. XIBD: software for inferring pairwise identity by descent on the X chromosome. *Bioinformatics.* 2016;32(15):2389-2391.

6. Buffalo V, Mount SM, Coop G. A genealogical look at shared ancestry on the X chromosome. *Genetics.* 2016;204(1):57-75.

7. Palamara PF, Lencz T, Darvasi A, Pe'er I. Length distributions of identity by descent reveal fine-scale demographic history. *American Journal of Human Genetics.* 2012;91(5):809-822.

8. Ségurel L, Martínez-Cruz B, Quintana-Murci L, et al. Sex-specific genetic structure and social organization in Central Asia: insights from a multi-locus study. *PLoS Genet.* 2008;4(9):e1000200.

9. Bryc K, Auton A, Nelson MR, et al. Genome-wide patterns of population structure and admixture in West Africans and African Americans. *Proceedings of the National Academy of Sciences.* 2010;107(2):786-791.

10. Heyer E, Chaix R, Pavard S, Austerlitz F. Sex-specific demographic behaviours that shape human genomic variation. *Mol Ecol.* 2012;21(3):597-612.

11. Goldberg A, Rosenberg NA. Beyond 2/3 and 1/3: the complex signatures of sex-biased admixture on the X chromosome. *Genetics.* 2015;201(1):263-279.

498   12.   Shringarpure SS, Bustamante CD, Lange K, Alexander DH. Efficient analysis of large datasets and

499   sex bias with ADMIXTURE. *BMC bioinformatics.* 2016;17(1):1-6.

500   13.   Hammer MF, Mendez FL, Cox MP, Woerner AE, Wall JD. Sex-biased evolutionary forces shape

501   genomic patterns of human diversity. *PLoS genetics.* 2008;4(9):e1000202.

502   14.   Bustamante CD, Ramachandran S. Evaluating signatures of sex-specific processes in the human

503   genome. *Nature genetics.* 2009;41(1):8-10.

504   15.   Clemente F, Gautier M, Vitalis R. Inferring sex-specific demographic history from SNP data. *PLoS*

505   *genetics.* 2018;14(1):e1007191.

506   16.   Musharoff S, Shringarpure S, Bustamante CD, Ramachandran S. The inference of sex-biased

507   human demography from whole-genome data. *PLoS genetics.* 2019;15(9):e1008293.

508   17.   Wang J, Santiago E, Caballero A. Prediction and estimation of effective population size. *Heredity.*

509   2016;117(4):193-206.

510   18.   Emery LS, Felsenstein J, Akey JM. Estimators of the human effective sex ratio detect sex biases

511   on different timescales. *The American Journal of Human Genetics.* 2010;87(6):848-856.

512   19.   File Formats Task Team. The Variant Call Format Specication: VCFv4.3 and BCFv2.2. 2020.

513   http://samtools.github.io/hts-specs/VCFv4.3.pdf.

514   20.   Zhou Y, Browning SR, Browning BL. A fast and simple method for detecting identity-by-descent

515   segments in large-scale data. *The American Journal of Human Genetics.* 2020;106(4):426-437.

516   21.   International HapMap Consortium. A second generation human haplotype map of over 3.1

517   million SNPs. *Nature.* 2007;449(7164):851.

518   22.   Halldorsson BV, Palsson G, Stefansson OA, et al. Characterizing mutagenic effects of

519   recombination through a sequence-level genetic map. *Science.* 2019;363(6425):eaau1043.

520   23.   Wright S. *Evolution and the genetics of populations: Vol. 2. The theory of gene frequencies.* 1969.

521   24.   Hartl DL, Clark AG. *Principles of population genetics.* Vol 116: Sinauer associates Sunderland;

522   1997.

523   25.   Haller BC, Messer PW. SLiM 3: forward genetic simulations beyond the Wright-Fisher model.

524   *Molecular biology and evolution.* 2019;36(3):632-637.

525   26.   Kelleher J, Etheridge AM, McVean G. Efficient coalescent simulation and genealogical analysis

526   for large sample sizes. *PLoS computational biology.* 2016;12(5):e1004842.

527   27.   Haller BC, Galloway J, Kelleher J, Messer PW, Ralph PL. Tree-sequence recording in SLiM opens

528   new horizons for forward-time simulation of whole genomes. *Molecular ecology resources.*

529   2019;19(2):552-566.

530   28.   Browning BL, Browning SR. Detecting identity by descent and estimating genotype error rates in

531   sequence data. *The American Journal of Human Genetics.* 2013;93(5):840-851.

532   29.   Bycroft C, Freeman C, Petkova D, et al. The UK Biobank resource with deep phenotyping and

533   genomic data. *Nature.* 2018;562(7726):203-209.

534   30.   Taliun D, Harris DN, Kessler MD, et al. Sequencing of 53,831 diverse genomes from the NHLBI

535   TOPMed Program. *Nature.* 2021;590(7845):290-299.

536   31.   Fry A, Littlejohns TJ, Sudlow C, et al. Comparison of sociodemographic and health-related

537   characteristics of UK Biobank participants with those of the general population. *American*

538   *journal of epidemiology.* 2017;186(9):1026-1034.

539   32.   Bhérer C, Campbell CL, Auton A. Refined genetic maps reveal sexual dimorphism in human

540   meiotic recombination at multiple scales. *Nat Commun.* 2017;8(1):1-9.

541   33.   Browning BL, Tian X, Zhou Y, Browning SR. Fast two-stage phasing of large-scale sequence data.

542   *The American Journal of Human Genetics.* 2021;108(10):1880-1890.

543   34.   Schaffner SF. The X chromosome in population genetics. *Nature Reviews Genetics.* 2004;5(1):43-

544   51.

35.   Ramachandran S, Rosenberg NA, Zhivotovsky LA, Feldman MW. Robustness of the inference of

human population structure: a comparison of X-chromosomal and autosomal microsatellites.

Human genomics. 2004;1(2):1-11.

36.   Pinto N, Gusmão L, Amorim A. X-chromosome markers in kinship testing: a generalisation of the

IBD approach identifying situations where their contribution is crucial. *Forensic Science*

International: Genetics. 2011;5(1):27-32.

37.   Keinan A, Mullikin JC, Patterson N, Reich D. Accelerated genetic drift on chromosome X during

the human dispersal out of Africa. *Nature genetics.* 2009;41(1):66-70.

 Figures

Figure 1. Estimates of autosome, X chromosome, and sex-specific N_e in simulation studies. Autosomal

N_e is shown in the left column, X chromosome N_e in the middle column, and sex-specific N_e in the right

column. The top two rows are the UK-like simulation, with equal sex ratio in the top row and 20%

females in the second row. The lower two rows are the US-like simulation, with equal sex ratio in the

third row and 20% females in the fourth row. Results for other choices of sex-ratio can be found in

Figures S1 and S2. Y-axes show N_e plotted on a log scale. In cases where the estimated sex-specific N_e is

negative (see Methods), it is not shown.

Figure 2. Estimates of population effective sex ratio (ESR) obtained from estimated sex-specific N_e.

The top row shows the UK-like simulation, while the bottom row shows the US-like simulation. The true

ratio is shown by the horizontal dashed line in each plot.

Figure 3. Effective population size of the UK Biobank White British group. From left to right, the top

row shows the estimated X chromosome N_e, the estimated autosome N_e, and a comparison of the

estimated X chromosome N_e with 75% of the estimated autosome N_e. The bottom row displays the

male-specific N_e, the female-specific N_e, and the ESR. For the N_e plots, the Y-axes show N_e on a log

scale. In cases where the estimated sex-specific N_e or its confidence band is negative (see Methods), the

negative values are not shown.

Figure 4. Effective population size of the UK Biobank Indian group. From left to right, the top row

shows the estimated X chromosome N_e, the estimated autosome N_e, and a comparison of the estimated

X chromosome N_e with 75% of the estimated autosome N_e. The bottom row displays the male-specific

N_e, the female-specific N_e, and the ESR. For the N_e plots, the Y-axes show N_e on a log scale. In cases

where the estimated sex-specific N_e or its confidence band is negative (see Methods), the negative

values are not shown.

Figure 5. Effective population size of the Black non-Hispanic group in the HyperGEN cohort. From left

to right, the top row shows the estimated X chromosome N_e, the estimated autosome N_e, and a

comparison of the estimated X chromosome N_e with 75% of the estimated autosome N_e. The bottom

row displays the male-specific N_e, the female-specific N_e, and the ESR. For the N_e plots, the Y-axes show

N_e on a log scale. In cases where the estimated sex-specific N_e or its confidence band is negative (see

Methods), the estimated values are not shown. Estimated ESR values outside the range $[0,1]$ are not

shown.

Supplementary materials

590 **Figure S1. Estimates of sex-specific population size and effective sex ratio in UK-like simulations.** From

591 left to right, the columns display results from a UK-like simulation with 80% , 60%, and 40% females.

592 Autosomal N_e is shown in the top row, X chromosome N_e in the second row, sex-specific N_e in the third

593 row, and the ESR in the bottom row. For the N_e plots the Y-axes are on a log scale.

594

595 **Figure S2. Estimates of sex-specific population size and effective sex ratio in US-like simulations.** From

596 left to right, the columns display results from a UK-like simulation with 80%, 60%, and 40% females.

597 Autosomal N_e is shown in the top row, X chromosome N_e in the second row, sex-specific N_e in the third

598 row, and the ESR in the bottom row. For the N_e plots the Y-axes are on a log scale. In cases where the

599 estimated sex-specific N_e is negative (see Methods), it is not shown.