The contraction-expansion behaviour in the demosponge *Tethya wilhelma* is diurnal and

light-controlled

Sarah B. Flensburg[1], Anders Garm[2], & Peter Funch[1]

[1] Aarhus University, Denmark

[2] University of Copenhagen, Denmark

Corresponding author:

Peter Funch

Contact details:

funch@bio.au.dk

Abstract

Sponges (phylum Porifera) are metazoans without muscles and nervous system. Still, they perform coordinated behaviours, such as whole body contrations. Previous studies have indicated diurnal variability in number of contractions, and in expression of circadian clock genes. Here we show that diurnal patterns are present in the contraction-expansion behaviour of the demosponge *Tethya wilhelma* using infrared videography and a simulated night/day-cycle including sunset and sunrise mimic. In addition, we show that this behaviour is at least strongly influenced by the ambient light intensity and therefore implicates light-sensing capabilities in this sponge species. This is backed by our finding that *T. wilhelma* consistently contracts at sunrise, and that this pattern disappears both when the sponge is kept in constant darkness and when in constant light.

Keywords:
Porifera, pre-nervous system, light reception, behaviour, time-lapse imaging

Introduction

Sponges (phylum Porifera) are aquatic animals with a world-wide distribution found at all water depths. The group comprises over 20,000 described species across four extant classes of which the class Demospongiae contains most of the species and is the most abundant (De Voogd et al., 2021; Van Soest et al., 2012). The sponge body plan is fundamentally organized around cavities and canals where water flows through the sponge (Asadzadeh et al., 2020). Despite their relatively simple body organization, sponges possess a number of different cell types but lack muscle and nerve cells (Lavrov and Kosevich, 2014; Musser et al., 2021). Still, they are able to perform coordinated behaviours such as whole-body contractions, and some sponges even have the ability to move (Bond, 2013; Nickel, 2006). Contractions can happen spontaneously. They can also be initiated by a range of stimuli including different chemicals (Elliott and Leys, 2010; Ellwanger et al., 2007; Ellwanger and Nickel, 2006; Goldstein et al., 2020), or by a mechanical stimulus locally on the exopinacoderm and spread as a contraction wave across the surface of the sponge (Elliott and Leys, 2007; Nickel et al., 2011). Also, shaking of the freshwater sponge *Ephydatia muelleri* causes a contraction, and exposure to high loads of ink particles causes it to contract and expel waste particles from its water canal system (Elliott and Leys, 2007). The mechanism initiating this behaviour has not been identified but primary cilia located in the oscular area have been suggested to function as mechanoreceptors involved in water flow regulation and sponge body contractions (Ludeman et al., 2014). Despite the lack of neurons, stimulation of one part of the sponge often result in changes in behaviour throughout the entire sponge, suggesting coordination, possibly facilitated by chemical messengers (Kealy et al., 2019). Light also influences sponge behaviours, including the release of larvae, which in some species is correlated with the lunar cycle (Amano, 1986; Amano, 1988; Hoppe and Reichert, 1987; Neely and Butler, 2020). Some sponge larvae likewise respond to changes in light conditions, suggesting that they possess a light-sensory system (Leys et al., 2002). However, genetic data suggest that sponges lack the photopigment opsin, most often involved in animal vision, and evidence from the demosponge *Amphimedon queenslandica* indicates that the larvae here use cryptochromes instead (Mah and Leys, 2017; Rivera et al., 2012). In addition to performing behaviours normally associated with a nervous system, sponges also possess many of the genes associated with the nervous system in other animals. This has lead to the hypothesis that sponges, while not having a nervous system *per se*, might have a kind of pre-nervous system able to perform some of the functions of a fully developed nervous system (Leys, 2015).

Diurnal activity patterns are commonly observed in many different types of

organisms. In animals, a typical pattern consists of alternating periods of inactivity, sleep,

which can be observed across a wide range of species (Siegel, 2008), including animals

without a conventional brain such as the cnidarian polyp *Hydra vulgaris* (Kanaya et al.,

2020) and the jellyfish *Cassiopea* sp. (Nath et al., 2017). Studies of diurnal rythms in animals

without a nervous system are very limited. In a long term *in situ* study Reiswig (1971)

showed a synchronized diurnal cycle of body contractions and expansions of the demosponge

Tectitethya crypta. Another demosponge *Tethya wilhelma* also seems to exhibit diurnal

variations in body contractions, as indicated by the results from two specimens tested under

laboratory conditions with an artifical 12h:12h light:dark cycle (Nickel, 2004). The control

mechanisms behind the diurnal rhythm were not tested, but molecular work from the

demosponge *Amphimedon queenslandica* revealed a diurnal expression of two metazoan

circadian clock genes suggesting that it could be at least partly intrinsically regulated

(Jindrich et al., 2017). Not surprisingly there seems to be a lot of variation between species

and an *in situ* study on the demosponge *Aplysina aerophoba* did not find any diurnal rhythms

in pumping activity (Pfannkuchen et al., 2009).

In this study we have assessed the indicated diurnal rhythm in the body contraction-

expansions of the sponge *T. wilhelma*, and tested the potential influence of the ambient light

intensity. We confirmed the diurnal rhythms in the contractions and through manipulations of

the light regime we also found evidence for the rhythm to at least largely be controlled

extrinsically by the light level.

Materials and methods

<u>Culturing of sponges</u>

Specimens of *Tethya wilhelma* Sarà et al., 2001 were collected at the National Aquarium of

Denmark, the Blue Planet, (Copenhagen, Denmark) in a coral reef display tank with sea

water of 24°C and 33 psu and transported to Aarhus University. The specimens had a size of

0.5-1.0 cm in diameter. After arrival, they were transferred to individual wells of a six well

culture plate with a 2.5 cm diameter black plastic disk placed at the bottom of each well for

sponge attachment. The well plate was then placed in a 25-litre glass tank with seawater

collected at Lemvig, Denmark, adjusted to 32-34 psu using artificial sea salt (Instant Ocean,

Aquarium Systems, France). The water temperature was kept at 22°C controlled by the room

temperature. The water was filtered and circulated using an EHEIM eXperience 150 external

filter pump incl. a biological filter (EHEIM GmbH & Co. KG, Germany). Light was supplied

with a Fluval Plant Nano Bluetooth LED light (Rolf C. Hagen Inc., Canada), which was

programmed to simulate sunrise from 6:00 AM to 7:00 AM, daylight from 7:00 AM to 5:00

PM, sunset from 5:00 PM to 6:00 PM, and dark from 6:00 PM to 6:00 AM. This corresponds

to a light regime of 12 hours of complete darkness, and 12 hours of varying light. The light

pre-set "Tropical River" was used at 1/10 of the original light intensity to prevent algal

growth. Approximately one third of the water in the tank was exchanged every three weeks

throughout the experimental period.

<u>Experimental setup</u>

The sponges were transferred to a submerged clear acrylic box (12/17/4.8 cm, H/W/D) inside

the holding tank. Light could enter directly from above and from the sides but the back was

covered with black fabric to provide a uniform dark background for video recordings. A

weight was placed on top of the box to make sure it wasn't moved by the water currents in

the tank during the video recordings. The sponges were video recorded from the outside of

the tank to document changes in the body sponge volume measured as changes in the

projected area. Recordings were made with a Panasonic HC-VX980 digital camcorder

(Panasonic Corp., Japan) using the night mode setting. Two infrared LED light sources (peak

emission at 940 nm) were used for illumination during the night (Fig. S1).

<u>Test of diurnal rhythm</u>

To characterize the contraction pattern of *Tethya wilhelma* under simulated natural dark/light

conditions, six specimens were recorded for 72 hours each. One at a time, the sponges were

transferred to the recording chamber at least three hours before the start of each recording

session to mimimize the influence of the handling. All recordings were started at 6:00 PM on

the first day and after 72 hours of recording, the sponge was transferred from the recording

chamber and back into the well plate. The influence of the light regime on contraction cycles

in the sponge *T. wilhelma* was investigated by subjecting the six specimens to 72 hours of

constant light and 72 hours of constant dark. These two treatments were separated by a

recovery period with a 12:12 hour dark:light cycle for seven days. During the recovery

period, the sponges were placed in another tank. Three sponges were tested at the same time

and they were again transferred to the recording chamber at least three hours before the start

of the experiment. Because the experimental period started at 6:00 PM, a simulated sunset

would be included at the onset of the constant darkness experiments but not the constant light

experiment. To ensure complete control of the light environment in the experimental tank it

was covered by a light proof box also including the camera and light sources.

<u>Processing of video material</u>

Single frames were grabbed from the videos with a time resolution of 100 s and transferred to

the software program ImageJ (Rasband, 2020) as an image stack, and a scale bar was added

using the plate the sponges were attached to as a reference. The images were converted into

8-bit greyscale, cropped, and had the brightness and contrast adjusted to optimize the contrast

between sponge and background. Based on the difference in contrast between the bright

sponge and the dark background, the part of the image containing the sponge could be

selected by converting the image to a binary image by setting an appropriate threshold and

applying it to the image stack. The projected area of the sponge was then measured using a

"Measure Stack" macro in ImageJ.

<u>Analysis of contraction patterns</u>

The area measuremensts of each sponge were normalized by the maximum area during the

72-hour period to obtain a relative projected area (RPA) (see Goldstein et al. (2020)). The

mean RPA for each 12-hour period (6:00 AM to 6:00 PM and 6:00 PM to 6:00 AM) was

calculated and the number of fully contracted states, indentified as RPA minima, were

counted for each 12-hour time period as well. The mean contraction amplitude was calculated

for each 12-hour time period as the mean change in RPA between the initiation of the

contractions and the fully contracted states during that time period. The mean contraction

area was calculated as the mean RPA of the fully contracted states during each 12-hour

period. Lastly, the minimum and maximum RPA during each 12-hour time period was

identified. If a contraction spanned both dark and light period then it was assigned to the

period where the contraction was initiated. The sponges are constantly making small changes

of their volume and accordingly a threshold of minimum 7% reduction in RPA was set before

it was considered a contraction. This threshold was choosen, as contractions with a greater

than 7% change in RPA displayed the typical shape of a contraction, with a rapid contraction,

followed by a slower expansion phase. A combined graph was produced for each of the three

experiments, combing the individal specimens (N=5/6) and the individual days (N=3) into

one graph to reveal possible consistent patterns in the timing of the individual contractions.

Here 95% confidence intervals were reported.

 Statistical analyses

The statistical analyses were performed in RStudio version 2022.02.1 (RStudio Team, 2022),

using the R packages "lme4" (Bates et al., 2015) and "lmerTest" (Kuznetsova et al., 2017).

Plots were made using the package "ggplot2" (Wickham, 2016) and Microsoft Excel. Linear

mixed-effect models (LMEM's) were fitted, that met the assumptions of normal distribution,

homoscedasticity, and independence of residuals. As response variables, mean RPA, mean

contraction amplitude, mean RPA of contractions, minimum RPA, and maximum RPA were

used. As explanatory variable, day period (either day or night) was added as a fixed factor,

and sponge ID and day number were added as random effects. For the analysis of the number

of fully contracted states, a generalized linear mixed model (GLMM) with poisson

distribution was fitted. This model met the same assumptions and contained the same

explanatory variables and random effects as the LMEMs. In the comparisons between

experiments (dark/light, constant dark, and constant light), experiment was added as a fixed

factor instead of period, and *post-hoc* tests were performed by multiple comparisons of

means using Tukey contrasts and single-step method for adjusting *p*-values using the 'glht'-

function from the R package "multcomp" (Hothorn et al., 2008). The critical p-value was set

to 0.05 for all tests. The standard deviations (± s.d.) are calculated based on number of

specimens with the experimental days pooled together.

Results

Contraction behaviour under normal conditions and constant dark/light

One sample had to be excluded from the analysis in the constant dark experiment because the

sponge moved out of the focus plane of the scale bar. The sample sizes were therefore as

follows; dark/light experiment (N=6), constant dark experiment (N=5), and constant light

experiment (N=6).

The summarized contractions-expansion diagrams showed a pattern of one contraction

happening consistently shortly after the time of sunrise in the simulated dark/light-experiment

(Fig. 1A, Fig. S2). This pattern was not observed in neither the constant dark, nor the

constant light experiment (Fig. 1B, C, Fig. S3, S4).

When sponges were kept under normal light conditions, the mean RPA was 0.873 (±0.026)

for the night periods and 0.803 (±0.050) for the day periods which is a significant difference

(LMEM, t=3.6, p=0.0011) (Fig. 2A). The mean number of fully contracted states also

differed significantly between night and day periods (LMEM, z=-2.4, p=0.016) and the mean

was 2.22 (±0.78) during night periods and 3.61 (±0.77) during day periods (Fig. 2B). The

relative contraction aplitude was 0.214 (±0.076) during the night and 0.190 (±0.058) during

the day (Fig. 2C). This difference was not significant (LMEM, t=1.2, p=0.239). The mean

contraction RPA was also not significantly different at night compared to daytime, 0.705

(±0.073) and 0.676 (±0.083) respectively (LMEM, t=1.6, p=0.135) (Fig. 2D). The mean

minimum RPA was 0.702 (±0.062) during the night period and 0.645 (±0.087) during the day

period respectively, which is a significant difference (LMEM, t=2.3, p=0.028) (Fig. 2E).

Finally, the maximum RPA did not differ between night period, 0.941 (±0.032), and day

period, 0.933 (±0.020) (LMEM, t=0.4, p=0.684) (Fig. 2F). Under constant darkness and

constant light conditions, no significant differences were found between night period and day

period in any of the examined parameters (Table 1, Fig. 2A-F).

Normal light regime vs constant dark/light

When comparing the timewise day periods between the experiments, the mean RPA was

significantly higher in the constant dark experiment compared to the dark/light according to

the linear mixed-effect model (LMEM, t=-2.3, p=0.040), but in the multiple comparisons test

it is only a trend (Tukey comparisons *post hoc*: p=0.054) (Fig. 2A). Also, minimum RPA was

significantly higher in the constant dark experiment compared to the dark/light experiment

according to the linear mixed-effect model (LMEM, t=-2.3, p=0.047), but again only a trend

according to the multiple comparisons test (Tukey comparisons *post hoc*: p=0.055) (Fig. 2E).

When comparing the timewise night periods between the three experiments, the number of

fully contracted states was significantly higher in constant dark experiment compared to the

dark/light experiment (Tukey comparisons *post hoc*: p<0.025) (Fig. 2B), and the mean

relative contraction amplitude was significantly lower in the constant dark experiment

compared to the dark/light experiment (Tukey comparisons *post hoc*: p=0.028) (Fig. 2C). The

constant light experiment did not differ significantly from the light/dark experiment for any

of the parameters (Fig. 2A-F).

Discussion

The present study shows a diurnal rhythm in the contraction patterns of *Tethya wilhelma*.

Specifically when comparing night and day periods, the mean RPA was significantly higher

during the night (Fig. 2A), the number of fully contracted states was significantly higher

during the day (Fig. 2B), and finally, the minimum RPA measurements during each 12-hour

time period was significantly higher during the night (Fig. 2E). These differences suggest that

T. wilhelma is more expanded during the night, and is more actively contracting during the

day. The constant light and constant dark experiments on *T. wilhelma* showed, that any

significant differences found in the dark/light experiment disappeared under these constant

light regimes. In complete darkness the number of fully contracted states is similar to daytime

in the dark/light experiment and significantly higher than at nighttime indicating that

prolonged darkness increases the frequency of contractions. In addition, the amplitude of the

contractions during the night period in the constant darkness experiment is significantly

smaller than the amplitude of the contractions during nighttime under normal light

conditions, showing that in addition to the contractions happening more frequently, they are

also smaller (Fig. 2C). The multiple comparisons *post-hoc*-tests failed to show differences

among experiments in mean RPA and minimum RPA during the timewise dayperiods. This is

likely due to low statistical power caused by unequal sample sizes between experiments, and

as the differences in means show, effects sizes are clearly large and confirm that the sponges

are more expanded during darkness, than during light (Fig. 2A, E).

The finding from the present study, that *T. wilhelma* has a diurnal variation in contraction

patterns is in line with another study on this species by Nickel (2004), where projected area

measurements were also used to count and compare the number of contractions during day

and night. However, the study by Nickel (2004) was limited to two specimens, so the present

study is the first to statistically prove this diurnal rhythm. In addition, our results show that

this diurnal rhythm is not restricted to the number of contractions alone but is also present in

two additional parameters; minimum and mean relative projected area. It should be noted,

though, that these three parameters are not completely independent, since more contractions

equal more minima, and more time spent in a contracted state leads to lower mean RPA. For

functional considerations of the contractions in sponges it is important that although

contractions result in changes of the overall volume of the sponge, these changes mainly

concern the volume of the water canal system (Nickel et al., 2011). Sponge contractions have

been suggested to serve different functions, including servering as a cleaning mechanism to

expel unwanted particles by flusing the water canal system (Elliott and Leys, 2007; Nickel,

2004). In this case diurnal periodicity in contraction-expansion behaviour could be related to

maintenance of the water canal system. Further studies are needed to reveal the relationship

between diurnal activity patterns and individual elements of the water canal system and to

test the hypothesis that *T. wilhelma* is more actively filtering during daytime and thus needs

more flushing in this period.

 Control of the diurnal rhythm

Since our study shows the absence of diurnal rhythms in *T. wilhelma* under constant dark and

constant light conditions, it implicates that contraction behaviour is, at least in part, regulated

exogenously by the ambient light intensity, and as a consequence, that this sponge has light

sensing capabilities. Nickel (2004) used an external flash to capture images of the contraction

states of *T. wilhelma*. Futher studies are needed to determine a possible behavioural impact of

exposure to short flashes of light during the night, and clarify if this could influence the

natural contraction patterns by, for instance, increasing the frequency of contractions. In our

study we used an infrared lightsource to avoid this possible impact. Also, our use of a light

capable of simulating sunrise/sunset results in more natural changes of light levels. Indeed,

we found a consistent pattern of a contraction happening shortly after simulated sunrise. This

behaviour was not reported by Nickel (2004) who used an artificial 12:12 light:dark cycle.

This further strengthens our hypothesis, that *T. wilhelma* has light sensing capabilities, and

that sunrise could serve as a 'guide' for when to increase activity i.e. contractions, potentially

flushing the canal system to be ready for increased daytime filtration.

The diurnal rhythm in *T. wilhelma* is in line with genomic studies on another demosponge

Amphimedon queenslandica. Here, diurnal variations were found in expression of two

circadian clock genes, but the variations became less pronounced when the sponge was

subjected to constant darkness (Jindrich et al., 2017). This suggest at least some endogenous

control of diurnal rhythms alongside the light control shown by our results. Because many

sponge species have a diverse microbiome including photosynthesizing symbionts

(Bickmeyer et al., 2019; Thomas et al., 2016), the diurnal rhythm of the sponge could be

influenced by this microbiome, and decreasing levels of oxygen inside the sponge at night

could account for the diurnal rhythm in behaviour. This however is contradicted by the very

similar patterns observed under constant darkness and constant light. Light seems to play a

similar role in the cnidarians *Hydra vulgaris, Tripedalia cystophora,* and *Copula sivickisi*

that all show diurnal rhythms in behaviour. Also, the diurnal activity patterns disappeared

under either constant light or constant dark conditions in *H. vulgaris* (Garm et al., 2012;

Kanaya et al., 2020). Changes in behaviour in response to light is also seen among the

sistergroup to animals, the choanoflagellates. Brunet et al. (2019) revealed that colonies of

the choanoflagellate *Choanoeca flexa* inverted when subjected to changing light levels,

showing that even simple holozoans are capable of modulating their behaviour in response to

light, demonstrating that such behaviour in sponges is not at all unexpected.

Acknowledgements

We are grateful to the National Aquarium of Denmark, the Blue Planet, for providing animals for our study, and to Dorthe Due Theilade for transport to our laboratory. In addition, we are grateful to Alexandra B. Flensburg for help with designing the experimental setup, and to Josephine Goldstein and Clémence Rose for helping with the statistical analyses.

Competing interests

The authors declare no competing or financial interests.

Author contributions

Conceptualization: S.B.F, A.G., P.F.; Methodology: S.B.F, A.G., P.F.; Formal analysis: S.B.F.; Investigation: S.B.F.; Resources: A.G., P.F.; Writing - original draft: S.B.F.; Writing - review and editing: S.B.F, A.G., P.F.; Visualization: S.B.F.; Supervision: A.G, P.F.; Funding acquisition: P.F.

Funding

This work was funded by a research grant (40834) from VILLUM FONDEN and by the Independent Research Fund Denmark (grant number 8021-00392B).

References

Amano, S. (1986). Larval release in response to a light signal by the intertidal sponge *Halichondria panicea*. *Biological Bulletin* **171**, 371-378.

Amano, S. (1988). Morning release of larvae controlled by the light in an intertidal sponge, *Callyspongia ramosa*. *Biological Bulletin* **175**, 181-184.

Asadzadeh, S. S., Kiorboe, T., Larsen, P. S., Leys, S. P., Yahel, G. and Walther, J. H. (2020). Hydrodynamics of sponge pumps and evolution of the sponge body plan. *Elife* **9**.

Bates, D., Machler, M., Bolker, B. M. and Walker, S. C. (2015). Fitting linear mixed-effects models using lme4. *Journal of Statistical Software* **67**, 1-48.

Bickmeyer, U., Thoms, S., Koch, F., Petety Mukagatare, L., Silalahi, R. and Sartoris, F. J. (2019). Enhancement of photosynthesis in *Synechococcus bacillaris* by sponge-derived Ageladine A. *Plos One* **14**, e0213771.

Bond, C. (2013). Locomotion and contraction in an asconoid calcareous sponge. *Invertebrate Biology* **132**, 283-290.

Brunet, T., Larson, B. T., Linden, T. A., Vermeij, M. J. A., McDonald, K. and King, N. (2019). Light-regulated collective contractility in a multicellular choanoflagellate. *Science* **366**, 326-+.

De Voogd, N. J., Alvarez, B., Boury-Esnault, N., Carballo, J. L., Cárdenas, P., Díaz, M.-C., Dohrmann, M., Downey, R., Hajdu, E., Hooper, J. N. A. et al. (2021). World Porifera Database. Accessed at http://www.marinespecies.org/porifera on September 30, 2021. doi: 10.14284/359

Elliott, G. R. D. and Leys, S. P. (2007). Coordinated contractions effectively expel water from the aquiferous system of a freshwater sponge. *Journal of Experimental Biology* **210**, 3736-3748.

Elliott, G. R. D. and Leys, S. P. (2010). Evidence for glutamate, GABA and NO in coordinating behaviour in the sponge, Ephydatia muelleri (Demospongiae, Spongillidae). *Journal of Experimental Biology* **213**, 2310-2321.

Ellwanger, K., Eich, A. and Nickel, M. (2007). GABA and glutamate specifically induce contractions in the sponge *Tethya wilhelma*. *Journal of Comparative Physiology a-Neuroethology Sensory Neural and Behavioral Physiology* **193**, 1-11.

Ellwanger, K. and Nickel, M. (2006). Neuroactive substances specifically modulate rhythmic body contractions in the nerveless metazoon *Tethya wilhelma* (Demospongiae, Porifera). *Frontiers in zoology* **3**, 1-14.

Garm, A., Bielecki, J., Petie, R. and Nilsson, D. E. (2012). Opposite patterns of

diurnal activity in the box jellyfish *Tripedalia cystophora* and *Copula sivickisi*. *Biological*

Bulletin **222**, 35-45.

Goldstein, J., Bisbo, N., Funch, P. and Riisgard, H. U. (2020). Contraction-

expansion and the effects on the aquiferous system in the demosponge *Halichondria panicea*.

Frontiers in Marine Science **7**.

Hoppe, W. F. and Reichert, M. J. M. (1987). Predictable annual mass release of

gametes by the coral-reef sponge *Neofibularia nolitangere* (Porifera, Demospongiae). *Marine*

Biology **94**, 277-285.

Hothorn, T., Bretz, F. and Westfall, P. (2008). Simultaneous inference in general

parametric models. *Biometrical Journal* **50**, 346-363.

Jindrich, K., Roper, K. E., Lemon, S., Degnan, B. M., Reitzel, A. M. and Degnan,

S. M. (2017). Origin of the animal circadian clock: diurnal and light-entrained gene

expression in the sponge *Amphimedon queenslandica*. *Frontiers in Marine Science* **4**.

Kanaya, H. J., Park, S., Kim, J. H., Kusumi, J., Krenenou, S., Sawatari, E., Sato,

A., Lee, J., Bang, H., Kobayakawa, Y. et al. (2020). A sleep-like state in *Hydra* unravels

conserved sleep mechanisms during the evolutionary development of the central nervous

system. *Science Advances* **6**.

Kealy, R. A., Busk, T., Goldstein, J., Larsen, P. S. and Riisgard, H. U. (2019).

Hydrodynamic characteristics of aquiferous modules in the demosponge *Halichondria*

panicea. *Marine Biology Research* **15**, 531-540.

Kuznetsova, A., Brockhoff, P. B. and Christensen, R. H. B. (2017). lmerTest

package: tests in linear mixed effects models. *Journal of Statistical Software* **82**, 1-26.

Lavrov, A. I. and Kosevich, I. A. (2014). Sponge cell reaggregation: Mechanisms

and dynamics of the process. *Russian Journal of Developmental Biology* **45**, 205-223.

Leys, S. P. (2015). Elements of a 'nervous system' in sponges. *Journal of*

Experimental Biology **218**, 581-591.

Leys, S. P., Cronin, T. W., Degnan, B. M. and Marshall, J. N. (2002). Spectral

sensitivity in a sponge larva. *Journal of Comparative Physiology a-Neuroethology Sensory*

Neural and Behavioral Physiology **188**, 199-202.

Ludeman, D. A., Farrar, N., Riesgo, A., Paps, J. and Leys, S. P. (2014).

Evolutionary origins of sensation in metazoans: functional evidence for a new sensory organ

in sponges. *Bmc Evolutionary Biology* **14**.

Mah, J. L. and Leys, S. P. (2017). Think like a sponge: The genetic signal of sensory

cells in sponges. *Developmental Biology* **431**, 93-100.

Musser, J. M., Schippers, K. J., Nickel, M., Mizzon, G., Kohn, A. B., Pape, C.,

Ronchi, P., Papadopoulos, N., Tarashansky, A. J. and Hammel, J. U. (2021). Profiling

cellular diversity in sponges informs animal cell type and nervous system evolution. *Science*

374, 717-723.

Nath, R. D., Bedbrook, C. N., Abrams, M. J., Basinger, T., Bois, J. S., Prober, D.

A., Stemberg, P. W., Gradinaru, V. and Goentoro, L. (2017). The jellyfish *Cassiopea*

exhibits a sleep-like state. *Current Biology* **27**, 2984-2990.

Neely, K. L. and Butler, C. B. (2020). Seasonal, lunar, and diel patterns in spawning

by the giant barrel sponge, *Xestospongia muta. Coral Reefs* **39**, 1511-1515.

Nickel, M. (2004). Kinetics and rhythm of body contractions in the sponge *Tethya*

wilhelma (Porifera : Demospongiae). *Journal of Experimental Biology* **207**, 4515-4524.

Nickel, M. (2006). Like a 'rolling stone': quantitative analysis of the body movement

and skeletal dynamics of the sponge *Tethya wilhelma. Journal of Experimental Biology* **209**,

2839-2846.

Nickel, M., Scheer, C., Hammel, J. U., Herzen, J. and Beckmann, F. (2011). The

contractile sponge epithelium sensu lato - body contraction of the demosponge *Tethya*

wilhelma is mediated by the pinacoderm. *Journal of Experimental Biology* **214**, 1692-1698.

Pfannkuchen, M., Fritz, G. B., Schlesinger, S., Bayer, K. and Brummer, F.

(2009). In situ pumping activity of the sponge *Aplysina aerophoba*, Nardo 1886. *Journal of*

Experimental Marine Biology and Ecology **369**, 65-71.

Rasband, W. S. (2020). ImageJ: U.S. National Institutes of Health, Bethesda,

Maryland, USA.

Rivera, A. S., Ozturk, N., Fahey, B., Plachetzki, D. C., Degnan, B. M., Sancar, A.

and Oakley, T. H. (2012). Blue-light-receptive cryptochrome is expressed in a sponge eye

lacking neurons and opsin. *Journal of Experimental Biology* **215**, 1278-1286.

RStudio Team. (2022). RStudio: Integrated Development for R: RStudio, Inc.,

Boston, Massachusetts, USA.

Sarà, M., Sarà, A., Nickel, M. and Brümmer, F. (2001). Three new species of

Tethya (Porifera: Demospongiae) from German aquaria. *Stuttgarter Beiträge zur Naturkunde*

(A, Biologie) **631**, 1-15.

Siegel, J. M. (2008). Do all animals sleep? *Trends in Neurosciences* **31**, 208-213.

Thomas, T., Moitinho-Silva, L., Lurgi, M., Björk, J. R., Easson, C., Astudillo-García, C., Olson, J. B., Erwin, P. M., López-Legentil, S. and Luter, H. (2016). Diversity, structure and convergent evolution of the global sponge microbiome. *Nature Communications* **7**, 1-12.

Van Soest, R. W. M., Boury-Esnault, N., Vacelet, J., Dohrmann, M., Erpenbeck, D., De Voogd, N. J., Santodomingo, N., Vanhoorne, B., Kelly, M. and Hooper, J. N. A. (2012). Global diversity of sponges (Porifera). *Plos One* **7**.

Wickham, H. (2016). ggplot2: Elegant Graphics for Data Analysis. New York: Springer.

Figures and tables

Time (hours) after 6:00 PM

Figure 1. Summarized contraction-expansion graphs of specimens of *T. wilhelma*. Mean
RPA over three days for all specimens are shown with thick lines and 95% confidence
interval with thin lines. A) Simulated night/day-cycle (N=6), B) constant dark (N=5) and C)
constant light (N=6). Dark grey indicates night period, light grey indicates periods with
sunrise/sunset.

Figure 2. Influence of light regime on contraction behaviour of *T. wilhelma*. Differences
between experiments, and between night period (dark grey) and light period (light grey) in
mean RPA (A), number of fully contracted states (B), relative contraction amplitude (C),
mean contraction RPA (D), minimum RPA (E), and maximum RPA (F). Asterisks indicate a
significant difference; *=p<0.05, **=p<0.01. Points are individual specimens (N=6 for DL
and DD, N=5 for LL), bars are means. DD = constant dark, DL = dark/light, and LL =
constant light.

462 **Table 1. Results from constant dark and constant light experiment in *T. wilhelma*.** Mean

463 (± s.d.) values for each of the six examined parameters during the night periods and the day

464 periods. For the comparisons between periods, *t*- and *p*-values are presented, exept for no.

465 fully contracted states, where *z*- and *p*-values are presented.

Constant darkness experiment				
	Night period	**Day period**	***t (z)***	***p***
Mean RPA	0.852 (±0.028)	0.875 (±0.028)	-1.5	0.155
No. Fully contracted states	3.8 (±0.61)	3.93 (±0.86)	-0.2	0.853
Rel. contraction amplitude	0.135 (±0.010)	0.150 (±0.020)	-1.0	0.314
Mean contraction RPA	0.755 (±0.032)	0.767 (±0.042)	-1.0	0.350
Min. RPA	0.723 (±0.039)	0.748 (±0.035)	-1.6	0.131
Max. RPA	0.926 (±0.024)	0.946 (±0.024)	-1.1	0.280
Constant light experiment				
	Night period	**Day period**	***t (z)***	***p***
Mean RPA	0.833 (±0.041)	0.841 (±0.055)	-0.5	0.652
No. Fully contracted states	3.39 (±0.49)	3.5 (±0.78)	-0.2	0.857
Rel. contraction amplitude	0.169 (±0.017)	0.168 (±0.013)	0.03	0.977
Mean contraction RPA	0.725 (±0.043)	0.734 (±0.046)	-0.7	0.485
Min. RPA	0.708 (±0.037)	0.705 (±0.040)	0.3	0.746
Max. RPA	0.931 (±0.037)	0.939 (±0.028)	-0.4	0.668