Oxytocin/vasopressin system of insects influences parental care in males but not females

Ahva L. Potticary*[1], Christopher B. Cunningham[1], Elizabeth C. McKinney[1], Patricia J. Moore[1], Amsale

T. Belay[2], and Allen J. Moore[1]

[1]Department of Entomology, University of Georgia, Athens, Georgia, 30602 USA

[2]Department of Ecology and Evolutionary Biology, University of Toronto, Toronto, Ontario, M5S 3B2

Canada

*Corresponding author

Ahva L. Potticary

Dept. of Entomology

120 Cedar Street

Biological Sciences Building #360

University of Georgia

Athens, GA 30602

+1.406.529.3416

ahva.potticary@uga.edu

ORCID identifier

Ahva L. Potticary: 0000-0002-1157-5315

Christopher B. Cunningham: 0000-0003-3965-2076

Patricia J. Moore:

Amsale T. Belay:

Allen J. Moore: 0000-0002-1498-3322

Keywords: gene duplication, parental care, burying beetle, inotocin

This PDF files includes:

Main Text

Figures 1 – 4

Supplementary Figure 1 - 2

Supplementary Table 1

Supplementary Information Appendix 1

Abstract

Gene duplication is thought to allow for the origin of novel gene functions. However, how duplication influences gene function is difficult to determine, as new adaptations may subsume previous traits and the ancestral state is often unknown. One solution is to investigate the functions of a gene that exists as a single and duplicated state in different species. Here, we leverage a system for which functions have been well-characterized in the duplicates – the oxytocin/vasopressin system in vertebrates – and that exists as a single copy – *inotocin* - in insects. To answer the question of how functions of a single copy are partitioned prior to gene duplication, we experimentally tested the relationship between *inotocin* gene expression and the known social and non-social functions of oxytocin/vasopressin in a subsocial beetle that expresses complex parental care. We found that *inotocin* was strongly related to water balance in both males and females, corroborating other evidence that water balance is an ancestral function. Moreover, we found that *inotocin* was associated with male, but not female, parental care states, influencing the expression of care behavior towards offspring in males. Lastly, using immunohistochemistry, we show that inotocin is broadly distributed in the brains of both female and male parents. These results suggest that the ancestral social function of the vasopressin system may have been to moderate flexibility in male, rather than female behavior, and the role of oxytocin/vasopressin in female-offspring interactions is more recent.

Significance Statement

Oxytocin/vasopressin are amongst the most studied neuropeptides in vertebrates, influencing water

balance, mating interactions, and most notably, social bonding. This gene pair evolved from a duplication

in the vertebrate lineage of an ancestral vasopressin-like gene. Are the multiple social effects in

vertebrates due to this duplication, or are social influences also ancestral? Here, we demonstrate that, in a

biparental social beetle with a single copy, inotocin influences social interactions between fathers and

offspring as well as having the ancestral function of water balance in both males and females. In

vertebrates, influences on social interactions with males is associated with both oxytocin and vasopressin.

This may indicate that gene duplication allowed for the evolution of new behavioral functions in females.

Introduction

Gene duplication, or the production of two copies of an existing gene, is broadly thought to enable the

evolution of gene function (1, 2). The fate of duplicated genes has been inferred in many ways, such as by

using variation of function of recently duplicated genes (3), reconstructing the hypothetical sequence of

the ancestor (4), and correlating the timing of whole genome duplication with the advent of new complex

behavior (e.g., anadromy; (5)). Yet, determining the evolutionary fate of duplicated genes is challenging,

as new adaptations can subsume previous traits and the ancestral state is often unknown, making it

difficult to link historical and current gene functions. This is especially true for behavior, which often

cannot be inferred from a fossil record. These constraints have hindered an understanding how behavior

may evolve following gene duplication. One solution to this problem is to investigate the function of a

gene that exists as a single copy in some species and as a duplicate in others.

Here, we leverage a system for functions have been well-characterized in the duplicates – the

oxytocin/vasopressin system in vertebrates – and that exists in a single copy – inotocin - in insects. The

nonapeptides oxytocin and vasopressin evolved from a tandem duplication event of a vasopressin-like

gene (6-8) that occurred approximately 500 million years ago in the common ancestor of vertebrates (9),

although the oxytocin and vasopressin group of homologs are part of an ancient family of conserved

peptides that have existed for over 700 million years (10, 11). In vertebrates, oxytocin/vasopressin are

best known for their influence on the regulation of complex social behaviors (such as parental care and

pair bonding) and non-behavioral functions such as water balance and myoactivity (Fig. 1). Moreover,

both peptides demonstrate great evolutionarily lability in how each function is influenced between males

and females (SI Table 1; 10, 12-14), and oxytocin/vasopressin are often noted for their sex-specificity

within species. For example, oxytocin and vasopressin can influence aggression in both sexes, yet there is

not a consistent link between either peptide, the sex it acts in, and the expression of aggression across

vertebrate species (SI Table 1). However, while the sex-specificity and functions of oxytocin/vasopressin

are well-understood, insects only possess a single homolog of oxytocin/vasopressin (15-18), providing a

unique opportunity to determine how the observed functions of the duplicates (oxytocin/vasopressin) are

influenced by a single homolog (inotocin). We particularly focus on the function of inotocin in males and

females, as oxytocin/vasopressin are known to show sex-specificity in their function across vertebra taxa

(SI Table 1). While some functions for inotocin are known (Fig. 1), it is unknown whether inotocin

influences the expression of parental care.

To address these questions, we first reviewed the most highly conserved functions of

oxytocin/vasopressin group in vertebrates and the sex in which they act, including water balance, parental

and social behaviors (SI Table 1). We then investigated how the phenotypic functions of a single copy

could be partitioned prior to gene duplication using the biparental subsocial beetle *Nicrophorus orbicollis*.

Burying beetles require small animal carcasses to breed, where they perform extensive elaborate parental

care. Upon finding a carcass, parents bury it, remove the exterior (e.g., fur), and coat it with an

antimicrobial exudate (19). Females lay eggs in the soil and upon hatching, larvae crawl to the carcass.

Burying beetles identify their larvae temporally; parents will accept and feed larvae only after a certain

amount of time has elapsed since egg laying has occurred (20). If larvae appear on the carcass too early,

the breeding pair will not provide care to the offspring and will eat them instead. Parental care in burying

beetles involves both direct (feeding of regurgitated food to their offspring) and indirect parental care

(maintenance of carcass while larvae are present, but does not involve social interactions). In general,

males trade-off indirect care for direct care while females are consistent in their expression of both (21,

22). Larvae of *N. orbicollis* cannot survive without parental care and while females are highly consistent

in their care, males show more environmentally dependent care behavior (e.g., loss of partner; 22, 23).

Parental behavior in *N. orbicollis* thus requires nesting behavior, egg laying, a transition from infanticide

to acceptance of larvae, and larval feeding, concordant with the behavioral functions of

oxytocin/vasopressin in vertebrates (Fig. 1).

We leveraged this variation across the lifetime of *N. orbicollis* to address whether *inotocin* and

inotocin receptor gene expression are associated with the functions of oxytocin/vasopressin in

vertebrates. *inotocin* and *inotocin receptor* were related to water balance in both males and females,

corroborating other work indicating that water balance was one ancestral function of this neuropeptide

group. Isolated male virgins showed an increase in *inotocin* and *inotocin receptor,* corresponding to the

age when males have reached sexual maturity. Moreover, we found that *inotocin* was related to parental

care behavior in males but not in females. Males that were actively caring for larvae had higher *inotocin*

expression than males than males without larvae. Males that spent more time feeding their larvae had

higher *inotocin* than males that fed their larvae less, and higher *inotocin* than males that spent more time

engaged in indirect (non-social) parental care behaviors. Lastly, immunohistochemistry reveals that

inotocin is broadly expressed in the brains of both male and female parents. Together, these results may

suggest that duplication of an ancestral vasopressin-like neuropeptide allowed for neofunctionalization of

the duplicate (oxytocin) and the subsequent elaboration of novel gene function in females.

Results

Inotocin Associated with Non-social Contexts.

In females, neither the expression of *inotocin* ($F_{5,54} = 1.27$, P = 0.30; Fig. 2a) nor *inotocin receptor* ($F_{5,54} =$

1.44, P = 0.22; Fig. 2A) changed with adult age. In contrast, in virgin males, *inotocin* ($F_{5,54} = 3.45$, P =

0.009; Fig. 2B) and *inotocin receptor* ($F_{5,54} = 3.90$, P = 0.004; Fig. 2B) expression both changed across

this timespan. This difference was primarily driven by lower *inotocin* and *inotocin receptor* expression in

6-day old males relative to post-reproductive ages (day 21 for *inotocin* and days 21 and 35 for *inotocin*

receptor, Tukey's HSD; Fig. 2B). A trend of lower expression of *inotocin* and *inotocin receptor* was seen

in both males and females in day 6, but the differences were not large in either sex. Six days is prior to

sexual maturity in both sexes.

The relative humidity of the two water balance treatments was lower and higher than the average

relative humidity experienced by beetles breeding with a carcass ($F_{2,57}$ = 1032.03, P < 0.0001); i.e., the

Low Humidity treatments averaged 15.39 ± 1.3 % RH, breeding beetles with a carcass had a RH of 72.99

± 0.79 %, and beetles in the High Humidity treatment experienced an average humidity of 84.8 ± 1.3 %

RH. Female beetles in Low Humidity treatments had trend towards higher *inotocin* ($t_{1,18}$ = -1.96, P < 0.07;

Fig. 2C), while Low Humidity females expressed higher *inotocin receptor* ($t_{1,18}$ = -2.12, P < 0.05; Fig.

2C), than females in the High Humidity treatment. Males in the Low Humidity treatment had higher

inotocin and *inotocin receptor* expression than males in the High Humidity treatment (*inotocin*: $t_{1,18}$ = -

2.99, P < 0.008; *inotocin receptor*: $t_{1,18}$ = -2.62, P < 0.02; Fig. 2D).

Inotocin in Social Contexts.

Females that accepted and cared for larvae did not differ in *inotocin* or *inotocin receptor* expression

relative to females that committed infanticide (*inotocin*: $t_{1,38}$ = 1.49, P = 0.14; *inotocin receptor*: $t_{1,38}$ =

0.34, P = 0.74). Female parental care states were not associated with *inotocin* ($F_{2,57}$ = 0.02, P = 0.98; Fig.

3A) or *inotocin receptor* expression ($F_{2,57}$ = 0.21, P = 0.81). Females parenting for the first time did not

have higher *inotocin* or *inotocin receptor* than females parenting for a second time (*inotocin*: $t_{1,38}$ = 0.37,

P = 0.71; *inotocin receptor*: $t_{1,37}$ = -0.14, P = 0.89). There was no relationship between the proportion of

time females spent feeding larva and *inotocin* or *inotocin receptor* expression (*inotocin*: $F_{1,32}$ = 0.17, r^2 =

0.005, P = 0.69, Fig. 3B; *inotocin receptor*: $F_{1,32}$ = 3.15, r^2 = 0.09, P = 0.09). There was no relationship

between indirect care provided by females and either *inotocin* ($F_{1,32}$ = 0.11, r^2 = 0.003, P = 0.74, Fig. 3B)

or *inotocin receptor* ($F_{1,33}$ = 0.0004, r^2 < 0. 0001, P = 0.98).

Males that committed infanticide did not differ in *inotocin* or *inotocin receptor* expression

compared to males that accepted larvae (*inotocin*: $t_{1,38}$ = -0.84, P = 0.41; *inotocin receptor*: $t_{1,38}$ = -0.08, P

= 0.94). However, males in the Care treatment had higher *inotocin* than males in the Nesting and Post

Care treatments ($F_{2,57}$ = 4.23, P = 0.02, Fig. 3c). We found no effect of male parental state on *inotocin*

receptor ($F_{2,57}$ = 0.24, P = 0.79). Males parenting for the first time did not have higher *inotocin* or *inotocin*

receptor than males parenting for a second time (*inotocin*: $t_{1,38}$ = -0.62, P = 0.54; *inotocin receptor*: $t_{1,39}$ =

0.56, P = 0.58). Moreover, males that spent more time feeding larva had higher *inotocin* expression than

males that fed less ($F_{1,14}$ = 8.91, r^2 = 0.39, P < 0.001; Fig. 3d). There was no relationship between the

amount of time males spent feeding larva and *inotocin receptor* ($F_{1,14}$ = 0.08, r^2 = 0.005, P = 0.79).

Amount of indirect care provided was negatively associated with *inotocin* expression in males ($F_{1,19}$ =

4.33, r^2 = 0.19, P = 0.05; Fig. 3d), but again there was no relationship between *inotocin receptor* and

amount of indirect care by males ($F_{1,19}$ = 0.06, r^2 = 0.003, P = 0.80).

Immunohistochemistry of inotocin in the brains of parents with larva showed diffused expression

in both male and females (Fig. 4). Staining was concentrated in the suboesophageal zone and adjacent to

the antennal lobes, in the vicinity of the mushroom bodies. There were no major differences observed in

either the pattern or density of staining in male versus female brains.

199   Discussion

Our results demonstrate that inotocin in *N. orbicollis* is associated with both non-social and social

contexts. Consistent with the water balance function of oxytocin/vasopressin across broad taxonomic

groups, we also found that *inotocin* is involved in water balance in both sexes (Fig. 1, SI Table 1). This

supports other research indicating that water balance was an ancestral function of the

oxytocin/vasopressin system (24, 25). However, the discovery that *inotocin* expression changes as beetles

age in isolated, male virgins, but not females, was unexpected (Fig. 2). In males, pre-reproductive males

had lower inotocin expression than post-reproductive stages. It is possible that *inotocin* is involved in

priming males for reproduction, but further research would be needed to determine how the development

of inotocinergic system may be related to reproduction.

Given the subsocial and complex parental care observed in this species, we predicted a social role

for inotocin as well. However, rather than a general role in parental care, we found that increased *inotocin*

expression was associated with male, but not female, parenting in *N. orbicollis*. This holds for both

transition from nesting to parental offspring provisioning, and back to a non-parenting state, as well as

overall amount of time attending to offspring and performing indirect care activities such as carcass

maintenance (Fig. 3). However, these social associations of *inotocin* did not hold for females. Overall

willingness to accept or commit infanticide of larvae was not associated with differences in *inotocin* or

inotocin receptor expression in either males or females, suggesting that inotocin has a role limited to

social interactions involved in male parenting behavior. Despite this, immunohistochemistry revealed that

brains of parenting males and females showed similar patterns of expression, with the majority of staining

occurring in the suboesophageal zone (SEZ) and the mushroom bodies (MB; Fig. 4).

Our data raise questions about the evolution of social influences in the oxytocin/vasopressin

system. There are three potential explanations for our results: first, that a role in male social interactions

involving parenting arose de-novo in burying beetles. This is possible but not parsimonious given the

distribution of social influences of oxytocin/vasopressin homologs across vertebrate taxa (SI Table 1).

Second, given both males and female social interactions are influenced by the oxytocin/vasopressin

system in vertebrates, it could be that the female role of inotocin was lost in burying beetles. However,

this is unlikely, given that other studies of female insects have not found a definitive influence of inotocin

on parenting behavior (26, 27). Finally, it may be that influences on male rather than female social

behavior is ancestral. A similar, male-specific role was found in *nematocin*, the single copy homolog for

oxytocin/vasopressin found in nematodes, where *nematocin* was found to function in the organization of

male reproductive behavior but not females (28). It is possible that *inotocin* has a similar function in

males, and acts to organize male parental care behavior or increase salience of offspring cues, and as

parental care is often far more variable in males than in females, the effects are only apparent in males.

Oxytocin/vasopressin have been found to increase the salience of social signals and group

formation in a context-dependent manner (29, 30), and are associated with plasticity in socially-mediated

behaviors (e.g., 31). In many insects that have parental care, there is little plasticity in performance of

care. For example, while workers transition from feeding and caring for larvae to foraging in eusocial

insects, they often do so in a temporal fashion moving from one task to the next in a developmental series.

In contrast, male burying beetles tend to be very flexible in care whilst females are more canalized (21,

22). It is possible that, in *N. orbicollis*, *inotocin* functions to draw male attention to offspring, but only

under specific conditions. Support for this idea is provided by the results that males with higher *inotocin*

fed their larva more (i.e., increased attention to larva) than males with lower *inotocin*, and males with

lower *inotocin* spent more time engaged in indirect care, where attention was focused on the maintenance

of the carcass instead of larva. This hints that there is some threshold of *inotocin* expression that

facilitates a transition from other kinds of care duties – e.g., nest defense, carcass preparation – to direct

care of larva, although future research is needed to disentangle this effect.

Immunohistochemistry shows diffuse staining for inotocin expression across the brains of both

males and female parents (Fig. 4), which does not exclude a social role for inotocin in females. Previous

work has indicated that inotocin is often produced in the SEZ, and inotocin in this location can be related

to water balance pathways (18, 27). Mushroom bodies have been associated with social behavior in

insects more broadly (32, 33), and for this reason, the expression of inotocin in the region of the

mushroom bodies may indicate that this is one mechanism by which parenting behavior is influenced.

However, future research would be needed to determine how the distribution of inotocin influences social

behavior in males and females.

Oxytocin was first discovered in female vertebrates in association with parturition (34), and

correspondingly, there has been a greater emphasis on the role of oxytocin in females with a lesser

emphasis on vasopressin and males (SI Table 1). For this reason, the discovery that an ancestral proxy

functions in male parental care behavior, but not female parental care, is contrary to initial expectations

for most researchers. Nevertheless, the results of this study support the idea that male influences are

ancestral, and that duplication of an ancestral vasopressin-like neuropeptide in vertebrates allowed for

neofunctionalization of the duplicate the elaboration and evolution of behavioral functions in females.

Multiple lines of evidence support the idea that the ancestral gene was more similar to *vasopressin* (6, 8)

and even *inotocin* is more structurally similar to the *vasopressin* group than *oxytocin*, despite its name

(17). The oxytocin/vasopressin system exhibits high levels of plasticity, sexual dimorphism, and species-

specificity across taxa (SI Table 1), hinting at sexually dimorphic functions of the ancestral peptide.

Methods

Insect Colony and Husbandry.

The *Nicrophorus orbicollis* used in this study were part of an outbred colony that is regularly

supplemented with wild-caught beetles from Whitehall Forest in Athens, GA, USA. All beetles are

maintained in an incubator (Percival, model no. I41VLC8, Perry, IA) at $22 \pm 1°C$ on a 14:10 hour

light:dark cycle to simulate summer breeding conditions. All individuals used for this study were bred in

the laboratory to ensure that parentage, age, and rearing conditions were known and standardized. We

housed beetles individually at larval dispersal in 9-cm-diameter by 4-cm-deep circular compostable deli

containers (EP-RDP5, Eco-Products) filled with ~2.5 cm of potting soil (Performance Organics All

Purpose In-Ground Soil, Miracle-Gro). Once larvae are placed into individual containers, they had no

further social interactions with other burying beetles until they were collected for the age series or until

they were allocated into the behavioral or physiological treatments. Individuals were defined as age Day 1

on the day of eclosion. Beetles were fed *ad libitum* organic ground beef twice a week following eclosion.

Inotocin and Non-social Contexts.

Age Series.

We collected whole heads early post-eclosion (Day 2-3), pre-sexual maturity (Day 6-7), after sexual

maturity and during an early reproductive age (Day 15-16), at a prime reproductive age (Day 21-23),

reproductive but late stage (Day 30-31), and reproductive but very late stage (Day 35-36). The post-

eclosion individuals were collected on Day 2-3 to ensure that the head was fully melanized, as individuals

are susceptible to water loss prior to hardening of the cuticle and *inotocin* is known to influence water

balance functions (18, 35). This age series corresponds to the timing when beetles were collected for each

behavior state and the water balance experiment (SI Fig. 1).

Water Balance.

To determine the relative humidity range that would be expected to induce water balance, we first

measured the relative humidity of 20 pairs of breeding *N. orbicollis* (10 at the nest/egg stage and 10 at the

larval stage) that had not been opened for one week to capture the range of humidity and temperature that

occurs over the breeding cycle on a decomposing mouse, using a handheld digital Preciva Humdity and

Temperature Reader (model no. HT154001). Based on these results, we ensured that our Low and High

Humidity treatments occurred at a relative humidity below and above this average relative humidity

(72.99 ± 1.16 % RH). The relative humidity of the Low Humidity treatments was maintained between 8-

25 % (15.39 ± 1.3 % RH) and High Humidity conditions were maintained between 80-95% (84.8 ± 1.3 %

RH; see Results). On Day 16, individuals were transferred to an individual container with approximately

2 cm of potting soil with perforations nested inside of a breeding box and sealed with parafilm. Low and

High humidity conditions were simulated by charging the outside box with either a) 12.5 g of $CaSO_4$

(Drierite, Hammond Drierite Co., Xena, OH) for the Low Humidity treatment, or b) a 3 x 4" cotton strip

soaked to saturation with deionized water for the High Humidity (after 36). Previous research employing

a similar set-up found that *Nicrophorus marginatus* began to die around 12 hours of exposure to a similar

CaSO$_4$-charged box (36). For this reason, beetles were left in Low Humidity or High Humidity treatments

for 8 hours to provide sufficient time for water stress to occur and to ensure that beetles survived until the

end of the treatment period. To account for the fact that burying beetles experiencing water stress may

bury themselves in soil to evade low humidity, individuals in the Low Humidity experiment were kept in

individual containers without soil to ensure that they could not mitigate exposure by burrowing into the

soil. The relative humidity and temperature of each set-up was measured at the end of the 8-hour period.

Inotocin and Social Behavior.

We randomly mated focal virgins by placing an unrelated male and female in a small plastic container

and observed to ensure that mating occurred at Day 14-15. The pair was then moved to a plastic breeding

box filled with approximately two cm of potting soil moistened with distilled water and containing a

freshly thawed mouse between 19-25 grams (Rodent Pro, Inglefield, IN, U.S.A.). Breeding boxes were

returned to the incubator and kept on a darkened shelf. Breeding boxes were checked for eggs every

twenty-four hours. Once eggs were observed, the focal female and the brood ball were moved to a new

breeding box filled with approximately two cm of potting soil. The focal male was kept separate from the

female and brood ball for 24 hours and then returned to the breeding box with the female. Once the male

and female were reunited, breeding boxes were again checked daily for eggs.

Affiliation and Infanticide.

To test the hypothesis that *inotocin* influences the transition from aggression to affiliation towards larvae,

we compared males that committed infanticide with those that accepted and cared for larvae arriving on

the mouse resource. *Nicrophorus* species identify their offspring using temporal cues, such that larva that

arrive too early on the carcass are eaten and those that arrive on time are accepted by the parents and fed

(20). These treatments took advantage of this temporal recognition to sample adults as they committed infanticide of the larva or accepted and fed offspring, respectively. For both treatments, the breeding pair were provided with ten larvae, at least 5 of which were their offspring from the first clutch of eggs and were supplemented with same-age larva from other clutches if insufficient numbers were available from the focal pair's clutch. Following the addition of larvae, the focal individuals was observed for at least thirty minutes to detect acceptance or infanticide of larva. If the focal individuals did not respond in this time, pairs were checked every 30 minutes to identify the onset of acceptance or infanticide. In some cases, one parent would accept larva while the other parent committed infanticide; in these cases, the male and female were sampled and categorized separately according to the phenotype that was observed for each individual. For the treatment where the male committed infanticide, focal individuals were collected while the focal individual was attacking/eating larvae. For the treatment where larvae were accepted, focal individuals were collected immediately after the parent made antennal contact with the larva and a feeding event was observed. This manipulation often led to a discrepancy in the temporal cues used by males versus females, as females base their window of larval acceptance on when eggs were laid (20) and males likely base their window on timing and frequency of female acceptance of mating (unpub. data). Thus, to reach sample size, most males in the treatment where larvae were accepted were moved with the brood ball and females were removed after the first batch of eggs were laid, which increased the likelihood of male acceptance of larva.

Parental Care.

To test the hypothesis that *inotocin* or *inotocin receptor* influences parental care, we performed four experimental treatments including Nesting (pair mated and have prepared the mouse "nest", no larvae), Care (pair mated, have nest, and are providing care to larvae), Post Care (pair mated, have nest, and have provided care until larvae disperse), and Double Care (pair mated, have nest and larvae for two breeding attempts).

For the Nesting treatment, we collected whole heads of males and females 24-36 hours after the

male had been reunited with the female. For individuals in the Care treatment, eggs from the second

clutch were allowed to hatch and crawl to the brood ball naturally to ensure female acceptance of larva.

Each breeding box was checked every 12-24 hours to determine when larvae arrived on the brood ball.

The pair was allowed to interact with larvae for 24 hours, and then parental care behavior was scored

during a behavioral observation (see below *Behavioral observations*). Whole heads of the focal male and

female were collected 24 hours after the larva were first observed on the carcass (at age Day 21-23),

while the focal individual was located on the brood ball and engaged in direct care behavior. Male

parental care is variable and often depends on the presence and activity of the female (23, 37). For males

that did not show direct parental care during the initial behavioral observation, females were removed to

induce male parental care, and then males were collected 24 hours later while showing direct care to

larva. Individuals in the Post Care treatment were allowed to complete a breeding cycle. Following the

dispersal of larvae away from the carcass and the cessation of parental care behavior, the male and female

were housed singly in individual containers with soil for 24 hours, at which point whole heads were

collected on approximately Day 30.

To determine whether changes of *inotocin* and *inotocin receptor* expression were directly related

to the expression of care, and/or organized the capacity for expression of parental behavior across the

lifetime, pairs were allowed to breed twice creating a Double Care treatment. Double Care beetles were

removed for 24 hours following larval dispersal, and then put with a new mouse under the same

conditions. For the second breeding attempt, the pair was allowed to breed normally and were monitored

every 24 hours to determine egg lay date and the arrival of larvae. Twenty-four hours after hatch, the pair

was observed for 30-minutes and whole heads were collected as they were engaged in direct care

behavior (approximately Day 35). Similar to the observations performed for Care treatments, if a male did

not show direct care behavior during the initial behavioral observation, the female was removed for

collection and males were collected 24 hours later while engaged in direct care behavior. When mortality

of one parent occurred during or immediately after first breeding attempt, a new mate was provided to the

focal individual (n = 7 of 20 breeding attempts).

Behavioral Observations of Parental Care.

We conducted focal behavioral observations for each pair in the Care and Double Care treatments to

quantify parental care. Before an observation was initiated, we removed lids to breeding boxes and the

pair was given at least 30 minutes to acclimate. All observations were conducted under red lights to

ensure no light effects on individual care behavior, as *N. orbicollis* are nocturnal (38). In each behavioral

observation, we recorded both direct and indirect care (37). Direct care is defined as when the male or

female fed larvae. Indirect care is defined as activities related to carcass cleaning and maintenance, which

allows for the preservation of the carcass for future offspring provisioning and thus represents a type of

parental care without direct interactions with offspring. Time spent showing "no care" was also recorded,

and is defined as when the parent was on the carcass but not engaged in direct or indirect care, or was off

of the carcass. Behaviors were recorded as a frequency (e.g., number of feedings) and a duration (length

of time engaged in the behavior). These data were collapsed to align with previous research using

instantaneous scan sampling, where the activity of each beetle every minute was recorded. This gave each

individual three scores (direct care, indirect care, and no care) which together summed to 30. This method

was used because specific behaviors are often difficult to disentangle, e.g., larval feeding is often

interrupted by the adults "self-feeding", which is often followed by direct provisioning of larvae. For this

reason, continuous observation allowed for the observer to categorize behavior into direct or indirect care

by placing the behavior at each time point into its functional context. For males that did not show either

indirect or direct parental care during the behavioral observations (n = 13), the female was collected and

removed to initiate care behavior in the males. For these males, direct care of larvae was verified prior to

collection by observing at least one larval feeding event.

RNA Sample Collection and Analysis.

Whole Head Collection.

We collected samples with dissecting scissors, flash freezing them in liquid nitrogen (39), and

immediately placed in storage at -80 °C until RNA extraction. A small subset of heads (n = 5), heads were

collected, bisected with scissors to expose brain tissue, placed into 500 ml RNAlater (Ambion, Grand

Island, NY, USA), and stored at -20°C until RNA extraction.

Gene Expression Analysis for Inotocin and Inotocin Receptor.

We homogenized whole heads in liquid nitrogen using a mortar and pestle (cat. no. 60310, CoorsTek,

Golden, CO, USA). Following homogenization, we extracted RNA from each sample using Qiagen's

RNAeasy Lipid Tissue Mini-Kit (cat. no. 74106) following (39), including the DNase I (Qiagen)

treatment. RNA was quantified using a Qubit 2.0 Fluorometer (Invitrogen Corporation, Carlsbad, CA,

USA) using the RNA Broad Range protocols per manufacturer's instructions. We synthesized cDNA with

Quanta Biosciences qScript reverse transcriptase master mix (Quanta Biosciences, Gaithersburg, MD,

USA) following the manufacturer's instructions from 500 ng total RNA. RNA was stored at -80 °C and

cDNA was stored at -20 °C.

We performed quantitative real-time PCR (qRT-PCR) for each of our genes of interest and

several endogenous control genes (*gapdh* and *tbp*) following methods of Cunningham et al. (40) and

using transcriptome of Benowitz et al. (41). We performed qRT-PCR with Roche LightCycler 480 SYBR

I Green Master Mix using a Roche LightCycler 480 (Roche Applied Science, Indianapolis, IN, USA)

following the manufacture's protocol with triple technical replicates of 10 µl reactions (n = 20 for each

parental experimental context, n = 10 for in each water treatment, and n = 10 for each age group, for both

males and females). Primers were at a working concentration of at 1.33 µmol/l. We used an annealing

temperature of 60 °C during the amplification cycles. We established the stability of the endogenous

reference gene by running replicates from each treatment and assessing the stability of endogenous

reference gene amplicons by visual inspection and analysis of C_T values found that *gapdh* did not vary

substantially across social, water, or age contexts. Additional information suggested by the Minimum

Information for Publication of Quantitative Real-Time PCR Experiments (MIQE) guidelines can be found

in Appendix 1.

Immunohistochemistry of Inotocin in the Brain.

We dissected brains from female and male beetles at the period of peak parental care behavior, as

described above, for immunofluorescence microscopy in ice-cold 1x PBS. We fixed brains immediately

in 4% paraformaldehyde in PBT (PBS with 0.5% Triton X-100) for 24 hours at 4° C. Following fixation,

we permeabilized the brains by first incubating in 2N HCl for 30 minutes at 4° C. Brains were then rinsed

3 times in PBT to remove the HCl and permeabilization was continued by treating the brains in PBS with

4% Triton X-100 for 1 hour at room temperature, after which we washed brains the brains 3 times for 10

minutes in PBT. We blocked non-specific binding sites by incubating the brains in 1% bovine serum

albumin in PBT overnight at 4° C.

The brains were incubated in a 1:300 dilution of primary polyclonal antibody that recognizes the

pre-pro-peptide of inotocin (27) for 48 hours at 4° C. For all treatments (male and female parental care), a

subset of brains was not treated with primary antibody to validate the specificity of inotocin staining. We

then washed the brains 5 times in PBT for 1 to 3 hours per wash step at room temperature and added

secondary antibody solution (Alexa Fluor 647 donkey anti-goat, 1:500). We added 0.5 mg/mL DAPI at a

1:2000 dilution to the secondary antibody solution in order to counterstain for DNA and incubated the

brains in secondary antibody overnight at 4° C. We washed the brains 5 times for 1 to 3 hours each step.

After washing, we infiltrated the brains in mounting medium (90% glycerol in 0.1 M Tris HCl, pH 8.7)

by taking the brains through a dilution series of 3:1, 1:1, 1:3 PBT: mounting medium, followed by 90%

mounting medium with incubation steps of at least two hours each. Brains were then mounted on

depression well slides and imaged with a Zeiss LSM 710 Confocal Microscope at the UGA Biomedical

Microscopy Core. Brains processed in the absence of primary antibody showed no background

fluorescence due to non-specific binding of the secondary antibody (SI Fig. 2).

Statistical Analyses.

All statistical analyses were performed using JMP PRO (v. 16.0.0, http://jmp.com) and all figures were

produced in SigmaPlot (v. 14.5, http://www.sigmaplot.co.uk). Results are means ± SE. We used the $\Delta\Delta C_T$

method (42) to convert raw expression data to normalized relative expression values. For all analyses,

males were used as the comparison group for male analyses, and females were used as the comparison

group for female analyses. For the age series, virgin D21 were used as the control group. For all

behavioral states (Accept vs. Infanticide, Nesting, Care, and Post Care, and Care vs. Double Care) we

used the behavioral state that occurred earliest within the experiment as the comparison group

(Infanticide, Nesting, and Care, respectively). For the Low vs. High Humidity experiment, we used High

Humidity as the comparison group. Data were visually inspected for outliers. We tested for the effect of

Low vs. High Humidity, Accept vs. Infanticide, and Care vs. Double Care using pooled t-tests, and for the

effect of age and parental care states using Analysis of Variance (ANOVA). We determined the direction

of significance for age analyses using post hoc Tukey's HSD test.

We also assessed if there was a direct association between *inotocin* and *inotocin receptor*

expression and parental care behaviors observed in behavioral observations. Data from individuals in the

Care and Double Care treatments were pooled as there was no difference in *inotocin* and *inotocin*

receptor expression between treatments (see Results). The influence of *inotocin* and *inotocin receptor*

expression on care behavior (proportion of observation points spent in direct care) was analyzed using

linear regression for individuals that were observed to feed larva at least once during the targeted

behavioral observation without removing the mate (females: n = 34, males: n = 16). The influence of

inotocin and *inotocin receptor* expression on indirect parental care was analyzed using linear regression

for individuals that interacted with the brood ball or larvae in any way (females: n = 34, males: n = 21).

For one behavioral observation, the pair buried the carcass and the observation time was thus only 25

minutes; for this reason, all behavioral data are presented as a proportion of the observation time. A

subset of observations was excluded (n = 6) because male and female behavior was more indicative of a

response to perceived competition or disruption (i.e., frequent stridulation and both individuals off of the

carcass exploring breeding box).

Acknowledgements and funding sources

We are indebted to Daniel J. Kronauer for providing the inotocin antibodies for the

immunohistochemistry work. We would also like to thank Kathryn Kollars for producing the beetle art

featured in this manuscript. We would like to thank Hans Otto, Kyle Benowitz, Shannon Harris, Emily

Shelby, and Kathryn Kollars for discussions and/or feedback on the manuscript. A.L.P. and A.J.M.

conceived and designed the experiments. A.L.P. conducted experiments. A.L.P., E.C.M., P.J.M., A.T.B.,

and C.B.C. conducted laboratory work. A.L.P. and A.J.M analyzed the data. A.L.P. and P.J.M. produced

figures. All authors contributed to the writing of the manuscript. A.T.B. was funded by a Natural Sciences

and Engineering Research Council of Canada discovery grant to Marla Sokolowski.

References

1.   S. Ohno, *Evolution by gene duplication* (Springer-Verlag, New York, USA, 1970).
2.   L. E. Flagel, J. F. Wendel, Gene duplication and evolutionary novelty in plants. *New*
Phytol. **183**, 557-564 (2009).
3.   R. Assis, D. Bachtrog, Neofunctionalization of young duplicate genes in Drosophila.
Proc. Natl. Acad. Sci. U.S.A **110**, 17409-17414 (2013).
4.   H. Escriva *et al.*, Neofunctionalization in vertebrates: The example of retinoic acid
receptors. *PLOS Genet.* **2**, e102 (2006).

5.   M. A. Alexandrou, B. A. Swartz, N. J. Matzke, T. H. Oakley, Genome duplication and
multiple evolutionary origins of complex migratory behavior in Salmonidae. *Molec.*
Phylogenet. Evol. **69**, 514-523 (2013).
6.   J. L. Goodson, "Nonapeptides and the evolutionary patterning of sociality" in *Progress in*
Brain Research, I. D. Neumann, R. Landgraf, Eds. (Elsevier, 2008), pp. 3-15.
7.   C. Theofanopoulou, G. Gedman, J. A. Cahill, C. Boeckx, E. D. Jarvis, Universal
nomenclature for oxytocin–vasotocin ligand and receptor families. *Nature* **592**, 747-755
(2021).
8.   R. E. Van Kesteren *et al.*, Structural and functional evolution of the vasopressin/oxytocin
superfamily: Vasopressin-related conopressin is the only member present in Lymnaea,
and is involved in the control of sexual behavior. *J. of Neurosci.* **15**, 5989-5998 (1995).
9.   K. Yamashita, T. Kitano, Molecular evolution of the oxytocin–oxytocin receptor system
in eutherians. *Molec. Phylogenet. Evol.* **67**, 520-528 (2013).
10.   Z. R. Donaldson, L. J. Young, Oxytocin, vasopressin, and the neurogenetics of sociality.
Science **322**, 900-904 (2008).
11.   P. Banerjee, K. P. Joy, R. Chaube, Structural and functional diversity of nonapeptide
hormones from an evolutionary perspective: A review. *Gen. Comp. Endocrinol.* **241**, 4-
23 (2017).
12.   J. L. Goodson, R. R. Thompson, Nonapeptide mechanisms of social cognition, behavior
and species-specific social systems. *Curr. Opin. Neurobiol.* **20**, 784-794 (2010).
13.   A. M. Kelly, J. L. Goodson, Hypothalamic oxytocin and vasopressin neurons exert sex-
specific effects on pair bonding, gregariousness, and aggression in finches. *Proc. Natl.*
Acad. Sci. U.S.A **111**, 6069-6074 (2014).
14.   J. Kohl *et al.*, Functional circuit architecture underlying parental behaviour. *Nature* **556**,
326-331 (2018).
15.   Z. Liutkevičiūtė *et al.*, Oxytocin-like signaling in ants influences metabolic gene
expression and locomotor activity. *FASEB J.* **32**, 6808-6821 (2018).
16.   E. Muratspahić *et al.*, Oxytocin/vasopressin-like neuropeptide signaling in insects. *Vitam.*
Horm. **113**, 29-53 (2020).
17.   E. Stafflinger *et al.*, Cloning and identification of an oxytocin/vasopressin-like receptor
and its ligand from insects. *Proc. Natl. Acad. Sci. U.S.A* **105**, 3262-3267 (2008).
18.   M. J. Aikins *et al.*, Vasopressin-like peptide and its receptor function in an indirect
diuretic signaling pathway in the red flour beetle. *Insect Biochem. Mol. Biol.* **38**, 740-748
(2008).
19.   M. P. Scott, The ecology and behavior of burying beetles. *Ann. Rev. Entomol.* **43**, 595-
618 (1998).
20.   J. A. Oldekop, P. T. Smiseth, H. D. Piggins, A. J. Moore, Adaptive switch from
infanticide to parental care: How do beetles time their behaviour? *J. Evol. Biol.* **20**, 1998-
2004 (2007).
21.   P. T. Smiseth, C. Dawson, E. Varley, A. J. Moore, How do caring parents respond to
mate loss? Differential response by males and females. *Anim. Behav.* **69**, 551-559 (2005).
22.   J. B. Moss, A. J. Moore, Constrained flexibility of parental cooperation limits adaptive
responses to harsh conditions. *Evolution* **75**, 1835-1849 (2021).

23. K. M. Benowitz, A. J. Moore, Biparental care is predominant and beneficial to parents in
the burying beetle Nicrophorus orbicollis (Coleoptera: Silphidae). *Biol. J. Linn. Soc.*
Lond. **119**, 1082-1088 (2016).

24. Y. Fujino *et al.*, Possible functions of oxytocin/vasopressin-superfamily peptides in
annelids with special reference to reproduction and osmoregulation. *J. Exp. Zool.* **284**,
401-406 (1999).

25. K. Ukena, E. Iwakoshi-Ukena, A. Hikosaka, Unique form and osmoregulatory function
of a neurohypophysial hormone in a urochordate. *Endocrinology* **149**, 5254-5261 (2008).

26. N. A. Nagy *et al.*, Inotocin, a potential modulator of reproductive behaviours in a
biparental beetle, Lethrus apterus. *J. Insect Physiol.* **132**, 104253 (2021).

27. I. Fetter-Pruneda *et al.*, An oxytocin/vasopressin-related neuropeptide modulates social
foraging behavior in the clonal raider ant. *PLOS Biol.* **19**, e3001305 (2021).

28. J. L. Garrison *et al.*, Oxytocin/vasopressin-related peptides have an ancient role in
reproductive behavior. *Science* **338**, 540-543 (2012).

29. M. Kosfeld, M. Heinrichs, P. J. Zak, U. Fischbacher, E. Fehr, Oxytocin increases trust in
humans. *Nature* **435**, 673-676 (2005).

30. A. R. Reddon, C. M. O'Connor, S. E. Marsh-Rollo, S. Balshine, Effects of isotocin on
social responses in a cooperatively breeding fish. *Anim. Behav.* **84**, 753-760 (2012).

31. J. R. Godwin, R. Sawby, R. R. Warner, D. Crews, M. S. Grober, Hypothalamic arginine
vasotocin mRNA abundance variation across sexes and with sex change in a coral reef
fish. *Brain Behav. Evol.* **55**, 77 - 84 (2000).

32. S. M. Farris, Insect societies and the social brain. *Curr. Opin. Insect Sci.* **15**, 1-8 (2016).

33. S. O'Donnell, M. Clifford, Y. Molina, Comparative analysis of constraints and caste
differences in brain investment among social paper wasps. *Proc. Natl. Acad. Sci. U.S.A*
108, 7107-7112 (2011).

34. H. H. Dale, The action of extracts of the pituitary body. *Biochem. J.* **4**, 427 (1909).

35. A. Koto *et al.*, Oxytocin/vasopressin-like peptide inotocin regulates cuticular
hydrocarbon synthesis and water balancing in ants. *Proc. Natl. Acad. Sci. U.S.A* **116**,
5597-5606 (2019).

36. J. C. Bedick, W. W. Hoback, M. C. Albrecht, High water-loss rates and rapid dehydration
in the burying beetle, Nicrophorus marginatus. *Physiol. Entomol.* **31**, 23-29 (2006).

37. C. A. Walling, C. E. Stamper, P. T. Smiseth, A. J. Moore, The quantitative genetics of
sex differences in parenting. *Proc. Natl. Acad. Sci. U.S.A*, 18430-18435 (2008).

38. D. S. Wilson, W. Knollenberg, J. Fudge, Species packing and temperature dependent
competition among burying beetles (Silphidae, *Nicrophorus*). *Ecol. Entomol.* **9**, 205-216
(1984).

39. E. M. Roy-Zokan, C. B. Cunningham, L. E. Hebb, E. C. McKinney, A. J. Moore,
Vitellogenin and vitellogenin receptor gene expression is associated with male and
female parenting in a subsocial insect. *Proc. Biol. Sci.* **282**, 20150787 (2015).

40. C. B. Cunningham, M. K. Douthit, A. J. Moore, Octopaminergic gene expression and
flexible social behaviour in the subsocial burying beetle *Nicrophorus vespilloides*. *Insect*
Mol. Biol. **23**, 391-404 (2014).

41. K. M. Benowitz, E. C. McKinney, C. B. Cunningham, A. J. Moore, Relating quantitative
variation within a behavior to variation in transcription. *Evolution* **71**, 1999-2009 (2017).

42.   K. J. Livak, T. D. Schmittgen, Analysis of relative gene expression data using real-time
605   quantitative PCR and the 2−ΔΔCT method. *Methods* **25**, 402-408 (2001).
606   43.   M. R. Elphick, O. Mirabeau, D. Larhammar, Evolution of neuropeptide signalling
607   systems. *J. Exp.Biol.* **221** (2018).

Figure legends

Figure 1. Oxytocin/vasopressin family evolved from duplication event. An ancestral vasopressin-like

neuropeptide was duplicated in the common ancestor of jawed vertebrates, and another duplication

occurred in the cephalopods. Duplication events are denoted by a red line. Neuropeptides of the

oxytocin/vasopressin group have been found across wide taxonomic groups (those represented on

phylogeny) although the functions of these molecules are not characterized for many taxa. Functions

associated with these neuropeptides include water balance, myoactivity (i.e., muscle activity, such as in

egg laying), nesting, offspring care, affiliation and aggression. Where known, functions for these

homologs are indicated next to the taxonomic group they function in. Note that the branch lengths in the

tree are arbitrary, as well as the marked timing of the duplication events in vertebrata and cephaloda.

Figure modified from (43) in BioRender.

Figure 2. Expression of *inotocin* and *inotocin receptor* genes is associated with non-social contexts.

Orange circles represent *inotocin* expression, while closed circles represent *inotocin receptor*. (***A***)

Inotocin and *inotocin receptor* does not change across the lifetime of females, while (***B***) *inotocin receptor*

but not *inotocin* was associated with water balance in females. (***C***) Both *inotocin* and *inotocin receptor*

change across the lifetime of males; this difference was primarily driven by lower *inotocin* and *inotocin*

receptor expression in 6-day old males relative to post-reproductive ages (letters above bars indicate

significantly different expression levels between ages). Lastly, (***D***) *inotocin* and *inotocin receptor* were

both associated with water balance in males. For *B & D*, p-values are denoted over the lines, with a P-

value less than 0.05 (*), and less than 0.01 (**). Results are mean ± SEM (n = 10 for each treatment).

Figure 3. Inotocin associated with male but not female parental care. (*A*) There was no difference in

inotocin for females across parental care states (n = 20 for each state), nor was there (*B*) a relationship

between *inotocin* and the proportion of time females spent either feeding larva (open circles and dashed

line) or engaging in indirect care (closed circles and solid line; n = 34). On the other hand, (*C*) males did

show an association between parental care states and *inotocin* (n = 20 for each), and (*D*) for the subset of

males that performed care behavior without removing the female, males that fed more had higher *inotocin*

(open circles, dashed line; n = 16) and males that spent more time engaged in indirect care had lower

inotocin (closed circles, solid line; n = 21). Discrepancy in sample sizes reflects that males that did not

show any direct or indirect care were excluded from those respective analyses. For *A & C*, significant P-

values are denoted over the lines, with a p-value less than 0.05 denoted*. Results are mean ± SEM.

Figure 4. Distribution of inotocin in the brain. (*A*) The whole brain of *Nicrophorus orbicollis* stained

with DAPI (blue), with the antennal lobes (AL) and suboesophageal zone (SEZ) indicated. Inotocin

staining was not found in the optic lobes, and for this reason, the blue rectangle indicates the regions

included in *B-D*. All images are of the frontal side of the brain and are at the same magnification as *A*

unless otherwise noted. Diffuse inotocin staining (purple) was observed in the SEZ and also adjacent to

the ALs, in both females (*B*) and males (*C*). (*D*) A cluster of inotocin-staining cells were frequently

observed in the lower SEZ in both sexes. (*E*) Inotocin staining is apparent in the dorsal view of the brain,

corresponding to the general location of the mushroom bodies (MB). Direction indicated as ventral (V) to

dorsal (D).

FEMALE

MALE