A minimum effective dose for (transcranial) alternating current stimulation

Ivan Alekseichuk[a,*], Miles Wischnewski[a], Alexander Opitz[a]

[a] Dept of Biomedical Engineering, University of Minnesota, Minneapolis, MN, USA

* Corresponding author. Dept of Biomedical Engineering, University of Minnesota, 312 Church St. SE, 7-105 Nils Hasselmo Hall, Minneapolis, MN 55455, USA.

E-mail address: ialeksei@umn.edu (I. Alekseichuk).

Abstract

Transcranial alternating current stimulation (tACS) is a popular method for non-invasive neuromodulation in human research and therapy. However, what constitutes an effective dose in tACS applications is still a matter of investigation and debate. Here, we examined available literature data regarding the effects of alternating current (AC)-induced electric fields on cellular-level neural activity. The literature search identified 17 relevant experimental reports that utilized brain slices, anesthetized rodents, and awake/behaving animal models. We implemented a probabilistic meta-analysis to estimate the minimum effective dose (MED) required for inducing detectable significant neural changes with AC stimulation. The results showed that AC stimulation at 0.3 mV/mm in awake/behaving mammals leads to an 80% probability of inducing minimum neural effects. A similar level of effectiveness in brain slices and anesthetized mammals required 0.7 mV/mm. In conclusion, (transcranial) alternating current stimulation is significantly more effective in awake than in anesthetized brains. The proposed dose targets can serve as a practical guideline for tACS in humans.

Introduction

The question of adequate dosing is fundamental to non-invasive brain stimulation. Selecting stimulation intensity that results in an electric field leading to a desired physiological response is crucial. For transcranial alternating current stimulation (tACS), the concept of dose refers to the induced electric fields in the brain [1,2]. Thus, we can ask: What is the minimum electric field strength that causes detectable changes in neural activity? This question is critical for understanding tACS dosing in humans necessary to modulate mental states and behavior. Many in-vitro and animal studies have investigated how intrinsic and induced oscillating electric fields modulate neural activity. However, a clear consensus on an adequate electric field strength for physiological effects is still lacking. Here, we use a meta-analytic probabilistic approach to determine the minimum effective dose (MED) at which alternating current (AC) modulates neural spiking.

Methods

To identify effective stimulation parameters, we manually collected data from 17 papers identified in a PubMed® search (see Supplementary Table S1). We included experimental mechanistic studies in brain slices and animals that applied AC stimulation at 0.1-100 Hz with simultaneous direct neural recordings. In addition, we only considered papers that measured the AC electric field. Our analysis summarized 11 studies in rodent brain slices and deeply anesthetized rodents (study group I) and 6 studies in awake or behaving nonhuman primates and rodents (group II). From them, we extracted the lowest electric field strength values at which neural changes were detected, which constitutes the upper MED boundary (a certainly effective dose per experiment). The highest electric field values at which no changes occurred defined the lower MED boundary (a certainly ineffective dose per experiment). We assumed that the MED lies between these boundaries and follows a Beta distribution (see Supplementary Fig. S1), commonly used in uncertainty analyses. We implemented a symmetric bell-shaped $\text{Beta}_{(3,3)}$ function for studies with known boundaries. In experiments with no measurements at a lower-than-effective dose, the lower boundary became zero. There, we used a more conservative, right-leaning $\text{Beta}_{(4,2)}$ function to deemphasize close-to-zero values. The meta-analytical distribution is a Gaussian function, satisfying the Central Limit Theorem, truncated to avoid negative values. The functions were fitted by minimizing their negative loglikelihood (see Supplementary Fig. S2). Moreover, we estimated 99% confidence intervals (CI_{99}) of the meta-analytical fit using a bootstrapping method (10,000

iterations), assuming that each underlying study has a statistical power of 80% (see Supplementary Fig. S3). Finally, study groups I and II were statistically compared using the nonparametric two-sample Kolmogorov-Smirnov (KS) test.

Fig. 1. Meta-analysis of alternating current (AC)-induced electric field magnitudes that led to minimum significant neural changes. The probability functions were fitted into the literature data from studies in brain slices and anesthetized rodents (blue) and awake/behaving rodents and nonhuman primates (red). The shaded areas correspond to CI_{99}.

Results and Discussion

Meta-analytic results show the mean probabilistic MED of 0.54 mV/mm (CI_{99}: 0.44-0.63 mV/mm) and standard deviation of 0.23 mV/mm for AC stimulation in brain slices and anesthetized rodents. This mean dose corresponds to the cumulative probability of 50% to elicit minimum significant

neural changes. For awake/behaving mammals, the MED's mean = 0.23 mV/mm (CI_{99}: 0.15-0.31 mV/mm) and SD = 0.10 mV/mm. By extension, the 80% probability of eliciting a significant neural response corresponds to 0.73 mV/mm (CI_{99}: 0.64-0.83 mV/mm) in the "disconnected" biological models and 0.31 mV/mm (CI_{99}: 0.23-0.40 mV/mm) in awake/behaving mammals. The required dose for the latter is 2.35 times lower than for the former, which is significant according to the KS test (p-value < 0.001).

Our analysis indicates that sub-millivolt per millimeter oscillating electric fields can elicit measurable neural modulation. Such electric field strengths can be achieved in humans using tACS at 2-4 mA [3,4] while mitigating tolerability issues with distributed multi-electrode stimulation montages. Crucially, the effective dose threshold here is twice lower for an active brain than for a "disconnected" brain, highlighting a major role of intrinsic neural states. This conclusion agrees with individual studies. Ozen et al. demonstrated that AC stimulation at ~0.67-1 mV/mm (recalculated from the reported 0.8-1.2 V) induced a response in rats under anesthesia, while ~0.33 mV/mm was already sufficient to modulate their brain during normal sleep [5]. Conversely, even ~1.67 mV/mm had no effects in rats during exploratory behavior. Furthermore, Fröhlich and McCormick showed that stimulating brain slices with their "natural" AC waveform (recorded and played back) makes the neurons twice more susceptible to entrainment in comparison to regular AC stimulation [2]. Huang et al. stimulated posterior alpha oscillations in ferrets and showed better neural susceptibility to tACS closer to intrinsic frequencies [6]. On the other hand, Johnson and Alekseichuk et al. found no straightforward relationships between the spiking rates of individual neurons and their response to a given stimulation frequency [7]. Noteworthy, the stimulation frequency itself can play a particular role. Several studies reported that stimulation at lower frequencies (<20 Hz) is more effective than at higher frequencies (50-100 Hz) [1,8,9]. Finally, different brain regions can have different innate thresholds to AC stimulation due to their cytoarchitecture. Francis et al. found a twice lower MED in CA1 vs. CA3 rat slices, all other factors equal [10].

Our analysis above addresses the minimum effective dose, but physiological responses at above threshold intensities are still under active investigation. Recently, Krause et al. reported that neurons intrinsically entrained to a particular oscillation can be desynchronized by tACS at the same frequency for low intensities before becoming re-entrained at higher intensities [11]. A full dose-response curve and its dependency on the AC frequency, cytoarchitecture, and baseline neural activity is beyond the scope of the present analysis, as the existing studies explored very

different subsets of stimulation parameters while extensive parametric experimental studies are yet to be performed.

In summary, we show that AC electric fields as low as 0.3 mV/mm can be sufficient to modulate neurons in awake and behaving mammals and can thus serve as practical guidance for tACS dosing. Notably, these magnitudes are twice lower than required for anesthetized rodents and brain slices, and future research is needed to determine if task engagement in humans further increases the brain's susceptibility to tACS.

Declaration of competing interest

All authors declare no competing interests.

Acknowledgements

This work was supported by the Brain & Behavior Research Foundation (Young Investigator Grant to I.A.), the University of Minnesota's MnDrive Initiative (M.W. and A.O.), and the National Institutes of Health (grant RF1MH124909 to A.O.).

Appendix A. Supplementary data

Supplementary data to this article can be found online.

References

[1] Anastassiou CA, Perin R, Markram H, Koch C. Ephaptic coupling of cortical neurons. Nature Neuroscience 2011;14:217–23. https://doi.org/10.1038/nn.2727.

[2] Fröhlich F, McCormick DA. Endogenous Electric Fields May Guide Neocortical Network Activity. Neuron 2010;67:129–43. https://doi.org/10.1016/j.neuron.2010.06.005.

[3] Opitz A, Falchier A, Yan C, Yeagle EM, Linn GS, Megevand P, et al. Spatiotemporal structure of intracranial electric fields induced by transcranial electric stimulation in humans

and nonhuman primates. Scientific Reports 2016;6:31236. https://doi.org/10.1038/srep31236.

[4] Huang Y, Liu AA, Lafon B, Friedman D, Dayan M, Wang X, et al. Measurements and models of electric fields in the in vivo human brain during transcranial electric stimulation. ELife 2017;6:e18834. https://doi.org/10.7554/eLife.18834.

[5] Ozen S, Sirota A, Belluscio MA, Anastassiou CA, Stark E, Koch C, et al. Transcranial Electric Stimulation Entrains Cortical Neuronal Populations in Rats. Journal of Neuroscience 2010;30:11476–85. https://doi.org/10.1523/JNEUROSCI.5252-09.2010.

[6] Huang WA, Stitt IM, Negahbani E, Passey DJ, Ahn S, Davey M, et al. Transcranial alternating current stimulation entrains alpha oscillations by preferential phase synchronization of fast-spiking cortical neurons to stimulation waveform. Nature Communications 2021;12:3151. https://doi.org/10.1038/s41467-021-23021-2.

[7] Johnson L, Alekseichuk I, Krieg J, Doyle A, Yu Y, Vitek J, et al. Dose-dependent effects of transcranial alternating current stimulation on spike timing in awake nonhuman primates. Science Advances 2020;6:eaaz2747. https://doi.org/10.1126/sciadv.aaz2747.

[8] Krause MR, Vieira PG, Csorba BA, Pilly PK, Pack CC. Transcranial alternating current stimulation entrains single-neuron activity in the primate brain. Proceedings of the National Academy of Sciences of the United States of America 2019;116:5747–55. https://doi.org/10.1073/pnas.1815958116.

[9] Deans JK, Powell AD, Jefferys JGR. Sensitivity of coherent oscillations in rat hippocampus to AC electric fields. The Journal of Physiology 2007;583:555–65. https://doi.org/10.1113/jphysiol.2007.137711.

[10] Francis JT, Gluckman BJ, Schiff SJ. Sensitivity of Neurons to Weak Electric Fields. The Journal of Neuroscience 2003;23:7255–61. https://doi.org/10.1523/JNEUROSCI.23-19-07255.2003.

[11] Krause MR, Vieira PG, Thivierge J-P, Pack CC. Brain stimulation competes with ongoing oscillations for control of spike timing in the primate brain. PLOS Biology 2022;20:e3001650. https://doi.org/10.1371/journal.pbio.3001650.