Posterodorsal medial amygdala urocortin-3, GABA and glutamate mediate suppression

of LH pulsatility in female mice

Deyana Ivanova[1], Xiao-Feng Li[1], Caitlin McIntyre[1], HongBei Xu[1] and Kevin T O'Byrne[1]

[1]Department of Women and Children's Health, School of Life Course and Population Sciences,

Faculty of Life Science and Medicine, King's College London, UK

Short title: MePD Ucn3 effect on LH and corticosterone release

Key words: Ucn3, GABA, glutamate, MePD, PVN, LH pulsatility, CORT secretion

Corresponding author's contact details: Deyana Ivanova (deyana.ivanova@kcl.ac.uk) and

Kevin T O'Byrne (kevin.obyrne@kcl.ac.uk), Department of Women and Children's Health,

School of Life Course and Population Sciences, Faculty of Life Science and Medicine, King's

College London, 2.92W Hodgkin Building, Guy's Campus, London SE1 1UL, UK.

Reprint requests: Deyana Ivanova, Department of Women and Children's Health, School of

Life Course Sciences, Faculty of Life Science and Medicine, King's College London, 2.92W

Hodgkin Building, Guy's Campus, London SE1 1UL, UK.

Grants supporting paper: UKRI: BBSRC (BB/S000550/1). DI is a PhD student funded by

MRC-DTP studentship.

The authors have nothing to disclose.

Abstract

The posterodorsal subnucleus of the medial amygdala (MePD) is an upstream modulator of the hypothalamic-pituitary-gonadal (HPG) and hypothalamic-pituitary-adrenal (HPA) axes. Inhibition of MePD urocortin-3 (Ucn3) neurons prevents psychological stress-induced suppression of LH pulsatility while blocking the stress-induced elevations in corticosterone (CORT) secretion in female mice. We explore the neurotransmission and neural circuitry suppressing the GnRH pulse generator by MePD Ucn3 neurons and we further investigate whether MePD Ucn3 efferent projections to the PVN control CORT secretion and LH pulsatility. Ucn3-cre-tdTomato female ovariectomised (OVX) mice were unilaterally injected with AAV-ChR2 and implanted with optofluid cannulae targeting the MePD. We optically activated Ucn3 neurons in the MePD with blue light at 10Hz and monitored the effect on LH pulses. Next, we combined optogenetic stimulation of MePD Ucn3 neurons with pharmacological antagonism of GABA$_A$ or GABA$_B$ receptors with bicuculline or CGP, respectively, as well as a combination of NMDA and AMPA receptor antagonists, AP5 and CNQX respectively, and observed the effect on pulsatile LH secretion. A separate group of Ucn3-cre-tdTomato OVX mice with oestradiol replacement were unilaterally injected with AAV-ChR2 in the MePD and implanted with fibreoptic cannulae targeting the PVN. We optically stimulated MePD Ucn3 neuron terminals with blue light at 20Hz and monitored the effect on CORT secretion and LH pulses. We reveal for the first time that activation of Ucn3 neurons in the MePD inhibits GnRH pulse generator frequency via GABA and glutamate signalling within the MePD, while MePD Ucn3 projections to the PVN modulate the HPG and HPA axes.

Introduction

Stress exerts a profound suppressive effect on reproduction in mammals, including rodents and humans. The amygdala, part of the limbic system, is a key emotional centre integrating incoming cues from the external environment, such as anxiogenic signals, with the reproductive and stress axes (1). The posterodorsal sub-nucleus of the medial amygdala (MePD) is an upstream modulator of pulsatile luteinizing hormone (LH) secretion (2–4) and exerts an inhibitory break on pubertal timing (5,6). The MePD sends direct projections, of an unknown phenotype, to the kisspeptin (kiss1) neuronal population in the hypothalamic arcuate nucleus (ARC) (7,8) known as KNDy because they co-express neurokinin B (NKB) and dynorphin A (Dyn) (9–11). The synchronous activity of the ARC KNDy network drives pulsatile gonadotropin-releasing hormone (GnRH) and LH release (11–14). The MePD also contains a kiss1 neuronal population and selective optogenetic activation of these neurons increases GnRH pulse generator frequency in mice (4) and unpublished observations show this is an effect involving both MePD GABA and glutamate signalling.

We have recently shown that urocortin-3 (Ucn3), a member of the corticotropin-releasing factor (CRF) stress neuropeptide family, and its receptor CRF type 2 (CRFR2) signalling in the MePD are involved in mediating the suppressive effect of psychosocial stress on LH pulsatility (3). However, the underlying neural mechanisms involved in mediating the inhibitory effect of MePD Ucn3 neurons on LH pulsatility remain to be established. GABA and glutamate signalling in the amygdala are implicated in the modulation of anxiety behaviour (15,16). The amygdala also modulates the stress response through the hypothalamic pituitary adrenal (HPA) axis. The medial amygdala activates the HPA axis in response to predator odor stress (17) and the MePD in particular has been shown to send stress-activated efferents to the PVN (18). Furthermore, MePD Ucn3 and CRFR2 positive neurons are implicated in regulating the stress response, with restraint stress increasing MePD Ucn3 (19) and social defeat elevating

MePD CRFR2 expression in rodents (20). Moreover, MePD Ucn3 and CRFR2 positive

neurons project to the PVN (21), and we have recently shown that chemogenetic inhibition of

MePD Ucn3 neurons prevents psychogenic stress-induced corticosterone (CORT) release in

female mice (3).

In this study, we aimed to determine whether activation of MePD Ucn3 neurons inhibits

pulsatile LH secretion via GABA and or glutamate signalling within this amygdala sub-

nucleus. To achieve this, we will combine selective optogenetic activation of MePD Ucn3

neurons with pharmacological antagonism of $GABA_A$ or $GABA_B$ receptors, or a combination

of N-methyl-D-aspartate (NMDA) and α-amino-3-hydroxy-5-methyl-4-isoxazole propionic

acid (AMPA) receptors, respectively, using chronically implanted intra-MePD optofluid

cannulae while collecting serial blood samples for LH pulse measurement in Ucn3-Cre-

tdTomato female mice. Additionally, we will investigate whether optogenetic stimulation of

MePD Ucn3 projection terminals in the PVN induce CORT release and suppress LH pulsatility.

Methods

Mice

Cryopreserved sperm of Ucn3-cre mice (strain Tg(Ucn3-cre)KF43Gsat/Mmucd; congenic on

C57BL/6 background) was acquired from MMRRC GENSAT and heterozygous transgenic

breeding pairs of Ucn3-Cre mice were recovered via insemination of female C57Bl6/J mice at

King's College London, as described previously (3). Ucn3-Cre mice were genotyped using

PCR for the detection of heterozygosity and Ucn3-cre mice were bred with td-Tomato mice

(strain   B6.Cg-Gt(ROSA)26Sortm9(CAG-tdTomato)Hze/J;   congenic   on   C57BL/6

background) acquired from The Jackson Laboratory, Bar Harbor, ME, USA to produce Ucn3-

cre-tdTomato reporter mice, as described previously (3,21). Female Ucn3-cre-tdTomato mice

weighing 19-23 g and aged 6-8 weeks were singly housed in individually ventilated cages

sealed with a HEPA-filter at 25 ± 1 °C in a 12:12 h light/dark cycle, lights on at 07:00 h

equipped with nesting material, wood-chip bedding and food and water ad libitum. All

procedures were carried out following the United Kingdom Home Office Regulations and

approved by the Animal Welfare and Ethical Review Body Committee at King's College

London.

Stereotaxic injection of adeno-associated-virus carrying the channelrhodopsin construct

and fibre optic or optofluid cannula implantation

All surgical procedures were carried out under aseptic conditions with general anaesthesia

using ketamine (Vetalar, 100 mg/kg, i.p.; Pfizer, Sandwich, UK) and xylazine (Rompun, 10

mg/kg, i.p.; Bayer, Leverkusen, Germany). Ucn3-Cre-tdTomato mice were secured in a David

Kopf stereotaxic frame (Kopf Instruments, Model 900) and either solely bilaterally

ovariectomised (OVX) or implanted with Silastic capsule containing 36 μg 17β-estradiol/mL

(E_2) in sesame oil capsule (22). These capsules produce circulating concentrations of E_2 within

the range observed during the diestrous phase of the estrous cycle (23,24). Estradiol is known

to enhance the suppressive effects of stress where the action of the gonadal steroids sensitizes

the HPA axis and augments stress-induced inhibition of the GnRH pulse generator in many

species, from rodents to primates (42–45). We had two experimental groups; OVX only and

OVX + E_2 capsule. The mouse brain atlas of Paxinos and Franklin (25) was used to obtain

target coordinates for the MePD (2.30 mm lateral, -1.55 mm from bregma, at a depth of -4.94

mm below the skull surface). To reveal the skull an incision was made of the scalp and one

small hole was drilled above the location of the right MePD. AAV9-EF1a-double floxed-

hChR2(H134R)-EYFP-WPRE-HGHpA (300 nL; $3x10^{11}$ GC/ml; Serotype:9; Addgene,

Massachusetts, USA) was unilaterally injected into the right MePD using a 2-μL Hamilton

micro syringe (Esslab, Essex, UK) over 10 min, using the robot stereotaxic system (Neurostar, Tubingen, Germany), performed for the targeted expression of ChR2-EYFP in MePD Ucn3 neurons. After injection, the needle was left in position for a further 5 min and lifted slowly over 2 min. Cre-positive mice received the AAV-ChR2 injection (test mice) or a control virus AAV-YFP (Addgene) (control mice). The control virus does not contain the ChR2 construct. The mice were then implanted with a dual optofluid cannula (Doric Lenses, Quebec, Canada) at the same AP and ML coordinates as the viral injection site, however a different DV was used such that the internal cannula targets the MePD and the fibre optic cannula is situated 0.2 mm above the MePD site. Once in position the optofluid cannula was secured on the skull using dental cement (Super-Bond Universal Kit, Prestige Dental, UK) and the incision of the skin was closed with suture. Mice were left to recover for one week and after recovery period, mice were handled daily to acclimatise to experimental procedures for a further 2 weeks. Thirteen mice received the AAV-ChR2 injection and were implanted with an optofluid cannula in the MePD. Five mice received the control AAV-YFP and were implanted with an optofluid cannula in the MePD. Eleven mice received the AAV-ChR2 injection in the MePD and were implanted with an optic fibre in the PVN. Five mice received the control AAV-YFP and were implanted with an optic fibre in the PVN.

In vivo optogenetic stimulation of MePD Urocortin-3 neurons

In order to test the effect of optogenetic stimulation of MePD Ucn3 neurons on LH pulsatility the ferrule of the implanted optofluid cannula, on the right side of the MePD, was attached via a ceramic mating sleeve to a multimode fibre optic rotatory joint patch cable (Thorlabs LTD, Ely, UK) at a length allowing for free movement of the mouse in their cage and blue light (473 nm wavelength; 10 mW) was delivered using a Grass SD9B stimulator-controlled laser (Laserglow Technologies, Toronto, Canada). The mice were left to acclimatise for 1 h.

Following the acclimatisation period blood samples (5 µL) were collected every 5 min for 2 h

where after 1 h of control blood sampling, Ucn3-cre-tdTomato mice received optic stimulation

at 10 Hz with a 10 ms pulse interval and pattern stimulation of 5 seconds on and 5 seconds off

for the remaining 1 h.

Intra-MePD administration of bicuculline (BIC), CGP-35348 or AP5 + CNQX during

optogenetic stimulation of MePD Urocortin-3 neurons

Neuropharmacological manipulation of GABA and glutamate receptor signalling in the MePD

with or without optogenetic stimulation of MePD Ucn3 neurons was done to test whether

MePD Ucn3 induced suppression of LH pulsatility involves GABA or glutamate receptor

signalling. OVX mice implanted with a optofluid cannula in the MePD were subjected to the

tail-tip blood collection procedure, as previously described (26). Infusion of drugs with or

without optogenetic stimulation and blood sampling were performed between 09:00-13:00 h,

where 5 µl blood was collected every 5 min for 2 h. Internal cannula (Doric Lenses) attached

to extension tubing (0.58 mm ID, 0.96 mm OD), preloaded with bicuculline (BIC) (GABA$_A$

receptor antagonist; Sigma-Aldrich), CGP-35348 (GABA$_B$ receptor antagonist; Sigma-

Aldrich), AP5 + CNQX cocktail (NMDA and AMPA receptor antagonists; Tocris) or aCSF as

vehicle control, was inserted into the guide cannula of the optofluid implant and mice were

connected to the laser as described above. The tubing for drug infusion extended beyond the

cage and the distal ends were attached to 10-µl Hamilton syringes (Waters Ltd, Elstress, UK)

fitted into a PHD 2000 Programmable syringe pump (Harvard Apparatus, Massachusetts,

USA), allowing for a continuous infusion of the drug at a constant rate and mice were kept in

the cage throughout the experiment, freely-moving with food and water *ad libitum*. After 55

min of control blood sampling, the mice were given an initial bolus injection, 0.30 µl at a rate

of 0.06 µl/min over 5 min, of BIC, CGP-35348, AP5 + CNQX or aCSF. The laser was turned

on at 60 min and the bolus injection was followed by a continuous infusion, 0.8 μl at a rate of

0.01 μl/min over 1 h, of BIC, CGP-35348, AP5 + CNQX or aCSF. BIC, CGP-35348, AP5 and

CNQX were dissolved in aCSF to reach bolus concentrations of 20 pmol, 4.5 nmol, 1.2 nmol

and 0.5 nmol respectively. The concentrations for the continuous infusion of BIC, CGP-35348,

AP5 and CNQX were 68 pmol, 15 nmol, 2nmol and 1 nmol respectively. In the absence of

optic stimulation, the same regimen was applied. The mice receive all treatments in a random

order, with at least 2 days between experiments.

167   MePD Urocortin-3 terminal optogenetic stimulation in the PVN

To test the effect of optogenetic stimulation of MePD Ucn3 neuronal terminals in the PVN on

LH pulsatility and CORT release, OVX + E_2 Ucn3-cre-tdTomato mice with AAV-ChR2

injection in the right MePD implanted with a fibre optic cannula in the right PVN were

subjected to the tail-tip blood collection procedure, as previously described (26). For LH

measurement, blood samples were collected every 5 min for 2 h, as described above. Mice

were connected to the laser as described above. Optogenetic stimulation was initiated after the

1 h control blood sampling period and Ucn3-cre-tdTomato mice received optic stimulation at

20 Hz, 15 mW with a 10 ms pulse interval and pattern stimulation of 5 seconds on and 5

seconds off for the remaining 1 h. For CORT measurements, 15 μl blood samples were

collected on a separate occasion and blood samples were stored in tubes containing 5 μl

heparinised saline (5 IU ml^{-1}). Optogenetic stimulation was initiated, as described above, 30

min into the experiment (time point, 0 min) and lasted for 1 h (time point 60 min). Remaining

blood samples were collected at 0, 30 and 60 min, with a final sample taken 1 h after the

termination of optogenetic stimulation (time point, 120 min). At the end of the experiment,

blood samples were centrifuged at 13,000 RPM for 20 min at 20°C and plasma stored at -20°C.

All experiments were performed between 9:00-12:00 h.

Validation of AAV injection and cannula implant site

Upon completion of experiments, mice were anaesthetised with a lethal dose of ketamine and transcardial perfusion was performed with heparinised saline for 5 min followed by ice-cold 4% paraformaldehyde (PFA) in phosphate buffer (pH 7.4) for 15 min using a pump (Minipuls, Gilson, Villiers Le Bel, France). Mice brains were collected immediately and post-fixed in 15% sucrose in 4% PFA at 4 °C and left to sink. Once sunk, they were transferred to 30% sucrose in phosphate-buffered saline (PBS) and left to sink. Brains were snap-frozen in isopropanol on dry ice and stored in -80°C until further processing. Every third coronal brain section (30-μm/section) through-out the MePD region (-1.34 mm to -2.70 mm from bregma) was collected using a cryostat (Bright Instrument Co., Luton, UK). Sections were mounted on microscope slides, air dried and covered with ProLong Antifade mounting medium (Molecular Probes, Inc. OR, USA). Cannula placement and AAV injection site was verified with Axioskop 2 Plus microscope equipped with Axiovision, version 4.7 (Zeiss) by determining whether cannula reach the MePD. For AAV-ChR2-EYFP injected Ucn3-cre-tdTomato mice we determined whether MePD Ucn3 neurons were infected by merging td-Tomato fluorescence of Ucn3 neurons with YFP fluorescence in the MePD, as described previously (3). Axioskop 2 Plus microscope (Carl Zeiss) equipped with axiovision, version 4.7 (Carl Zeiss) was used to take images. Only mice with correct AAV injection and cannula placement in the MePD were analysed.

LH and CORT measurement

Mice were handled daily to acclimatise to the tail-tip blood sampling procedure for LH measurement (3). Blood samples collected for LH measurement were processed using LH ELISA, as previously reported (3). The capture antibody (monoclonal antibody, anti-bovine LHβ subunit, AB_2665514) was purchased from Department of Animal Science at the

University of California, Davis. The mouse LH standard (AFP-5306A) and primary antibody

(polyclonal antibody, rabbit LH antiserum, AB_2665533) were obtained from Harbour-UCLA

(California, USA). The secondary antibody (Horseradish-Peroxidase (HRP)-linked donkey

anti-rabbit IgG polyclonal antibody, AB_772206) was purchased from VWR International

(Leicestershire, UK). Intra-assay and inter-assay variations were 4.6% and 10.2%, respectively

and the assay sensitivity was 0.0015 ng/mL. DynPeak algorithm was used for the detection of

LH pulses (3). The effect of optogenetic stimulation of MePD Ucn3 neurons and

neuropharmacology studies was established by comparing the LH inter-pulse interval (IPI)

from the 1 h control period (pre-stimulation) or drug administration control period to the 1 h

experimental period, as previously described (3).

Blood samples were collected for CORT measurement were processed using a commercially

available enzyme immunoassay (EIA) kit (sheep polyclonal antibody specific for

corticosterone, AB_2877626; DetectX® Corticosterone Enzyme Immunoassay Kit, K014;

Arbor Assays, Michigan, USA), as described previously (3).

Statistics

Mice were compared between groups using a one-way ANOVA with a Tukey post-hoc and

data was presented as mean ± SEM. Statistics were performed using Igor Pro 7, Wavemetrics,

Lake Oswego, OR, USA. Data was represented as mean ± SEM and +p<0.05, ++p<0.001 and

+++p<0.0001 were considered to be significant.

Results

Effects of optogenetic stimulation of MePD Urocortin-3 neurons on LH pulse frequency

Optogenetic stimulation of MePD Ucn3 neurons in control mice had no effect on LH pulse

interval (Fig. 1A; AAV-YFP: n = 3) and administration of aCSF alone (AAV-YFP with no

stimulation: n = 2) had no effect on LH pulse interval. Optogenetic stimulation of MePD Ucn3

neurons with blue light at 10 Hz, 10 mW suppressed pulsatile LH secretion (Fig. 1B; AAV-

ChR2: n = 9). These data are summarized in the figure 1C.

Effects of intra-MePD administration of bicuculline (BIC), a GABA$_A$R antagonist, and

CGP, a GABA$_B$R antagonist, during optical stimulation of MePD Urocortin-3 neurons

on LH pulsatility

Delivery of BIC alone had no significant effect on LH pulsatility (Fig. 1D; n = 8). Intra-MePD

infusion of BIC during optogenetic stimulation of MePD Ucn3 neurons with blue light at 10

Hz, 10 mW completely blocked the suppressive effect of MePD Ucn3 neuronal activation on

LH pulsatility (Fig. 1E; n = 9). The results of this experiment are summarised in the figure 1F.

Effects of intra-MePD administration of CGP, a GABA$_B$R antagonist, during optical

stimulation of MePD Urocortin-3 neurons on LH pulsatility

Delivery of CGP alone had no significant effect on LH pulsatility (Fig. 1G; n = 7). Intra-MePD

infusion of CGP during optogenetic stimulation of MePD Ucn3 neurons with blue light at 10

Hz, 10 mW completely blocked the suppressive effect of MePD Ucn3 neuronal activation on

LH pulsatility (Fig. 1H; n = 7). The results of this experiment are summarised in the figure 1I.

Effects of intra-MePD glutamate antagonism during optogenetic stimulation of MePD

Urocortin-3 neurons on LH pulsatility

Delivery of AP5 + CNQX alone had no significant effect on LH pulsatility (Fig. 1J; n = 6).

Intra-MePD infusion of AP5 (selective NMDA antagonist) + CNQX (selective AMPA

antagonist) during optogenetic stimulation of MePD Ucn3 neurons with blue light at 10 Hz, 10

mW completely blocked the suppressive effect of MePD Ucn3 neuronal activation on LH

pulsatility (Fig. 1K; n = 6). The results of this experiment are summarised in the figure 1L.

254 **Effect of MePD Urocortin-3 terminal stimulation in the PVN on dynamic CORT secretion**

255 Optogenetic stimulation of MePD Ucn3 neuron terminals in the PVN with blue light at 20 Hz,

256 15 mW elevated CORT secretion compared to controls (Fig. 2; AAV-ChR2: n = 6).

257 Optogenetic stimulation of MePD Ucn3 neurons and terminal projections in the PVN in control

258 mice had no effect on dynamic CORT secretion (AAV-YFP; n = 4).

259 **Effect of MePD Urocortin-3 terminal stimulation in the PVN on LH pulsatility**

260 Optogenetic stimulation of MePD Ucn3 projections in the PVN in control mice had no effect

261 on LH pulse interval (Fig. 3A; AAV-YFP, n = 3; wild-type with AAV-ChR2, n = 2) and no

262 stimulation of the MePD Ucn3 projections in the PVN (no stimulation, n = 2) had no effect on

263 LH pulse interval. Optogenetic stimulation of MePD Ucn3 neuron terminals in the PVN with

264 blue light at 20 Hz, 15 mW reduced LH pulsatility compared to controls (Fig. 3B; AAV-ChR2:

265 n = 8). The results of this experiment are summarised in the figure 3C.

266 **Selective expression of ChR2 in MePD Ucn3 neurons and MePD Ucn3 neuron**

267 **projections in the PVN, and validation of cannula position**

268 Evaluation of AAV9-EF1a-double floxed-hChR2(H134R)-EYFP-WPRE-HGHpA expression

269 in tdTomato labelled neurons from AAV-injected Ucn3-cre-tdTomato mice revealed that 87.00

270 ± 6.00% of MePD Ucn3 neurons expressed AAV-ChR2 and the number of tdTomato labelled

271 Ucn3 neurons per side per slice in the MePD was counted at 49.00 ± 5.37 (mean ± SEM) with

272 the number of AAV-hM4D-mCitrine infected neurons being 43.01 ± 3.65 (mean ± SEM)

273 (n=8). A representative example is shown in Fig 4, A-C. The optofluid cannula was also

274 verified to be correctly positioned in the MePD in all of 8 mice. Evaluation of AAV9-EF1a-

275 double floxed-hChR2(H134R)-EYFP-WPRE-HGHpA expression in Ucn3 projections from

276 the MePD to the PVN showed YFP labelled fibres in the PVN Fig 5, A-Aii. Local Ucn3

277 neurons in the PVN are labelled with tdTomato. The optic fibre cannula was also verified to

278 be correctly positioned in the PVN in all of 8 mice (Fig. 5A).

Discussion

We show optogenetic stimulation of MePD Ucn3 neurons inhibits pulsatile LH secretion, an

effect that was completely blocked by intra-MePD GABA and glutamate antagonism.

Moreover, optogenetic stimulation of MePD Ucn3 projections in the hypothalamic PVN also

suppresses LH pulsatility while inducing release of CORT. These results demonstrate that

inhibition of the GnRH pulse generator by Ucn3 neurons within the MePD relies on GABA

and glutamate signalling in the MePD, *per se*. Moreover, MePD Ucn3 efferent projections in

the PVN modulate the activity of the HPA and HPG axes.

The MePD is a major upstream regulator of gonadotropic hormone secretion with a significant

population of GABAergic neurons projecting to reproductive neural centres in the

hypothalamus (27). Although classic lesioning studies suggest an inhibitory output of the

MePD over reproductive function (28,29), selective optical stimulation of MePD kiss1 neurons

increases LH pulse frequency (4), which our unpublished observations show relies upon both

GABA and glutamate signalling in the MePD. Due to this interesting dichotomy, we have

proposed a disinhibitory system, which is consistent with the pallidal origin of this limbic

structure (30), whereby MePD kiss1 activation excites inhibitory GABAergic interneurons,

which in turn reduces the inhibitory tone of the GABAergic projections from the MePD to the

ARC KNDy network allowing for an increase in GnRH pulse generator frequency (4).

Contrastingly, kisspeptin antagonism, reducing MePD kiss1 signalling decreases GnRH pulse

generator frequency (31).

We have recently shown that stress-induced activation of Ucn3 and CRFR2 signalling within

the MePD inhibits GnRH pulse generator activity (3) and activation of MePD Ucn3 neurons

delays pubertal timing (32). It has also been shown that the majority of MePD CRFR2

expressing neurons are GABAergic (21). In the MePD, Ucn3 fibres overlap with sites

expressing its cognate receptor CRFR2 (33,34) and some local Ucn3 neurons within the MePD

are shown to have an interneuron-like appearance (35), thus Ucn3 neurons may form

connections with and signal via GABAergic CRFR2 positive neurons in this region. In the

present study, we show that optogenetic activation of MePD Ucn3 neurons inhibits pulsatile

LH secretion. Therefore, we propose that activated MePD Ucn3 neurons may signal through

inhibitory GABAergic CRFR2 interneurons, upstream to the MePD kiss1 neurons,

downregulating MePD kiss1 signalling resulting in decreased GnRH pulse generator

frequency.

In our proposed model, we hypothesised that GABA signalling is an important factor mediating

the suppressive effect of MePD Ucn3 activation on GnRH pulse generator activity. Therefore,

we tested the effect of intra-MePD administration of $GABA_A$ or $GABA_B$ receptor antagonists

during optical stimulation of MePD Ucn3 neurons on LH pulsatility, and indeed, we found that

both $GABA_A$ and $GABA_B$ receptor antagonism blocks the suppression of LH pulses by MePD

Ucn3 neuronal activation. From our previous studies, we know MePD Ucn3 neurons mediate

psychosocial stress-induced inhibition of LH pulsatility (3). Based on our proposed neuronal

circuit, $GABA_A$ and $GABA_B$ receptor antagonism may be working upstream and downstream

of MePD kiss1 activity to cancel both the inhibitory GABAergic CRFR2 interneurons upstream

to and the inhibitory GABAergic interneurons downstream to MePD kiss1 neurons. Therefore,

the inhibitory output of the MePD GABAergic projection neurons to the ARC KNDy network

is cancelled, thus, negating the suppressive effect of MePD Ucn3 activation on the GnRH pulse

generator. Essentially, terminating the ability of the GABA interneuron, upstream and

downstream of MePD kiss1 activity, to take part in the disinhibition by antagonising GABA

receptors on the MePD kiss1 neurons as well as post-synaptic GABA projection neurons

removing the inhibitory control over the GnRH pulse generator from the MePD.

We have identified GABA signalling as a key part of the MePD Ucn3 neural circuitry

associated with suppression of the GnRH pulse generator, however glutamate signalling in the

amygdala is also known to be involved in modulating reproduction and anxiety-behaviour (36).

Administration of NMDA into the MePD disrupts ovarian cycles in rats, while NMDA receptor

antagonism in the MePD delays pubertal timing (29). Urocortins can function endogenously

as modulators of excitatory glutamatergic transmission in the amygdala, with *in vitro* studies

showing modulation of glutamatergic excitatory post-synaptic potentials via pre- and

postsynaptic CRFR2-mediated mechanisms (37). Moreover, Ucn3 neurons may be

glutamatergic, as shown in the hypothalamic perifornical area, where they are activated during

infant-directed aggression (38) and increase anxiety-like behaviour (39), but this awaits future

study for the MePD Ucn3 population. Therefore, we hypothesised that glutamate

neurotransmission may form a critical part of the MePD Ucn3 neural circuitry involved in

stress-induced inhibition of GnRH pulse generator activity. In the present study, we show that

intra-MePD AMPA and NMDA receptor antagonism blocks the inhibitory effect of MePD

Ucn3 neuronal activation on LH pulsatility. These observations suggest the involvement of

similar neural circuitry as in the neighbouring posteroventral medial amygdala which exhibits

functional glutamatergic signalling onto inhibitory GABA neurons (30). Moreover, inhibition

of both GABA and glutamate signalling is equally effective at blocking suppression of LH

pulses suggesting they are key components in the circuit. Building upon our proposed neuronal

circuit, MePD Ucn3 expressing neurons may be glutamatergic modulating excitatory post-

synaptic signals to activate inhibitory GABAergic CRFR2 positive interneurons, which in turn

downregulate MePD kiss1 activity, resulting in suppression of GnRH pulse generator activity.

Thus, NMDAR and AMPAR antagonism possibly cancels MePD Ucn3 glutamatergic

activation of inhibitory GABAergic CRFR2 inter-neurons preventing the decrease in GnRH pulse generator activity.

Stress elevates Ucn3 mRNA (19) and c-fos expression in the MePD (18) while chemogenetic inhibition of MePD Ucn3 neurons prevents stress-induced suppression of LH pulsatility (3). The MePD sends stress-activated efferents to the PVN (18) and more recently MePD Ucn3 neurons have been shown to project directly to the PVN (21). In the current study, we demonstrate for the first time that optical stimulation of MePD Ucn3 projections in the PVN inhibits pulsatile LH secretion. Moreover, chemogenetic activation of PVN CRF neuron perikarya rapidly suppresses LH pulse frequency, however optogenetic activation of their nerve terminals in the ARC does not affect the firing of KNDy neurons (40) and directly injecting CRF in the ARC also has no effect on LH secretion (41), suggesting the involvement of an indirect mechanism modulating GnRH pulse generator activity. Since MePD Ucn3 neurons receive inputs from the PVN (21) and the MePD sends direct efferent projections to KNDy neurons of an unknown neurochemical phenotype (8), this amygdala structure may function as a relay centre between the PVN and ARC, to regulate stress signals to the GnRH pulse generator as well as contributing to cross talk between the HPA and HPG axes.

Central administration of Ucn3 augments the HPA axis response to restraint stress in rats (19) and increases hypothalamic CRF concentrations and circulating levels of CORT in mice (42). Overexpression of Ucn3 in the hypothalamic perifornical area increases anxiety-like behaviour (39) and activation of the CRFR2 in the lateral septum is involved in stress-induced persistent anxiety in mice (43). In the medial amygdala, Ucn3 neuronal populations are shown to consist of both projection neurons extending to the PVN, the bed nucleus of the stria terminalis and the suprachiasmatic nucleus (21,35) and local interneurons where Ucn3 neuronal fibres overlap with CRFR2-expression (33,34). We have previously shown chemogenetic inhibition targeting both local and efferent MePD Ucn3 neurons prevents psychosocial stress-induced

release of CORT (3). In the present study, we show that selective optogenetic activation of

MePD Ucn3 projections in the PVN elevates CORT secretion, thus MePD Ucn3 projections to

the PVN are critical for the regulation of CORT secretion. Further work is required to identify

whether only the MePD Ucn3 projecting efferents to the PVN are crucial for regulating CORT

secretion. The presence of estradiol is known to up-regulate basal CRF mRNA levels in the

PVN in rodents and induce high levels of circulating corticosterone providing a sensitizing

effect to the activity of the HPA axis (44,45). Therefore, we implanted the OVX mice with

estradiol capsules producing circulating concentrations of estradiol within the range observed

during the diestrous phase of the estrous cycle for HPA axis sensitization in mice similarly to

what has been reported in previous studies (23,24,46). PVN-CRF neurons express mRNA

transcripts for CRFR2 (47) and CRFR2 primarily activate the Gαs adenylyl cyclase/cAMP

signalling pathway (48) indicating that signalling via the CRFR2 located on PVN-CRF neurons

may result in activation of this neuronal population and hence the HPA axis. Therefore, Ucn3

released following optical stimulation of MePD Ucn3 projections in the PVN may directly

activate the PVN-CRF neurons, signalling via CRFR2, to stimulate CORT release.

Selective CRFR2 antagonism has been shown to reverse the stress-induced inhibition of LH

pulses (49) where central administration of astressin2-B, a specific CRFR2 antagonist, blocked

the restraint-stress induced suppression of pulsatile LH secretion in rats (50). We have shown

specific antagonism of CRFR2 in the MePD blocked the inhibitory effects of predator odor on

the GnRH pulse generator in mice (3) and activation of MePD Ucn3 neurons delayed puberty

in mice (32). Therefore, we know CRFR2 neurons are involved in processing stressful

information and coordinating the activity of the GnRH pulse generator accordingly.

Additionally, CRF is also implicated in reducing pulsatile LH secretion via CRFR1 expressing

neurons in mice (51). Chronic stress levels of CORT decrease LH pulsatility (46) and recently

an acute rise in CORT sustained for 30-60 min was shown to inhibit pulsatile LH release, but

only in OVX mice with estrogen replacement (52) and after a delay of approximately 30 min

following the CORT rise. From our observations, inhibition of LH pulsatility occurs

immediately after the stimulation of MePD Ucn3 projections in the PVN in OVX estradiol

replaced mice, thus it is unlikely the result of negative feedback from CORT release but rather

a central suppression of GnRH pulse generator activity possibly from the activation of PVN

CRF neurons, although the mechanism involved has yet to be established. The PVN CRF

neuronal population was not traditionally considered important for mediating stress-induced

suppression of pulsatile LH secretion, as PVN lesion studies have shown that this region is not

critical in mediating the stress-induced suppression of the reproductive axis (53). However,

with the recent observations showing that chemogenetic activation of PVN CRF neurons

indirectly inhibits the GnRH pulse generator it is plausible that the stress neural circuitry

includes upstream signalling via the MePD. Therefore, a possible route may involve stressful

information processed by the PVN CRF neurons relayed via the MePD Ucn3 neuronal circuitry

to the ARC KNDy neurons to modulate their activity. The neighbouring central nucleus of the

amygdala is known to act as a relay centre in pain control where the central lateral part receives

nociceptive input from the parabrachial nucleus and the central medial part is involved in

regulating the output of defensive behaviour and nociception (54). Local and efferent MePD

Ucn3 signalling plays a critical dual role in regulating GnRH pulse generator activity and

CORT secretion, thus MePD Ucn3 neuronal populations and their projections may represent a

nodal centre in the interaction between the reproductive and stress axes.

Our findings show for the first time that stimulation of local Ucn3 neurons in the MePD inhibits

GnRH pulse generator activity involving an interplay between Ucn3, GABA and glutamate

signalling within the MePD. Furthermore, we interrogate the MePD Ucn3 neuron projection

terminals in the PVN demonstrating that stimulation of the MePD Ucn3 efferent projections in

the PVN elevates CORT secretion to stress levels while suppressing the GnRH pulse generator.

How the local neural circuit within the MePD engages with efferent projections from the MePD to the PVN and the interplay between Ucn3, GABA and glutamate neurons to mediate the suppressive effect on LH pulsatility and the stimulatory effect on the HPA axis remains an exciting question for the future.

Bibliography

1. Ressler KJ. Amygdala Activity, Fear, and Anxiety: Modulation by Stress. Biol Psychiatry. 2010;67(12):1117–9.

2. Lin Y, Li XF, Lupi M, Kinsey-Jones JS, Shao B, Lightman SL, O'Byrne KT . The role of the medial and central amygdala in stress-induced suppression of pulsatile LH secretion in female rats. Endocrinology. 2011;152(2):545–55.

3. Ivanova D, Li XF, McIntyre C, Liu Y, Kong L, O'Byrne KT . Urocortin3 in the Posterodorsal Medial Amygdala Mediates Stress-induced Suppression of LH Pulsatility in Female Mice. Endocrinology. 2021;162(12):bqab206.

4. Lass G, Li XF, de Burgh RA, He W, Kang Y, Hwa-Yeo S, Sinnett-Smith LC, Manchishi SM, Colledge WH, Lightman SL, O'Byrne KT . Optogenetic stimulation of kisspeptin neurones within the posterodorsal medial amygdala increases luteinising hormone pulse frequency in female mice. J Neuroendocrinol. 2020;32(2):e12823.

5. Li XF, Adekunbi DA, Alobaid HM, Li S, Pilot M, Lightman SL, O'Byrne KT . Role of the posterodorsal medial amygdala in predator odour stress-induced puberty delay in female rats. J Neuroendocrinol. 2019;31(6):e12719.

6. Adekunbi DA, Li XF, Li S, Adegoke OA, Iranloye BO, Morakinyo AO, Lightman SL, Taylor PD, Poston L, O'Byrne KT . Role of amygdala kisspeptin in pubertal timing in female rats. PLoS One. 2017;12(8):e0183596.

7.   Moore AM, Coolen LM, Lehman MN. Kisspeptin/Neurokinin B/Dynorphin (KNDy)

cells as integrators of diverse internal and external cues: evidence from viral-based

monosynaptic tract-tracing in mice. Sci Rep. 2019;9(1):1–5.

8.   Yeo SH, Kyle V, Blouet C, Jones S, Colledge WH. Mapping neuronal inputs to Kiss1

neurons in the arcuate nucleus of the mouse. PLoS One. 2019;14(3):e0213927.

9.   Qiu J, Nestor CC, Zhang C, Padilla SL, Palmiter RD, Kelly MJ, Rønnekleiv OK .

High-frequency stimulation-induced peptide release synchronizes arcuate kisspeptin

neurons and excites GnRH neurons. Elife. 2016;5:e16246.

10.   Clarkson J, Han SY, Piet R, McLennan T, Kane GM, Ng J, Porteous RW, Kim JS,

Colledge WH, Iremonger KJ HA. Definition of the hypothalamic GnRH pulse

generator in mice. Proc Natl Acad Sci U S A. 2017;114(47):e10216–23.

11.   Voliotis M, Li XF, De Burgh R, Lass G, Lightman SL, O'Byrne KT, Tsaneva-

Atanasova K . The Origin of GnRH pulse generation: An integrative mathematical-

experimental approach. J Neurosci. 2019;39(49):9738–47.

12.   Moore AM, Coolen LM, Lehman MN. In vivo imaging of the GnRH pulse generator

reveals a temporal order of neuronal activation and synchronization during each pulse.

Proc Natl Acad Sci U S A. 2022;119(6).

13.   Han SY, McLennan T, Czieselsky K, Herbison AE. Selective optogenetic activation of

arcuate kisspeptin neurons generates pulsatile luteinizing hormone secretion. Proc Natl

Acad Sci U S A. 2015;112(42):13109–14.

14.   Voliotis M, Li XF, De Burgh RA, Lass G, Ivanova D, McIntyre C, O'Byrne K,

Tsaneva-Atanasova K . Modulation of pulsatile GnRH dynamics across the ovarian

cycle via changes in the network excitability and basal activity of the arcuate

kisspeptin network. Elife. 2021;10.

15. Martin EI, Ressler KJ, Binder E, Nemeroff CB. The Neurobiology of Anxiety

Disorders: Brain Imaging, Genetics, and Psychoneuroendocrinology. Psychiatr Clin

North Am. 2009;32(3):549.

16. Nuss P. Anxiety disorders and GABA neurotransmission: A disturbance of

modulation. Neuropsychiatr Dis Treat. 2015;11:165–75.

17. Takahashi LK. Olfactory systems and neural circuits that modulate predator odor fear.

Front Behav Neurosci. 2014;8:72.

18. Van-Hover C, Li C. Stress-activated afferent inputs into the anterior parvicellular part

of the paraventricular nucleus of the hypothalamus: Insights into urocortin 3 neuron

activation. Brain Res. 2015;1611:29–43.

19. Jamieson PM, Li C, Kukura C, Vaughan J, Vale W. Urocortin 3 modulates the

neuroendocrine stress response and is regulated in rat amygdala and hypothalamus by

stress and glucocorticoids. Endocrinology. 2006;147(10):4578–88.

20. Fekete ÉM, Zhao Y, Li C, Sabino V, Vale WW, Zorrilla EP. Social defeat stress

activates medial amygdala cells that express type 2 corticotropin-releasing factor

receptor mRNA. Neuroscience. 2009;162(1):5–13.

21. Shemesh Y, Forkosh O, Mahn M, Anpilov S, Sztainberg Y, Manashirov S,

Shlapobersky T, Elliott E, Tabouy L, Ezra G, Adler ES . Ucn3 and CRF-R2 in the

medial amygdala regulate complex social dynamics. Nat Neurosci. 2016;19(11):1489–

96.

22. Ström JO, Theodorsson A, Ingberg E, Isaksson IM, Theodorsson E. Ovariectomy and

17β-estradiol Replacement in Rats and Mice: A Visual Demonstration. J Vis Exp.

2012;(64):4013.

23. Phumsatitpong C, Moenter SM. Estradiol-Dependent stimulation and suppression of

gonadotropin-Releasing hormone neuron firing activity by corticotropin-Releasing

hormone in female mice. Endocrinology. 2018;159(1):414–25.

24. Czieselsky K, Prescott M, Porteous R, Campos P, Clarkson J, Steyn FJ, Campbell RE,

Herbison AE . Pulse and Surge Profiles of Luteinizing Hormone Secretion in the

Mouse. Endocrinology. 2016;157(12):4794–802.

25. Paxinos G, Franklin KB . Mouse Brain in Stereotaxic Coordinates. Acad Press.

2019;5:246.

26. McCosh RB, Kreisman MJ, Breen KM. Frequent tail-tip blood sampling in mice for

the assessment of pulsatile luteinizing hormone secretion. J Vis Exp. 2018;(137).

27. Pardo-Bellver C, Cádiz-Moretti B, Novejarque A, Martínez-García F, Lanuza E.

Differential efferent projections of the anterior, posteroventral, and posterodorsal

subdivisions of the medial amygdala in mice. Front Neuroanat. 2012;6(33):1–26.

28. Stephens SBZ, Raper J, Bachevalier J, Wallen K. Neonatal amygdala lesions advance

pubertal timing in female rhesus macaques. Psychoneuroendocrinology. 2015;51:307–

17.

29. Li XF, Hu MH, Hanley BP, Lin YS, Poston L, Lightman SL, O'Byrne KT . The

posterodorsal medial amygdala regulates the timing of puberty onset in female rats.

Endocrinology. 2015;156(10):3725–36.

30. Keshavarzi S, Sullivan RKP, Ianno DJ, Sah P. Functional properties and projections of

neurons in the medial amygdala. J Neurosci. 2014;34(26):8699–715.

31. Comninos AN, Anastasovska J, Sahuri-Arisoylu M, Li XF, Li S, Hu M, Jayasena CN,

Ghatei MA, Bloom SR, Matthews PM, O'Byrne KT . Kisspeptin signaling in the

amygdala modulates reproductive hormone secretion. Brain Struct Funct.

2016;221(4):2035–47.

32. Ivanova D, Li X, Liu Y, McIntyre C, Fernandes C, Lass G, Kong L, O'Byrne KT .

Role of Posterodorsal Medial Amygdala Urocortin-3 in Pubertal Timing in Female

Mice. Front Endocrinol (Lausanne). 2022;0:910.

33. Cavalcante JC, Sita LV, Mascaro MB, Bittencourt JC, Elias CF. Distribution of

urocortin 3 neurons innervating the ventral premammillary nucleus in the rat brain.

Brain Res. 2006;1089(1):116–25.

34. Li C, Vaughan J, Sawchenko PE, Vale WW. Urocortin III-immunoreactive projections

in rat brain: Partial overlap with sites of type 2 corticotrophin-releasing factor receptor

expression. J Neurosci. 2002;22(3):991–1001.

35. Deussing JM, Breu J, Kühne C, Kallnik M, Bunck M, Glasl L, Yen YC, Schmidt MV,

Zurmühlen R, Vogl AM, Gailus-Durner V . Urocortin 3 modulates social

discrimination abilities via corticotropin-releasing hormone receptor type 2. J

Neurosci. 2010;30(27):9103–16.

36. Hegoburu C, Parrot S, Ferreira G, Mouly AM. Differential involvement of amygdala

and cortical NMDA receptors activation upon encoding in odor fear memory. Learn

Mem. 2014;21(12):651–5.

37. Liu J, Yu B, Neugebauer V, Grigoriadis DE, Rivier J, Vale WW, Shinnick-Gallagher

P, Gallagher JP . Corticotropin-releasing factor and Urocortin I modulate excitatory

glutamatergic synaptic transmission. J Neurosci. 2004;24(16):4020–9.

38. Autry AE, Wu Z, Kapoor V, Kohl J, Bambah-Mukku D, Rubinstein ND, Marin-

Rodriguez B, Carta I, Sedwick V, Tang M, Dulac C . Urocortin-3 neurons in the

mouse perifornical area promote infant-directed neglect and aggression. Elife.

2021;10.

39. Kuperman Y, Issler O, Regev L, Musseri I, Navon I, Neufeld-Cohen A, Gil S, Chen A

. Perifornical urocortin-3 mediates the link between stress-induced anxiety and energy

homeostasis. Proc Natl Acad Sci U S A. 2010;107(18):8393–8.

40. Yip SH, Liu X, Hessler S, Cheong I, Porteous R, Herbison AE. Indirect Suppression of

Pulsatile LH Secretion by CRH Neurons in the Female Mouse. Endocrinology.

2021;162(3).

41. Rivest S, Plotsky PM, Rivier C. CRF alters the infundibular LHRH secretory system

from the medial preoptic area of female rats: possible involvement of opioid receptors.

Neuroendocrinology. 1993;57(2):236–46.

42. Bagosi Z, Csabafi K, Karasz G, Jászberényi M, Földesi I, Siska A, Szabó G, Telegdy

G . The effects of the urocortins on the hypothalamic-pituitary-adrenal axis -

similarities and discordancies between rats and mice. Peptides. 2019;112:1–13.

43. Anthony TE, Dee N, Bernard A, Lerchner W, Heintz N, Anderson DJ. Control of

stress-induced persistent anxiety by an extra-amygdala septohypothalamic circuit.

Cell. 2014;156(3):522–36.

44. Li XF, Mitchell JC, Wood S, Coen CW, Lightman SL, O'Byrne KT . The effect of

oestradiol and progesterone on hypoglycaemic stress-induced suppression of pulsatile

luteinizing hormone release and on corticotropin-releasing hormone mRNA expression

in the rat. J Neuroendocrinol. 2003;15(5):468–76.

45. Zavala E, Voliotis M, Zerenner T, Tabak J, Walker JJ, Li XF, Terry JR, Lightman SL,

O'byrne K, Tsaneva-Atanasova K . Dynamic Hormone Control of Stress and Fertility.

Front Physiol. 2020;11:1457.

46. Kreisman MJ, McCosh RB, Tian K, Song CI, Breen KM. Estradiol Enables Chronic

Corticosterone to Inhibit Pulsatile Luteinizing Hormone Secretion and Suppress Kiss1

Neuronal Activation in Female Mice. Neuroendocrinology. 2020;110(6):501–16.

47.   Dabrowska J, Hazra R, Guo JD, DeWitt S, Rainnie DG. Central CRF neurons are not

569   created equal: Phenotypic differences in CRF-containing neurons of the rat

570   paraventricular hypothalamus and the bed nucleus of the stria terminalis. Front

571   Neurosci. 2013;0(7 AUG):156.

572   48.   Deussing JM, Chen A. The corticotropin-releasing factor family: Physiology of the

573   stress response. Vol. 98, Physiological Reviews. 2018. p. 2225–86.

574   49.   Li XF, Bowe JE, Kinsey-Jones JS, Brain SD, Lightman SL, O'Byrne KT . Differential

575   role of corticotropin-releasing factor receptor types 1 and 2 in stress-induced

576   supression of pulsatile luteinising hormone secretion in the female rat. J

577   Neuroendocrinol. 2006;18(8):602–10.

578   50.   Li XF, Bowe JE, Lightman SL, O'Byrne KT . Role of corticotropin-releasing factor

579   receptor-2 in stress-induced suppression of pulsatile luteinizing hormone secretion in

580   the rat. Endocrinology. 2005;146(1):318–22.

581   51.   Raftogianni A, Roth LC, García-González D, Bus T, Kühne C, Monyer H, Spergel DJ,

582   Deussing JM, Grinevich V . Deciphering the Contributions of CRH Receptors in the

583   Brain and Pituitary to Stress-Induced Inhibition of the Reproductive Axis. Front Mol

584   Neurosci. 2018;11:305.

585   52.   Kreisman MJ, McCosh RB, Breen KM . Role of CORT Duration and Estradiol

586   Dependence for Stress-level of CORT to Inhibit Pulsatile LH Secretion in Female

587   Mice. J Endocr Soc. 2021;Apr;5(Supplement_1):A552-.

588   53.   Serge S, Rivier C. Influence of the paraventricular nucleus of the hypothalamus in the

589   alteration of neuroendocrine functions induced by intermittent footshock or

590   interleukin. Endocrinology. 1991;129(4):2049–58.

54.   Wank I, Pliota P, Badurek S, Kraitsy K, Kaczanowska J, Griessner J, Kreitz S, Hess A,

Haubensak W . Central amygdala circuitry modulates nociceptive processing through

differential hierarchical interaction with affective network dynamics. Commun Biol.

2021;4(1):1–10.

Figure 1. Optogenetic stimulation of posterodorsal medial amygdala (MePD) urocortin-3

(Ucn3) neurons with blue light at 10 Hz, 10 mW suppressed pulsatile LH secretion in adult

ovariectomised (OVX) Ucn3-cre-tdTomato female mice. Infusion of bicuculline (BIC),

GABA$_A$ receptor antagonist, CGP, GABA$_B$ receptor antagonist, and AP5 and CNQX, NMDA

and AMPA receptor antagonists, respectively, into the posterodorsal medial amygdala (MePD)

during optogenetic stimulation of MePD urocortin-3 (Ucn3) neurons with blue light at 10 Hz,

10 mW completely blocked the suppressive effect of MePD Ucn3 neuronal activation on LH

pulsatility. Representative LH pulse profiles showing the effects of (A) control AAV-YFP

mice administered with aCSF receiving optical stimulation of 10 Hz, 10 mW, (B) AAV-ChR2

mice administered with aCSF receiving optical stimulation of 10 Hz, 10 mW, (C) Summary of

LH pulse interval for the pre-stimulation control period (1 h) and optical stimulation/infusion

period (1 h), (D) AAV-ChR2 mice administered with BIC, (E) AAV-ChR2 mice administered

with BIC receiving optical stimulation of 10 Hz, 10 mW, (F) Summary of LH pulse interval,

(G) AAV-ChR2 mice administered with CGP, (H) AAV-ChR2 mice administered with CGP

receiving optical stimulation of 10 Hz, 10 mW, (I) Summary of LH pulse interval, (J) AAV-

ChR2 mice administered with a combination of AP5 and CNQX, (K) AAV-ChR2 mice

administered with with a combination of AP5 and CNQX receiving optical stimulation of 10

Hz, 10 mW, (L) Summary of LH pulse interval. LH pulses detected by the DynePeak algorithm

are indicated with an asterisk located above each pulse on the representative LH pulse profiles.

For A-C +++$p<0.0001$ control period vs optic stimulation + aCSF (AAV-ChR2: n = 9);

##$p<0.001$ optic stimulation vs control group (AAV-YFP: n = 3; no stimulation: n = 2). For

D-F n = 8-9 per group. For G-I n = 7 per group. For J-L n = 6 per group.

Figure 2. Optogenetic stimulation of posterodorsal medial amygdala (MePD) urocortin-3

(Ucn3) neuronal terminals in the hypothalamic paraventricular nucleus (PVN) with blue light

at 20 Hz, 15 mW elevated corticosterone (CORT) secretion in adult ovariectomised (OVX) +

E_2 Ucn3-cre-tdTomato female mice. CORT secretion time-course for mice receiving optical

stimulation initiated at 0 min and terminated at 60 min (1 h duration), followed by a 1 h

recovery period (60-120 min) for the AAV-ChR2 group (blue line, circles) and control group

(red line, squares). +$p < 0.05$ AAV-ChR2 (n = 6) at time point 30 min and 60 min vs. control

(AAV-YFP: n = 3) at time point 30 min and 60 min.

Figure 3. Optogenetic stimulation of posterodorsal medial amygdala (MePD) urocortin-3

(Ucn3) neuronal terminals in the hypothalamic paraventricular nucleus (PVN) with blue light

at 20 Hz, 15 mW reduced LH pulsatility in adult ovariectomised (OVX) + E_2 Ucn3-cre-

tdTomato female mice. Representative LH pulse profiles showing the effects of (A) AAV-

YFP/WT mice receiving optical stimulation of 20 Hz, 5 mW, (B) AAV-ChR2 mice receiving

optical stimulation of 20 Hz, 5 mW, (C) Summary of LH pulse interval for the pre-stimulation

control period (1 h) and optic stimulation period (1 h). LH pulses detected by the DynePeak

algorithm are indicated with an asterisk located above each pulse on the representative LH

pulse profiles. +++$p < 0.0001$ control period vs optic stimulation for AAV-ChR2 mice (n = 8);

###$p < 0.0001$ AAV-YFP/WT (AAV-YFP, n = 3; wild-type with AAV-ChR2, n = 2; no

stimulation, n = 2) vs AAV-ChR2 mice during optic stimulation.

Figure 4. Expression of AAV9-EF1a-double floxed-hChR2(H134R)-EYFP-WPRE-HGHpA

in posterodorsal medial amygdala (MePD) Urocortin-3 (Ucn3) neurons. (A-C) Representative

dual fluorescence photomicrographs of the MePD from a Ucn3-cre-tdTomato female OVX

mouse injected with AAV9-EF1a-double floxed-hChR2(H134R)-EYFP-WPRE-HGHpA.

Ucn3 neurons labelled with YFP (A) and tdTomato (B) appear yellow/orange (C). Scale bars

represent A, B, C, 20 μm.

Figure 5. Expression of AAV9-EF1a-double floxed-hChR2(H134R)-EYFP-WPRE-HGHpA

in Urocortin-3 (Ucn3) neuronal projection in the paraventricular nucleus of the hypothalamus

(PVN) originating from the posterodorsal medial amygdala (MePD). (A-Aii) Representative

dual fluorescence photomicrographs of the MePD from a Ucn3-cre-tdTomato female OVX +

E_2 mouse injected with AAV9-EF1a-double floxed-hChR2(H134R)-EYFP-WPRE-HGHpA.

Ucn3 neurons and projections labelled with (A) YFP and tdTomato merged, (Ai) higher power

view of YFP labelled MePD Ucn3 projections in the PVN and (Aii) higher power view of

tdTomato labelled MePD Ucn3 projections in the PVN. Scale bars represent A, 100 μm, Ai

and Aii, 10 μm. Blue box shows the YFP labelled MePD Ucn3 projections in the PVN, white

arrows point to YFP labelled MePD Ucn3 projections in the PVN and yellow arrow points to

optic cannula implantation site. 3V; third ventricle.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5