Posterodorsal medial amygdala urocortin-3, GABA and glutamate mediate suppression

of LH pulsatility in female mice

Deyana Ivanova[1], Xiao-Feng Li[1], Caitlin McIntyre[1] and Kevin T O'Byrne[1]

[1]Department of Women and Children's Health, School of Life Course and Population Sciences,

Faculty of Life Science and Medicine, King's College London, UK

Short title: MePD Ucn3 effect on LH and corticosterone release

Key words: Ucn3, GABA, glutamate, MePD, PVN, LH pulsatility, CORT secretion

Corresponding author's contact details: Deyana Ivanova (deyana.ivanova@kcl.ac.uk) and

Kevin T O'Byrne (kevin.obyrne@kcl.ac.uk), Department of Women and Children's Health,

School of Life Course and Population Sciences, Faculty of Life Science and Medicine, King's

College London, 2.92W Hodgkin Building, Guy's Campus, London SE1 1UL, UK.

Reprint requests: Deyana Ivanova, Department of Women and Children's Health, School of

Life Course Sciences, Faculty of Life Science and Medicine, King's College London, 2.92W

Hodgkin Building, Guy's Campus, London SE1 1UL, UK.

Grants supporting paper: UKRI: BBSRC (BB/S000550/1 & BB/W005913/1). DI is a PhD

student funded by MRC-DTP studentship.

The authors have nothing to disclose.

Abstract

The posterodorsal subnucleus of the medial amygdala (MePD) is an upstream modulator of the hypothalamic-pituitary-gonadal (HPG) and hypothalamic-pituitary-adrenal (HPA) axes. Inhibition of MePD urocortin-3 (Ucn3) neurons prevents psychological stress-induced suppression of LH pulsatility while blocking the stress-induced elevations in corticosterone (CORT) secretion in female mice. We explore the neurotransmission and neural circuitry suppressing the GnRH pulse generator by MePD Ucn3 neurons and we further investigate whether MePD Ucn3 efferent projections to the PVN control CORT secretion and LH pulsatility. Ucn3-cre-tdTomato female ovariectomised (OVX) mice were unilaterally injected with AAV-ChR2 and implanted with optofluid cannulae targeting the MePD. We optically activated Ucn3 neurons in the MePD with blue light at 10Hz and monitored the effect on LH pulses. Next, we combined optogenetic stimulation of MePD Ucn3 neurons with pharmacological antagonism of $GABA_A$ or $GABA_B$ receptors with bicuculline or CGP, respectively, as well as a combination of NMDA and AMPA receptor antagonists, AP5 and CNQX respectively, and observed the effect on pulsatile LH secretion. A separate group of Ucn3-cre-tdTomato OVX mice with 17ß-estradiol (E_2) replacement were unilaterally injected with AAV-ChR2 in the MePD and implanted with fibreoptic cannulae targeting the PVN. We optically stimulated the MePD Ucn3 efferent projections in the PVN with blue light at 20Hz and monitored the effect on CORT secretion and LH pulses. We reveal for the first time that activation of Ucn3 neurons in the MePD inhibits GnRH pulse generator frequency via GABA and glutamate signalling within the MePD, while MePD Ucn3 projections to the PVN modulate the HPG and HPA axes.

Introduction

Stress exerts a profound suppressive effect on reproduction in mammals, including rodents and humans. The amygdala, part of the limbic system, is a key emotional centre integrating incoming cues from the external environment, such as anxiogenic signals, with the reproductive and stress axes (1). The posterodorsal sub-nucleus of the medial amygdala (MePD) is an upstream modulator of pulsatile luteinizing hormone (LH) secretion (2–4) and exerts an inhibitory break on pubertal timing (5,6). The MePD sends direct projections, of an unknown phenotype, to the kisspeptin (kiss1) neuronal population in the hypothalamic arcuate nucleus (ARC) (7,8) known as KNDy because they co-express neurokinin B (NKB) and dynorphin A (Dyn) (9–11). The synchronous activity of the ARC KNDy network drives pulsatile gonadotropin-releasing hormone (GnRH) and LH release (11–14). The MePD also contains a kiss1 neuronal population and selective optogenetic activation of these neurons increases GnRH pulse generator frequency in mice (4) and unpublished observations show this is an effect involving both MePD GABA and glutamate signalling.

We have recently shown that urocortin-3 (Ucn3), a member of the corticotropin-releasing factor (CRF) stress neuropeptide family, and its receptor CRF type 2 (CRFR2) signalling in the MePD are involved in mediating the suppressive effect of psychosocial stress on LH pulsatility (3). However, the underlying neural mechanisms involved in mediating the inhibitory effect of MePD Ucn3 neurons on LH pulsatility remain to be established. GABA and glutamate signalling in the amygdala are implicated in the modulation of anxiety behaviour (15,16). The amygdala also modulates the stress response through the hypothalamic pituitary adrenal (HPA) axis. The medial amygdala activates the HPA axis in response to predator odor stress (17) and the MePD in particular has been shown to send stress-activated efferents to the paraventricular nucleus of hypothalamus (PVN) (18). Furthermore, MePD Ucn3 and CRFR2-positive neurons are implicated in regulating the stress response, with restraint stress increasing

MePD Ucn3 (19) and social defeat elevating MePD CRFR2 expression in rodents (20).

Moreover, MePD Ucn3 and CRFR2-positive neurons project to the PVN (21), and we have

recently shown that chemogenetic inhibition of MePD Ucn3 neurons prevents psychogenic

stress-induced corticosterone (CORT) release in female mice (3).

In this study, we aimed to determine whether activation of MePD Ucn3 neurons inhibits

pulsatile LH secretion via GABA and or glutamate signalling within this amygdala sub-

nucleus. To achieve this, we will combine selective optogenetic activation of MePD Ucn3

neurons with pharmacological antagonism of GABA$_A$ or GABA$_B$ receptors, or a combination

of N-methyl-D-aspartate (NMDA) and α-amino-3-hydroxy-5-methyl-4-isoxazole propionic

acid (AMPA) receptors, respectively, using chronically implanted intra-MePD optofluid

cannulae while collecting serial blood samples for LH pulse measurement in Ucn3-Cre-

tdTomato female mice. Additionally, we will investigate whether optogenetic stimulation of

MePD Ucn3 projection terminals in the PVN induce CORT release and suppress LH pulsatility.

Methods

Mice

Cryopreserved sperm of Ucn3-cre mice (strain Tg(Ucn3-cre)KF43Gsat/Mmucd; congenic on

C57BL/6 background) was acquired from MMRRC GENSAT and heterozygous transgenic

breeding pairs of Ucn3-Cre mice were recovered via insemination of female C57Bl6/J mice at

King's College London, as described previously (3). Ucn3-Cre mice were genotyped using

PCR for the detection of heterozygosity and Ucn3-cre mice were bred with td-Tomato mice

(strain   B6.Cg-Gt(ROSA)26Sortm9(CAG-tdTomato)Hze/J;   congenic   on   C57BL/6

background) acquired from The Jackson Laboratory, Bar Harbor, ME, USA to produce Ucn3-

cre-tdTomato reporter mice, as described previously (3,21). Female Ucn3-cre-tdTomato mice

weighing 19-23 g and aged 6-8 weeks were singly housed in individually ventilated cages

sealed with a HEPA-filter at 25 ± 1 °C in a 12:12 h light/dark cycle, lights on at 07:00 h

equipped with nesting material, wood-chip bedding and food and water ad libitum.   All

procedures were carried out following the United Kingdom Home Office Regulations and

approved by the Animal Welfare and Ethical Review Body Committee at King's College

London.

Stereotaxic injection of adeno-associated-virus carrying the channelrhodopsin construct

and fibre optic or optofluid cannula implantation

All surgical procedures were carried out under aseptic conditions with general anaesthesia

using ketamine (Vetalar, 100 mg/kg, i.p.; Pfizer, Sandwich, UK) and xylazine (Rompun, 10

mg/kg, i.p.; Bayer, Leverkusen, Germany).   Ucn3-Cre-tdTomato mice were secured in a David

Kopf stereotaxic frame (Kopf Instruments, Model 900) and either solely bilaterally

ovariectomised (OVX) or implanted with an 17ß-estradiol (E_2) silastic capsule. E_2 was

dissolved in sesame oil to reach a final concentration of 36 µg E_2/mL and filled in 14 mm

capsules of inner/outer diameter: 1.575/3.175 mm, as previously described (22). The reference

study reported that at 21 days of implantation, serum E_2 concentrations were within

physiological concentrations equivalent to those found in mice in the diestrous stage of the

cycle. However, by 35 days of implantation the serum E_2 concentrations had slowly declined

but still within the range found in diestrous mice. Due to the technically demanding nature of

our experiments the E_2 capsules were implanted for longer than 35 days post-surgery during

the experimental period. Therefore, E_2 levels may be lower than expected for the diestrous

phase. E_2 is known to enhance the suppressive effects of stress where the action of the gonadal

steroids sensitizes the HPA axis and augments stress-induced inhibition of the GnRH pulse

generator in many species, from rodents to primates (42–45).   We had two experimental

groups; OVX only and OVX + E_2 capsule. The mouse brain atlas of Paxinos and Franklin (23)

was used to obtain target coordinates for the MePD (2.30 mm lateral, -1.55 mm from bregma,

at a depth of -4.94 mm below the skull surface). To reveal the skull an incision was made of

the scalp and one small hole was drilled above the location of the right MePD. AAV9-EF1a-

double floxed-hChR2(H134R)-EYFP-WPRE-HGHpA (300 nL; $3x10^{11}$ GC/ml; Serotype:9;

Addgene, Massachusetts, USA) was unilaterally injected into the right MePD using a 2-µL

Hamilton micro syringe (Esslab, Essex, UK) over 10 min, using the robot stereotaxic system

(Neurostar, Tubingen, Germany), performed for the targeted expression of ChR2-EYFP in

MePD Ucn3 neurons. After injection, the needle was left in position for a further 5 min and

lifted slowly over 2 min. Cre-positive mice received the AAV-ChR2 injection (test mice) or a

control virus AAV-YFP (Addgene) (control mice). The control virus does not contain the

ChR2 construct. The mice were then implanted with a dual optofluid cannula (Doric Lenses,

Quebec, Canada) at the same AP and ML coordinates as the viral injection site, however a

different DV was used such that the internal cannula targets the MePD and the fibre optic

cannula is situated 0.2 mm above the MePD site. Once in position the optofluid cannula was

secured on the skull using dental cement (Super-Bond Universal Kit, Prestige Dental, UK) and

the incision of the skin was closed with suture. Mice were left to recover for one week and

after recovery period, mice were handled daily to acclimatise to experimental procedures for a

further 2 weeks. Thirteen mice received the AAV-ChR2 injection and were implanted with an

optofluid cannula in the MePD. Five mice received the control AAV-YFP and were implanted

with an optofluid cannula in the MePD. Eleven mice received the AAV-ChR2 injection in the

MePD and were implanted with an optic fibre in the PVN. Five mice received the control

AAV-YFP and were implanted with an optic fibre in the PVN.

In vivo optogenetic stimulation of MePD Urocortin-3 neurons

In order to test the effect of optogenetic stimulation of MePD Ucn3 neurons on LH pulsatility

the ferrule of the implanted optofluid cannula, on the right side of the MePD, was attached via

a ceramic mating sleeve to a multimode fibre optic rotary joint patch cable (Thorlabs LTD,

Ely, UK) at a length allowing for free movement of the mouse in their cage and blue light (473

nm wavelength; 10 mW) was delivered using a Grass SD9B stimulator-controlled laser

(Laserglow Technologies, Toronto, Canada). The mice were left to acclimatise for 1 h.

Following the acclimatisation period blood samples (5 µL) were collected every 5 min for 2 h

where after 1 h of control blood sampling, Ucn3-cre-tdTomato mice received optic stimulation

at 10 Hz with a 10 ms pulse interval and pattern stimulation of 5 seconds on and 5 seconds off

for the remaining 1 h.

Intra-MePD administration of bicuculline (BIC), CGP-35348 or AP5 + CNQX during

optogenetic stimulation of MePD Urocortin-3 neurons

Neuropharmacological manipulation of GABA and glutamate receptor signalling in the MePD

with or without optogenetic stimulation of MePD Ucn3 neurons was done to test whether

MePD Ucn3 induced suppression of LH pulsatility involves GABA or glutamate receptor

signalling. OVX mice implanted with a optofluid cannula in the MePD were subjected to the

tail-tip blood collection procedure, as previously described (24). Infusion of drugs with or

without optogenetic stimulation and blood sampling were performed between 09:00-13:00 h,

where 5 µl blood was collected every 5 min for 2 h. Internal cannula (Doric Lenses) attached

to extension tubing (0.58 mm ID, 0.96 mm OD), preloaded with bicuculline (BIC) ($GABA_A$

receptor antagonist; Sigma-Aldrich), CGP-35348 ($GABA_B$ receptor antagonist; Sigma-

Aldrich), AP5 + CNQX cocktail (NMDA and AMPA receptor antagonists; Tocris) or aCSF as

vehicle control, was inserted into the guide cannula of the optofluid implant and mice were

connected to the laser as described above. The tubing for drug infusion extended beyond the

cage and the distal ends were attached to 10-μl Hamilton syringes (Waters Ltd, Elstress, UK)

fitted into a PHD 2000 Programmable syringe pump (Harvard Apparatus, Massachusetts,

USA), allowing for a continuous infusion of the drug at a constant rate and mice were kept in

the cage throughout the experiment, freely-moving with food and water *ad libitum*. After 55

min of control blood sampling, the mice were given an initial bolus injection, 0.30 μl at a rate

of 0.06 μl/min over 5 min, of BIC, CGP-35348, AP5 + CNQX or aCSF. The laser was turned

on at 60 min and the bolus injection was followed by a continuous infusion, 0.8 μl at a rate of

0.01 μl/min over 1 h, of BIC, CGP-35348, AP5 + CNQX or aCSF. BIC, CGP-35348, AP5 and

CNQX were dissolved in aCSF to reach bolus concentrations of 20 pmol, 4.5 nmol, 1.2 nmol

and 0.5 nmol respectively. The concentrations for the continuous infusion of BIC, CGP-35348,

AP5 and CNQX were 68 pmol, 15 nmol, 2nmol and 1 nmol respectively. In the absence of

optic stimulation, the same regimen was applied. Data collection was conducted over a period

of 4-6 weeks. The mice receive all treatments in a random order, with at least 2 days between

experiments.

MePD Urocortin-3 terminal optogenetic stimulation in the PVN

To test the effect of optogenetic stimulation of MePD Ucn3 neuronal terminals in the PVN on

LH pulsatility and CORT release, OVX + E_2 Ucn3-cre-tdTomato mice with AAV-ChR2

injection in the right MePD implanted with a fibre optic cannula in the right PVN were

subjected to the tail-tip blood collection procedure, as previously described (24). For LH

measurement, blood samples were collected every 5 min for 2 h, as described above. Mice

were connected to the laser as described above. Optogenetic stimulation was initiated after the

1 h control blood sampling period and Ucn3-cre-tdTomato mice received optic stimulation at

20 Hz, 15 mW with a 10 ms pulse interval and pattern stimulation of 5 seconds on and 5

seconds off for the remaining 1 h. For CORT measurements, 15 μl blood samples were

collected on a separate occasion and blood samples were stored in tubes containing 5 µl

heparinised saline (5 IU ml^{-1}). Optogenetic stimulation was initiated, as described above, 30

min into the experiment (time point, 0 min) and lasted for 1 h (time point 60 min). Remaining

blood samples were collected at 0, 30 and 60 min, with a final sample taken 1 h after the

termination of optogenetic stimulation (time point, 120 min). At the end of the experiment,

blood samples were centrifuged at 13,000 RPM for 20 min at 20°C and plasma stored at -20°C.

Data collection was conducted over a period of 4-6 weeks. All experiments were performed

between 9:00-12:00 h.

Validation of AAV injection and cannula implant site

Upon completion of experiments, mice were anaesthetised with a lethal dose of ketamine and

transcardial perfusion was performed with heparinised saline for 5 min followed by ice-cold

4% paraformaldehyde (PFA) in phosphate buffer (pH 7.4) for 15 min using a pump (Minipuls,

Gilson, Villiers Le Bel, France). Mice brains were collected immediately and post-fixed in

15% sucrose in 4% PFA at 4 °C and left to sink. Once sunk, they were transferred to 30%

sucrose in phosphate-buffered saline (PBS) and left to sink. Brains were snap-frozen in

isopropanol on dry ice and stored in -80°C until further processing. Every third coronal brain

section (30-µm/section) through-out the MePD region (-1.34 mm to -2.70 mm from bregma)

was collected using a cryostat (Bright Instrument Co., Luton, UK). Sections were mounted on

microscope slides, air dried and covered with ProLong Antifade mounting medium (Molecular

Probes, Inc. OR, USA). Cannula placement and AAV injection site was verified with

Axioskop 2 Plus microscope equipped with Axiovision, version 4.7 (Zeiss) by determining

whether cannula reach the MePD. For AAV-ChR2-EYFP injected Ucn3-cre-tdTomato mice

we determined whether Ucn3 neurons were infected in the MePD region by merging td-Tomato

fluorescence of Ucn3 neurons with YFP fluorescence in the MePD. The number of Ucn3

mCitrine-positive neurones co-localised with td-Tomato fluorescence in the MePD of each

animal was counted using 4 sections and the average number of neurons presented is per section

per MePD. The group mean percent was calculated by taking the average number of Ucn3

mCitrine-positive neurons out of the average number of Ucn3 neurons expressing td-Tomato

fluorescence per 4 sections and presented as mean ± SEM %, as described previously (3).

Axioskop 2 Plus microscope (Carl Zeiss) equipped with axiovision, version 4.7 (Carl Zeiss)

was used to take images. Only mice with correct AAV injection and cannula placement in the

MePD were analysed.

LH and CORT measurement

Mice were handled daily to acclimatise to the tail-tip blood sampling procedure for LH

measurement (3). Blood samples collected for LH measurement were processed using LH

ELISA, as previously reported (3). The capture antibody (monoclonal antibody, anti-bovine

LHβ subunit, AB_2665514) was purchased from Department of Animal Science at the

University of California, Davis. The mouse LH standard (AFP-5306A) and primary antibody

(polyclonal antibody, rabbit LH antiserum, AB_2665533) were obtained from Harbour-UCLA

(California, USA). The secondary antibody (Horseradish-Peroxidase (HRP)-linked donkey

anti-rabbit IgG polyclonal antibody, AB_772206) was purchased from VWR International

(Leicestershire, UK). Intra-assay and inter-assay variations were 4.6% and 10.2%, respectively

and the assay sensitivity was 0.0015 ng/mL. DynPeak algorithm was used for the detection of

LH pulses (3).   The effect of optogenetic stimulation of MePD Ucn3 neurons and

neuropharmacology studies was established by comparing the LH inter-pulse interval (IPI)

from the 1 h control period (pre-stimulation) or drug administration control period to the 1 h

experimental period, as previously described (3). The LH pulse amplitude was calculated as

the difference between the peak of an LH pulse and the baseline LH level before the onset of

the pulse. The means were compared between the pre-treatment and treatment periods for all

experiments. The LH pulse amplitude was also compared between experimental groups within

the treatment period. The mean LH value for each LH pulse profile was quantified by summing

all values from one LH pulse profile and dividing by the number of values recorded in 120

mins.

Blood samples were collected for CORT measurement were processed using a commercially

available enzyme immunoassay (EIA) kit (sheep polyclonal antibody specific for

corticosterone, AB_2877626; DetectX® Corticosterone Enzyme Immunoassay Kit, K014;

Arbor Assays, Michigan, USA), as described previously (3).

Statistics

Mice were compared between groups using a two-way ANOVA with a Tukey post-hoc and

data was presented as mean ± SEM. Statistics were performed using Igor Pro 7, Wavemetrics,

Lake Oswego, OR, USA. Data was represented as mean ± SEM and +$p<0.05$, ++$p<0.001$ and

+++$p<0.0001$ were considered to be significant.

Results

Effects of optogenetic stimulation of MePD Urocortin-3 neurons on LH pulse frequency

Optogenetic stimulation of MePD Ucn3 neurons in control mice had no effect on LH pulse

interval (Fig. 1A; AAV-YFP: n = 3) and administration of aCSF alone (AAV-YFP with no

stimulation: n = 2) had no effect on LH pulse interval. Optogenetic stimulation of MePD Ucn3

neurons with blue light at 10 Hz, 10 mW suppressed pulsatile LH secretion (Fig. 1B; AAV-

ChR2: n = 9). These data are summarized in the figure 1C. LH pulse amplitude was quantified

and data is provided in supplemental table (25).

Effects of intra-MePD administration of bicuculline (BIC), a GABA$_A$R antagonist, and CGP, a GABA$_B$R antagonist, during optical stimulation of MePD Urocortin-3 neurons on LH pulsatility

Delivery of BIC alone had no significant effect on LH pulsatility (Fig. 1D; n = 8). Intra-MePD infusion of BIC during optogenetic stimulation of MePD Ucn3 neurons with blue light at 10 Hz, 10 mW completely blocked the suppressive effect of MePD Ucn3 neuronal activation on LH pulsatility (Fig. 1E; n = 9). The results of this experiment are summarised in the figure 1F. LH pulse amplitude was quantified and data is provided in supplemental table (25).

Effects of intra-MePD administration of CGP, a GABA$_B$R antagonist, during optical stimulation of MePD Urocortin-3 neurons on LH pulsatility

Delivery of CGP alone had no significant effect on LH pulsatility (Fig. 1G; n = 7). Intra-MePD infusion of CGP during optogenetic stimulation of MePD Ucn3 neurons with blue light at 10 Hz, 10 mW completely blocked the suppressive effect of MePD Ucn3 neuronal activation on LH pulsatility (Fig. 1H; n = 7). The results of this experiment are summarised in the figure 1I. LH pulse amplitude was quantified and data is provided in supplemental table (25).

Effects of intra-MePD glutamate antagonism during optogenetic stimulation of MePD Urocortin-3 neurons on LH pulsatility

Delivery of AP5 + CNQX alone had no significant effect on LH pulsatility (Fig. 1J; n = 6). Intra-MePD infusion of AP5 (selective NMDA antagonist) + CNQX (selective AMPA antagonist) during optogenetic stimulation of MePD Ucn3 neurons with blue light at 10 Hz, 10 mW completely blocked the suppressive effect of MePD Ucn3 neuronal activation on LH pulsatility (Fig. 1K; n = 6). The results of this experiment are summarised in the figure 1L. LH pulse amplitude was quantified and data is provided in supplemental table (25).

Effect of MePD Urocortin-3 terminal stimulation in the PVN on dynamic CORT secretion

Optogenetic stimulation of MePD Ucn3 neuron terminals in the PVN with blue light at 20 Hz, 15 mW elevated CORT secretion compared to controls (Fig. 2; AAV-ChR2: n = 6). Optogenetic stimulation of MePD Ucn3 neurons and terminal projections in the PVN in control mice had no effect on dynamic CORT secretion (AAV-YFP; n = 5).

Effect of MePD Urocortin-3 terminal stimulation in the PVN on LH pulsatility

Optogenetic stimulation of MePD Ucn3 projections in the PVN in control mice had no effect on LH pulse interval (Fig. 3A and C; AAV-YFP, n = 3; wild-type with AAV-ChR2, n = 2) and no stimulation of the MePD Ucn3 projections in the PVN (no stimulation, n = 2) had no effect on LH pulse interval. Optogenetic stimulation of MePD Ucn3 neuron terminals in the PVN with blue light at 20 Hz, 15 mW reduced LH pulsatility compared to controls (Fig. 3B and D; AAV-ChR2: n = 8). These data are summarised in the figure 3E. LH pulse amplitude was quantified and data is provided in supplemental table (25).

Selective expression of ChR2 in MePD Ucn3 neurons and MePD Ucn3 neuron projections in the PVN, and validation of cannula position

Evaluation of AAV9-EF1a-double floxed-hChR2(H134R)-EYFP-WPRE-HGHpA expression in tdTomato labelled neurons from AAV-injected Ucn3-cre-tdTomato mice revealed that 87.00 ± 6.00% of MePD Ucn3 neurons expressed AAV-ChR2 and the number of tdTomato labelled Ucn3 neurons per side per slice in the MePD was counted at 49.00 ± 5.37 (mean ± SEM) with the number of AAV-ChR2 infected neurons being 43.01 ± 3.65 (mean ± SEM) (n=8). Non-specific expression of ChR2 in the MePD (YFP single labelling) has been quantified to be limited to 6 ± 4.63 (mean ± SEM) in all of 8 mice. A representative example is shown in figure 4, A-F. The optofluid cannula was also verified to be correctly positioned in the MePD in all

of 8 mice. Evaluation of AAV9-EF1a-double floxed-hChR2(H134R)-EYFP-WPRE-HGHpA

expression in Ucn3 projections from the MePD to the PVN showed YFP labelled fibres in the

PVN figure 5, A-Aiii. Local Ucn3 neurons in the PVN are labelled with tdTomato. We cannot

however exclude the possibility that a small portion of fibres may be present that are not

colocalized with tdTomato, which may reflect unspecific expression of ChR2 in nerve

terminals other than Ucn3 in the PVN. The optic fibre cannula was also verified to be correctly

positioned in the PVN in all of 8 mice.

Discussion

We show optogenetic stimulation of MePD Ucn3 neurons inhibits pulsatile LH secretion, an

effect that was completely blocked by intra-MePD GABA and glutamate antagonism.

Moreover, optogenetic stimulation of MePD Ucn3 projections in the hypothalamic PVN also

suppresses LH pulsatility while inducing the release of CORT. These results demonstrate that

inhibition of the GnRH pulse generator by Ucn3 neurons within the MePD relies on GABA

and glutamate signalling in the MePD, *per se*. Moreover, MePD Ucn3 efferent projections in

the PVN modulate the activity of the HPA and HPG axes.

The MePD is a major upstream regulator of gonadotropic hormone secretion with a significant

population of GABAergic neurons projecting to reproductive neural centres in the

hypothalamus (26). Although classic lesioning studies suggest an inhibitory output of the

MePD over reproductive function (27,28), selective optical stimulation of MePD kiss1 neurons

increases LH pulse frequency (4), which our unpublished observations show relies upon both

GABA and glutamate signalling in the MePD. Due to this interesting dichotomy, we have

proposed a disinhibitory system, which is consistent with the pallidal origin of this limbic

structure (29), whereby MePD kiss1 activation excites inhibitory GABAergic interneurons,

which in turn reduces the inhibitory tone of the GABAergic projections from the MePD to the

ARC KNDy network allowing for an increase in GnRH pulse generator frequency (4).

Contrastingly, kiss1R antagonism, reducing MePD kiss1 signalling decreases GnRH pulse

generator frequency (30).

We have recently shown that stress-induced activation of Ucn3 and CRFR2 signalling within

the MePD inhibits GnRH pulse generator activity (3) and activation of MePD Ucn3 neurons

delays pubertal timing (31). It has also been shown that the majority of MePD CRFR2

expressing neurons are GABAergic (21). In the MePD, Ucn3 fibres overlap with sites

expressing its cognate receptor CRFR2 (32,33) and some local Ucn3 neurons within the MePD

are shown to have an interneuron-like appearance (34), thus Ucn3 neurons may form

connections with and signal via GABAergic CRFR2-positive neurons in this region. In the

present study, we show that optogenetic activation of MePD Ucn3 neurons inhibits pulsatile

LH secretion. Therefore, we propose that activated MePD Ucn3 neurons may signal through

inhibitory GABAergic CRFR2-positive interneurons, upstream to the MePD kiss1 neurons,

downregulating MePD kiss1 signalling resulting in decreased GnRH pulse generator

frequency, as illustrated in figure 6A.

In our proposed model, we hypothesised that GABA signalling is an important factor mediating

the suppressive effect of MePD Ucn3 activation on GnRH pulse generator activity. Therefore,

we tested the effect of intra-MePD administration of $GABA_A$ or $GABA_B$ receptor antagonists

during optical stimulation of MePD Ucn3 neurons on LH pulsatility, and indeed, we found that

both $GABA_A$ and $GABA_B$ receptor antagonism blocks the suppression of LH pulses by MePD

Ucn3 neuronal activation. Indeed, it has been shown that in $GABA_B$ receptor knock out mice

Kiss1 mRNA expression increases in the MePD suggesting $GABA_B$ regulates Kiss1 expression

in this region (35). Whether the MePD kiss1 neurons also contain $GABA_A$ receptors is

unknown. Selective optical stimulation of Ucn3 neurons may activate inhibitory GABAergic

CRFR2-positive interneurons (35), which in turn downregulate MePD kiss1 signalling, which

we have previously shown results in suppression of GnRH pulse generator frequency (30) (see

Fig. 6A). Although, based on our proposed neuronal circuit, GABA$_A$ and GABA$_B$ receptor

antagonism may be working upstream and downstream of MePD kiss1 activity to cancel both

the inhibitory GABAergic CRFR2 interneurons upstream to and the inhibitory GABAergic

interneurons downstream to MePD kiss1 neurons. Therefore, the inhibitory output of the

MePD GABAergic projection neurons to the ARC KNDy network is cancelled, thus, negating

the suppressive effect of MePD Ucn3 activation on the GnRH pulse generator. Essentially,

terminating the ability of the GABA interneuron, upstream and downstream of MePD kiss1

activity, to take part in the disinhibition by antagonising GABA receptors located on the MePD

kiss1 neurons as well as post-synaptic GABA projection neurons removing the inhibitory

control over the GnRH pulse generator from the MePD (see Fig. 6B).

We have identified GABA signalling as a key part of the MePD Ucn3 neural circuitry

associated with suppression of the GnRH pulse generator, however glutamate signalling in the

amygdala is also known to be involved in modulating reproduction and anxiety-like behaviour

(36). Administration of NMDA into the MePD disrupts ovarian cycles in rats, while NMDA

receptor antagonism in the MePD delays pubertal timing (28). We hypothesised that glutamate

neurotransmission may form a critical part of the MePD Ucn3 neural circuitry involved in

stress-induced inhibition of GnRH pulse generator activity. In the present study, we show that

intra-MePD AMPA and NMDA receptor antagonism blocks the inhibitory effect of MePD

Ucn3 neuronal activation on LH pulsatility. These observations suggest the involvement of

similar neural circuitry as in the neighbouring posteroventral medial amygdala which exhibits

functional glutamatergic signalling onto inhibitory GABA neurons (29). Moreover, inhibition

of both GABA and glutamate signalling is equally effective at blocking suppression of LH

pulses suggesting they are key components in the circuit.

We know the MePD Ucn3 neurons are interconnected with each other as well as CRFR2

expressing neurons, forming a Ucn3-CRFR2 neuronal circuit (21). The effect of glutamate

antagonism illustrated in figure 6C, suggests that glutamate transmission may be mediated via

glutamatergic Ucn3 neurons. This has been shown in the hypothalamic perifornical area,

where they are activated during infant-directed aggression (37) and increase anxiety-like

behaviour (38), but this awaits future study for the MePD Ucn3 population. Our previous data

shows that infusing Ucn3 in the MePD blocks LH pulses indicating their activity in this brain

region is sufficient to suppress GnRH pulse generator activity (3). We have also shown that

infusion of CRFR2 antagonists in the MePD blocks the stress-induced suppression of LH

pulses suggesting MePD CRFR2 expressing neurons are also involved in modulating the

GnRH pulse generator under stressful conditions. Our current data expands on our previous

observations determining whether neurotransmission modulating pulsatile LH secretion is

involved in this MePD circuit. Our results suggest that glutamate may be necessary for the

inhibition of LH pulses induced by activation of MePD Ucn3 neurons, thus we show that

glutamate signalling is an indispensable part of this circuit. Based on our observations, we

propose that Ucn3 neurons in the MePD may be glutamatergic similarly to the Ucn3 neurons

in the hypothalamic perifornical area. Glutamate is possibly required to trigger network

activity of the MePD Ucn3 neurons similar to the role glutamate plays in the ARC KNDy

network (14). This allows the network to reach a certain level of excitability that promotes

signalling via Ucn3 to activate CRFR2 expressing GABAergic neurons, which in turn

downregulate MePD kiss1 activity, resulting in suppression of GnRH pulse generator activity.

Thus, NMDA and AMPA receptor antagonism possibly cancels MePD Ucn3 glutamatergic

activation of inhibitory GABAergic CRFR2 expressing inter-neurons preventing the decrease

in GnRH pulse generator activity (Fig. 6C). However, fast synaptic activity occurs due to

synaptic neurotransmitter release and modulation of this activity is known to be a key target

for neuropeptides. Indeed, urocortins can also function endogenously as modulators of

excitatory glutamatergic transmission in the amygdala, with *in vitro* studies showing

modulation of glutamatergic excitatory post-synaptic potentials via pre- and postsynaptic

CRFR2-mediated mechanisms (39). Therefore, we cannot exclude the possibility that Ucn3

activity in the MePD is augmenting glutamatergic transmission to activate downstream

inhibitory GABA neurons.

The fact that GABAergic and glutamatergic receptor antagonism within the MePD did not alter

LH pulsatility without optogenetic stimulation of Ucn3 neurons is an important factor in our

proposed mechanisms and neural circuits within the MePD. This observation indicates that

under basal non-stress conditions MePD Ucn3 neurons are relatively quiescent, thus solely

pharmacologically blocking the upstream GABA and glutamate inputs to the GABAergic

MePD projections without a corresponding increase in Ucn3 activity would make little

difference to the net effect of the MePD over the KNDy neuronal system.

Stress elevates Ucn3 mRNA (19) and c-fos expression in the MePD (18) while chemogenetic

inhibition of MePD Ucn3 neurons prevents stress-induced suppression of LH pulsatility (3).

The MePD sends stress-activated efferents to the PVN (18) and MePD Ucn3 neurons have

been shown to project directly to the PVN (21). In the current study, we demonstrate for the

first time that optical stimulation of MePD Ucn3 projections in the PVN inhibits pulsatile LH

secretion. Recently, chemogenetic activation of PVN CRF neuron perikarya was shown to

rapidly suppress LH pulse frequency, however optogenetic activation of their nerve terminals

in the ARC does not affect the firing of KNDy neurons (40) and directly injecting CRF in the

ARC also has no effect on LH secretion (41), suggesting the involvement of an indirect

mechanism inhibiting GnRH pulse generator activity. The MePD has been shown to send

Ucn3 neuronal projections to the PVN and receive Ucn3 projections from the PVN, indicating

bidirectional communication between these two brain regions via Ucn3 signalling (21).

Moreover, the MePD sends direct efferent projections to KNDy neurons of an unknown

neurochemical phenotype (8). Therefore, stressful stimuli may be processed by PVN CRF or

Ucn3 neurons and this information may be relayed via the MePD Ucn3 neuronal circuitry to

the ARC KNDy neurons to modulate their activity (see Fig. 6D). Ultimately, the MePD may

function as a novel relay centre between the PVN and ARC to regulate stress signals to the

GnRH pulse generator and contribute to the crosstalk between the reproductive and stress axes.

Central administration of Ucn3 augments the HPA axis response to restraint stress in rats (19)

and increases hypothalamic CRF concentrations and circulating levels of CORT in mice (42).

Overexpression of Ucn3 in the hypothalamic perifornical area increases anxiety-like behaviour

(38) and activation of the CRFR2 in the lateral septum is involved in stress-induced persistent

anxiety in mice (43). In the medial amygdala, Ucn3 neuronal populations are shown to consist

of both projection neurons extending to the PVN, the bed nucleus of the stria terminalis and

the suprachiasmatic nucleus (21,34) and local Ucn3 interneurons (32,33). We have previously

shown chemogenetic inhibition targeting both local and efferent MePD Ucn3 neurons prevents

psychosocial stress-induced release of CORT (3). In the present study, we show that selective

optogenetic activation of MePD Ucn3 projections in the PVN elevates CORT secretion, thus

MePD Ucn3 projections to the PVN are critical for the regulation of CORT secretion. Further

work is required to identify whether only the MePD Ucn3 projecting efferents to the PVN are

crucial for regulating CORT secretion. The presence of E_2 is known to up-regulate basal CRF

mRNA levels in the PVN in rodents and induce high levels of circulating corticosterone

providing a sensitizing effect to the activity of the HPA axis (44,45). Moreover, the CRFR2

promoter contains a classical estrogen receptor response element (46), indicating that CRFR2

activity is upregulated in the presence of E_2. Therefore, we implanted the OVX mice with E_2

capsules for HPA axis sensitization in mice similarly to what has been reported in previous

studies (44,47–49). However, E_2 levels may have fallen lower than expected, to low or

potentially below the physiological range observed in diestrous mice, as evidenced by little or

no effect on LH pulse frequency. However, an effect on LH pulse amplitude and mean LH

levels was evident in the E_2 replaced animals (25). Although there was weak E_2 negative

feedback on LH pulses in our model, our main reason for E_2 replacement was to sensitizes the

HPA axis (44,47–49). PVN-CRF neurons express mRNA transcripts for CRFR2 (50) and

CRFR2 primarily activate the Gαs adenylyl cyclase/cAMP signalling pathway (51) indicating

that signalling via the CRFR2 located on PVN-CRF neurons may result in activation of this

neuronal population and hence the HPA axis (see Fig. 6D). Therefore, Ucn3 released following

optical stimulation of MePD Ucn3 projections in the PVN may directly activate the PVN-CRF

neurons, signalling via CRFR2, to stimulate CORT release as observed in the present study.

However, this hypothesis requires further experimental interrogation.

Selective CRFR2 antagonism has been shown to reverse the stress-induced inhibition of LH

pulses (52) where central administration of astressin2-B, a specific CRFR2 antagonist, blocked

the restraint-stress induced suppression of pulsatile LH secretion in rats (53). We have shown

specific antagonism of CRFR2 in the MePD blocked the inhibitory effects of predator odor on

the GnRH pulse generator in mice (3) and activation of MePD Ucn3 neurons delayed puberty

in mice (31). Therefore, we know CRFR2 neurons are involved in processing stressful

information and coordinating the activity of the GnRH pulse generator accordingly.

Additionally, CRF is also implicated in reducing pulsatile LH secretion via CRFR1 expressing

neurons in mice (54). Chronic stress levels of CORT decrease LH pulsatility (55) and recently

an acute rise in CORT sustained for 30-60 min was shown to inhibit pulsatile LH release, but

only in OVX mice with estrogen replacement (56) and after a delay of approximately 30 min

following the CORT rise. From our observations, inhibition of LH pulsatility occurs

immediately after the stimulation of MePD Ucn3 projections in the PVN in OVX E_2 replaced

mice, thus it is unlikely the result of negative feedback from CORT release but rather a central

suppression of GnRH pulse generator activity possibly from the activation of PVN CRF

neurons, although the mechanism involved has yet to be established. Moreover, acute

psychosocial stress exposure reduces LH pulse frequency in OVX mice (3,57), suggesting E_2

is not necessary for the inhibition of the GnRH pulse generator in response to psychosocial

stress (3,57). Therefore, acute stress pathways possibly act via CORT and E_2 independent

pathways to suppress LH pulses. In our study, optical stimulation of MePD Ucn3 local and

efferent neurons is supposed to mimic acute exposure to psychosocial stress as this has been

shown to be processed by Ucn3 activity in the MePD, thus supporting the idea of a central

suppression of GnRH pulse generator activity mediated by MePD Ucn3 neurons.

The PVN CRF neuronal population was not traditionally considered important for mediating

stress-induced suppression of pulsatile LH secretion, as PVN lesion studies have shown that

this region is not critical in mediating the stress-induced suppression of the reproductive axis

(58). However, with the recent observations showing that chemogenetic activation of PVN

CRF neurons indirectly inhibits the GnRH pulse generator it is plausible that the stress neural

circuitry includes upstream signalling via the MePD. Therefore, a possible route may involve

stressful information processed by the PVN CRF neurons relayed via the MePD Ucn3 neuronal

circuitry to the ARC KNDy neurons to modulate their activity (see Fig. 6D). The neighbouring

central nucleus of the amygdala is known to act as a relay centre in pain control where the

central lateral part receives nociceptive input from the parabrachial nucleus and the central

medial part is involved in regulating the output of defensive behaviour and nociception (59).

Local and efferent MePD Ucn3 signalling plays a critical dual role in regulating GnRH pulse

generator activity and CORT secretion, thus MePD Ucn3 neuronal populations and their

projections may represent a novel nodal centre in the interaction between the reproductive and

stress axes.

Our findings show for the first time that stimulation of local Ucn3 neurons in the MePD inhibits

GnRH pulse generator activity involving an interplay between Ucn3, GABA and glutamate

signalling within the MePD. Furthermore, we interrogate the MePD Ucn3 neuron projection

terminals in the PVN demonstrating that stimulation of the MePD Ucn3 efferent projections in

the PVN elevates CORT secretion to stress levels while suppressing the GnRH pulse generator.

How the local neural circuit within the MePD engages with efferent projections from the MePD

to the PVN and the interplay between Ucn3, GABA and glutamate neurons to mediate the

suppressive effect on LH pulsatility and the stimulatory effect on the HPA axis remains an

exciting question for the future.

Bibliography

1.   Ressler KJ. Amygdala Activity, Fear, and Anxiety: Modulation by Stress. Biol

Psychiatry. 2010;67(12):1117–9.

2.   Lin Y, Li XF, Lupi M, Kinsey-Jones JS, Shao B, Lightman SL, O'Byrne KT . The role

of the medial and central amygdala in stress-induced suppression of pulsatile LH

secretion in female rats. Endocrinology. 2011;152(2):545–55.

3.   Ivanova D, Li XF, McIntyre C, Liu Y, Kong L, O'Byrne KT . Urocortin3 in the

Posterodorsal Medial Amygdala Mediates Stress-induced Suppression of LH

Pulsatility in Female Mice. Endocrinology. 2021;162(12):bqab206.

4.   Lass G, Li XF, de Burgh RA, He W, Kang Y, Hwa-Yeo S, Sinnett-Smith LC,

Manchishi SM, Colledge WH, Lightman SL, O'Byrne KT . Optogenetic stimulation of

kisspeptin neurones within the posterodorsal medial amygdala increases luteinising

hormone pulse frequency in female mice. J Neuroendocrinol. 2020;32(2):e12823.

5.   Li XF, Adekunbi DA, Alobaid HM, Li S, Pilot M, Lightman SL, O'Byrne KT . Role

of the posterodorsal medial amygdala in predator odour stress-induced puberty delay

in female rats. J Neuroendocrinol. 2019;31(6):e12719.

6.   Adekunbi DA, Li XF, Li S, Adegoke OA, Iranloye BO, Morakinyo AO, Lightman SL,

Taylor PD, Poston L, O'Byrne KT . Role of amygdala kisspeptin in pubertal timing in

female rats. PLoS One. 2017;12(8):e0183596.

7.   Moore AM, Coolen LM, Lehman MN. Kisspeptin/Neurokinin B/Dynorphin (KNDy)

cells as integrators of diverse internal and external cues: evidence from viral-based

monosynaptic tract-tracing in mice. Sci Rep. 2019;9(1):1–5.

8.   Yeo SH, Kyle V, Blouet C, Jones S, Colledge WH. Mapping neuronal inputs to Kiss1

neurons in the arcuate nucleus of the mouse. PLoS One. 2019;14(3):e0213927.

9.   Qiu J, Nestor CC, Zhang C, Padilla SL, Palmiter RD, Kelly MJ, Rønnekleiv OK .

High-frequency stimulation-induced peptide release synchronizes arcuate kisspeptin

neurons and excites GnRH neurons. Elife. 2016;5:e16246.

10.   Clarkson J, Han SY, Piet R, McLennan T, Kane GM, Ng J, Porteous RW, Kim JS,

Colledge WH, Iremonger KJ, Herbison AE . Definition of the hypothalamic GnRH

pulse generator in mice. Proc Natl Acad Sci U S A. 2017;114(47):e10216–23.

11.   Voliotis M, Li XF, De Burgh R, Lass G, Lightman SL, O'Byrne KT, Tsaneva-

Atanasova K . The Origin of GnRH pulse generation: An integrative mathematical-

experimental approach. J Neurosci. 2019;39(49):9738–47.

12.   Moore AM, Coolen LM, Lehman MN. In vivo imaging of the GnRH pulse generator

reveals a temporal order of neuronal activation and synchronization during each pulse.

Proc Natl Acad Sci U S A. 2022;119(6).

13.   Han SY, McLennan T, Czieselsky K, Herbison AE. Selective optogenetic activation of

arcuate kisspeptin neurons generates pulsatile luteinizing hormone secretion. Proc Natl

Acad Sci U S A. 2015;112(42):13109–14.

545 14. Voliotis M, Li XF, De Burgh RA, Lass G, Ivanova D, McIntyre C, O'Byrne K,

546   Tsaneva-Atanasova K . Modulation of pulsatile GnRH dynamics across the ovarian

547   cycle via changes in the network excitability and basal activity of the arcuate

548   kisspeptin network. Elife. 2021;10.

549 15. Martin EI, Ressler KJ, Binder E, Nemeroff CB. The Neurobiology of Anxiety

550   Disorders: Brain Imaging, Genetics, and Psychoneuroendocrinology. Psychiatr Clin

551   North Am. 2009;32(3):549.

552 16. Nuss P. Anxiety disorders and GABA neurotransmission: A disturbance of

553   modulation. Neuropsychiatr Dis Treat. 2015;11:165–75.

554 17. Takahashi LK. Olfactory systems and neural circuits that modulate predator odor fear.

555   Front Behav Neurosci. 2014;8:72.

556 18. Van-Hover C, Li C. Stress-activated afferent inputs into the anterior parvicellular part

557   of the paraventricular nucleus of the hypothalamus: Insights into urocortin 3 neuron

558   activation. Brain Res. 2015;1611:29–43.

559 19. Jamieson PM, Li C, Kukura C, Vaughan J, Vale W. Urocortin 3 modulates the

560   neuroendocrine stress response and is regulated in rat amygdala and hypothalamus by

561   stress and glucocorticoids. Endocrinology. 2006;147(10):4578–88.

562 20. Fekete ÉM, Zhao Y, Li C, Sabino V, Vale WW, Zorrilla EP. Social defeat stress

563   activates medial amygdala cells that express type 2 corticotropin-releasing factor

564   receptor mRNA. Neuroscience. 2009;162(1):5–13.

565 21. Shemesh Y, Forkosh O, Mahn M, Anpilov S, Sztainberg Y, Manashirov S,

566   Shlapobersky T, Elliott E, Tabouy L, Ezra G, Adler ES . Ucn3 and CRF-R2 in the

567   medial amygdala regulate complex social dynamics. Nat Neurosci. 2016;19(11):1489–

568   96.

22. Ingberg E, Theodorsson A, Theodorsson E, Strom JO . Methods for long-term 17β-

estradiol administration to mice. Gen Comp Endocrinol. 2012;175(1):188–93.

23. Paxinos G, Franklin KB . Mouse Brain in Stereotaxic Coordinates. Acad Press.

2019;5:246.

24. McCosh RB, Kreisman MJ, Breen KM. Frequent tail-tip blood sampling in mice for

the assessment of pulsatile luteinizing hormone secretion. J Vis Exp. 2018;(137).

25. Ivanova D. Supplementary Data. OSF. 2022;September. osf.io/vem8y

26. Pardo-Bellver C, Cádiz-Moretti B, Novejarque A, Martínez-García F, Lanuza E.

Differential efferent projections of the anterior, posteroventral, and posterodorsal

subdivisions of the medial amygdala in mice. Front Neuroanat. 2012;6(33):1–26.

27. Stephens SBZ, Raper J, Bachevalier J, Wallen K. Neonatal amygdala lesions advance

pubertal timing in female rhesus macaques. Psychoneuroendocrinology. 2015;51:307–

17.

28. Li XF, Hu MH, Hanley BP, Lin YS, Poston L, Lightman SL, O'Byrne KT . The

posterodorsal medial amygdala regulates the timing of puberty onset in female rats.

Endocrinology. 2015;156(10):3725–36.

29. Keshavarzi S, Sullivan RKP, Ianno DJ, Sah P. Functional properties and projections of

neurons in the medial amygdala. J Neurosci. 2014;34(26):8699–715.

30. Comninos AN, Anastasovska J, Sahuri-Arisoylu M, Li XF, Li S, Hu M, Jayasena CN,

Ghatei MA, Bloom SR, Matthews PM, O'Byrne KT . Kisspeptin signaling in the

amygdala modulates reproductive hormone secretion. Brain Struct Funct.

2016;221(4):2035–47.

31. Ivanova D, Li X, Liu Y, McIntyre C, Fernandes C, Lass G, Kong L, O'Byrne KT .

Role of Posterodorsal Medial Amygdala Urocortin-3 in Pubertal Timing in Female

Mice. Front Endocrinol (Lausanne). 2022;0:910.

32.   Cavalcante JC, Sita LV, Mascaro MB, Bittencourt JC, Elias CF. Distribution of

urocortin 3 neurons innervating the ventral premammillary nucleus in the rat brain.

Brain Res. 2006;1089(1):116–25.

33.   Li C, Vaughan J, Sawchenko PE, Vale WW. Urocortin III-immunoreactive projections

in rat brain: Partial overlap with sites of type 2 corticotrophin-releasing factor receptor

expression. J Neurosci. 2002;22(3):991–1001.

34.   Deussing JM, Breu J, Kühne C, Kallnik M, Bunck M, Glasl L, Yen YC, Schmidt MV,

Zurmühlen R, Vogl AM, Gailus-Durner V . Urocortin 3 modulates social

discrimination abilities via corticotropin-releasing hormone receptor type 2. J

Neurosci. 2010;30(27):9103–16.

35.   Di Giorgio NP, Semaan SJ, Kim J, López PV, Bettler B, Libertun C, Lux-Lantos VA,

Kauffman AS . Impaired GABAB receptor signaling dramatically up-regulates Kiss1

expression selectively in nonhypothalamic brain regions of adult but not prepubertal

mice. Endocrinology. 2014;155(3):1033–44.

36.   Hegoburu C, Parrot S, Ferreira G, Mouly AM. Differential involvement of amygdala

and cortical NMDA receptors activation upon encoding in odor fear memory. Learn

Mem. 2014;21(12):651–5.

37.   Autry AE, Wu Z, Kapoor V, Kohl J, Bambah-Mukku D, Rubinstein ND, Marin-

Rodriguez B, Carta I, Sedwick V, Tang M, Dulac C . Urocortin-3 neurons in the

mouse perifornical area promote infant-directed neglect and aggression. Elife.

2021;10.

38.   Kuperman Y, Issler O, Regev L, Musseri I, Navon I, Neufeld-Cohen A, Gil S, Chen A

. Perifornical urocortin-3 mediates the link between stress-induced anxiety and energy

homeostasis. Proc Natl Acad Sci U S A. 2010;107(18):8393–8.

39.   Liu J, Yu B, Neugebauer V, Grigoriadis DE, Rivier J, Vale WW, Shinnick-Gallagher

619   P, Gallagher JP . Corticotropin-releasing factor and Urocortin I modulate excitatory

620   glutamatergic synaptic transmission. J Neurosci. 2004;24(16):4020–9.

621   40.   Yip SH, Liu X, Hessler S, Cheong I, Porteous R, Herbison AE. Indirect Suppression of

622   Pulsatile LH Secretion by CRH Neurons in the Female Mouse. Endocrinology.

623   2021;162(3).

624   41.   Rivest S, Plotsky PM, Rivier C. CRF alters the infundibular LHRH secretory system

625   from the medial preoptic area of female rats: possible involvement of opioid receptors.

626   Neuroendocrinology. 1993;57(2):236–46.

627   42.   Bagosi Z, Csabafi K, Karasz G, Jászberényi M, Földesi I, Siska A, Szabó G, Telegdy

628   G . The effects of the urocortins on the hypothalamic-pituitary-adrenal axis -

629   similarities and discordancies between rats and mice. Peptides. 2019;112:1–13.

630   43.   Anthony TE, Dee N, Bernard A, Lerchner W, Heintz N, Anderson DJ. Control of

631   stress-induced persistent anxiety by an extra-amygdala septohypothalamic circuit.

632   Cell. 2014;156(3):522–36.

633   44.   Li XF, Mitchell JC, Wood S, Coen CW, Lightman SL, O'Byrne KT . The effect of

634   oestradiol and progesterone on hypoglycaemic stress-induced suppression of pulsatile

635   luteinizing hormone release and on corticotropin-releasing hormone mRNA expression

636   in the rat. J Neuroendocrinol. 2003;15(5):468–76.

637   45.   Zavala E, Voliotis M, Zerenner T, Tabak J, Walker JJ, Li XF, Terry JR, Lightman SL,

638   O'byrne K, Tsaneva-Atanasova K . Dynamic Hormone Control of Stress and Fertility.

639   Front Physiol. 2020;11:1457.

640   46.   Catalano RD, Kyriakou T, Chen J, Easton A, Hillhouse EW. Regulation of

641   Corticotropin-Releasing Hormone Type 2 Receptors by Multiple Promoters and

642   Alternative Splicing: Identification of Multiple Splice Variants. Mol Endocrinol.

643   2003;17(3):395–410.

644 47. Cagampang FR, Cates PS, Sandhu S, Strutton PH, McGarvey C, Coen CW, O'Byrne

645   KT . Hypoglycaemia-induced inhibition of pulsatile luteinizing hormone secretion in

646   female rats: role of oestradiol, endogenous opioids and the adrenal medulla. J

647   Neuroendocrinol. 1997;9(11):867–72.

648 48. Chen MD, O'Byrne KT, Chiappini SE, Hotchkiss J, Knobil E . Hypoglycemic "stress"

649   and gonadotropin-releasing hormone pulse generator activity in the rhesus monkey:

650   role of the ovary. Neuroendocrinology. 1992;56(5):666–73.

651 49. McCosh RB, Breen KM, Kauffman AS . Neural and Endocrine Mechanisms

652   Underlying Stress-Induced Suppression of Pulsatile LH Secretion. Mol Cell

653   Endocrinol. 2019;498:110579.

654 50. Dabrowska J, Hazra R, Guo JD, DeWitt S, Rainnie DG. Central CRF neurons are not

655   created equal: Phenotypic differences in CRF-containing neurons of the rat

656   paraventricular hypothalamus and the bed nucleus of the stria terminalis. Front

657   Neurosci. 2013;0(7 AUG):156.

658 51. Deussing JM, Chen A. The corticotropin-releasing factor family: Physiology of the

659   stress response. Vol. 98, Physiological Reviews. 2018. p. 2225–86.

660 52. Li XF, Bowe JE, Kinsey-Jones JS, Brain SD, Lightman SL, O'Byrne KT . Differential

661   role of corticotropin-releasing factor receptor types 1 and 2 in stress-induced

662   supression of pulsatile luteinising hormone secretion in the female rat. J

663   Neuroendocrinol. 2006;18(8):602–10.

664 53. Li XF, Bowe JE, Lightman SL, O'Byrne KT . Role of corticotropin-releasing factor

receptor-2 in stress-induced suppression of pulsatile luteinizing hormone secretion in

the rat. Endocrinology. 2005;146(1):318–22.

54. Raftogianni A, Roth LC, García-González D, Bus T, Kühne C, Monyer H, Spergel DJ,

Deussing JM, Grinevich V . Deciphering the Contributions of CRH Receptors in the

Brain and Pituitary to Stress-Induced Inhibition of the Reproductive Axis. Front Mol

Neurosci. 2018;11:305.

55. Kreisman MJ, McCosh RB, Tian K, Song CI, Breen KM. Estradiol Enables Chronic

Corticosterone to Inhibit Pulsatile Luteinizing Hormone Secretion and Suppress Kiss1

Neuronal Activation in Female Mice. Neuroendocrinology. 2020;110(6):501–16.

56. Kreisman MJ, McCosh RB, Breen KM . Role of CORT Duration and Estradiol

Dependence for Stress-level of CORT to Inhibit Pulsatile LH Secretion in Female

Mice. J Endocr Soc. 2021;Apr;5(Supplement_1):A552-.

57. Yang JA, Song CI, Hughes JK, Kreisman MJ, Parra RA, Haisenleder DJ, Kauffman

AS, Breen KM . Acute psychosocial stress inhibits LH pulsatility and kiss1 neuronal

activation in female mice. Endocrinology. 2017;158(11):3716–23.

58. Serge S, Rivier C. Influence of the paraventricular nucleus of the hypothalamus in the

alteration of neuroendocrine functions induced by intermittent footshock or

interleukin. Endocrinology. 1991;129(4):2049–58.

59. Wank I, Pliota P, Badurek S, Kraitsy K, Kaczanowska J, Griessner J, Kreitz S, Hess A,

Haubensak W . Central amygdala circuitry modulates nociceptive processing through

differential hierarchical interaction with affective network dynamics. Commun Biol.

2021;4(1):1–10.

Figure 1. Optogenetic stimulation of posterodorsal medial amygdala (MePD) urocortin-3

(Ucn3) neurons with blue light at 10 Hz, 10 mW suppressed pulsatile LH secretion in adult

ovariectomised (OVX) Ucn3-cre-tdTomato female mice. Infusion of bicuculline (BIC),

$GABA_A$ receptor antagonist, CGP, $GABA_B$ receptor antagonist, and AP5 and CNQX, NMDA

and AMPA receptor antagonists, respectively, into the posterodorsal medial amygdala (MePD)

during optogenetic stimulation of MePD urocortin-3 (Ucn3) neurons with blue light at 10 Hz,

10 mW completely blocked the suppressive effect of MePD Ucn3 neuronal activation on LH

pulsatility. Representative LH pulse profiles showing the effects of (A) control AAV-YFP

mice administered with aCSF receiving optical stimulation of 10 Hz, 10 mW, (B) AAV-ChR2

mice administered with aCSF receiving optical stimulation of 10 Hz, 10 mW, (C) Summary of

LH pulse interval for the pre-stimulation control period (1 h) and optical stimulation/infusion

period (1 h), (D) AAV-ChR2 mice administered with BIC, (E) AAV-ChR2 mice administered

with BIC receiving optical stimulation of 10 Hz, 10 mW, (F) Summary of LH pulse interval,

(G) AAV-ChR2 mice administered with CGP, (H) AAV-ChR2 mice administered with CGP

receiving optical stimulation of 10 Hz, 10 mW, (I) Summary of LH pulse interval, (J) AAV-

ChR2 mice administered with a combination of AP5 and CNQX, (K) AAV-ChR2 mice

administered with a combination of AP5 and CNQX receiving optical stimulation of 10 Hz, 10

mW, (L) Summary of LH pulse interval. LH pulses detected by the DynePeak algorithm are

indicated with an asterisk located above each pulse on the representative LH pulse profiles. For

A-C +++$p < 0.0001$ control period vs optic stimulation + aCSF (AAV-ChR2: $n = 9$); ##$p < 0.001$

optic stimulation vs control group (AAV-YFP: $n = 3$; no stimulation: $n = 2$). For D-F $n = 8$-9

per group. For G-I $n = 7$ per group. For J-L $n = 6$ per group.

Figure 2. Optogenetic stimulation of posterodorsal medial amygdala (MePD) urocortin-3

(Ucn3) neuronal terminals in the hypothalamic paraventricular nucleus (PVN) with blue light

at 20 Hz, 15 mW elevated corticosterone (CORT) secretion in adult ovariectomised (OVX) +

17ß-estradiol (E_2) Ucn3-cre-tdTomato female mice. CORT secretion time-course for mice

receiving optical stimulation initiated at 0 min and terminated at 60 min (1 h duration),

followed by a 1 h recovery period (60-120 min) for the AAV-ChR2 group (blue line, circles)

and control group (red line, squares). +p<0.05 AAV-ChR2 (n = 6) at time point 30 min and

60 min vs. control (AAV-YFP: n = 5) at time point 30 min and 60 min.

Figure 3. Optogenetic stimulation of posterodorsal medial amygdala (MePD) urocortin-3

(Ucn3) neuronal terminals in the hypothalamic paraventricular nucleus (PVN) with blue light

at 20 Hz, 15 mW reduced LH pulsatility in adult ovariectomised (OVX) + 17ß-estradiol (E_2)

Ucn3-cre-tdTomato female mice. Representative LH pulse profiles showing the effects of (A)

AAV-YFP/WT mice receiving optical stimulation of 20 Hz, 5 mW, (B) AAV-ChR2 mice

receiving optical stimulation of 20 Hz, 5 mW, (C) Another example of AAV-YFP/WT mice

receiving optical stimulation of 20 Hz, 5 mW, (D) Another example of AAV-ChR2 mice

receiving optical stimulation of 20 Hz, 5 mW, (E) Summary of LH pulse interval for the pre-

stimulation control period (1 h) and optic stimulation period (1 h). LH pulses detected by the

DynePeak algorithm are indicated with an asterisk located above each pulse on the

representative LH pulse profiles. +++p<0.0001 control period vs optic stimulation for AAV-

ChR2 mice (n = 8); ###p<0.0001 AAV-YFP/WT (AAV-YFP, n = 3; wild-type with AAV-

ChR2, n = 2; no stimulation, n = 2) vs AAV-ChR2 mice during optic stimulation.

Figure 4. Expression of AAV9-EF1a-double floxed-hChR2(H134R)-EYFP-WPRE-HGHpA

in posterodorsal medial amygdala (MePD) Urocortin-3 (Ucn3) neurons. (A-E) Representative

dual fluorescence photomicrographs of the MePD from a Ucn3-cre-tdTomato female

ovariectomised (OVX) mouse injected with AAV9-EF1a-double floxed-hChR2(H134R)-

EYFP-WPRE-HGHpA. Ucn3 neurons labelled with tdTomato (A) and YFP (C) appear

yellow/orange (E). (B), (D) and (F) are a higher power view of (A), (C) and (E) respectively.

Scale bars represent A, C, E 50 μm and B, D, F 20 μm. White arrows show examples of Ucn3

neurons labelled with tdTomato and YFP. OT, Optic tract (blue line).

Figure 5. Expression of AAV9-EF1a-double floxed-hChR2(H134R)-EYFP-WPRE-HGHpA

in Urocortin-3 (Ucn3) neuronal projection in the paraventricular nucleus of the hypothalamus

(PVN) originating from the posterodorsal medial amygdala (MePD). (A-Aiii) Representative

dual fluorescence photomicrographs of the PVN from a Ucn3-cre-tdTomato female

ovariectomised (OVX) + 17ß-estradiol (E$_2$) mouse injected with AAV9-EF1a-double floxed-

hChR2(H134R)-EYFP-WPRE-HGHpA. Ucn3 neurons and projections labelled with (A) YFP

and tdTomato expression in PVN, (Ai) higher power view of YFP labelled MePD Ucn3

projections in the PVN, (Aii) higher power view of tdTomato labelled MePD Ucn3 projections

in the PVN and (Aiii) YFP and tdTomato merged. Scale bars represent A, 100 μm, Ai, Aii and

Aiii, 10 μm. Blue box shows the YFP labelled MePD Ucn3 projections in the PVN, yellow

arrows point to parts of the YFP labelled MePD Ucn3 projection in the PVN. 3V; third

ventricle.

Figure 6. Schematic illustration of the MePD neurocircuits and their pharmacological

manipulation. (A) Optical stimulation of local MePD Ucn3 neurons releases Ucn3 and the co-

transmitter glutamate, which activate inhibitory GABAergic CRFR2-positive interneurons

located upstream to the kiss1 neurons suppressing their activity. The reduced kiss1 signalling

deactivates its downstream inhibitory GABAergic interneuron causing disinhibition of the

GABAergic projection efferents from the MePD, resulting in an increased inhibitory tone over

the ARC KNDy network, thus decreasing GnRH pulse generator frequency. (B) Antagonism

of both GABA$_A$ and GABA$_B$ receptors in the MePD during optical stimulation of Ucn3 neurons

blocks the downregulation of kiss1 neuronal activity, preventing the inhibitory effect of Ucn3

signalling on the KNDy network. (C) Antagonism of both NMDA and AMPA receptors in

the MePD during optical stimulation of Ucn3 neurons blocks the activation of the downstream

GABAergic CRFR2-positive interneurons, preventing the inhibitory effect of Ucn3 signalling

on the KNDy network and suggesting glutamate is necessary for the inhibition of pulsatile LH

secretion. (D) High frequency photoactivation of MePD Ucn3 projection terminals in the PVN

may activate Ucn3 neurons and/or other neurons of an unknown phenotype in the PVN

resulting in stimulation of the HPA axis and CORT release. Additionally, there may be

collateral activation of reciprocal PVN Ucn3 projections to the MePD, which in turn relay via

the MePD Ucn3 neuronal circuitry an inhibitory influence on the ARC KNDy neuronal

network to suppress GnRH pulse generator frequency. Urocortin3 (Ucn3), Corticotropin-

releasing factor type 2 receptor (CRFR2), glutamate receptor (GlutR), the posterodorsal

subnucleus of the medial amygdala (MePD), paraventricular nucleus of hypothalamus (PVN),

the hypothalamic arcuate nucleus (ARC).

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6