PACT - Prediction of Amyloid Cross-interaction by Threading

Jakub W. Wojciechowski[1*], **Witold Szczurek**[1], **Małgorzta Kotulska**[1*]

*For correspondence:
jakub.wojciechowski@pwr.edu.pl
(JWW);
malgorzata.kotulska@pwr.edu.pl
(MK)

[1]Department of Biomedical Engineering, Faculty of Fundamental Problems of Technology, Wrocław University of Science and Technology, 50-370 Wrocław, Poland

Abstract

Amyloids are protein aggregates usually associated with their contribution to several diseases e.g., Alzheimer's and Parkinson's. However, they are also beneficially utilized by many organisms in physiological roles, such as microbial biofilm formation or hormone storage. Recent studies showed that an amyloid aggregate can affect aggregation of another protein. Such cross-interactions may be crucial for understanding comorbidity of amyloid diseases or influence of microbial amyloids on human amyloidogenic proteins. However, due to demanding experiments, understanding of interaction phenomenon is still limited. Moreover, no dedicated computational method to predict potential amyloid interactions has been available until now. Here, we present PACT - a computational method for prediction of amyloid cross-interactions. The method is based on modeling a heterogenous fibril formed by two amyloidogenic peptides. Stability of the resulting structure is assessed using a statistical potential that approximates energetic stability of a model. Importantly, the method can work with long protein fragments and, as a purely physicochemical approach, it relies very little on training data. PACT was evaluated on data collected in AmyloGraph database and it achieved high values of AUC (0.91) and F1 (0.81). The new method opens a possibility of high throughput studies of amyloid interactions. We used PACT to study interactions of CsgA, a bacterial biofilm protein from several bacterial species inhabiting human intestines, and human Alpha-synuclein protein which is involved in the onset of Parkinson's disease. We show that the method correctly predicted the interactions and highlighted the importance of specific regions in both proteins.
The tool is available as a web server at: https://pact.e-science.pl/pact/. The local version can be downloaded from: https://github.com/KubaWojciechowski/PACT

Introduction

Pathological misfolding and aggregation of proteins is a hallmark of a number of devastating disorders, including major public health challenges like Alzheimer's and Parkinson's diseases (*Ross and Poirier, 2004*), type II diabetes (*Haque et al., 2017*), as well as some cancers (*Yang-Hartwich et al., 2015*). These diseases not only share a similar molecular mechanism, but they also often co-occur in the same patients. Among others, comorbidities were observed between Alzheimer's disease and type II diabetes (*Janson et al., 2004*; *Sims-Robinson et al., 2010*) and Alzheimer's and Parkinson's diseases (*Rajput et al., 1993*). One of the possible explanations of this phenomenon could be related to amyloid cross-interactions. Amyloids are insoluble protein aggregates characterized by an exceptional stability due to the tight packing of monomers, resulting in characteristic pattern in X-ray diffraction experiments. (*Eisenberg and Sawaya, 2017*). Despite significant structural similarities shared by all amyloids, their sequences are surprisingly diverse and have little homology (*Sipe et al., 2010*). More recent studies have shown that in some cases presence of amyloid

aggregates can affect the aggregation rate of other proteins (*Lundmark et al., 2005*). Later, it was observed that interacting proteins can form heterogeneous fibers consisting of molecules of both interaction partners. Such cross-interactions were then identified between numerous proteins, including those involved in type II diabetes and neurodegenerative diseases. This shed new light on potentially new aspects regarding the origin of comorbidity of these disorders (*Ren et al., 2019*). Despite the importance of this process, its mechanisms are still poorly understood, which can be attributed to a limited number of experimental data. As a result, interactions of only a few well-described proteins, such as Amyloid-beta (Abeta), islet amyloid polypeptide, or Alpha-synuclein, have been very extensively studied and they contributed a majority of data. This may introduce a bias in available data. Despite the difficulties, it was shown that proteins with similar sequences are more likely to interact, however, many counterexamples were also shown (*Ren et al., 2019*). The studies highlight the importance of the structural compatibility of amyloid cores.

The main limitation regarding experimental studies of amyloid aggregation and their interactions is that they require expensive and time consuming methods. In practice, biochemical assays based on binding of Congo Red (*Howie and Brewer, 2009*), Thioflavin T (*Nielsen et al., 2001*) and infrared spectroscopy are frequently used. Especially the last method is widely applied, due to its simplicity and efficiency (*Szulc et al., 2021b*).Another approach involves direct observation of fibers using high resolution imaging techniques, such as electron microscopy (*Shirahama and Cohen, 1967*) and atomic force microscopy (*Wang et al., 2003*). Since different methods rely on different approaches, their results might differ in some cases. More importantly, all of them are expensive and time consuming. Experiments are hampered by difficulties in handling amyloids, including their low solubility, rapid aggregation, and need for their high-purity (*Gąsior-Głogowska et al., 2022*). Currently, the use of experimental methods for identification of all amyloids in genome wide studies would be impossible. To address this problem several computational methods have been proposed based on different approaches (reviewed in (*Kotulska and Wojciechowski, 2022*)), starting from structural modeling (*Thompson et al., 2006*), statistical analysis of the sequence including FoldAmyloid (*Garbuzynskiy et al., 2010*) and FishAmyloid (*Gasior and Kotulska, 2014*), physico-chemical models like PASTA 2.0 (*Walsh et al., 2014*), machine learning techniques such as APPNN (*Família et al., 2015*) and AmyloGram (*Burdukiewicz et al., 2017*). Furthermore, there are methods combining both approaches such as PATH (*Wojciechowski and Kotulska, 2020*) and Cordax (*Louros et al., 2020*). Finally, some methods, like Aggrescan 3D (*Kuriata et al., 2019*), utilize information about protein structure. It was also shown that bioinformatics techniques are quite robust and capable of identification even some misannotated data despite being trained on them (*Szulc et al., 2021a*). Unfortunately, none of these methods can predict amyloid interactions.

Here, we present a new computational method PACT (Prediction of Amyloid Cross-interaction by Threading) designed for identification of potentially interacting amyloid pairs. The method is based on the molecular threading applied to the potential complex of interacting amyloids and assessment of the stability of obtained molecular model.

Results

The main assumption of the method is that interactions between amyloidogenic fragments that cross-interact, threaded into an amyloid fiber structure, would result in a heterogeneous aggregate which is more stable, thus energetically more favorable than a non-interacting pair. In PACT, we use the Modeller software for threading a query sequence on the structure of amyloid fiber formed by Islet Amyloid Polypeptide (IAPP) (*Luca et al., 2007*). To asses obtained models we have proposed *ndope* score, which is a normalized version of DOPE statistical potential implemented in the Modeller software.

PACT correctly identifies amyloid prone peptides

In the first step, we focused on prediction of homoaggregation, which can be considered a special case of heteroaggregation. We compared *ndope* scores obtained for models of potential ho-

Figure 1. A) Distribution of *ndope* score for models of amyloidogenic and non-amyloidogenic peptides. B) ROC curve for amyloid vs non-amyloid classification. The orange line represents the distance between perfect classification point (0,1) and the chosen threshold

moagregates of amyloidogenic and non-amyloidogenic peptides, for which the sequences were obtained from the AmyLoad database (*Wozniak and Kotulska, 2015*). The majority of models obtained for amyloidogenic peptides showed much lower scores (meaning more stable structures) in comparison with non-amyloids and their first quartiles of the scores were well separated (Fig.1). Considering energy difference, we built a threshold based classifier. The classification threshold was chosen based on the Receiver Operating Characteristic (ROC) curve, as a point on the curve closest to the point (0,1), representing perfect classification (1B). The chosen value of *ndope* score was -231. If used merely for distingushing amyloids from non-amyloids, such a classifier was able to achieve Area Under ROC Curve (AUC) of 0.76 and *Accuracy* of 0.74. Moreover, well-balanced values of *Sensitivity* (0.73) and *Specifictity* (0.76) were obtained. Such results are comparable with state-of-the-art amyloid predictors on the same data set (Sup Tab S1).

We also tested if the method is capable of recognizing amyloid propensity in functional amyloids, which pose a major problem for most predictors due to their under-representation in databases of amyloids. We tested the performance of the method on imperfect repeats of CsgA protein from *Escherichia coli* and *Salmonella enterica* (*Szulc et al., 2021b*) (Sup Fig S1). Aggregation prone regions of this protein (R1, R3, and R5) scored much lower than non-amyloidogenic regions (R2 and R4) from *Escherichia coli*. On this data, PACT achieved the accuracy of 0.9. Furthermore, the observed difference between *ndope* score for R4 fragments from *Escherichia coli* and *Salmonella enterica* corresponds very well to the difference in their aggregation propensities (*Szulc et al., 2021b*).

The results showed that the method can accurately predict aggregation prone peptides of varying lengths. Furthermore, it can be utilized to detect functional amyloids.

PACT predicts amyloid cross-interactions

We used similar methodology to predict cross-interactions of amyloid peptides, which is the main purpose of the method. The *ndope* scores of heteroaggregates consisting of pairs of peptides whose cross-interactions resulted in faster aggregation were compared with non-amyloidogenic pairs of peptides (Fig.2). A similar analysis was performed for pairs of peptides whose cross-interactions resulted in slower aggregation (Sup Fig S3). In both cases, models of heterologous aggregates resulting from cross-interactions showed lower values of *ndope* scores than non-amyloids, and well separated first quartiles of their scores (Fig.2). Therefore, we built the threshold based classifier using the approach described in the previous section.

To assess the performance and choose the optimal threshold value, ROC curves were calculated

Figure 2. The score *ndope* for models of interacting identical non-amyloidogenic peptides (negative set) and interacting pairs resulting in increased aggregation rates (*faster* set).

for both cases; *faster* rate vs *negative* (Fig.2) and *slower* vs *negative* (Sup Fig S3) on both training and test sets. To minimize the impact of the data choice, we performed k-folds cross-validation with k=5 on the training set and calculated several metrics describing the performance of the method (Tab. 1). The same metrics were then calculated on an independent test set. The same analysis was performed for the case of prediction of interactions resulting in slower aggregation (Sup Tab S2). Chosen *ndope* thresholds were very similar in both scenarios, namely -236 and -235 for faster vs negative and slower vs negative respectively. PACT performed well on both cross-validation and independent test set. It achieved the *Accuracy* of 0.85 and 0.82 on test sets of *faster* vs *negative* and *slower* vs *negative* cases, respectively. In all cases, the results obtained on the test set were within the value of one standard deviation range from the mean values obtained with the cross-validation procedures. The method performance was quite similar in both *faster* vs *negative* and *slower* vs *negative* scenarios. However, due to the smaller data set size, larger standard deviation was obtained for *slower* vs *negative* scenario (Tab.1). The results show that the method can predict whether two peptides can cross-interact but cannot distinguish between types of interactions with regard to their rate.

PACT is robust to bias in data

A serious problem with data regarding interacting amyloids, available in the literature and, therefore, in our dataset is large overrepresentation of interactions concerning Abeta peptide. This may cause overfitting of the method to this group of sequences. To assess its effect, we analyzed scores for interactions between different Abeta variants (Sup Fig S4). Observed scores for Abeta pairs are within a range of values observed for the remaining pairs, and therefore they should not have a significant effect on the performance of the method. Those pairs showed relatively narrow distribution of *ndope* values, centered slightly below *ndope* value of -250, which is relatively close to identified classification threshold of -236, while the remaining interacting pairs showed even lower

Figure 3. ROC curves for classification of non-aggregating and cross interacting pairs resulting in faster aggregation on (A) training and (B) test set.

Performance of PACT on cross validation and independent test set for classification of non-aggregating and cross interacting pairs resulting in faster aggregation

	Acc [std]	Sens [std]	Spec [std]	F1 [std]	MCC [std]
Cross-validation	0.82 [0.06]	0.90 [0.09]	0.77 [0.08]	0.82 [0.07]	0.66 [0.12]
Test set	0.85	0.81	0.92	0.87	0.71

147 scores.

Interactions between bacterial amyloids and Alpha-synuclein

In recent years plenty of studies highlighted the connection between gut microbiome composition
and the onset of many diseases including neurodegenerative ones like Alzheimer's and Parkinson's
diseases (*Cani, 2018*). Despite extensive studies, understanding of molecular mechanisms of this
connection remains elusive. One of the possible explanations for this relates functional amyloids
from bacteria and human disease related amyloids through the cross-interaction theory. The ag-
gregation of the bacterial ones could speed up the aggregation of the disease related proteins,
leading to the disorder (*Friedland and Chapman, 2017*). This hypothesis seems consistent with re-
sults obtained by Chen and co-workers, who discovered increased production and aggregation of
Alpha-synuclein in rats exposed to bacterial strains producing biofilm related functional amyloids
(*Chen et al., 2016*).

In order to better understand this connection we studied possible interactions between bac-
terial functional amyloid CsgA and human Alpha-synuclein, whose aggregation is a hallmark of
Parkinson's disease. We used PACT to predict interactions of CsgA protein from five different or-
ganisms found in human microbiome; *Escherichia coli* (EC), *Hafnia alvei* (HA), *Yokenella regensburgei*
(YR), *Citrobacter youngae* (CY), and C*edecea davisae* (CD) with human Alpha-synuclein which were
recently studied experimentally by Bhoite and coworkers (*Bhoite et al., 2022*).

The sequence of Alpha-synuclein was divided into overlapping fragments of length 20 amino
acids and their interactions with R1-R5 repeats of each of CsgA proteins were tested. Consistently
with experimental results, all of the studied CsgA variants were predicted to interact with Alpha-
synuclein. Among CsgA proteins' fragments, R1, R3 and R5 were predicted to interact, with R5
showing the best scores (Fig4). These results are consistent with our current state of knowledge
about CsgA as the most aggregation prone regions in these proteins are R1, R3 ad R5. Furthermore,

Figure 4. Lowest *ndope* scores for interactions of R1-R5 repeats from each of CsgA proteins with Alpha-synuclein.

R5 fragment which showed the lowest *ndope* scores, is located at the protein surface, therefore it can interact without a need for major conformational changes. On the Alpha-synuclein part the best scoring region was located between positions 32 and 56 (Sup Fig S5). This region was recently shown to be of crucial importance for the aggregation of the protein (*Ulamec et al., 2022*).

Code Availability

PACT was implemented as an open source Python module, available at GitHub repository: https://github.com/KubaWojciechowski/PACT. For users' convenience we prepared a docker container for the application, as well as the web server: https://pact.e-science.pl/pact/

Discussion

High throughput identification of amyloid cross-interaction is an important step toward our understanding of this mechanism. It can allow for a better understanding of the principles governing the process but it can also be used to identify novel cases of amyloid interactions. Such capabilities can shed light on a possible mechanism responsible for the comorbidity of devastating disorders.

We proposed the first computational method for predicting amyloid cross-interactions. It is based on a highly interpretable and well-established physicochemical model which is not heavily dependent on training data. This feature is especially important since the available data contains a strong interest bias toward interactions of a few popular amyloids related to neurodegenerative diseases, such as Abeta. We have carefully studied the effect of this overrepresentation and showed that it does not affect the performance of the method. Furthermore, good performance on functional amyloids suggests that the method is very robust and can be effectively used on a wide range of sequences. In total, PACT achieved a high accuracy of 0.85 and 0.82 on the independent test sets of interactions increasing and decreasing aggregation rates. On both sets the method

achieved high AUC values of 0.91 and F1 values of 0.81 and 0.80. On the other hand, since both cases were characterized by similar interaction energies, the method cannot distinguish between enhancement and inhibition of aggregation cases. These results suggest that both processes may be driven by similar mechanisms. This issue was addressed in a recently published work by Louros and coworkers (*Louros et al., 2022*), who applied a somewhat similar approach to study the effect of point mutations on aggregation characteristics.

We used PACT to predict interactions of bacterial functional amyloid CsgA from different species with human Alpha-synuclein. Not only our results are in a good agreement with recently published data but they can also indicate which regions can drive the cross-interactions between both proteins. Identification of potentially interacting regions can provide an important insight into possible mechanism of the process and guide the future experiments.

Apart from identification of amyloid cross-interactions, the proposed method is also capable of reliably predicting amyloid prone regions in proteins with comparable accuracy to the state-of-the-art techniques. Furthermore, it overcomes their major limitations regarding the identification of functional amyloids. Unlike most of the currently available amyloid predictors, it does not rely on scanning of a query sequence with a short sliding window.

Methods

The main assumption of the method is that interactions between amyloidogenic fragments that cross-interact, threaded into an amyloid fiber structure, would result in a heterogeneous aggregate which is more stable, thus energetically more favorable than a non-interacting pair. A somewhat similar assumption was successfully applied in our previous work to predict the aggregation of short amyloidogenic fragments (*Wojciechowski and Kotulska, 2020*), although the current approach differs in other aspects of the method and the objectives. In PACT, we use the Modeller software for threading a query sequence on the structure of amyloid fiber formed by Islet the Amyloid Polypeptide (IAPP) (*Luca et al., 2007*). To asses obtained models we have proposed *ndope* score, which is a normalized version of the DOPE statistical potential implemented in the Modeller software.

Data sets

To build and test the method we used the following datasets:

- the set of 90 amyloidogenic (*amyloid*) and 55 non-amyloidogenic (*non-amyloid*) peptides of lengths between 14 and 45 from the AmyLoad database (*Wozniak and Kotulska, 2015*).
- the set of 119 pairs of peptides, which enhance (*faster* dataset) and 73 which slow down (*slower* dataset) the aggregation of each other. Both from AmyloGraph database (*Burdukiewicz et al., 2022*).

The first two sets (*amyloid* and *non-amyloid*) were used to test the method on cases of homoaggregation i.e., identifying amyloid prone peptides.

For the prediction of cross-interactions, we used *faster*, *slower* and non-amyloid sets. The use of the set of non-aggregating peptides as the negative set in the interaction study was caused by the lack of a sufficient number of negative examples of non-interacting amyloid pairs. This is a common problem in studies of protein-protein interactions since negative results are rarely published, which often creates a strong bias in biological data (*Nowakowska and Kotulska, 2022*).

Modeling

A query pair of sequences was threaded on the structure of amyloid fiber formed by Islet Amyloid Polypeptide (IAPP) (*Luca et al., 2007*). In order to allow the method to deal with sequences of varying lengths, sequences shorter than the sequence of a template use only the main part of the template structure. In such cases, a shorter sequence is aligned to the middle of the template

Figure 5. Schematic representation of the modeling procedure. A) In case when a query sequence is shorter than the template, only a part of it is used in modeling. B) The model of heterogenous fibril consists of two chains of each interacting peptide.

sequence (Fig 4A). This choice can be justified considering that most of the currently known amyloid
fragments, which are longer than a few amino acids, share a similar beta-sheet turn architecture,
commonly known as the beta arch. This assumption was previously successfully applied by Ahmed
and coworkers to build the ArchCandy method for amyloidogenic region prediction (***Ahmed et al.,***
2015). PACT allows sequences to be marginally longer than the template and, as a result, can
be used to study cross-interactions between peptides of lengths between 14 and 45. For each
of the tested pairs, 10 different models, consisting of two chains of each interacting peptide (Fig
4/B) were built using Modeller 9.24 model-multichain.py procedure with default parameters (***Šali***
and Blundell, 1993). Then, the model with the lowest DOPE value was chosen for further analysis.
Since the dataset consisted of fragments of varying lengths, we proposed a normalized DOPE score
(*ndope*) defined as follows:

$$ndope = \frac{DOPE}{L} \tag{1}$$

where L is an average length of sequences used to build a given model. Then, *ndope* scores were
compared between amyloids and non-amyloids, as well as between pairs of amyloids interacting
with non-amyloids.
To choose the *ndope* threshold for the classification, Receiver Operating Characteristics (ROC)

curve was calculated by applying different score thresholds and recording False Positive Ratio (FPR)
and True Positive Ratio (TPR). The threshold closest to the (0,1) point (representing perfect classifi-
cation) was chosen.
We also tested a variant of the method which utilized three different structural templates, how-
ever it did not improve the accuracy of the method but significantly increased the computational
time. Therefore, this approach was finally abandoned.

Assessment of performance and data analysis

All the data analysis was performed using Python 3.8 with Matplotlib (*Hunter, 2007*), NumPy (*Harris
et al., 2020*), Pandas (*McKinney et al., 2010*), Scikit-Learn (*Pedregosa et al., 2011*), and Seaborn
(*Waskom, 2021*) packages.
To test the performance of the proposed method, a dataset was randomly split into a training
set and test set, which consisted of 30% of the data. Additionally, k-folds cross-validation (with k=5)
was performed on the training data. Area Under ROC curve (AUC), *Accuracy* (ACC), *Sensitivity* (Sens),
Specificity (Spec) and Matthew Correlation Coefficient (MCC) were used to assess the performance
of the method

Effect of Amyloid-beta variants

270 For the analysis of the effect of over represented amyloid beta pairs, we divided the *faster* data
271 set into two subsets: one containing only pairs were both interacting peptides were variants of
272 amyloid beta (16 pairs) (*abeta*), and the set of remaining pairs (39 pairs) (*no abeta*).

Interactions between bacterial amyloids and Alpha-synuclein

Modeling the interactions between Alpha-synuclein and CsgA proteins was performed using hu-
man Alpha-synuclein sequence (Uniprot id: P37840) and CsgA protein from five different organ-
isms found in human microbiome; *Escherichia coli* (EC) (Uniprot id:), *Hafnia alvei* (HA) (Uniprot id:
G9Y7N6) , *Yokenella regensburgei* (YR) (Uniprot id: A0A6H0K4L9), *Citrobacter youngae* (CY) (Uniprot
id: A0A549VPM7), and C*edecea davisae* (CD) (Uniprot id: S3IYN9). The sequence of Alpha-synuclein
was divided into overlaping subsequences of length 20. CsgA variants were split into five non-
overlaping fragments R1-R5 corresponding to five imperfect repeats observed in their sequences.
Interactions of each of CsgA fragments with all Alpha-synuclein fragments were studied.

Acknowledgments

283 We would like to thank the team of AmyloGraph developers for their effort in preparing the database
284 used in this project.
285   This work was partially supported by the National Science Centre, Poland, Grant 2019/35/B/NZ2/03997.
286 Access to Wroclaw Centre for Networking and Supercomputing is greatly acknowledged.

References

Ahmed AB, Znassi N, Château MT, Kajava AV. A structure-based approach to predict predisposition to amyloi-
dosis. Alzheimer's & Dementia. 2015; 11(6):681–690.

Bhoite SS, Han Y, Ruotolo BT, Chapman MR. Mechanistic insights into accelerated α-synuclein aggregation
mediated by human microbiome-associated functional amyloids. Journal of Biological Chemistry. 2022; p.
102088.

Burdukiewicz M, Rafacz D, Barbach A, Hubicka K, Bąkała L, Lassota A, Stecko J, Szymańska N, Wojciechowski
JW, Kozakiewicz D, et al. AmyloGraph: a comprehensive database of amyloid–amyloid interactions. Nucleic
Acids Research. 2022; .

Burdukiewicz M, Sobczyk P, Rödiger S, Duda-Madej A, Mackiewicz P, Kotulska M. Amyloidogenic motifs re-
vealed by n-gram analysis. Scientific reports. 2017; 7(1):1–10.

Cani PD. Human gut microbiome: hopes, threats and promises. Gut. 2018; 67(9):1716–1725.

299 **Chen SG**, Stribinskis V, Rane MJ, Demuth DR, Gozal E, Roberts AM, Jagadapillai R, Liu R, Choe K, Shivakumar
300 B, et al. Exposure to the functional bacterial amyloid protein curli enhances alpha-synuclein aggregation in
301 aged Fischer 344 rats and Caenorhabditis elegans. Scientific reports. 2016; 6(1):1–10.

302 **Eisenberg DS**, Sawaya MR. Structural studies of amyloid proteins at the molecular level. Annu Rev Biochem.
303 2017; 86(1):69–95.

304 **Família C**, Dennison SR, Quintas A, Phoenix DA. Prediction of peptide and protein propensity for amyloid
305 formation. PloS one. 2015; 10(8):e0134679.

306 **Friedland RP**, Chapman MR. The role of microbial amyloid in neurodegeneration. PLoS pathogens. 2017;
307 13(12):e1006654.

308 **Garbuzynskiy SO**, Lobanov MY, Galzitskaya OV. FoldAmyloid: a method of prediction of amyloidogenic regions
309 from protein sequence. Bioinformatics. 2010; 26(3):326–332.

310 **Gasior P**, Kotulska M. FISH Amyloid–a new method for finding amyloidogenic segments in proteins based on
311 site specific co-occurence of aminoacids. BMC bioinformatics. 2014; 15(1):1–8.

312 **Gąsior-Głogowska ME**, Szulc N, Szefczyk M. Challenges in Experimental Methods. In: *Computer Simulations of*
313 *Aggregation of Proteins and Peptides* Springer; 2022.p. 281–307.

314 **Haque E**, Kamil M, Irfan S, Hasan A, Sheikh S, SSM M. Protein aggregation: A new challenge in type-II diabetes.
315 Adv Biotech & Micro. 2017; .

316 **Harris CR**, Millman KJ, Van Der Walt SJ, Gommers R, Virtanen P, Cournapeau D, Wieser E, Taylor J, Berg S, Smith
317 NJ, et al. Array programming with NumPy. Nature. 2020; 585(7825):357–362.

318 **Howie AJ**, Brewer DB. Optical properties of amyloid stained by Congo red: history and mechanisms. Micron.
319 2009; 40(3):285–301.

320 **Hunter JD**. Matplotlib: A 2D graphics environment. Computing in science & engineering. 2007; 9(03):90–95.

321 **Janson J**, Laedtke T, Parisi JE, O'Brien P, Petersen RC, Butler PC. Increased risk of type 2 diabetes in Alzheimer
322 disease. Diabetes. 2004; 53(2):474–481.

323 **Kotulska M**, Wojciechowski JW. Bioinformatics Methods in Predicting Amyloid Propensity of Peptides and
324 Proteins. In: *Computer Simulations of Aggregation of Proteins and Peptides* Springer; 2022.p. 1–15.

325 **Kuriata A**, Iglesias V, Pujols J, Kurcinski M, Kmiecik S, Ventura S. Aggrescan3D (A3D) 2.0: prediction and engi-
326 neering of protein solubility. Nucleic acids research. 2019; 47(W1):W300–W307.

327 **Louros N**, Orlando G, De Vleeschouwer M, Rousseau F, Schymkowitz J. Structure-based machine-guided map-
328 ping of amyloid sequence space reveals uncharted sequence clusters with higher solubilities. Nature com-
329 munications. 2020; 11(1):1–13.

330 **Louros N**, Ramakers M, Michiels E, Konstantoulea K, Morelli C, Garcia T, Moonen N, D'Haeyer S, Goossens V,
331 Thal DR, et al. Mapping the sequence specificity of heterotypic amyloid interactions enables the identification
332 of aggregation modifiers. Nature communications. 2022; 13(1):1–20.

333 **Luca S**, Yau WM, Leapman R, Tycko R. Peptide conformation and supramolecular organization in amylin fibrils:
334 constraints from solid-state NMR. Biochemistry. 2007; 46(47):13505–13522.

335 **Lundmark K**, Westermark GT, Olsén A, Westermark P. Protein fibrils in nature can enhance amyloid protein
336 A amyloidosis in mice: Cross-seeding as a disease mechanism. Proceedings of the National Academy of
337 Sciences. 2005; 102(17):6098–6102.

338 **McKinney W**, et al. Data structures for statistical computing in python. In: *Proceedings of the 9th Python in*
339 *Science Conference*, vol. 445 Austin, TX; 2010. p. 51–56.

340 **Nielsen L**, Frokjaer S, Brange J, Uversky VN, Fink AL. Probing the mechanism of insulin fibril formation with
341 insulin mutants. Biochemistry. 2001; 40(28):8397–8409.

342 **Nowakowska AW**, Kotulska M. Topological analysis as a tool for detection of abnormalities in protein-protein
343 interaction data. Bioinformatics. 2022; .

Pedregosa F, Varoquaux G, Gramfort A, Michel V, Thirion B, Grisel O, Blondel M, Prettenhofer P, Weiss R, Dubourg V, et al. Scikit-learn: Machine learning in Python. the Journal of machine Learning research. 2011; 12:2825–2830.

Rajput A, Rozdilsky B, Rajput A. Alzheimer's disease and idiopathic Parkinson's disease coexistence. Journal of geriatric psychiatry and neurology. 1993; 6(3):170–176.

Ren B, Zhang Y, Zhang M, Liu Y, Zhang D, Gong X, Feng Z, Tang J, Chang Y, Zheng J. Fundamentals of cross-seeding of amyloid proteins: an introduction. Journal of Materials Chemistry B. 2019; 7(46):7267–7282.

Ross CA, Poirier MA. Protein aggregation and neurodegenerative disease. Nature medicine. 2004; 10(7):S10–S17.

Šali A, Blundell TL. Comparative protein modelling by satisfaction of spatial restraints. Journal of molecular biology. 1993; 234(3):779–815.

Shirahama T, Cohen AS. High-resolution electron microscopic analysis of the amyloid fibril. The Journal of cell biology. 1967; 33(3):679–708.

Sims-Robinson C, Kim B, Rosko A, Feldman EL. How does diabetes accelerate Alzheimer disease pathology? Nature Reviews Neurology. 2010; 6(10):551–559.

Sipe JD, Benson MD, Buxbaum JN, Ikeda SI, Merlini G, Saraiva MJ, Westermark P. Amyloid fibril protein nomenclature: 2010 recommendations from the nomenclature committee of the International Society of Amyloidosis. Amyloid. 2010; 17(3-4):101–104.

Szulc N, Burdukiewicz M, Gąsior-Głogowska M, Wojciechowski JW, Chilimoniuk J, Mackiewicz P, Šneideris T, Smirnovas V, Kotulska M. Bioinformatics methods for identification of amyloidogenic peptides show robustness to misannotated training data. Scientific reports. 2021; 11(1):1–11.

Szulc N, Gąsior-Głogowska M, Wojciechowski JW, Szefczyk M, Żak AM, Burdukiewicz M, Kotulska M. Variability of amyloid propensity in imperfect repeats of csga protein of salmonella enterica and escherichia coli. International journal of molecular sciences. 2021; 22(10):5127.

Thompson MJ, Sievers SA, Karanicolas J, Ivanova MI, Baker D, Eisenberg D. The 3D profile method for identifying fibril-forming segments of proteins. Proceedings of the National Academy of Sciences. 2006; 103(11):4074–4078.

Ulamec SM, Maya-Martinez R, Byrd EJ, Dewison KM, Xu Y, Willis LF, Sobott F, Heath GR, van Oosten Hawle P, Buchman VL, et al. Single residue modulators of amyloid formation in the N-terminal P1-region of α-synuclein. Nature communications. 2022; 13(1):1–16.

Walsh I, Seno F, Tosatto SC, Trovato A. PASTA 2.0: an improved server for protein aggregation prediction. Nucleic acids research. 2014; 42(W1):W301–W307.

Wang Z, Zhou C, Wang C, Wan L, Fang X, Bai C. AFM and STM study of β-amyloid aggregation on graphite. Ultramicroscopy. 2003; 97(1-4):73–79.

Waskom ML. Seaborn: statistical data visualization. Journal of Open Source Software. 2021; 6(60):3021.

Wojciechowski JW, Kotulska M. Path-prediction of amyloidogenicity by threading and machine learning. Scientific Reports. 2020; 10(1):1–9.

Wozniak PP, Kotulska M. AmyLoad: website dedicated to amyloidogenic protein fragments. Bioinformatics. 2015; 31(20):3395–3397.

Yang-Hartwich Y, Bingham J, Garofalo F, Alvero AB, Mor G. Detection of p53 protein aggregation in cancer cell lines and tumor samples. In: *Apoptosis and Cancer* Springer; 2015.p. 75–86.