A fully phased interspecific grapevine rootstock genome sequence representing *V. riparia* and *V. cinerea* and allele-aware annotation of the phylloxera resistance locus *Rdv1*

Bianca Frommer[1], Ludger Hausmann[2], Daniela Holtgräwe[1], Prisca Viehöver[1], Bruno Hüttel[3], Richard Reinhardt[3], Reinhard Töpfer[2], Bernd Weisshaar[1]

[1] Bielefeld University, Chair of Genetics and Genomics of Plants, Faculty of Biology & Center for Biotechnology (CeBiTec), Bielefeld, Germany
[2] Julius Kuehn Institute (JKI), Institute for Grapevine Breeding Geilweilerhof, Siebeldingen, Germany
[3] Max Planck Genome Centre Cologne, Max Planck Institute for Plant Breeding Research, Cologne, Germany

Email addresses:
BF: frommer@CeBiTec.Uni-Bielefeld.DE / ORCID: 0000-0002-5792-0102
LH: ludger.hausmann@julius-kuehn.de / ORCID: 0000-0002-4046-9626
DH: dholtgra@CeBiTec.Uni-Bielefeld.DE / ORCID: 0000-0002-1062-4576
PV: viehoeve@CeBiTec.Uni-Bielefeld.DE
BH: huettel@mpipz.mpg.de / ORCID: 0000-0001-7165-1714
RR: reinhardt@mpipz.mpg.de / ORCID: 0000-0001-9376-2132
RT: reinhard.toepfer@julius-kuehn.de / ORCID: 0000-0003-1569-2495
BW: bernd.weisshaar@Uni-Bielefeld.DE / ORCID: 0000-0002-7635-3473

Abstract

The phylloxera resistant rootstock cultivar 'Börner' is an interspecific hybrid derived from *Vitis riparia* and *V. cinerea* and a valuable resource for *Vitis* disease resistances. We created a fully phased, high-quality 'Börner' genome sequence named BoeRC using long PacBio reads. Comprehensive gene annotation of both 'Börner' haplotypes, designated BoeRip and BoeCin, was applied to describe the phylloxera resistance locus *Rdv1*. Using a mapping population derived from a susceptible *V. vinifera* breeding line and 'Börner', the *Rdv1* locus was further delimited. *Rdv1*, which is derived from *V. cinerea* and included in the haplotype BoeCin, was compared with sequences of phylloxera-susceptible and phylloxera-tolerant cultivars. Between flanking regions that display high synteny, we detected and precisely characterized a diverse sequence region that covers between 202 to 403 kbp in different haplotypes. In BoeCin, five putative disease resistance genes were identified that represent likely candidates for conferring resistance to phylloxera.

Keywords

Assembly, PacBio, SMRT sequencing, grapevine, rootstock, genome, Canu, annotation, resistance, resistance gene analogs, phylloxera, Rdv1

Background

Vitis vinifera Linné subspecies vinifera is the most cultivated Vitis species worldwide. It is economically highly important and appreciated for wine production, table grapes, dry fruits like raisins, and other products. The species is endemic to Europe, and domestication and development of cultivars has a very long history [1]. V. vinifera is highly susceptible to many diseases and pests including downy and powdery mildew as well as to the root aphid phylloxera (Daktulosphaira vitifoliae Fitch) [2]. In the 19th century, phylloxera caused a crisis of unprecedented dimension for viticulture and questioned the survival of viticulture in general. Vitis vinifera plants died quickly after phylloxera attack of the roots and huge cultivation areas were destroyed. Only the grafting of susceptible V. vinifera cultivars as scions to phylloxera-tolerant rootstocks derived from American wild Vitis species or their interspecific hybrids rescued viticulture [3, 4]. The phylloxera plague remains as a serious case even if it is controlled at the moment.

The selection of a good rootstock that provides resistances or tolerances to the almost ubiquitous phylloxera is crucial for nowadays viticulture. One newly developed rootstock variety is the cultivar 'Börner', an interspecific F1 hybrid derived from a cross of the American Vitis genotypes Vitis riparia GM183 and Vitis cinerea Arnold. 'Börner' inherited full resistance to phylloxera on roots from V. cinerea and develops no nodosities or tuberosities [5] upon infestation with phylloxera (Fig. 1). V. cinerea is an American wild species known to often provide high resistance at the root and very high resistance at the leaf against phylloxera [6]. A quantitative trait locus (QTL), located on chromosome (chr) 13, was identified to provide phylloxera resistance [6] and is referred to as Rdv1 (resistance to D. vitifoliae) [7]. In addition, 'Börner' inherited positive viticultural traits including tolerance to drought, resistance to black rot caused by the fungus Guignardia bidwellii [8], and resistance to downy mildew caused by Plasmopara viticola [9, 10].

71
Fig. 1: Root balls of susceptible and resistant grapevine plants attacked by
phylloxera. Some phylloxera-induced nodosities on the roots are marked with red
arrows. The pictures show roots of the grapevines indicated, the two F1 plants
selected with the phenotypes as indicated are derived from the mapping population
V3125 x 'Börner' [6].

Disease resistance, usually conferred through resistance genes, makes plants capable
of surviving attacks from a broad variety of pathogens and pests [11]. So called resis-
tance gene analogues (RGAs) are identified by the protein domains they encode and
structural features which are similar to validated resistance genes [12]. Based on the
protein domain structure that also indicates the localisation and/or site of action within
the cell, they are broadly classified as nucleotide-binding site and leucine rich repeat
domain receptor (NLRs) or pattern-recognition receptors (PRRs) [11]. Usually, PRRs

act at the cell surface because of a transmembrane (TM) domain, while NLRs cover a

nucleotide-binding site (NBS) domain and usually act intracellularly.

Vitis genomes cover a haploid set of 19 chromosomes, and the haploid genome size

varies around 500 Mbp [13, 14]. The inbred cultivar 'PN40024' has been sequenced [15, 16]

and the resulting assembly serves as a reference genome sequence (referred to as

PN40024). However, since *Vitis* species and cultivars like almost all perennial plants

are highly heterozygous and genetically divers [10, 17, 18], there is a need for more *Vitis*

genome sequences. Several additional *Vitis* genome sequences have become available

which were generated from long reads, including those of Cabernet Sauvignon [19, 20],

Carménère [21], Chardonnay [14, 22], *Muscadinia rotundifolia* cultivar Trayshed [23] and *V.*

riparia Gloire de Montpellier (referred to here as VitRGM) [24]. These assemblies each

cover a primary and an alternate pseudo-haplotype of which the alternate pseudo-

haplotype is usually of reduced contiguity. The cultivar *V. riparia* Gloire de Montpel-

lier is tolerant against root infestation by phylloxera [25], but to the best of our knowl-

edge it is unknown if this tolerance is linked to the *Rdv1* locus or to another region of

the genome.

Long read DNA sequencing technology, like "Single Molecule, Real Time" (SMRT)

sequencing [26] provided by Pacific Biosciences (PacBio), is one option to generate

high quality genome sequence assemblies. The long reads more likely span problem-

atic genomic regions and thus pave the way for more contiguous assemblies [27]. Also,

information of the haplotype and phase differences are more completely contained in

a single read. This fact, together with the transmission of exactly one haplotype from

two parents to a single offspring according to Mendel's laws, is exploited by the "trio

binning" approach for generating fully phased genome sequence assemblies [28].

To access and resolve the *Rdv1* locus, we set out to generate a haplotype-resolved

genome sequence of 'Börner' with the goal to overcome complications caused by

high heterozygosity and the complexity of resistance gene clusters. Using SMRT se-

quencing to generate long reads and additional Illumina short read data from both

parents of 'Börner', we assembled both haplotype sequences of 'Börner' at chromo-

some level. The two truly phased haplotypes of the diploid interspecific hybrid

'Börner' represent the genome sequences of the two species *V. riparia* and *V. cinerea*.

Structural and functional gene annotation was performed for both haplotype se-

quences, and RGAs were studied. The new sequence and annotation data as well as

additional mapping results were used to dissect the *Rdv1* locus of 'Börner' at the gene

level.

Results

Sequencing data of 'Börner' and its parents

To assemble the 'Börner' genome sequence, ~66 Gbp of raw SMRT sequencing data comprising 7,328,737 subreads with an average read length of 9,016 bp and an N50 length of 12,963 bp were generated. The expected diploid genome size (2n) of 'Börner' is 2x500 Mbp; thus the calculated coverage with long reads for a genome sequence with merged haplotypes would be 132 times, or more than 60-fold coverage for each individual haplotype. To allow k-mer-based binning of the long reads according to the two haplotypes that 'Börner' inherited from its parents, the parental genotypes were sequenced with Illumina technology to about 140-fold coverage.

A 'Börner' genome sequence assembly in two phases

To generate a 'Börner' genome sequence with two separated haplotypes, the trio-binning approach was applied [28]. The two haplotype-specific 'Börner' read subsets or bins, designated Vrip and Vcin, contain approximately 50 % of all reads and bases. These were assembled with Canu (see Methods). Statistics of the two resulting haplotype assemblies are shown in Table 1. Both assemblies reached the expected genome size of ~500 Mbp. The BoeRip and BoeCin haplotype assemblies represent "1n double haploid" genome sequences of the two species *V. riparia* and *V. cinerea*, respectively.

Table 1 Assembly statistics for the raw and scaffolded haplotype assemblies BoeRip and BoeCin of 'Börner'.

	Scaffold Level[a]		Total	
	BoeRip	BoeCin	Contigs[b]	Scaffolds
Sequences	971	767	1,843	1,738
Size (bp)	495,882,484	501,563,116	997,019,452	997,445,600
Largest seq. (bp)	14,614,105	16,784,012	16,784,012	16,784,012
N50 (Mbp)	5.29	5.34	4.79[c]	5.29
N90 (Mbp)	0.72	0.51	0.49[c]	0.55

[a]scaffolded haplotype assemblies
[b]raw haplotype assemblies, numbers include contigs shorter than 10 kbp
[c]see Supplementary File 1 Fig. S1 for details

Pseudochromosome construction and validation by genetic map

The assignment of scaffolds to pseudochromosomes was mainly based on the

PN40024 reference sequence and was achieved by using RBHs (see Methods). For

BoeRip, 446 scaffolds accounting for 95.21 % of all bases were assigned to pseudo-

chromosomes. For BoeCin, 320 scaffolds holding 93.89 % of all bases were assigned.

Since the reference contains a pseudochromosome chr00 for unassigned sequences,

this has been created as well and is referred to as "chrUn". In addition, some scaffolds

find no clear corresponding sequence in the reference and result in a pseudochromo-

some "chrNh" (for no hit, see Fig. 2A). As a result, the haplotype assemblies each

consist of 21 pseudochromosome sequences representing the 19 true chromosomes

and two artificial sequences.

To validate the pseudochromosomes, 314 SSR markers that show unequivocal chro-

mosome mappings on the haplotype assemblies were evaluated relative to their posi-

tion on PN40024. A total of 309 markers confirm the pseudochromosome assignment.

Conflicting markers were checked manually and corrected if possible (see Methods).

The completeness of the assemblies was evaluated by detection of plant core genes

with BUSCO (Fig. 2B). For both BoeRip and BoeCin, about 96.5 % of the conserved

single copy gene set were detected. This indicated that both haplotype assemblies are

more complete than the assemblies PN40024 and VitRGM (Supplementary File 2

Table S2).

Fig. 2 Comparison of the 'Börner' genome sequence assembly with other *Vitis* assemblies. (A) Pseudochromosome lengths comparison of both 'Börner' haplotypes, PN40024 and VitRGM. The bars represent the pseudochromosomes of the different assemblies. Red, BoeRip; dark red, VitRGM; grey, PN40024; blue, BoeCin. ChrUn holds sequences that were assigned to chr00 of PN40024 or VitRGM, and chrNh collects sequences with no assignment (Supplementary File 2 Table S1). **(B)** Plant core gene content (2,326 eudicots genes in reference set) of the assemblies in comparison to the PN40024 and VitRGM genome assemblies. Note that the bar graph is truncated at the left and focusses on only the duplicated, fragmented and missing BUSCO genes. The track labelled BoeRC represents the whole 'Börner' assembly (combined BoeRip and BoeCin sequences; merged results from the two haplotypes).

Evaluation of assembly quality and phasing

For validation of the phasing, bacterial artificial chromosome (BAC) sequences (Supplementary File 2 Table S3) with known haplotype/parental origin were mapped on BoeRip and BoeCin (Supplementary File 2 Table S4). The BACs were selected to cover regions on chr01 and chr14 and were sequenced with a read coverage of about 1,000-fold (Supplementary File 2 Table S5). BAC contigs of the same haplotype map

with almost no sequence difference, while BAC contigs derived from the other haplo-
type display a wide range of mismatches and InDels (Supplementary File 1 Fig. S2).
On both haplotypes, ~3.6 Mbp (chr01) and ~5.5 Mbp (chr14) were covered by BAC
contigs of both phases and support a correct phasing over 9 Mbp.

In addition, the assembly quality evaluation tool Merqury (phasing assessment for
genome sequence assemblies) was used to assess the quality of both haplotype assem-
blies. The overall base quality value (QV, consensus quality value, representing the
log-scaled probability of error for consensus base calls) of 37.42 found for BoeRC
indicates a very high level of correctness (see Supplementary File 2 Table S2 for
comparisons to other assemblies). It should be noted that Merqury requires Illumina
data for the complete trio (both parents and F1 as available for 'Börner' and its ances-
tors) to calculate the complete set of quality values.

Also the phasing accuracy was evaluated with Merqury (Supplementary File 1 Fig.
S3). Based on haplotype-specific k-mers (referred to as hap-mers in Mercury), 2,759
phase blocks with an N50 of 2.25 Mbp were calculated for BoeRip and 1,852 blocks
with an N50 of 2.42 Mbp for BoeCin. Almost all bases were covered by a block and
the switch error rate was estimated to be less than 0.05 % (Supplementary File 2 Ta-
ble S6; Supplementary File 1 Fig. S4). Overall, the quality of the 'Börner' genome
sequence as well as its separation into haplotypes derived from *V. riparia* GM183
(BoeRip) and *V. cinerea* Arnold (BoeCin) is considered to be high.

Sequence similarities and variations between BoeRip, BoeCin, PN40024 and
VitRGM

All-versus-all alignments between the pseudochromosomes of the two 'Börner' haplo-
types, and between each haplotype and PN40024 as well as VitRGM, were computed
and visualised as dot plots (Fig. 3, Supplementary File 1 Fig. S5 to S8). All homolo-
gous pseudochromosomes show strong synteny, yet some rearrangements and smaller
and larger gaps indicating insertions, deletions and/or missing sequence were de-
tected.

On average 90.20 % of all bases with an identity of 95.52 % were aligned between the
pseudochromosomes of the two haplotypes except of chrUn and chrNh (see Methods).
The SNP and InDel frequency for the protein coding fraction was 1/1,001 bp and 1/34
bp for the non-coding fraction of the genome sequence. When aligning with

PN40024, on average 88.59 % and 87.51 % of all bases were aligned with an identity
of 94.82 % (BoeRip) and 94.94 % (BoeCin) over all pseudochromosomes, respec-
tively. The alignments resulted in a SNP and InDel frequency of 1/932 bp (BoeRip)
and 1/935 bp (BoeCin) for the coding regions and of 1/32 bp for the non-coding re-
gions.
An alignment of BoeRip and BoeCin with VitRGM resulted in 93.22 % (BoeRip) and
88.45 % (BoeCin) aligned bases with an identity of 98.53 % (BoeRip) and 95.29 %
(BoeCin). The SNP and InDel frequency was 1/3,128 bp (BoeRip) and 1/1,019 bp
(BoeCin) for coding regions and 1/94 bp (BoeRip) and 1/35 bp (BoeCin) for non-
coding regions.

Fig. 3 All-versus-all dot plot of both 'Börner' haplotypes. The graphic shows the
dot plot between concatenated pseudochromosomes of BoeRip and BoeCin.

Gene annotation

In total, ~274 Mbp (55.24 %, BoeRip) and ~278 Mbp (55.46 %, BoeCin) repetitive sequences were identified with the MAKER pipeline (Supplementary File 2 Table S7). The final gene annotation comprises 27,738 and 27,995 protein-coding genes (see Data availability for links to the genome sequence annotation of the *V. cinerea* and the *V. riparia* haplotypes) and 525 and 419 tRNA genes for BoeRip and BoeCin, respectively (Supplementary File 2 Table S7). Of the protein-coding genes, 19,500 genes of each haplotype were captured in RBHs. Based on RBHs, BoeRip shares 17,922 genes with VCost.v3 (PN40024) and 18,928 genes with VitRGM; BoeCin shares 17,837 genes with VCost.v3 (PN40024) and 18,126 genes with VitRGM.

Resistance genes

The annotated genome sequence of 'Börner' was used to identify RGAs based on their typical protein domain structure. Resistance gene annotation with RGAugury revealed 2,081 RGAs for BoeRip and 2,009 RGAs for BoeCin (Fig. 4). Of these, 1,509 were classified as RBHs between BoeRip and BoeCin.

Fig. 4 Distribution of genes to RGA classes based on characteristic protein domains. The red bars display the number of RGA genes of BoeRip and the blue bars the RGA genes of BoeCin. RGAs were classified according to the domains encoded by the predicted genes. Designations were adapted according to [29]. NBS, Nucleotide Binding Site; CC-NBS-LRR with CC for Coiled-Coil and LRR for Leucine Rich Repeat; TIR-NBS-LRR with TIR for Toll/Interleukin-1 Receptor like; TIR-X with X for unknown domain; Other including TIR-CC-NBS-LRR or TIR-NBS-LRR-CC-TM

249 with TM for TransMembrane region; RLP, Receptor Like Protein; RLK, Receptor
250 Like Kinase.

Candidate genes for Rdv1

At the sequence level, the region between GF13-01 and GF13-11 that contains *Rdv1*
displays a highly diverged structure between BoeRip, BoeCin, PN40024 and VitRGM
(Fig. 5A, Supplementary File 1 Fig. S9, S10).
The interval flanked by these genetic markers includes 2,978,547 bp with 179 genes
for BoeRip and 2,538,570 bp with 173 genes for BoeCin, according to the BoeRC
assembly and its annotation, as well as 2,796,801 bp with 273 genes for PN40024 and
5,854,243 bp with 450 genes for VitRGM according to published data. The interval
between GF13-01 and GF13-11 in VitRGM, which has been generated with a FAL-
CON-Unzip and which is representing a pseudo-haplotype, is twice as large as in the
other assemblies.
To further delimitate *Rdv1*, new genetic markers were designed and used to establish
a local map. F1 individuals with different flanking haplotypes were selected from the
mapping population V3125 x 'Börner', genotyped and tested for phylloxera resis-
tance. Since marker assay design tuned out to be difficult for gene-level resolution,
genotyping-by-sequencing (GBS) was applied to five crucial F1 genotypes. Mapping
of SSR and GBS markers as well as the presence or absence of the resistance encoded
by BoeCin led to delimitation of *Rdv1* to a region between the markers GBS-04 and
GBS-11 (Fig. 5B-D, Supplementary File 2 Table S8). The GBS markers were used to
precisely locate recombination sites that were identified between SSR markers. Since
the exact positions of the GBS markers are unknown for PN40024 and VitRGM, the
markers GF13-04.2 and GF13-26, located very close to the markers GBS-04 and
GBS-11, were used to specify the interval size.

Fig. 5 Integration of genetic map and haplotype-specific physical map of the
***Rdv1* region on pseudochromosome 13. (A)** Dot plot of the *Rdv1* region between
BoeRip and BoeCin. Genetic markers mapping on BoeRip are represented as red
arrows and markers mapping on BoeCin as blue arrows. The *Rdv1* locus is located

between GBS-04 and GBS-11. (**B-D)** Fine mapping of the phylloxera resistance locus *Rdv1*. (**B)** Genomic region of the genetically mapped *Rdv1* locus on chromosome 13 of 'Börner'. The grey bar represents the interval between the delimiting SSR markers GF13-01 and GF13-09 [6], the numbers indicate the genetic distance between adjacent markers in cM. **(C)** Physical position of the SSR and GBS markers used for fine mapping of this region. The bar depicts the enlarged region taken from (B). Numbers refer to the physical distance of adjacent markers in kbp based on BoeCin (haplotype conferring resistance). **(D)** Local map of seven relevant F1 individuals from the V3125 x 'Börner' population revealed after haplotype-specific genotyping, only the 'Börner' haplotypes are shown. F1 genotypes analysed by GBS are highlighted in bold. The phenotypes are indicated on the right side: R, resistant; S, susceptible. Red and blue bars indicate the BoeRip and BoeCin haplotypes, respectively. Regions with unlocalised recombination sites are shown in purple.

Based on the gene annotation data generated for the assemblies, the RGA studies and intensive manual curation (see Methods), the genetically delimited *Rdv1* locus was evaluated in detail at the gene level (Fig. 6). The locus has a size of 223,738 bp with 20 genes and eight potential resistance genes in BoeCin, a size of 402,698 bp with 25 genes and 13 resistance genes in BoeRip, a size of 202,484 bp with 15 genes and seven resistance genes in PN40024 and 208,610 bp with 16 genes and four resistance genes (all detected as RGAs) in VitRGM. Complete gene structures were counted as "normal" genes even if the gene belongs to a TE. The *Rdv1* locus is covered by a single contig in both 'Börner' haplotypes.

A resistance gene cluster that displays a very divergent structure and different gene content among the three analysed haplotypes and the pseudo-haplotype VitRGM is present between the cytokinin dehydrogenase encoding gene (dark blue in Fig. 6, annotation ID *BoeCin13g18380*, tagged by GF13-04.2/GBS-04) and the BYPASS1-related gene (purple in Fig. 6, annotation ID *BoeCin13g18400*, tagged by GF13-26/GBS-11). The haplotype conferring dominant resistance (BoeCin) contains eight potential resistance genes (Supplementary File 2 Table S9). Three of them, located in the southern part of the *Rdv1* locus (*BoeCin13g18393*, CC-NBS-LRR class; *BoeCin13g18396*, TM-CC class; and *BoeCin13g18399*, TM-CC class), are located in a syntenic region that contains similar alleles of the genes detected in all four haplotypes (see Discussion). In the following, the distinction between true haplotype sequences (PN40024, BoeRip and BoeCin) and the pseudo-haplotype VitRGM, that may contain merged sequences from the two phases of *V. riparia* Gloire de Montpel-

lier, is neglected. Due to presence of these three genes in susceptible genotypes with

highly similar alleles, they were considered not to be relevant for phylloxera resis-

tance. The remaining five potential resistance genes (*BoeCin13g18381*;

BoeCin13g18382, CC-NBS-LRR/RLP; *BoeCin13g18388*; *BoeCin13g18389,* NBS-

LRR; *BoeCin13g18390*) can be considered as candidate genes for causing phylloxera

resistance. *BoeCin13g18381, BoeCin13g18382* and *BoeCin13g18389* are putative

disease resistance genes encoding RPP13-like proteins. The other two

(*BoeCin13g18388*, *BoeCin13g18390*) are similar to the *Arabidopsis thaliana* resis-

tance gene *AtLRRAC1*. To visualise the similarities between the protein-coding genes,

especially the putative disease resistance genes among and within the haplotypes, a

similarity matrix was calculated between BoeCin and itself as well the three other

haplotypes (Supplementary File 2 Table S10 to S13). The similarities detected con-

firm the relations mentioned above, including the relatedness of, for example, the

similarity of the putative disease resistance proteins encoded by *BoeCin13g18388* and

BoeCin13g18390.

Fig. 6 Genes of BoeCin, BoeRip, PN40024 and VitRGM at and surrounding the

***Rdv1* locus.** The genes (alleles) were clustered according to similarity and are shown

in one colour for each orthogroup. The grey box represents the *Rdv1* locus delineated

by the genetic markers GF13-04.2/GBS-04 and GF13-26/GBS-11; additional markers

are included upstream and downstream of the *Rdv1* locus. The kbp values on the axis

were adapted such that the start of the first gene is bp zero (relative coordinates). At

the top, the colour code for each group of related genes/alleles and the corresponding

functional annotation is given. Grey coloured genes are potential resistance genes

according to their functional annotation; genes with a red edging were identified as

RGA (see text). Above each RGA, the domain type classification of the encoded

protein is mentioned. See legend to Fig. 4 for acronyms, CC-NBS-LRR abbreviated as
CNL, NBS-LRR as NL. Three genes qualified as TE genes of different types, these
are marked with roman numbers; I: mutator-like element (MULE), II: similar to
transposon TX1 protein, III: similar to retrovirus-related polymerase polyprotein.

Discussion

Final 'Börner' genome sequence assembly

For the fully phased genome assembly, the sequence reads were binned into parental

subsets prior to assembly. The binning process produced two main read sets, both

containing approximately half of all reads. An additional small read set with relatively

short reads that were probably derived from homozygous regions was also generated.

The binning process provided an excellent basis for generating high quality, haplo-

type-separated genome assemblies, with sufficient sequence coverage. Two fully

phased genome assemblies of the grapevine cultivar 'Börner' were computed. About

90 % of all bases were included in contigs equally or larger than 530 kbp (BoeRip)

and 450 kbp (BoeCin). In contrast to other published grapevine genome assemblies,

most of them generated with FALCON and FALCON-Unizp, the genome assembly

size was not over- or underestimated by Canu and thus reaches for both haplotypes

approximately 500 Mbp (Supplementary File 2 Table S2). As noted before [14, 19, 24],

FALCON-Unzip sometimes overestimates the primary contigs and underestimates the

haplotigs because it orders some alternative sequences of heterozygous regions to the

primary contigs. Regarding genome assembly, TrioCanu/Canu [28, 30] turned out to be a

very good choice for generating the genome sequence assembly BoeRC of the inter-

specific grapevine cultivar 'Börner'. The N50 value of both 'Börner' haplotype as-

semblies outreaches most of the other *Vitis* genome assemblies. Comparing the N50,

QV and overall k-mer completeness with some available chromosome level *Vitis* as-

semblies, BoeRC has with distance the highest N50 value, the highest quality values

(estimated accuracy of 99.9998 %) and the highest degree of completeness (98.97 %)

(Supplementary File 2 Table S2). To further demonstrate that the 'Börner' haplotype

assemblies are fairly complete, the assemblies were investigated for the plant core

genes. As in other high-quality genome sequence assemblies of grapevine like that of

Cabernet Sauvignon [20], more than 98 % of complete core genes (BUSCOs) were iden-

tified.

Phasing

The haplotype assemblies can be considered as fully and correctly phased. Through mapping of phased BAC sequences derived from parts of chr01 and chr14 of 'Börner', it was shown that several Mbp of chr01 and chr14 are truly phased. The haplotype-specific BAC sequences coincide with the correct haplotype and share plenty of mismatches with the other haplotype. Also, the mapping validated correctness of contiguity and chromosome assignment in those regions.

Additionally, the phasing was analysed through k-mer analysis with Merqury. The k-mer completeness of BoeRC over both haplotypes was rated to 99.04 % (BoeRip) and 99.51 % (BoeCin), while haplotype precision was estimated to 99.07 % (BoeRip) and 99.92 % (BoeCin) (Supplementary File 2 Table S2). Thus, the haplotype assemblies can be considered as completely and correctly phased. The purity of the haplotypes was also expressed through the very low switch error rate of 0.046 % (BoeRip) and 0.035 % (BoeCin) and the high block N50 value of 2.25 Mbp (BoeRip) and 2.42 Mbp (BoeCin). None of the scaffolds showed a notable amount of k-mers from the other haplotype (Supplementary File 1 Fig. S2).

There would potentially be room for improvement of the assemblies BoeRip and BoeCin by methods like chromosome conformation capture sequencing (Hi-C). Even if there are indications that the organization of chromosomes into territories within the nucleus of diploid organisms could allow to extract data supporting correct haplotype separation [31], we did not follow this route and anticipate that the quality of the BoeRC assembly is sufficient for almost any application with relevance to viticulture.

Similarity of 'Börner' haplotypes and comparison with PN40024 and VitRGM

The assemblies BoeRip and BoeCin show overall similarity to each other and to PN40024. Even if the haplotypes among themselves have almost the same SNP frequency as with PN40024, more precisely 1 SNP in 33 bp compared to 1 SNP in 31 bp, more bases were aligned between the haplotypes with a higher identity than with PN40024. The difference in SNP frequency becomes more specific when comparing the SNP frequency of the coding regions. Here, BoeRip and BoeCin show less SNPs (1/1,001 bp) than with PN40024 (1/932 bp BoeRip & 1/935 bp BoeCin). Because of the high heterozygosity between haplotypes of grapevine [32], it is not surprising that the SNP frequency between 'Börners' haplotypes is almost the same as to another

Vitis species. Reported SNP frequencies for grapevines range from 1 SNP per 64 bp [33]

to 1 SNP per 200 bp [17]. The SNP frequency between *V. riparia* and *V. vinifera* was

determined to 1 SNP per 78 bp [34]. Generally, it seems that SNP frequencies between

and within *Vitis* species are not that different. However, the SNP frequency between

BoeRip and VitRGM, both representing *V. riparia* genome sequences, was signifi-

cantly lower than between BoeCin and VitRGM, for coding as well as non-coding

regions. This may indicate that *V. riparia* GM183 and *V. riparia* Gloire de Montpel-

lier were derived from related populations, or that the species *V. riparia* generally

covers less variation.

High sequence identities and a good chromosome assignment were revealed through

dot plots between homologous pseudochromosomes of the haplotypes of BoeRC and

PN40024 or VitRGM. The "point clouds" often observed in the dot plots between

homologous pseudochromosomes can be assumed to pinpoint the location of the cen-

tromere as centromeric regions are often highly repetitive and thus hard to sequence

and assemble. The gaps detected in comparison with PN40024 may represent missing

bases in the PN40024 genome sequence assembly. Almost all identified rearrange-

ments between the haplotypes of BoeRC and PN40024 were found between both hap-

lotypes and PN40024, and most of them are inversions. This calls for an improvement

of the 'PN40024' reference genome sequence based on long read sequence data.

Rdv1 locus

The *Rdv1* locus has been genetically mapped to a region of 224 kbp in size in the

BoeCin haplotype of the 'Börner' genome. The genome regions orthologous to the

Rdv1 locus from BoeCin and different *Vitis* species are flanked on both sides by large

syntenic regions. The high variability and divergence that is known for resistance

gene clusters [35] was also detected within the *Rdv1* locus.

In syntenic regions, alleles may vary significantly in lengths between haplotypes, still

they encode the same protein even among different *Vitis* species. For example, the

PN40024 gene *Vitvi13g01613* which encodes a putative glycerol-3-phosphate dehy-

drogenase (dark blue gene in Fig. 6, northern part of the locus), is 3,600 bp longer

than its BoeCin ortholog. The 466 amino acid long protein sequence is identical be-

tween all haplotypes analysed except one amino acid substitution each in PN40024

and BoeRip. Increased gene length of *Vitvi13g01613* in PN40024 is caused by a lar-

ger third intron. Several TE fragments and especially one larger LTR/Copia-like TE

explain the difference in the length of intron three.

In addition to SNPs, quite some presence/absence variation (PAV) contributes to the

divergence of the *Rdv1* region in *Vitis*. The various PAVs were, at least in several

cases, explained by detection of TEs (either DNA transposons or retrotransposons) in

the haplotypes (Fig. 6, Supplementary File 1 Fig. S11). For example, two intact

LTR/Copia TEs and two intact DNA/hAT TEs were detected in the *Rdv1* region of

BoeRip in the context of the tandem array of seven RGAs of the CC-NBS-LRR

(CNL) type that is only present in BoeRip. Another genic PAV is the E3 ubiquitin

ligase coding gene (*BoeRip13g18552*) that was detected in BoeRip and VitRGM

which might be specific for *V. riparia*, although this hypothesis needs to be tested

with more data from *V. riparia*. The resistance gene *BoeCin13g18388* encodes a LRR

domain and is about 2,400 bp longer in BoeCin than in BoeRip (*BoeRip13g18553*).

The reason is again an intron in the 5'UTR region of BoeCin which includes an

LTR/Gypsy TE. This TE insertion causes a shifted translation start and the protein of

the BoeCin allele is 49 amino acids longer at the N-terminus than the protein encoded

by the BoeRip allele. In case of the MULE transposon (TE with similarity to mutator-

like elements, marked with I in Fig. 6), a gene coding for a transposase protein was

identified in BoeCin, BoeRip and VitRGM. There are hints from the TE annotation

(Supplementary File 2 Table S7) that the MULE transposon might contain two ORFs

in opposite orientation which is characteristic for some MULEs. However, the termi-

nal inverted repeats were not identified. Because of the great diversity within the same

and between transposon superfamilies, generally and also with regard to the structure

and length of MULEs [36, 37], further work is needed to precisely identify all TEs and

TE fragments that contribute to PAVs at this and other loci.

The *Rdv1* region is characterized by one extended resistance gene cluster. Ortholo-

gous regions in the other studied haplotypes derived from different *Vitis* species dis-

play very divers cluster structures with respect to length in kbp and number of resis-

tance genes. In the BoeCin haplotype that confers the dominant resistance to phyllox-

era and which segregates in offspring of 'Börner', the cluster has a length of about

350 kbp and includes 10 potential resistance genes. The cluster extends at the south-

ern end beyond the genetically delimited *Rdv1* locus. In the BoeRip haplotype that

confers recessive susceptibility to phylloxera, the cluster includes 14 potential resis-
tance genes.

In both 'Börner' haplotypes as well as in VitRGM, the cluster shows an insertion of a
conserved array of four to six genes compared to PN40024 representing *V. vinifera* in
the comparison. The inserted genes (or the genes deleted in PN40024) encode a tRNA
ligase, a protein of unknown function, a mutator-like element (MULE) DNA transpo-
son, a DEAD-box ATP-dependent RNA helicase, a ribonucleoside-diphosphate re-
ductase small chain and a ubiquitin ligase. These genes display no features of resis-
tance genes and are found with high identity values in the susceptible haplotype
BoeRip. They are, therefore, no candidate genes for *Rdv1*.

Within the southern part of the locus, RGAugury detected two RGAs with TM-CC
domains in the encoded proteins that are potential false positive results (orange and
pink in Fig. 6, with red edging in BoeRip and BoeCin due to the RGAugury output).
Based on the functional annotation (Supplementary File 2 Table S9), the two or-
thogroups code for homologs of SEC31A (a component of the coat protein complex II
(COPII) involved in the formation of transport vesicles) and homologs of AGL15
(agamous-like MADS-box factor potentially involved in control of development). The
genes in the syntenic block that starts at its northern end with the homolog of *SLX1*
(encoding a subunit of a structure-specific endonuclease complex involved in process-
ing diverse DNA damage intermediates, light ochre in Fig. 6) and which extends to
the south beyond the delimiting markers GF13-04.2/GBS-04, are also very unlikely to
be candidates for *Rdv1*.

To the north, the synteny extends for two more genes and terminates at a TE gene
annotated as TX1-like non-LTR retrotransposon (*BoeCin13g18391*, marked II in Fig.
6) that is only detected in BoeCin. Southern to this TX1-like TE, a syntenic or-
thogroup (including *BoeCin13g18393* in BoeCin) encodes a CC-NBS-LRR (CNL)
resistance protein related to the wild potato resistance protein RGA4. Since this gene
is represented by closely related alleles in BoeCin and the haplotypes conferring sus-
ceptibility (amino acid sequence identity: BoeRip 99.9 %, PN40024 97.1 %), it is not
considered to be a candidate for *Rdv1*.

Based on this evaluation, the most promising genes mediating phylloxera root resis-
tance are the five resistance genes northern and southern to the array of four to six
genes conserved in BoeCin, BoeRip and VitRGM (*BoeCin13g18381*;

BoeCin13g18382, CNL/RLP; *BoeCin13g18388*; *BoeCin13g18389,* NL;

BoeCin13g18390). Ideally, an F1 genotype from the V3125 x 'Börner' cross with a

recombination site between the BoeCin and BoeRip haplotypes should have been

identified. However, such an event was not detected. It is possible that the high diver-

gence and sequence dissimilarity throughout the *Rdv1* locus causes a suppression or at

least reduction of recombination events at the locus, which could explain this failure.

A reduction of recombination frequency was, for example, also described for the

Rpv3.1 locus that confers resistance to *P. viticola* in *Vitis* [38].

The five remaining resistance genes can be divided into two types with respect to

functional annotation. The resistance genes *BoeCin13g18388* and *BoeCin13g18390*

were not detected as RGAs by RGAugury, but encode proteins similar to the *A.*

thaliana resistance gene *AtLRRAC1* (*At3g14460*). *AtLRRAC1* has been reported to be

involved in the defence against the biotrophic fungus *Golovinomyces orontii* as well

as the hemi-biotrophic bacteria *Pseudomonas syringae*, and might be relevant for sig-

nalling via cAMP-dependent defence pathways [39]. The three other resistance genes,

namely *BoeCin13g18381*, *BoeCin13g18382* CC-NBS-LRR/RLP and

BoeCin13g18389 NBS-LRR (NL), encode proteins similar to *RPPL1* (Recognition of

Peronospora Parasitica 13-Like 1). *AtRPPL1* (*At3g14470*) received its annotation

from *AtRPP13* (*At3g46530*), a gene that confers resistance to the biotrophic oomycete

Peronospora parasitica and which encodes an NBS-LRR protein [40, 41]. In *A. thaliana*,

AtLRRAC1 and *AtRPPL1* form a cluster of two neighbouring genes, a feature that is

conserved in *Vitis* (Fig. 6). It will be interesting to figure out if the conserved co-

localisation of two different resistance genes has functional implications for resistance

to phylloxera in *Vitis* species.

In BoeCin and BoeRip, one of the resistance genes is classified as encoding an NBS-

LRR (NL) protein. Despite an identical NBS domain with only very few substitutions,

the LRR region varies greatly between the two alleles (*BoeCin13g18389* and

BoeRip13g18554) and is shorter in BoeRip because of an early stop codon. Also for

the other four candidate resistance genes, differences between the alleles in BoeCin,

BoeRip and PN40024 were detected. For example, the protein encoded by

BoeCin13g18382 was detected as RGA based on CC-NBS-LRR domains in addition

to features of an RLP. However, the genes detected at syntenic positions (alleles) of

BoeRip and PN40024 lack these features either in part or fully.

Since the PN40024 sequence is derived from the most susceptible line in our compari-
son, the protein sequences of the five *Rdv1* candidate resistance genes of BoeCin were
compared with all PN40024 protein sequences. No identical PN40024 protein se-
quence or proteins with identical resistance domain sequences were identified. Thus,
the five BoeCin resistance genes are promising candidate genes for conferring resis-
tance to phylloxera.

Conclusions

The fully phase-separated genome sequence assembly BoeRC from the phylloxera-
resistant rootstock 'Börner', and its structural and functional gene annotation, build a
cornerstone for the investigation of loci from 'Börner' that are linked to various valu-
able traits. Here, the focus was on phylloxera resistance. The sequences of the haplo-
types BoeCin conferring resistance to phylloxera and BoeRip conferring susceptibility
allowed to precisely map recombination sites by GBS in crucial genotypes of a map-
ping population of V3125 x 'Börner'. Subsequently, detailed examination and com-
parative genomics of the gene content of the *Rdv1* locus allowed to delimit the locus
to a region of 123 kb in BoeCin haplotype with five resistance genes as potential can-
didates involved in mediating phylloxera resistance at the root. The resources gener-
ated will allow to study additional traits and have the potential to support resistance
breeding for the benefit of viticulture.

Methods

Reference data sets

In   this   study,   the   'PN40024'   genome   sequence   [16]   assembly   12X.v2
(urgi.versailles.inra.fr/Species/Vitis/Data-Sequences/Genome-sequences) and its an-
notation,   the   VCost.v3   gene   annotation
(https://urgi.versailles.inra.fr/Species/Vitis/Annotations) [42] as well as the *V. riparia*
Gloire de Montpellier genome sequence assembly VitRGM and its annotation
(https://ncbi.nlm.nih.gov/assembly/GCF_004353265.1) [24] were used for comparisons.

Plant material and DNA and RNA extraction

'Börner' is listed in the *Vitis* International Variety Catalogue with the variety number *VIVC* 1499, its parents *V. riparia* GM183 and *V. cinerea* Arnold with variety numbers *VIVC* 4686 and *VIVC* 13645, respectively. All three cultivars do not belong to an endangered species and were obtained and are grown at the Institute for Grapevine Breeding Geilweilerhof at Siebeldingen (JKI Siebeldingen) in accordance with German legislation.

Dormant wood cuttings of the interspecific rootstock variety 'Börner' were raised in a growth chamber on soil at 18°-24°C (64°F-75°F), 70 % relative humidity and 16 h light and 8 h darkness. Prior to harvest the plants were cultivated in the dark for 72 hours. Small, young leaves were harvested and stored on ice until use. Extraction of high molecular DNA for SMRT sequencing was performed starting with 1.5 g fresh weight and was carried out with the NucleoSpin Plant II Maxi kit for DNA (Macherey and Nagel, Düren; Germany) according to the recommendations of the manufacturer for plants. DNA integrity, quality and quantity was checked using a TapeStation (Agilent) and was found to peak at a fragment size higher than 48.5 kbp.

Leaf material of 'Börners' parents was harvested from cuttings grown in the green house, frozen in liquid nitrogen and stored at -80°C. The DNA extraction was carried out with the DNeasy Plant Maxi Kit (Qiagen, Hilden; Germany) according to the instructions of the manufacturer. Tissue samples of leaves and tendrils of 'Börner' were collected from grown in the field. Total RNA was isolated using the Spectrum Plant RNA-Kit (Sigma, Taufkirchen; Germany).

Plant material for the local map of *Rdv1* was selected from the previously described mapping population V3125 x 'Börner' [6, 43] and another set of 310 F1 genotypes obtained after repeating the cross in 2006. To search for recombinant F1 lines, SSR markers associated with *Rdv1* (Supplementary File 2 Table S14) were used for genotyping as described [43]. For GBS, young leaf material was harvested from selected individual F1 genotypes as well as from the maternal genotype V3125 and used for DNA extraction. Genomic DNA was prepared using an established CTAB-based protocol [44]. Subsequently, the DNA obtained was treated with RNAse and quantified using PicoGreen.

Library construction for 'Börner' genome and transcriptome sequencing

PacBio Sequel libraries were prepared according to the SMRTbell Template Prep Kit 1.0 and then size-selected with BluePippin for a target insert size of greater than 10 kb. Seventeen 1Mv3 SMRT cells were run on a Sequel I sequencer using the Binding Kit 1.0 and the Sequencing Chemistry version 1.0 (all from PacBio) for six hours per cell. The output BAM files of the sequencing step were loaded into SMRT Link 5.0.1 according to the recommendation of the manufacturer (PacBio Reference Guide 2018) and converted to FASTA format for downstream processing. The length distribution over all reads was calculated with the assembler Canu [30] and is provided in Supplementary File 1 Fig. S12.

The RNA samples from leaves and tendrils of 'Börner' were paired-end sequenced using Illumina technology on a HiSeq-2000 essentially as described [45]. The new RNA-Seq read data (ENA/SRA study accession no. PRJEB46079) were processed together with data submitted before [45]. These already existing RNA-Seq reads of 'Börner' (PRJEB34983) cover leaves (ERR3894001), senescent leaves (winter leaves, ERR3895010), inflorescences (ERR3894002), tendrils (ERR3894003) and roots (ERR3895007). Read data were trimmed with Trimmomatic-0.38 [46] allowing reads larger than 80 nt (new data) and 90 nt (published data) to be kept. The trimming statistics of the RNA-Seq data is provided in Supplementary File 2 Table S15.

Illumina sequencing of the parental genomes of 'Börner'

Library preparation for the *V. riparia* GM183 and *V. cinerea* Arnold samples was performed according to the Illumina TruSeq DNA Sample Preparation v2 Guide. Genomic DNA (1500 ng each) was fragmented by nebulisation. After end repair and A-tailing, individual adaptors were ligated to the fragments for PE sequencing. The adaptor ligated fragments were purified on a 2 % low melt agarose gel and size selected by excising a band ranging from 500-800 bp. After enrichment PCR of DNA fragments that carry adaptors on both ends, the final libraries were quantified by using the Quant-iT PicoGreen dsDNA assay on a FLUOstar Optima Platereader (BMG Labtech) and qualified on a BioAnalyzer High Sensitivity DNA Chip (Agilent). The *V. riparia* GM183 library was sequenced on a MiSeq PE run (2 x 250 nt) and on two lanes of a 2 x 100 PE run on a HiSeq-1500 in high output mode. The *V. cinerea* Arnold library was sequenced in one 2 x 150 bp PE run on a HiSeq-1500 in rapid mode.

624 The read data were submitted to ENA (ENA/SRA study accession no. PRJEB45595).

625 All genomic short read data were quality trimmed with Trimmomatic-0.36 [46] allowing

626 reads equal or longer 80 nt to be kept. The trimming statistics is provided in Supple-

627 mentary File 2 Table S16.

Sequencing of pools of BACs from a BAC library of 'Börner'

For the creation of phased sequences of 'Börner', 8 pools of BAC constructs contain-

ing a total of 440 mapped BACs from two loci (one on chr01, the other on chr14)

were selected from a 'Börner' BAC library [10]. The BACs were selected and distrib-

uted to the pools according to the positions of BAC end sequences on the correspond-

ing regions of PN40024. Mapping to the reference assembly was carried out with the

CLC Genomics Workbench v8.0 toolkit (https://www.qiagenbioinformatics.com/).

Pools were arranged by excluding overlap of BAC inserts to allow assembly of indi-

vidual BAC insert sequences. The BAC Pools1-4 were sequenced on a 454 Life Sci-

ences Genome Sequencer GS FLX and PE (2 x 250 nt) on the Illumina MiSeq plat-

form. The BAC Pools5-8 were sequenced on the MiSeq platform only. The read data

were submitted to ENA (ENA/SRA study accession no. PRJEB46081). The MiSeq

raw data were quality trimmed with Trimmomatic-0.32 using 'ILLUMINACLIP:

2:40:15 LEADING:3 TRAILING:3 SLIDINGWINDOW:4:15 MINLEN:36' [46]. The

trimming statistics for the BAC pools is provided in Supplementary File 2 Table S5.

Library construction and sequencing for GBS

The barcoded libraries were prepared using Illumina library preparation kits using

TrueSeq technology and sequenced on a NextSeq500. Sequencing was performed in

150 nt single-end (SE) modus and the read data were submitted to ENA. The

ENA/SRA study accession no. is PRJEB53997. Information on genotypes, sequenc-

ing strategy, data volume and run IDs are summarized in Supplementary File 2 Table

S17.

Phenotyping of phylloxera resistance

The test system used to assess phylloxera resistance of selected F1 individuals was

essentially described previously [6, 47]. The phylloxera population used for the artificial

inoculation of potted plants was derived from leaves of naturally infested rootstock

cultivars from the germplasm repository at JKI Siebeldingen. The phylloxera resis-

tance tests started at the beginning of the vegetation season when enough infested leaf

material was available as inoculum. Leaf pieces with 8 to 12 (3-5 mm) galls contain-

ing mature eggs were inserted in the soil of rooted cuttings of the F1 individuals. The

inoculation procedure was repeated after 3 to 4 weeks. Approximately 4 weeks after

the second inoculation the plants were rated as susceptible if nodosities were visible,

or resistant if the roots developed normally. Each F1 genotype was tested in triplicate

and the assays were repeated at least twice.

Diploid genome sequence assembly

To compute a phased genome assembly, a binning and assembly approach was used.

The binning prior to assembly was performed with the tool TrioCanu of the assembler

Canu v1.7 [28, 30]. TrioCanu uses short reads of the parents and long reads of the F1 as

input. It bins the long reads of the F1 into parental subsets based on k-mer compari-

sons. The binning step was performed on a compute cluster using TrioCanu default

parameters (k-mer size of 20). Binning resulted in one read subset for each parent and

an additional small subset with reads that could not be assigned to either of the two

parental haplotypes, referred to as UAR subset. In the following, the parental subsets

are referred to as Vrip and Vcin read subset according to 'Börners' parents and the

two different haplotypes combined into 'Börner', respectively. The UAR subset con-

tained only 0.13 % of all bases with an average read length of 1.5 kbp.

After binning, the read subsets were used to compute the two haplotype assemblies

BoeRip and BoeCin of 'Börner' with the assembler Canu v1.7 [30]. To compute the

BoeRip haplotype assembly, the Vrip and the UAR subsets were used in order to in-

clude identical and/or homozygous regions. Likewise, the BoeCin haplotype assembly

was generated from the Vcin read subset and the UAR subset. Therefore, the basis of

the haplotype assemblies, BoeRip and BoeCin, are approximately 32 to 34 Gbp read

data with an estimated genome coverage of more than 60-fold each. For both sepa-

rately computed haplotype assemblies, Canu was run on a compute cluster utilizing

the parameters 'genomeSize=550m', 'corMhapSensitivity=normal', 'correctedError-

Rate=0.065, 'canuIteration=1' and 'stopOnReadQuality=false'. During the assembly

processes about 2.5 % of the corresponding input data (bp) remained unassembled. A

search with blastn of the Basic Local Alignment Search Tool (BLAST) package+

v2.8.1 [48, 49] and the grapevine chloroplast [50] and mitochondrial sequences [51] was car-

ried out on the haplotype assemblies and detected contigs were removed. Finally, the

haplotype assemblies were polished twice with *arrow* v2.2.2 (smrtlink-

release_5.1.0.26412) and the corresponding read subsets. Assembly statistics were

computed with QUAST v4.6.3 [52].

Scaffolding with 'Börner' BAC end sequences

To validate the quality of the two genome sequence assemblies, and also to scaffold a

few smaller contigs, high-quality paired BAC end sequences from the 'Börner' ge-

nome (ENA/GenBank accession numbers KG622866 - KG692309 [10]) were used. The

69,444 BAC end sequences cover 39,360,203 bp, have an average sequence length of

566 bp, and represent an eight-fold coverage of the 'Börner' genome based on aver-

age BAC insert length. First, the 62,498 BAC end sequences that represented pairs

from one BAC were mapped to BoeRip and BoeCin with HISAT2 v2.1.0 [53] using the

parameters '--maxins 2000000000', '--secondary', '--no-softclip' and '--no-spliced-

alignment'. Alignments with a mapping quality >10 were filtered out and alignment

statistics were collected with SAMtools v1.8 [54]. After mapping and quality filtering,

the BAC end sequences were used to scaffold the haplotype assemblies with SSPACE

(standard) v3.0 [55]. SSPACE was run with the parameters '-x 0', '-m 50', '-k 3', '-n

500' and '-T 10' and by using BWA-SW [56] for mapping. Contig extension was dis-

abled ('-x 0') as the BAC end sequences are not phased and thus would eventually

contaminate the haplotypes with a sequence from the other haplotype. However, a test

run with enabled extension showed that the BAC end sequences were anyway not

suitable for gap filling.

Scaffold to pseudochromosome assignment

To build pseudochromosomes (the nucleotide sequence representation of chromo-

somes), an *ad hoc* gene prediction was generated with AUGUSTUS v3.3 [57]. Prior to

gene prediction, the RNA-Seq reads from 'Börner' (see above) were processed to cre-

ate a *de novo* transcriptome assembly with Trinity v2.8.5 [58] with default settings. Hint

files with information about exon and intron positions for the haplotype assemblies

were created through mapping all trimmed RNA-Seq reads and the transcriptome as-

sembly on the haplotype assemblies. The RNA-Seq reads were mapped with HISAT2

v2.1.0 [59] and soft-clipping disabled ('--no-softclip'). The transcriptome sequences

were mapped with BLAT v36x2 [60] and the parameters '-stepSize=5' and '-
minIdentity=93'. Hints were generated according to the AUGUSTUS documentation.
Additionally, AUGUSTUS was trained with the CRIBI v2.1 gene annotation of
PN40024 [17] to retrieve parameter sets for grapevine. Finally, genes were predicted on
both haplotype assemblies BoeRip and BoeCin with the generated *Vitis* parameter set
and the hint files.

The protein sequence of the primary transcript variant was extracted from the
PN40024 CRIBI v2.1 annotation and from the *ad hoc* annotation of BoeRip and
BoeCin. Reciprocal best BLAST hits (RBHs) between the protein sets of each of both
haplotypes and the filtered proteins of PN40024 were computed with blastp utilizing
an e-value threshold of 0.0001. For RBHs, two rounds of BLAST searches were per-
formed. Here, one round was run with the proteins of one haplotype as input query
and one with the PN40024 proteins as input query. Only RBHs with a percentage of
identical matches and a query coverage per subject ≥80 % for at least one direction
(e.g. PN40024 protein sequences as input query) were kept. Based on RBHs encoded,
the scaffolds were assigned to the pseudochromosomes. Scaffolds with less than 10
RBHs in total and scaffolds that contain a significant number of RBHs linking a dif-
ferent pseudochromosome (30 % distance to the 2nd rank in terms of RBH numbers
required) were filtered out for later assignment or moved to the chromosomally unas-
signed part of the respective assembly.

Additionally, reciprocal hits between the scaffold sequences and the pseudochromo-
some sequences of PN40024 were computed with blastn. The e-value, identity and
query coverage filters remained the same as above. The nucleotide RBHs were itera-
tively computed and scaffolds assigned to pseudochromosomal positions. If the classi-
fications based on protein and nucleotide level contradicted, the protein RBH classifi-
cation was preferred.

To further refine the pseudochromosomes of the haplotypes, an additional assignment
based on VitRGM was performed. Also, reciprocal hits between the scaffold se-
quences of one haplotype and the pseudochromosomes of the other haplotype and *vice*
versa were calculated as described above. After manual construction of pseudochro-
mosomes for the BoeRip and BoeCin phases, the refinement with the corresponding
other haplotype was repeated until no further scaffolds could be assigned. The pseu-
dochromosomes were constructed with gaps of 100 bp length between the scaffolds

and with 10 kbp as minimal scaffold length. Due to the length filter, 61 (BoeRip) and
22 (BoeCin) scaffolds were discarded.

The orientation and order of the scaffolds on the pseudochromosomes was verified
with dot plots and manually adapted if necessary. Thus, DNAdiff v1.3 of the MUM-
mer package v4.0.0beta2 [61] was run pairwise on homologous pseudochromosomes of
BoeRip or BoeCin and PN40024 with default settings and the resulting 1-to-1 align-
ments were visualized with mummerplot v3.5.

The pseudochromosomes were validated with 340 simple sequence repeat (SSR) ge-
netic markers from [43] and the 4 SSR markers ATP3 [62], Gf13_11a, VMC8E6 [6] and
Gf14-42 [8] (Supplementary File 2 Table S14). Primer sequences were mapped to the
pseudochromosomes with primersearch of the EMBOSS v6.6.0.0 package [63] and with
a blastn search. A total of 314 markers were assigned to unequivocal sequence posi-
tions, the remaining markers do not map, show non-evaluable multi-mappings or map
with too many mismatches (Supplementary File 2 Table S18). Several markers did
map only to one of the haplotypes (16 to only BoeRip and 30 to only BoeCin, see
Supplementary File 2 Table S19). Emerging disagreements between the marker posi-
tion on PN40024 and the BoeRip and BoeCin phases were further investigated
through all-versus-all dot plots computed with the webtool D-Genies v1.2.0 [64] and by
using long read mapping to detect sequence positions that are not or only very weakly
supported by continuously mapping reads. The haplotype specific read subsets to-
gether with the unassigned read subset were mapped with minimap2 [65] v2.17 ('-ax
map-pb --secondary=no') to the corresponding haplotype assembly and the coverage
values per base were calculated with SAMtools. The haplotype assemblies are both
covered at about 54x (Supplementary File 1 Fig. S13). The back-mapping results of
the read subsets were consulted to reveal miss-assemblies if indicated by a genetic
marker. A sequence location was further investigated if five reads end in a 10 bp re-
gion and if the read coverage five bp around the region drops to ≤ five.

Conflicting marker VVMD28 maps on chr03 of PN40024 and BoeCin, yet on chr13
of BoeRip. However, eight markers allocate the sequence to chr13 of BoeRip and
neither a significant breakpoint was found in the read mappings nor in the all-versus-
all dot plot (Supplementary File 1 Fig. S3). VCHR16B maps on chr16 of PN40024
and BoeCin, but on chr03 of BoeRip. Here, one SSR marker assigns the sequence to
chr03 and no significant breakpoint was found in the read mapping or in the all-

versus-all dot plot. The three SSR marker VVS4, VMC5G6.1 and UDV-126 map on

chr08 of PN40024 and BoeRip, but on chr04 of BoeCin. The corresponding sequence

was assigned to chr04 according to 376 protein RBHs, because only 141 protein

RBHs assign it to chr08. Moreover, three SSR markers allocate the sequence to chr04,

too. Through investigating the all-versus-all dot plot, an approximately four Mbp

large alignment between chr04 and chr08 was found. However, since no position for a

split was detected in the read mappings, the sequence remained on chr04 (Supplemen-

tary File 1 Fig. S6). GF10-06, VMC3D7 and UDV-073 mapped wrongly on chr02 of

BoeRip instead of chr10; in this case the investigation of the respective contig resulted

in a split of the sequence and a corrected assignment of the contig fragment in ques-

tion. Due to 348 protein RBHs the sequence was initially assigned to chr02. The se-

quence showed the second most protein RBHs (171) with chr10. On marker side, 13

markers assign the sequence to chr02 and only three to chr10. However, the all-

versus-all dot plot showed a large alignment between chr02 of BoeRip and chr10 of

PN40024 (Supplementary File 1 Fig. S14), and the read mappings indicated a break-

point position with a coverage drop to ≤ 5. Thus, the 12.3 Mbp sequence was split into

a 7.7 Mbp large sequence that was assigned to chr10 and a 4.6 Mbp large sequence

that remained at chr02. Another case was detected in the all-versus-all dot plots be-

tween chr13 of BoeCin and chr03 of PN40024 (Supplementary File 1 Fig. S6). The

corresponding sequence was assigned to chr13 through 485 protein RBHs. Only 36

protein RBHs would allocate this sequence to chr03. Furthermore, 10 SSR marker

support the assignment to chr13 and none to chr03. No coverage drop was found in

the read mappings, the sequence remained on chr13 of BoeCin. Finally, only five ge-

netic markers remained unresolved (VVMD28, VCHR16B, VVS4, VMC5G6.1,

UDV-126).

To estimate the completeness of the assemblies, the plant core genes were localized

with the program Benchmarking Universal Single-Copy Orthologs (BUSCO) v5.1.2

utilizing the database 'eudicots_odb10' (2,326 genes) [66, 67]. For comparison, the

BUSCOs of PN40024 and of VitRGM were also determined with identical parame-

ters. The pseudochromosome lengths were visualized with cvit v1.2.1 [68].

Validation of phasing

To validate the phasing of the 'Börner' haplotypes, the BAC sequence data of pool1-4 were *de novo* assembled with Newbler v2.6. The data of pool5-8 were assembled with CLC Genomics Workbench 8.0 using a word size of 30. Remaining vector sequences were removed with vectorstrip of the EMBOSS package v6.2. Assembled sequences shorter than 500 bp were discarded. The short reads of the parents *V. riparia* GM183 and *V. cinerea* Arnold were mapped with CLC's map reads to reference algorithm to the assembled BAC sequences (linear gap cost, length & similarity fraction 1.0). Based on these mapping results regarding read coverage and percent of covered bases, the BAC sequences were assigned to either *V. riparia* GM183 or *V. cinerea* Arnold. BAC sequences with no read coverage were discarded. The assembled and phase-separated BAC contig sequences are available at https://doi.org/10.4119/unibi/2962639.

The 'Börner' BAC sequences with known haplotype allocation were mapped against the BoeRip and BoeCin haplotype assemblies with minimap2 v2.17 [65], sorted with SAMtools and the mappings visualized with the Integrative Genomics Viewer (IGV) v2.5.2 [69, 70]. The number of base calls of the aligned BAC sequences on each haplotype were determined with SAMtools' mpileup. Continuously and correct aligned BAC sequences show high base calls and low numbers of SNP/InDels (small nucleotide polymorphism, insertion-deletion polymorphism) to one of the two haplotypes and the opposite result with the alternative haplotype.

k-mer based reference-free assembly evaluation with Mercury

Merqury v1.3 [71] was used to evaluate the diploid BoeRC assembly. First, the best k-mer size was determined with Merqurys best_k.sh script for the expected 500 Mbp haploid genome size resulting in 19-mers. Consequently, 19-mer databases were computed for the *V. riparia* GM183 and *V. cinerea* Arnold Illumina reads as well as for existing Illumina reads from 'Börner' [10]. With these, hap-mer (haplotype-specific k-mers as defined by Mercury [71]) databases were generated and the haplotype assemblies evaluated using the scaffolds of both haplotypes. For building of phased blocks, the parameter 'num_switch 10' and 'short_range 20,000' were applied. For comparison, Merqury was also run on PN40024, on VitRGM, on both Cabernet Sauvignon haplotype assemblies v1.1 and on both *M. rotundifolia* cultivar 'Trayshed' haplotype

assemblies v2.0 [23]. The 19-mer profiles were computed from PN40024 WGS reads

SRR5627797, from VitRGM WGS reads SRR8379638, from Cabernet Sauvignon

WGS reads SRR3346861 and SRR3346862 and from *M. rotundifolia* WGS reads

SRR6729333. Prior to analyses, the reads were trimmed with Trimmomatic-v0.39

allowing a minimum read length of 80 or 90 (PN40024 reads).

Gene annotation

The 'Börner' haplotypes were annotated *ab initio* with MAKER v3.01.03 following

the MAKER-P pipeline [72, 73]. As input data, haplotype-specific repeat libraries were

created according to the MAKER advanced repeat library protocol. The monocotyle-

dons repeat library of RepBase (RepeatMaskerEdition-20181026,

model_org=monocotyledons) [74], the transposable element (TE) sequences from

MAKER, the 'Börner' *de novo* transcriptome assembly described above, *Vitis* protein

sequences (NCBI, Protein DB "Vitis"[Organism]), plant protein sequences (UniProt,

release 2020_02, "Viridiplantae [33090]" AND reviewed:yes) and *Vitis* full-length

cDNAs (NCBI, Nucleotide DB "Vitis" [Organism] AND complete cds[Title]) were

used. The gene predictors SNAP v2006-07-28 (three rounds) [75] and GeneMark v3.60

[76] were trained on both, BoeRip and BoeCin. AUGUSTUS v3.2.3, tRNAscan-SE

v1.3.1 and EvidenceModeler v1.1.1 [77], SNAP and GeneMark together were ran

through MAKER to compute the final gene annotation. The option 'split_hit' was set

to 20,000 and the previously generated *Vitis* parameter set was adjusted for AUGUS-

TUS.

The gene models from BoeRip and BoeCin were refined with PASA v2.4.1 [78]. As

evidence data, the *de novo* and reference-guided 'Börner' transcriptome assemblies,

the *Vitis* full-length cDNAs and *Vitis* EST data (NCBI, txid3603[Organism:exp] AND

is_est[filter]) were used. The parameter file is available as Supplementary File 3 Data

S1. The refinement with PASA was iteratively applied three times on the gene mod-

els. The reference-guided assembly was computed with Trinity v2.10.0 using the

'Börner' RNA-Seq data aligned with HISAT2 v2.2.0. Only alignments with MQ >10

were given to Trinity and '--genome_guided_max_intron' was set to 20,000.

After structural gene annotation, a functional gene annotation was added through a

BLAST search (package+ v2.10.0) against the UniProt/Swiss-Prot database (release

2020_02), through domain identification with InterProScan5 v5.42-78.0 [79] and the

PFAM database v32.0. Gene models with Annotation Edit Distance (AED) > 0.5 were

filtered from the final gene model set. Also, predicted genes expected to encode a

polypeptide with less than 50 amino acids and no functional annotation were dis-

carded.   The   annotation   data   for   BoeRC   are   available   at

(https://doi.org/10.4119/unibi/2962793). For comparison, functional annotation was

also carried out for the PN40024 VCost.v3 structural gene annotation with identical

resources.

Prediction of resistance genes

Resistance gene analogs (RGAs) were predicted for both haplotypes with the pipeline

RGAugury v2.1.7 [29]. RGAugury employed ncoil [80], PfamScan v1.6 [81], InterProScan5

v5.45-80.0 [79] and Phobius-1.01 [82] on the protein sequences to predict known resis-

tance domains. The initial filtering against the RGAdb was disabled since the database

has not been updated since 2016, unknown RGAs not included in the database were of

interest and the identification of (resistance) domains for none RGAs was enabled.

For InterProScan5 domain search through RGAugury, the databases Gene3D-4.2.0 [83],

Pfam-33.1 [84], SMART-7.1 [85] and SUPERFAMILY-1.75 [86] were utilized.

The RGA annotations of the two haplotypes were correlated through determination of

RBHs based on protein sequences. Only RBHs with an identity and coverage ≥ 90

were considered.

Assembly wide variant detection

898   The SNP and InDel positions of the SNP output from the DNAdiff analysis (see

899   above) were compared with the positions of the coding and non-coding gene regions

900   and classified accordingly.

Genetic mapping

The annotated genes and nucleotide sequences of the phylloxera resistance QTL on

chr13, referred to as *Rdv1*, of BoeRip, BoeCin, PN40024 and VitRGM were com-

pared. Before this study, the QTL *Rdv1* was delimited with the associated genetic

markers GF13-01 and GF13-12 [6]; the marker positions were identified based on the

respective marker assay primer sequences with primersearch (EMBOSS package [63]).

To narrow down the location of *Rdv1*, additional genetic fine mapping was carried out

with a mapping population of in total 498 F1 genotypes (see above) and newly de-

signed sequence-tagged SSR (STS) markers. The additional markers were derived

from PN40024 [16] (Supplementary File 2 Table S14) and used to screen the F1 geno-

types from the mapping population.

Five relevant genotypes with marker data indicating recombination events within the

initial *Rdv1* QTL were subjected to GBS to locate the recombination points between

the two haplotypes of 'Börner'. V3125 was included as a control for comparative read

mapping. The SE reads were quality trimmed using Trimmomatic-0.39[46] with the fol-

lowing parameters: ILLUMINACLIP: 4:30:15 LEADING:30 TRAILING:30 SLID-

INGWINDOW:4:15 MINLEN:60. Trimmed reads of each of the six studied geno-

types as well as reads from 'Börner' [10] were separately mapped to BoeCin and

BoeRip using the HISAT2 version 2.2.0 [59] with the following parameters: --no-

softclip –no-spliced-alignment. The obtained SAM files were converted to sorted

BAM files as well as indexed using SAMtools. Variant calling against BoeRC and

estimation of genomic recombination sites was supported by QIAGEN CLC Genom-

ics Workbench 22.0 (QIAGEN, Aarhus, Denmark). SNPs detected in the *Rdv1* region

on chr13 for each genotype were filtered with respect to read coverage and frequency.

Homozygous variants were counted if > 90 % of the reads support the variation, for

heterozygous variants a frequency between 25 % and 75 % was required. An example

case (GBS-04) for the assessment of GBS markers is shown in Supplementary File 1

Fig. S15. Recombination sites were determined by monitoring the switch from either

the BoeCin haplotype to the BoeRip haplotype or *vice versa* between a set of at least

two variant positions.

Allele sequence comparison and TE detection

The genes (alleles, orthologs) of the *Rdv1* region of BoeRip, BoeCin, PN40024 and

VitRGM between the markers GF13-03 and GF13-07 were placed into groups with

OrthoFinder v2.3.11 [87] using the longest protein sequence annotated for the splicing

variants of each gene. Missing alleles (or, alternatively phrased, missing predictions

of orthologous genes at syntenic positions) were investigated and re-annotated by

aligning the protein sequences of the other cultivars against the genome sequences

with exonerate v2.4.0 [88]. Only alignments with at least 70 % coverage of the protein

sequence were considered. To obtain gene annotations for remaining poorly annotated

sequences despite existing RNA-Seq support in the genomic region studied,

BRAKER was run with strictly filtered RNA-Seq mappings (only uniquely mapped
PE reads on both 'Börner' haplotypes offered as combined target, both reads of a pair
must map); adding untranslated regions (UTRs) and prediction of alternative tran-
scripts from RNA-Seq data was enabled. Since the focus here is on phylloxera resis-
tance at the root, an additional AUGUSTUS species model was trained on BoeRC
with the uniquely mapped root RNA-Seq reads, the above described *Vitis* protein and
plant protein sequences and eudicot protein sequences from OrthoDB v10.1 [89]. The
results were manually evaluated and restored gene models of BoeRip and BoeCin
were refined two times with PASA. Through investigation of the RNA-Seq mappings
from all available tissues to BoeCin, the gene predictions in the *Rdv1* region of all
four sequence regions studied were manually screened for low supported gene mod-
els. Transcripts encoding for protein sequences with less than 50 amino acids, tagged
as 'protein of unknown function' (PUF) and with no ortholog were removed. RGA
classes were determined for the manually curated genes/alleles with RGAugury as
described above. TEs were identified with the Extensive *de novo* TE Annotator
(EDTA) [90]. Finally, the results were manually collected in allelic representations of
the genes of the *Rdv1* region of BoeRip, BoeCin, PN40024 and VitRGM. An ex-
tended version of the gene overview at the *Rdv1* locus that includes TEs has also been
prepared (Supplementary File 1 Fig. S11).
960

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

Sampling and phenotyping were done at the JKI (Julius Kuehn Institute, Institute for Grapevine Breeding Geilweilerhof, Siebeldingen, Germany). Sequencing and data analysis were performed at Bielefeld University, Faculty of Biology & Center for Biotechnology (CeBiTec) and MPIPZ (Cologne). The design of the experiments was set up by LH, DH and BF. Genotype selection, sampling and DNA isolations were carried out by LH and DH. PV, BH and RR accomplished library preparation and sequencing. LH performed the SSR analysis. BW supervised the work at Bielefeld University. RT supervised the work at JKI. RT and BW acquired project funding and wrote the project proposal. All bioinformatic data analyses, creation of figures, tables and writing of the initial draft of the manuscript were performed by BF and DH. BF and BW finalised the manuscript. All authors have read and approved the final manuscript.

Acknowledgements

The authors like to thank all members of the Genetics and Genomics of Plants team at Bielefeld University as well as the members of the Julius Kuehn Institute for Grapevine Breeding at Geilweilerhof.

The project was supported by funds from the Federal Ministry of Food and Agriculture (BMEL), based on a decision by the Parliament of the Federal Republic of Germany via the Federal Office for Agriculture and Food (BLE) under the innovation support program (project acronym MureViU, 28-1-82.066-15), as well as by the EU COST Action INTEGRAPE (CA 17111). We acknowledge support for the article processing charge by the Deutsche Forschungsgemeinschaft and the Open Access Publication Fund of Bielefeld University.

This work was supported by the BMBF-funded de.NBI Cloud within the German Network for Bioinformatics Infrastructure (de.NBI).

Data availability

The SMRT sequencing reads of 'Börner' (ERR6182799), the WGS reads of 'Börners' parents *V. riparia* GM18 (ERR6182967, ERR6183009) and *V. cinerea* Arnold (ERR6183041) as well as the 'Börner' haplotype assemblies (ERS6637871) were deposited in ENA/GenBank/DDJ under project no. PRJEB45595. The haplotype assemblies BoeRip and BoeCin at chromosome level with structural and functional annotation are available at (https://doi.org/10.4119/unibi/2962793). The RNA-Seq data of 'Börner' leaf (ERR6182801) and tendril (ERR6182801) were deposited in ENA/GenBank/DDJ under project no. PRJEB46079, the raw sequencing reads of 'Börner' BACs (ERR6183016-ERR6183021, ERR6183023-ERR6183025, ERR6183030-ERR6183032) under project no. PRJEB46081. The assembled and phase-separated BAC contigs are available at (https://doi.org/10.4119/unibi/2962639). DNA sequence read data from the F1 genotypes of the mapping population and the maternal parent V3125 generated for GBS were deposited in ENA/GenBank/DDJ under project no. PRJEB53997.

Supplementary information

Supplementary File 1 Fig. S1: Cumulative length of both 'Börner' haplotype assemblies

Supplementary File 1 Fig. S2: Sequence region with BAC sequences of both haplotypes

Supplementary File 1 Fig. S3: Spectra-copy number and assembly spectrum plot of the 'Börner' genome sequence assembly

Supplementary File 1 Fig. S4: Analysis of phasing with Mercury

Supplementary File 1 Fig. S5: All-versus-all dot plot between pseudochromosomes of the *V. riparia* haplotype of 'Börner' and the *V. vinifera* reference sequence.

Supplementary File 1 Fig. S6: All-versus-all dot plot between pseudochromosomes of the *V. cinerea* haplotype of 'Börner' and the *V. vinifera* reference sequence

Supplementary File 1 Fig. S7: All-versus-all dot plot between pseudochromosomes of the *V. riparia* haplotype of 'Börner' and those of *V. riparia* Gloire de Montpellier

Supplementary File 1 Fig. S8: All-versus-all dot plot between pseudochromosomes of the *V. cinerea* haplotype of 'Börner' and those of *V. riparia* Gloire de Montpellier

Supplementary File 1 Fig. S9: Dot plot of the extended *Rdv1* sequence region on pseudochromosome 13 between BoeRip and PN40024

Supplementary File 1 Fig. S10: Dot plot of *Rdv1* on pseudochromosome 13 between BoeCin and PN40024

Supplementary File 1 Fig. S11: Extended version of Fig. 6, "Genes of BoeCin,
BoeRip, PN40024 and VitRGM at and surrounding the *Rdv1* locus", transposable
elements/TEs included
Supplementary File 1 Fig. S12: Length distribution calculated over all subreads
Supplementary File 1 Fig. S13: Coverage plot of the 'Börner' haplotype assemblies
Supplementary File 1 Fig. S14: Display of initial assembly analysis of chr02 of
BoeRip
Supplementary File 1 Fig. S15: Example for determination of allelic constitution in
selected F1 genotypes
1033
Supplementary File 2 Table S1: Pseudochromosome sizes of the haplotype assemblies
BoeRip and BoeCin
Supplementary File 2 Table S2: Quality statistics of *Vitis* genome assemblies
Supplementary File 2 Table S3: Assembly statistics of the BAC contigs
Supplementary File 2 Table S4: Mapping statistics of the BAC contigs on the BoeRip
and BoeCin haplotype assembly
Supplementary File 2 Table S5: Statistics of the trimmed BAC sequences
Supplementary File 2 Table S6: Haplotype blocks of BoeRip and BoeCin
Supplementary File 2 Table S7: Repeat content of the haplotypes
Supplementary File 2 Table S8: GBS marker sites for the parents 'Börner' and V3125
and selected F1 offspring
Supplementary File 2 Table S9: BoeCin, BoeRip, PN40024 and VitRGM genes in
Rdv1 region
Supplementary File 2 Table S10: Amino acid sequence similarity matrix BoeCin to
BoeCin deduced from genes in *Rdv1* region
Supplementary File 2 Table S11: Amino acid sequence similarity matrix BoeCin to
BoeRip deduced from genes in *Rdv1* region
Supplementary File 2 Table S12: Amino acid sequence similarity matrix BoeCin to
PN40024 deduced from genes in *Rdv1* region
Supplementary File 2 Table S13: Amino acid sequence similarity matrix BoeCin to
VitRGM deduced from genes in *Rdv1* region
Supplementary File 2 Table S14: Primer sequences for SSR marker
Supplementary File 2 Table S15: RNA-Seq read trimming statistics
Supplementary File 2 Table S16: Trimming statistics of *V. riparia* and *V. cinerea* ge-
nomic reads
Supplementary File 2 Table S17: Description of samples applied to GBS
Supplementary File 2 Table S18: Mapping positions of the genetic SSR marker on the
PN40024 genome assembly and on the 'Börner' haplotype assemblies BoeRip and
BoeCin
Supplementary File 2 Table S19: Distribution of the 344 genetic marker used for
pseudochromosome validation over BoeRip and BoeCin.
1065
1066   Supplementary File 3 Data S1: Parameter file for PASA gene model refinement

1067

References

1. Ramos-Madrigal J*, et al.* Palaeogenomic insights into the origins of French grapevine diversity. *Nature Plants* **5**, 595-603 (2019).

2. Vezzulli S*, et al.* Genomic Designing for Biotic Stress Resistant Grapevine. In: *Genomic Designing for Biotic Stress Resistant Fruit Crops* (ed Kole C). Springer Nature Switzerland AG (2022).

3. Granett J, Walker MA, Kocsis L, Omer AD. Biology and management of grape phylloxera. *Annual Review of Entomology* **46**, 387-412 (2001).

4. Tello J, Mammerler R, Cajic M, Forneck A. Major Outbreaks in the Nineteenth Century Shaped Grape Phylloxera Contemporary Genetic Structure in Europe. *Scientific Reports* **9**, 17540 (2019).

5. Forneck A, Kleinmann S, Blaich R, Anvari SF. Histochemistry and anatomy of phylloxera (Daktulosphaira vitifoliae) nodosities on young roots of grapevine (Vitis spp). *Vitis* **41**, 93-97 (2002).

6. Zhang J, Hausmann L, Eibach R, Welter LJ, Töpfer R, Zyprian EM. A framework map from grapevine V3125 (Vitis vinifera 'Schiava grossa' x 'Riesling') x rootstock cultivar 'Börner' (Vitis riparia x Vitis cinerea) to localize genetic determinants of phylloxera root resistance. *Theoretical and Applied Genetics* **119**, 1039-1051 (2009).

7. Töpfer R, Hausmann L, Harst M, Maul E, Zyprian E, Eibach R. New horizons for grapevine breeding. In: *Fruit, vegetable and cereal science and biotechnology,*). Global Science Books (2011).

8. Rex F, Fechter I, Hausmann L, Töpfer R. QTL mapping of black rot (Guignardia bidwellii) resistance in the grapevine rootstock 'Börner' (V. riparia Gm183 × V. cinerea Arnold). *Theoretical and Applied Genetics* **127**, 1667-1677 (2014).

9. Ochssner I, Hausmann L, Töpfer R. Rpv14, a new genetic source for Plasmopara viticola resistance conferred by Vitis cinerea. *Vitis* **55**, 79-81 (2016).

10. Holtgräwe D*, et al.* A Partially Phase-Separated Genome Sequence Assembly of the Vitis Rootstock 'Borner' (Vitis riparia x Vitis cinerea) and Its Exploitation for Marker Development and Targeted Mapping. *Frontiers in Plant Science* **11**, 156 (2020).

11. Han GZ. Origin and evolution of the plant immune system. *New Phytologist* **222**, 70-83 (2019).

12. Sekhwal MK, Li P, Lam I, Wang X, Cloutier S, You FM. Disease Resistance Gene Analogs (RGAs) in Plants. *International Journal of Molecular Sciences* **16**, 19248-19290 (2015).

13. Lodhi MA, Reisch BI. Nuclear DNA content of Vitis species, cultivars, and other genera of the Vitaceae. *Theoretical and Applied Genetics* **90**, 11-16 (1995).

14. Zhou Y*, et al.* The population genetics of structural variants in grapevine domestication. *Nature Plants* **5**, 965-979 (2019).

15. Adam-Blondon A-F, Jaillon O, Vezzulli S, Zharkikh A, Troggio M, Velasco R. Genome Sequence Initiatives. In: *Genetics, Genomics and Breeding of Grapes* (eds Adam-Blondon A-F, Martinez-Zapater JM, Kole C) (2011).

16. Jaillon O, *et al.* The grapevine genome sequence suggests ancestral hexaploidization in major angiosperm phyla. *Nature* **449**, 463-467 (2007).

17. Vitulo N, *et al.* A deep survey of alternative splicing in grape reveals changes in the splicing machinery related to tissue, stress condition and genotype. *BMC Plant Biology* **14**, 99 (2014).

18. Riaz S, *et al.* Genetic diversity analysis of cultivated and wild grapevine (Vitis vinifera L.) accessions around the Mediterranean basin and Central Asia. *BMC Plant Biology* **18**, 137 (2018).

19. Chin CS, *et al.* Phased diploid genome assembly with single-molecule real-time sequencing. *Nature Methods* **13**, 1050-1054 (2016).

20. Massonnet M, *et al.* The genetic basis of sex determination in grapes. *Nature Communications* **11**, 2902 (2020).

21. Minio A, Massonnet M, Figueroa-Balderas R, Castro A, Cantu D. Diploid Genome Assembly of the Wine Grape Carmenere. *G3 Genes Genomes Genetics* **9**, 1331-1337 (2019).

22. Roach MJ, *et al.* Population sequencing reveals clonal diversity and ancestral inbreeding in the grapevine cultivar Chardonnay. *PLoS Genetics* **14**, e1007807 (2018).

23. Cochetel N, Minio A, Massonnet M, Vondras AM, Figueroa-Balderas R, Cantu D. Diploid chromosome-scale assembly of the Muscadinia rotundifolia genome supports chromosome fusion and disease resistance gene expansion during Vitis and Muscadinia divergence. *G3 Genes Genomes Genetics* **11**, jkab033 (2021).

24. Girollet N, Rubio B, Lopez-Roques C, Valiere S, Ollat N, Bert PF. De novo phased assembly of the Vitis riparia grape genome. *Scientific Data* **6**, 127 (2019).

25. Ollat N, *et al.* Rootstocks as a component of adaptation to environment. In: *Grapevine in a changing environment: a molecular and ecophysiological perspective* (eds Geros H, Chaves MM, Gil HM, Delrot S). John Wiley and Sons (2016).

26. Eid J, *et al.* Real-time DNA sequencing from single polymerase molecules. *Science* **323**, 133-138 (2009).

27. Michael TP, VanBuren R. Building near-complete plant genomes. *Current Opinion in Plant Biology* **54**, 26-33 (2020).

28. Koren S, *et al.* De novo assembly of haplotype-resolved genomes with trio binning. *Nature Biotechnology* **36**, 1174-1182 (2018).

29. Li P, Quan X, Jia G, Xiao J, Cloutier S, You FM. RGAugury: a pipeline for genome-wide prediction of resistance gene analogs (RGAs) in plants. *BMC Genomics* **17**, 852 (2016).

30.   Koren S, Walenz BP, Berlin K, Miller JR, Bergman NH, Phillippy AM. Canu: scalable and accurate long-read assembly via adaptive k-mer weighting and repeat separation. *Genome Research* **27**, 722-736 (2017).

31.   Xu Z, Dixon JR. Genome reconstruction and haplotype phasing using chromosome conformation capture methodologies. *Briefings in Functional Genomics* **19**, 139-150 (2020).

32.   Sweetman C, Wong DC, Ford CM, Drew DP. Transcriptome analysis at four developmental stages of grape berry (Vitis vinifera cv. Shiraz) provides insights into regulated and coordinated gene expression. *BMC Genomics* **13**, 691 (2012).

33.   Lijavetzky D, Cabezas JA, Ibanez A, Rodriguez V, Martinez-Zapater JM. High throughput SNP discovery and genotyping in grapevine (Vitis vinifera L.) by combining a re-sequencing approach and SNPlex technology. *BMC Genomics* **8**, 424 (2007).

34.   Salmaso M*, et al.* Genome diversity and gene haplotypes in the grapevine (Vitis vinifera L.), as revealed by single nucleotide polymorphisms. *Molecular Breeding* **14**, 385-395 (2004).

35.   Hulbert SH, Webb CA, Smith SM, Sun Q. Resistance gene complexes: evolution and utilization. *Annual Review of Phytopathology* **39**, 285-312 (2001).

36.   Dupeyron M, Singh KS, Bass C, Hayward A. Evolution of Mutator transposable elements across eukaryotic diversity. *Mobile DNA* **10**, 12 (2019).

37.   Kapitonov VV, Jurka J. A universal classification of eukaryotic transposable elements implemented in Repbase. *Nature Reviews Genetics* **9**, 411-414 (2008).

38.   Foria S*, et al.* Gene duplication and transposition of mobile elements drive evolution of the Rpv3 resistance locus in grapevine. *The Plant Journal* **101**, 529-542 (2020).

39.   Bianchet C*, et al.* An Arabidopsis thaliana leucine-rich repeat protein harbors an adenylyl cyclase catalytic center and affects responses to pathogens. *Journal of Plant Physiology* **232**, 12-22 (2019).

40.   Bittner-Eddy PD, Beynon JL. The Arabidopsis downy mildew resistance gene, RPP13-Nd, functions independently of NDR1 and EDS1 and does not require the accumulation of salicylic acid. *Molecular Plant Microbe Interactions* **14**, 416-421 (2001).

41.   Allen RL*, et al.* Host-parasite coevolutionary conflict between Arabidopsis and downy mildew. *Science* **306**, 1957-1960 (2004).

42.   Canaguier A*, et al.* A new version of the grapevine reference genome assembly (12X.v2) and of its annotation (VCost.v3). *Genomics Data* **14**, 56-62 (2017).

43.   Fechter I*, et al.* QTL analysis of flowering time and ripening traits suggests an impact of a genomic region on linkage group 1 in Vitis. *Theoretical and Applied Genetics* **127**, 1857-1872 (2014).

44.   Rosso MG, Li Y, Strizhov N, Reiss B, Dekker K, Weisshaar B. An *Arabidopsis thaliana* T-DNA mutagenised population (GABI-Kat) for flanking sequence tag based reverse genetics. *Plant Molecular Biology* **53**, 247-259 (2003).

45.   Frommer B, *et al.* Genome Sequences of Both Organelles of the Grapevine Rootstock Cultivar 'Börner'. *Microbiology Resource Announcement* **9**, e01471-01419 (2020).

46.   Bolger AM, Lohse M, Usadel B. Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics* **30**, 2114-2120 (2014).

47.   Forneck A, Walker MA, Blaich R, Yvon M, Leclant F. Interaction of phylloxera (Daktulosphaira vitifoliae Fitch) with grape (Vitis spp.) in simple isolation chambers. *American Journal of Enology and Viticulture* **52**, 28-34 (2001).

48.   Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. Basic local alignment search tool. *Journal of Molecular Biology* **215**, 403-410 (1990).

49.   Camacho C, *et al.* BLAST+: architecture and applications. *BMC Bioinformatics* **10**, 421 (2009).

50.   Jansen RK, *et al.* Phylogenetic analyses of Vitis (Vitaceae) based on complete chloroplast genome sequences: effects of taxon sampling and phylogenetic methods on resolving relationships among rosids. *BMC Evolutionary Biology* **6**, 32 (2006).

51.   Goremykin VV, Salamini F, Velasco R, Viola R. Mitochondrial DNA of Vitis vinifera and the issue of rampant horizontal gene transfer. *Molecular Biology and Evolution* **26**, 99-110 (2009).

52.   Mikheenko A, Valin G, Prjibelski A, Saveliev V, Gurevich A. Icarus: visualizer for de novo assembly evaluation. *Bioinformatics* **32**, 3321-3323 (2016).

53.   Kim D, Langmead B, Salzberg SL. HISAT: a fast spliced aligner with low memory requirements. *Nature Methods* **12**, 357-360 (2015).

54.   Li H, *et al.* The Sequence Alignment/Map format and SAMtools. *Bioinformatics* **25**, 2078-2079 (2009).

55.   Boetzer M, Henkel CV, Jansen HJ, Butler D, Pirovano W. Scaffolding pre-assembled contigs using SSPACE. *Bioinformatics* **27**, 578-579 (2011).

56.   Li H, Durbin R. Fast and accurate short read alignment with Burrows-Wheeler transform. *Bioinformatics* **25**, 1754-1760 (2009).

57.   Stanke M, Waack S. Gene prediction with a hidden Markov model and a new intron submodel. *Bioinformatics* **19**, ii215–ii225 (2003).

58.   Grabherr MG, *et al.* Full-length transcriptome assembly from RNA-Seq data without a reference genome. *Nature Biotechnology* **29**, 644-652 (2011).

59.   Kim D, Paggi JM, Park C, Bennett C, Salzberg SL. Graph-based genome alignment and genotyping with HISAT2 and HISAT-genotype. *Nature Biotechnology* **37**, 907-915 (2019).

60.   Kent WJ. BLAT--the BLAST-like alignment tool. *Genome Research* **12**, 656-664 (2002).

61. Marcais G, Delcher AL, Phillippy AM, Coston R, Salzberg SL, Zimin A. MUMmer4: A fast and versatile genome alignment system. *PLoS Computational Biology* **14**, e1005944 (2018).

62. Fechter I, *et al.* Candidate genes within a 143 kb region of the flower sex locus in Vitis. *Molecular Genetics and Genomics* **287**, 247-259 (2012).

63. Rice P, Longden I, Bleasby A. EMBOSS: The European Molecular Biology Open Software Suite. *Trends in Genetics* **16**, 276-277 (2000).

64. Cabanettes F, Klopp C. D-GENIES: dot plot large genomes in an interactive, efficient and simple way. *PeerJ* **6**, e4958 (2018).

65. Li H. Minimap2: pairwise alignment for nucleotide sequences. *Bioinformatics* **34**, 3094-3100 (2018).

66. Simão FA, Waterhouse RM, Ioannidis P, Kriventseva EV, Zdobnov EM. BUSCO: assessing genome assembly and annotation completeness with single-copy orthologs. *Bioinformatics* **31**, 3210-3212 (2015).

67. Waterhouse RM, *et al.* BUSCO Applications from Quality Assessments to Gene Prediction and Phylogenomics. *Molecular Biology and Evolution* **35**, 543-548 (2018).

68. Cannon EK, Cannon SB. Chromosome visualization tool: a whole genome viewer. *International Journal of Plant Genomics* **2011**, 373875 (2011).

69. Robinson JT, *et al.* Integrative genomics viewer. *Nature Biotechnology* **29**, 24-26 (2011).

70. Thorvaldsdottir H, Robinson JT, Mesirov JP. Integrative Genomics Viewer (IGV): high-performance genomics data visualization and exploration. *Briefings in Bioinformatics* **14**, 178-192 (2013).

71. Rhie A, Walenz BP, Koren S, Phillippy AM. Merqury: reference-free quality, completeness, and phasing assessment for genome assemblies. *Genome Biology* **21**, 245 (2020).

72. Holt C, Yandell M. MAKER2: an annotation pipeline and genome-database management tool for second-generation genome projects. *BMC Bioinformatics* **12**, (2011).

73. Campbell MS, *et al.* MAKER-P: a tool kit for the rapid creation, management, and quality control of plant genome annotations. *Plant Physiology* **164**, 513-524 (2014).

74. Bao W, Kojima KK, Kohany O. Repbase Update, a database of repetitive elements in eukaryotic genomes. *Mobile DNA* **6**, 11 (2015).

75. Korf I. Gene finding in novel genomes. *BMC Bioinformatics* **5**, 59 (2004).

76. Lomsadze A, Burns PD, Borodovsky M. Integration of mapped RNA-Seq reads into automatic training of eukaryotic gene finding algorithm. *Nucleic Acids Research* **2014**, e119 (2014).

77.   Haas BJ, *et al.* Automated eukaryotic gene structure annotation using EVidenceModeler and the Program to Assemble Spliced Alignments. *Genome Biology* **9**, R7 (2008).

78.   Haas BJ*, et al.* Improving the Arabidopsis genome annotation using maximal transcript alignment assemblies. *Nucleic Acids Research* **31**, 5654-5666 (2003).

79.   Jones P*, et al.* InterProScan 5: genome-scale protein function classification. *Bioinformatics* **30**, 1236-1240 (2014).

80.   Lupas A, Van Dyke M, Stock J. Predicting coiled coils from protein sequences. *Science* **252**, 1162-1164 (1991).

81.   Mistry J, Bateman A, Finn RD. Predicting active site residue annotations in the Pfam database. *BMC Bioinformatics* **8**, 298 (2007).

82.   Kall L, Krogh A, Sonnhammer EL. A combined transmembrane topology and signal peptide prediction method. *Journal of Molecular Biology* **338**, 1027-1036 (2004).

83.   Sillitoe I*, et al.* CATH: expanding the horizons of structure-based functional annotations for genome sequences. *Nucleic Acids Research* **47**, D280-D284 (2019).

84.   Mistry J*, et al.* Pfam: The protein families database in 2021. *Nucleic Acids Research* **49**, D412-D419 (2021).

85.   Letunic I, Bork P. 20 years of the SMART protein domain annotation resource. *Nucleic Acids Research* **46**, D493-D496 (2018).

86.   Gough J, Karplus K, Hughey R, Chothia C. Assignment of homology to genome sequences using a library of hidden Markov models that represent all proteins of known structure. *Journal of Molecular Biology* **313**, 903-919 (2001).

87.   Emms DM, Kelly S. OrthoFinder: phylogenetic orthology inference for comparative genomics. *Genome Biology* **20**, 238 (2019).

88.   Slater GS, Birney E. Automated generation of heuristics for biological sequence comparison. *BMC Bioinformatics* **6**, 31 (2005).

89.   Kriventseva EV*, et al.* OrthoDB v10: sampling the diversity of animal, plant, fungal, protist, bacterial and viral genomes for evolutionary and functional annotations of orthologs. *Nucleic Acids Research* **47**, D807-D811 (2019).

90.   Ou S*, et al.* Benchmarking transposable element annotation methods for creation of a streamlined, comprehensive pipeline. *Genome Biology* **20**, 275 (2019).