1

Cortico-cerebellar coordination facilitates neuroprosthetic learning

Aamir Abbasi[1], Andrew W. Fealy[1], Nathan P. Danielsen[1], Tanuj Gulati[1,2, #]

[1] Center for Neural Science and Medicine, Departments of Biomedical Sciences and Neurology, Cedars-Sinai Medical Center, Los Angeles, CA.

[2] Department of Medicine, David Geffen School of Medicine, and Department of Bioengineering, Henry Samueli School of Engineering, University of California-Los Angeles, Los Angeles, CA.

[#] Correspondence: tanuj.gulati@csmc.edu

Abstract

Temporal coordination among neurons and development of functional neuronal assemblies is central to nervous system function and purposeful behavior. Still, there is a paucity of evidence about how functional coordination emerges in neuronal assemblies in cortical and subcortical regions that are directly related to the control of functional output. We investigated emergent neural dynamics between primary motor cortex (M1) and the contralateral cerebellar cortex as rats learned a neuroprosthetic/ brain-machine interface (BMI) task. BMIs offer a powerful tool to causally test how distributed neural networks achieve specific neural patterns. During neuroprosthetic learning, actuator movements are causally linked to primary motor cortex (M1) neurons, i.e., "direct" neurons, that project to the decoder and whose firing is required to successfully perform the task. However, it is unknown how such direct M1 activity interacts with cerebellar activity. We observed a striking 3-6 Hz coherence that emerged between these regions' local-field potentials (LFPs) with neuroprosthetic learning which also modulated task-related spiking. We found a robust task-related indirect modulation in the cerebellum, and we found that it became synchronous with M1-direct activity with learning. We also performed optogenetic inhibition of cerebellar activity and found that this led to performance impairments in neuroprosthetic control. Together, these results demonstrate that coordinated neural dynamics emerge in cortico-cerebellar regions during neuroprosthetic learning which supports task-relevant activity in M1 direct neuronal populations, and further, that cerebellar influence is necessary for M1-driven rapid direct neural control.

Keywords: Cerebellum, motor cortex, brain-machine interface

Introduction

To accomplish even the simplest of tasks, the nervous system coordinates activity across distant brain regions. For example, holding a bottle full of water requires several neurons to produce well-calibrated muscle forces for grasping and monitoring sensory feedback. Parallel processes in several sensorimotor regions underlie even the most trivial tasks such as above. Dense reciprocal connectivity between these regions supports this processing which likely need to be configured rapidly and flexibly to support a repertoire of behavior and afford learning of new skills. Two motor regions, the primary motor cortex (M1) and the cerebellum, have dense reciprocal connections, and are known to be involved in motor learning[1,2]. Studies have shown how learning alters local activity in the cerebellum or M1[3–6], yet learning also alters task-related cross area coordination[2,7]. Both M1 and the cerebellum have dense connections with other cortical and subcortical regions, and hence it is difficult to ascertain if changes in interaction are because of direct interactions between M1 and cerebellum[8–10], or because of their roles coordinating the upper limb.

To isolate the effects of M1 and cerebellum to movement control versus the direct task-related coordination, we have used a brain-machine interface task, where select M1 neurons (*'direct'* neurons) modulate their activity to control an external disembodied actuator. This offers experimenters a powerful paradigm where they can dictate/set the neuron-behavior relationship. During "brain control", direct M1 neurons change their firing properties during neuroprosthetic learning[11–18]. In addition, other neurons in the local M1 network also become task-coupled (i.e., *'indirect'* neurons)[12,13,15,19–23]. It remains

unknown what activity emerges in the cerebellum with M1-driven neuroprosthetic learning

and what role it plays. Here, we hypothesized that cerebellum neurons will demonstrate

task-related firing during M1-driven neuroprosthetic learning. We also predicted that

modulating the cerebellum optogenetically will impact M1-driven neuroprosthetic

learning.

Another focus of our investigations was on co-emergent synchronous activity across M1

and cerebellum with neuroprosthetic learning. Recent theories have proposed that

alterations in the pattern of synchronous activity across regions can serve to coordinate

network activity for natural and neuroprosthetic behaviors[20,24]. Such transient local-field

potential (LFP) activity can modulate excitability of cell groups across varying

spatiotemporal scales[25,26]. This helps in achieving precise temporal control in neural

networks that can enhance information transfer in specific cell populations[27] and can

influence spike-timing-dependent plasticity[28]. Such temporally coordinated activity among

ensembles underlies diverse neural processes ranging from perception, decision making,

action, memory and attention[6,29–33]. Importantly, spiking in one region becomes

coordinated with LFP in another region, indicative of synchrony[30,34]. Despite the evidence

that synchronous LFP is related to learning[6,35], there is a paucity of evidence that this

selectively modulates task-relevant activity of neurons across brain areas. Using the

neuroprosthetic task paradigm, where we can control the neurons that are linked to

behavioral output, we aimed to disentangle the synchronous activity of "direct" neurons

within M1 locally and with task-relevant indirect activity in the cerebellum to understand

how these diverse classes of cells were modulated by synchronous LFP.

We have used a BMI paradigm in which a small set of M1-direct neurons controlled a

simple 1-D actuator[12,18,21]. We recorded additional neural activity from neighboring

indirect neurons in M1, as well as distant cerebellar cortex. First, we found that cerebellar

neurons developed strong 'task-related' indirect modulation, and M1 and cerebellum

developed a 3-6Hz LFP coherence with neuroprosthetic learning. We also found that

direct M1 neurons and task-related indirect neurons in both M1 and cerebellum enhanced

their phase-locking to this 3-6Hz oscillation in the LFPs. Next, we also found that

synchronous spiking between M1 direct neurons and cerebellar indirect neurons

increased with neuroprosthetic learning, and fine timescale coordination between these

regions also increased as evaluated through canonical correlation analyses. In our last

set of experiments, we optogenetically inhibited the cerebellum and found that this led to

performance impairments in the neuroprosthetic task and weakening of M1-direct activity.

Together, our results support a necessary role of cross-area interactions between M1 and

cerebellum that underlie neuroprosthetic skill learning.

Results

We implanted microwire electrode arrays in the primary motor cortex (M1) and silicon probes in a tetrode/ polytrode configuration in the cerebellar posterior lobes (see Methods for details). After neural implant surgery, we trained 8 animals to exert direct neural control on the angular velocity of a mechanical actuator that can also deliver water. A linear decoder with randomized weights typically converted the firing rates of two groups of units in M1 (randomly selected and assigned positive or negative weights; hereafter referred to as *'direct'* units; M1 TR_d's) into the angular velocity of the actuator. We also recorded multiple other units in both M1 and cerebellum that were not causally linked to actuator movements but showed task-related modulation (hereafter referred to as *'indirect'* units). The decoder weights were held constant during the session to exclusively rely on neural learning mechanisms. Each trial started with the simultaneous delivery of an auditory tone and opening of a door to allow access to the tube (**Fig. 1 a,b**). At the start of each trial, the angular position of the tube was set to resting position, P_1. If the angular position of the tube was held for >400 ms at target position P_2, a defined amount of water was delivered (that is, successful trial). A trial was stopped if this was not achieved within 15s (that is, unsuccessful trial). At the end of a trial, the actuator was returned to position P_1 and the door was closed.

Direct control of BMI by M1 units

We observed that over the course of a typical 1-2h practice session, animals showed improvements in task performance with a significant reduction in the time to successful

trial completion and decrease in the proportion of unsuccessful trials. We report results

here from 23 sessions, where we saw significant reductions in both metrics.

Overall we observed that rats showed improvement in task-performance with a significant

reduction in the time to a successful trial completion (**Fig. 1c,d**; 7.69 ± 0.42 sec to

4.09 ± 0.3 sec, mixed-effects model: $t(44) = -7.13$, $P = 7.3 \times 10^{-9}$) and a decrease in the

percentage of unsuccessful trials (**Fig. 1d**; 28.12 ± 3.65 % to 2.71 ± 0.97 %, mixed-effects

model: $t(44) = -7.01$, $P = 1.0 \times 10^{-8}$).

In a subset of sessions, we performed video recording of rats during the BMI training and

quantified the proficiency of brain control by ruling out movement of the implanted limb.

Consistent with previous reports of lack of muscle contractions during neuroprosthetic

control, we did not observe limb movements to systematically predict feeding tube

movements[15]. Specifically, we analyzed whether limb movements measured using the

video recording correlated with movements of the feeding tube. Across multiple sessions

we observed significant reduction of this correlation (**Supplementary Fig 3a,b**;

0.014 ± 0.002 to 0.006 ± 0.001, mixed-effects model: $t(26) = -3.52$, $P = 1 \times 10^{-3}$).

We also analyzed the proportion of TR_d's that developed robust task-related modulation

(see Methods). **Figure 1e**, depicts the increase in the activity of M1 direct units from early

to late trials. Since the direct units were causally linked to the movement of the actuator,

we found that a high proportion of these units developed task-related modulation (96% of

50 TR_d's), as has been reported in the past[11].

Indirect modulation of neurons in M1 and cerebellum

We also analyzed indirect units in the M1 and cerebellum and checked for their task-

related modulation. We further sub-classified indirect units as either task-related (TR_i's)

or task-unrelated (*TU's*) based on changes in modulation depth with learning. Consistent

with previous reports of indirect units in the M1 likely contributing to neuroprosthetic

control[11,13,15,19,21], we found strong indirect modulation in M1 (**Fig. 1f**). Additionally, we

found robust indirect modulation of units in the cerebellum (**Fig. 1h**). Majority of units in

M1 and cerebellum developed strong indirect task-related modulation with learning (**Fig.**

1j). We also looked at the change in modulation depth (MD_Δ) and found that M1 TR_d

undergoes greater MD_Δ from early to late trials as compared to M1 and Cb TR_i units (M1

TR_d: 130.46 ± 47.8%; M1 TR_i: 51.075 ± 15.5%; cerebellum TR_i: 25.88 ± 9.5 %, Kruskal-

Wallis H-test, $F_{2,713}$ = 195.23, P = 4.0 x 10^{-43}, *post hoc t*-test showed significant difference

among M1 TR_d, TR_i and cerebellum TR_i MD_Δ *'s; P* < 0.05). Our next analyses focused

on these task-related cells.

Figure 1. Direct and indirect modulation of M1 and cerebellar activity with neuroprosthetic learning. **a,** Direct neural control of a feeding tube (θ = angular position). Each trial started with the tube at P_1. **b,** Trial started with an audio tone cue and opening of the door. A successful trial required movement of the tube to P_2 within 15 seconds. **c,** Change in task completion time as a function of trial number. Line shows moving average of 5 trials. Dots show raw task completion times across all trials. **d,** Change in time to trial completion (*left*) and reduction in the percentage of unsuccessful trials (*right*) from *early* to *late* trials across all sessions. **e,** Peri-event histogram (PETH) from *early* and *late* trials from a single session are shown in *left* and *right* panels respectively, M1 TR_d task-related direct units. **f,** M1 TR_i units. **g,** Cerebellum TR_i units **h,** Distribution of units classified as TR_d, TR_i and TU in M1 (*top*) and cerebellum (*bottom*). ***p<0.001.

Emergence of coordinated task-related activity in M1 and cerebellar LFPs

Interestingly, as the rats became proficient in M1-driven neuroprosthetic control, we

observed that a coordinated low-frequency activity (approximately 3–6 Hz, **Fig. 2a**)

emerged with learning in both M1 and cerebellum. The task-related LFP power between

3–6 Hz increased from early to late trials in both M1 (**Fig. 2b,c**; 0.44 ± 0.07 to 0.80 ± 0.075,

mixed-effects model: $t(44) = 4.9$, $P = 1.0 \times 10^{-5}$) and the cerebellum (**Fig. 2d,e**; 0.18 ± 0.03

to 0.37 ± 0.04, mixed-effects model: $t(44) = 2.5$, $P = 1.0 \times 10^{-2}$). Task-related LFP

coherence also increased in the 3−6 Hz frequency range from early to late trials (**Fig. 2f,**

g; 0.16 ± 0.01 to 0.22 ± 0.013, mixed-effects model: $t(44) = 4.6$, $P = 2.5 \times 10^{-5}$). This

emergence of cross-region low-frequency activity is consistent with other observations of

emergence of low-frequency activity during motor skill learning across reciprocally

connected neural networks[7].

Figure 2. Co-ordinated task-related oscillations emerge in M1 and cerebellar LFPs *a,* Raw
and filtered LFP trace from a representative trials showing increase in 3–6 Hz oscillations after
task-start during *late* trials in both M1 and cerebellum. ***b,*** Spectrogram from a representative M1
channel showing increase in 3–6 Hz power during *late* trials. ***c,*** Increase in 3–6 Hz power in M1
LFP from *early* to *late* trials across sessions. ***d,*** Same as ***b*** but from a representative cerebellum
LFP channel. ***e,*** Same as ***c*** for 3–6 Hz cerebellum LFP power across sessions. ***f,*** Coherogram
from a representative pair of M1–cerebellum LFP channel showing increase in 3–6 Hz coherence
during *late* trials. ***g,*** Change in 3–6 Hz coherence from *early* to *late* trials across sessions. *p<0.05,
***p<0.001.

Increase in locking between spiking and LFP after neuroprosthetic learning

We next investigated the relationship between spiking activity and the low frequency LFP oscillations that we observed during robust task engagement. We performed spike-triggered averaging (STA) of the LFP in *early* and *late* learning time-locked to spikes occurring either in the same region (*i.e.,* M1 spikes to M1 LFP and cerebellum spikes to cerebellum LFP) or in the other region (*i.e.,* M1 spikes to cerebellum LFP and cerebellum spikes to M1 LFP). If spiking activity was independent of low-frequency LFP activity, then fluctuations would cancel and produce a flat average LFP, and hence, the STA can provide an intuitive estimate of how neural spiking is modulated by the low-frequency LFP oscillations. We observed a clear increase in the amplitude of mean LFP oscillations in both regions around action potentials of M1 TR_d, TR_i and cerebellum TR_i units (**Fig. 3a**). This increase was not present for these classes of cells when we did the STA analysis during task-unrelated inter-trial interval (**Fig. 3b**). M1 TR_d and TR_i units, and cerebellum TR_i units experienced 86.15 ± 11.50%, 90.97 ± 3.78% and 83.28 ± 5.95% increases in STA amplitude with M1 LFP during task periods, respectively. During inter-trial interval these units experience a decrease of −3.52 ± 2.68%, −0.25 ± 2.55% and −2.02 ± 2.39%, respectively (**Fig. 3c**; Kruskal-Wallis H-test, $F_{5,137}$ = 73.74, P = 3.0 x 10^{-20}, *post hoc t*-test showed significant difference for M1 TR_d, TR_i and cerebellum TR_i during task-relevant period and inter-trial period; P < 0.001). When STA was performed with cerebellum LFP, M1 TR_d and TR_i units, and cerebellum TR_i units experienced 80.87 ± 11.28%, 78.52 ± 6.83% and 113.75 ± 22.10% increases in STA amplitude (during the task period), respectively. Similar to M1 LFP, these units experienced significantly less change of 2.55 ± 1.73%, −0.81 ± 2.85% and 1.44 ± 3.52%, respectively, when STA was performed with

cerebellum LFP during inter-trial interval (**Fig. 3d**; Kruskal-Wallis H-test, $F_{5,137}$ = 23.63, P

= 5.6 x 10^{-21}, *post hoc t*-test showed significant difference for M1 TR_d, TR_i and cerebellum

TR_i during task-relevant period and inter-trial periods; P < 0.001).

Figure 3. M1 and cerebellum spike-LFP locking increases with learning. *a,* The mean M1
LFP (*top* row) or cerebellum LFP (*bottom* row) time locked to occurrences of spikes from M1 (left
column) or cerebellum (right column) during task-period from a representative session. ***b,*** Same
as ***a*** during inter-trial interval. ***c,*** Percentage change in STA amplitude for M1 LFP in each of the
categories of units (mean ± s.e.m.). ***d,*** Same as ***c*** for STA of cerebellum LFP. ***p<0.001.
234

Increase in neural synchrony between M1 and cerebellum

We subsequently assessed whether there are changes in the correlated spiking among

the recorded M1 and cerebellum neural ensembles with neuroprosthetic learning. To

analyze this, we looked at how the magnitude of cross-correlation between pairs of M1

TR_d's and cerebellum TR_i's changed. An example of an increase in cross-correlation is

shown in **Fig. 4a**. At the population level, we calculated the magnitude of the difference

between the peak average early trials and late trials correlogram from a shuffled

correlogram (*ΔCCH*, see "Methods"). We found a significant increase in correlated firing

(calculated as an increase in above-mentioned difference) during late trials (**Fig. 4b**;

4.52 ± 0.46 to 6.30 ± 0.60, mixed-effects model: $t(44) = 5.15$, $P = 5.8 \times 10^{-6}$). Thus, our

findings show that *'direct'* units in M1 (M1 TR_d's) and *'indirect'* cerebellar TR_i's fire more

coincidently with neuroprosthetic learning.

We also performed cross-correlation between M1 TR_d's and M1 TR_i's. Consistent with

previous reports[11,15], we found a significant increase in the magnitude of cross-correlation

(**Fig. 4c**; 2.86 ± 0.41 to 4.18 ± 0.46, mixed-effects model: $t(44) = 2.64$, $P = 0.01$). We also

performed cross-correlation between M1 TR_i's and cerebellum TR_i's (see **Fig. 4d**).

Surprisingly, we found no significant change in the magnitude of cross-correlation in this

pair with learning (**Fig. 4e**; 3.56 ± 0.69 to 3.11 ± 0.54, mixed-effects model: $t(44) = -1.92$,

$P = 0.06$).

Figure 4. M1 *TR_d* and cerebellar *TR_i* neural synchrony increases with learning. *a,* Representative cross-correlogram of a pair of M1 TR_d and cerebellum TR_i unit in early (*left*) and late (*right*) trials. ***b,*** Change in magnitude of cross-correlogram from early to late trials across all sessions for all M1 TR_d and cerebellum TR_i units (mean ± s.e.m.). ***c,*** Same as ***b*** for M1 TR_d and M1 TR_i units pairs. ***d,*** Representative cross-correlogram of a pair of M1 TR_i and cerebellum TR_i unit in early (*left*) and late (*right*) trials. ***e,*** Same as ***b*** for M1 TR_i and cerebellum TR_i unit pairs. *p<0.05 ***p<0.001

Fine timescale coordination of M1 and cerebellum activity with task learning

While our analyses so far show a coordinated activity in M1 and cerebellum with task learning; it doesn't necessarily indicate that the neural activity patterns in the two structures are coordinated across trials. Recent studies have explored fine-timescale coordination at the level of spiking[36,37]. Such methods use statistical methods to measure 'communication subspaces' based on ensemble patterns. Here, we used canonical correlation analysis (CCA) to assess fine-timescale coordination between M1 and cerebellum. CCA has been used in various neuroscience studies to extract correlated

population activity between two areas[36–40]. Specifically, CCA finds a linear combination

of units in M1 and cerebellum that project to M1 and cerebellum subspaces, respectively,

and where activity in these subspaces is maximally correlated (**Fig. 5a**). We used

concatenated single-trial spiking activity binned at 100ms as in two recent papers[37,40].

The top component produced by CCA is the axis of the M1 and cerebellar subspaces that

has the maximum correlation between the two areas. We found that this maximum

correlation increased with neuroprosthetic learning only for M1 TR_d and cerebellum TR_i

units (**Fig. 5b** shows CCA changes in an example session from a single animal, and **Fig.**

5c shows all sessions from all animals; *early* canonical correlation: 0.24 ± 0.032, *late*

canonical correlation 0.54 ± 0.035, mixed-effects model: $t(44) = 7.55$, $P = 1.7 \times 10^{-9}$).

Moreover, we found that the subspace activity in the two structures became more

precisely temporally correlated with learning (**Fig. 5d)**. For M1 TR_i's and cerebellar TR_i

's, maximal correlation produced by CCA didn't show a significant increase

(**Supplementary Fig. 4**; 0.37 ± 0.042 to 0.39 ± 0.045, mixed-effects model: $t(44) = 0.52$,

$P = 0.604$).

Figure 5. Increase in neural subspace correlation between M1 and cerebellum. *a,*
Description of canonical correlation analysis (CCA). CCA finds a linear combination of binned
spike counts from M1 units (x_1, x_2, … x_n) and cerebellum units (y_1, y_2, …, y_n) that maximizes the
correlation between M1 and cerebellum. *b,* M1 TR_d and cerebellum TR_i subspace activity (from
the first canonical component) around task-start (−2 to 2 s) for an example session. Each dot
represents one time bin of *early* or *late* trials from the session. Canonical correlation score is given
by *r*. *c,* Change in the canonical correlation score from *early* to *late* trials across all sessions.
***p<0.001. *d,* Relationship between CCA correlation score and average time to a succesful trial
across all sessions.

Cerebellar indirect modulation leads M1 response after task onset

With recent reports of M1 activity being input driven and secondary motor cortex's (M2) modulatory influence over M1 during M1-driven neuroprosthetic task[37,41], we wanted to check how did the cerebellar TR_i activity 'integrate' with M1 TR_d activity. Interestingly, both early and late in learning, the temporal profiles of modulation (i.e., **Fig. 6a)** for M1 TR_d and cerebellar TR_i were different, suggesting that these two distinct groups of task-related units became functionally and temporally coupled with learning and successful task performance. Additionally, we also observed that cerebellum TR_i activity tended to peak before M1 TR_d during early trials (**Fig. 6b**; time to peak for M1 activity: 252.27 ± 15.53 ms and time to peak for cerebellum activity: 228.08 ± 14.18 ms, mixed-effects model: $t(332) = -1.20$, $P = 0.23$) and late trials (**Fig. 6c**; time to peak for M1 activity: 249.92 ± 5.99 ms and time to peak for cerebellum activity: 231.51 ± 5.86 ms, mixed-effects model: $t(332) = -2.19$, $P = 0.02$).

Figure 6. Cerebellum TR_i **activity peaks before M1.** ***a,*** PETH showing the difference of time-to-peak for a M1 TR_d and cerebellum TR_i units during early trials. ***b,*** Same as ***a*** for late trials. ***c,*** Difference between time to peak across all M1 TR_d and cerebellum TR_d units during *early* trials. ***d,*** Same as ***c*** for *late* trials. n.s. indicates p>0.05 *p<0.05.

Cerebellar optogenetic inhibition impairs neuroprosthetic performance

Next, we wanted to see the effects of optogenetic inhibition of cerebellum and the effects

on neuroprosthetic skill learning. We used a red-light shifted halorhodopsin- JAWS for

inhibiting neural activity in the cerebellum (see Methods). First, we found that JAWS was

robustly expressed in cerebellar cortical neurons **(Fig. 7a)**. When we looked at the activity

of cerebellar neurons under optical illumination acutely, we found that JAWS activation

led to strong inhibition of cerebellar neurons (**Supplementary Fig. 5a,b**). Optogenetic inhibition significantly reduced firing across cerebellar neurons (**Supplementary Fig. 5c**; $Stim_{pre}$: 27.30 ± 2.88 Hz, $Stim_{ON}$: 4.99 ± 0.67 Hz and $Stim_{post}$: 20.16 ± 2.09 Hz; $Stim_{pre}$ vs $Stim_{ON}$ mixed-effects model: $t(88) = -7.69$, $P = 1.9 \times 10^{-11}$; $Stim_{post}$ vs $Stim_{ON}$ mixed-effects model: $t(88) = 7.05$, $P = 3.8 \times 10^{-10}$) , with a reduction in 84.44% of recorded cells during $Stim_{ON}$: (n = 45).

Next, we performed optogenetic inhibition of the cerebellum in chronically implanted rats during the BMI task practice (see Methods). When we inhibited the cerebellum in rats that had already gained proficiency in neuroprosthetic task performance (*i.e.,* during *late* trials), we found that time to successful completion of neuroprosthetic task increased (**Fig. 7b**; 5.95 ± 0.68s without cerebellum inhibition and 8.85 ± 0.69 s with cerebellum inhibition, mixed-effects model: $t(22) = 3.24$, $P = 3.7 \times 10^{-3}$). Furthermore, we also found that cerebellar inhibition decreased the MD_Δ of M1 TR_d units during trials when cerebellum was optogenetically inhibited (**Fig. 7c,d**; 190.11 ± 49.86% without cerebellum inhibition and 112.11 ± 17.42% with cerebellum inhibition, mixed-effects model: $t(34) = -1.71$, $P = 1 \times 10^{-2}$). Interestingly, this effect of cerebellar inhibition was not as strong on M1 TR_i units as it was on M1 TR_d units (**Supplementary Fig. 6**; MD_Δ for M1 TR_i units: 48.67 ± 6.90% without cerebellum inhibition and 47.13 ± 4.77% with cerebellum inhibition, mixed-effects model: $t(391) = -0.15$, $P = 0.87$). Cerebellar inhibition also impacted M1 3−6 Hz LFP power. During late trials, 3−6 Hz M1 power was reduced under cerebellar inhibition (**Fig. 7e,f**; 1.29 ± 0.14 z-scored M1 power without cerebellum inhibition and

0.73 ± 0.11 M1 power with cerebellum inhibition, mixed-effects model: $t(22) = -3.30$,

$P = 3.2 \times 10^{-3}$).

While these experiments spoke about loss of performance on the neuroprosthetic task

when cerebellum was optogenetically inhibited, we also observed that the effects of

cerebellar inhibition on neuroprosthetic learning was not related to the order in which it

was performed, i.e., effects of inhibition of cerebellum in first BMI block versus the second

BMI block within a day with the same M1 TR_d units (see Methods for details on the order

of two BMI blocks within a day), **Supplementary Fig. 7b**; time to task completion with

cerebellum inhibition in first BMI block: 9.33 ± 0.86s versus time to task completion

without cerebellum inhibition in second BMI block: 6.34 ± 1.02s, mixed-effects model:

$t(12) = -2.58$, $P = 2 \times 10^{-2}$. **Supplementary Fig. 7c**; Time to task completion without

cerebellum inhibition in first BMI block: 5.54 ± 0.9s versus time to task completion with

cerebellum inhibition in second BMI block: 7.98 ± 1.49s, mixed-effects model: $t(8) = 1.74$,

$P = 1 \times 10^{-2}$). Hence, our results show that cerebellum was actively involved in supporting

rapid M1-driven neuroprosthetic control, whether the animals were new to the BMI control

scheme or when they had become proficient with a given control scheme.

373
374
Figure 7. BMI performance gets impaired with cerebellar inhibition ***a,*** Fluorescence image
of a coronal brain section showing neurons expressing JAWS (green) in the cerebellum cortex.
b, Cerebellar inhibition increases time to task completion. ***p<0.001. ***c,*** PETH of an example M1
TR_d unit, during *late* trials, with (Laser ON) and without (Laser OFF) cerebellar inhibition. ***d,***
Change in modulation depth (MD_Δ) of M1 TR_d units from *early* to *late* trials, with and without
cerebellar inhibition. *p<0.05. ***e,*** Spectrograms of an example M1 LFP channel showing an
absense of 3-6 Hz power during cerebellar inhibition (left) in *late* trials. The 3-6 Hz power emerges
during *late* trials in the same day session where cerebellar inhibition was not done. ***f,*** The 3-6 Hz
power emerges during *late* trials on the same day session where cerebellar inhibition was not
done.

Discussion

In this study, we found an emergent 3-6Hz activity in the M1 and cerebellum LFPs that also modulated task-related direct and indirect spiking in these regions. Additionally, we found that neuroprosthetic learning led to increased synchronous spiking between M1 TR_d and cerebellar TR_i units, and this pair also had a stronger spiking coordination across trials as indicated by heightened canonical correlation with learning. Furthermore, optogenetic inhibition of the cerebellum led to performance impairments (with increased time to task completion) and weakening of M1 TR_d activity. At population spiking level, we also found that cerebellar activity was leading M1 direct activity. These findings suggest that cerebellum plays an important hierarchical role by providing influence on M1 neural patterns that are related to behavioral output. Furthermore, we found that cerebellar indirect activity became more synchronous with M1 direct activity and canonical correlation also increased between M1 TR_d and cerebellum TR_i's, but this was not observed for M1 TR_i and cerebellum TR_i units suggesting that cerebellum had a more privileged relationship with effector-specific neurons of M1. Although we found that M1 TR_i's correlated activity increased with M1 TR_d units, indicating that same-area interactions also participate in achieving neuroprosthetic skill learning, cerebellum TR_i's preferential coordinated activity with only M1 direct units suggests that cerebellum has a strong selective online modulatory influence on M1, as well as instantiating behaviorally relevant patterns. This role of cerebellum that we found in neuroprosthetic skill learning is consistent with cerebellum's role in fine-tuning movement, as well as co-emergent activity patterns that have been reported with learning new skills[2,42]. Our studies here have helped elucidate cerebellar contributions to M1-driven neuroprosthetic learning and

can help improve BMI functionality in the future. For example, BMI paradigms could

incorporate cerebellar indirect signals for improving BMI controllers.

Emergent mesoscopic dynamics across M1-cerebellum

One of our first findings was an emergence of 3-6 Hz coherence in M1- cerebellum LFPs

associated with learning, and neurons in both these regions also showed enhanced

phase-locking to this oscillation during task-relevant periods as revealed through STA.

Similar observations have been observed in a neuroprosthetic study that looked at task-

related cells in cortico-striatal networks[20]. Such coherence can serve to enhance

communication during task period between task-relevant cell populations across the

larger motor networks that should integrate signals for optimal behavioral output. Such

coherent activity may allow for flexible use of task-relevant cells in either region. The

synchrony that we observed in 3-6 Hz band is consistent with other work that has showed

low-frequency coherence between M1 and other motor regions during learning[6,7].

Using BMIs to study cross-region coordination in motor control.

BMI offers investigations of connected regions by selecting target neural patterns (that

dictate output) in one region and concurrent examination of another region as task

performance improves. Implementing this strategy, we aimed to disentangle cerebello-

cortical communication as M1 direct control was learned. Importantly, both M1 and

cerebellum have direct connections to the spinal cord[8–10], and are implicated in movement

control[3,4,6]. In our BMI paradigm, we randomly selected target neural pattern in M1

(enforced by the decoder), which is unlikely to be correlated with processes in other brain

regions. This permitted us to examine how cross-area communication during BMI control

facilitates control. M1 and cerebellum are reciprocally connected[1,2], and while some

extent of coupling of task-related activity in M1 and cerebellum is not surprising, it is

unknown how the indirect cerebellum activity interacts with behavioral output specific

direct activity versus indirect or non-task specific activity of M1 neurons.

One of the striking findings of our work is that canonical correlation increased between

M1 TR_d and cerebellum TR_i units. Similarly, cross-correlograms between these unit pairs

also showed more coincident firing. However, this was not the case for M1 TR_i's and

cerebellum TR_i's. Cerebellum optogenetic inhibition also weakened modulation of M1 TR_d

's which suggests that the cerebellum communicated more strongly with task-specific

pools of M1 direct activity. These ideas are consistent with theoretical studies that

suggest that cerebellar cortex helps in task-relevant dimensionality expansion that aids

in learning[43].

Roles of multiplexed cross-area interactions

Motor control involves signals at longer time scales appropriately integrating with shorter

time scales across several spatially segregated regions to deliver movement precision[44].

Little is known about how these signals at varying spatiotemporal scales are interacting

during motor control. Simultaneous recordings are best suited to understand these

interactions[6,36,36,45–47]. Further, a majority of studies that have used simultaneous

recordings in motor regions use extensively trained animals performing natural motor

tasks[46–49]. Understanding M1-cerebellum interactions in such context raises important

concerns: *(i)* since both M1 and cerebellum are directly controlling movement, it is difficult

to ascertain modulatory influence of one area over the other; and *(ii)* overtrained animals

may have transitioned to an 'automatic' state[50], and it may no longer be suitable to

investigate emergent dynamics in interacting structures. These confounds limit the

inferences about M1-cerebellum interactions in experiments with extensively trained

animals.

Here we have shown that in an M1-driven BMI task, cerebellum activity leads M1 direct

activity and hence, may play an instructional role in the beginning. This idea is consistent

with the notion of cerebellar role in skill acquisition[51]. We find that cerebellar indirect

activity leads later in learning as well, indicative of an ongoing modulatory influence over

M1 to sustain proficiency in the task. This is consistent with the notions of cerebellar

contributing to fine-tuning effector movements[42,52]. We also find that M1 indirect activity

is correlated to M1 direct activity, but not to cerebellar indirect activity. This leads us to

conclude that M1 output-specific neurons contain multiplexed signals coordinated with

both local and distant-area activity and that cerebellar activity might be somatotopically

organized; thus, the cerebellar activity is retained in the cortico-cerebellar loop[53]. Our

cross-correlation between M1 TR_d and cerebellum TR_i units also showed a broader

timescale influence of cerebellum on M1 activity (this is different from the shorter time

scale influences seen between M1 TR_d and M1 TR_i units[11]). Such broader influence of

cerebellar activity may be related to coordination in the larger motor network. Larger

network activity may represent attention regulation[54], motivation[55] or coordination of the

motor task with sensory feedback[56].

Neural dynamics over the course of BMI learning

Natural motor learning is known to involve an early phase marked exploration and high

variability with a transition to late stage when the skill is consolidated[57–60]. Our paradigm

here is focused on early exploratory BMI learning by using mostly single-sessions. BMI

studies that allow for sleep consolidation[11,12,18] or use multiple days of learning[15,61] find

that M1 TR_i units weaken their modulation through the course of extensive training. It is

possible that cerebellar TR_i's may exhibit similar weakening over time. However, local

versus cross-area interactions may differ in the long-term. One recent study had focused

on cross-area activity through multiple days of training, and they found that indirect task-

related modulation persists in several cortical areas[23]. Work that has looked at extensively

trained mice on motor task has shown sustained activity in the cerebellum[42]. There is

further evidence from studies of natural learning that emergent activity in cortico-

cerebellar networks becomes stronger as task proficiency increases[7].

To summarize, our studies leveraged a multiarea BMI paradigm to probe M1-cerebellar

cross-area interactions. We demonstrated that oscillatory dynamics emerged as seen

through LFPs across these regions that also modulated task-related spiking in both areas.

Finer time scale analyses of spiking revealed that cerebellar indirect activity selectively

influenced our imposed artificial target activity pattern within M1. This paradigm allowed

us to manufacture an output-specific pattern, with a relatively small neural footprint to

examine internal motor networks dynamics, removing the constraints of movement

performance. Thus, multiarea BMIs, through such impositions, allows for a more natural

inside-out investigation of cross-area interactions.

Methods

Animal preparation

Adult male Long-Evans rats were used in this study (n = 13, 250–400 g, ~ 8 weeks old, Charles River Laboratories). All animal procedures were performed according to the protocol approved by the Institutional Animal Care and Use Committee at Cedars-Sinai Medical Center, Los Angeles. This ensured that the animals that were used in this research were acquired, cared for, housed, used, and disposed of in compliance with the applicable federal, state, and local laws and regulations, institutional policies and with international conventions to which the United States is a party. Animals were housed on a 14 h light and 10 h dark cycle (Photoperiod is from 6 am to 8 pm) in a climate-controlled vivarium. Out of 13 rats, 8 were used in BMI with simultaneous M1 and cerebellum recordings, 3 rats were used for BMI with cerebellar optogenetic inhibition and the remaining (n = 2) were used in acute cerebellar recording under optogenetic inhibition. Neural probes were implanted during a recovery surgery performed under isofluorane (1–3%) anesthesia. The analgesic regimen included the administration of 0.1 mg per kg body weight buprenorphine, and 5 mg per kg body weight carprofen. Post-operatively, rats were also administered 2 mg per kg body weight dexamethasone and 33 mg per kg body weight sulfatrim for 5 days. Ground and reference screws were implanted posterior to lambda contralateral to the recorded cerebellum, contralateral to the neural recordings. For M1 recordings, 32-channel arrays (33-μm polyamide-coated tungsten microwire arrays) were lowered to a depth of ~1,200–1,500 μm in either the left or right M1 depending on handedness. These were implanted centered at 0.5 mm anterior and 3 mm lateral to the bregma (Ramanathan et al., 2015; Lemke et al., 2019). For cerebellar

recordings we used 32-64 channel tetrodes (Neuronexus, MI) or shuttle-mounted

polytrodes (Cambridge Neurophysiology, UK). The probes were lowered into the

cerebellar cortex through a craniotomy centered at 12.5 mm posterior and 2.5-3 mm

lateral to bregma. Shuttle mounted probes were moved across days and recorded from

depths of 1.5-4 mm. Our target regions were Simplex/ Crus I and Crus II areas of the

cerebellum (Atkins and Apps, 1997; Baker et al., 2001; Heck et al., 2007). Activity in these

areas has shown modulation during upper limb motor behaviors in response to

corticofugal fiber and forelimb stimulation. We did not perform subject-specific

implantation based on motor mapping. However, a subset of rats also performed reaching

tasks and had robust activation during reaching[7].

Viral injections

We used a red-shifted halorhodopsin, Jaws (AAV8-hSyn-Jaws-KGC-GFP-ER2, UNC

Viral Core) for neural silencing in 5 rats for optogenetic experiments[12,18,62]. Viral injections

were done at least 3 weeks before chronic implant surgeries. Rats were anesthetized, as

stated before and body temperature was maintained at 37 °C with a heating pad. Burr

hole craniotomies were performed over injection sites, and the virus was injected using a

Hamilton Syringe with 34G needle. 500-nl injections (100 nl per min) were made at two

sites in the cerebellum (11.5 mm posterior, 2.5 mm lateral to bregma and 11.5 mm

posterior and 3.5 mm lateral to bregma; depth of 1-3 mm). After the injections, the skin

was sutured, and the animals were allowed to recover with same regimen as stated

above. Viral expression was confirmed with fluorescence imaging.

Electrophysiology

Units and LFP activity were recorded using a 128-channel TDT-RZ2 system (Tucker-Davis Technologies). Spike data were sampled at 24,414 Hz and LFP data at 1,017.3 Hz. ZIF (zero insertion force) clip-based digital head stages from TDT were used that interface the ZIF connector and the Intan RHD2000 chip that uses 192x gain. Only clearly identifiable units with good waveforms and high signal-to-noise were used. The remaining neural data was recorded for offline analysis. Behavior related timestamps (i.e. trial onset, trial completion) were sent to the RZ2 analog input channel using an Arduino digital board and synchronized to neural data.

We have used the term *'unit'* to refer to the sorted spike recordings from both the MEA and silicon probe recordings. For both, we initially used an online sorting tool (Synapse, TDT) for neuroprosthetic control. We used waveform shape and the presence of an absolute/relative refractory period in the inter-spike interval (ISI) to judge quality of isolation. Specifically, a voltage-based threshold was set based on visual inspection for each channel that allowed for best separation between putative spikes and noise; typically, this threshold was at least 4 standard deviations (SD) away from the mean. Events were time-stamped and waveforms for each event were peak aligned. K-means clustering was then performed across the entire data matrix of waveforms. Automated sorting was performed by: (1) first over-clustering waveforms using a K-means algorithm (i.e., split into many mini-clusters), (2) followed by a calculation of interface energy (a nonlinear similarity metric that allows for an automated decision of whether mini-clusters

are actually part of the same cluster), and (3) followed by aggregation of similar clusters.

We conducted offline spike sorting in Plexon or Spyking Circus[63].

Behavior

After recovery, animals were typically acclimated for 1-2 day to a custom plexiglass

behavioral box (**Fig. 1a**) prior to the start of experimental sessions. After acclimatization,

rats were water restricted for BMI training. We monitored body weights daily to ensure

that the weight did not drop below 95% of the initial weight.

Behavioral sessions were typically conducted for 1-2 hours. Recorded neural data was

entered in real-time to custom routines in Matlab (R2018b; Mathworks, Natick, MA).

These then served as control signals for the angular velocity of the feeding tube. The rats

performed ~120 trials on an average in a session (e.g. **Fig. 1c**). In a subset of sessions

(n = 14), we also video-monitored the rat during the BMI training using a 30-fps camera

(TDT, USA).

Neural control of the feeding tube

During the BMI training sessions, we typically selected one or two M1 channels' units to

be selected as 'direct' (TR_d) units, and their neural activity was used to control the angular

velocity of the feeding tube. If one channel was choosen for neuroprosthetic control, its

neurons were associated with positive weight and if two channels were chosen, one

channels neurons were associated with a positive weight and another's with a negative

weight. These units maintained their stability throughout the recording as evidenced by

593 stability of waveform shape and interspike-interval histograms (**Supplementary Fig. 1**).

594 We binned the spiking activity into 100ms bins. We then established a mean firing rate

595 for each neuron over a 3–5 minute baseline period. The mean firing rate was then

596 subtracted from its current firing rate at all time points. The specific transform that we

597 used was:

599           $\theta_v = C \times [G_1 \times r_1(i) + G_2 \times r_2(i)]$

601 where θ_v was the angular velocity of the feeding tube, $r_1(i)$ and $r_2(i)$ were firing rates of the

602 direct units. G_1 and G_2 were randomized coefficients that ranged from +1 to −1 and were

603 held constant after initialization. C was a fixed constant that scaled the firing rates to

604 angular velocity. The animals were then allowed to control the feeding tube via modulation

605 of neural activity. The tube started at the same position at the start of each trial (P_1 in **Fig.**

606 **1a**). The calculated angular velocity was added to the previous angular position at each

607 time step (100 ms). During each trial, the angular position could range from −45 to +180

608 degrees. If the tube stayed in the 'target zone' (P_2 in **Fig. 1a;** spanned 10° area) for a

609 period of 400 ms, then a water reward was delivered. In the beginning of a session, most

610 rats were unsuccessful at bringing the feeding tube to position P_2. Most rats steadily

611 improved control and reduced the time to completion of the task during a session. Multiple

612 learning sessions were obtained from each animal. Consistent with past studies, we also

613 found that incorporation of new units into the control scheme required new learning[11,64–

614 66].

Optogenetics

Optogenetic experiments was carried out in JAWS injected rats using a high-power laser (50 W/mm^2: Laserglow Technologies, USA) emitting a 625 nm beam. A subset of rats (n = 3) was implanted with a cannula over Crus I/Crus II region of the cerebellar cortex and a 32-channel microwire array (TDT Florida) was implanted in M1. During the behavior experiments, laser was turned on at every 10 ms at 50% duty cycle to inhibit cerebellar activity after the trial onset for the total duration of the trial (15s). We performed two sessions each day with ~100 trials in each session. Each day, we alternated between turning the laser on in the first session or the second session (**Supplementary Fig. 6a**). Another subset of rats (n = 2) was implanted with fiber-optic cannula mounted on a silicon probe (Cambridge Neurotech, UK) in the cerebellum, and the recordings were performed under isofluorane anesthesia. In every trial, we recorded 5s of baseline followed by 5s for which the laser was on and another 5 s thereafter with laser off. We performed 30 such repetitions.

Histology

After the experiment, rats were deeply anaesthetized with isoflurane (4-5%), then exsanguinated and perfused with 4 % paraformaldehyde (PFA). The brains were extracted and stored in 4% PFA for up to 72 hours. The brains were then transferred to a solution of 30% sucrose and stored for sectioning. We performed sagittal or coronal sections of the brain using a cryostat (Leica, Germany) and stored them in phosphate-buffered saline for imaging. Images were mounted on slices and imaged using a

microscope (Keyence, Japan). The location and depth of the silicon probe in the brain

were traced by DiI depositing on the electrodes prior to their implantation and by looking

afterwards at the fluorescent dye present in the histological slices (**Supplementary Fig.**

2).

Data Analysis

Sessions and changes in performance

Analysis was performed in Matlab (R2020b) with custom-written routines. A total of 23

training sessions recorded from 8 rats were used for our initial analysis. In addition, we

analyzed 12 separate sessions across 3 rats where optogenetic inhibition of the

cerebellum was performed. For **Fig. 1b,c** we compared changes in task performance

across sessions. Specifically, we compared the performance change by calculating the

mean and standard error of the mean (s.e.m) of the time to completion during the first

and last 30% of trials (referred as early and late trials respectively). Furthermore, we also

compared the best performance in early and late trials by calculating the percentage of

unsuccessful trials.

Task-related activity

The distinction between TR_d, TR_i and TU units was based on whether significant

modulation of baseline firing activity after trial onset (i.e., peak of modulation at the time

> 2.5 SD above the baseline period).

LFP analysis

Artifact rejection was first performed on LFP signals to remove broken channels and noisy

trials. LFPs were then z-scored, median referenced and evoked-activity was subtracted

separately for M1 and cerebellum. LFP power was calculated on a trial-by-trial basis and

then averaged across channels and animals, with wavelet decomposition using the

EEGLAB function *newtimef*[67]. M1-Cb LFP coherence was calculated for each pair of

channels using the EEGLAB function *newcrossf*[67]. All the comparisons were done

between early and late trials. For this analysis, across all the early trials, only trials where

time to task completion was over 10 seconds were included. Across all the late trials, only

trials where time to task completion was under 5 seconds were included.

Spike-triggered averaging (STA)

We calculated the STA to measure how spikes locked to the 3-6 Hz LFP oscillations, both

in the M1 and the cerebellum. We used filtered (3–6Hz) median LFP from each region for

this analysis. For filtering, we used the EEGLAB function *eegfilt*. We used the first 4

seconds after the start of the trial to calculate these STAs. For every unit, we

concatenated the spikes from early trials in a spike vector and from late trials in another

vector. Before STA calculation, we equaled the length these vectors. Then, we extracted

2 sec of LFP around every spike-time in those vectors and average it to get early and late

STAs for a given unit. To calculate the change in modulation for every unit, we looked at

the difference between the minimum and maximum peak in a 300 ms window around a

spike in the averaged STA of early and late trails and then calculated this change from

early to late trials in percentage. We also calculated STA during inter-trial interval. Here,

we used from 2 to 6 seconds after the end of the trial and applied the same steps to

compute the STA as described above.

Spike cross-correlation

We computed the cross-correlation histogram (CCH) for M1 TR_d – cerebellar TR_i, M1 TR_i

– cerebellar TR_i and M1 TR_d -M1 TR_i unit pairs. For every trial, the spike counts were

equalized between the pairs before computing the CCH using 10 ms bins. Furthermore,

we constructed pseudo-random spike train CCHs with simulated spike counts of equal

length for every pairing. Such simulations were run 300 times (Monte Carlo

Simulations)[11,68]. To quantify pairwise coincident firing from early to late trials, we took

the mean of the simulated CCH magnitude within ± 400 ms around the center and

subtracted it from the mean of real CCH magnitude in the same window for early and late

trials, respectively (ΔCCH).

Canonical correlation analysis

700 CCA identifies maximally correlated linear combinations between two groups of variables.

701 Unit spiking data in M1 and cerebellum from -2 s to + 2 s around task-start for each trial

702 were binned at 100ms and concatenated. CCA models were then fit using the MATLAB

703 function *canoncorr*. Only the top canonical component was used in this paper.

Video tracking & trajectory analysis

We performed automated tracking of the tip of the feeding tube and the implanted forepaw

of the rats using DeepLabCut[69]. We did cross-correlation between the trajectories of the

feeding tube and the forepaw using *xcorr* function of MATLAB and looked at the average

cross-correlation magnitude ±30ms around zero lag for every trial.

Statistical analysis

All statistical analyses were implemented within MATLAB. The linear mixed-effects model

(implemented using MATLAB *fitlme*) was used to compare the differences in behavior,

trial to mean correlation, speed decoding and canonical correlation unless specified

otherwise. This model accounts for the fact that units or sessions from the same animal

are more correlated than those from different animals and is more stringent than

computing statistical significance over all units and sessions[6,40,70]. We fitted random

intercepts for each rat and reported the *p*-values for the regression coefficients associated

with sessions, channels, or units. We also performed Kruskal-Wallis H-test with multiple

comparisons in **Fig. 3c** and **d**.

Supplementary Figures

727   **Supplementary Figure 1. Single Unit Waveforms and Inter-Spike Interval (ISI).** *a,* Left
728   panel shows waveforms of a representative M1 TR_d unit. Right panel shows ISI
729   histograms for the same unit. *b,* Left panel shows waveforms of a representative task-
730   related M1 TR_i unit. Right panel shows ISI histograms for the corresponding units in the
731   left panel.
732

Supplementary Figure 2. Cerebellum electrode localization. The location of the electrode in lobule 7 of the cerebellum is indicated by a white arrow in the image of a brain section 4.75 mm lateral to the midline.

740
741
742
743

Supplementary Figure 3. Movement correlation reduces between forepaw and feeding tube *a,* Decrease in movement correlation in an example session ***b,*** Correlation reduces from early to late trials. **p<0.01.

Supplementary Figure 4. Canonical correlation analysis for M1 and cerebellum units. Change in CCA correlation shown between M1 and cerebellum task-related indirect (TR_i) units. (n.s.; p>0.5)**.**

Supplementary Figure 5. Inhibition of cerebellar neural activity with optogenetic illumination *a,* A raw trace from an example channel showing reduction in acttvity when the laser was turned on. ***b,*** PSTH of an example unit showing reduction in firnig rate with laser on. ***c,*** Change in the firing rate of all the recorded units before, during and after cerebellar inhibition. ***p<0.001.

Supplementary Figure 6. Effect of cerebellar inhibition on *'indirect'* M1 units
Change in modulation depth (MD_Δ) of M1 TR_i units from *early* to *late* trials, with and
without cerebellar inhibition. [n.s.]p>0.05.

786
787
788

789
790

Supplementary Figure 7. Cerebellar inhibition impairs BMI performace during task acquisition and maintanence. ***a,*** Schematic of session structure. On a day, initial session were performed with cerebellar inhibition trials (Laser ON) followed by subsequent session without cerebellar inhibition (top). Next day, the order was inverted with initial sessions' trials without cerebellar inhibition performed first followed by subsequent trials with cerebellar inhibition (bottom). ***b*** and ***c,*** An increase in the time to task completion was observed irrespective of the order of cerebellar stimulation.

Acknowledgements

We thank Celine Riera and Joshua Burda for resources and reagents for histology, and Pierson Fleischer for insightful feedback during discussion. This research is supported by American Heart Association (postdoctoral fellowship 897265 to A.A. and career development award 847486 to T.G.), National Institutes of Health (R00NS097620 to T.G.), National Science Foundation (CAREER award 2048231 to T.G.) and Cedars-Sinai Medical Center. A.A. also received support from a Cedars-Sinai's Center for Neural Science and Medicine postdoctoral fellowship.

Author Contributions

A.A. and T.G. designed the study. A.A., A.W.F. and N.P.D. carried out the electrophysiology experiments. A.A., A.W.F., N.P.D. and T.G. performed the surgical procedures. A.A. carried out the analysis. A.W.F. and N.P.D. assisted with analysis. A.W.F. and N.P.D. carried out histology. A.A. and T.G. wrote the manuscript. A.W.F. edited the manuscript.

References

1. Kelly, R. M. & Strick, P. L. Cerebellar Loops with Motor Cortex and Prefrontal Cortex of a Nonhuman Primate. *J. Neurosci.* **23**, 8432–8444 (2003).
2. Wagner, M. J. *et al.* Shared Cortex-Cerebellum Dynamics in the Execution and Learning of a Motor Task. *Cell* vol. 177 669-682 e24 (2019).
3. Bloedel, J. R., Bracha, V., Milak, M. & Shimansky, Y. Cerebellar contributions to the acquisition and execution of learned reflex and volitional movements. *Prog Brain Res* vol. 114 499–509 (1997).
4. Heck, D., Kummell, F., Thach W. T. & Aertsena, A. Dynamic Correlation of Neuronal Activity in Rat Cerebellar Cortex Modulated by Behavior. *Annals of the New York Academy of Sciences* vol. 978 156–163 (2002).
5. Ramanathan, D. S., Gulati, T. & Ganguly, K. Sleep-Dependent Reactivation of Ensembles in Motor Cortex Promotes Skill Consolidation. *PLoS Biol* vol. 13 e1002263 (2015).
6. Lemke, S. M., Ramanathan, D. S., Guo, L., Won, S. J. & Ganguly, K. Emergent modular neural control drives coordinated motor actions. *Nat Neurosci* vol. 22 1122–1131 (2019).
7. Fleischer, P. J. *et al.* Emergent low-frequency activity in cortico-cerebellar networks with motor skill learning. 2022.05.18.491978 (2022) doi:10.1101/2022.05.18.491978.
8. Dum, R. P. & Strick, P. L. Motor areas in the frontal lobe of the primate. *Physiol. Behav.* **77**, 677–682 (2002).
9. Nudo, R. J. & Masterton, R. B. Descending pathways to the spinal cord, III: Sites of origin of the corticospinal tract. *J. Comp. Neurol.* **296**, 559–583 (1990).
10. Pompeiano, O. Functional Organization of the Cerebellar Projections to the Spinal Cord. in *Progress in Brain Research* (eds. Fox, C. A. & Snider, R. S.) vol. 25 282–321 (Elsevier, 1967).
11. Gulati, T., Ramanathan, D. S., Wong, C. C. & Ganguly, K. Reactivation of emergent task-related ensembles during slow-wave sleep after neuroprosthetic learning. *Nat Neurosci* vol. 17 1107–13 (2014).
12. Gulati, T., Guo, L., Ramanathan, D. S., Bodepudi, A. & Ganguly, K. Neural reactivations during sleep determine network credit assignment. *Nature Neuroscience* vol. 20 1277–1284 (2017).
13. Arduin, P. J., Fregnac, Y., Shulz, D. E. & Ego-Stengel, V. 'Master' neurons induced by operant conditioning in rat motor cortex during a brain-machine interface task. *J Neurosci* vol. 33 8308–8320 (2013).
14. Sadtler, P. T. *et al.* Neural constraints on learning. *Nature* vol. 512 423–6 (2014).
15. Ganguly, K., Dimitrov, D. F., Wallis, J. D. & Carmena, J. M. Reversible large-scale modification of cortical networks during neuroprosthetic control. *Nat Neurosci* vol. 14 662–667 (2011).
16. Taylor, D. M., Tillery, S. I. & Schwartz, A. B. Direct cortical control of 3D neuroprosthetic devices. *Science* vol. 296 1829–1832 (2002).
17. Athalye, V. R., Ganguly, K., Costa, R. M. & Carmena, J. M. Emergence of Coordinated Neural Dynamics Underlies Neuroprosthetic Learning and Skillful Control. *Neuron* **93**, 955-970.e5 (2017).

862   18.   Kim, J., Gulati, T. & Ganguly, K. Competing Roles of Slow Oscillations and Delta
863   Waves in Memory Consolidation versus Forgetting. *Cell* vol. 179 514-526 e13
864   (2019).
865   19.   Fetz, E. E. Volitional control of neural activity: implications for brain-computer
866   interfaces. *J Physiol* vol. 579 571–579 (2007).
867   20.   Koralek, A. C., Costa, R. M. & Carmena, J. M. Temporally precise cell-specific
868   coherence develops in corticostriatal networks during learning. *Neuron* vol. 79 865–
869   872 (2013).
870   21.   Koralek, A. C., Jin, X., Long 2nd, J. D., Costa, R. M. & Carmena, J. M. Corticostriatal
871   plasticity is necessary for learning intentional neuroprosthetic skills. *Nature* vol. 483
872   331–335 (2012).
873   22.   Clancy, K. B., Koralek, A. C., Costa, R. M., Feldman, D. E. & Carmena, J. M.
874   Volitional modulation of optically recorded calcium signals during neuroprosthetic
875   learning. *Nature Neuroscience* (2014) doi:10.1038/nn.3712.
876   23.   Liu, Z. & Schieber, M. H. Neuronal Activity Distributed in Multiple Cortical Areas
877   during Voluntary Control of the Native Arm or a Brain-Computer Interface. *eNeuro*
878   **7**, (2020).
879   24.   Canolty, R. T. *et al.* Oscillatory phase coupling coordinates anatomically dispersed
880   functional cell assemblies. *Proc Natl Acad Sci U S A* vol. 107 17356–17361 (2010).
881   25.   Buzsaki, G. & Draguhn, A. Neuronal oscillations in cortical networks. *Science* vol.
882   304 1926–1929 (2004).
883   26.   Fröhlich, F. & McCormick, D. A. Endogenous Electric Fields May Guide Neocortical
884   Network Activity. *Neuron* **67**, 129–143 (2010).
885   27.   Tiesinga, P. H. E., Fellous, J.-M., José, J. V. & Sejnowski, T. J. Optimal information
886   transfer in synchronized neocortical neurons. *Neurocomputing* **38–40**, 397–402
887   (2001).
888   28.   Harris, K. D., Csicsvari, J., Hirase, H., Dragoi, G. & Buzsáki, G. Organization of cell
889   assemblies in the hippocampus. *Nature* **424**, 552–556 (2003).
890   29.   Rodriguez, E. *et al.* Perception's shadow: long-distance synchronization of human
891   brain activity. *Nature* **397**, 430–433 (1999).
892   30.   Pesaran, B., Nelson, M. J. & Andersen, R. A. Free choice activates a decision circuit
893   between frontal and parietal cortex. *Nature* **453**, 406–409 (2008).
894   31.   Pesaran, B., Pezaris, J. S., Sahani, M., Mitra, P. P. & Andersen, R. A. Temporal
895   structure in neuronal activity during working memory in macaque parietal cortex.
896   *Nat. Neurosci.* **5**, 805–811 (2002).
897   32.   Meer, M. A. A. van der & Redish, A. D. Theta Phase Precession in Rat Ventral
898   Striatum Links Place and Reward Information. *J. Neurosci.* **31**, 2843–2854 (2011).
899   33.   Fries, P., Womelsdorf, T., Oostenveld, R. & Desimone, R. The Effects of Visual
900   Stimulation and Selective Visual Attention on Rhythmic Neuronal Synchronization in
901   Macaque Area V4. *J. Neurosci.* **28**, 4823–4835 (2008).
902   34.   Minxha, J., Adolphs, R., Fusi, S., Mamelak, A. N. & Rutishauser, U. Flexible
903   recruitment of memory-based choice representations by the human medial frontal
904   cortex. *Science* **368**, eaba3313 (2020).
905   35.   DeCoteau, W. E. *et al.* Learning-related coordination of striatal and hippocampal
906   theta rhythms during acquisition of a procedural maze task. *Proc. Natl. Acad. Sci.*
907   **104**, 5644–5649 (2007).

36. Semedo, J. D., Zandvakili, A., Machens, C. K., Yu, B. M. & Kohn, A. Cortical Areas
Interact through a Communication Subspace. *Neuron* **102**, 249-259.e4 (2019).
37. Veuthey, T. L., Derosier, K., Kondapavulur, S. & Ganguly, K. Single-trial cross-area
neural population dynamics during long-term skill learning. *Nat. Commun.* **11**, 4057
(2020).
38. Gallego, J. A. *et al.* Cortical population activity within a preserved neural manifold
underlies multiple motor behaviors. *Nat. Commun.* **9**, 4233 (2018).
39. Sussillo, D., Churchland, M. M., Kaufman, M. T. & Shenoy, K. V. A neural network
that finds a naturalistic solution for the production of muscle activity. *Nat. Neurosci.*
18, 1025–1033 (2015).
40. Guo, L., Kondapavulur, S., Lemke, S. M., Won, S. J. & Ganguly, K. Coordinated
increase of reliable cortical and striatal ensemble activations during recovery after
stroke. *Cell Rep.* **36**, 109370 (2021).
41. Sauerbrei, B. A. *et al.* Cortical pattern generation during dexterous movement is
input-driven. *Nature* **577**, 386–391 (2020).
42. Becker, M. I. & Person, A. L. Cerebellar Control of Reach Kinematics for Endpoint
Precision. *Neuron* vol. 103 335-348 e5 (2019).
43. Litwin-Kumar, A., Harris, K. D., Axel, R., Sompolinsky, H. & Abbott, L. F. Optimal
Degrees of Synaptic Connectivity. *Neuron* **93**, 1153-1164.e7 (2017).
44. Doya, K. Complementary roles of basal ganglia and cerebellum in learning and
motor control. *Curr. Opin. Neurobiol.* **10**, 732–739 (2000).
45. Perich, M. G. & Rajan, K. Rethinking brain-wide interactions through multi-region
'network of networks' models. *Curr. Opin. Neurobiol.* **65**, 146–151 (2020).
46. Perich, M. G., Gallego, J. A. & Miller, L. E. A Neural Population Mechanism for Rapid
Learning. *Neuron* **100**, 964-976.e7 (2018).
47. Stringer, C. *et al.* Spontaneous behaviors drive multidimensional, brainwide activity.
Science **364**, eaav7893 (2019).
48. Kaufman, M. T., Churchland, M. M., Ryu, S. I. & Shenoy, K. V. Cortical activity in the
null space: permitting preparation without movement. *Nat. Neurosci.* **17**, 440–448
(2014).
49. Musall, S., Kaufman, M. T., Juavinett, A. L., Gluf, S. & Churchland, A. K. Single-trial
neural dynamics are dominated by richly varied movements. *Nat. Neurosci.* **22**,
1677–1686 (2019).
50. Kawai, R. *et al.* Motor Cortex Is Required for Learning but Not for Executing a Motor
Skill. *Neuron* **86**, 800–812 (2015).
51. Cantarero, G. *et al.* Cerebellar Direct Current Stimulation Enhances On-Line Motor
Skill Acquisition through an Effect on Accuracy. *J. Neurosci.* **35**, 3285–3290 (2015).
52. Guo, J.-Z. *et al.* Disrupting cortico-cerebellar communication impairs dexterity. *eLife*
10, e65906 (2021).
53. van der Zwaag, W. *et al.* Digit somatotopy in the human cerebellum: A 7T fMRI
study. *NeuroImage* **67**, 354–362 (2013).
54. Ravizza, S. M. & Ivry, R. B. Comparison of the Basal Ganglia and Cerebellum in
Shifting Attention. *J. Cogn. Neurosci.* **13**, 285–297 (2001).
55. Wagner, M. J. & Luo, L. Neocortex–Cerebellum Circuits for Cognitive Processing.
Trends Neurosci. **43**, 42–54 (2020).

56. Wolpert, D., Ghahramani, Z. & Jordan, M. An internal model for sensorimotor integration. *Science* vol. 269 1880–1882 (1995).

57. Hikosaka, O., Nakamura, K., Sakai, K. & Nakahara, H. Central mechanisms of motor skill learning. *Curr. Opin. Neurobiol.* **12**, 217–222 (2002).

58. Kleim, J. A. *et al.* Cortical Synaptogenesis and Motor Map Reorganization Occur during Late, But Not Early, Phase of Motor Skill Learning. *J. Neurosci.* **24**, 628–633 (2004).

59. Floyer-Lea, A. & Matthews, P. M. Distinguishable Brain Activation Networks for Short- and Long-Term Motor Skill Learning. *J. Neurophysiol.* **94**, 512–518 (2005).

60. Luft, A. R. & Buitrago, M. M. Stages of motor skill learning. *Mol. Neurobiol.* **32**, 205–216 (2005).

61. Silversmith, D. B. *et al.* Plug-and-play control of a brain-computer interface through neural map stabilization. *Nat Biotechnol* (2020) doi:10.1038/s41587-020-0662-5.

62. Chuong, A. S. *et al.* Noninvasive optical inhibition with a red-shifted microbial rhodopsin. *Nature Neuroscience* vol. 17 1123–1129 (2014).

63. Yger, P. *et al.* A spike sorting toolbox for up to thousands of electrodes validated with ground truth recordings in vitro and in vivo. *eLife* **7**, e34518 (2018).

64. Moritz, C. T., Perlmutter, S. I. & Fetz, E. E. Direct control of paralysed muscles by cortical neurons. *Nature* vol. 456 639–642 (2008).

65. Gilja, V. *et al.* A high-performance neural prosthesis enabled by control algorithm design. *Nat Neurosci* vol. 15 1752–1757 (2012).

66. Ganguly, K. & Carmena, J. M. Emergence of a stable cortical map for neuroprosthetic control. *PLoS Biol* vol. 7 e1000153 (2009).

67. Delorme, A. & Makeig, S. EEGLAB: an open source toolbox for analysis of single-trial EEG dynamics including independent component analysis. *J. Neurosci. Methods* **134**, 9–21 (2004).

68. Abbasi, A. *et al.* Epidural cerebellar stimulation drives widespread neural synchrony in the intact and stroke perilesional cortex. *J. NeuroEngineering Rehabil.* **18**, 89 (2021).

69. Mathis, A. *et al.* DeepLabCut: markerless pose estimation of user-defined body parts with deep learning. *Nat Neurosci* vol. 21 1281–1289 (2018).

70. Aarts, E., Verhage, M., Veenvliet, J. V., Dolan, C. V. & van der Sluis, S. A solution to dependency: using multilevel analysis to accommodate nested data. *Nat Neurosci* vol. 17 491–6 (2014).

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

a Canonical Correlation Analysis

x_1 x_2 ... x_n

M1 subspace

Maximally correlated

Cb subspace

y_1 y_2 ... y_n

b

Early Trials

r = 0.29

Late Trials

r = 0.63

Cerebellar suspace activity

M1 supspace activity

c *M1 TR$_d$ - Cerebellar TR$_i$*

***

CCA correlation

E   L

d

R^2 = 0.27
P < 0.01

Time to task completion (s)

CCA correlation

Figure 6

Figure 7

a

1mm

b

Time to task completion (s)

***

Laser OFF   Laser ON

c

M1 TR_d

Firing rate (Hz)

—— Laser OFF
—— Laser ON

Task Start   0.5 s

d

M1 TR_d

Change in MD$_\Delta$ (%)

*

Laser OFF   Laser ON

e

M1 Freq (Hz)

Laser ON   Laser OFF

Power

0.5 s

f

Power

***

Laser OFF   Laser ON

Figure S1

Figure S2

Figure S3

a

b

Figure S4

Figure S5

a

Laser ON

100 μV

1s

b

Laser ON

Firing rate (Hz)

0   Time (s)   15

c

Firing rate (Hz)

***   ***

$Stim_{pre}$ $Stim_{ON}$ $Stim_{post}$

Figure S6

Figure S7

b

Time to task completion (s)

*

Laser ON — Laser OFF

c

Time to task completion (s)

*

Laser OFF — Laser ON

0