Faecal carriage of ESBL-producing *Escherichia coli* in a remote region of Niger

Running Title: Faecal carriage of ESBL-*E. coli* in Niger

Hervé Jacquier[1,2], Bachir Assao[3], Françoise Chau[1], Ousmane Guindo[3], Bénédicte Condamine[1], Mélanie

Magnan[1], Antoine Bridier-Nahmias[1], Nathan Sayingoza-Makombe[3], Aissatou Moumouni[3], Anne-Laure

Page[4], Céline Langendorf[4], Matthew E Coldiron[4], Erick Denamur[1,5,*], Victoire de Lastours[1,6,*]

1. Université Paris Cité, IAME UMR 1137, INSERM, 75018 Paris, France

2. Assistance Publique – Hôpitaux de Paris, Département de Prévention, Diagnostic et Traitement des
Infections, Hôpital Henri Mondor, Créteil, France

3. Epicentre, Médecins Sans Frontières, Maradi, Niger

4. Epicentre, Médecins Sans Frontières, 75019 Paris, France

5. Assistance Publique – Hôpitaux de Paris, Laboratoire de Génétique Moléculaire, Hôpital Universitaire Bichat,
75018 Paris, France

6. Assistance Publique – Hôpitaux de Paris, Service de Médecine Interne, Hôpital Universitaire Beaujon, 92110
Clichy, France

* These authors contributed equally to this work.

Corresponding authors: Hervé Jacquier: herve.jacquier@aphp.fr;

Summary

Objective

Whole genome sequencing (WGS) of extended-spectrum β-lactamase-producing *Escherichia coli*
(ESBL-*E. coli*) in developing countries is lacking. Here we describe the population structure and
molecular characteristics of ESBL-*E. coli* faecal isolates in rural Southern Niger.

Methods

Stools of 383 healthy participants were collected among which 92.4% were ESBL-*E. coli* carriers; 90 of
these ESBL-*E. coli* containing stools (109 ESBL-*E. coli* isolates) were further analysed by WGS, using
short- and long-reads.

Results

Most isolates belonged to the commensalism-adapted phylogroup A (83.5%), with high clonal
diversity. The $bla_{CTX-M-15}$ gene was the major ESBL determinant (98.1%), chromosome-integrated in
approximately 50% of cases, in multiple integration sites. When plasmid-borne, $bla_{CTX-M-15}$ was found
in IncF (57.4%) and IncY plasmids (26.2%). Closely related plasmids were found in different genetic
backgrounds. Genomic environment analysis of $bla_{CTX-M-15}$ in closely related strains argued for
mobilisation between plasmids or from plasmid to chromosome.

Conclusions

Massive prevalence of community faecal carriage of CTX-M-15-producing *E. coli* was observed in a
rural region of Niger due to the spread of highly diverse A phylogroup commensalism-adapted
clones, with frequent chromosomal integration of $bla_{CTX-M-15}$. Plasmid spread was also observed.
These data suggest a risk of sustainable implementation of ESBL in community faecal carriage.

41
42 Keywords: ESBL – *E. coli*, community, developing country, faecal carriage, chromosomal integration

Introduction

Extended-Spectrum β-Lactamase-producing (ESBL) Enterobacterales, and particularly *Escherichia coli* (ESBL-*E. coli*) have spread worldwide and become endemic, especially in low-and middle-income countries [1–5]. The diffusion of ESBL-*E. coli* is mainly the result of the epidemiological success of *bla*$_{CTX-M}$ genes in the 2000s, which are ESBL-encoding genes often carried by mobile genetic elements such as plasmids. *E. coli* are commensals of the gut of vertebrates, including humans [6], and constitute the main reservoir of resistance genes in Enterobacterales, leading to ESBL-*E. coli* dissemination in communities [7]. *E. coli* are also opportunistic pathogens responsible for severe infections such as bloodstream infections, urinary tract infections and in some cases diarrhea [8]. Infections with ESBL-*E. coli* present a clinical challenge due to limited therapeutic options, which is especially crucial in low-income countries [9]. Globally, ESBL-*E. coli* infections represent the biggest burden of bacterial antimicrobial resistance (AMR) morbidity and mortality [10].

Prevalence of ESBL-*E. coli* faecal carriage in humans is a good indicator of the global burden of antibacterial resistance in a given community [11]. In Africa, the few available studies showed a prevalence of ESBL-*E. coli* community faecal carriage ranging from 15% to 92% (higher than in most high income countries) depending on the country, the population studied and the year of stool collection [2–5,12–14]. However, little is known concerning the molecular epidemiology and evolution of resistance mechanisms in low-income countries [13,15]. Yet, this level of understanding is essential to unravel the evolutionary mechanisms as well as the transmission mechanisms enabling the success of ESBL-*E. coli* in community settings. It is crucial to determine specific strategies to prevent further dissemination of these resistant bacteria. We previously reported that 92.4% of healthy individuals carried ESBL-*E. coli* carried in their stools as part of a trial of ciprofloxacin prophylaxis for meningococcal meningitis performed in a rural region of South Niger [14]. Faced with such a high prevalence, we sought to describe the population structure of the strains and the molecular basis of this resistance to understand the mechanisms of resistance spread.

Patients and Methods

Study design

This work was an ancillary study of a clinical trial on ciprofloxacin prophylaxis against bacterial meningitis carried out between April 22, 2017 and May 18, 2017 during a meningococcal meningitis outbreak in the Madarounfa district in rural southern Niger near the border with Nigeria [14].

Population characteristics

Briefly, a subset of 10 villages in the control arm and 10 villages randomized to receive village-wide

prophylaxis in the primary trial were randomly selected to participate in a sub-study examining faecal

carriage of ciprofloxacin-resistant and ESBL-producing *Enterobacterales*, at baseline and after

interventions [14]. In each village, 20 households were randomly selected from a census list. One

household member in each of the selected households was then selected using a simple lottery.

Villages were assigned an identification number and GPS coordinates were collected. Villages were

further described by the presence/absence of a market or a health center, as well as the distance to

the nearest paved road.

Strain collection

In the primary trial, stool samples of 383 individuals were collected at day 0 (prior to ciprofloxacin

administration in the intervention arm). Samples were collected at home by participants and

inoculated into Cary-Blair media (Copan, Brescia, Italy) by study staff. Of them, 99 faecal samples

were randomly selected and were sent to the Infection Antimicrobials Modelling and Evolution

laboratory (IAME, Université Paris Cité) for quality control in agar deep tubes [14]. In Paris, faeces were

plated on cefotaxime containing plates, and colonies of different morphologies were identified by

mass spectrometry (MALDI-TOF-MS Microflex Bruker Daltonics/ BioTyper™ version 2.0). When

colonies with different morphologies on the same plate were identified as *E. coli*, each isolate was

studied independently. Antimicrobial susceptibility testing was performed by the disk diffusion

method according to the EUCAST guidelines, and ESBL detection using the synergy test. Among the

99 samples, 90 samples from 90 subjects carrying at least one confirmed ESBL-*E. coli*, constitute the

material for the analyses performed in this work.

Global genomic analysis

For each *E. coli* isolate, DNA was extracted using NucleoMag DNA Bacteria for DNA purification

(Macherey Nagel, Hoerdt, France) and the whole genome was sequenced using Illumina Hiseq 2x100

bp after NextEra XT library preparation (Illumina, San Diego, CA, USA).

Sequences were analysed with a previously described in-house bioinformatic pipeline

[16](https://github.com/iame-researchCenter/Petanc). Briefly, *de novo* assembling was performed

using SPAdes 3.10.0 [17] and genomes were ordered with Ragout after genetic distance estimation

using Mash [18,19]. Genome typing was performed with *in silico* methods including phylogrouping [20],

multi-locus sequence typing (MLST) (according to the Warwick University and Pasteur Institute)[21],

serotyping [22], and *fim*H typing [23]. Plasmid replicons were determined using PlasmidFinder [24], and the

contig location (plasmid or chromosome) using Plascope [25]. Resistome was analyzed with Resfinder

v3.2 [26], and the virulome with a database based on Virulence Finder [27] and VFDB [28]. The main

virulence factors associated with extra-intestinal or intra-intestinal pathogenic *E. coli* (respectively
ExPEC and InPEC) were as in [29].

A core genome was determined using Roary v3.12 with default parameters [30], and a SNP distance
matrix was computed from the core genome (available in Supplementary Table 1). Then, a phylogeny
based on a maximum likelihood approach using a core genome alignment was performed using Iqtree
v1.6.12 [31]. Patristic distances obtained were used to design the phylogenetic tree with Itol (v6.3)[32].

ESBL-encoded gene support determination
For strains with ESBL-encoded genes embedded in chromosome, the integration sites of $bla_{CTX-M-15}$
was mapped on the linkage map of *E. coli* K-12[33], a A phylogroup strain ST10, assuming synteny wih
the studied isolates. Coordinates established by the completed sequence were expressed as 100
minutes for the entire circular map[33].

For strains with ESBL-encoded genes embedded in plasmids, an additional long-read sequencing was
performed. One representative isolate was selected for isolates with close genetic distances
according to Mash. The Nanopore sequencing (Oxford, UK) was performed using Ligation Sequencing
Kit (SQK-LSK109) with Native Barcoding Expansion 1-12 or 13-24, a FLO--min106 R9.4.1 flowcell, and
a MinION device Mk1B. The reads were filtered using Filtlong (v0.2.0)
(https://github.com/rrwick/Filtlong) to keep reads longer than 1000 nucleotides. The selected reads
were then assembled with the combination of minimap2 (v2.15) and miniasm (v0.3)[34]. The resulting
assembly was then polished through 2 rounds of minipolish (v0.1.0), a convenient wrapper for Racon
(V1.4.11)[35,36]. Ilumina and nanopore contigs were subsequently assembled using Unicycler[37].
Plasmids were compared with BRIG[38] when a scaffold was already available, or circlize R-package[39] in
absence of previously described reference. Plasmids with less than 1% difference in the nucleotide
sequences or ≤2 large insertions or deletions (indels) were considered as highly similar. Genome
annotation was performed using RAST[40].

Ethics

The parent study protocol, which included this sub-study, was reviewed and approved by the
National Consultative Ethics Committee of Niger (Ref: 003/2016/CCNE) and the Ethics Review Board
of Médecins Sans Frontières (Ref: 1603)[14]. The trial is registered at ClinicalTrials.gov (NCT02724046).
Written informed consent was obtained from individual participants.

Results

All raw data concerning antibiotic resistance and genome typing are available in Supplementary

Table 2.

Population characteristics

In this ancillary study, the median age of the 90 participants was 10.5 years [IQR 4-24.8], and the F:M

sex-ratio was 1.5. Only 3 subjects (3.3%) reported antibiotic exposure or hospitalization in the three

months prior to sample collection.

ESBL-*E. coli* isolation and resistance to antibiotics

A total of phenotypically distinct 109 ESBL-*E. coli* were recovered from the 90 samples. Seventy-two

participants carried only one ESBL-*E. coli,* 17 carried two distinct ESBL-*E. coli* and one participant

carried 3 distinct ESBL-*E. coli* by colony morphology (see above).

Among the 109 ESBL-*E. coli* isolates, 58 (53.2%) were resistant to nalidixic acid, 56 (51.4%) to

ciprofloxacin, 23 (21.1%) to gentamicin, 76 (69.7%) to cotrimoxazole, and 85 (78.0%) to tetracycline.

All were susceptible to amikacin and imipenem.

Genomic diversity of ESBL-*E. coli*

The genomic diversity of ESBL-*E. coli* is shown in Table 1 and Figure 1. The ESBL-*E. coli* mainly

belonged to phylogroup A (91/109, 83.5%), phylogroup B1 (8/109, 7.3%) and phylogroup C (6/109,

5.5%). One single isolate was found in each of phylogroups D, F, B2 and clade I.

ST typing using Warwick University scheme highlighted 44 different STs. The principal STs identified

were ST46 (15/109, 13.8%), ST10 (10/109, 9.2%), ST6359 (7/109, 6.4%), ST617 (6/109, 5.5%), and

ST410 (6/109, 5.5%). *In silico* serotyping approaches identified 42 different serotypes, including 11

serotypes with unknown O antigens and new O antigens waiting for assignment for 24/109 (22%)

isolates. The main serotypes were O101:H9 (11/109, 10.1%), O101:H4 (10/109, 9.1%), O101:H10

(9/109, 8.3%), and O9:H9 (8/109, 7.3%). A total of 21 different *fimH* alleles were identified. For

34/109 (31.2%) isolates, no *fimH* gene was found. The main *fimH* alleles identified belonged to

fimH54 (15/109, 13.8%), *fimH34* (11/109, 10.1%), *fimH24* (7/109, 6.4%), and *fimH41* (6/109, 5.5%).

Clone definition and global epidemiology

The definition of clone within the *E. coli* species is not trivial and has been refined at each

technological advance from serotype, MLST, *fimH* typing, to more recently whole genome

sequencing. The ST entity is heterogeneous in terms of divergence, and can encompass several

clones, as exemplified by the emerging ST131 with clades A, B and C having high levels of diversity[8].

Here, the clones were defined according the following strategy. First, isolates were grouped by their

haplogroup, defined as a combination of their sequence types (ST) according to the Warwick

University and Institute Pasteur schemes, their serotype (O:H), and their *fimH* allele. Second, the

distribution of SNPs among isolates was determined, and displayed a SNP cut-off around 120 where

few values were observed (Supplementary Figure 1). Third, SNP distances within and between

haplogroups were calculated and confronted to the cut-off. By using a cut-off at 120 SNPs, most

isolates belonging to a given haplogroup had a SNP distance below this cut-off (Supplementary Figure

1A). A diversity > 120 SNPs was found for few isolates belonging to a given haplogroup: clones 4/5

(683 SNPs), clones 12/14 (2565 SNPs) and clones 34/35 (173 SNPs)(Figure 1). Conversely, the clone

36 was associated with two different haplogroups, due to differences in ST assignation. It should be

noted that strain ST10479 is a single locus variant of strain ST46, highlighting a high degree of

similarity. Similar results have been found with patristic distances (Supplementary Figure 1B). We

thus defined the clone as encompassing isolates with less than 120 SNPs on their core genomes.

Based on this definition, 57 distinct clones were identified among the 109 isolates. Of 109 isolates, 74

(67.9%) belonged to clones represented by two or more isolates. Isolates mainly belonged to clones

36 (10/109, 9.2%), 37 (5/109, 4.6%) and 26 (5/109, 4.6%). Of note, *E. coli* strains with different

morphologies from the same individual were always genetically distinct (Table1, Figure 1).

In summary, despite belonging almost uniquely to phylogroup A, a huge genomic diversity of clones

was observed among the ESBL-*E. coli* isolates.

ESBL-encoding genes and genomic location

The ESBL phenotype was conferred by the $bla_{CTX-M-15}$ gene for 107/109 (98.1%) isolates. The $bla_{CTX-M-15}$

gene was located on the chromosome for 48/107 (44.9%) isolates and on a plasmid for 59/107

(55.1%) isolates. The remaining two isolates had the ESBL phenotype conferred by bla_{SHV-2a} (1 isolate)

and $bla_{CTX-M-55}$ genes (1 isolate), and were both located on a plasmid.

The 48 isolates with a chromosomal $bla_{CTXM-15}$ gene were phylogenetically diverse, belonging to 23

clones. We identified 19 different chromosome integration sites (Figure 1 and Supplementary Table

3). For most isolates, a given chromosome integration site was associated with a given clone (Figure

1). For a single clone (clone 36), we observed two different integration sites (Supplementary Table 3).

Altogether, these data indicate multiple integration events in diverse genetic backgrounds. The

distribution of integration sites on the chromosome of *E. coli* K-12 (phylogroup A) has been mapped,

assuming global synteny between studied isolates and K-12 (Figure 2). Integration sites have been

found in different chromosomal regions, but are not evenly distributed across the genome. Most of

the integration sites are within the Ter macrodomain whereas no integration has been identified in
the Ori macrodomain [41]

For isolates with ESBL-encoding gene located on a plasmid, 10 different replicon types were
identified. The plasmids with bla_{SHV-2a} and $bla_{CTX-M-55}$ genes belonged to IncN and IncX1 incompatibility
groups, respectively. The plasmids harbouring $bla_{CTX-M-15}$ belonged mainly to IncF (IncFIA (5/107),
IncFIB (16/107), IncFIB (H89 phagemid plasmid) (6/107), IncFIC(FII) (8/107)) and IncY (16/107) and, to
a lesser extent, by IncI1 (6/107) incompatibility groups (Table 1, Figure 3).

To better understand the relationships between the clones and the plasmids, we studied the isolates
for which the plasmids showed a strong similarity (see above). In six cases, we found highly similar
plasmids in different genetic backgrounds, whereas 20 plasmids were strictly associated with a single
genetic background (for plasmid alignments, see Supplementary Figure 2). For incFIA plasmids, we
identified similar plasmids, with insertion of 26.7 kbp in the plasmid hosted by clone 8 compared to
that of clone 9. For IncFIB plasmids, we identified similar plasmids in clones 28 and 50, with insertion
of 26.7 kbp in the plasmid hosted by clone 28, and an insertion of 8.8 kbp in the plasmid hosted by
clone 50. Moreover, the plasmid hosted by clone 29 presented a 27.9 kbp insertion compared to that
of clone 45. For IncI1 plasmids, plasmids hosted by clones 16 and 25 presented more than 99%
identity, and the ones hosted by the clones 17, 30, and 56 presented two indels of 2.14 and 1.34 kbp.
For IncY plasmids, we identified similar plasmids, with insertion of 17 kbp in the plasmid hosted by
clone 32 compared to that of clone 37. Altogether, these data suggest plasmid transmission between
clones.

Focus on evolutionary scenarios of $bla_{CTX-M-15}$ gene spread in closely related clones

If our analyses suggest both clonal and plasmid spreads, we wanted to go further in the unravelling
of putative evolutionary scenarios leading to the diffusion of $bla_{CTX-M-15}$ gene in closely related
isolates. Here we focus on two clusters: one with clones 1 to 7, and a second with clones 36 and 37.

Focus on clones 1 to 7

For isolates with $bla_{CTX-M-15}$ gene embedded in the integration site "A" (clones 1 to 5 and 7), we
assume that $bla_{CTX-M-15}$ gene acquisition would have followed the most parsimonious scenario:
isolates with chromosomal $bla_{CTX-M-15}$ gene have diversified after a single event of integration in a
common ancestor. This assumption is based on the fact that the insertion site was the same for all
isolates with chromosomal $bla_{CTX-M-15}$ and that IS*EcpI* upstream $bla_{CTX-M-15}$ is largely deleted (data not
shown). In addition, in the closely-related clone 6, $bla_{CTX-M-15}$ gene was acquired through an IncFIA
plasmid.

Focus on clones 36 and 37

Conversely, in clone 36/ST46, $bla_{CTX-M-15}$ is hosted in two different chromosome integration sites, J (six

isolates) and K (two isolates), suggesting two independent integration events in the same clone. In

addition, in the closely-related clones 36/ST10479 and clone 37, $bla_{CTX-M-15}$ gene was acquired by the

IncFIB and IncY plasmids, respectively.

Analysis of synteny in the regions flanking the $bla_{CTX-M-15}$ gene (Figure 4) displays highly conserved

genetic environment, suggesting transfer between plasmid and chromosome and/or between

plasmids.

Virulence associated genes

We focused on the main virulence factors associated with extra-intestinal or intra-intestinal

pathogenic *E. coli* (respectively ExPEC and InPEC)(19). The average number of ExPEC virulence genes

was low (1.24), the most prevalent being *fuyA* and *irp2* (25/109, 22.9%) (Supplementary Table 2).

These two genes encode the yersiniabactin siderophore and its transporter, and are known to be

part of the high pathogenicity island (HPI)[42]. For phylogroup A isolates, these two genes were

unevenly distributed among the strains, and were mostly found in a cluster of clones (clones 1 to 6,

and clone 8). In the phylogroup C, most isolates (4/6) harboured the adhesin encoding *pap* genes in

addition to *fuyA* and *irp2* genes. One enterotoxinogenic *E. coli* (ETEC) (*eltB* positive) from clade I and

two enteropathogenic *E. coli* (EPEC) from B1 and B2 phylogroups were identified, the latter belonging

to the ST2346 and being a typical EPEC (*eae* and *bfp* positive)[43].

Geographical origin of ESBL-*E. coli* carriers

The geographical distribution of the villages of the carriers of the 109 ESBL-*E. coli* is represented in

Figure 5.

Most isolates belonging to the same clone were not limited to a given geographical area, but were

widely spread across the different villages (such as clones 1, 8, 15, 20, 36, 37, 40, 41, and 42).

Nevertheless, few clones, had a local spread limited to 1 or 2 close villages (such as clones 3, 18, 19,

24, and 26). Last, some clones widely spread in the eastern area, in villages connected by a paved

road and which depend on the same healthcare center (clones 11, 30, 31, 32, 38, 51, and 53).

The isolates 3_13 (belonging to clone 50) and the isolates 3_12 and 3_27 (belonging to clone 28)

harbour plasmids with strong similarity, and were identified in the same village (village 3), suggesting

a putative plasmid transfer between isolates with different genetic backgrounds (Figure 1 and Figure

4).

In sum, we did not find any correlation between geographical localization of the subjects and the
epidemiological genomic data of ESBL-*E. coli*, which suggests a large spread of the most frequent
clones.

Discussion

After finding an unexpectedly high prevalence of ESBL-*E. coli* faecal carriage in healthy subjects
(>90%) in rural South Niger [14], we analysed the population structure, molecular support of resistance
and mode of spread of the resistance in these commensal strains, using state-of-the-art sequencing
techniques with both Illumina WGS and an additional long-read sequencing by Nanopore to
specifically study strains with ESBL-encoded genes embedded in plasmids. Previous studies of ESBL
carriage in African settings has varied by the region, the community and the age of the subjects [2–
5,12,13]. Reported prevalence in Africa has been high, ranging from 46 % in district hospital patients in
South Africa in 2018 [3], 51% in hospitalized patients and 38% in healthy healthcare workers in Chad in
2019 [13], and 59% in healthy children in the community in the Central African Republic [12]. When
resistance mechanisms and phylogenetic groups were described, $bla_{CTX-M-15}$ was the most commonly
encountered gene encoding for resistance to β-lactams (ranging from 75% to 90% of cases), and the
clonal group ST131 was predominant among phylogroup B2 *E. coli* [12,13,15,44].

The main features of our work are the following. First, most ESBL-*E. coli* belonged to phylogroup A
(83.4%), but with a high diversity in terms of STs, serotypes, *fimH* alleles and finally clones, even
though some clones were overrepresented. *E. coli* strains belonging to phylogroup A are known to be
adapted to human gut commensalism lifestyle [6] and have been described as predominant in low-and
middle-income countries [45]. As expected, these strains carry few virulence genes, with the exception
of a cluster of clones (clones 1 to 6, and clone 8), which have accumulated many virulence genes,
especially iron capture systems such as yersiniabactin (*fyuA, irp2*), borne on the HPI (Supplementary
Table 2). ST410 (phylogroup C), an emerging high-risk clone [46], is also well represented, with strains
harbouring the *pap* adhesin encoding genes in addition to the HPI. Both gene clusters have been
associated with ESBL-*E. coli* strains with a longer duration of faecal carriage in travellers returning
from developing countries [47]. On the other hand, B2 phylogroup was represented by a single isolate,
a typical EPEC belonging to the EPEC9 cluster [43,48,49]. This is in contrast to what has been observed in
Western countries, where the proportion of B2 strains among commensal *E. coli* has dramatically
increased in the last four decades, probably under the pressure of several factors such as changes in
dietary habits and improved hygiene [6,50]. Of note, no ST131-*E. coli* isolate (also a B2) was found here,
despite the clone's global epidemiological success [12,13,51]. The pronounced diversity among A strains
and the low prevalence of B2 strains supports the hypothesis that the strains described here are
almost individual-specific, and aligns with the expected composition of commensal *E. coli* strains

from humans living in developing countries. Along the same line, a recent study performed in healthy

pregnant women in Madagascar showed a high prevalence of ESBL-*E. coli* (34%), composed mostly of

isolates belonging to various phylogroup A clones (72%)[52].

Second, we found that the ESBL phenotype was mainly associated with the $bla_{CTX-M-15}$ gene (98%),

which was embedded in the chromosome in approximately 50% of cases, at different integration

sites. The spread of the $bla_{CTX-M-15}$ gene has been well-described in the last decades, even though

other bla_{CTX-M} alleles such as $bla_{CTX-M-14}$ or $bla_{CTX-M-27}$ have had some epidemiological success [2]. As with

most AMR genes, the spread of bla_{CTX-M} genes has been associated with plasmid dissemination,

particularly the pC15-1a plasmid [53]. However, chromosomal locations have increasingly been

described. Chromosome-located bla_{CTX-M} genes were found in 4.8% of ESBL-*E. coli* isolates in a

veterinary collection from Germany, the Netherlands, and the UK [54]. This rate increases to 8.3% and

14.6%, respectively, in clinical isolates recovered from hospitals in Zambia [55] and China [56]. In the

study from Madagascar cited above [52], the bla_{CTX-M} gene was located on the chromosome in 31% of

strains. The highest incidence of chromosomal bla_{CTX-M} gene described so far was observed in Japan,

where $bla_{CTX-M-14}$ was found on the chromosome in 45.9% of cases, and in 16.2% both on

chromosome and plasmid in community commensal strains [57]. The location of bla_{CTX-M} genes on both

chromosome and plasmid probably reflects the ability of their genetic environment to mobilize these

resistance genes. In their model, Hamamoto *et al.* display high rates of $bla_{CTX-M-14}$ mobilization, and

integration in various sites [58], as was observed here with our isolates. Whether a chromosomal or

plasmid location confers a selective advantage for the resistance genes is a matter of debate. In a

mathematical model, Lethtinen *et al.* suggested that plasmid location could be preferred under

selective pressure [59], suggesting that a chromosomal location is more adapted to a lower levels of

selective pressure, as would be the case in community carriage. This hypothesis is also supported by

the fitness cost associated with plasmid-carried resistance genes [60] and by the conservation of

vertical transfer of chromosome-located genes. These assumptions remain challenged by other

authors, as compensatory mutations alleviating plasmid fitness costs and high plasmid-chromosome

beneficial interactions have also been described [60]. From an epidemiological point-of-view, a recent

work suggested that chromosome integration of $bla_{CTX-M-15}$ in ST131 isolates could have contributed

to the clonal expansion of a rare C2 lineage [61].

Third, for isolates with plasmid-located $bla_{CTX-M-15}$, we showed a predominance of IncF plasmids,

known to be well adapted to *Enterobacterales*, followed by IncY plasmids, also reported in

Madagascar. Interestingly, a high similarity was found for some plasmids in different genetic

backgrounds, suggesting both isolate and plasmid diffusion. This observation here in a human

community is in line with a previous study of temporal intestinal carriage of ESBL-*E. coli* in calf farms

[62]. Additionally, we found a similarity between IncFIb plasmid backbones and pCFSAN061762, a

plasmid described in Egypt in an isolate of *E. coli* ST38 (data not shown).

Fourth, careful genetic analysis of the $bla_{CTX-M-15}$ gene in the ST46 and ST46 single locus variant clones

(Figure 4) argues for mobilisation of the ESBL gene with variable numbers of flanking genes between

plasmids and from plasmids to chromosome. This process has been recently suggested in A

phylogroup strains isolated during a "One Health" study on Reunion island [63].

Finally, we did not find any correlation between geographical localization of the subjects (the villages

they live in or their usual social interactions, such as marketplaces) and the epidemiological genomic

data of ESBL-*E. coli*, which suggests a large spread of the most frequent clones. These results may

unveil an underestimated mix of populations in this area, or contamination by a common source of

food or water that could not be captured by mapping health centers, distance to an asphalt road or

marketplaces, as represented in Figure 4.

Our work has some limitations: (i) only a subset of stool samples from the overall study was analysed,

potentially introducing biases, yet the random selection should have reduced this risk; (ii) only the

predominant cefotaxime-resistant *E. coli* isolates present in the faeces were studied, which may not

represent the diversity of *E. coli* isolates in the gut, but does represent the quantitatively

predominant *E. coli* isolates and therefore the ones that constitute the biggest threat in terms of

infectious risks and dissemination [64]; (iii) the search of an ESBL-*E. coli* reservoir in other

environments, such as the food chain (poultry/livestock), water or sediments has not been planned,

impeding the understanding of the routes of dissemination in a "One Health" approach [11].

Despite these limitations, this epidemiologic study in a remote region of Niger is unique and shows

the unprecedented dissemination of ESBL-*E. coli* in the gut of a geographically localized human

population. It suggests the widespread dissemination of multiple resistant clones, possibly fostered

by underestimated inadequate antibiotic treatments and multiple population exchanges that have

probably lasted for years already. From a molecular point of view, this work highlights the diversity of

pathways followed by $bla_{CTX-M-15}$. The high frequency of chromosomal $bla_{CTX-M-15}$ in isolates of *E. coli*

belonging to the commensal phylogroup A is of particular concern because it could be associated

with their sustainable implementation in the community. Understanding the evolutionary forces that

select for such a high prevalence of ESBL-*E. coli* should be major objective.

Altogether, antibiotic resistance in commensal *E. coli* in this remote part of Africa appears to have all

the characteristics of long-term adaptation and all the tools for further dissemination. This is

especially worrying as commensal *E. coli* are responsible for clinical infections and antimicrobial

resistant *E. coli* infections represent the biggest threat in terms of AMR globally [10]. Noteworthy,

community-acquired bacteremia in children hospitalized for severe acute malnutrition was recently

followed in an intensive nutritional rehabilitation center in the same rural district. Approximately

two-thirds of the *E. coli* and *K. pneumoniae* strains (16/26, 61.5%) were ESBL producers, reinforcing

the link between the widespread carriage of ESBL-producing *E. coli* strains and the risk of severe

infection from these same microorganisms [65]. Infections with these strains will be increasingly

difficult to treat, with the need for broader spectrum antibiotics, which in turn may select more

resistant strains such as carbapenemase producers. This work may just be exposing the tip of the

iceberg of an AMR epidemiological catastrophe well on its way.

Acknowledgments

We offer sincere thanks to all the participants of the study, as well as the staff from Maradi
laboratory for their excellent fieldwork, both in terms of sample collection and phenotypic
microbiological analyses. We thank "Médecins Sans Frontières" teams for their trust and
involvement in the parent study as well as this ancillary study. We thank Epicentre and Rebecca
Grais, for trusting our laboratory for the quality control and genetic analyses from the beginning of
this work and for the opportunity to gather an impressive collection of faecal samples from such a
remote region. We are grateful to Olivier Clermont for his expertise in the analysis of the recently
annotated STs.

Financial support

This work was partially supported by a grant from the "Fondation pour la Recherche Médicale"
(Equipe FRM 2016, grant number DEQ20161136698). The parent study was funded by "Médecins
Sans Frontières".
396

397 **Declaration of interest statement**

398 None

References

Karanika Styliani, Karantanos Theodoros, Arvanitis Marios, Grigoras Christos, Mylonakis Eleftherios. Fecal Colonization With Extended-spectrum Beta-lactamase–Producing Enterobacteriaceae and Risk Factors Among Healthy Individuals: A Systematic Review and Metaanalysis. *Clin Infect Dis* 2016;**63**(3):310–8. Doi: 10.1093/cid/ciw283.

Bevan Edward R., Jones Annie M., Hawkey Peter M. Global epidemiology of CTX-M β-lactamases: temporal and geographical shifts in genotype. *J Antimicrob Chemother* 2017;**72**(8):2145–55. Doi: 10.1093/jac/dkx146.

Founou Raspail Carrel, Founou Luria Leslie, Essack Sabiha Yusuf. Extended spectrum beta-lactamase mediated resistance in carriage and clinical gram-negative ESKAPE bacteria: a comparative study between a district and tertiary hospital in South Africa. *Antimicrob Resist Infect Control* 2018;**7**:134. Doi: 10.1186/s13756-018-0423-0.

Letara Nuru, Ngocho James Samwel, Karami Nahid, Msuya Sia E., Nyombi Balthazar, Kassam Nancy A., et al. Prevalence and patient related factors associated with Extended-Spectrum Beta-Lactamase producing *Escherichia coli* and *Klebsiella pneumoniae* carriage and infection among pediatric patients in Tanzania. *Sci Rep* 2021;**11**(1):22759. Doi: 10.1038/s41598-021-02186-2.

Diriba Kuma, Awulachew Ephrem, Tekele Lami, Ashuro Zemachu. Fecal Carriage Rate of Extended-Spectrum Beta-Lactamase-Producing *Escherichia coli* and *Klebsiella pneumoniae* Among Apparently Health Food Handlers in Dilla University Student Cafeteria. *Infect Drug Resist* 2020;**13**:3791–800. Doi: 10.2147/IDR.S269425.

Tenaillon Olivier, Skurnik David, Picard Bertrand, Denamur Erick. The population genetics of commensal *Escherichia coli*. *Nat Rev Microbiol* 2010;**8**(3):207–17. Doi: 10.1038/nrmicro2298.

Woerther Paul-Louis, Burdet Charles, Chachaty Elisabeth, Andremont Antoine. Trends in human fecal carriage of extended-spectrum β-lactamases in the community: toward the globalization of CTX-M. *Clin Microbiol Rev* 2013;**26**(4):744–58. Doi: 10.1128/CMR.00023-13.

Denamur Erick, Clermont Olivier, Bonacorsi Stéphane, Gordon David. The population genetics of pathogenic *Escherichia coli*. *Nat Rev Microbiol* 2021;**19**(1):37–54. Doi: 10.1038/s41579-020-0416-x.

Ouedraogo A. S., Jean Pierre H., Bañuls A. L., Ouédraogo R., Godreuil S. Emergence and spread of antibiotic resistance in West Africa : contributing factors and threat assessment. *Med Sante Trop* 2017;**27**(2):147–54. Doi: 10.1684/mst.2017.0678.

Murray Christopher JL, Ikuta Kevin Shunji, Sharara Fablina, Swetschinski Lucien, Aguilar Gisela Robles, Gray Authia, et al. Global burden of bacterial antimicrobial resistance in 2019: a systematic analysis. *The Lancet* 2022;**399**(10325):629–55. Doi: 10.1016/S0140-6736(21)02724-0.

444 11 Puspandari Nelly, Sunarno Sunarno, Febrianti Tati, Febriyana Dwi, Saraswati Ratih Dian,
445   Rooslamiati Indri, et al. Extended spectrum beta-lactamase-producing *Escherichia coli* surveillance
446   in the human, food chain, and environment sectors: Tricycle project (pilot) in Indonesia. *One*
447   *Health* 2021;**13**:100331. Doi: 10.1016/j.onehlt.2021.100331.
448

449 12 Farra A., Frank T., Tondeur L., Bata P., Gody J. C., Onambele M., et al. High rate of faecal carriage
450   of extended-spectrum β-lactamase-producing Enterobacteriaceae in healthy children in Bangui,
451   Central African Republic. *Clin Microbiol Infect* 2016;**22**(10):891.e1-891.e4. Doi:
452   10.1016/j.cmi.2016.07.001.
453

454 13 Ouchar Mahamat Oumar, Tidjani Abdelsalam, Lounnas Manon, Hide Mallorie, Benavides Julio,
455   Somasse Calèbe, et al. Fecal carriage of extended-spectrum β-lactamase-producing
456   Enterobacteriaceae in hospital and community settings in Chad. *Antimicrob Resist Infect Control*
457   2019;**8**:169. Doi: 10.1186/s13756-019-0626-z.
458

459 14 Coldiron Matthew E., Assao Bachir, Page Anne-Laure, Hitchings Matt D. T., Alcoba Gabriel,
460   Ciglenecki Iza, et al. Single-dose oral ciprofloxacin prophylaxis as a response to a meningococcal
461   meningitis epidemic in the African meningitis belt: A 3-arm, open-label, cluster-randomized trial.
462   *PLoS Med* 2018;**15**(6):e1002593. Doi: 10.1371/journal.pmed.1002593.
463

464 15 Maataoui Naouale, Langendorf Céline, Berthe Fatou, Bayjanov Jumamurat R., van Schaik Willem,
465   Isanaka Sheila, et al. Increased risk of acquisition and transmission of ESBL-producing
466   Enterobacteriaceae in malnourished children exposed to amoxicillin. *J Antimicrob Chemother*
467   2020;**75**(3):709–17. Doi: 10.1093/jac/dkz487.
468

469 16 Bourrel Anne Sophie, Poirel Laurent, Royer Guilhem, Darty Mélanie, Vuillemin Xavier, Kieffer
470   Nicolas, et al. Colistin resistance in Parisian inpatient faecal *Escherichia coli* as the result of two
471   distinct evolutionary pathways. *J Antimicrob Chemother* 2019;**74**(6):1521–30. Doi:
472   10.1093/jac/dkz090.
473

474 17 Bankevich Anton, Nurk Sergey, Antipov Dmitry, Gurevich Alexey A., Dvorkin Mikhail, Kulikov
475   Alexander S., et al. SPAdes: A New Genome Assembly Algorithm and Its Applications to Single-Cell
476   Sequencing. *J Comput Biol* 2012;**19**(5):455–77. Doi: 10.1089/cmb.2012.0021.
477

478 18 Ondov Brian D., Treangen Todd J., Melsted Páll, Mallonee Adam B., Bergman Nicholas H., Koren
479   Sergey, et al. Mash: fast genome and metagenome distance estimation using MinHash. *Genome*
480   *Biol* 2016;**17**(1):132. Doi: 10.1186/s13059-016-0997-x.
481

482 19 Kolmogorov Mikhail, Raney Brian, Paten Benedict, Pham Son. Ragout-a reference-assisted
483   assembly tool for bacterial genomes. *Bioinforma Oxf Engl* 2014;**30**(12):i302-309. Doi:
484   10.1093/bioinformatics/btu280.
485

20 Beghain Johann, Bridier-Nahmias Antoine, Le Nagard Hervé, Denamur Erick, Clermont Olivier.
ClermonTyping: an easy-to-use and accurate in silico method for *Escherichia* genus strain
phylotyping. *Microb Genomics* 2018;**4**(7). Doi: 10.1099/mgen.0.000192.

21 Jolley Keith A., Bray James E., Maiden Martin C. J. Open-access bacterial population genomics:
BIGSdb software, the PubMLST.org website and their applications. *Wellcome Open Res*
2018;**3**:124. Doi: 10.12688/wellcomeopenres.14826.1.
492

493   22 Ingle Danielle J., Valcanis Mary, Kuzevski Alex, Tauschek Marija, Inouye Michael, Stinear Tim, et al.
494   In silico serotyping of E. coli from short read data identifies limited novel O-loci but extensive
495   diversity of O:H serotype combinations within and between pathogenic lineages. *Microb*
496   *Genomics* 2016;**2**(7):e000064. Doi: 10.1099/mgen.0.000064.
497

498   23 Roer Louise, Tchesnokova Veronika, Allesøe Rosa, Muradova Mariya, Chattopadhyay Sujay,
499   Ahrenfeldt Johanne, et al. Development of a Web Tool for *Escherichia coli* Subtyping Based on
500   fimH Alleles. *J Clin Microbiol* 2017;**55**(8):2538–43. Doi: 10.1128/JCM.00737-17.
501

502   24 Carattoli Alessandra, Zankari Ea, García-Fernández Aurora, Voldby Larsen Mette, Lund Ole, Villa
503   Laura, et al. In silico detection and typing of plasmids using PlasmidFinder and plasmid multilocus
504   sequence typing. *Antimicrob Agents Chemother* 2014;**58**(7):3895–903. Doi: 10.1128/AAC.02412-
505   14.
506

507   25 Royer G., Decousser J. W., Branger C., Dubois M., Médigue C., Denamur E., et al. PlaScope: a
508   targeted approach to assess the plasmidome from genome assemblies at the species level. *Microb*
509   *Genomics* 2018;**4**(9). Doi: 10.1099/mgen.0.000211.
510

511   26 Zankari Ea, Hasman Henrik, Cosentino Salvatore, Vestergaard Martin, Rasmussen Simon, Lund Ole,
512   et al. Identification of acquired antimicrobial resistance genes. *J Antimicrob Chemother*
513   2012;**67**(11):2640–4. Doi: 10.1093/jac/dks261.
514

515   27 Joensen Katrine Grimstrup, Scheutz Flemming, Lund Ole, Hasman Henrik, Kaas Rolf S., Nielsen Eva
516   M., et al. Real-time whole-genome sequencing for routine typing, surveillance, and outbreak
517   detection of verotoxigenic *Escherichia coli*. *J Clin Microbiol* 2014;**52**(5):1501–10. Doi:
518   10.1128/JCM.03617-13.
519

520   28 Chen Lihong, Zheng Dandan, Liu Bo, Yang Jian, Jin Qi. VFDB 2016: hierarchical and refined dataset
521   for big data analysis--10 years on. *Nucleic Acids Res* 2016;**44**(D1):D694-697. Doi:
522   10.1093/nar/gkv1239.
523

524   29 Clermont Olivier, Condamine Bénédicte, Dion Sara, Gordon David M., Denamur Erick. The E
525   phylogroup of *Escherichia coli* is highly diverse and mimics the whole *E. coli* species population
526   structure. *Environ Microbiol* 2021;**23**(11):7139–51. Doi: 10.1111/1462-2920.15742.
527

528   30 Page Andrew J., Cummins Carla A., Hunt Martin, Wong Vanessa K., Reuter Sandra, Holden
529   Matthew T. G., et al. Roary: rapid large-scale prokaryote pan genome analysis. *Bioinforma Oxf*
530   *Engl* 2015;**31**(22):3691–3. Doi: 10.1093/bioinformatics/btv421.
531

532 31 Minh Bui Quang, Schmidt Heiko A., Chernomor Olga, Schrempf Dominik, Woodhams Michael D.,
533   von Haeseler Arndt, et al. IQ-TREE 2: New Models and Efficient Methods for Phylogenetic
534   Inference in the Genomic Era. *Mol Biol Evol* 2020;**37**(5):1530–4. Doi: 10.1093/molbev/msaa015.
535

536 32 Letunic Ivica, Bork Peer. Interactive Tree Of Life (iTOL) v5: an online tool for phylogenetic tree
537   display and annotation. *Nucleic Acids Res* 2021;**49**(W1):W293–6. Doi: 10.1093/nar/gkab301.
538

539 33 Berlyn Mary K. B. Linkage Map of *Escherichia coli* K-12, Edition 10: The Traditional Map. *Microbiol*
540   *Mol Biol Rev* 1998;**62**(3):814–984.
541

542 34 Li Heng. Minimap2: pairwise alignment for nucleotide sequences. *Bioinforma Oxf Engl*
543   2018;**34**(18):3094–100. Doi: 10.1093/bioinformatics/bty191.
544

545 35 Vaser Robert, Sović Ivan, Nagarajan Niranjan, Šikić Mile. Fast and accurate de novo genome
546   assembly from long uncorrected reads. *Genome Res* 2017;**27**(5):737–46. Doi:
547   10.1101/gr.214270.116.
548

549 36 Wick Ryan R., Holt Kathryn E. Benchmarking of long-read assemblers for prokaryote whole
550   genome sequencing. *F1000Research* 2019;**8**:2138. Doi: 10.12688/f1000research.21782.4.
551

552 37 Wick Ryan R., Judd Louise M., Gorrie Claire L., Holt Kathryn E. Unicycler: Resolving bacterial
553   genome assemblies from short and long sequencing reads. *PLoS Comput Biol*
554   2017;**13**(6):e1005595. Doi: 10.1371/journal.pcbi.1005595.
555

556 38 Alikhan Nabil-Fareed, Petty Nicola K., Ben Zakour Nouri L., Beatson Scott A. BLAST Ring Image
557   Generator (BRIG): simple prokaryote genome comparisons. *BMC Genomics* 2011;**12**:402. Doi:
558   10.1186/1471-2164-12-402.
559

560 39 Gu Zuguang, Gu Lei, Eils Roland, Schlesner Matthias, Brors Benedikt. circlize Implements and
561   enhances circular visualization in R. *Bioinforma Oxf Engl* 2014;**30**(19):2811–2. Doi:
562   10.1093/bioinformatics/btu393.
563

564 40 Brettin Thomas, Davis James J., Disz Terry, Edwards Robert A., Gerdes Svetlana, Olsen Gary J., et
565   al. RASTtk: a modular and extensible implementation of the RAST algorithm for building custom
566   annotation pipelines and annotating batches of genomes. *Sci Rep* 2015;**5**:8365. Doi:
567   10.1038/srep08365.
568

569 41 Valens Michèle, Penaud Stéphanie, Rossignol Michèle, Cornet François, Boccard Frédéric.
570   Macrodomain organization of the *Escherichia coli* chromosome. *EMBO J* 2004;**23**(21):4330–41.
571   Doi: 10.1038/sj.emboj.7600434.
572

42 Diard Médéric, Garry Louis, Selva Marjorie, Mosser Thomas, Denamur Erick, Matic Ivan.
Pathogenicity-associated islands in extraintestinal pathogenic *Escherichia coli* are fitness elements
involved in intestinal colonization. *J Bacteriol* 2010;**192**(19):4885–93. Doi: 10.1128/JB.00804-10.

43 Hazen Tracy H., Sahl Jason W., Fraser Claire M., Donnenberg Michael S., Scheutz Flemming, Rasko
David A. Refining the pathovar paradigm via phylogenomics of the attaching and effacing
Escherichia coli. Proc Natl Acad Sci 2013;**110**(31):12810–5. Doi: 10.1073/pnas.1306836110.
580

581 44 Moremi Nyambura, Claus Heike, Vogel Ulrich, Mshana Stephen E. Faecal carriage of CTX-M
582   extended-spectrum beta-lactamase-producing Enterobacteriaceae among street children dwelling
583   in Mwanza city, Tanzania. *PloS One* 2017;**12**(9):e0184592. Doi: 10.1371/journal.pone.0184592.
584

585 45 Bailey Jannine K., Pinyon Jeremy L., Anantham Sashindran, Hall Ruth M. Distribution of human
586   commensal *Escherichia coli* phylogenetic groups. *J Clin Microbiol* 2010;**48**(9):3455–6. Doi:
587   10.1128/JCM.00760-10.
588

589 46 Roer Louise, Overballe-Petersen Søren, Hansen Frank, Schønning Kristian, Wang Mikala, Røder
590   Bent L., et al. *Escherichia coli* Sequence Type 410 Is Causing New International High-Risk Clones.
591   *MSphere* 2018;**3**(4):e00337-18. Doi: 10.1128/mSphere.00337-18.
592

593 47 Armand-Lefèvre Laurence, Rondinaud Emilie, Desvillechabrol Dimitri, Mullaert Jimmy, Clermont
594   Olivier, Petitjean Marie, et al. Dynamics of extended-spectrum beta-lactamase-producing
595   Enterobacterales colonization in long-term carriers following travel abroad. *Microb Genomics*
596   n.d.;**7**(7):000576. Doi: 10.1099/mgen.0.000576.
597

598 48 Ingle Danielle J., Levine Myron M., Kotloff Karen L., Holt Kathryn E., Robins-Browne Roy M.
599   Dynamics of antimicrobial resistance in intestinal *Escherichia coli* from children in community
600   settings in South Asia and sub-Saharan Africa. *Nat Microbiol* 2018;**3**(9):1063–73. Doi:
601   10.1038/s41564-018-0217-4.
602

603 49 Hernandes Rodrigo T., Hazen Tracy H., dos Santos Luís F., Richter Taylor K. S., Michalski Jane M.,
604   Rasko David A. Comparative genomic analysis provides insight into the phylogeny and virulence of
605   atypical enteropathogenic *Escherichia coli* strains from Brazil. *PLoS Negl Trop Dis*
606   2020;**14**(6):e0008373. Doi: 10.1371/journal.pntd.0008373.
607

608 50 Massot Méril, Daubié Anne-Sophie, Clermont Olivier, Jauréguy Françoise, Couffignal Camille,
609   Dahbi Ghizlane, et al. Phylogenetic, virulence and antibiotic resistance characteristics of
610   commensal strain populations of *Escherichia coli* from community subjects in the Paris area in
611   2010 and evolution over 30 years. *Microbiol Read Engl* 2016;**162**(4):642–50. Doi:
612   10.1099/mic.0.000242.
613

614 51 Nicolas-Chanoine Marie-Hélène, Bertrand Xavier, Madec Jean-Yves. *Escherichia coli* ST131, an
615   intriguing clonal group. *Clin Microbiol Rev* 2014;**27**(3):543–74. Doi: 10.1128/CMR.00125-13.
616

617 52 Milenkov Milen, Rasoanandrasana Saida, Rahajamanana Lalaina Vonintsoa, Rakotomalala Rivo
618   Solo, Razafindrakoto Catherine Ainamalala, Rafalimanana Christian, et al. Prevalence, Risk Factors,
619   and Genetic Characterization of Extended-Spectrum Beta-Lactamase *Escherichia coli* Isolated From

620   Healthy Pregnant Women in Madagascar. *Front Microbiol* 2021;**12**:786146. Doi:
621   10.3389/fmicb.2021.786146.
622

623   53 Lavollay M., Mamlouk K., Frank T., Akpabie A., Burghoffer B., Ben Redjeb S., et al. Clonal
624   dissemination of a CTX-M-15 beta-lactamase-producing *Escherichia coli* strain in the Paris area,
625   Tunis, and Bangui. *Antimicrob Agents Chemother* 2006;**50**(7):2433–8. Doi: 10.1128/AAC.00150-06.
626

627   54 Rodríguez I., Thomas K., Van Essen A., Schink A.-K., Day M., Chattaway M., et al. Chromosomal
628   location of blaCTX-M genes in clinical isolates of *Escherichia coli* from Germany, The Netherlands
629   and the UK. *Int J Antimicrob Agents* 2014;**43**(6):553–7. Doi: 10.1016/j.ijantimicag.2014.02.019.
630

631   55 Shawa Misheck, Furuta Yoshikazu, Mulenga Gillan, Mubanga Maron, Mulenga Evans, Zorigt
632   Tuvshinzaya, et al. Novel chromosomal insertions of ISEcp1-blaCTX-M-15 and diverse
633   antimicrobial resistance genes in Zambian clinical isolates of Enterobacter cloacae and *Escherichia
634   coli*. *Antimicrob Resist Infect Control* 2021;**10**(1):79. Doi: 10.1186/s13756-021-00941-8.
635

636   56 Zeng Shihan, Luo Jiajun, Li Xiaoyan, Zhuo Chao, Wu Aiwu, Chen Xiankai, et al. Molecular
637   Epidemiology and Characteristics of CTX-M-55 Extended-Spectrum β-Lactamase-Producing
638   *Escherichia coli* From Guangzhou, China. *Front Microbiol* 2021;**12**:3046. Doi:
639   10.3389/fmicb.2021.730012.
640

641   57 Hamamoto Kouta, Ueda Shuhei, Toyosato Takehiko, Yamamoto Yoshimasa, Hirai Itaru. High
642   Prevalence of Chromosomal blaCTX-M-14 in *Escherichia coli* Isolates Possessing blaCTX-M-14.
643   *Antimicrob Agents Chemother* 2016;**60**(4):2582–4. Doi: 10.1128/AAC.00108-16.
644

645   58 Hamamoto Kouta, Hirai Itaru. Characterisation of chromosomally-located blaCTX-M and its
646   surrounding sequence in CTX-M-type extended-spectrum β-lactamase-producing *Escherichia coli*
647   isolates. *J Glob Antimicrob Resist* 2019;**17**:53–7. Doi: 10.1016/j.jgar.2018.11.006.
648

649   59 Lehtinen Sonja, Huisman Jana S., Bonhoeffer Sebastian. Evolutionary mechanisms that determine
650   which bacterial genes are carried on plasmids. *Evol Lett* 2021;**5**(3):290–301. Doi:
651   10.1002/evl3.226.
652

653   60 Rodríguez-Beltrán Jerónimo, DelaFuente Javier, León-Sampedro Ricardo, MacLean R. Craig, San
654   Millán Álvaro. Beyond horizontal gene transfer: the role of plasmids in bacterial evolution. *Nat Rev
655   Microbiol* 2021;**19**(6):347–59. Doi: 10.1038/s41579-020-00497-1.
656

657   61 Ludden Catherine, Decano Arun Gonzales, Jamrozy Dorota, Pickard Derek, Morris Dearbhaile,
658   Parkhill Julian, et al. Genomic surveillance of *Escherichia coli* ST131 identifies local expansion and
659   serial replacement of subclones. *Microb Genomics* n.d.;**6**(4):e000352. Doi:
660   10.1099/mgen.0.000352.
661

662   62 Massot Méril, Châtre Pierre, Condamine Bénédicte, Métayer Véronique, Clermont Olivier, Madec
663   Jean-Yves, et al. Interplay between Bacterial Clones and Plasmids in the Spread of Antibiotic

664   Resistance Genes in the Gut: Lessons from a Temporal Study in Veal Calves. *Appl Environ Microbiol*
665   2021;**87**(24):e0135821. Doi: 10.1128/AEM.01358-21.
666

667 63 Miltgen Guillaume, Martak Daniel, Valot Benoit, Kamus Laure, Garrigos Thomas, Verchere
668   Guillaume, et al. One Health compartmental analysis of ESBL-producing *Escherichia coli* on
669   Reunion Island reveals partitioning between humans and livestock. *J Antimicrob Chemother*
670   2022;**77**(5):1254–62. Doi: 10.1093/jac/dkac054.
671

672 64 Ruppé Etienne, Lixandru Brandusa, Cojocaru Radu, Büke Cagri, Paramythiotou Elisabeth,
673   Angebault Cécile, et al. Relative fecal abundance of extended-spectrum-β-lactamase-producing
674   *Escherichia coli* strains and their occurrence in urinary tract infections in women. *Antimicrob*
675   *Agents Chemother* 2013;**57**(9):4512–7. Doi: 10.1128/AAC.00238-13.
676

677 65 Andersen Christopher T., Langendorf Céline, Garba Souna, Sayinzonga-Makombe Nathan,
678   Mambula Christopher, Mouniaman Isabelle, et al. Risk of community- and hospital-acquired
679   bacteremia and profile of antibiotic resistance in children hospitalized with severe acute
680   malnutrition in Niger. *Int J Infect Dis* 2022:S1201-9712(22)00184-9. Doi:
681   10.1016/j.ijid.2022.03.047.
682
683

Figure 1. Core genome SNP-based phylogenetic tree of 109 ESBL-*E. coli*. The tree is rooted on *E. coli* clade I (isolate 9_02 (clone 47)). From center to periphery, the different layers correspond to the isolate name, the phylogenetic group, the *fimH* allele, the serotype, the sequence type (according the Warwick University scheme), the clone number, the ESBL-type, the chromosome integration site, and the CTX-M carrying plasmid replicon type. In the layer dedicated to ST type, ST10470 and ST13827 (clones 36 and 37) are single locus variant of ST46 and annotated with a star. Squares and stars

are used alternatively to separate isolates. Chromosome integration sites (A to S) are described in Supplementary Table 3.

Figure 2. Location of integration sites of *bla*CTX-M-15 on the linkage map of *E. coli* K-12. Coordinates established by the completed sequence are expressed as 100 minutes for the entire circular map. The height of each bar is proportional to the number of isolates with a given integration site.

Macrodomains are indicated by different colors : Left (13'-26') in yellow, Ter (26'-47') in green, Right (47'-62') in grey and Ori (81'-1') in blue, as described by Valens *et al.* [41].

Figure 3. Prevalence of plasmid replicons harboring ESBL-encoding genes.

Figure 4. Genetic environments of CTX-M-15 encoding genes for four isolates representative of different genetic patterns of ST46 and ST46 single locus variants

For the four isolates, the number of the isolate, the single variant locus of ST46, the clone number, and the chromosome integration site or plasmid type are indicated on the right of the figure.

Figure 2. Geographical location of different clones of ESBL-*E. coli*

House icons represent the locations of the villages. No isolates were recovered from village 10. Villages with a market are bordered in red. Healthcare centres are indicated by a green box with an H. Villages belonging to the same healthcare district appear in the light blue shapes. Distance to the nearest paved road are indicated in blue. For each village, circles represent clones whose area is proportional to the number of isolates. Circles coloured with different colours correspond to clones represented by more than two isolates, those in grey and black to clones represented by two or one isolate, respectively.

Table 1. Characteristics of ESBL-*E. coli* according to genome typing and geographical location

Phylogroup (n)	ST[a] (n)	Serotype[b] (n)	*fimH* allele[c] (n)	Clone[d] (n)	ESBL-typing	Genomic location[e]	Village code[f] (n)
A (91)	ST6359 (7)	O101:H9 (6)	fimH41 (4)	1 (4)	CTX-M-15	chr	1(2), 6, 11
			NF (2)	3 (2)	CTX-M-15	chr	15, 19
		O101:H4 (1)	NF (1)	7 (1)	CTX-M-15	chr	7
	ST617 (6)	O101:H10 (4)	fimH29 (3)	8 (3)	CTX-M-15	IncFIA	6, 8, 13
			NF (1)	9 (1)	CTX-M-15	IncFIA	4
		O101:H9 (2)	NF (2)	4 (1)	CTX-M-15	chr	18
				5 (1)	CTX-M-15	chr	7
	ST13643 (1)	O101:H9 (1)	fimH24 (1)	2 (1)	CTX-M-15	chr	6
	ST1284 (1)	O101:H9 (1)	NF (1)	6 (1)	CTX-M-15	IncFIA	14
	ST167 (1)	O101:H10 (1)	NF (1)	10 (1)	CTX-M-15	IncFIC_FII	3
	ST10 (3)	O101:H10 (3)	fimH215 (3)	11 (3)	CTX-M-15	chr	13, 15, 16
	ST10 (1)	O9:H9 (1)	fimH54 (1)	12 (1)	SHV-12	IncN	5
	ST10 (1)	O101:H9 (1)	fimH54 (1)	13 (1)	CTX-M-15	IncFIC_FII	12
	ST10 (1)	O9:H9 (1)	fimH54 (1)	14 (1)	CTX-M-15	IncFIB	17
	ST44 (2)	O101:H4 (2)	fimH340 (2)	15 (2)	CTX-M-15	IncFIC_FII	8, 15
	ST227 (1)	Ouk:H10 (1)	NF (1)	16 (1)	CTX-M-15	IncI1	12
	ST6143 (1)	Ouk:H10 (1)	NF (1)	17 (1)	CTX-M-15	IncI1	12
	ST34 (2)	O62:H30 (2)	NF (2)	18 (2)	CTX-M-15	IncY	3, 5
	ST1290 (3)	Ouk:H19 (3)	NF (3)	19 (3)	CTX-M-15	chr	7(2), 20
	ST10 (3)	O107:H27 (3)	fimH54 (3)	20 (3)	CTX-M-15	IncFIB_H89	9, 18, 19
	ST10 (1)	OgN12:H32 (1)	fimH23 (1)	21 (1)	CTX-M-15	chr	5
	ST1312 (1)	O25:H16 (1)	fimH198 (1)	22 (1)	CTX-M-15	chr	15
	ST450 (1)	O45:H16 (1)	fimH54 (1)	23 (1)	CTX-M-15	Col156	19
	ST206 (2)	Ouk:H12 (2)	fimH41 (2)	24 (2)	CTX-M-15	IncY	4(2)
	ST215 (1)	O62:H12 (1)	fimH23 (1)	25 (1)	CTX-M-15	IncI1	15
	ST761 (5)	Onv6:H4 (5)	NF (5)	26 (5)	CTX-M-15	IncFIB	2(4), 3
	ST993 (1)	O100:H30 (1)	fimH604 (1)	27 (1)	CTX-M-15	chr	19
	ST165 (4)	Ouk:H26 (4)	fimH54 (3)	28 (3)	CTX-M-15	IncFIB	6, 8, 13
			fimH398 (1)	29 (1)	CTX-M-15	IncFIB	16
	ST1114 (2)	Ouk:H45 (2)	fimH398 (2)	30 (2)	CTX-M-15	IncI1	15, 16
	ST3331 (4)	O155:H9 (4)	fimH25 (4)	31 (4)	CTX-M-15	chr	4, 16(2), 18
	ST540 (2)	O9:H30 (2)	fimH54 (2)	32 (2)	CTX-M-15	IncY	16, 20
	ST361 (1)	O9:H30 (1)	fimH54 (1)	33 (1)	CTX-M-15	chr	1
	ST685 (2)	O8:H43 (2)	fimH25 (2)	34 (1)	CTX-M-15	chr	9
				35 (1)	CTX-M-15	IncY	3
	ST46 (8)	O8:H4 (8)	fimH34 (8)	36 (8)	CTX-M-15	chr	9, 11, 12, 13, 14, 17, 18(2)
	ST10479*(2)	O8:H4 (2)	fimH34 (2)	36 (2)	CTX-M-15	chr	1 (2)
	ST13837*(5)	O9:H4 (5)	NF (5)	37 (5)	CTX-M-15	IncY	3, 8(2), 12, 18
	ST1598 (4)	O9:H4 (4)	NF (4)	38 (4)	CTX-M-15	IncFIB	12, 18, 19(2)
	ST2450 (1)	Ouk:H4 (1)	NF (1)	39 (1)	CTX-M-15	chr	6
	ST2325 (4)	O66:H25 (4)	fimH40 (4)	40 (4)	CTX-M-15	chr	2(2), 11, 19
	ST8406 (2)	O51:H10 (2)	fimH54 (2)	41 (2)	CTX-M-15	chr	5, 13
	ST216 (4)	H55:H4 (4)	fimH69 (4)	42 (4)	CTX-M-15	chr	1, 3, 14, 18
B1 (8)	ST602 (1)	Onv4:H21 (1)	fimH86 (1)	43 (1)	CTX-M-15	IncFIC_FII	17
	ST40 (1)	O21:H21 (1)	NF (1)	44 (1)	CTX-M-15	IncY	20
	ST11075 (1)	O174:H19 (1)	NF (1)	45 (1)	CTX-M-15	IncFIB	6
	ST1431 (1)	O8:H19 (1)	fimH32 (1)	46 (1)	CTX-M-15	IncY	1
	ST448 (1)	Onv12:H8 (1)	fimH35 (1)	47 (1)	CTX-M-15	IncFIB_H89	14
	ST940 (1)	Ouk:H2 (1)	NF (1)	48 (1)	CTX-M-15	chr	4
	ST155 (1)	Onv28:H19 (1)	fimH121 (1)	49 (1)	CTX-M-15	chr	11
	ST58 (1)	O75:H9 (1)	fimH34 (1)	50 (1)	CTX-M-15	IncFIB	3
C (6)	ST410 (3)	Onv14:H9 (2)	fimH24 (2)	51 (2)	CTX-M-15	IncFIC_FII	14, 20
		Onv1:H9 (1)	fimH24 (1)	51 (1)	CTX-M-15	IncFIC_FII	7

	ST410 (1)	O8:H9 (1)	fimH24 (1)	52 (1)	CTX-M-15	chr	18
	ST410 (2)	O8:H21 (2)	fimH24 (2)	53 (2)	CTX-M-15	IncY	12, 14
D (1)	ST38 (1)	O99:H30 (1)	fimH5 (1)	54 (1)	CTX-M-15	chr	5
F (1)	ST648 (1)	O1:H6 (1)	NF (1)	55 (1)	CTX-M-55	IncX1	13
B2 (1)	ST2346 (1)	O142:H34 (1)	fimH57 (1)	56 (1)	CTX-M-15	IncI1	1
cladeI (1)	ST3910 (1)	O153vr2:H33 (1)	fimH552 (1)	57 (1)	CTX-M-15	IncBOKZ	9

[a]STs are assignated according to the Warwick scheme. STNF: ST not found. ST10479* and ST13837* correspond to single locus variants of ST46 in *icd*8 and *fumC*8 alleles, respectively.

[b]Ouk : unknown O antigen, Onv : new O antigen waiting for assignment.

[c]NF : not found.

[d]Isolates that belong to the same clone are designated by a dedicated number.

[e]plas : plasmid, chr : chromosome. When located on the plasmid, the incompatibility group is noticed

[f]villages are designated by a number, and are mapped in Figure 5.