A population modification gene drive targeting both *Saglin* and *Lipophorin*
impairs *Plasmodium* transmission in *Anopheles* mosquitoes

Emily Green, Etienne Jaouen, Dennis Klug, Roenick Proveti Olmo, Amandine Gautier,
Stéphanie Blandin, Eric Marois*

Inserm U1257, CNRS UPR9022, Université de Strasbourg, Strasbourg, France

*Corresponding author: e.marois@unistra.fr

Abstract

Lipophorin is an essential, highly expressed lipid transport protein that is secreted and
circulates in insect hemolymph. We hijacked the *Anopheles coluzzii Lipophorin* gene to
make it co-express a single-chain version of antibody 2A10, which binds sporozoites of
the malaria parasite *Plasmodium falciparum*. The resulting transgenic mosquitoes show
a markedly decreased ability to transmit *Plasmodium berghei* expressing the *P.*
falciparum circumsporozoite protein. To force the spread of this anti-malarial transgene
in a mosquito population, we designed and tested several CRISPR/Cas9-based gene
drives. One of these is installed in, and disrupts, the pro-parasitic gene *Saglin* and also
cleaves wild-type *Lipophorin,* causing the anti-malarial modified *Lipophorin* version to
replace the wild type and hitch-hike together with the *Saglin* drive. Although generating
drive-resistant alleles and showing instability in its gRNA-encoding multiplex array, the
Saglin-based gene drive reached high levels in caged mosquito populations and
efficiently promoted the simultaneous spread of the antimalarial *Lipophorin::Sc2A10*
allele. This combination is expected to decrease parasite transmission via two different
mechanisms. This work contributes to the design of novel strategies to spread
antimalarial transgenes in mosquitoes, and illustrates some expected and unexpected
outcomes encountered when establishing a population modification gene drive.

Introduction

Malaria-transmitting mosquitoes cause over 245 million malaria cases and about
620,000 deaths annually (WHO, 2022). Vector control, mainly in the form of insecticide
indoor residual spraying and insecticide-treated bed nets, has permitted a steady
decrease in malaria cases during the last decades. However, this decline is stalling as
mosquitoes become increasingly resistant to insecticides and as malaria parasites
continue to evolve resistance to drugs. These ongoing genetic changes in vectors and
parasites, under the selection pressure of two major pillars of malaria control, jeopardize
the hard-won successes of the fight against malaria. To address the need for
complementary malaria control approaches, intervention strategies based on intentional
genetic changes in mosquitoes to decrease malaria transmission are under
development. Recent advances in the field of gene drive (GD) provide a mean to push
desirable transgenes to invade a target population of insects (Carballar-Lejarazú and

James, 2017; Gantz and Akbari, 2018; Quinn and Nolan, 2020). GD interventions against malaria mosquitoes are still in the laboratory phase of their development, and can be subdivided in two classes: (a) population suppression GDs that spread female sterility in mosquito populations with the aim to eliminate them (Hammond et al., 2016; Kyrou et al., 2018); (b) population modification GDs that leave mosquito populations in place, but genetically alter them to reduce their disease transmission capacity (Gantz et al., 2015; Pham et al., 2019; Carballar-Lejarazú et al., 2020). Both approaches raise ethical, safety and ecological concerns, and trigger controversy. One concern raised by elimination approaches is their possible consequences on the food web and ecosystems, even where dozens of sympatric mosquito species are potentially available to compensate for the loss of the target species. In some regions, elimination of one species could favor the installation of competing species that could also act as vectors, as can be feared for the *Aedes aegypti* / *Ae. albopictus* mosquito dyad vectoring dengue, Zika and chikungunya viruses. Modification approaches alleviate these concerns by leaving the target species in its ecological niche, but in turn, the indefinite persistence of transgenes utilized in modification drives could have long-term secondary effects, so that both approaches should be planned with utmost caution. However, the option of rejecting gene drive technologies as a complementary tool to fight mosquito-borne pathogens also raises ethical issues, by ignoring a chance to curb the immense mortality and morbidity caused by mosquito-borne infections.

Anopheles gambiae (*sensu lato*), the major mosquito vector of malaria in Sub-Saharan Africa, is widely distributed, locally abundant during rainy seasons and its biomass might significantly contribute to the food chain, in a temporal fashion. Besides their possible negative impact on food webs, elimination approaches for this mosquito are likely to be complicated by re-colonization from residual pockets of surviving isolated populations, resulting in chasing dynamics of gene drive (Champer et al., 2021). For this species in its native ecosystem, population modification to decrease vector competence and preserve ecological functions could be a meaningful gene drive approach. Here, we present the engineering of the *Anopheles coluzzii Lipophorin* (*Lp*) essential gene, which naturally encodes an abundant lipid transport protein secreted in mosquito hemolymph. We modified *Lp* to co-express a single-chain antibody (ScFv) derived from 2A10, a well-established mouse monoclonal antibody that binds the *Plasmodium falciparum* circumsporozoite protein (CSP) (Zavala et al., 1983). Sporozoites are the parasite stage that mosquito females inject in vertebrate hosts while aquiring a blood meal. Once in the host, sporozoites quickly infect the liver and multiply to produce merozoites with a tropism for red blood cells. Through its binding to CSP (the major sporozoite surface antigen), 2A10 was shown to decrease sporozoite infectivity to cultured human hepatocytes (Hollingdale et al., 1984). Transgenic expression of a 2A10 ScFv, fused to the antimicrobial peptide Cecropin A and expressed under the control of the *Vitellogenin* promoter, decreased the number of salivary gland sporozoites in *Anopheles stephensi* (Isaacs et al., 2011; Isaacs et al., 2012). Expression of a 2A10 ScFv directly in the salivary glands of *An. stephensi* reduced sporozoite infectivity to cultured hepatocytes, and transmission to mice of chimeric *Plasmodium berghei* expressing CSP from *P. falciparum* (Sumitani et al., 2013). Likewise, expression of a different CSP-binding ScFv in transgenic *A. coluzzii* decreased the sporozoite's ability to infect cultured hepatocytes and to cause malaria in mice (Triller et al., 2017). Here, we inserted the 2A10 ScFv

coding sequence within the endogenous open reading frame of the essential *Lp* gene, preserving Lp function and exploiting a natural proteolytic cleavage site to separate the fusion protein moieties during their secretion. Using a mouse model of malaria, in which a transgenic rodent *Plasmodium berghei* strain expresses CSP from *P. falciparum*, we show a strong decrease in the ability of the genetically engineered mosquitoes to transmit *Plasmodium*. In a second step, we tested different CRISPR/Cas9-based GD designs to force the spread of the *Lp::Sc2A10* transgene in laboratory mosquito populations. A first approach aimed to boost *Lp::Sc2A10* allelic frequency indirectly, using a suppression GD removing wild-type (WT) *Lp*. This failed, as both WT *Lp* and *Lp::Sc2A10* proved to be largely haploinsufficient in the presence of the *Lp* loss-of-function GD allele. In a second approach we inserted the GD cassette into, and disrupting, the *Saglin* gene, which encodes a *Plasmodium* agonist (Okulate et al., 2007; Ghosh et al., 2009; O'Brochta et al., 2019; Klug et al., 2023). In contrast to *Lp*, the loss of *Saglin* does not by itself impose a large fitness cost on laboratory mosquitoes (Klug et al., 2023). The *Saglin* GD also expressed *Lp* guide RNAs promoting homing of *Lp::Sc2A10* into WT *Lp*. These gRNAs targeting the essential *Lp* gene confer lethality to the *Saglin* GD and prevent it from spreading, unless the cleavage-resistant *Lp::Sc2A10* transgene is present. We show that this GD stimulates dual homing at both the *Lp* and *Saglin* loci, resulting in the simultaneous spread of the two transgenes in caged mosquito populations. These combined modifications are expected to decrease sporozoite transmission by two distinct mechanisms, i.e., depriving *Plasmodium* from the Saglin agonist plus attacking sporozoites with the 2A10 ScFv. GDs are susceptible to the emergence of GD-refractory mutant versions of their target sites (Hammond et al., 2017; Champer et al., 2018; Carballar-Lejarazú et al., 2022). Because our design employs multiplexes of three gRNAs for each of the *Saglin* and the *Lp* loci, we expected most failed homing events to result in loss-of-function *Lp* mutations that would be naturally counter-selected, and/or loss-of-function, GD-resistant *Saglin* alleles that should still contribute to decreasing vector competence. However, at least one functional allele that is refractory to homing was observed at the *Lp* locus.

Results

Development of an engineered Lp allele encoding an anti-sporozoite factor while preserving Lp function

The two most abundant proteins secreted in *Anopheles* hemolymph are the nutrient transporters Lipophorin (Lp) and Vitellogenin (Vg), as illustrated by Coomassie staining of electrophoresis gels revealing both proteins as the major visible bands in hemolymph extracts (Fig. 1A and Rono et al., 2010). While Vg is induced in mosquito females only upon blood feeding and wanes within 3 days as eggs develop, Lp is expressed constitutively, with a further spike of expression after blood feeding. We reasoned that Lp protein abundance, secretion in hemolymph and persistence of expression outside gonotrophic cycles make *Lp* a particularly attractive host gene to co-express and secrete high levels of synthetic anti-*Plasmodium* factors. Here we tested a 2A10 scFv version without fusion to Cecropin as antimalarial effector, and hereinafter refer to this effector as Sc2A10. Using the CRISPR-Cas9 system, we knocked-in codon-optimized *Sc2A10* in

frame after the endogenous first *Lp* coding exon, which encodes the Lp secretion signal
peptide (Fig. 1B, for sequence of the plasmid used for knock-in see Suppl. file 1A). As a
result, transcription of the chimeric *Lp::Sc2A10* mRNA encodes the natural Lp signal
peptide followed by Sc2A10 fused to the remainder of the Lp protein sequence. A
natural proteolytic cleavage site located between the ApoLpII and ApoLpI subunits of the
Lp polypeptide allows their separation during maturation in the Golgi apparatus. We
duplicated this motif to detach Sc2A10 from ApoLpII during protein maturation. The
knock-in design also introduced silent mutations in the majority of possible Cas9 sgRNA
target sites within *Lp* exon 1 and the beginning of intron 1, to facilitate selection of
multiple sgRNAs targeting WT *Lp*, but not its engineered allele, when designing
subsequent GDs. Finally, the 1644-bp first intron immediately following *Lp* exon 1
offered an ideal location to host a fluorescent selection marker to identify and track the
modified gene. We placed a *GFP* marker gene under control of the synthetic *3xP3*
promoter (Horn and Wimmer, 2000) and *D. melanogaster Tubulin56D* terminator in this
intron, with care taken to preserve the intron splice junctions and in reverse orientation
relative to *Lp* transcription so that the *Tub56D* terminator would not cause premature
arrest of *Lp* mRNA transcription. The *Lp* promoter is active in the fat body, while the
3xP3 promoter is active in the nervous system. These synthetic modifications in the *Lp*
locus are schematized in Figure 1B and annotated in the provided sequence file (Suppl.
file 1B).
Using automated flow cytometry-based (COPAS) selection of live neonate larvae, based
on differential GFP expression between homozygotes and heterozygotes (Marois et al.,
2012), we established two stocks of *Lp::Sc2A10* mosquitoes, one heterozygous, the
other homozygous. The homozygous stock was fertile and viable, devoid of obvious
fitness cost. This indicates that the transgenic insertion into *Lp* did not disrupt its
essential functions in development, physiology, flight or reproduction. A more careful
evaluation of *Lp::Sc2A10* fitness costs was conducted by following the dynamics of the
transgene in the heterozygous population over 17 generations. This revealed slow
disappearance of the transgene over time (Figure 1C and Suppl. Table 1A) indicative of
a subtle fitness cost that decreased transgene frequency by an average of 2.3% each
generation. In fertility assays comparing WT and homozygous transgenic females, the
latter consistently engendered fewer progeny than their WT sisters, pointing to a
negative impact of the *Lp* modification on their reproductive capacity (Suppl. Table 1B).

The Lp::Sc2A10 *transgene directs Sc2A10 secretion into mosquito hemolymph*

To test whether the Sc2A10 single-chain antibody was expressed and secreted in
hemolymph via the Lp signal peptide, we extracted hemolymph from transgenic female
mosquitoes and analyzed its protein content by mass spectrometry. Because the
Lp::Sc2A10 gene fusion produces a single mRNA, with the endogenous Lp signal
peptide addressing the entire fusion protein to the secretory pathway, the Sc2A10
protein fragment must initially be stoechiometric to the Lp subunits ApoLpI and ApoLpII.
Upon proteolytic cleavage dissociating Sc2A10, ApoLpII and ApoLpI and after their
secretion into the hemolymph, each moiety possibly follows a different fate. In two mass
spectrometry samples from the hemolymph of homozygous transgenic mosquitoes, for a

total of 2629 peptides from the largest Lp subunit ApoLpI, we detected 672 peptides from smaller ApoLpII and 18 from Sc2A10 (Table 1). Corrected for size, this suggests that Sc2A10 molecules are 10 to 20-fold less abundant in the hemolymph compared to ApoLpI and ApoLpII. This observation suggests that while Sc2A10 is made and secreted into hemolymph, its disappearance by degradation, uptake in cells or stickiness to tissue, is faster than that of its sister Lp proteins. Of note, western blots using anti-ApoLpII antibodies on hemolymph samples from homozygous transgenic mosquitoes showed no size upshift for ApoLpII, confirming that Sc2A10 is properly released from ApoLpII (Fig. 1D). Overall we observed higher Lp concentrations in heterozygous compared to homozygous mosquitoes, and a 3 to 4-fold lower Sc2A10 : Lp ratio in heterozygotes compared to homozygotes, indicating that the chimeric proteins from *Lp::Sc2A10* are less expressed than from WT *Lp*.

Sample	ApoLpI	ApoLpII	Sc2A10
WT control	2075 (5.8)	486 (5.1)	0 (0)
hom Sc2A10, sample 1	1627 (5.1)	405 (4.8)	11 (0.4)
hom Sc2A10, sample 2	1002 (4.4)	267 (4.4)	7 (0.3)
het Sc2A10, sample 1	1557 (10.1)	351 (8.7)	2 (0.1)
het Sc2A10, sample 2	2290 (6.0)	664 (6.7)	2 (0.1)

Table 1: Mass spectrometry identification of peptides from Lipophorin and Sc2A10 in hemolymph from transgenic mosquitoes and wild-type sibling controls. The table shows the total number of peptide spectra detected in each hemolymph sample for the two Lp subunits and for Sc2A10. Hemolymph was collected from homozygous (hom) and heterozygous (het) transgenic mosquitoes. Relative, normalized abundance of these peptides in the protein sample, calculated by dividing their spectral count by the respective protein's molecular weight (kDa) and by the total spectral counts from known abundant hemolymph proteins found in the sample (ApoLpIII, APL1C, LRIM1, Nimrod, Phenoloxidase, and TEP1), is indicated in parentheses. Resulting values were multiplied by a constant factor equal to the average number of spectra from these control proteins across samples.

The Lp::Sc2A10 *transgene decreases* Plasmodium *transmission*

To assess whether the amount of Sc2A10 present in hemolymph can affect *Plasmodium* transmission, we infected homozygous transgenic mosquitoes and their control wild-type siblings and allowed parasites to develop for 16-20 days, before exposing naïve mice to be bitten by groups of 10 infected mosquitoes per mouse. The *Plasmodium berghei* strain we employed, *Pb-PfCSP*[hsp70-GFP], expresses GFP at a high level under control of the constitutive *hsp70* promoter and has its native CSP substituted with that from *P. falciparum* (Manzoni et al., 2014; Triller et al., 2017 and see Methods), which is the antigen recognized by 2A10. One day before biting naïve mice, the cold-anesthetized mosquitoes were examined under 488-nm light. Those lacking visible GFP parasites (oocysts in the abdomen, sporozoites visible through the cuticle in wing veins or salivary glands) were discarded to ensure that mice were exposed only to *Plasmodium*-infected

mosquitoes. We controlled that each mouse was bitten by at least 6 infected mosquitoes, with exceptions indicated in Suppl. Table 2, by counting the blood-engorged females.

Consistent with previous reports (Isaacs et al., 2011; Isaacs et al., 2012; Sumitani et al., 2013), Sc2A10 significantly reduced transmission of *Pb-PfCSP* (p<0.0001). Only 29.7% (11/37) of mice exposed to *Lp::Sc2A10* homozygous infected mosquitoes developed parasitemia, compared to 97.1% (33/34) exposed to infected control mosquitoes (Fig. 2A and Suppl. Table 2). Additionally, 6 of the 11 mice that did become infected by transgenic mosquitoes experienced a 1-day delay in the development of infection (Suppl. Table 2), indicative of reduced liver infection (Reuling et al., 2020).

We examined if the transmission-reducing activity of the *Lp::Sc2A10* transgene is dependent on the dosage of Sc2A10, i.e., whether a single copy of the transgene would be as efficient as two. For this we exposed naïve mice to groups of 10 infected mosquitoes carrying zero, one or two *Lp::Sc2A10* copies. While 17 out of 18 mice bitten by control mosquitoes became infected, only 4 out of 12 mice bitten by heterozygous mosquitoes became infected (Fig. 2B and Suppl. Table 2). Two-tailed Fisher's exact test shows a statistically significant reduction when comparing wild-type and heterozygous (p=0.0006) mosquitoes regarding their *Plasmodium*-transmitting capacity.

To verify that the transmission-blocking phenotype of the *Lp::Sc2A10* transgene was specific to CSP from *P. falciparum*, we performed a control bite-back experiment comparing *Pb-PfCSP*[hsp70-GFP] to its *Pb-hsp70-GFP* parental parasite strain expressing endogenous *PbCSP*. While *Lp::Sc2A10* and wild-type mosquitoes transmitted the control *P. berghei* strain to mice with equal efficiencies (n=5/5 and 4/4 mice infected, respectively), only the WT mosquitoes transmitted *Pb-PfCSP*[hsp70-GFP] (n=3/3 mice infected), whereas *Lp::Sc2A10* mosquitoes blocked transmission of the *Pf*CSP-expressing strain (n=0/4 mice infected). This indicates that the transmission blocking capacity of the *Sc2A10* transgene is indeed *Pf*CSP-specific (Fig. 2C).

Testing a suppression gene drive homing into wild-type Lp *to push* Lp::Sc2A10 *toward fixation*

To push the *Lp::Sc2A10* anti-*Plasmodium* gene towards fixation in a mosquito population, our first strategy was to employ a transiently acting Cas9-based GD destroying the wild-type *Lp* gene (Suppl. Fig. 1A). This aimed to exploit the natural tendency of suppression GDs that target an essential gene to initially spread in the population only to be supplanted, over time, by GD-immune functional mutants of the target gene (Hammond et al., 2017). Here *Lp::Sc2A10* itself, lacking the gRNA target sites, would play the role of a GD-immune, functional *Lp* mutant allele. The advantage of this approach was that a *Cas9* and gRNA-carrying transgenic GD cassette inserted in *Lp* need not persist indefinitely in the population, being supplanted by the technologically more benign *Lp::Sc2A10* transgene. To minimize the spontaneous emergence of undesirable additional GD-immune and functional *Lp* alleles, we used a multiplex of 4 gRNAs targeting *Lp*. The rationale was that in cases of failed GD homing, multiple Cas9

cuts repaired by non-homologous end joining (NHEJ) would mainly generate loss-of-
function alleles in the essential *Lp* gene (e.g., deletions between Cas9 cut sites,
frameshifts at several individual cut sites) doomed to be eliminated by natural selection.
We constructed a DsRed$_{NLS}$-marked *Lp* suppression GD (Suppl. Fig. 1A and annotated
sequence in Supplementary File 1C), in which an array of 4 gRNAs separated by tRNAs
(Xie et al., 2015; Port and Bullock, 2016), targeting *Lp* but not *Lp::Sc2A10*, were
expressed under control of a single U6 promoter. *Cas9* was placed under the control of
zpg regulatory elements, currently considered best for use in suppression GDs due to
tighter restriction of Cas9 activity to the germ line (Kyrou et al., 2018; Hammond et al.,
2021). Knocking-in this GD into the *Lp* locus was achieved by injecting this plasmid into
>800 mosquito eggs carrying a heterozygous copy of GFP-marked *Lp::Sc2A10*, in order
to retain a protected functional copy of *Lp* during the disruption of wild-type *Lp* by Cas9.
Genomic insertion in injected embryos relied on expression of the construct's own
CRISPR/Cas9 components supplemented with an additional heterozygous copy of a
YFP-marked *vasa-Cas9* transgene in the embryos.

By screening for the presence of red fluorescence in the G1 progeny of about 140
surviving injected G0 mosquitoes, we recovered several dozens of knock-in G1 larvae
(most of which were DsRed$_{NLS}$ and GFP positive) and examined transgene inheritance
by crossing them to wild-type mosquitoes. In this cross, DsRed$_{NLS}$ marking the *Lp-GD*
construct was expected to strictly segregate away from GFP marking the *Lp::Sc2A10*
allele. Puzzlingly however, the majority of knock-in larvae contained a compound *Lp*
locus harboring the DsRed$_{NLS}$-marked GD cassette strictly linked to the GFP-marked
Lp::Sc2A10 transgene. Sequencing PCR products spanning parts of this insertion
confirmed that, instead of the expected insertion of the *Lp-GD* cassette, a complex
chromosomal rearrangement (Suppl. Fig. 1B) had repaired the Cas9-cleaved wild-type
chromosome by incorporating the entire injected plasmid, and by recombining with the
intact sister chromosome bearing the *Lp::Sc2A10* transgene. The resulting unintended
transgenic modification at the *Lp* locus harbored all the components of a classical
payload-bearing GD that we hereinafter refer to as *GFP-RFP*. Similarly to the
Lp::Sc2A10 allele, the *GFP-RFP Lp* allele was homozygous viable, functional Lp protein
being expressed from its *Lp::Sc2A10* moiety. In spite of the presence of the *zpg-Cas9*
and *gRNA*-encoding cassettes, the *GFP-RFP* locus showed little homing activity after
crosses to WT, except for the appearance of rare GFP-only or RFP-only progeny larvae,
raising concern about the efficiency of the *zpg-Cas9* and/or the gRNA-expressing
cassettes. Providing an independent copy of *vasa-Cas9* on a different chromosome
rescued this defect and resulted in high homing rates of various segments from the
GFP-RFP cassette (mostly GFP alone, more rarely DsRed$_{NLS}$ alone or both) (Suppl. Fig.
2A), indicating that at least some of the gRNAs in the tRNA-spaced array were active
and that the defect lied in *zpg-Cas9* expression.

Fortunately, we also recovered a minority of knock-in larvae that had integrated the *Lp-*
GD construct as expected, marked with DsRed$_{NLS}$ only (Suppl. Fig. 1A). However, no
evidence of gene drive was observed in further generations, as DsRed inheritance did
not exceed 50% when crossing heterozygotes to WT. On the contrary, DsRed frequency
dropped rapidly in F2 and F3 generations, indicating that the non-functional GD cassette
inserted in *Lp* conferred a strong fitness cost. Most heterozygous mosquitoes carrying

one copy of *Lp-GD* and either *GFP-RFP* or *Lp::Sc2A10* (both of which should be refractory to homing of the GD construct) showed a striking developmental delay and died before adulthood (Suppl. Fig. 2B). Heterozygous mosquitoes carrying *Lp-GD* / WT survived to adulthood but were small and short-lived. These observations indicate that WT *Lp* is largely haplo-insufficient in presence of a *Lp* loss-of-function allele, even more so if WT *Lp* is replaced by *Lp::Sc2A10*. This last point is consistent with the observed fitness cost and lower *Lp* expression from the *Lp::Sc2A10* allele, that already suggested that albeit functional, this allele is inferior to the wild-type.

From the haplo-insufficiency of *Lp* we concluded that it was impossible to create a loss-of function GD in this locus, precluding an indirect same-locus GD approach. In addition, the *zpg* promoter appeared to have low activity when integrated in the *Lp* locus.

Driving Lp::Sc2A10 distantly from Saglin with a zpg-Cas9 gene drive

To host a GD promoting the spread of *Lp::Sc2A10*, we turned our attention to the mosquito *Saglin* locus. Indeed, while not leading to complete *Plasmodium* transmission blockage, the loss of *Saglin* reduces parasite prevalence and loads, reduces transmission, is homozygous viable and has no fitness cost in laboratory mosquitoes (Klug et al., 2023). Similar to an antimicrobial combination therapy, combining Sc2A10 expression with the knockout of *Saglin* can be expected to reinforce the transmission-blocking properties of each strategy taken alone. Therefore, we sought to disrupt *Saglin* with a gene drive cassette that would home into wild-type *Saglin* and concomitantly promote *Lp::Sc2A10* homing into the *Lp* locus (Fig. 3A).

The *zpg* promoter still seemed an attractive choice to build a GD dually targeting *Saglin* and *Lp*, to avoid drawbacks from other characterized germline promoters (high maternal Cas9 deposition resulting in failed homing events in the zygote accompanied by the formation of GD refractory alleles; possible somatic mosaic loss-of-function of wild-type *Lp* leading to haplo-insufficiency in heterozygous [*Lp::Sc2A10* /+] mosquitoes). To exclude that the low *zpg-Cas9* activity we previously observed at the *Lp* locus was due to overlooked mutations in our construct, we subcloned the *zpg-Cas9* cassette from a validated GD construct (Kyrou et al., 2018) and incorporated it in a *Saglin* knock-in plasmid expressing Cas9, DsRed$_{NLS}$, 3 gRNAs directed against *Saglin* and 4 gRNAs against *Lp* (Fig. 3, and annotated sequence of this plasmid in Suppl. File 1D). From injection of this plasmid into heterozygous *Lp::Sc2A10* / WT embryos, we crossed 36 G0 male survivors to WT, and recovered more than 60 transgenic G1 individuals. The sole source of Cas9 in this injection was the knock-in GD plasmid itself, showing that *zpg-Cas9* was efficiently active at least for initial insertion. 100% of the G1 transgenics arising from injected males were females, an indication that the construct had correctly homed onto the X chromosome, where *Saglin* is located. In addition, all G1 transgenics had also inherited a copy of *Lp::Sc2A10* (GFP positive), instead of the expected 50%. This suggests that the GD construct acted on the *Lp* locus, causing either efficient *Lp::Sc2A10* homing already in the germ line of G0 males, and/or the death of larvae that did not inherit a protected *Lp* copy due to the toxicity of the construct's *Lp* gRNAs.

In spite of the efficiency of *zpg-Cas9* for initial integration, this first *Saglin* GD (termed
SagGD^{zpg}) showed disappointingly modest homing levels at both the *Saglin* and *Lp* loci
in the G2 generation (Fig. 4A). Using COPAS flow cytometry, we examined GFP and
DsRed inheritance in 2327 larvae that arose from the pooled G1 females crossed to WT
males. It showed 57.7% GFP inheritance (*Lp::Sc2A10* locus) compared to 50%
expected in the absence of homing. DsRed inheritance was more difficult to assess, as
the low intensity of red fluorescence did not allow accurate gating to separate red
fluorescent from negative larvae on COPAS diagrams. Approximate gating completed
with visual examination of a sample of 200 larvae suggested 54.1% DsRed inheritance
at the *Saglin* locus, an even more modest inheritance bias than for GFP at the *Lp* locus.
Mild homing could be explained by position effects affecting the activity of the *zpg*
promoter, by a lack of recombination at the target loci, or by inefficiency of gRNAs. To
distinguish among these possibilities, we generated females carrying single
heterozygous copies of the three unlinked transgenes *SagGD^{zpg}*, *Lp::Sc2A10* and a non-
fluorescent (puromycin resistance-marked) *vasa-Cas9*. The progeny of these triple-
transgenic females crossed to WT males not only showed markedly better homing rates
(85% GFP inheritance) but also evidence of homing in the zygote (6% progeny
appeared to be GFP homozygous) (Fig. 4B). Thus, *vasa-Cas*9 was able to significantly
rescue the modest homing rate of *zpg-Cas9* from *SagGD^{zpg}*, resulting in an efficient split
GD. This suggests that the cloned *zpg* promoter sequence is sensitive to positional
effects, being poorly active when inserted in *Saglin* or *Lp* in contrast to other previously
reported host loci such as *DSX* or *Nudel* (Kyrou et al., 2018; Hammond et al., 2021).

Tracking one mosquito population of *Lp::Sc2A10* combined with *SagGD^{zpg}* (initial allele
frequencies: 25% and 33%, respectively) over 7 generations showed an increase in the
frequency of both alleles (G7: GFP allelic frequency=59.2%, phenotypic expression of
DsRed in >90% of larvae, n=4282 larvae), indicating that the modest GD strength of
SagGD^{zpg} was still able to increase the frequency of the *SagGD^{zpg}* allele over time and
counteract the *Lp::Sc2A10* fitness cost.

Driving Lp::Sc2A10 *distantly from* Saglin *with a* vasa-Cas9 *gene drive*

In order to achieve a stronger GD that may push *Lp::Sc2A10* to fixation, we re-built the
Saglin GD construct, replacing *zpg* with the *vasa* promoter to control *Cas9* expression
(*SagGD^{vasa}*), in spite of this promoter being known for causing maternal deposition of
Cas9 mRNA and/or protein, potentially resulting in undesired zygotic and somatic
mutation at gRNA target loci.

We initially made two versions of *SagGD^{vasa}*: one carrying the complex array of seven
gRNAs described above for *SagGD^{zpg}* (as in Fig. 3A) ; another carrying six tandem U6-
gRNA units (Fig. 3B; sequence provided in Suppl. File 1E). Both yielded similar results
in early generations of the new mosquito lines. We focused all subsequent analyses on
the latter version expressing three gRNAs against *Saglin* and three against *Lp*, because
its fluorescent marker (*3xP3-DsRed*) proved easier to track in neonate larvae than
OpIE2-DsRed marking the tRNA-containing version (the *OpIE2* promoter integrated in
the *Saglin* locus only became active late in larval development).

About 35 surviving G0 injected male mosquitoes (which were homozygous for *Lp::Sc2A10* to prevent any premature damage on *Lp*), outcrossed to WT females, yielded about 20 DsRed positive G1 females and no positive male, again an indication of proper insertion of the GD cassette onto the X chromosome, where *Saglin* is located. These G1 females (heterozygous at both *SagGD^vasa* on the X chromosome and *Lp::Sc2A10* on the third chromosome) developed normally to adulthood and were fertile when crossed to wild-type males. High homing rates at both loci were recorded in G2 by examining the progeny of the individual G1 females backcrossed to WT males, a large majority of progeny inheriting both a *SagGD^vasa* copy (DsRed_{NLS}) and a *Lp::Sc2A10* copy (GFP) (Table 2).

Female #	Total larvae	Negatives	GFP+ only	DsRed+ only	GFP+ DsRed+	GFP inheritance (*Lp* locus)	DsRed inheritance (*Saglin* locus)
1	89	0	0	0	89	100 %	100 %
2	52	0	4	0	48	100 %	84.6 %
3	59	0	0	0	59	100 %	100 %
4	59	0	6	0	53	100 %	79.6 %
5	50	0	5	0	45	100 %	80 %
6	57	0	3	0	54	100 %	89.4 %

Table 2: The *SagGD^vasa* GD shows high homing rates in the G2 generation at both the *Lp* and *Saglin* loci. Six individual *SagGD^vasa* G1 females mated to WT males oviposited in individual tubes and their larval progeny was scored visually for GFP and DsRed fluorescence.

We then sought to examine the homing rates at the two loci at a larger scale, using COPAS flow cytometry to analyse the progeny from [*SagGD^vasa*/Y ; *Lp::Sc2A10*/+] G2 males backcrossed *en masse* to wild-type females, and from [*SagGD^vasa*/+ ; *Lp::Sc2A10*/+] G2 females backcrossed *en masse* to wild-type males. The male cross yielded only 405 larvae, of which 391 were GFP positive (GFP inheritance: 96.5%, corresponding to a homing rate of 93%). *DsRed* on the X chromosome was, as expected, passed on to 50% of the progeny (daughters). The female cross yielded 5197 larvae showing 95.1% *GFP* inheritance and 80.7% *DsRed* inheritance, corresponding to homing rates of 90.2% at the *Lp* locus and 81.4% at the *Saglin* locus.

[*SagGD^vasa*/Y ; *Lp::Sc2A10*/+] G3 males from this female cross were used as G0 founders of Populations 1 and 2 further monitored below (Fig. 5).

Thus, in contrast to *zpg*, the *vasa* promoter was highly active in germ cells when inserted in the *Saglin* locus. The selected gRNAs, or homing at the *Lp* locus, appeared to be more effective than at the *Saglin* locus. Efficient homing lifted doubt on the activity of *Lp* and *Saglin* gRNAs in the GD construct, or on the propensity of the *Saglin* locus to undergo homologous recombination.

To verify that *SagGD^vasa* abolished *Saglin* expression, we dissected salivary glands from control and homozygous DsRed+ mosquito females, and subjected them to western

blotting using anti-Saglin antibodies. As expected, Saglin was undetectable in
SagGD^vasa mosquitoes (Fig. 3C).

Temporal dynamics of the SagGD^vasa *and* Lp::Sc2A10 *transgenes*

We selected DsRed and GFP double-positive [*SagGD^vasa*/Y ; *Lp::Sc2A10*/+] males from
the female backcross analyzed above and crossed them to WT females, to establish two
independent mosquito populations with a starting (G0) frequency of the autosomal
Lp::Sc2A10 transgene of 25%, whereas X-linked *SagGD^vasa* frequency was 33.3%. We
maintained the two populations for >31 generations and, after 10 and 16 generations,
derived six additional populations from the basal ones by outcrossing to WT. To
examine the long-term dynamics of both transgenes, we exploited GFP fluorescence
from *Lp::Sc2A10* and DsRed fluorescence from *SagGD^vasa* to COPAS-analyze
successive generations of all populations, with large sample sizes, over a long period of
time (>2 years). COPAS analyses allowed not only to distinguish negative from
transgenic larvae but also, at least partially, heterozygotes from homozygotes for each
transgene. Visual examination of the successive COPAS diagrams provides an accurate
sense of the temporal dynamics of both transgenes generation after generation, and an
appreciation of sample sizes (see snapshots of COPAS diagrams in Suppl. File 2).
Figure 5 offers a graphical representation of the same data, after extracting approximate
percentages of larvae of each genotype from the COPAS diagrams. In all populations,
mosquitoes lacking *Lp::Sc2A10* disappeared rapidly, and all young populations rapidly
consisted of a majority of homozygous *Lp::Sc2A10* mosquitoes (Fig. 5 and Suppl. File
2). Despite the absence of physical linkage between the two transgenes, DsRed-only
individuals were strikingly absent from all early populations, indicative of the lethality of
the *SagGD^vasa* transgene in the absence of a protected *Lp* allele. Importantly, toxicity of
SagGD^vasa was due to its destructive action on the *Lp* locus rather than to a fitness cost
of its disrupted *Saglin* host locus, since individuals carrying both DsRed and GFP thrived
in all populations, and consistent with the reported absence of obvious fitness cost of
Saglin mutants (Klug et al., 2023). Several populations (notably populations 1, 3, 4, 5, 6,
7) rapidly seemed fixed for *Lp::Sc2A10*. A trend for reversal, with *Lp::Sc2A10*
progressively giving way to GFP negative individuals (presumably functional *Lp* mutants
refractory to homing), was observed from G14 on for population 2, although GFP
positive individuals still represented a large majority of the population (>90% from G1
on), while GFP allele frequency reached 0.98 at G8 before decreasing and oscillating
between 0.71 – 0.92 after G14. For the *Saglin* locus, COPAS diagram interpretation
must take into account the position of this gene on the X chromosome, with male
mosquitoes bearing a single copy and appearing in the same larval cloud as
heterozygous females (dosage compensation of the transgene not being apparent at
this locus). Drive will be slower as it can occur only in females. While the majority of
mosquitoes of all populations rapidly carried at least one copy of *SagGD^vasa*, we
observed that a substantial fraction of DsRed negative individuals (up to 30%), likely
carrying GD refractory *Saglin* mutants, persisted in some populations (Fig. 5 and Suppl.
File 2).

Detection of homing-refractory Saglin *mutants*

As a non-essential gene, *Saglin* is likely to accumulate mutations in evolving mosquito populations due to failed homing events of the GD construct associated with NHEJ repair at the three gRNA target sites. Indeed, DsRed negative mosquitoes (lacking the GD construct in *Saglin*) persisted at a relatively stable frequency of approximately 30 % in populations 2 and 3 (Fig. 5). To characterize *Saglin* mutants, we COPAS-extracted 750 and 150 DsRed-negative larvae from a surplus of larvae from generation 4 of population 1 and generation 3 of population 2. DsRed-negative individuals represented 18 and 31 % of these populations, respectively, with the *Saglin* gene on their X chromosome having been exposed to between one and four rounds of Cas9 activity. We PCR amplified a *Saglin* fragment spanning the 3 gRNA target sites, and subjected the amplicon to high-throughput sequencing. While the majority of sequenced DsRed negative alleles still corresponded to wild-type *Saglin* having thus far escaped GD activity, a wide variety of *Saglin* mutations, different between the two sampled populations, were readily identified (Fig. 6a,b). Interestingly, a majority of these *Saglin* haplotypes were mutated only at the target site of gRNA2, suggesting that this gRNA was the most active of the three and that Cas9, confronted to a pool of three *Saglin* and three *Lp* gRNAs, will not necessarily cleave all target sites in a given germ cell. However, combinations of mutations at gRNA1, gRNA2 and/or gRNA3 target sites were also identified. Several haplotypes were consistent with iterative action of Cas9 in lineages of haplotypes, a specific mutation occurring either singly, or associated to one or two additional mutations at the other gRNA target sites. Contrarily to our expectation, deletions spanning two gRNA target sites were rare in our samples (one instance of such a haplotype was detected, carrying a deletion between gRNA1 and gRNA2). Overall, Cas9 was not highly active at the *Saglin* locus, some target sites being missed at each generation and combinations of mutations accumulating iteratively as germ cells were exposed to Cas9 in successive generations. Therefore, the target locus of a multiplex gRNA GD experiencing failed homing at a given generation resulting in mutation of one target site may still undergo successful homing in later generations, as long as all gRNA targets are not yet mutated. Inexorably however, *Saglin* mutants fully refractory to the GD will form in mosquito populations; we recovered at least one haplotype with mutations in all three target sites after 4 generations (Fig. 6a).

Appearance of a GD-refractory Lp mutant

NHEJ mutants in *Lp* were expected to emerge at much lower frequency than *Saglin* mutants in evolving populations, given that *Lp* is an essential gene under strong selective pressure. While tracking the frequency of transgenes in [*SagGD^vasa*; *Lp::Sc2A10*] mosquitoes, we noted the appearance of a few GFP negative larvae in populations 2 and 3. These amounted to 2.4% of total larvae in generation 14 of population 2. We COPAS-isolated a sample of these larvae and grew them to adulthood, expecting that they might die during development due to unrescued deleterious mutations in *Lp*. However, these GFP negative mosquitoes developed to healthy and fertile adults. Sequencing the *Lp* gene in 12 individuals revealed that all were homozygous for the same array of mutations within two of the three Lp gRNA target

sequences (Fig. 6c). One mutation deleted three nucleotides before the PAM of gRNA2, removing a single valine from the Lp signal peptide (MWVLGGRRLLWSFLVSLVLIQSVSA, missing valine underlined). The other is a 5-base deletion within the target of gRNA3 in the first Lp intron. Viability of the homozygous mutant indicates that this combination of mutations, while rendering Lp refractory to Cas9-mediated gene conversion, preserved Lp function. This suggested that (i) the first of the three gRNA expressed from the gRNA array did not act; and (ii) combinations of mutations preserving the function of target essential genes can emerge despite the cumulative activity of two gRNAs, which in this case was certainly facilitated by the fact that one of them targeted an intron, more permissive to change.

Genetic instability of the gRNA array

The gRNA array in the *SagGD^vasa* construct being a repetitive tandem of six U6-gRNA units, genetic instability may be expected and observable after several generations of mosquito breeding. We examined transgene integrity in 48 individual mosquitoes of the 32nd generation after transgenesis, by genotyping DsRed-positive males, which possess a single gene drive copy on their X chromosome. PCR primers were chosen to span the gRNA array. Resulting amplicons were examined on agarose gels (Fig. 7) and sequenced. Interestingly, two derivative alleles of the gene drive construct dominated the 4 sampled mosquito populations: a longer version that still retained five of the original six gRNAs, and a shorter version that retained only two gRNAs. The original transgenesis vector used for microinjection was resequenced and verified to contain all six gRNA expression units; therefore, deletions occurred post integration in the mosquito genome. gRNA deletions probably began to occur early in the evolution of the transgenic populations, since the same deletion of the first *Lp* gRNA was detected in both populations 1 and 2, separated at the third generation following transgenesis and subsequently bred independently for 29 generations before sampling. The larger deletion of 4 gRNAs was not observed among the 12 sampled males from population 2, whereas it was the most frequent allele in population 1 and its two daughter populations 5 and 7. This larger deletion may have arisen from a further loss of gRNA units in the allele carrying 5 gRNAs, since *Lp* gRNA1 is missing in both alleles, its position occupied by *Lp* gRNA3 (Fig. 7). The early loss of *Lp* gRNA1 likely explains the lack of any mutation at its target site in the homing-refractory *Lp* mutant identified above. Interestingly, both deleted versions in principle retain the capability to promote homing at both *Saglin* and *Lp* target loci, as even the short form carries one active gRNA against each gene. However, since the single remaining *Lp* gRNA is targeting the *Lp* intron, this shorter GD derivative would easily select for novel homing-refractory *Lp* mutants and this derivative gene drive is likely to be much less durable.

*Vector competence of [*SagGD^vasa*; Lp::Sc2A10] mosquitoes*

We expected that the combination of Sc2A10 expression with *Saglin* loss-of-function would enhance the *Plasmodium* transmission-blocking phenotype. To assess this, we

mixed *[SagGD^vasa; Lp::Sc2A10]* mosquito larvae from the 20^th generation of population 2
with an equal number of non-fluorescent wild-types. The resulting cage of adults was
blood-fed on a mouse infected with *Pb-PfCSP*^hsp70-GFP. Seventeen days later, females
carrying visible GFP parasites were selected and split into six cups of ten transgenic,
and four cups of ten WT. All four mice exposed on day 18 to the infected control females
developed parasitemia, whereas all six mice exposed to infected transgenics remained
negative (Suppl. File 2). Six additional mice were exposed to 8-12 bites from
*[*SagGD^vasa; Lp::Sc2A10*]* oocyst-carrying females of the 4^th generation of population 6,
corresponding to the 23^d generation since transgenesis. Only two of these six mice
became infected (Suppl. Table 2). While the small scale of these experiments cannot
strictly confirm whether the *Saglin* knockout enhanced protection against transmission
(especially as in these experiments we discarded less-infected mosquitoes, which are
expected to result from *Saglin* loss-of-function), the absence of infection in 10/12 mice
showed that *[SagGD^vasa; Lp::Sc2A10]* mosquitoes maintained a markedly decreased
ability to transmit *Plasmodium* after >20 generations of existence.

DISCUSSION

We hijacked an essential gene, *Lipophorin* (*Lp*), to force *Anopheles coluzzii* mosquitoes
to co-express an anti-malarial factor. The choice of the *Lp* gene was motivated by its
exceptionally strong expression level (Lp being the most abundant constitutively
expressed protein in mosquito hemolymph) and by its endogenous signal peptide, which
we could exploit to obtain concomitant secretion of Sc2A10 with Lp protein. The
existence of an endogenous proteolytic cleavage site within the Lp protein, that we
duplicated to detach Sc2A10 from Lp, was also advantageous, as well as the natural
intron immediately following the signal peptide-encoding exon 1 that could accept a
visual selection marker for easy tracking of the genetic modification. This approach is
related to the integral gene drive strategy (Hoermann et al., 2021) in that anti-malarial
factors are expressed together with an endogenous gene, but here the modified
endogenous gene does not comprise any component of a gene drive —in the case of
integral gene drives, gRNAs are inserted in a synthetic intron. The genetic modification
does not disrupt expression or function of the *Lp* essential gene. On the contrary, Lp
protein expression is intended to remain as unaffected as possible, but obligatorily
proceeds through translation of the Sc2A10 antimalarial factor inserted between the Lp
signal peptide and the rest of the Lp protein. This design minimizes the risk of
spontaneous transgene loss by mutation, since frameshift or nonsense mutations in the
Sc2A10 coding sequence would prevent *Lp* expression, resulting in their elimination by
natural selection. Furthermore, the absence of repeated motifs in the synthetic sequence
decreases the risk of in-frame deletions. The possibility of a loss of the transgene's anti-
Plasmodium activity should therefore be limited to missense mutations or in-frame
deletions arising in the *Sc2A10* coding sequence, or to *Plasmodium* evolution, notably
under selection pressure imposed by the transgene itself (Marshall et al., 2019).
However, the 2A10 antibody targets a highly repeated epitope on the CSP protein (the
NANP motif), that, in addition to being repetitive, is also conserved across *P. falciparum*
strains. A combination of mutations leading to the complete loss of this epitope in

Plasmodium therefore seems unlikely, and would probably cause a dramatic loss of
parasite fitness as CSP is required in several processes in the sporozoite's journey
between two hosts (Coppi et al., 2011; Balaban et al., 2021). Hijacking Lp and its signal
peptide is an approach that could be widely applicable to other possible factors acting in
the hemolymph to attack various parasite stages, such as ookinetes, which represent a
particularly vulnerable stage of parasite development in the mosquito. To make
mosquitoes that express and combine additional secreted factors against parasites,
Vitellogenin (AGAP004203) could also be engineered in the same manner, taking
advantage of its own signal peptide, extremely high expression level but also transient
induction following a blood meal. Blood meal inducibility would be particularly suitable for
antimalarial factors specifically targeting ookinetes, as they emerge on the hemolymph-
bathed side of the midgut exactly within the short time window when *Vitellogenin* is
expressed. Mosquitoes would not be exposed during their development to a factor
expressed from *Vg*, limiting potential fitness costs. As proposed by Hoermann et al.
(2021) *Carboxipeptidase* (*Cp*) represents an additional attractive host locus for
antimalarial factors, sharing many of the advantages of *Vg*. A single GD expressing
multiplexed gRNAs could stimulate the simultaneous spread of *Lp, Vg and Cp* modified,
anti-parasitic alleles.

In spite of the high Lp protein secretion in mosquito hemolymph, we only detected less
than 1/10 of this molecular amount for Sc2A10 in hemolymph by mass spectrometry.
Apart from technical limitations in detection sensitivity (e.g., biased trypsin digestion of
some peptides) this could be explained by faster removal from the hemolymph and /or
faster degradation of Sc2A10 compared to Lp. Different, more stable anti-*Plasmodium*
peptides that could accumulate to higher levels may show even better efficacy as
transmission-blocking agents compared to Sc2A10. This also highlights the benefit of
choosing a strong promoter such as that of *Lp* to ensure maximal production of
antimalarial effectors in the hemolymph. A single copy of the *Lp::Sc2A10* transgene still
showed a reduction in transmission. Of note, the rodent model of malaria used here (*P.*
berghei expressing CSP from *P. falciparum*) is characterized by very high infection
rates, typically producing dozens of oocysts per midgut, each of which can release
hundreds of sporozoites in mosquito hemolymph. The ability of secreted Sc2A10 to
decrease sporozoite transmission even in this high infection context is highly
encouraging, given the much lower levels of *P. falciparum* (typically 1 to a few oocysts)
in field infections. An even greater anti-*Plasmodium* activity of Sc2A10 may be reached
by fusing it to antimicrobial peptides such as CecropinA, as achieved by Isaacs et al. in
conventional transgenic *An. stephensi* mosquitoes (Isaacs et al., 2011; Isaacs et al.,
2012) and subsequently in the first published GD for *An. stephensi* (Gantz et al., 2015).
In these studies, the Sc2A10::CecropinA fusion was expressed intermittently under the
control of the blood-meal inducible *Vitellogenin* promoter, and combined with a second
scFv targeting *Plasmodium falciparum* ookinete Chitinase 1 also expressed after blood
meals under the control of the *Cp* promoter. While the insertion of *Sc2A10::CecropinA*
within *Lp* coding sequences may further improve the efficiency of our approach, there is
a risk that CecropinA expressed at all developmental stages, and an increased
complexity of the Lp fusion protein, worsen the fitness cost already observed in
Lp::Sc2A10 mosquitoes.

To force the spread of the *Lp::Sc2A10* transgene in mosquito populations, we tested
several gene drive (GD) designs. A same-locus suppression drive was also installed in
the *Lp* gene, creating a loss-of-function allele. The *Lp::Sc2A10* allele was planned to act
as an R1-type, gene drive refractory functional allele (Champer et al., 2018) that would
supplant the suppression gene drive and invade the population. This was unsuccessful,
as *Lp* proved largely haplo-insufficient. In contrast, distant-locus GDs disrupting *Saglin*,
a non-essential but pro-parasitic gene, were efficient at promoting invasion of both the
Saglin, and particularly the *Lipophorin,* modified alleles. Unlike the first approach, this
design allows *Cas9* and gRNA-coding genes to persist indefinitely within the invaded
mosquito population. Although for this reason we initially preferred the same-locus
approach in which *Lp* function disruption by the insertion of CRISPR/Cas9 components
caused their gradual disappearance by natural selection, the added benefit of a GD
installed in *Saglin* is that it should further reduce the transgenic mosquitoes' vectorial
capacity, a combination of two distinct mechanisms to decrease sporozoite transmission
being more robust to combat *Plasmodium* durably. The genetic design established here
can be viewed as a two-component GD in which one component (*Lp::Sc2A10*) has no
drive capacity by itself but (i) rescues the lethality, and (ii) hitchhikes on the drive
capacity of the other (*SagGD^vasa*) ensuring its own hyper-Mendelian spread. The two
transgenic loci are, therefore, reciprocally dependent on each other: *Lp::Sc2A10*
depends on *SagGD* for its long-term persistence and spread in a population, *SagGD*
depends on *Lp::Sc2A10* for its survival. Importantly, we observed that same-locus or
distant-locus constructs carrying guide RNAs that target wild-type *Lp* are inviable, and
rapidly go extinct in the absence of a recoded *Lp* allele such as *Lp::Sc2A10* (or
spontaneous *Lp* mutants immune to GD that may arise in GD populations). Since
SagGD^vasa will destroy mosquitoes not carrying *Lp::Sc2A10,* this system could initiate
population replacement with a large proportion of laboratory-derived genetic
background, rather than population modification by introgression of the transgenes in a
field genetic background. This is tempered by (i) the viability of [*SagGD^vasa*/+ *;*
Lp::Sc2A10/+] dually heterozygous hybrids and (ii) the moderate fitness cost of
Lp::Sc2A10 that can allow it to diffuse in the field in a Mendelian way independently of
SagGD^vasa, leading to an influx of field genetic background in the final population.
Because *Saglin* knockout mutants bear no fitness cost at least in laboratory conditions,
GD-refractory *Saglin* alleles generated by failed homing events are certain to
accumulate over time (all the more so as their lack of toxicity towards the WT *Lp* allele
compared to *Sag*GD confers them a high selective advantage). If the demise of the
Saglin GD occurs sufficiently late, *Lp-Sc2A10* could still reach high levels or even
fixation in a mosquito population. If, however, GD refractory mutations in *Saglin*
accumulate rapidly and no longer support *Lp::Sc2A10* maintenance, the elimination of
both *SagGD^vasa* and *Lp::Sc2A10* will be inexorable. Thus wild-types alleles are likely to
re-establish at both loci once the frequency of active *Saglin* GDs drops, especially under
conditions of strong mosquito immigration from regions adjacent to the initial GD spread
range. Still, even as genetic resistance to the GD develops, the resulting high frequency
of *Saglin* loss-of-function mutant alleles arising from failed homing events, with their
transient selective advantage while *SagGD^vasa* and wild *Lp* alleles co-exist, should
decrease global vector competence. Once a GD installed in the *Saglin* locus is defeated
by the accumulation of GD-refractory mutations, new generations of GD installed at
different loci could be launched to re-ignite the spread of the *Lp-Sc2A10* allele. For

instance, GDs built for the same mosquito species in the frame of other projects could
also incorporate gRNAs re-igniting the spread of *Lp::Sc2A10*, promoting antimalarial
approaches similar to combination therapies. In this respect, transgenic systems
permitting the expression of multiplexes of many gRNAs would be instrumental. Some of
the constructs presented here incorporate multiplex gRNAs combining individual U6-
gRNA cassettes with a U6-gRNA::tRNA quadruple gRNA expression cassette, with the
drawback of displaying many repeated motifs (U6 promoters, multiple units of
tRNAglycine, gRNA invariable region). Genetic instability of these constructs, with loss
of gRNA expression segments, was prominent in our experiments. This is illustrated by
the high prevalence, in our tracked mosquito populations, of a gene drive derivative
allele carrying only two residual U6-gRNA expression units, from the original tandem of
six. Luckily, the identified allele still retains a driving potential for both the *Saglin* and *Lp*
loci, but it would take only one additional recombination event to reduce the array to a
single gRNA. This would result either in a *Saglin*-only non-lethal GD allele, which may
continue to reduce vectorial capacity of the mosquito population by mutating *Saglin*, or in
a non-driving *Saglin* mutant still able to promote some *Lp::Sc2A10* spread (but
particularly able to generate *Lp* R1-type mutants given the intronic target of the single
residual gRNA). We are currently working to improve gRNA multiplex design by
increasing the number and variety of tRNA spacers, so that the 76-bp constant region of
each gRNA would be the only remaining repeated element.

Apart from the stabilization of multiplex gRNA arrays, other paths to improvement of the
system presented here include avoiding gRNAs targeting intronic sequences, and the
use of promoters better restricted to the germline to control Cas9 expression and limit
GD refractory mutation emergence. Interestingly, the *zpg* promoter that in theory
possesses this advantage showed low activity once inserted in the *Lp* and *Saglin* loci, in
contrast to the *DSX* or *Nudel* loci (Hammond et al., 2016; Kyrou et al., 2018). This
suggests that the *zpg* promoter is more sensitive to the local chromatin context than the
vasa2 promoter, and that its usability in GD constructs may be restricted to a subset of
genomic loci. The *nanos* promoter (Gantz et al., 2015; Terradas et al., 2021; but see
Hammond et al., 2021) may represent an interesting alternative.

A simpler design to promote the spread of antimalarial factors such as Sc2A10 in
mosquito populations would be a single-locus GD (for example installed in *Saglin* or
other pro-parasitic genes if the idea of consolidating the anti-*Plasmodium* activity via two
mechanisms is retained). Few loci, however, can provide the benefits offered here by *Lp*
that include a suitable expression pattern, high expression, an endogenous secretion
signal, endogenous proteolytic cleavage, a natural intron ideally positioned to host a
fluorescent marker, and the status of essential gene that a translational fusion exploits to
limit spontaneous transgene loss by mutation. Besides, adding the *Cas9* and gRNA-
coding cassettes to an intron of an essential gene chosen as GD host would add >5 kb
of foreign DNA sequence, thereby significantly increasing the complexity of the
modification and the risk of affecting functionality of the essential gene. For example,
inadvertent intron splice sites in synthetic sequences can preclude proper mRNA
splicing (Hoermann et al., 2022); we also encountered this problem when inserting a
different anti-*Plasmodium* ScFv in *Lp* (Green, 2019).

In conclusion, the *Sag-GD^vasa* transgene presented here constitutes a viable GD only in
the presence of an engineered *Lp* allele, which by itself has no drive capacity but will
hitchhike with *Sag-GD^vasa* to spread at hyper-Mendelian rates. In caged mosquito
populations, both genetic loci initially spread together and decrease mosquito capacity to
transmit *Plasmodium* parasites expressing CSP from *P. falciparum*. Mutations at both
loci, especially in *Saglin* that is under low selection pressure, arise due to failed homing
events, have a selective advantage over the toxic *SagGD^vasa* allele, and over time
should replace the GD, so that the system carries the seeds of its own removal. The
anti-*Plasmodium Lp-Sc2A10* allele will likely slowly disappear due to its fitness cost,
unless its spread is re-ignited by updated distant-locus GDs. Though such a reversibility
could be seen as a safety advantage for modification GDs (and may be enhanced by
intentionally releasing laboratory-selected resistance alleles), future work should focus
on establishing GD constructs in *Saglin* that limit the emergence of GD refractory alleles
to prolong their potential field lives.

Materials and Methods

Plasmid construction

We used Golden Gate Cloning (Engler and Marillonnet, 2014) to assemble all parts of
the *Lp* (AGAP001826) knock-in plasmid shown in Fig. 1B in destination vector
pENTRR4-ATCC-LacZ-GCTT (Addgene# 173668). The full annotated plasmid
sequence is provided in Suppl. File 1B. The recoded beginning of *Lp*, the Sc2A10
coding sequence and intronic 3xP3-GFP-Tub56D marker cassette were ordered as
synthetic DNA gBlocks (IDT DNA, Belgium) flanked by appropriate *Bsa*I restriction sites.
The Lp endogenous Furin cleavage site (RFRR) was replicated along with 3 N-terminal
and 4 C-terminal flanking amino acids between the Sc2A10 and ApoLpII amino acid
sequences. This, and retaining 7 endogenous nucleotides upstream of the intron 5'
splice junction after the *Sc2A10* coding sequence, resulted in the addition of 7 non-
natural amino acids (GIRESAA) to the N-terminus of the ApoLpII moiety. 1520 bp of 5'
and 1027 bp of 3' flanking sequence were cloned on either side of the modified *Lp*
sequence to be knocked-in. Adjacent to the 3' *Lp* homology arm, we cloned three guide
RNA (gRNA) expression modules (U6 promoter — gRNA coding sequence —
terminator) that specify Cas9 cleavage within the first exon and intron of the genomic
(but not recoded) *Lp* sequence. The three gRNA-expressing modules were prepared in
pKSB-sgRNA1—3 (Addgene #173671—173673) as described (Dong et al., 2018) with
gRNAs targeting the motifs: GAATCAGCACTAGCGACACCAGG,
GCGTGAAATATCGTCAGGGATGG and GGAACGATGTGGGTCCTCGGTGG (PAMs
underlined) in *Lp*. One gRNA target site (with PAM) was included on the 3' flanking
homology arm's distal extremity to favor linearization of the donor plasmid after egg
microinjection. This donor plasmid was called *pENTR R4 3xgRNA Lp::Sc2A10*.
Plasmids encoding gene drive components for insertion into the *Lp* and *Saglin*
sequences were assembled by Golden Gate Cloning in the same manner, including
promoter-Cas9-terminator and *3xP3* or *OpIE2-DsRed_{NLS}*-SV40 modules, the sequences
of which are provided in the complete plasmid sequence (Suppl. File 1A-D). The *zpg-*
Cas9 cassette in *SagGD^zpg* was subcloned from p17410 (Kyrou et al., 2018).

Cas9-expressing mosquito strains

Three transgenic mosquito strains with germline *Cas9* expression under control of the
vasa2 promoter (Papathanos et al., 2009) were used as a source of eggs for
microinjection or in test crosses described in the text. They were constructed by
assembling *vasa* and *Cas9* modules by Golden Gate Cloning into plasmids pDSAY,
pDSARN and pDSAP, marked with *3xP3-YFP*, *3xP3-DsRed$_{NLS}$* and puromycin
resistance, respectively. The pDSARN-vas2-eSpCas9 plasmid encodes the eSpCas9
variant with reduced off-target activity (Slaymaker et al., 2016). Each plasmid was
inserted in the X1 *att*P docking site on chromosome 2 as described (Volohonsky et al.,
2015). The DsRed$_{NLS}$-marked, *vas2-eSpCas9* transgene was then introgressed into the
Ngousso background of *A. coluzzii* by 8 successive backcrosses. In the case of the non
fluorescent, puromycin-resistant Cas9 transgene, homozygosity was achieved by first
balancing the transgene by crossing to the YFP *vasa-Cas9* line, with puromycin
selection of the F1 progeny as described (Volohonsky et al., 2015). Counter-selecting
YFP fluorescence in the F2 progeny of the self-crossed puromycin resistant F1 yielded
the non-fluorescent, vas2-Cas9 homozygous line.

Mosquito egg microinjection and recovery of transgenics

Embryo microinjection to obtain the *Lp::Sc2A10* transgene by CRISPR-Cas9 knockin
was performed as for classical transgenesis (Volohonsky et al., 2015) with 400 ng/µl of
donor plasmid DNA and addition of 2 µM of the drug Scr7 (APExBio) to the injected
plasmid mix. eSpCas9-expressing Ngousso mosquitoes were used as a source of eggs
for microinjection (*vas2-eSpCas9, 3xP3-DsRed$_{NLS}$* strain). We injected the knock-in
plasmid into 400 eggs, recovered 140 surviving G0 adults and out-crossed them *en*
masse to wild-type mosquitoes. In the G1 progeny of the injected G0 males, about 100
GFP positive, putative knock-in individuals were recovered. We established independent
families from ten GFP positive G1 females outcrossed to wild-type, and eliminated the
Cas9 transgene (by counter-selecting *3xP3-DsRed$_{NLS}$*) in these families. All 10 founder
females were confirmed by PCR to carry the intended transgene insertion in the target
Lp locus. We pooled two families to yield the *Lp::Sc2A10* line. To avoid confusion
between two *DsRed* markers, injection of the *Lp-GD* suppression GD construct
(DsRed$_{NLS}$-marked) was performed in eggs expressing wild type Cas9 (vasa-Cas9,
3xP3-YFP G3 background strain). Injection of the *SagGD^{zpg}* and *SagGD^{vasa}* constructs
encoding their own source of *Cas9* was performed in embryos lacking any Cas9
transgene, but respectively carrying a heterozygous or homozygous copy of
Lp::Sc2A10. Numbers of injected eggs and recovered transgenics are provided in the
main text.

COPAS analyses

810
A COPAS Select instrument (Union Biometrica) was used as described (Marois et al.,

2012; Bernardini et al., 2014) to quantify fluorescent transgenes in populations of
neonate mosquito larvae or to establish mixed cultures of defined proportions. The use
of unfed neonate larvae in clean water was crucial for precise gating and sorting. For
sorting, flow rate was kept under 20 larvae/second in Pure mode allowing superdrops.
PMTs were set at 500 (GFP), 600-900 (RFP). Delay, Width, signal threshold and
minimum TOF parameters were set on 8, 6, 100 and 150, respectively.
818

Plasmodium berghei Pb-PfCSP[hsp70-GFP] parasite strain

We generated a novel *P. berghei* strain combining the strong GFP fluorescence of
Δp230p-GFP (Manzoni et al., 2014) with the substitution of endogenous *P. berghei* CSP
for CSP from *P. falciparum* from strain *Pb-PfCSP* (Triller et al., 2017), in which GFP
fluorescence was too faint for selecting live infected mosquitoes examined under 488-
nm light. For this, 150 mosquito females were allowed to blood feed on a mouse co-
infected with both parental parasite strains at a ratio of 1 strongly fluorescent *Δp230p-*
GFP for 40 *Pb-PfCSP*. Sexual reproduction of *P. berghei* in the mosquito generates
hybrid haploid sporozoites, some of which inherit both *Δp230p*-GFP and *Pb-PfCSP*. 17
days after their infective blood meal, live female mosquitoes were screened under 488-
nm light to select those displaying visible GFP sporozoites trapped at the base of their
wing veins. 20 positive females were offered a blood meal on a naïve mouse. When
parasitemia reached 0.1%, 3000 strongly GFP positive blood stage parasites were
sorted by flow cytometry and injected intravenously into two naïve mice, only one of
which developed parasitemia 11 days after passage. Its blood was then passaged into a
new mouse. When parasitemia reached 0.4%, 10 new mice were injected with a blood
dilution corresponding to 1 parasite each, although this number was probably
underestimated as all 10 mice developed parasitemia. Of these, four mice tested PCR
positive for *PfCSP* (PCR primers GGCCTTATTCCAGGAATACCAGTGCT /
GGATCAGGATTACCATCCGCTGGTTG) and negative for PbCSP (PCR primers
GAAGAAGTGTACCATTTTAGTTGTAGCGTC / TGGGTCATTTGGGTTTGGTGGTG).
We selected one clone for passage into naive mice, confirmed its PCR negativity for
PbCSP and positivity for PfCSP, this clone was called *Pb-PfCSP*[hsp70-GFP].

Mosquito infections and bite-back experiments

Neonate larvae from the heterozygous stock of *Lp::Sc2A10* mosquitoes were sorted by
COPAS flow cytometry (Marois et al., 2012) to assemble a mix of equal numbers of
homozygous GFP+ larvae (or heterozygous, in some experiments) and negative control
siblings. Maintaining the transgene in a heterozygous population ensured the absence
of genetic divergence that otherwise may change vector competence between control
and transgenic mosquitoes over time. Following COPAS sorting, transgenic and control
larvae were co-cultured, and adults kept in the same cage, to equalize potential
environmental influences (including unequal microbial communities) that could
differentially affect the vector competence of the two genotypes if grown separately.
Mosquitoes were infected together by blood-feeding from the same mouse with a
parasitemia between 2 and 4%. 16-19 days after mosquito infection, they were

anesthetized and *Lp::Sc2A10* females separated from negative control females based on fluorescence of the transgenesis marker in their eyes. In addition, only *Plasmodium*-carrying mosquitoes were retained, by screening for mosquitoes with GFP-positive midgut oocysts and wing vein or salivary gland sporozoites visible through their cuticle. On day 17-20 after infection, naïve mice were individually exposed to groups of 10 *Lp::Sc2A10* or negative control infected mosquitoes. The number of engorged mosquitoes was recorded after infectious feeding, only mice bitten by at least 6 mosquitoes were included in subsequent analyses with the exception of two mice bitten by 5 and 4 highly infected transgenic and control mosquitoes, respectively, as in this experiment the mouse bitten by the 4 control mosquitoes became infected. Mouse parasitemia was monitored by flow cytometry using an Accuri™ C6 SORP flow cytometer between day 4 and 12 after bite-back. Mice reaching >1.5% parasitemia were sacrificed, and mice remaining negative 12 days post infectious feeding were considered uninfected.

Mass spectrometry analysis of hemolymph

Hemolymph from 25-30 cold-anesthetized female mosquitoes was collected directly into 1x Laemmli buffer 48 hours post-blood feeding by clipping the proboscis and gently pressing their abdomen. Samples were precipitated and digested with trypsin, and 1/5 of the digestion product was analyzed on a Q Exactive Plus Mass Spectrometer coupled to an Easy-nanoLC1000 (Thermo). The acquired data was searched against the *Anopheles* UniProt database plus the Sc2A10 protein sequence using Mascot and the total number of spectra corresponding to ApoLpI, ApoLpII, Sc2A10 and a selection of additional hemolymph proteins that served for normalization were counted.

Protein gels and Western blotting

Mosquito salivary glands or carcass were dissected in 20 µl Laemmli buffer, crushed with a pestle and denatured at 65 °C for 5 minutes. Samples were centrifuged for 3 min at 15,800 g, loaded on a Mini-PROTEAN TGX Stain-Free Precast Gel (BioRad) along with 6 µL of PageRuler Plus Prestained Protein Ladder (ThermoFisher) as standard and electrophoresed at 170 V using the Mini Trans-Blot cell system (BioRad). Gels were blotted on PVDF membranes (Trans-Blot Turbo Mini 0.2 µm PVDF Transfer Pack; BioRad) using the mid-range program of a Pierce Fast blotter (Thermo Fisher Scientific). Membranes were blocked for one hour in PBS + 0.1% Tween (PBST) supplemented with 5% fat-free milk powder, incubated overnight at 4°C with primary antibody, washed three times in PBST and incubated one hour at room temperature in secondary antibody conjugated to horseradish peroxidase (HRP). Antibodies were diluted in PBST, 3% milk. Membranes were then washed three times for 10 minutes with PBS. Antibody binding was revealed using the Super signal WestPico Plus kit (ThermoFisher). After a 1-2 minute incubation, images were acquired using the Chemidoc software (Biorad). Before incubation with further primary antibodies to visualize additional proteins, membranes were stripped for 20 to 30 minutes in Restore PLUS Western Blot Stripping Buffer

(ThermoFisher) and washed three times for 10 minutes in PBST followed by a new
blocking incubation. Saglin polyclonal antibody and recombinant Saglin protein sample
were a kind gift from Eric Calvo (NIAID, Rockville, MD). Human antibodies used to
control protein loading of salivary gland samples were obtained from 2 ml venous blood
taken from a human volunteer arm-feeding *Anopheles* cages over several years. Blood
was incubated at room temperature for 1 hr to allow coagulation and centrifuged for 10
min at ~16.000 g. Serum supernatant was aliquoted and stored at -20°C until use. For
Coomassie staining of electrophoresed hemolymph samples, hemolymph was collected
42h post blood feeding by clipping the proboscis of cold-anesthetized mosquitoes and
gently pressing their abdomen, collecting drops of hemolymph into a pipet tip containing
protein sample buffer (1 µl of sample buffer per mosquito). Samples were denatured for
2 minutes at 90°C and hemolymph from 10 mosquitoes was loaded on each lane of a 4 -
20% 1.5mm polyacrylamide gel (Invitrogen). After electrophoresis, gels were rinsed in
water and stained in SimplyBlue safeStain solution (ThermoFisher).

911
912 Amplicon sequencing
913
A 494-bp region of *Saglin* encompassing the 3 gRNA target sites was amplified from
genomic DNA extracted from 150 and 750 COPAS-sorted, DsRed negative neonate
larvae with the Blood and Tissue kit (Qiagen) using PCR primers
<u>ACACTCTTTCCCTACACGACGCTCTTCCGATCT</u>GCAGAAGCAGCTCGACGC and
<u>GACTGGAGTTCAGACGTGTGCTCTTCCGATCT</u>GCAGCTGCCGGAAGTGCT (Illumina
adapters underlined) and Phire DNA polymerase (Thermofisher Scientific), an annealing
temperature of 68°C and 32 amplification cycles. Resulting amplicons were gel-purified
and sent for sequencing using the Genewiz AmpliconEZ service. Returned paired-end
reads (250 nt length each) were assembled to form single contigs using PANDAseq
version 2.11 (Masella et al., 2012); unmerged reads were manually combined pairwise
and added to the assembled contigs. Data was analysed using a Docker Desktop
Personal version 20.10.16 (https://www.docker.com) -based CRISPResso2 version
2.2.8 (Clement et al., 2019) followed by manual curation.

927
928
929 PCR genotyping of the *SagGD^vasa* gRNA array
930
Genomic DNA was extracted from 48 individual pupae taken from populations 1 and 2
G29 and from populations 5 and 7 G14 (corresponding to >32 generations following
initial transgenesis). A forward primer (GAAGGCGCTGCAGAAGCAGCTCG) was
designed in the Saglin 5' region and a reverse primer (GCCCTCCATGCGCACCTTGAA)
in the *DsRed* cassette (Fig. 7). PCR was performed in a total volume of 15µL using
GoTaq (Promega) (94°C for 5 min then 45 cycles at 95°C for 1 min, 67°C for 20 sec,
72°C for 3.5 min) or Phire polymerase (Thermofisher) (98°C for 2 min then 45 cycles at
98°C for 15 sec, 71°C for 10 sec, 72°C for 1.3 min), with identical results. PCR products
were resolved on 1% agarose gels and the major bands at 3.2 or 1.4 kb were excised
from the gel, purified and sent for Sanger sequencing (Eurofins GATC, Germany) using
one PCR primer. Sequence alignments to the theoretical original plasmid used for
knock-in were performed with SnapGene software. The unique, 20-bp gRNA

protospacer sequences in each of the 6 gRNA coding units allowed to pinpoint which of
the gRNAs had been deleted.

945
946
Acknowledgements

We thank the laboratory of Chris Janse for providing the parental *P. berghei-PfCSP*
strain from which the *Pb-PfCSP*hsp70-GFP strain was derived. We thank Andrew
Hammond, Kyrous Kyrou and Andrea Crisanti for the gift of plasmid p17410 to subclone
the *zpg-Cas9* expression cassette, and Eric Calvo for the gift of the recombinant Saglin
protein and polyclonal Saglin antibodies. We thank Lauriane Kuhn, Johana Chicher and
Philippe Hammann of the Strasbourg IBMC Proteomics Platform for the mass
spectrometry sample preparation and analysis. We thank Muriel Philipps and Claudine
Ebel from the Illkirch IGBMC flow cytometry platform for sorting *P. berghei*-infected red
blood cells. We thank Mallory Kastner and Julie Fimeyer for help during the project. This
work was supported by Agence Nationale de la Recherche, through research grant
#ANR-19-CE35-0007-01 GDaMO to E.M, the Laboratoire d'Excellence (LabEx)
ParaFrap #ANR-11-LABX-0024 to S.A.B., equipment grant #ANR-11-EQPX-0022 for
insectarium operation and by funding from CNRS, Inserm, and the University of
Strasbourg. Additional funding was awarded to D.K. by the DFG as a postdoctoral
fellowship (#KL 3251/1-1).

Agreements

Work on mice was evaluated by the CREMEAS Ethics committee and authorized by
Ministère de l'Enseignement Supérieur et de la Recherche (MESRI) under reference
APAFIS #20562-2019050313288887v3. Work with genetically modified mosquitoes was
evaluated by Haut Conseil des Biotechnologies and authorized by MESRI (agréments
d'utilisation d'OGM en milieu confiné #3243 and #3912).

Figure legends

Figure 1. Characterization and modification of the endogenous *Lipophorin* gene.
A: Coomassie staining of electrophoresed hemolymph protein. Mosquitoes were injected
with the indicated double-stranded RNA to silence *Vg* or *Lp* as described (Rono et al.,
2010) and offered a blood meal after 2 days. Hemolymph was collected 42h post blood
feeding. B: scheme (not to scale) depicting the insertion of synthetic sequences into the
natural Lp gene. The Signal Peptide (SP) was recoded to remove gRNA target sites.
Sc2A10 immediately follows the SP and ends with a RFRR furin cleavage site (red star)
for separation from ApoLpII in the secretory pathway. This site is duplicated from the
ApoLpI/II natural furin cleavage site found downstream. Fat body cells of mosquitoes
bearing the *Lp::Sc2A10* transgene are thus expected to constitutively secrete Sc2A10,
ApoLpII and ApoLpI in the hemolymph. C: spontaneous dynamics of *Lp::Sc2A10*

transgene frequency over 17 generations. Between 1200 and 9100 neonate larvae in
generations 1, 2, 3, 4, 7, 10, 11, 12, and 17, and 522 larvae in generation 16, were
analysed by COPAS flow cytometry. The proportions of homozygotes (Hom, brightest
GFP), heterozygotes (Het, intermediate GFP) and GFP negative (Neg) were estimated
for each analyzed generation by gating each larval population in the COPAS software.
Values are provided in Suppl. Table 1. D: western blot with anti-ApoLpII antibody on
hemolymph samples from 10 female mosquitoes. First and last lanes: protein size ladder
(molecular weights indicated), second lane: hemolymph from WT mosquitoes. Lanes c,
d: hemolymph from homozygous *Lp::Sc2A10* mosquitoes. Failed cleavage between
scFv and ApoLpII would result in a 13 kDa molecular weight upshift compared to the
control. Lanes a, b: hemolymph from mosquitoes expressing a distinct *Lp::ScFv* fusion
not further discussed in this work.

Figure 2: infection status of mice bitten by Sc2A10-expressing vs. WT mosquitoes
infected with *Plasmodium*. Mosquito genotype and number of mice exposed to
infectious mosquito bites is indicated below the bar for each condition. A. Comparison of
the percentages of infected mice after exposure to the bites of homozygous transgenic
Lp::Sc2A10 mosquito females or their wild-type siblings, infected with a *Plasmodium*
berghei strain expressing *P. falciparum* CSP. Aggregated data from 11 independent
experiments performed over a period of two years were analyzed using Fisher's exact
test with a two-tailed p-value (p<0.0001). B. Comparison of mouse infection upon
exposure to heterozygous transgenic *Lp::Sc2A10* mosquito females or their wild-type
siblings, aggregated data from 5 independent experiments (p=0.0006). C. Mice exposed
to WT or homozygous *Lp::Sc2A10* mosquitoes infected with either *P. berghei* expressing
P. falciparum CSP or *P. berghei* expressing endogenous *P. berghei* CSP (single
experiment, p=0.0286).

1010

Figure 3: Designs of *Saglin*-based gene drives that also promote *Lp* locus
modification. A. Gene drive cassette comprising *Cas9* (under control of the *zpg*
promoter) and an array of 7 gRNA-coding modules, inserted disruptively in the
endogenous *Saglin* open reading frame on chromosome X along with a *3XP3-DsRed$_{NLS}$*
fluorescence marker used to track the genetic modification. Three gRNAs, each
expressed under control its own U6 promoter, target wild-type Saglin (purple arrow) and
promote homing of the gene drive cassette. Four gRNAs separated by a repeated
glycine tRNA are expressed under control of one U6 promoter, and target wild-type
Lipophorin on chromosome III (symbolized by green arrow) and promote *Lp::Sc2A10*
homing. B. Updated *SagGD^{vasa}* gene drive construct comprising six gRNAs, each under
control of its own U6 promoter, and Cas9 under control of the *vasa* promoter. C.
Western-blot using Saglin antibodies showing the absence of Saglin protein in the
salivary glands of dissected *SagGD^{vasa}* homozygous females. The same membrane was
re-probed with serum from a human volunteer regularly bitten by mosquitoes, providing
a loading control with salivary and carcass protein signals.

1026

Figure 4: *Sag-GD^{zpg}* triggers modest homing at the *Lp* locus, which can be

rescued by *vasa-Cas9*. A. COPAS analysis of the progeny from [*SagGD^zpg*/+;
*Lp::Sc2A*10/+] females crossed to WT. Weak DsRed fluorescence does not allow
accurate separation of DsRed+ and DsRed- larvae. Note that GFP inheritance is only
slightly higher than the 50% expected in the absence of homing. **B**. COPAS analysis of
the progeny from triple transgenic [*SagGD^zpg* /+; *Lp::Sc2A*10/+; *vasa-Cas9*/+] females
crossed to WT. Note the higher GFP homing rate and the existence of GFP
homozygotes revealing homing in early embryos rather than in the germ line, due to
maternal Cas9 deposition. DsRed positive and negative larvae were indistinguishable
with the COPAS settings used for this experiment.
1037
Figure 5: Temporal dynamics of the *Sag-GD^vasa* and *Lp::Sc2A10* transgenes in 8
mosquito populations. Transgenic mosquitoes carrying both transgenes were crossed
to wild-type (populations 1, 2, 5, 6, 7, 8) or mixed with wild-types (populations 3, 4) and
the frequency of each transgene in 1000-4000 neonate larvae of successive generations
was tracked by flow cytometry (COPAS) using the GFP fluorescent marker of
Lp::Sc2A10 and the DsRed fluorescent marker of *Sag-GD^vasa*. Calculated transgene
frequency, taking homozygous and heterozygous larvae into account (continous lines),
and percentage of fluorescent mosquitoes (dotted lines) are shown on the graphs.
Frequency of each genotype in the population was calculated after gating corresponding
clouds of larvae on COPAS diagrams (see Suppl. File 2) and recording the associated
percentage measured by COPAS software, or by opening COPAS files and gating in
WinMDI software. Gating was approximate due to partial overlap between different
clouds of larvae. Supplementary File 2 shows strips of all COPAS diagrams that served
to extract this data, which provide a more accurate sense of transgene dynamics over
generations.
1053
Figure 6: (a, b) Characterization of *Saglin* failed homing mutations. A PCR product
spanning the three *Saglin* gRNA target sites was amplified from 150 DsRed-negative
larvae from generation 4 of population 1 (a) or from 750 DsRed-negative larvae from
generation 3 of population 2 (b) and subjected to high-throughput amplicon sequencing.
The different *Saglin* mutant haplotypes discovered in each sample are aligned to the WT
sequence (top) with target site and protospacer-adjacent motifs (PAMs) indicated.
Arrows point to Cas9 cleavage sites. Deleted nucleotides are highlighted in green,
inserted nucleotides in blue. // indicates portions of WT sequence not represented on
the figure. The distance between two gRNA target sites (number of nucleotides) is
indicated above the WT sequence. **(c) gRNA target sites and failed homing-induced**
mutations in the *Lp* gene. The WT sequence of the 5' region of the *Lp* gene shows
exon and intron sequences, the position of gRNA target sites with PAMs (red), and the
ATG initiator codon (underlined). Nucleotides shown in orange were deleted in twelve
sequenced homozygous mutant mosquitoes. An additional, 11-bp deletion in the intron,
distant from gRNA target sites (red) was unexpected.
1069
Figure 7: PCR genotyping of 31^st generation individual *Sag-GD^vasa* mosquitoes
reveals deletions in the gRNA array. PCR amplicons spanning the six initial gRNAs in
Sag-GD^vasa were generated from 12 individual male mosquitoes (carrying a single
transgene copy) from each of populations 1, 2, 5 and 7, 31 generations after initial
transgene integration (corresponding to the indicated generation number for each

tracked population) and resolved on 1% agarose gels with 1 kb+ DNA ladder
(Thermofisher) as a size marker (upper panels). Arrows point to PCR products specific
of the transgene as confirmed by Sanger sequencing; other bands are PCR artefacts.
Sanger sequencing of amplicons A and B showed that they contain only five and two
residual gRNA coding units, respectively, as schematized in the bottom panel. PCR was
replicated on the same samples with different polymerases (*Taq* and Phire) with
identical results.
1082
Suppl. File 2: Temporal dynamics of the *Sag-GD^vasa* and *Lp::Sc2A10* transgenes in
8 mosquito populations. Each strip of panels on successive pages corresponds to an
independent caged mosquito population. Each panel is a COPAS analysis snapshot of
neonate larvae at the indicated generation, obtained as shown on the first page.
Successive panels provide a global view of each transgene's evolution on samples of
1000-4000 neonate larvae at each generation, with fluorescence intensity correlating
with transgene copy number (GFP marks the *Lp::Sc2A10* transgene on the Y axis,
DsRed marks *Sag-GD^vasa* on the X axis). Up to 9 partially overlapping distinct larval
populations corresponding to the 9 possible genotypes (0, 1, or 2 copies of each
fluorescence, sketched on page 1 and on selected panels duplicated below the original
panel) can be resolved on COPAS diagrams. Dots indicated in population 1 panels
correspond to debris (dead larvae, egg shells) and are difficult to distinguish from non-
fluorescent larvae unbless sorting them for verification under the micrscope. Yellow
arrow in population 2, G14 indicates the appearance of GFP negative larvae that
represent a *Lp* homing refractory mutant (see text). Tracking Populations 1 and 2 started
shortly after initial transgenesis, by crossing G2 [*Sag-GD^vasa* / Y ; *Lp::Sc2A10 / +*] males
to WT females (considered the G0 cross for temporal tracking of genotypes). Population
3 was assembled by mixing 20 transgenics from generation 12 of population 2 with 190
WT. Population 4 was assembled by mixing 60 transgenics from generation 12 of
population 2 with 189 WT. In these new G0 mixes reproductive success of transgenics
proved higher than that of the WT (due to differences in the quality of the parental
mosquitoes), as G1 pupae consisted of 189 GFP transgenics for 358 negatives = 34.5%
instead of expected 10% (population 3) or 346 GFP transgenics for 304 negatives =
53.2% instead of expected 24.1% (population 4). In all populations, note the initial rapid
convergence of genotypes towards *Lp::Sc2A10* (GFP) homozygosity, the rapid initial
disappearance of *Sag-GD^vasa* (DsRed) positives lacking GFP, and the persistence of a
fraction of DsRed negative larvae.
1110
Supplementary Figure 1: Insertion of a *Lipophorin* suppression gene drive and its
expected and unexpected outcomes. A: scheme of the suppression drive construct.
Upon chromosome cleavage by Cas9 (shown by scissors), homologous recombination
(gray shading) via 5' and 3' regions of homology (HR) is expected to insert the
intervening elements (gRNA-encoding array, zpg-Cas9 expression cassette, DsRed$_{NLS}$
marker cassette) into the beginning of the Lp gene, disrupting its function. Note that this
recombination event knocking out the essential *Lp* gene was designed to occur in the
presence of an intact *Lp::Sc2A10* sister chromosome, not shown in the scheme but that
contributed to the unexpected outcome shown in B. B: most knock-in events
unexpectedly arose from compound recombinations between the cleaved WT

chromosome and both the repair template plasmid and the intact *Lp::Sc2A10* sister
chromosome, resulting in insertion of the gene drive elements genetically linked to the
Lp::Sc2A10 transgene. A hypothetical scheme interpreting the compound insertion is
shown. It was inferred from the genetic linkage observed between GFP and DsRed$_{NLS}$,
and from sequencing a PCR product obtained with primers AG105 and EM890
(indicated). ∆ indicates a 500-bp deletion in the cloned Lp 3'region, which arose
following micro-injection of the plasmid because one of the gRNA target sites was
retained by error in this part of the construct (indicated by scissors). Presence of the
gRNA-tRNA array (green boxes + yellow arrows) was confirmed by drive activity in
presence of a *vasa-Cas9* transgene.

Supplementary Figure 2: Characterization of the two types of Lp-GD integration
events. A: The GFP-RFP line showed high homing rates when complemented with an
independent vasa-Cas9 transgene. COPAS diagrams show the fluorescence of progeny
neonate larvae from heterozygous GFP-RFP mosquitoes (third chromosome) that also
carried a non-fluorescent, puromycin resistance-marked vasa-Cas9 transgene (second
chromosome, heterozygous), crossed to non-fluorescent partners (WT* indicates non-
fluorescent partners actually carrying one copy of the puromycin resistant vasa-Cas9
transgene, not influencing the crossing outcome). N indicates the total number of larvae
analyzed in the diagrams; percentages of larvae of each fluorescence are indicated.
Non-fluorescent progeny would amount to 50% in the absence of homing. Note that the
percentage of GFP-RFP is close to 50%, mainly reflecting Mendelian inheritance of the
parental transgene. Thus most homing events involved only a GFP-containing segment
from the parental GFP-RFP transgene. Rarer instances of RFP-only segment homing
(arrows) were accompanied by a decrease of DsRed fluorescence intensity, indicating
that the *3xP3* promoter is less active than in the GFP-RFP context. **B**: Haploinsufficiency
at the *Lipophorin* locus. Photograph shows larval and pupal progeny from heterozygous
Lp-GD crossed to heterozygous *Lp::Sc2A10.* Right panel focuses on a subset of larvae
observed under fluorescent light. Larvae that inherited one copy of the *Lp-GD* loss-of-
function allele (red eyes) and one copy of *Lp::Sc2A10* (green eyes/brain) are
developmentally delayed and will die before adulthood. Individuals that inherited a WT
Lp copy and either modified allele will complete development, with strong loss of fitness
for WT/*Lp-GD* individuals (red-eyed pupa).

1153
Supplementary Table 2: mouse parasitemia after infection by mosquito bites.
Mice are grouped by categories according to the genotype of the *Plasmodium*-carrying
mosquitoes biting them or by the strain of parasite infecting them. Groups of 10 infected
mosquito females (transgenic or their wild-type siblings grown in the same culture), were
allowed to bite a mouse. Colored cells indicate the stage at which infected mice were
sacrificed to prevent the appearance of malaria symptoms. n.d= not determined. 'D' in
the third column indicates a mouse bitten by transgenic mosquitoes that showed a
clearly delayed parasitemia compared to the controls. In two cases (indicated in third
column), the mouse was bitten by fewer than 6 WT mosquitoes.

1163
Bibliography

Balaban, A.E., Kanatani, S., Mitra, J., Gregory, J., Vartak, N., Sinnis-Bourozikas, A.,
Frischknecht, F., Ha, T., Sinnis, P., 2021. The repeat region of the
circumsporozoite protein is an elastic linear spring with a functional role in
Plasmodium sporozoite motility. https://doi.org/10.1101/2021.05.12.443759
Bernardini, F., Galizi, R., Menichelli, M., Papathanos, P.-A., Dritsou, V., Marois, E.,
Crisanti, A., Windbichler, N., 2014. Site-specific genetic engineering of the
Anopheles gambiae Y chromosome. Proc. Natl. Acad. Sci. U. S. A. 111, 7600–
7605. https://doi.org/10.1073/pnas.1404996111
Carballar-Lejarazú, R., James, A.A., 2017. Population modification of Anopheline
species to control malaria transmission. Pathog. Glob. Health 111, 424–435.
https://doi.org/10.1080/20477724.2018.1427192
Carballar-Lejarazú, R., Ogaugwu, C., Tushar, T., Kelsey, A., Pham, T.B., Murphy, J.,
Schmidt, H., Lee, Y., Lanzaro, G.C., James, A.A., 2020. Next-generation gene
drive for population modification of the malaria vector mosquito, Anopheles
gambiae. Proc. Natl. Acad. Sci. U. S. A. 117, 22805–22814.
https://doi.org/10.1073/pnas.2010214117
Carballar-Lejarazú, R., Tushar, T., Pham, T.B., James, A.A., 2022. Cas9-mediated
maternal effect and derived resistance alleles in a gene-drive strain of the African
malaria vector mosquito, Anopheles gambiae. Genetics iyac055.
https://doi.org/10.1093/genetics/iyac055
Champer, J., Kim, I.K., Champer, S.E., Clark, A.G., Messer, P.W., 2021. Suppression
gene drive in continuous space can result in unstable persistence of both drive
and wild-type alleles. Mol. Ecol. 30, 1086–1101.
https://doi.org/10.1111/mec.15788
Champer, J., Liu, J., Oh, S.Y., Reeves, R., Luthra, A., Oakes, N., Clark, A.G., Messer,
P.W., 2018a. Reducing resistance allele formation in CRISPR gene drive. Proc.
Natl. Acad. Sci. U. S. A. 115, 5522–5527.
https://doi.org/10.1073/pnas.1720354115
Champer, J., Liu, J., Oh, S.Y., Reeves, R., Luthra, A., Oakes, N., Clark, A.G., Messer,
P.W., 2018b. Reducing resistance allele formation in CRISPR gene drive. Proc.
Natl. Acad. Sci. 115, 5522–5527. https://doi.org/10.1073/pnas.1720354115
Clement, K., Rees, H., Canver, M.C., Gehrke, J.M., Farouni, R., Hsu, J.Y., Cole, M.A.,
Liu, D.R., Joung, J.K., Bauer, D.E., Pinello, L., 2019. CRISPResso2 provides
accurate and rapid genome editing sequence analysis. Nat. Biotechnol. 37, 224–
226. https://doi.org/10.1038/s41587-019-0032-3
Coppi, A., Natarajan, R., Pradel, G., Bennett, B.L., James, E.R., Roggero, M.A.,
Corradin, G., Persson, C., Tewari, R., Sinnis, P., 2011. The malaria
circumsporozoite protein has two functional domains, each with distinct roles as
sporozoites journey from mosquito to mammalian host. J. Exp. Med. 208, 341–
356. https://doi.org/10.1084/jem.20101488
Dong, Y., Simões, M.L., Marois, E., Dimopoulos, G., 2018. CRISPR/Cas9 -mediated
gene knockout of Anopheles gambiae FREP1 suppresses malaria parasite
infection. PLoS Pathog. 14, e1006898.
https://doi.org/10.1371/journal.ppat.1006898
Engler, C., Marillonnet, S., 2014. Golden Gate cloning. Methods Mol. Biol. Clifton NJ
1116, 119–131. https://doi.org/10.1007/978-1-62703-764-8_9

Gantz, V.M., Akbari, O., 2018. Gene editing technologies and applications for insects. Curr. Opin. Insect Sci. 28. https://doi.org/10.1016/j.cois.2018.05.006

Gantz, V.M., Jasinskiene, N., Tatarenkova, O., Fazekas, A., Macias, V.M., Bier, E., James, A.A., 2015. Highly efficient Cas9-mediated gene drive for population modification of the malaria vector mosquito Anopheles stephensi. Proc. Natl. Acad. Sci. U. S. A. 112, E6736-6743. https://doi.org/10.1073/pnas.1521077112

Ghosh, A.K., Devenport, M., Jethwaney, D., Kalume, D.E., Pandey, A., Anderson, V.E., Sultan, A.A., Kumar, N., Jacobs-Lorena, M., 2009. Malaria parasite invasion of the mosquito salivary gland requires interaction between the Plasmodium TRAP and the Anopheles saglin proteins. PLoS Pathog. 5, e1000265. https://doi.org/10.1371/journal.ppat.1000265

Green, E., 2019. Development of genetic control strategies for insect pests using CRISPR/Cas9 (phdthesis). Université de Strasbourg.

Hammond, A., Galizi, R., Kyrou, K., Simoni, A., Siniscalchi, C., Katsanos, D., Gribble, M., Baker, D., Marois, E., Russell, S., Burt, A., Windbichler, N., Crisanti, A., Nolan, T., 2016. A CRISPR-Cas9 gene drive system targeting female reproduction in the malaria mosquito vector Anopheles gambiae. Nat. Biotechnol. 34, 78–83. https://doi.org/10.1038/nbt.3439

Hammond, A., Karlsson, X., Morianou, I., Kyrou, K., Beaghton, A., Gribble, M., Kranjc, N., Galizi, R., Burt, A., Crisanti, A., Nolan, T., 2021. Regulating the expression of gene drives is key to increasing their invasive potential and the mitigation of resistance. PLoS Genet. 17, e1009321. https://doi.org/10.1371/journal.pgen.1009321

Hammond, A.M., Kyrou, K., Bruttini, M., North, A., Galizi, R., Karlsson, X., Kranjc, N., Carpi, F.M., D'Aurizio, R., Crisanti, A., Nolan, T., 2017. The creation and selection of mutations resistant to a gene drive over multiple generations in the malaria mosquito. PLoS Genet. 13, e1007039. https://doi.org/10.1371/journal.pgen.1007039

Hoermann, A., Habtewold, T., Selvaraj, P., Corsano, G.D., Capriotti, P., Inghilterra, M.G., Kebede, T.M., Christophides, G.K., Windbichler, N., 2022. Gene Drive Mosquitoes Can Aid Malaria Elimination by Retarding Plasmodium Sporogonic Development. https://doi.org/10.1101/2022.02.15.480588

Hoermann, A., Tapanelli, S., Capriotti, P., Del Corsano, G., Masters, E.K., Habtewold, T., Christophides, G.K., Windbichler, N., 2021. Converting endogenous genes of the malaria mosquito into simple non-autonomous gene drives for population replacement. eLife 10, e58791. https://doi.org/10.7554/eLife.58791

Hollingdale, M.R., Nardin, E.H., Tharavanij, S., Schwartz, A.L., Nussenzweig, R.S., 1984. Inhibition of entry of Plasmodium falciparum and P. vivax sporozoites into cultured cells; an in vitro assay of protective antibodies. J. Immunol. 132, 909–913.

Horn, C., Wimmer, E.A., 2000. A versatile vector set for animal transgenesis. Dev. Genes Evol. 210, 630–637. https://doi.org/10.1007/s004270000110

Isaacs, A.T., Jasinskiene, N., Tretiakov, M., Thiery, I., Zettor, A., Bourgouin, C., James, A.A., 2012. Transgenic Anopheles stephensi coexpressing single-chain antibodies resist Plasmodium falciparum development. Proc. Natl. Acad. Sci. U. S. A. 109, E1922-1930. https://doi.org/10.1073/pnas.1207738109

Isaacs, A.T., Li, F., Jasinskiene, N., Chen, X., Nirmala, X., Marinotti, O., Vinetz, J.M., James, A.A., 2011. Engineered resistance to Plasmodium falciparum development in transgenic Anopheles stephensi. PLoS Pathog. 7, e1002017. https://doi.org/10.1371/journal.ppat.1002017

Klug, D., Gautier, A., Calvo, E., Marois, E., Blandin, S.A., 2023. The salivary protein Saglin facilitates efficient midgut colonization of Anopheles mosquitoes by malaria parasites. PLoS Pathog. 19, e1010538. https://doi.org/10.1371/journal.ppat.1010538

Kyrou, K., Hammond, A.M., Galizi, R., Kranjc, N., Burt, A., Beaghton, A.K., Nolan, T., Crisanti, A., 2018. A CRISPR-Cas9 gene drive targeting doublesex causes complete population suppression in caged Anopheles gambiae mosquitoes. Nat. Biotechnol. 36, 1062–1066. https://doi.org/10.1038/nbt.4245

Manzoni, G., Briquet, S., Risco-Castillo, V., Gaultier, C., Topçu, S., Ivănescu, M.L., Franetich, J.-F., Hoareau-Coudert, B., Mazier, D., Silvie, O., 2014. A rapid and robust selection procedure for generating drug-selectable marker-free recombinant malaria parasites. Sci. Rep. 4, 4760. https://doi.org/10.1038/srep04760

Marois, E., Scali, C., Soichot, J., Kappler, C., Levashina, E.A., Catteruccia, F., 2012. High-throughput sorting of mosquito larvae for laboratory studies and for future vector control interventions. Malar. J. 11, 302. https://doi.org/10.1186/1475-2875-11-302

Marshall, J.M., Raban, R.R., Kandul, N.P., Edula, J.R., León, T.M., Akbari, O.S., 2019. Winning the Tug-of-War Between Effector Gene Design and Pathogen Evolution in Vector Population Replacement Strategies. Front. Genet. 10, 1072. https://doi.org/10.3389/fgene.2019.01072

Masella, A.P., Bartram, A.K., Truszkowski, J.M., Brown, D.G., Neufeld, J.D., 2012. PANDAseq: paired-end assembler for illumina sequences. BMC Bioinformatics 13, 31. https://doi.org/10.1186/1471-2105-13-31

O'Brochta, D.A., Alford, R., Harrell, R., Aluvihare, C., Eappen, A.G., Li, T., Chakravarty, S., Sim, B.K.L., Hoffman, S.L., Billingsley, P.F., 2019. Is Saglin a mosquito salivary gland receptor for Plasmodium falciparum? Malar. J. 18, 2. https://doi.org/10.1186/s12936-018-2634-5

Okulate, M.A., Kalume, D.E., Reddy, R., Kristiansen, T., Bhattacharyya, M., Chaerkady, R., Pandey, A., Kumar, N., 2007. Identification and molecular characterization of a novel protein Saglin as a target of monoclonal antibodies affecting salivary gland infectivity of Plasmodium sporozoites. Insect Mol. Biol. 16, 711–722. https://doi.org/10.1111/j.1365-2583.2007.00765.x

Papathanos, P.A., Windbichler, N., Menichelli, M., Burt, A., Crisanti, A., 2009. The vasa regulatory region mediates germline expression and maternal transmission of proteins in the malaria mosquito Anopheles gambiae: a versatile tool for genetic control strategies. BMC Mol. Biol. 10, 65. https://doi.org/10.1186/1471-2199-10-65

Pham, T.B., Phong, C.H., Bennett, J.B., Hwang, K., Jasinskiene, N., Parker, K., Stillinger, D., Marshall, J.M., Carballar-Lejarazú, R., James, A.A., 2019. Experimental population modification of the malaria vector mosquito, Anopheles stephensi. PLoS Genet. 15, e1008440. https://doi.org/10.1371/journal.pgen.1008440

Port, F., Bullock, S.L., 2016. Augmenting CRISPR applications in Drosophila with tRNA-flanked sgRNAs. Nat. Methods 13, 852–854. https://doi.org/10.1038/nmeth.3972

Quinn, C.M., Nolan, T., 2020. Nuclease-based gene drives, an innovative tool for insect vector control: advantages and challenges of the technology. Curr. Opin. Insect Sci. 39, 77–83. https://doi.org/10.1016/j.cois.2020.03.007

Reuling, I.J., Mendes, A.M., de Jong, G.M., Fabra-García, A., Nunes-Cabaço, H., van Gemert, G.-J., Graumans, W., Coffeng, L.E., de Vlas, S.J., Yang, A.S.P., Lee, C., Wu, Y., Birkett, A.J., Ockenhouse, C.F., Koelewijn, R., van Hellemond, J.J., van Genderen, P.J.J., Sauerwein, R.W., Prudêncio, M., 2020. An open-label phase 1/2a trial of a genetically modified rodent malaria parasite for immunization against Plasmodium falciparum malaria. Sci. Transl. Med. 12, eaay2578. https://doi.org/10.1126/scitranslmed.aay2578

Rono, M.K., Whitten, M.M.A., Oulad-Abdelghani, M., Levashina, E.A., Marois, E., 2010. The major yolk protein vitellogenin interferes with the anti-plasmodium response in the malaria mosquito Anopheles gambiae. PLoS Biol. 8, e1000434. https://doi.org/10.1371/journal.pbio.1000434

Slaymaker, I.M., Gao, L., Zetsche, B., Scott, D.A., Yan, W.X., Zhang, F., 2016. Rationally engineered Cas9 nucleases with improved specificity. Science 351, 84–88. https://doi.org/10.1126/science.aad5227

Sumitani, M., Kasashima, K., Yamamoto, D.S., Yagi, K., Yuda, M., Matsuoka, H., Yoshida, S., 2013. Reduction of malaria transmission by transgenic mosquitoes expressing an antisporozoite antibody in their salivary glands. Insect Mol. Biol. 22, 41–51. https://doi.org/10.1111/j.1365-2583.2012.01168.x

Terradas, G., Hermann, A., James, A.A., McGinnis, W., Bier, E., 2021. High-resolution in situ analysis of Cas9 germline transcript distributions in gene-drive Anopheles mosquitoes. G3 Bethesda Md jkab369. https://doi.org/10.1093/g3journal/jkab369

Triller, G., Scally, S.W., Costa, G., Pissarev, M., Kreschel, C., Bosch, A., Marois, E., Sack, B.K., Murugan, R., Salman, A.M., Janse, C.J., Khan, S.M., Kappe, S.H.I., Adegnika, A.A., Mordmüller, B., Levashina, E.A., Julien, J.-P., Wardemann, H., 2017. Natural Parasite Exposure Induces Protective Human Anti-Malarial Antibodies. Immunity 47, 1197-1209.e10. https://doi.org/10.1016/j.immuni.2017.11.007

Volohonsky, G., Terenzi, O., Soichot, J., Naujoks, D.A., Nolan, T., Windbichler, N., Kapps, D., Smidler, A.L., Vittu, A., Costa, G., Steinert, S., Levashina, E.A., Blandin, S.A., Marois, E., 2015. Tools for Anopheles gambiae Transgenesis. G3 Bethesda Md 5, 1151–1163. https://doi.org/10.1534/g3.115.016808

WHO, 2021. World Malaria report 2021.

Xie, K., Minkenberg, B., Yang, Y., 2015. Boosting CRISPR/Cas9 multiplex editing capability with the endogenous tRNA-processing system. Proc. Natl. Acad. Sci. U. S. A. 112, 3570–3575. https://doi.org/10.1073/pnas.1420294112

Zavala, F., Cochrane, A.H., Nardin, E.H., Nussenzweig, R.S., Nussenzweig, V., 1983. Circumsporozoite proteins of malaria parasites contain a single immunodominant region with two or more identical epitopes. J. Exp. Med. 157, 1947–1957. https://doi.org/10.1084/jem.157.6.1947

Figure 1

Figure 2

Figure 3

Figure 4

A   Progeny from:
♀ SagGD^{zpg} /+; Lp::Sc2A10/+
x ♂ WT

DsRed fluorescence

B   Progeny from:
♀ SagGD^{zpg} /+; Lp::Sc2A10/+; vasa-Cas9/+
x ♂ WT

DsRed fluorescence

Figure 5

Figure 6

Figure 7