5   Partial protection from fluctuating selection leads to evolution toward amplified fluctuation

6   in population size and a novel mechanism of balancing selection

8   Yuseob Kim

9   Division of EcoScience and Department of Life Science, Ewha Womans University

10   Ewhayeodae-gil 52, Seodaemun-gu, Seoul, Korea 03760

11   Email: yuseob@ewha.ac.kr

13   **Keywords:** fluctuating selection, balancing selection, storage effect, absolute fitness, population size

2022-12-04

Abstract

Cyclic or random fluctuation of population density is prevalent in nature and is not generally considered

adaptive but primarily the result of fluctuation in environmental conditions. Classical theory predicted that

natural selection favors a variant whose fitness fluctuation is smaller, thus causing reduced fluctuation of

population density. It is also known that if population is partially protected from fluctuating selection, for

example in the presence of seed bank, the variance of fitness is further reduced and therefore the reproductive

success of population is ensured. This study, exploring a mathematical model for coupled demographic and

evolutionary dynamics based on absolute fitness, finds that such a 'refuge' from fluctuating selection can even

allows positive selection for a variant increasing the amplitude of population size fluctuation under weak or

moderate regulation of population density. Under strong density regulation and constant carrying capacity, the

model recreates the condition for long-term maintenance of polymorphism, which is known as the storage effect.

However, if the carrying capacity is changing either cyclically or randomly and the density regulation is strong,

variants whose fitness fluctuates in the same direction as the population size are positively selected to either

fixation or oscillation at intermediate frequencies. The latter dynamics, arising when fitness fluctuates as

expected under a simple life-history trade-off, is a novel form of balancing selection that may explain the recent

discovery of allele frequency oscillation at many polymorphic sites in North American *Drosophila* populations.

Introduction

Natural populations face cyclic or random fluctuations of their abiotic and biotic environment that lead to fluctuation in individuals' fitness (GRANT AND GRANT 2002; BELL 2010; THOMPSON 2013). Evolution under fluctuating selection has been a major topic in population genetics. Theoretical models have explored the effect of fluctuating selection on various aspects of evolution such as the level of both neutral and non-neutral genetic variation (DEMPSTER 1955; HALDANE AND JAYAKAR 1963; HEDRICK 1976; HEDRICK 1986; GILLESPIE 1994; BARTON 2000; HUERTA-SANCHEZ *et al.* 2008). Recently, balancing selection under seasonal fluctuation of fitness has received particular attention as a potential explanation for the apparent seasonal oscillation of allele frequencies at thousands of nucleotide sites in temperate *Drosophila melanogaster* populations (BERGLAND et al. 2014; WITTMANN et al. 2017; MACHADO et al. 2021). One area of theoretical development focuses on the effect of a subset of the population or a particular life cycle stage protected from fluctuating selection in increasing species or allelic diversity (CHESSON AND WARNER 1981; ELLNER AND HAIRSTON 1994; GULISIJA AND KIM 2015; SVARDAL et al. 2015; BERTRAM AND MASEL 2019), which is commonly called the storage effect. Simple mathematical models showed that negative frequency-dependent selection emerges in the presence of a refuge from selection, such as a seed bank (TURELLI et al. 2001). The storage effect can generate stable long-term polymorphism at multiple loci (PARK AND KIM 2019), although it requires strong selection (a large magnitude of fitness oscillation; BERTRAM AND MASEL (2019)).

Partial protection from fluctuating selection is expected not only with seed banks but also in other conditions commonly encountered by plants and animals. The survival of individuals in diapause, for example in the insect pupal stage, is probably unaffected by a variable environment (HEDRICK 1995). In species with overlapping generations, only a portion of a population will be subject to selection on a trait that is expressed at a particular stage of the life-cycle (CHESSON AND WARNER 1981; ELLNER AND HAIRSTON 1994). In dioecious species, alleles affecting female reproductive traits, for example, are protected from selection if they are carried by males (REINHOLD 2000). If an allele conferring genetic robustness or phenotypic plasticity is segregating in a population, it constitutes a genetic background on which new mutants are protected from selection (GULISIJA et al. 2016). In all of these cases, a species effectively consists of a subpopulation exposed to selection and another subpopulation, the refuge, protected from selection leading to the storage effect. Individuals (i.e., alleles) move between these subpopulations with a certain probability per unit time.

Temporally variable environments most likely cause a fluctuation not only in the relative fitness among

alleles but also their absolute fitness, the per-capita rate of reproduction per unit time. Namely, the evolutionary

change of population under fluctuating environment must occur while population size is changing concurrently.

If population density regulation, which is defined as the force that pushes the population size toward a carrying

capacity set by environment conditions at a given time, is not very strong, population size can also change (e.g.

overshoot or undershoot the carrying capacity) as its genetic composition changes. However, theoretical models

for the storage effect have considered an infinitely large or constant-sized population, under which the

evolutionary dynamics are fully described by the relative fitness and the relative frequencies of alleles. In this

study, the storage effect is modeled with absolute fitness, which allows joint changes in demography and allele

frequencies. Then, it can be investigated whether and how evolutionary changes interact with fluctuations in

population size. First, it remains to be examined whether the maintenance of polymorphism by the storage effect

still occurs under variable population sizes due to relaxed density regulation. Second, one may ask whether the

nature of population size fluctuation is altered by evolutionary process. The proximate cause of population size

fluctuation, observed in a wide range of species in nature, is probably the fluctuation of abiotic and biotic factors

in the environment. However, the degree of such demographic fluctuation depends on the phenotypes of

individuals that are the product of evolution.

Classical studies in population genetics found that the long-term reproductive success of a population

or an allele under fluctuating environment is given by the geometric mean of its fitness over time. The geometric

mean decreases as fitness fluctuates in time even if its arithmetic mean remains constant. Therefore, it is

predicted that a genotype causing less fluctuation in absolute fitness, for example by allocating resource for

survival during a harsh period, is favored under fluctuating environment (GILLESPIE 1977; LANDE et al. 2009). It

should be noted that seed bank and other means of partial protection from fluctuating selection were proposed

as a means of increasing the geometric mean fitness of a population by reducing the variance of absolute fitness,

a mechanism known as the evolutionary bet-hedging (SEGER AND BROCKMANN 1987; PHILIPPI AND SEGER 1989).

Assuming that there are two alleles whose absolute fitness fluctuate differently but yield the same

arithmetic or geometric mean fitness outside the refuge, I found that an allele with a wider fluctuation in

absolute fitness can be positively selected under weak or moderate regulation of population density. Namely,

directional, not negative frequency-dependent, selection occurs and results in the amplification of population

size fluctuation. More surprisingly, preexisting oscillation in population size was found to generate positive

selection for an allele whose relative fitness oscillates in phase with population size oscillation, leading to its

fixation or long-term oscillation at intermediate frequencies. The latter is a novel mechanism of balancing

selection that may explain the seasonal oscillation of allele frequencies at multiple loci in BERGLAND *et al.* (2014).

The main results will be given in two steps. The full model will be first explored using numerical

iterations, which reveals various patterns of eco-evolutionary dynamics. Then, theoretical explanations for these

results will be sought by mathematical analyses on reduced models.

Modeling the fitness of a variant under fluctuating selection

Consider a population of haploid individuals reproducing in discrete generations in an environment that

cyclically (e.g., seasonally) changes with a period of λ generations. In the following, analyses will focus on the

fate of a focal allele, A_2, at a locus that arose recently by mutation so that its frequency, q, is low initially. The

ancestral (wild-type) allele of the locus is A_1 with frequency $p = 1 - q$. A trade-off is assumed in the effect of A_2

on fitness over different phases of an environmental cycle. For example, due to the allocation of finite resource

over time, if A_2 increases fitness in one season, it inevitably leads to a decline in fitness in the other season. Here,

the fitness of an A_2–carrying individual can be defined relative to that of an A_1-carrying individual or as the

expected number of offspring per capita (the absolute fitness). In the (implicit) assumption of strictly constant-

sized population, one only needs to specify the relative fitness of alleles in the model. However, in order to model

the joint dynamics of evolutionary and demographic changes, absolute fitness needs to be specified. Here the

model of constant (or infinite) sized population will be first considered.

First, the fitness of A_2 relative to A_1 may be given by $w_2 = e^{s_t}$ ($w_1 = 1$), where

$$s_t = s_{max} \sin \left[\frac{2\pi t}{\lambda} \right]. \tag{1}$$

This ensures that the geometric mean fitness of A_2 is equal to that of A_1, because the relative frequencies of A_1

and A_2 after one cycle of fluctuating environment is given by p' and q', where

$$\frac{q'}{p'} = \left(\prod_{t=1}^{\lambda} \frac{w_2}{w_1} \right) \frac{q}{p} = \exp\left[\sum_{t=1}^{\lambda} s_t \right] \frac{q}{p} = \frac{q}{p}.$$

Therefore, allele frequencies are not expected to change in this population. In this case, A_1 and A_2 may be called

quasi-neutral to each other. There are other ways to specify the fitness of A_1 and A_2 so that their geometric mean

fitness are identical. For example, GULISIJA AND KIM (2015) assumed $w_1 = 1 - s_t$ and $w_2 = 1 + s_t$ and BERTRAM

AND MASEL (2019) also used an analogous model where the ratio of two alleles' fitness is simply reversed upon

a seasonal change. This class of fitness assignments may be called geometric mean-preserving fitness-

fluctuation, or GMF in short. In the following a mutant allele having this fitness effect will be called a GMF allele.

Second, one may assume $w_1 = 1$ and $w_2 = 1 + s_t$, making the arithmetic mean fitness identical. Here,

fitness ratio is not reversed in the opposite phase of a cycle (e.g., $1 + s < 1/(1 - s)$). In this case, it can be easily

shown that $q'/p' < q/p$ after one environmental cycle with $s_{max} > 0$. Namely, the derived allele A_2 will be

eliminated from the population since its geometric mean fitness is lower than that of A_1. This fitness assignment

is called arithmetic mean-preserving fitness-fluctuation or AMF. A mutant allele with this fitness effect will be

called an AMF allele.

Which fitness model, GMF or AMF, should be assumed is not a trivial matter as the outcomes of

following models critically depend on it. Which one is more realistic and thus should arise more often needs to

be addressed as well. Under the strict density regulation in a constant-sized population, w_2/w_1 is

approximately the absolute fitness of A_2 when it is rare. Then, AMF means that a fitness gain, i.e. an excess in the

number of offspring, in one phase is just enough to compensate the loss in another phase of the environmental

cycle. This may happen naturally if a finite resource limits the total number of offspring born in one

environmental cycle. In case of GMF, the arithmetic mean of w_2/w_1 over the cycle is greater than 1 (e.g.,

$(e^s + e^{-s})/2 > 1$ for $s > 0$). Thus, modeling with a GMF allele means that the carrier of a rare derived allele is

assumed to produce more offspring, averaged over a cycle, than that of the wild-type allele. One may argue that

such a positive effect on fitness may not be achieved by simply changing resource allocation over time. However,

there can be many other ways in which fitness fluctuation arises (see more in Discussion). Why the frequency

of A_2 does not increase with GMF, despite an increased production of offspring, was well explained by GILLESPIE

(1977) and others who showed that the outcome of fluctuating selection is determined by the geometric mean

fitness, which decreases as the variance of fitness fluctuation increases: A_2 at low frequency has higher mean

but larger variance in fitness compared to A_1.

Storage effect under the fluctuation of relative fitness

Next, I will review the theory of the storage effect (CHESSON AND WARNER 1981; GULISIJA AND KIM 2015; BERTRAM

AND MASEL 2019). A GMF allele, A_2, is initially assumed in the fluctuating environment: $w_2 = e^{s_t}$ and $w_1 = 1$.

The population is now divided into two with fractions 1-*r* and *r*. Only the first subpopulation resides in an

environment subject to the fluctuating selection, which I will call the "field". In the second subpopulation,

individuals have identical fitness regardless of alleles, and thus are protected from fluctuating selection. While

this "refuge" from selection primarily models a seed bank, it may also correspond to a certain stage of a life cycle

in species with overlapping generations, a spatially separated population with relaxed selection, or one sex that

is not the target of sex-limited selection. Many of such scenarios are true for diploid species, but the current

haploid model should be applicable to diploids with incomplete dominance. It is also assumed that individuals

are redistributed into the field and the refuge, with probability 1-*r* and *r*, respectively, during the migration step

of each generation. Let q_t be the frequency of A_2 at generation *t*. It is again assumed that A_2 recently arose by

mutation and therefore q_t is close to 0. After one cycle of fluctuating selection, frequency $q_{t+\lambda}$ is expected to be

larger than q_t because

$$q_{t+\lambda} = q_t \prod_{i=1}^{\lambda} \left((1-r)\frac{w_2}{\overline{w}} + r \right)$$

$$\approx q_t \prod_{i=1}^{\lambda} \left((1-r)\frac{w_2}{w_1} + r \right) = q_t \exp\left[\sum_{i=1}^{\lambda} \log[(1-r)\frac{w_2}{w_1} + r] \right]$$

$$\geq q_t \exp\left[\sum_{i=1}^{\lambda} \left((1-r)\log\frac{w_2}{w_1} + r\log 1 \right) \right] = q_t \left(\prod_{i=1}^{\lambda} \frac{w_2}{w_1} \right)^{1-r} = q_t. \tag{2}$$

The mean fitness in the field $\overline{w} = pw_1 + qw_2$ is approximated by w_1 with $q \ll p$ and Jensen's inequality was

applied above. No change in allele frequency ($q_{t+\lambda} = q_t$) occurs if fluctuating selection is turned off ($s_{max} = 0$).

Therefore, the presence of a refuge confers an advantage to a rare allele under fluctuating selection, thus

generating the storage effect.

Furthermore, the above analysis suggests that, to generate the storage effect, the change of

environment in the field (fluctuation of w_2 versus w_1) does not have to be periodic as long as the product of $\frac{w_2}{w_1}$

over successive generations converges to 1 with GMF. To confirm this, the iteration of allele frequencies

according to the model above was made using $s_t = s_{max}\sin[2\pi t^*/\lambda]$, where t^* is a random integer between 1

and λ drawn each generation. Interestingly, the rare allele increased in frequency slightly faster on average

under this mode of random fluctuating selection than under the periodic ("unscrambled") fluctuation (Fig. 1).

Figure 1. Frequency (q_t) of the A_2, a GMF allele, over time t (in generations) under periodic (gray curve; using $s_t = s_{max}\sin[2\pi t/\lambda]$) or random (dark curve; using $s_t = s_{max}\sin[2\pi t^*/\lambda]$, averaged over 300 replicates) fluctuation of fitness in a population subdivided into the field and the refuge, under the relative fitness-based model. Other parameters: $r = 0.5$, $\lambda = 10$, $q_0 = 0.01$, $w_1 = 1$, $w_2 = e^{st}$, and $s_{max} = 0.3$.

The storage effect is intriguing because a rare allele is positively selected simply by moving between the field and refuge randomly, even though it is not favored in either subpopulation. It is even possible that a slightly deleterious allele (i.e., having a smaller geometric mean fitness than the other allele in the field) is positively selected when it is rare (GULISIJA AND KIM 2015). A plain explanation for this phenomenon was that the loss of an allele from the field during an unfavorable period is buffered by its presence in the refuge; then, during the next period when the same allele is now favored in the field, the refuge supplies this allele to the field. However, the buffering effect of a refuge should apply to the other (common) allele as well.

A better explanation may be given by the theory of evolutionary bet-hedging (SEGER AND BROCKMANN 1987; PHILIPPI AND SEGER 1989), according to which a refuge such as a seedbank reduces the variance of reproductive success (of the total population) in a variable environment, thus increasing the geometric mean fitness. This effect of refuge should be larger for the carriers of the rare allele; as mentioned earlier, with GMF, the arithmetic mean of A_2's fitness is higher than that of A_1 but its fluctuation larger than that of A_1 prevents it from attaining higher geometric mean fitness than A_1. Eq. (2) suggests that, if the variance of fitness in the field is V, that of the total population is reduced to $(1 - r)^2 V$ in the presence of refuge. Although the mean fitness of A_2 also decreases due to refuge (e.g., fitness advantage over A_1 decreases by a factor r), this effect must be smaller compared to the decrease of variance; the net effect is to elevate the geometric mean fitness of A_2 over 1. On the other hand, refuge has little effect on the geometric mean fitness of A_1, the major allele, because its absolute

fitness fluctuates negligibly. This explains how negative frequency-dependent selection, the storage effect, arises

due to partial protection from selection if GMF is assumed.

The storage effect does not arise for an AMF allele ($w_1 = 1$ and $w_2 = 1 + s_t$). Again, a refuge reduces

the variance of A_2's fitness and thus increases its geometric mean fitness when A_1 is rare. However, the geometric

mean fitness of A_2 relative to A_1 will approach only up to its arithmetic mean fitness relative to A_1, which cannot

exceed 1 under the assumption of AMF. Therefore, the proposal of storage effect as a mechanism of balancing

selection in the previous studies critically depended on assuming a GMF, not AMF, allele in modeling fluctuating

selection.

Cyclic fluctuation of absolute fitness: Constant carrying capacity

The above model based on relative fitness and relative allele frequencies does not assume any particular

population dynamics. However, a fluctuating environment is likely to generate fluctuation in population size as

well if population density is not strictly regulated. Furthermore, the dynamics of allele frequency changes may

depend on whether population size varies in time or not. I therefore expanded the above model to allow the

sizes of subpopulations to change according to the absolute fitness of individuals under varying degrees of

population density regulation.

Let n_{1f} and n_{2f} (n_{1r} and n_{2r}) be the numbers of A_1 and A_2 individuals in the field (in the refuge).

In the absence of density regulation, n_{1f} and n_{2f} are assumed to increase to $W_{1f}n_{1f}$ and $W_{2f}n_{2f}$ over a

generation in the field. Namely, W_{1f} and W_{2f} are the absolute fitness of A_1 and A_2, which may change over

time. Absolute fitness for both alleles in the refuge is given by W_r. For density regulation, I define the carrying

capacities of the field and refuge to be K_f and K_r, which may also change over time. In addition, it is assumed

that, after steps of growth and density regulation, individuals in the field migrate to the refuge with probability

m_{fr} and those in the refuge migrate to the field with probability m_{rf}. Then, the numbers of A_1 and A_2

individuals in the field and refuge in the next generation become

$$n'_{1f} = (1 - m_{fr})W_{1f}n_{1f}R[W_{1f}n_{1f} + W_{2f}n_{2f}, g_f, K_f] + m_{rf}W_r n_{1r}R[W_r(n_{1r} + n_{2r}), g_r, K_r], \qquad (3a)$$

$$n'_{2f} = (1 - m_{fr})W_{2f}n_{2f}R[W_{1f}n_{1f} + W_{2f}n_{2f}, g_f, K_f] + m_{rf}W_r n_{2r}R[W_r(n_{1r} + n_{2r}), g_r, K_r], \qquad (3b)$$

$\qquad n'_{1r} = m_{fr}W_{1f}n_{1f}R[W_{1f}n_{1f} + W_{2f}n_{2f}, g_f, K_f] + (1 - m_{rf})W_r n_{1r}R[W_r(n_{1r} + n_{2r}), g_r, K_r],$ \hfill (3c)

and

$\qquad n'_{2r} = m_{fr}W_{2f}n_{2f}R[W_{1f}n_{1f} + W_{2f}n_{2f}, g_f, K_f] + (1 - m_{rf})W_r n_{2r}R[W_r(n_{1r} + n_{2r}), g_r, K_r],$ \hfill (3d)

where

$\qquad R[N, g, K] = \dfrac{1+g}{1+g\frac{N}{K}}$ \hfill (3e)

determines the strength of density regulation. If individuals carry only neutral alleles and the population size N

is much smaller than its carrying capacity K, the population will grow at rate g (≥ 0). Therefore, a large value of

g means strong density regulation and the relative fitness-based model above is restored with $g \to \infty$. Different

strengths of regulation, g_f and g_r, are applied to the field and the refuge.

$\qquad$ How a population evolves under this absolute fitness-based model was explored by the recursion of

Eq. (3). Constant carrying capacity for the field ($K_f = 1000$) and immediate redistribution over subpopulations

($m_{fr} = m_{rf} = 0.5$) were initially assumed. Four values of g_f that represent the increasing strength of density

regulation in the field were used (0.02, 0.2, 2, and 20). For the refuge, $g_r = 0$ is assumed (as the model of seed

bank is primarily considered), which makes it meaningless to specify K_r.

$\qquad$ The dynamics of a GMF allele was first explored. The fitness of A_1 and A_2 was given by $W_{1f} = e^{-0.5s_t}$,

$W_{2f} = e^{0.5s_t}$, and $W_r = 1$, where s_t is given by Eq. (1). Namely, the oscillation in absolute fitness of the two alleles

is symmetric in the field. Under this fitness scheme n_1, which initially oscillated around $2K_f$ (= 2,000),

decreased towards an oscillation around K_f (Fig. 2A). On the other hand, n_2 increased from a small to large

number that oscillates around K_f. This approach of both alleles toward the relative frequency of 0.5 was faster

with increasing g. (In Fig. 2A, the ranges of oscillation in n_1 and n_2 may appear to change little over time with g

= 0.02 but they do converge close to 0.5 if iterations are done for a sufficiently large number of generations.)

This observation is consistent with the storage effect arising in the relative fitness-based model, which is

equivalent to this absolute fitness-based model with $g \to \infty$.

$\qquad$ Next, the fitness scheme is changed to $W_{1f} = 1$ and $W_{2f} = e^{s_t}$, which keeps the relative fitness

identical to the above scenario but lets the absolute fitness of A_2 fluctuate more than that of A_1. This dramatically

changed the course of evolution (Fig. 2B). With strong density regulation ($g = 20$), the relative frequency of A_2

approached 0.5, as before. However, under weaker density regulation, the absolute count of A_2 continued to

increase toward $2K_f$, while that of A_1 decreased toward zero. Similar results are obtained when the absolute

fitness of both alleles fluctuate but the fluctuation of A_2 is larger (e.g., $W_{1f} = e^{0.5s_t}$ and $W_{2f} = e^{1.5s_t}$). Namely,

an allele increases in frequency by directional, not negative frequency-dependent, selection if its absolute fitness

fluctuates more than that of the other allele. Switching the fitness oscillation of A_1 and A_2 ($W_{1f} = e^{-s_t}$ and

$W_{2f} = 1$) confirmed this conclusion (Fig. 2C). Consequently, unless density regulation is strong, the population

evolves toward experiencing a greater fluctuation in size.

Figure 2. Changes in the absolute frequencies of A_1 ($n_1 = n_{1f} + n_{1r}$; gray curves) and A_2 alleles ($n_2 = n_{2f} + n_{2r}$; dark curves) in a population subdivided into the field and the refuge ($m_{fr} = m_{rf} = 0.5$), according to deterministic recursion of Eq. (3) starting from $n_{1f} = n_{1r} = 990$ and $n_{2f} = n_{2r} = 10$. Note that, as these graphs are unable to resolve curves oscillating very rapidly on the scale of time (in generations) shown above, they appear in the shape of bands. The field maintains a constant carrying capacity ($K_f = 1,000$) with varying strengths (g_f) of density regulation. Other parameters: $\lambda = 10$, $W_r = 1$, $g_r = 0$, $s_{max} = 0.3$.

This result – directional selection on an allele with larger fluctuation in absolute fitness – is again the

consequence of assuming a GMF allele and the presence of refuge. Briefly, in the field, the arithmetic mean of

$W_{2f} = e^{st}$ becomes increasingly larger than 1 as the oscillation of s_t becomes larger in magnitude, while its

geometric mean remains at 1. However, refuge reduces variance in the fitness of A_2 in the total population,

elevating its geometric mean (absolute) fitness above 1.

According to a simple mathematical analysis of this model at the limit of no density regulation ($g_f =$

0), given in Appendix A, if the proportion of refuge is r in the total population and haploids are redistributed

between the field and the refuge each generation ($m_{fr} = r$ and $m_{rf} = 1 - r$), the increase in the number of an

allele whose absolute fitness in the field fluctuates according to $W_f = e^{s_t}$ is given by

$$n_T \approx n_0 \exp\left[\frac{1}{4}{s_{max}}^2 r(1-r)T\right] \tag{5}$$

where n_0 is the initial number and T is a multiple of λ. This approximates well the initial increase of A_2 under

weak density regulation given by exact recursion (Fig. S1). Eq. (5) shows that the geometric mean fitness of a

GMF allele is maximized when the size of refuge is equal to that of the field ($r = 0.5$), and that, as the rate of

increase is in the order of ${s_{max}}^2$, a significant change is expected only if the amplitude of fitness fluctuation is

large (BERTRAM AND MASEL 2019). It is also shown in Appendix A that this increase does not occur for mutants

with AMF.

Figure S1. Comparison between the changes of n_1 according to exact recursion (Eq. [3]; dark curves) and approximation by $n_0 \exp\left[\frac{{s_{max}}^2}{16}t\right]$ (red curves) in populations with $m_{fr} = m_{rf} = 0.5$ under weak density regulation ($g_f = 0$ or 0.02). Fitness is given by GMF ($W_{1f} = \exp(s_t)$ and $W_{2f} = 1$), with $s_{max} = 0.3$. The changes of n_2 by Eq. (3) are shown in gray curves.

The results shown in Fig. 2 can now be understood in terms of the "intrinsic growth rate" of a GMF

allele under no density regulation described by Eq. (5). With positive (albeit weak) density regulation ($g_f > 0$),

the number of A_2 cannot increase indefinitely. However, A_2 outcompetes A_1, whose absolute fitness fluctuates

less ($W_{1f} = 1$ in Fig. 2B). Namely, the two alleles change in number according to their intrinsic growth rates

before density regulation is applied, and their ratio determines the change in their relative frequencies. Under

strong density regulation ($g_f \gg 1$), the relative frequency of A_2 approaches a stable fluctuation around 0.5, at

which n_1 and n_2 must fluctuate symmetrically so that their intrinsic rates of growth suggested by Eq. (5) become

equal.

Cyclic fluctuation of absolute fitness: oscillating carrying capacity

Next, to investigate whether the pre-existing fluctuation in population size affects the evolutionary dynamics of

an allele with fluctuating fitness, the carrying capacity of the field was set to change cyclically, again with period

λ: $K_f = K_{f0}\exp\left[\beta\sin\left[\frac{2\pi t}{\lambda}\right]\right]$, assuming $\beta > 0$ without the loss of generality. This models many species'

population cycles, which are commonly found to be the symmetric oscillations of density in the logarithmic scale

(MYERS 1988). Again, a GMF allele A_2 whose fitness fluctuation is in or out of phase of the population size

fluctuation arises at $t = 0$. The fitness scheme of $W_{1f} = e^{-0.5st}$ and $W_{2f} = e^{0.5st}$, for example, means that the

A_2 (A_1) allele acts to amplify (dampen) the oscillation of population size beyond the level specified by the

ecological factor β under weak density regulation. The recursion of Eq. (3) under this parameter set revealed

that the A_2 allele is positively selected to reach fixation, surprisingly, even under strong density regulation (g_f

$= 20$) (Fig. 3A). Similar rates of increase in n_2 over n_1 with $g_f > 1$ were observed with other fitness schemes:

$\left(W_{1f}, W_{2f}\right) = (1, e^{st})$ or $(e^{-st}, 1)$ (Fig. 3B and 3C). On the other hand, if A_2's fitness fluctuation is out of phase

with population size fluctuation ($W_{1f} = 1$ and $W_{2f} = e^{-st}$; Fig. 3D) positive selection for this allele under

strong density regulation did not happen. Therefore, under strong density regulation, an allele is positively

selected if its fluctuation in relative fitness is in phase with population size (carrying capacity) fluctuation.

Further exploration revealed that the fixation of A_2 allele however requires strong the fluctuation of K_f for a

given magnitude of fitness fluctuation, namely $\beta > s_{\max}$ (Fig. S2). With relatively weaker fluctuation of

population size, the long-term maintenance of polymorphism, the storage effect, was observed.

With weak density regulation ($g_f = 0.02$ and 0.2), the allele with a larger amplitude in absolute fitness

fluctuation outcompetes the other, as in the case of constant carrying capacity (Fig. 2B and 3B), probably

because the effect of (oscillating) carrying capacity on evolutionary dynamics diminishes as density regulation

weakens. It should be noted again that, unless density regulation is very strong, the fixation of A_2 leads to the

increased magnitude of population size fluctuation compared to that when the population was fixed for A_1 (Fig.

3).

Figure 3. Changes in the absolute frequencies (n_1 and n_2) of A_1 (gray curves) and A_2 (dark curves) alleles under the cyclic fluctuation of carrying capacity in the field ($K_{f0} = 1,000$ and β = 0.5). Four fitness schemes that are classified as GMF [$(W_{1f}, W_{2f}) = (e^{-0.5s_t}, e^{0.5s_t})$ (A), $(1, e^{s_t})$ (B), $(e^{-s_t}, 1)$ (C), and $(1, e^{-s_t})$ (D)] were used. Other parameters are identical to those in Fig. 2.

Figure S2. The relative frequency, q, of A_2 allele $(W_{1f} = 1, W_{2f} = e^{s_t})$ at the 10,000th generation of iterating eq. (3) with $g_f = 20$. For s_{max} = 0.1, 0.3, and 0.5, increasing values of β, from 0 to 0.5 with increment of 0.05 were used. Other parameter values are identical to those in Fig. 3.

Assuming an AMF allele also resulted in diverse evolutionary outcomes with oscillating carrying

capacity. As confirmed earlier, A_2 failed to increase in frequency when $W_{1f} = 1$ and $W_{2f} = 1 + s_t$ with a

constant carrying capacity of the field (Fig. 4A), as its geometric mean fitness in the field alone is less than 1.

However, under cyclic oscillation of K_f ($\beta = 0.5$) and strong density regulation ($g_f = 2$ or 20), the frequency of

A_2 increased by positive selection. With $s_{max} = 0.3$, A_2 did not reach fixation but remained polymorphic with A_1

in the long run (Fig. 4B), suggesting that a form of balancing selection arose. With smaller s_{max} (= 0.2), however,

A_2 finally reached fixation (Fig. 4C). Again, these results depend on the condition that the fitness oscillation of

A_2 is in phase with population size oscillation in the field. If fitness oscillation was out of phase ($W_{1f} = 1$ and

$W_{2f} = 1 - s_t$), positive selection on A_2 did not occur (Fig. 4D).

Figure 4. Changes in the absolute frequencies (n_1 and n_2) of A_1 (gray curves) and A_2 (dark curves) alleles when the fitness scheme of A_2 is classified as AMF: $(W_{1f}, W_{2f}) = (1, 1 + s_t)$ (A, B, C), and $(1, 1 - s_t)$ (D). Carrying capacity in the field is either constant ($\beta = 0$; A) or oscillating ($\beta = 0.5$; B, C, D). The magnitude of fitness fluctuation, s_{max}, is either 0.3 (A, B, D) or 0.2 (C). Other parameters are identical to those in Fig. 3. When much longer iteration (> 20,000) was done for the parameter set shown in panel B with g_f = 20, n_1 and n_2 were observed to be oscillating at values that have changed little from those observed at generation 2,000.

Two-season model for fitness and population size fluctuation

For the general understanding of positive selection generated under the oscillation of carrying capacity, revealed

above by numerical iterations of eq. (3) for a limited set of parameters, a simpler model at the limit of strict

density regulation is analyzed. A cyclic environment with $\lambda = 2$ is assumed, in which each period is made up of

a "good" season (the first generation) and a "bad" season (the second generation). The carrying capacity of the

field then alternates between K_G and K_B, where $K_G \geq K_B$. The absolute fitness of A_1 and A_2 individuals in the

field are assumed to be W_{1G} and W_{2G} in a good season and W_{1B} and W_{2B} in a bad season. In the refuge, $W_r = 1$.

Strict density regulation in the field but no density regulation in the refuge ($g_f \to \infty$ and $g_r = 0$) is assumed.

Then, assuming $m_{fr} = r$ and $m_{rf} = 1 - r$, this population reaches a demographic equilibrium in which the

total population size alternates between N_G (at the end of a good season) and N_B (at the end of a bad season).

In Appendix B, it is shown that the number of haploid individuals carrying A_2 allele changes from n_2 to n''_2 in

one cycle, where

$$\frac{n''_2}{n_2} = \left((1-r)\frac{K_G}{(1-r)N_B}\frac{W_{2G}}{W_{1G}} + r\right)\left((1-r)\frac{K_B}{(1-r)N_G}\frac{W_{2B}}{W_{1B}} + r\right) \tag{6}$$

approximately, if n_2 is far smaller than n_1. Eq. (6) shows that, as expected under strict density regulation, the

evolutionary change of n_2 is dependent on the relative, not absolute, fitness of alleles. However, because the

absolute, not relative, frequency of n_2 is tracked in the derivation, the growth rates of population size in the field

should be combined with the relative fitness to determine its evolutionary change; the field grows from $(1-$

$r)N_B$ to K_G during the good season and shrinks from $(1-r)N_G$ to K_B during the bad season.

Now, A_2 is assumed to be a GMF allele. The relative fitness between two alleles is reversed each season:

$\frac{W_{2G}}{W_{1G}} = \frac{W_{1B}}{W_{2B}} = \omega$. Then, the change in the number of A_2 when it is rare is given by

$$f(\omega) \equiv \frac{n''_2}{n_2} = \left(\frac{K_G}{N_B}\omega + r\right)\left(\frac{K_B}{N_G}\frac{1}{\omega} + r\right).$$

330 It can be easily shown that $f(1) = 1$, as expected for a neutral allele. Solving $f(\omega) = 1$ and analyzing the

331 derivative of $f(\omega)$ shows that

$$f(\omega) \geq 1 \text{ if } r \geq 0 \text{ and } \omega \leq \frac{1+rk}{k^2+rk} \text{ or } \omega \geq 1,$$

333 where $k = \frac{K_G}{K_B}$, and $f(\omega) < 1$ otherwise (Appendix B). Therefore, positive selection on A_2 allele requires the

presence of refuge. If $K_G = K_B$ (no change in population size), $f(\omega) > 1$ for all $\omega \neq 1$. Since $f(\omega^{-1})$ describes the change of A_1 frequency when A_1 is rare, this result means that both A_1 and A_2 alleles are positively selected when they are rare. This symmetry creates negative frequency-dependent selection—the storage effect. However, if $K_G > K_B$ and $1 < \omega < \frac{k^2+rk}{1+rk}$, A_2 is positively selected but A_1 is not, even if A_1 is rare. This asymmetry means that A_2, whose fitness fluctuates in the same direction as (i.e., in phase with) population size fluctuation, is under directional selection and will eventually reach fixation. For example, if $K_G = 2K_B$, an allele that is between 1 and 2.5 times advantageous over the other allele during a good season but inversely disadvantageous during a bad season is positively selected until it reaches fixation. Lastly, if $\omega > \frac{k^2+rk}{1+rk}$, both A_1 and A_2 alleles are positively selected when they are rare, thus yielding the storage effect again: fitness fluctuation is much stronger than population size fluctuation so that the qualitative behavior of system is identical to that with no population size fluctuation.

In case of an AMF allele, we may assume $\frac{W_{2G}}{W_{1G}} = 1 + s$ and $\frac{W_{2B}}{W_{1B}} = 1 - s$. The change in the number of A_2 when it is rare is given by

$$h_2(s) \equiv \frac{n''_2}{n_2} = \left(\frac{K_G}{N_B}(1+s) + r\right)\left(\frac{K_B}{N_G}(1-s) + r\right).$$

Solving $h_2(s) \geq 1$ gives $0 \leq s \leq \frac{r}{1-r^2}\left(\frac{K_G}{K_B} - \frac{K_B}{K_G}\right)$. Therefore, unless s is too large relative to the magnitude of population size fluctuation (K_G/K_B), A_2 allele whose fitness fluctuation is in phase with population size fluctuation invades the population. At the same time, the condition for the A_1 allele invading a population fixed with A_2 is given by

$$h_1(s) \equiv \frac{n''_1}{n_1} = \left(\frac{K_G}{N_B}\frac{1}{1+s} + r\right)\left(\frac{K_B}{N_G}\frac{1}{1-s} + r\right).$$

For $0 \leq s \leq \frac{r(k+1)(k-1)}{(k+r)(rk+1)}$, where $k = \frac{K_G}{K_B}$, $h_1(s) \leq 1$. Note that $\frac{r(k+1)(k-1)}{(k+r)(rk+1)}$ is an increasing function of k (> 1). Therefore, with s that is small for a given magnitude of population size fluctuation, the condition for A_2 increasing in frequency to fixation is satisfied. However, with $s > \frac{r(k+1)(k-1)}{(k+r)(rk+1)}$, $h_1(s) > 1$: A_1 can invade the population. Therefore, with an intermediate strength of fluctuating selection $(\frac{r(k+1)(k-1)}{(k+r)(rk+1)} < s < \frac{r}{1-r^2}\left(k - \frac{1}{k}\right))$, polymorphism with A_1 and A_2 should be maintained. Parameter ranges leading to different evolutionary

358   outcomes under the assumption of AMF are graphically summarized in Fig. 5.

Figure 5. Domains of s and K_G/K_B in which negative selection ($h_2(s)$ <1), balancing selection ($h_1(s)$ >1, $h_2(s)$ >1), and directional selection ($h_1(s)$ <1, $h_2(s)$ >1) on the derived allele A_2 occurs in the two-season model (r = 0.5) of AMF and population size fluctuation.

Random fluctuation of absolute fitness

Finally, it was examined whether evolutionary dynamics produced in the absolute fitness-based model also

arise when the absolute fitness fluctuates randomly, rather than cyclically. Iteration of Eq. (3) was performed

using the same set of parameters above. However, at each generation a random integer t^* was drawn from 1

to λ. Then, the carrying capacity and selection coefficient were given by $K_f = K_{f0}\exp\left[\beta\sin[2\pi t^*/\lambda]\right]$ and $s_t =$

$s_{max}\sin[2\pi t^*/\lambda]$. Therefore, if β > 0 and the fitness of the A_2 allele is $W_{2f} = e^{s_t}$ or $1 + s_t$, K_f and W_{2f}

fluctuate in the same direction; it models an allele that, for example, leads to the immediate consumption of all

available resources, the amount of which changes randomly, rather than even allocation of resources over

successive generations. For each parameter set, 100 iterations were made and the mean frequency trajectory

was obtained (Fig. S2).

Random fitness fluctuation yielded all major results observed with cyclic fitness fluctuation: GMF

alleles positively selected and then maintained at intermediate frequency under strong density regulation with

constant carrying capacity (Fig S2A, B), GMF alleles with wider fluctuation in absolute fitness positively selected

toward fixation under weak density regulation with or without population size fluctuation (Fig. S2B, C), GMF

and AMF alleles whose fitness fluctuates in the same direction as population size positively selected toward

fixation under strong density regulation (Fig. S2C, D), and, with a smaller magnitude of population size

fluctuation (β = 0.3 instead of 0.5), AMF alleles positively selected and then maintained polymorphic under

strong density regulation (Fig. S2E). The polymorphic equilibria reached in the last cases were confirmed in

longer iterations (Fig. S3).

Figure S2. Changes in the absolute frequencies (n_1 and n_2) of A_1 (gray curves) and A_2 (dark curves) alleles under the random fluctuation of carrying capacity on the field ($K_{f0} = 1,000$, $K_f = K_{f0}\exp[\beta\sin[2\pi t^*/\lambda]]$, and $s_t = s_{max}\sin[2\pi t^*/\lambda]$, where t^* is a random integer between 1 and λ drawn each generation). Other parameters are identical to those in Fig. 3.

Figure S3. Paired histograms showing the distribution of A_2 frequency (q) at the 5,000th and 10,000th generation in the iteration of eq. (3) with random fitness fluctuation, corresponding to $g_f = 2$ (A) and 20 (B) of Fig. S2E. 1,000 iterations in total were performed for each parameter set.

DISCUSSION

The model of fluctuating selection in the presence of refuge was modified in terms of absolute fitness to accommodate the joint dynamics of population size and allele frequency changes. The general importance of investigating the coupled demographic and evolutionary dynamics was already well demonstrated. The outcome of natural selection under a changing population size can differ from that under a constant population size, as shown in the case of the fixation probability of beneficial alleles (OTTO AND WHITLOCK 1997) or the nature of fitness that is maximized (SÆTHER AND ENGEN 2015). Conversely, various demographic changes can occur as the result of natural selection, as in the case of evolutionary rescue or trapping (CARLSON et al. 2014). Furthermore, as shown in studies using simple mathematical models (PIMENTEL 1961; CHARLESWORTH 1971; ROUGHGARDEN 1971; LION 2018), demographic and evolutionary dynamics occur in a tightly-coupled process, which is described as an eco-evolutionary feedback (KOKKO AND LÓPEZ-SEPULCRE 2007; GOVAERT et al. 2019).

This study found that the model of fluctuating selection with refuge makes drastic changes in prediction if it is built upon absolute instead of relative fitness. First, under weak density regulation, a GMF allele that causes a wider fluctuation in population size is under positive directional selection instead of negative frequency-dependent selection. The latter, referred to as the storage effect, has long been proposed as a mechanism for promoting species or genetic diversity (CHESSON AND WARNER 1981; BERTRAM AND MASEL 2019). It was also clarified that AMF alleles are not subject to either balancing or directional selection in the presence of refuge if the carrying capacity of the population (the field) is constant over time. These results therefore calls for significant revisions in conclusions drawn from previous studies on the storage effect.

Second, more surprisingly, preexisting fluctuation in population size was found to select an allele,

regardless of whether it is GMF or AMF, if its relative fitness fluctuates in the same direction with population

size under strong density regulation. In case of derived AMF alleles, either negative, balancing, or directional

selection on them occurs depending on the magnitude of fitness fluctuation relative to population size

fluctuation. To my knowledge this type of positive selection, acting on either quasi-neutral (GMF) or deleterious

(AMF) alleles outside the refuge, induced by demographic changes was not discovered before. The emergence

of balancing selection – leading to the long-term oscillation of an AMF allele at intermediate frequencies – driven

by population size fluctuation is particularly remarkable. Although this mechanism also requires the presence

of a refuge, it should be distinguished from the storage effect because the latter arises as a larger arithmetic

mean of a derived allele's relative fitness is converted into a larger geometric mean due to the variance-reducing

effect of refuge. Here, the derived allele is not conferred a larger arithmetic mean. Mutations to AMF alleles may

arise frequently as many life history traits are subject to trade-off given a finite resource. Then, as the whole

genome is affected by population size fluctuation, positive selection on AMF alleles can arise at many loci

independently as long as their fitness fluctuation is in phase with demographic fluctuation. This is a potential

explanation for the seasonal oscillation of allele frequencies at more than 1,000 loci in North American

Drosophila melanogaster populations (BERGLAND *et al.* 2014). While further refinement of theory is likely needed

and the ecology of these fruit flies needs to be elucidated before confirming whether this novel mechanism of

balancing selection can operate in the actual population, the major requirements for the theory – clear

seasonality in the temperate region and the pupal stage that can serve as a refuge – appear to be satisfied in

these populations.

The fluctuation of population size in response to either a randomly or cyclically fluctuating

environment is observed in many plant and animal species, particularly those reproducing in multiple

generations over one seasonal cycle. Classical results in population genetics predict that a genotype causing less

fluctuation in fitness is favored under such an environment (GILLESPIE 1977; LANDE et al. 2009). Namely, studies

suggested that, given a mutation increasing fitness in one phase of a temporally changing environment and

decreasing it in another phase as a trade-off, it is positively selected if the trade-off occurs in a direction that

decreases the amplitude of population size fluctuation. Such a mutation, for example, would change the

allocation of resources toward a season unfavorable for the survival and/or reproduction of a species. This study,

however, showed that natural selection can occur in the opposite direction if a population is already partially

protected from selection. Under weak or moderate density regulation, even if the population size is initially

constant over time, a GMF allele causing population size fluctuation is positively selected toward fixation (Fig.

2B). If the population size is already fluctuating due to external forces (a variable environment) and/or due to

the fixation of above-mentioned allele at another locus, an allele causing a wider fluctuation of absolute fitness

can invade and therefore amplify the fluctuation in population size under moderate density regulation (cases

with g_f = 0.02 and 0.2 in Figs. 3 and 4). Such an allele, for example, allocates further resources to a period during

which the rate of growth and reproduction is already high, which inevitably causes further deterioration of

fitness during other periods. Assuming a large number of loci at which mutations affecting life history traits can

arise, one can imagine that selection at multiple loci can reinforce each other until a very wide fluctuation in

population size arises.

In the theory of evolutionary bet-hedging (SEGER AND BROCKMANN 1987; PHILIPPI AND SEGER 1989), a

refuge from selection such as a seed bank emerges in a reproductive strategy that may decrease the arithmetic

mean but increase the geometric mean of fitness by reducing the variance of reproductive success. Namely,

plants' delayed germination or insects' diapause can be the result of natural selection toward less severe

fluctuations in fitness. In this study, however, refuge from selection is given as a fixed demographic parameter

in the model under which selection acts on variants causing different patterns of fitness fluctuations. Given the

current results, findings in the previous investigation on evolutionary bet-hedging may be interpreted

differently. For example, by fitting the field observation of annual desert plants to their mathematical model,

which closely resembles Eq. (3), GREMER AND VENABLE (2014) showed that species experiencing wider fluctuation

in fitness after germination have greater proportions of their population remaining in the seed bank. This

supports the hypothesis that delayed germination evolved as a bet-hedging strategy. However, it could be other

life history traits that evolved in those species; in species with a larger probability of staying in the refuge (larger

seed bank), variants causing wider fluctuations in fitness in the field (after germination) could have been

selected, as suggested in this study.

The storage effect discovered in the previous studies and evolution toward amplified fluctuation in

population size discovered in this study depend on the occurrence of mutations that cause fitness fluctuations

without decreasing their geometric mean relative to the ancestral allele. Because these GMF alleles are required

to produce more offspring than their ancestral alleles within an environmental cycle, they may be considered as

a type of beneficial allele. Such mutations probably arise less frequently than mutations generating AMF alleles,

which can arise for example by simply changing resource allocation over different seasons. However, there can

be many biological scenarios under which mutants' fluctuating fitness arise in a way to preserve the geometric

mean. For example, imagine that the body size of a haploid individual carrying A_2 allele relative to A_1 is 1 + x (>

1) in one season and 1 - x in the next season, compatible with the trade-off of finite resource. Assume further

that, when they encounter a predator with probability y for both seasons, the predator may always choose the

smaller individual to attack. Then, the relative fitness of A_2 becomes $1/(1 - y)$ in the first season and $1 - y$ in

the second season, making A_2 the GMF allele.

Acknowledgement

This study was supported by the Research Foundation (NRF) grant 2020R1A2C1009261 funded by the Korean

government.

Conflict of interest statement

The author declares no conflict of interest

References

Barton, N. H., 2000 Genetic hitchhiking. Philosophical Transactions of the Royal Society of London. Series B:
Biological Sciences 355**:** 1553-1562.

Bell, G., 2010 Fluctuating selection: the perpetual renewal of adaptation in variable environments. Philos Trans
R Soc Lond B Biol Sci 365**:** 87-97.

Bergland, A. O., E. L. Behrman, K. R. O'Brien, P. S. Schmidt and D. A. Petrov, 2014 Genomic evidence of rapid and
stable adaptive oscillations over seasonal time scales in Drosophila. PLoS Genet 10**:** e1004775.

Bertram, J., and J. Masel, 2019 Different mechanisms drive the maintenance of polymorphism at loci subject to
strong versus weak fluctuating selection. Evolution 73**:** 883-896.

Carlson, S. M., C. J. Cunningham and P. A. Westley, 2014 Evolutionary rescue in a changing world. Trends Ecol
Evol 29**:** 521-530.

Charlesworth, B., 1971 Selection in density-regulated populations. Ecology 52**:** 469-474.

Chesson, P. L., and R. R. Warner, 1981 Environmental variability promotes coexistence in lottery competitive
systems. The American Naturalist 117**:** 923-943.

Dempster, E. R., 1955 Maintenance of genetic heterogeneity, pp. 25-32 in *Cold Spring Harbor Symposia on*
Quantitative Biology. Cold Spring Harbor Laboratory Press.

Ellner, S., and N. G. Hairston, 1994 Role of overlapping generations in maintaining genetic variation in a
fluctuating environment. The American Naturalist 143: 403-417.

Gillespie, J. H., 1977 Natural selection for variances in offspring numbers: a new evolutionary principle. The
American Naturalist 111: 1010-1014.

Gillespie, J. H., 1994 *The causes of molecular evolution*. Oxford University Press.

Govaert, L., E. A. Fronhofer, S. Lion, C. Eizaguirre, D. Bonte *et al.*, 2019 Eco-evolutionary feedbacks—Theoretical
models and perspectives. Functional Ecology 33: 13-30.

Grant, P. R., and B. R. Grant, 2002 Unpredictable evolution in a 30-year study of Darwin's finches. Science 296:
707-711.

Gremer, J. R., and D. L. Venable, 2014 Bet hedging in desert winter annual plants: optimal germination strategies
in a variable environment. Ecol Lett 17: 380-387.

Gulisija, D., and Y. Kim, 2015 Emergence of long-term balanced polymorphism under cyclic selection of spatially
variable magnitude. Evolution 69: 979-992.

Gulisija, D., Y. Kim and J. B. Plotkin, 2016 Phenotypic plasticity promotes balanced polymorphism in periodic
environments by a genomic storage effect. Genetics 202: 1437-1448.

Haldane, J., and S. Jayakar, 1963 Polymorphism due to selection of varying direction. Journal of Genetics 58: 237-
242.

Hedrick, P. W., 1976 Genetic variation in a heterogeneous environment. II. Temporal heterogeneity and
directional selection. Genetics 84: 145-157.

Hedrick, P. W., 1986 Genetic polymorphism in heterogeneous environments: a decade later. Annual review of
ecology and systematics: 535-566.

Hedrick, P. W., 1995 Genetic polymorphism in a temporally varying environment: effects of delayed germination
or diapause. Heredity 75: 164-170.

Huerta-Sanchez, E., R. Durrett and C. D. Bustamante, 2008 Population genetics of polymorphism and divergence
under fluctuating selection. Genetics 178: 325-337.

Kokko, H., and A. López-Sepulcre, 2007 The ecogenetic link between demography and evolution: can we bridge
the gap between theory and data? Ecology Letters 10: 773-782.

Lande, R., S. Engen and B.-E. Saether, 2009 An evolutionary maximum principle for density-dependent
population dynamics in a fluctuating environment. Philosophical Transactions of the Royal Society B:
Biological Sciences 364: 1511-1518.

Lion, S., 2018 Theoretical approaches in evolutionary ecology: environmental feedback as a unifying perspective.
The American Naturalist 191: 21-44.

Machado, H. E., A. O. Bergland, R. Taylor, S. Tilk, E. Behrman *et al.*, 2021 Broad geographic sampling reveals the
shared basis and environmental correlates of seasonal adaptation in Drosophila. eLife 10.

Myers, J. H., 1988 Can a general hypothesis explain population cycles of forest Lepidoptera?, pp. 179-242 in
Advances in Ecological Research. Elsevier.

Otto, S. P., and M. C. Whitlock, 1997 The probability of fixation in populations of changing size. Genetics 146**:**
723-733.

Park, Y., and Y. Kim, 2019 Partial protection from cyclical selection generates a high level of polymorphism at
multiple non-neutral sites. Evolution 73**:** 1564-1577.

Philippi, T., and J. Seger, 1989 Hedging one's evolutionary bets, revisited. Trends in ecology & evolution 4: 41-44.

Pimentel, D., 1961 Animal population regulation by the genetic feed-back mechanism. The American Naturalist
95**:** 65-79.

Reinhold, K., 2000 Maintenance of a genetic polymorphism by fluctuating selection on sex-limited traits. Journal
of Evolutionary Biology 13**:** 1009-1014.

Roughgarden, J., 1971 Density-dependent natural selection. Ecology 52**:** 453-468.

Seger, J., and H. J. Brockmann, 1987 Oxford surveys in evolutionary biology. Oxford Surveys in Evolutionary
Biology 4**:** 182-211.

Sæther, B.-E., and S. Engen, 2015 The concept of fitness in fluctuating environments. Trends in Ecology &
Evolution 30**:** 273-281.

Svardal, H., C. Rueffler and J. Hermisson, 2015 A general condition for adaptive genetic polymorphism in
temporally and spatially heterogeneous environments. Theoretical Population Biology 99**:** 76-97.

Thompson, J. N., 2013 *Relentless evolution*. University of Chicago Press.

Turelli, M., D. W. Schemske and P. Bierzychudek, 2001 Stable two-allele polymorphisms maintained by
fluctuating fitnesses and seed banks: protecting the blues in Linanthus parryae. Evolution 55: 1283-
1298.

Wittmann, M. J., A. O. Bergland, M. W. Feldman, P. S. Schmidt and D. A. Petrov, 2017 Seasonally fluctuating
selection can maintain polymorphism at many loci via segregation lift. Proceedings of the National
Academy of Sciences 114**:** E9932-E9941.

Appendix A. Growth rate under fluctuating absolute fitness in the presence of refuge

At the limit of no density regulation ($g_f = 0$) the number, n, of haploid individuals carrying a given allele changes independently of the other allele. Assuming that $m_{fr} = r$ and $m_{rf} = 1 - r$ and that this allele has the absolute fitness of W_f and W_r in the field and in the refuge, respectively, n changes to $n((1 - r)W_f + rW_r)$ over a generation. Starting from n_0 the number of individuals after a single cycle of fitness fluctuation is given by

$$n_\lambda = n_0 \prod_{t=1}^{\lambda} \big((1-r)W_f + rW_r\big) = n_0 \exp\big(\sum_{t=1}^{\lambda} \log\big[(1-r)W_f + rW_r\big]\big). \tag{A1}$$

First, the absolute fitness of a GMF allele can be given by $W_f = \exp\left[\alpha \sin\left(\frac{2\pi t}{\lambda}\right)\right]$, where α determines the amplitude of fluctuation, and $W_r = 1$. Then,

$$\log\big[(1-r)W_{f,t} + rW_r\big] = \log\left[(1-r)\exp\left[\alpha \sin\left[\frac{2\pi t}{\lambda}\right]\right] + r\right] \approx \alpha(1-r)\sin\left[\frac{2\pi t}{\lambda}\right] + \frac{\alpha^2 r(1-r)}{4}\left(1 - \cos\left[\frac{4\pi t}{\lambda}\right]\right)$$

after Taylor expansion around $\alpha = 0$ up to the second order. Plugging the above into eq. (A1) yields,

$$n_\lambda \approx n_0 \exp\left[\frac{1}{4}\alpha^2 r(1-r)\lambda\right] \approx n_0(1 + \frac{1}{4}\alpha^2 r(1-r)\lambda). \tag{A2}$$

Namely, the absolute number of individuals increases as long as their absolute fitness fluctuates in the field, when no such change is expected in the field or refuge in isolation ($\prod_t W_f = 1$ and $W_r = 1$).

Second, in the case of an AMF allele, absolute fitness can be given by $W_f = 1 + \alpha \sin\left(\frac{2\pi t}{\lambda}\right)$ and $W_r = 1$. Using Taylor approximation again,

$$\log\big[(1-r)W_{f,t} + rW_r\big] = \log\left[(1-r)(1 + \alpha \sin\left[\frac{2\pi t}{\lambda}\right]) + r\right] \approx \alpha(1-r)\sin\left[\frac{2\pi t}{\lambda}\right] - \frac{\alpha^2(1-r)^2}{4}\left(1 - \cos\left[\frac{4\pi t}{\lambda}\right]\right).$$

Then,

$$n_\lambda \approx n_0 \exp\left[-\frac{1}{4}\alpha^2(1-r)^2\lambda\right] \approx n_0(1 - \frac{1}{4}\alpha^2(1-r)^2\lambda). \tag{A3}$$

A refuge therefore cannot turn the negative growth rate of a subpopulation carrying an AMF allele from a negative to a positive value. A wider fluctuation of fitness accelerates this decrease.

573 **Appendix B. Demography-driven selection in the two-season model under strict density regulation**

574 One environmental cycle consists of two generations, which correspond to two seasons, good and bad, with the

575 carrying capacity of the field given by K_G and K_B, respectively. At the start of each generation (season), haploid

576 individuals migrate to the field and the refuge with probability 1 - r and r regardless of their current location. At

577 the end of bad season there are N_B individuals. During the good season, the population size of the field changes

578 from $(1-r)N_B$ to K_G while that of the refuge remains at rN_B since $W_r = 1$ is assumed. Total population

579 size at the end of a good season is therefore $N_G = K_G + rN_B$. Similarly, $N_B = K_B + rN_G$. Therefore, $N_G = (K_G +$

580 $rK_B)/(1-r^2)$ and $N_B = (K_G + rK_B)/(1-r^2)$. Let n_1 and n_2 be the number of A_1 and A_2 individuals at the end

581 of a bad season (therefore, $n_1 + n_2 = N_B$). Then, n_2 increases to

582
$$n'_2 = (1-r)n_2 W_{2G} \frac{K_G}{(1-r)n_1 W_{1G} + (1-r)n_2 W_{2G}} + rn_2 = n_2 \left(\frac{K_G}{n_1/\omega_G + n_2} + r\right) = n_2 \left(\frac{K_G}{N_B} \frac{\omega_G}{p+q\omega_G} + r\right) \qquad \text{(B1)}$$

583 at the end of good season, where $\omega_G = W_{2G}/W_{1G}$ is the relative fitness of A_2 in the field in good season and $q =$

584 $1 - p = n_2/(n_1 + n_2)$. Using also

585
$$n'_1 = n_1 \left(\frac{K_G}{n_1 + n_2\omega_G} + r\right),$$

586 the number of A_2 individuals at the end of the following winter generation changes to

587
$$n''_2 = n'_2 \left(\frac{K_B}{n'_1/\omega_B + n'_2} + r\right) = n'_2 \left(\frac{K_B\omega_B(n_1 + n_2\omega_G)}{K_G(n_1 + n_2\omega_G\omega_B) + r(n_1 + n_2\omega_B)(n_1 + n_2\omega_G)} + r\right)$$

588
$$= n'_2 \left(\frac{K_B\omega_B(1-q+q\omega_G)}{K_G(1-q+q\omega_G\omega_B) + rN_B(1-q+q\omega_B)(1-q+q\omega_G)} + r\right), \qquad \text{(B2)}$$

589 where $\omega_B = W_{2B}/W_{1B}$. When A_2 is rare ($q \to 0$) the change in its absolute frequency after one cycle is obtained

590 from eqs. (B1) and (B2):

591
$$\frac{n''_2}{n_2} = \frac{n'_2}{n_2}\frac{n''_2}{n'_2} = \left(\frac{K_G}{N_B}\omega_G + r\right)\left(\frac{K_B}{K_G + rN_B}\omega_B + r\right) = \left(\frac{K_G}{N_B}\omega_G + r\right)\left(\frac{K_B}{N_G}\omega_B + r\right).$$

592 If $\omega_G = \omega_B = 1$, the above becomes

593
$$\left(\frac{K_G}{N_B} + r\right)\left(\frac{K_B}{N_G} + r\right) = \frac{(K_G + rK_B)(K_B + rK_G)}{N_B N_G} = 1.$$

594 Then, if $\omega_G = 1/\omega_B = \omega$, the condition for A_2 allele invading the population $\left(\frac{K_G}{N_B}\omega + r\right)\left(\frac{K_B}{N_G}\frac{1}{\omega} + r\right) > 1$ is

595 rewritten as

$$r\frac{K_\mathrm{G}}{N_\mathrm{B}}\omega^2 + \left(\frac{K_\mathrm{G}K_\mathrm{B}}{N_\mathrm{B}N_\mathrm{G}} + r^2 - 1\right)\omega + r\frac{K_\mathrm{B}}{N_\mathrm{G}} = r\left(\frac{K_\mathrm{G}}{N_\mathrm{B}}\omega - \frac{K_\mathrm{B}}{N_\mathrm{G}}\right)(\omega - 1) > 0.$$

If $\omega_\mathrm{G} = 1 + s$ and $\omega_\mathrm{B} = 1 - s$, the condition for the invasion of A_2 is

$$\left(\frac{K_\mathrm{G}}{N_\mathrm{B}}(1 + s) + r\right)\left(\frac{K_\mathrm{B}}{N_\mathrm{G}}(1 - s) + r\right) = -\frac{K_\mathrm{G}K_\mathrm{B}}{N_\mathrm{B}N_\mathrm{G}}s^2 + r\left(\frac{K_\mathrm{G}}{N_\mathrm{B}} - \frac{K_\mathrm{B}}{N_\mathrm{G}}\right)s + 1 > 1,$$

which is satisfied for

$$0 < s < r\left(\frac{N_\mathrm{G}}{K_\mathrm{B}} - \frac{N_\mathrm{B}}{K_\mathrm{G}}\right) = \frac{r}{1 - r^2}\left(\frac{K_\mathrm{G}}{K_\mathrm{B}} - \frac{K_\mathrm{B}}{K_\mathrm{G}}\right).$$

At the same time, the condition for the invasion of A_1, $\dfrac{n''_1}{n_1} = \left(\frac{K_\mathrm{G}}{N_\mathrm{B}}\frac{1}{1+s} + r\right)\left(\frac{K_\mathrm{B}}{N_\mathrm{G}}\frac{1}{1-s} + r\right) > 1$ is rearranged to

$$(1 - r^2)N_\mathrm{B}N_\mathrm{G}s^2 + r(N_\mathrm{B}K_\mathrm{B} - N_\mathrm{G}K_\mathrm{G})s = (1 - r^2)N_\mathrm{B}N_\mathrm{G}s\left(s - \frac{r(K_\mathrm{G}+K_\mathrm{B})(K_\mathrm{G}-K_\mathrm{B})}{(K_\mathrm{B}+rK_\mathrm{G})(K_\mathrm{G}+rK_\mathrm{B})}\right) > 0.$$