Possible role of left–right asymmetry in the sensory system and behavior during adaptation to food-sparse cave environments

Vânia Filipa Lima Fernandes[1], Yannik Glaser[2], Motoko, Iwashita[1], and Masato Yoshizawa[1]

[1] School of Life Sciences, University of Hawai'i at Mānoa, Honolulu, Hawai'i, USA

[2] Department of Information and Computer Sciences, University of Hawai'i at Mānoa, Honolulu, Hawai'i, USA

Corresponding author: Masato Yoshizawa (yoshizaw@hawaii.edu)

Abstract

Laterality in relation to behavior and sensory systems is found commonly in a variety of animal taxa. Despite the advantages conferred by laterality (e.g., the startle response and complex motor activities), little is known about the evolution of laterality and its plasticity in response to ecological demands. In the present study, a comparative study model, the Mexican tetra (*Astyanax mexicanus*), composed of two morphotypes, i.e., riverine surface fish and cave-dwelling cavefish, was used to address the relationship between environment and laterality. The use of a machine learning-based fish posture detection system and sensory ablation revealed that the left cranial lateral line significantly supports one type of foraging behavior, i.e., vibration attraction behavior, in one cave population. Additionally, left–right asymmetric approaches toward a vibrating rod became symmetrical after fasting in one cave population but not in the other populations. Based on these findings, we propose a model explaining how the observed sensory laterality and behavioral shift could help adaptations in terms of the tradeoff in energy gain and loss during foraging according to differences in food availability among caves.

Conflict of Interest: The authors declare no competing financial interests.

Acknowledgement: We thank L Espinasa for constructive comments regarding insights on variation of laterarities in *Astyanax* cavefish. We also thank to H Yoshizawa for supporting the repeatability experiments and fish care assistance. We are grateful to H Hernandez, N Doeden, C Balaan, V Crystal, J Choi, L Lu, J Nguyen, K Lactaoen, M Worsham, J Kato, M Ito, R Balmilero-Unciano, E Doy, A Martinez, and D Mones for fish care assistance. We gratefully acknowledge support from the National Institute of Health (P20GM125508) to MY, Hawaii Community Foundation (18CON-90818) to MY.

Author contribution:

VFLF: designed and performed experiments, wrote original draft, edited the manuscript, performed all
experiments, and analyzed results.
YG: designed the detection and analysis algorithm and helped with actual analysis related to
DeepLabCut-based aanalysis, consulted the experimental procedure, and edited the manuscript.
MI: performed experiments, edited the manuscript, analyzed the results, and developed a mathematical
model.
MY: designed the experiment, edited the manuscript,co-developed DeepLabCut-based analysis method
and algorithm, and analyzed the results.
45

Introduction

Laterality, which is common among various animal taxa, is expressed, for example, as the asymmetrical use of the left or right hand or foot (Tommasi, 2009), the biased use of the left or right eye, and the use of the left or right brain hemispheres for different processing tasks (Heilman and Abell, 1980; Nottebohm, 1977, 1971; Nottebohm and Nottebohm, 1976; Rogers, 1981; Rogers and Workman, 1989; Schwartz et al., 1975; Wernicke, 1874). Laterality confers several advantages in recognizing and escaping from predators (Chivers et al., 2017, 2016; Dadda et al., 2010), facilitating complex and intricate motor activities (Magat and Brown, 2009) and enabling spatial learning (Sovrano et al., 2005). Phylogenetic studies have shown that left–right (L–R) asymmetric traits have repeatedly and independently evolved among different animal taxa (Güntürkün et al., 2020), possibly due to the need for laterality in diverse environments. However, the influence of environments on variation in the laterality and its plasticity among populations has not been well-studied.

As an entry point to solve this evolutionary question, we explored the environment–laterality relationship in *Astyanax mexicanus*, the Mexican tetra, which has two interfertile morphotypes, i.e., surface fish and cavefish (Keene et al., 2016; Protas and Jeffery, 2012; Rétaux et al., 2013; Wilkens et al., 1988) and is known as a model for studying the evolution of behavior. Cavefish, the blind cave-dwelling form of *A. mexicanus*, were separated from their surface-dwelling ancestors about 2,000–20,000 years ago (Fumey et al., 2018; Herman et al., 2018). In total, 32 *A. mexicanus* cave populations have been described in Mexico (Espinasa et al., 2020), and many of these populations have evolved morphological, physiological, and behavioral traits independently, likely via adaptation (Borowsky, 2008; Mitchell et al., 1977; Ornelas-García et al., 2008; Strecker et al., 2012; Wilkens et al., 1988).

One type of laterality has been reported in the Pachón cave population of *A. mexicanus* cavefish, i.e., a preference for using the right-side of the body to detect novel immotile objects in the environment (de Perera et al., 2005). However, it is not yet known which environmental factors determine this type of laterality (unclear ecological relevance) or whether it evolved in a surface fish-like ancestor because such laterality in surface fishes has not been investigated. In the present study, we investigated a well-characterized type of foraging behavior, vibration attraction behavior (VAB), which is evoked in response to 35–40 Hz vibrations (Yoshizawa et al., 2010) within the range potentially produced by soil crustaceans and percolating water (Lang, 1980; Montgomery and Macdonald, 1987). VAB is evolutionarily enhanced in cavefish, and its advantage was demonstrated as they hunt small prey in the dark (Yoshizawa et al.,

2010). Different degrees of increased VAB levels have been reported in lab cave populations

and in the original wild cave populations (Espinasa et al., 2021; Yoshizawa et al., 2010), likely

due to differences in ecological factors in these caves. Outside of cave environments, VAB

may be detrimental because of nocturnal predators, such as prawn fish (Wilson et al., 2004;

Yoshizawa and Jeffery, 2011).

VAB is mediated by the mechanosensory lateral line system, which is composed of sensory

units known as superficial neuromasts (SNs) particularly located at the eye orbit and the third

infraorbital bone (IO3) (Yoshizawa et al., 2012b, 2010). We reported previously that, despite

L–R symmetry in the number of SNs in surface and cave morphs, a positive correlation existed

between the number of left SNs and VAB level in Pachón cavefish (Fernandes et al., 2018).

This asymmetric sensory usage (hereafter termed "sensory laterality") was not observed in

surface fish, suggesting that this sensory laterality emerged through an evolutionary process

that conferred some advantage to fish that colonized cave environments (Fernandes et al.,

2018).

In the current study, we investigated the asymmetric L–R responses to a vibrating stimulus in

A. mexicanus populations. In contrast to the Pachón cavefish, independent cavefish

populations from the Los Sabinos and Tinaja caves did not show detectable coupling between

the left or right SN and VAB levels, indicating that asymmetric sensory–behavior coupling

could be adaptive in the Pachón cave. SN ablation revealed that the left SN in Pachón

cavefish indeed regulate the number of approaches (NOA) level variation. In Tinaja cavefish,

we found alternative L–R asymmetry in their approaches toward a vibrating rod that did not

depend on the number of L–R SNs but was dependent on a fasting condition. Specifically,

Tinaja individuals that exhibited asymmetric L–R approaches prior to fasting showed L–R

balanced approaches after fasting. Such fasting-dependent L–R plasticity was not detected in

the Pachón cavefish or surface fish populations.

Our results indicate that different cave populations of *A. mexicanus* have evolved distinct

laterality strategies. Our study also suggests that L–R sensory usage and the L–R asymmetric

approach have different underlying mechanisms. Based on these findings, we propose a

model that explains the advantages of laterality according to diet availability and energy gain

and loss. This model implies that the laterality of Pachón cavefish saves energy most

effectively among the studied cave and surface populations and may have allowed Pachón

cavefish to adapt successfully to the fluctuating availability of food in the Pachón cave.

Results

Sensory laterality among different cave populations of *A. mexicanus*

In a previous study, we reported a positive correlation between the VAB level and the number of left- but not right-side SNs in Pachón cavefish (Fernandes et al., 2018). This L–R asymmetric association between the number of SNs and VAB likely evolved during the cave adaptation process. However, whether such left SN–VAB coupling exists in other discrete cave populations in the context of convergent/parallel evolution was unclear. Thus, we investigated other VAB-positive cave populations in Tinaja and Los Sabinos (Espinasa et al., 2021; Yoshizawa et al., 2015, 2010) (Figure 1). First, to detect left or right approaches toward a vibrating rod, we developed a method using DeepLabCut (DLC), a machine-learning suite (Mathis et al., 2018; see Materials and Methods), with which we accurately measured the X–Y coordinates of the left and right craniums of fish, which was not possible with a method we developed previously (Yoshizawa et al., 2015, 2010). To validate the new method, we measured the number of approaches toward the vibrating rod (NOA) and the duration around the vibrating rod (DIR) in one surface fish and three cavefish populations (see Materials and Methods). Consistent with previous results (Yoshizawa et al., 2015, 2012a, 2010), Pachón cavefish showed the highest level of VAB in comparison to Los Sabinos and Tinaja cavefish and surface fish, both in terms of NOA and DIR (Figure 1C; Supplementary Table 1), indicating that the new detection method is consistent with the method developed previously (Yoshizawa, 2015; Yoshizawa et al., 2012a, 2010).

We then explored whether an L–R bias existed in the couplings between the number of SNs and NOA or DIR. In the surface population, no correlation was detected between the left SNs and the L–R summed NOA/DIR (i.e., total NOA/DIR) or the right SNs and the total NOA/DIR (Figure 1D; statistics in Supplementary Table 2; all NOA and DIR in Figs. 1 and 2 are L–R summed). In contrast, a positive correlation was found between the left SNs and NOA in the Pachón cave population (τ = 0.329, P = 0.047; Figure 1G; Supplementary Table 2) but not between the right SNs and NOA (τ = 0.258, P = 0.118; Figure 1G; Supplementary Table 2). This result is also consistent with our previous study (Fernandes et al., 2018); thus, we tested the other cave populations with confidence. Both the Los Sabinos and Tinaja cave populations showed results similar to those of the surface population, i.e., no correlation was detected between the left SNs and NOA/ DIR or between the right SNs and NOA/ DIR (Figure 1E and F; Supplementary Table 2). In summary, our machine-learning-based detection method generated results consistent with our previous studies, showing that the left SN number is

151   coupled with the NOA level in the Pachón cave population. However, this method failed to

152   detect such correlations in other cave populations.

Coupling between the left SNs and VAB is based on inputs to the left SNs

To investigate the underlying mechanism of the coupling between the left SNs and VAB, we

ablated the SNs on either the left or right-side at the IO3 area in the Pachón cave populations

using a well-developed noninvasive method (Yoshizawa et al., 2010). To highlight SN–VAB

coupling, we selected VAB-positive individuals [NOA ≥ 6; (Yoshizawa et al., 2010)]. The

ablation was confirmed using DASPEI staining after the behavior assay (Figure 2A and B).

After ablation, Pachón cavefish individuals with a higher percentage of ablated SNs exhibited

lower levels of VAB (τ = −0.342, P = 0.009; Figure 2B; Supplementary Table 3). However, the

change in VAB was not observed when the right-side SNs were ablated (τ = −0.064, P =

0.635; Figure 2D; Supplementary Table 3). We also performed SN ablation in surface fish and

detected no correlation between NOA and the proportion of ablated SNs on the left or right

sides (Figure 2—figure supplement 1; Supplementary Table 3). These results reinforce our

previous findings (Fernandes et al., 2018) regarding the significant role of the left-side SNs in

the VAB of the Pachón cave population.

For DIR (i.e., the time spent located within 1.3 cm of the vibrating rod), no correlation was

found with the proportions of left or right ablated SNs in Pachón cavefish (Supplementary

Table 3), suggesting a possibility that the NOA and DIR are regulated differently, i.e., in

Pachón cavefish, NOA could be mediated by the SNs, whereas DIR could be regulated via

non-SN means, such as auditory and/or tactile sensing (see the Discussion).

Regarding population averages, for Pachón cavefish, the ablation of the left, right, or left and

right SNs induced no detectable changes in NOA or DIR (although the ablation of left-side SNs

increased NOA significantly: V = 74.5, P = 0.032; Figure 2—figure supplement 2;

Supplementary Table 4; See also Supplementary file 1—extended result and discussion).

In summary, for Pachón cavefish, the left SN number raises NOA but not DIR, and the right

SNs were not significantly associated with NOA.

Behavioral laterality across different populations of *A. mexicanus*

Our results showed that the left SNs promoted VAB in Pachón cavefish, whereas the other

populations did not show L–R-biased coupling between the SNs and VAB (i.e., sensory

laterality) (Figs. 1 and 2). Given that VAB evolved as a foraging behavior and that diet type and

availability differ among caves [e.g., from decomposed plants to soil or cave-adapted
arthropods as well as seasonal flooding (Culver, 1982; Culver and Pipan, 2009; Espinasa et
al., 2017; Keene et al., 2016; Mitchell et al., 1977; Yoshizawa et al., 2010; Yoshizawa and
Jeffery, 2011)], we hypothesized that alternative forms of laterality in VAB exist in surface
and/or cave populations. Thus, we explored two types of L–R bias in VAB (i.e., left/right NOA
and left/right DIR; Figure 3A) among the surface population and Pachón, Tinaja, and Los
Sabinos cave populations. To quantify the lateral preference of individual fish, the right NOA or
DIR was divided by the total NOA or total DIR, respectively (i.e., ratio scores of >0.67 and
<0.33 indicate right and left preferences, respectively; Figure 3B and 3C). We measured the
L–R biases of NOA and DIR separately because our results indicated that they were likely
regulated differently (see the previous section and Discussion). Individuals from each of the
cave and surface populations showed a preference for one side or the other while approaching
(NOA) and touching the rod (DIR) (Figure 3B and 3C, respectively). This L–R preference in
each individual was not repeatable or only fairly repeatable in surface fish and Pachón
cavefish when we conducted assays 6 days apart and repeated three times (Table 1)
[respective NOA intraclass correlation coefficients (ICCs): κ = 0.0 (none), P = 0.48; κ = 0.39
(fair), P = 0.00025. Respective DIR ICCs: κ = 0.0 (none), P = 0.48; κ = 0.085 (none), P = 0.20].
In contrast, Tinaja cavefish individuals exhibited a moderate level of repeatability in NOA (κ =
0.54, P = 0.0021; repeated three times; Table 1), suggesting that these fish tended to
consistently approach the vibrating rod unidirectionally.
204
205 Table 1. Intraclass correlation coefficient (ICC) under Cohen's suggestion (Landis and Koch, 1977;
206 Zegers et al., 2010) (N = 3 repeats; N= 22 for surface, N = 9 for Pachón cave and N = 11 for Tinaja
207 cave fish).

Population	Measurement	Average L–R ratio	Statistics	Kappa	P-values
Surface fish	NOA	0.459 ± 0.034	$F_{21,42}$ = 1.0	0.0 (none)	0.48
Surface fish	DIR	0.481 ± 0.028	$F_{21,42}$ = 1.0	0.0 (none)	0.48
Pachón cavefish	NOA	0.464 ± 0.036	$F_{28,56}$ = 3.0	0.39 (fair)	0.00025***
Pachón cavefish	DIR	0.473 ± 0.024	$F_{28,56}$ = 1.3	0.085 (none)	0.20
Tinaja cavefish	NOA	0.482 ± 0.052	$F_{10,20}$ = 4.5	0.54 (moderate)	0.0021**
Tinaja cavefish	DIR	0.478 ± 0.046	$F_{10,20}$ = 2.4	0.32 (fair)	0.043*

208
As shown in the histograms in Figure 3B and C, there was no detectable bias between the left
and right-side NOA ($F_{1,152}$ = 0.149, P = 0.700) or between the left and right DIR among the
populations ($F_{1,152}$ = 3.378, P = 0.068) (Figure 3—figure supplement 1; Supplementary Table
1). Additionally, there was no detectable bias between the duration of left and right facing

outside the 1.3-cm radius around the vibrating rod, suggesting that there was no tendency to

face left or right in relation to the vibrating rod from a distance ($F_{3,152}$ = 0.016, P = 0.997;

Supplementary Table 1).

The above results suggested that laterality in SN usage (sensory: Pachón cavefish) and

laterality in approaches (behavior: Tinaja cavefish) existed in some *A. mexicanus* populations.

Therefore, we investigated whether relationships existed between the lateralities in the number

of left or right SNs and left or right NOA or left or right DIR. Detailed correlation analyses were

performed in the ablation study. In Pachón cavefish, the ablation ratio of the left SNs mirrored

the reduction of left approaches (left NOA), but no correlation was detected between the left

SN ablation ratio and the right NOA, the right SN ablation ratio and the left or right NOA, and

the left or right SN ablation ratio and the left or right DIR (Supplementary Table 3). Additionally,

there was no detectable effect of any left or right SN ablation on the L–R approaches (NOA or

DIR) of surface fish. These results indicate that the number of left SNs (inputs) affects the left

approaches of Pachón cavefish but not those of surface fish. As in the intact SN condition,

there were some positive correlations between the number of L–R SNs and the L–R NOA

and/or DIR in surface fish and Pachón cavefish (but not in other cave populations; see

Supplementary Table 2). However, the SN–behavior relationship of surface fish in the intact

state may not be vital because it was not detected in the ablation study.

Collectively, these results indicate that sensory laterality was found only in Pachón cavefish

and was associated with left-side approaches. Individuals in the Tinaja cave population but not

in the surface or Pachón cave populations showed a moderately consistent preference for left-

or right-side approaches, and L–R bias was not detectable at the whole population level.

Fasting increased sensory responses and reduced approach laterality

Previous prey-capture experiments indicated that VAB promotes the food capture rate in dark

and food-sparse environments (Yoshizawa et al., 2010). VAB plasticity according to the

environment has also been suggested (Espinasa et al., 2021). Thus, we hypothesized that

fasting induces shifts in laterality. Because younger fish are more sensitive to fasting in

general, we used 1-year-old (young adult) surface fish, Pachón cavefish, and Tinaja cavefish

rather than the 2–3-years-old (adult) fish used in the earlier experiments (Figs. 1–3).

First, we investigated the effect of fasting on sensory laterality (i.e., the number of left or right

SNs vs. the total NOA or DIR) among the surface, Pachón, and Tinaja populations. The 6-day

fasting reduced the total NOA and total DIR in surface fish but increased the total DIR in Tinaja cavefish (Figure 4A and 4B; Supplementary Table 5); however, no changes were detected in Pachón cavefish (Figure 4A and 4B; Supplementary Table 5). To investigate L–R SN–behavior coupling, we determined the correlations between either the number of left or right SNs and the total NOA after fasting with subtraction of the prefasting NOA (Figure 4C; Supplementry Table 7). Pachón individuals with fewer "left" SNs exhibited a higher increase in total NOA (Figure 4C; Supplementry Table 7). Pachón individuals with fewer "right" SNs showed a higher increase in NOA but not at a significant level (Figure 4C; Supplementry Table 7). Surface fish or Tinaja cavefish did not show fasting-dependent changes in SN–approach coupling (Figure 4C; Supplementry Table 7). In summary, fasting yielded different responses in the surface and Pachón populations, i.e., NOA was reduced in surface fish, and Pachón individuals with fewer left SNs showed a higher increase in NOA relative to that of individuals with more left SNs. Second, we studied the effect of fasting on laterality in approaches. The surface and Tinaja populations had more number of younger fishes (1-year-old fish vs. 2–3-years-old fish) that showed L–R-biased approaches in the prefasting condition (NOA; broader standard variations; Figure 5A and 5B). In contrast, Pachón cavefish individuals showed L–R symmetrical NOA with a narrower standard variation than that observed in the surface and Tinaja populations in the prefasting condition (Figure 5A and 5B). After 6 days of food deprivation, the distributions of the L–R preference in surface fish became more uniform for both NOA and DIR (Figure 5A, 5B, 5C, and 5D). Interestingly, Tinaja cavefish shifted from a right-biased approach (for both NOA and DIR) to a symmetrical approach in NOA (~0.6 in a bias index; Figure 5C) and a reduced right bias in DIR (Figure 5). Regarding population averages, none of surface fish, Tinaja, or Pachón cavefish individuals showed detectable L–R bias in NOA or DIR in the prefasting or postfasting condition, except DIR in Tinaja cavefish (right bias; Figure 5—figure supplement 1A and 1B), which mirrors the observation in Figure 5A and 5B. Fasting also induces alternative responses among surface fish, Tinaja, and Pachón cavefish individuals; surface fish decreased the overall NOA and DIR after fasting, while Tinaja cavefish increased DIR (Figure 5—figure supplement 1A and 1B). Pachón cavefish did not show detectable changes in the overall NOA and DIR levels.

In summary, food deprivation drove behavioral shifts in the surface and Tinaja cave population but not in the Pachón cave population, i.e., surface fish showed reduced VAB, whereas Tinaja cavefish showed increased VAB (Figure 5, and Figure 5—figure supplement 1). Furthermore,

278 Tinaja cavefish showed a postfasting plastic change in the L–R bias of NOA toward a more L–

279 R symmetric approach (Figure 5, and Figure 5—figure supplement 1).

Discussion

We addressed laterality in sensory–behavior coupling and rod approaches among different

populations of *A. mexicanus*. Using a newly developed DLC- and machine learning-based

video analysis method, we found the following: (1) only the Pachón cave population showed a

bias in SN–VAB coupling, i.e., the left SNs promote VAB, but not detectable in other

populations; (2) no L–R bias detected in approaches (NOA) and adherence (DIR) in the

surface, Pachón, Tinaja, and Los Sabinos cave populations at the population levels, but some

individuals, especially those in the Tinaja cavefish population, repeatedly showed a preference

for left or right approaches; and (3) fasting for 6 days reduced the L–R-biased in approaches in

the Tinaja cave population, indicating that VAB plasticity is present in this population. Thus, L–

R asymmetry in foraging behavior seems to have evolved flexibly under different caves, and

our findings provide insights into the evolutionary dynamics of L–R asymmetry under

ecological pressure (see below). At the end, we discuss the energy-saving advantage

associated with foraging laterality.

Introducing a machine-learning method to assess adaptive foraging behavior

With the aid of a DLC neural network (trained with 807 images), we developed a new video

analysis method that achieved a detection accuracy of >99%, which was confirmed using a

manual survey of the labeled videos. In comparison with our previously reported method

(Yoshizawa et al., 2010), this trained network yielded higher resolution for the approach (NOA)

and adherence (DIR) events of VAB, and left or right facing in the direction of the vibrating rod

could be monitored. The new method was also confirmed to be consistent with the ImageJ-

based method developed previously (Figure 1), which supports the findings of the present

study as an expansion of the previous results.

Coupling between the left SNs and NOA but not DIR in Pachón cavefish

Pachón cavefish showed a significant correlation between the number of left SNs and NOA.

This coupling was observed repeatedly in different cohorts from different Pachón parents

(Fernandes et al., 2018). Furthermore, this sensoory laterality in NOA in Pachón cavefish was

confirmed in our ablation experiment, i.e., as more left SNs were ablated, the NOA was

reduced. Notably, a correlation between the left SNs and DIR was not detected in Pachón

cavefish. Such variation in sensory–behavior coupling suggests that differences exist in the

sensory mechanisms underlying NOA and DIR, i.e., given the correlation between the SNs and

NOA and our SN ablation results, NOA in Pachón cavefish is likely enhanced by SNs.

Contrastingly, DIR was neither correlated with the number of SNs nor changed after SN

ablation, indicating that other sensory systems, e.g., tactile and/or auditory sensors, affect DIR.

The auditory inner ear can sense near field particle motion based on the motion lag between

the fish body and the otolith (Kalmijn, 1988; Popper and Fay, 2011; Tricas, 2020). Overlapping

functions between auditory, lateral line, and tactile senses have been noted but not studied in

detail in the context of foraging (e.g., Braun and Sand, 2013; Kohashi et al., 2012). VAB thus

emerges as a good model behavior of the modality switch between these three sensory

systems because it can be evoked from the far field (>5 m in the field: auditory; see below;

Espinasa et al., 2021) and near field (possibly; >1.3 cm: lateral line; <1.3 cm: tactile).

Laterality in cranial morphology and behavior

There are reports of cranial bending in adult Pachón and Tinaja cavefish but not in adult

surface fish or younger individuals of the Pachón, Tinaja, and surface populations (Powers

et al., 2017). We did not detect left cranial bending in our experimental fish (2–3 cm

standard length; 1–3 years old), which excludes the confounding effect of head bending on

lateralities in sensory–behavior coupling and approaches to the rod. Other morphological

asymmetries have been reported previously, e.g., L–R asymmetry in the suborbital (i.e.,

infraorbital) bone fragmentations in Pachón and Tinaja cavefish but not in surface fish

(Gross et al., 2016); and L–R asymmetry in cranial SN distribution but not the number of

SNs, which corresponds to asymmetric bone fragmentation (Gross et al., 2016; Gross and

Powers, 2016). In the present study, we investigated the L–R asymmetric "number" but not

"distribution" of SNs. It will be interesting to investigate whether the SN distribution pattern

plays a role in VAB expression in future studies.

Laterality of approaches at the population and individual levels, and the effect of

starvation

In our L–R approach study, there was no detectable left or right bias for NOA and DIR at the

population level among the four tested populations (with the exception of 1-year-old Tinaja

cavefish). However, some individual fish showed one side bias in NOA and/or DIR, and these

asymmetric biases were consistent within each individual. Laterality at the individual level has been largely overlooked (Rogers, 1989); nonetheless, studies in chicks and fish showed that behavior lateralization was random in a population but existed and seemed advantageous (e.g., by minimizing the use of brain resources) at the individual level in complex tasks, including social interactions (Bisazza et al., 2000; Bisazza and Dadda, 2005; Rogers and Workman, 1989). In the current study, we also revealed plastic laterality in a cave population responding to fasting. Because VAB evolved as a foraging behavior under food-limited conditions (Yoshizawa et al., 2010; Yoshizawa and Jeffery, 2011), we propose that cavefish laterality evolved to conserve energy.

Many caves in North-eastern Mexico experience 6-months wet and 6-months dry seasons; thus, cavefish experience food-sparse conditions for ~6 months per year (Espinasa et al., 2017). We tested the effect of food deprivation on L–R asymmetry, and removing food intake for 6 days increased both DIR and L–R symmetric NOA in the starvation resistant Tinaja cavefish (Aspiras et al., 2015). The increases in DIR and symmetric NOA could help boost food intake while saving energy (see below). In contrast, the surface fish, which are less tolerant to starvation, showed decreased NOA and DIR, seemingly saving energy in a simpler manner by reducing the number of foraging attempts. The most derived Pachón cave population (Herman et al., 2018) is also starvation resistant but showed a nonsignificant change in symmetric NOA and DIR, perhaps because these cavefish have optimized their foraging under food-sparse conditions over the course of evolution (see below). In the Supplementary file 1—extended results and discussion, we also discuss age-dependent L–R plasticity in Tinaja cavefish (c.f., Figure 3B and C; Figure 5A and B) and the VAB levels of Los Sabinos and Tinaja cavefish.

Advantages of laterality in the cave ecosystem
To understand the evolution of laterality and its plasticity in the context of cave ecology, we propose a model that explains the potential advantages of such lateralities and its plasticity in terms of calorie gain and consumption in food-sparse caves (Figure 6). This model assumes that (1) only one side of SNs accounts for VAB evocation and (2) starvation induces L–R symmetric NOA. Based on these assumptions, we hypothesized that sensory laterality and non-L–R-biased (i.e., symmetrical) approaches are beneficial in food-limited environments at the population level due to the tradeoff between calorie gain and calorie use in foraging.

The model, including food/energy gain, energy loss by thrasting, and the change in the total
amount of calories gained in a population (= a sum of fish individuals), can be described as the
following function:

$\frac{dE}{dt} = E_g - E_c$ (equation 1; Supplementary file 1),

where E represents the total calories/energy of an average fish in a population gained per unit
of time t ($E = \sum_{i=1}^{N} E_i$, where i represents each individual fish and E_i represents the total
calories each fish gains per unit time), E_g represents the average total calories the population
gains from the diet per unit of time, and E_c is the average total calories the population uses
during foraging per unit of time.
Food capturing movements are assumed to use more calories, as the fish perform rapid and
short turns toward their food, than are used in a typical swimming motion (personal
observation). If the energy gain, E_g, is higher than the energy consumption, E_c (eq. 1), the
current fish in the population can survive in the given environment. If the opposite is true ($\frac{dE}{dt} <$
0; eq. 1), the number of fish in the population is reduced. We expect that unilateral sensing has
an advantage over bilateral sensing in terms of the tradeoff between E_g and E_c.
Depending on two scenarios, i.e., (1) only a unilateral input or (2) the inputs of both sides are
accounted for in VAB expression, the probability of sensing the food (p_d) is defined as follows:

$p_d = \begin{cases} \frac{1}{4} & \textit{for unilateral sensing} \\ \frac{1}{2} & \textit{for bilateral sensing} \end{cases}$ (equation 5; Supplementary file 1; Figure 6B and C).

When a diet is available in a given cell area and two fish exist in the same cell area, two
possibilities exist (Figure 6A): (1) in unilateral sensing, only one fish can sense and acquire the
diet ($\frac{E_i}{dt} > 0$), whereas the other fish does not evoke VAB, i.e., no calories are lost in the other
fish ($\frac{E_{i'}}{dt} = 0$) (Figure 6A and B); (2) in bilateral sensing, the two individuals compete for the diet
unit (which only supplies one fish), resulting in one fish capturing the diet and gaining a
positive E value ($\frac{E_{i'}}{dt} > 0$) and the other losing calories due to their foraging activity ($\frac{E_{i'}}{dt} < 0$),
which results in many fish losing energy in the population (Figure 6C). The total $\frac{E_i}{dt} + \frac{E_{i'}}{dt}$ in
unilateral sensing (1) is larger than that in bilateral sensing (2). This example shows that the
number of fish reacting to and capturing food will differ in bilateral- or unilateral sensing

situations, resulting in differences in E based on E_c in equation 2 (Supplementary file 1): E_c (calories consumed) depends on p_d (eqs. 2, 4, and 5; Supplementary file 1), so the E_c for bilateral sensing is twice as high as that for unilateral sensing (Figure 6C). As shown in Figure 6D and E, in a diet-rich environment, the bilateral sensing population has an advantage because its calorie gain is higher than that of the unilateral sensing population. Under food-sparse conditions, the advantage is reversed because the unilateral sensing population maintains $\frac{dE}{dt} > 0$, whereas the bilateral sensing population does not achieve $\frac{dE}{dt} > 0$ because many failed foraging attempts increase E_c (Figure 6D and F). In summary, the unilateral sensing population can survive better than the bilateral sensing population by reducing the number of failed foraging attempts in food-limited conditions. This model suggests that the sensory laterality of Pachón cavefish could be beneficial in food-limited environments.

The laterality in approaches (NOA) is defined as a preference for using the left or right-side of the head/body when approaching the vibration source. Our video observations indicated that fish exhibit multiple quick turns against the vibrating rod when they are in its vicinity (within a 1.3-cm radius of the rod). We assume that sharp turns cost more calories than typical swimming. We also assume that the fish that use unilateral approaches make steeper turns more often than the fish that use bilateral approaches (Figure 6G). The different levels of calorie consumption in fish using unilateral and bilateral approaches are implemented in the parameter 'a', the coefficient of E_c, i.e., the average calorie consumption per time unit (eq. 2 in Supplementary file 1). Accordingly, the parameter 'a' value of the fish that use unilateral approaches is higher than that of the fish that use bilateral approaches (hatched lines in Figure 6H and I; Supplementary file 1). With this setting, under food-rich conditions, the $\frac{dE}{dt}$ values of fish that use either unilateral or bilateral approaches are positive (Figure 6H); thus, both types of fish are expected to survive. In contrast, a population that uses bilateral approaches has a higher chance of survival ($\frac{dE}{dt} > 0$) than a population that uses unilateral approaches in a food-limited environment ($\frac{dE}{dt} < 0$; Figure 6I). Our starvation experiment showed that Tinaja cavefish shifted their approach from unilateral to L–R balanced (Figure 5A and 5C); therefore, these fish seem to adjust their L–R approach depending on food availability.

According to our model, Pachón cavefish is the most optimized population as it shows L–R symmetrical approaches (Figure 5A and 5C) as well as left-biased unilateral sensing (Figs. 1 and 2). The degree of laterality in Pachón and Tinaja cavefish populations fits well with their ecological conditions where Pachón cavefish experiences seasonal fluctuation of animated

diets, crustaceans, which seems to be available more in the ~6-month dry season than in the

6-month wet season (Espinasa et al., 2017; Mitchell et al., 1977). In contrast, several Tinaja

cave pools receive percolating water most of the time and have thick organic matter covering

the bottom, distinct from Pachón cave, and the plastic laterality may be sufficient for adaptation

to cave environment (Mitchell et al., 1977). There might be a possibility that Pachón cavefish

reside in a more diet-limited condition than Tinaja, yielding further diversified laterality to adapt

to cave environment.

In summary, laterality in sensory–behavior coupling and symmetrical approaches may have

advantages in food-sparse environments and may have promoted adaptation processes in

Pachón cave. Over the course of evolution, Tinaja cavefish evolved a plastic foraging strategy

according to food availability, whereas Pachón cavefish evolved a robust SN sensory laterality

with balanced approaches to vibrating objects. These results represent new insights that help

us better understand the evolution of laterality, and they may lead to further insights into the

diversification of laterality among animals.

Materials and Methods

Astyanax mexicanus husbandry and ethics statement

The lab-raised surface fish population used in this study is descended from an original collection from Balmorhea Springs State Park (Texas, USA), whereas the lab-raised cave populations used in this study are descendants of an original collection from Cueva de El Pachón in Tamaulipas collected by Richard Burowski (2009/2011), and collections from Sótano de la Tinaja and Cueva de Los Sabinos in San Luis Potosí, Mexico collected by Bill Jeffery in (2000 - 2001). The adult fish (1–3 years old) used in the experiments were raised in groups of 20–25 individuals in 0.74-L Ziploc containers (S.C. Johnson & Son Inc., Racine, WI, USA) and 1.4-L Zebrafish Aquatic Housing System tanks (Aquaneering Inc., San Diego, CA, USA). Fish were fed twice per day *ad libitum* with live *Artemia salina* larvae (Brine Shrimp Direct, Ogden, UT, USA).

All animal procedures were performed under protocol 17-2560 approved by the Institutional Animal Care and Use Committee at the University of Hawai'i.

Recording VAB

VAB assays were performed as described previously by Yoshizawa et al. (2010) and Worsham et al. (2019). Briefly, fish were acclimated for 4–5 days in a cylindrical chamber (10 cm in diameter; 5 cm in height; Pyrex 325 mL; Corning, NY, USA) filled with conditioned water (pH 6.8–7.0; conductivity of ~700 µS). Fish were fed once per day *ad libitum* with live *A. salina* larvae. After acclimation, the behavior of fish in response to vibrations was recorded for 3 min immediately after the introduction of the vibration rod into the bowl. The vibration stimulus was generated using a 7.5-mm diameter glass rod at 40 Hz with a GW Instek SFG-1003 DDS Function Generator (Good Will Instrument Co., Ltd., New Taipei City, Taiwan) connected to an audio speaker (ProSpeakers, Apple, Cupertino, CA, USA). Video footage of behavior was recorded under infrared illumination (880-nm wavelength; BL41192-880 blacklight; Advanced Illumination, Rochester, VT, USA) using a customized Microsoft MAIN-31891 LifeCam Studio installed with a Zoom lens (Zoom 7000, Navitar, Rochester, NY) and VirtualDub2 video capture software (licensed under the GNU General Public License: https://sourceforge.net/projects/vdfiltermod/).

The NOA was analyzed using ImageJ macro plugins (ImageJ 1.53a; NIH, Bethesda, MD, USA) and a homemade script published previously (Worsham et al., 2019; Yoshizawa et al.,

2010). Using this method, we measured NOA and selected Pachón cavefish that exhibited

NOA ≥ 6 for the ablation study (see "Ablation experiments" below).

VAB video analysis using DLC

Videos were analyzed using the DLC toolbox version 2.0.6 (Mathis et al., 2018; Nath et al.,

2019) in Python 3.6. DLC provides a robust suite of tools for deep learning-based markerless

pose estimation. The approach is based on transfer learning, wherein a deep neural network is

initially trained on an initial task with large amounts of available training data to achieve robust

initial parameterization and then fine-tuned on a target task (ideally as closely related to the

initial task as possible; in this case, pose estimation) using a smaller dataset. DLC uses a

ResNet50 (He et al., 2016) backbone pretrained on the ImageNet (Deng et al., 2009) dataset

to classify 1000 object categories in a dataset of over one million hand-labeled images. This

network parameterization is then transferred and fine-tuned, during which the last layer of the

network used originally for classification is replaced with transposed convolutional layers for

pose estimation. These new layers predict individual probability densities over all pixels of a

given input image for each pose marker, and such predictions are repeated independently for

each frame in the video. DLC was also used to simplify data selection, data labeling, training,

evaluation, and retraining processes. In this study, we specified a pose estimation task to track

both the fish and rods in the VAB videos. For the fish, three head markers were tracked (right-

side, left-side, and center) as well as one marker in the caudal fin; additionally, one marker

was placed in the center of the vibrating rod. Because some videos contained two Petri dishes

and some contained one, two separate sets of the same markers were allocated. The initial

training set included frames selected randomly from 88 different VAB videos, and 807 hand-

labeled training frames (images) were used to train our final model. The ResNet50 model was

fine-tuned until the DLC early-stopping criterion was reached while using the default DLC

training parameters as described by Mathis et al. (2018). All weights in the network were

adjusted during training. The trained model was used to predict marker locations in all videos,

outputting the x and y coordinates associated with the highest probability pixel for each

marker. In frames with at least one DLC marker below a predefined confidence threshold, all

marker locations were interpolated based on the markers' positions in the preceding and

following frames, which ensured that fish tracking was smooth throughout the duration of each

video. Specifically, for a sequence of one or more subsequent frames where each frame had

at least one marker below the confidence threshold, all markers' locations were linearly

interpolated between each of the respective markers' locations in the frame immediately

preceding and the frame immediately following that sequence. For videos including multiple

recording arenas, markers for each Petri dish were treated independently. The confidence

threshold was set empirically to 0.92 after manual inspection of the DLC outputs. The resulting

marker locations were used to track several behaviors of interest in the videos: time spent

within a 1.3-cm radius of the rod (DIR), time spent being outside of this radius (DOR), left- and

right-sided approaches (NOA), and total swimming distance. The 1.3-cm radius from the

vibrating rod was determined as the cutoff of VAB because it was the distance that best

discriminated between the surface fish and cavefish approaches across the tested radii of 0.5–

2.5 cm (in 0.1-cm increments) from the vibrating rod. A left-sided approach was defined as

entering the 1.3-cm radius threshold with the head-center marker while the head-left marker

was closer than the head-right marker to the rod, and vice versa for a right-sided approach

(Figure 3A). Accordingly, DIR, DOR, and NOA were separated as right- or left-facing based on

the proximity of the head-right and head-left markers to the rod (Figure 3A).

Repeatablity test

Repeatablity test was performed as described previously (Yoshizawa et al., 2010). Briefly, we

first acclimated 22 surface, 9 Pachón cave and 11 Tinaja cave fish in cylindrical chambers for

4–5 days. Fish were fed once per day *ad libitum* with live *A. salina* larvae and recorded their

movement as described in the Recording VAB section. After the recording, the cylindrical

chambers were cleaned and fish were maintained in the same chambers for 6 days, and

rerecorded in 7[th] days of the second acclimation (6 days a part). This procedure was repeated

for the 3[rd] recording. The conditioned water was replaced every two days during this period.

After the behavior analysis using DLC, interclass correlation repeatability (3 repeated

measurements) was tested as referring Single_random_raters by the 'psych' R package.

SN imaging

Neuromast vital staining was performed as described previously (Jørgenson and Jørgensen,

1989; Worsham et al., 2019; Yoshizawa et al., 2010). Fish were immersed in 25 µg/mL of 4-(4-

(dimethylaminostyryl)−1-methylpyridiniumiodide (4-Di-1-ASP; MilliporeSigma, Burlington, MA)

dissolved in conditioned water for 40–60 min, after which they were anesthetized in 66.7

µg/mL of ice-cold ethyl 3-aminobenzoate methanesulfonate salt (MS222; MilliporeSigma) in

conditioned water (pH 6.8–7.0; conductivity of ~700 µS). Fish were then visualized using a

fluorescence microscope [BX61WI Olympus microscope with a 2.5× MPlanFL N lens, a Rhodamine filter set, and an ORCA-Flash 4.0 digital camera (Olympus Corp.)]. SNs were counted from digital images of 4-Di-1-ASP-stained fish using the "Analyze Particles" function of ImageJ 1.53a. SNs were counted both in the left and right sides of the third infraorbital/suborbital bone.

Ablation experiments

Ablation experiments were performed using 1-year old surface fish (n = 47) and VAB-positive cavefish (Pachón population, n = 67). The VAB level was measured before ablation, and VAB-positive cavefish (NOA ≥ 6 according to ImageJ-based analysis) were selected strategically to determine the association between IO3 SNs and VAB level.

SN ablations were performed using Vetbond (3M, St. Paul, MN, USA), a nontoxic tissue adhesive, on the left (surface fish, n = 14; Pachón cavefish, n = 30), right (surface fish, n = 13; Pachón cavefish, n = 25), and both left and right (surface fish, n = 10; Pachón cavefish, n = 12) sides (Figure 2A and B). Fish were anesthetized in 66.7 µg/mL of ice-cold ethyl 3-aminobenzoate methanesulfonate salt (MS222, MilliporeSigma) in conditioned water (pH 6.8–7.0; conductivity of ~700 µS). Excess water was removed carefully using Kimtech wipes, and the adhesive (0.5 µL) was applied to the skin surface. After being exposed to the air for 10 s, treated fish were placed in bowls (10 cm in diameter; 5 cm in height; Pyrex 325 mL, Corning) filled with conditioned water at room temperature. Within a day of active swimming, the adhesive usually peeled off, and VAB was assayed. 4-Di-1-ASP staining was performed to count the number of SNs in the IO3 region before and after ablation.

Fasting

Pachón cavefish (n = 20), Tinaja cavefish (n = 20) and surface fish (n = 16) individuals (~1 year old) were placed in cylindrical chambers (10 cm in diameter; 5 cm in height; Pyrex 325 mL, Corning) and acclimated for 4–5 days in conditioned water (pH 6.8–7.0; conductivity of ~600 µS) in which they were fed once a day (weight or approximate number of *Artemia*) with live *A. salina* larvae. After acclimation, VAB was recorded and analyzed. Following this first assessment, fish were starved for 6 days and VAB was recorded and analyzed once again.

Statistical analysis

All statistical analyses were performed in RStudio 4.0.3 (RStudio, Boston, MA, USA). The R

packages used included lme4, lmerTest, car, coin, yarrr, AICcmodavg, and ggpubr. Linear or

generalized linear models were selected using Akaike's information criterion function to identify

the best fit models for analysis of the NOA and DIR in all experiments. Analysis of variance

(ANOVA) was performed using generalized linear model fitting functions (glm or glmer in the

lme4 package). Post-hoc tests were performed using the Wilcoxon signed-rank (paired) test

followed by Holm's multiple-test correction. Nonparametric Kendall's t correlation was used to

evaluate associations between neuromasts and behavior outputs. In the field experiments, the

total number of turns within the circle were pooled for each cave pool, and the left-side turns

vs. the right-side turns were tested by fitting a generalized linear mixed-effects model using a

binomial distribution.

References

Aspiras AC, Rohner N, Martineau B, Borowsky RL, Tabin CJ. 2015. Melanocortin 4 receptor mutations contribute to the adaptation of cavefish to nutrient-poor conditions. *Proc Natl Acad Sci U S A* **112**:9668–9673. doi:10.1073/pnas.1510802112

Bisazza A, Cantalupo C, Capocchiano M, Vallortigara G. 2000. Population lateralisation and social behaviour: A study with 16 species of fish. *Laterality Asymmetries Body, Brain Cogn* **5**:269–284. doi:10.1080/713754381

Bisazza A, Dadda M. 2005. Enhanced schooling performance in lateralized fishes. *Proc R Soc B Biol Sci* **272**:1677–1681. doi:10.1098/rspb.2005.3145

Borowsky R. 2008. Restoring sight in blind cavefish. *Curr Biol* **18**:R23–R24.

Braun CB, Sand O. 2013. Functional Overlap and Nonoverlap Between Lateral Line and Auditory Systems In: Coombs S, Bleckmann H, Fay RR, Popper AN, editors. The Lateral Line System. Springer Handbook of Auditory Research 48. New York: Springer New York. pp. 281–312. doi:10.1007/2506_2013_19

Chivers DP, McCormick MI, Allan BJM, Mitchell MD, Gonçalves EJ, Bryshun R, Ferrari MCO. 2016. At odds with the group: changes in lateralization and escape performance reveal conformity and conflict in fish schools. *Proceedings Biol Sci* **283**:20161127. doi:10.1098/rspb.2016.1127

Chivers DP, McCormick MI, Warren DT, Allan BJM, Ramasamy RA, Arvizu BK, Glue M, Ferrari MCO. 2017. Competitive superiority versus predation savvy: the two sides of behavioural lateralization. *Anim Behav* **130**:9–15. doi:10.1016/J.ANBEHAV.2017.05.006

Culver DC. 1982. Cave life, evolution and ecology. Cambridge: Harvard University Press.

Culver DC, Pipan T. 2009. The biology of caves and other subterranean habitats, The biology of habitats series. Oxford: Oxford University Press.

Dadda M, Koolhaas WH, Domenici P. 2010. Behavioural asymmetry affects escape performance in a teleost fish. *Biol Lett* **6**:414–417. doi:10.1098/rsbl.2009.0904

de Perera TB, Braithwaite VAA, Burt de Perera T, Braithwaite VAA. 2005. Laterality in a non-visual sensory modality - the lateral line of fish. *Curr Biol* **15**:R241–R242. doi:10.1016/j.cub.2005.03.035

Deng J, Dong W, Socher R, Li L-J, Li K, Fei-Fei L. 2009. Imagenet: A large-scale hierarchical image database2009 IEEE Conference on Computer Vision and Pattern Recognition. pp. 248–255.

Espinasa L, Bonaroti N, Wong J, Pottin K, Queinnec E, Rétaux S. 2017. Contrasting feeding habits of post-larval and adult *Astyanax* cavefish. *Subterr Biol* **21**:1–17. doi:10.3897/subtbiol.21.11046

Espinasa L, Heintz C, Rétaux S, Yoshisawa M, Agnès F, Ornelas-Garcia P, Balogh-Robinson R. 2021. Vibration attraction response is a plastic trait in blind Mexican tetra (*Astyanax mexicanus*), variable within subpopulations inhabiting the same cave. *J Fish Biol* **98**:304–316. doi:10.1111/jfb.14586

Espinasa L, Ornelas-García CP, Legendre L, Rétaux S, Best A, Gamboa-Miranda R, Espinosa-Pérez H, Sprouse P. 2020. Discovery of Two New Astyanax Cavefish Localities Leads to Further Understanding of the Species Biogeography. *Diversity* **12**:368. doi:10.3390/d12100368

Fernandes VFL, Macaspac C, Lu L, Yoshizawa M. 2018. Evolution of the developmental plasticity and a coupling between left mechanosensory neuromasts and an adaptive foraging behavior. *Dev Biol* **441**:262–271. doi:10.1016/j.ydbio.2018.05.012

Fumey J, Hinaux H, Noirot C, Thermes C, Rétaux S, Casane D. 2018. Evidence for late Pleistocene origin of Astyanax mexicanus cavefish. *BMC Evol Biol* **18**:43.

doi:10.1186/s12862-018-1156-7

Gross JB, Gangidine A, Powers AK. 2016. Asymmetric facial bone fragmentation mirrors
asymmetric distribution of cranial neuromasts in blind Mexican cavefish. *Symmetry (Basel)*
8. doi:10.3390/sym8110118

Gross JB, Powers AK. 2016. The Evolution of the Cavefish Craniofacial Complex In: Keene
AC, Yoshizawa M, McGaugh SE, editors. Biology and Evolution of the Mexican Cavefish.
Amsterdam: Academic Press. pp. 193–207. doi:10.1016/B978-0-12-802148-4.00010-4

Güntürkün O, Ströckens F, Ocklenburg S. 2020. Brain Lateralization: A Comparative
Perspective. *Physiol Rev* **100**:1019–1063. doi:10.1152/physrev.00006.2019

He K, Zhang X, Ren S, Sun J. 2016. Deep Residual Learning for Image
RecognitionProceedings of the IEEE Conference on Computer Vision and Pattern
Recognition (CVPR).

Heilman KM, Abell TVD. 1980. Right hemisphere dominance for attention: The mechanism
underlying hemispheric asymmetries of inattention (neglect). *Neurology* **30**:327–327.
doi:10.1212/WNL.30.3.327

Herman A, Brandvain Y, Weagley J, Jeffery WR, Keene AC, Kono TJY, Bilandžija H,
Borowsky R, Espinasa L, O'Quin K, Ornelas-García CP, Yoshizawa M, Carlson B,
Maldonado E, Gross JB, Cartwright RA, Rohner N, Warren WC, McGaugh SE. 2018. The
role of gene flow in rapid and repeated evolution of cave-related traits in Mexican tetra,
Astyanax mexicanus. *Mol Ecol* **27**:4397–4416. doi:10.1111/mec.14877

Jørgenson JM, Jørgensen JM. 1989. Evolution of octavolateralis sensory cells In: Coombs S,
Görner P, Münz H, editors. The Mechanosensory Lateral Line. New York: Springer-
Verlag. pp. 115–145.

Kalmijn AJ. 1988. Hydrodynamic and Acoustic Field DetectionSensory Biology of Aquatic
Animals. New York, NY: Springer New York. pp. 83–130. doi:10.1007/978-1-4612-3714-
3_4

Keene AC, Yoshizawa M, McGaugh SE. 2016. Biology and Evolution of the Mexican Cavefish,
Biology and Evolution of the Mexican Cavefish. Amsterdam: Elsevier Inc.
doi:10.1016/C2014-0-01426-8

Kohashi T, Nakata N, Oda Y. 2012. Effective sensory modality activating an escape triggering
neuron switches during early development in zebrafish. *J Neurosci* **32**:5810–5820.
doi:10.1523/JNEUROSCI.6169-11.2012

Landis JR, Koch GG. 1977. The measurement of observer agreement for categorical data.
Biometrics **33**:159–74.

Lang HH. 1980. Surface wave discrimination between prey and nonprey by the back
swimmerNotonecta glauca L. (Hemiptera, Heteroptera). *Behav Ecol Sociobiol* **6**:233–246.
doi:10.1007/BF00569205

Magat M, Brown C. 2009. Laterality enhances cognition in Australian parrots. *Proc R Soc B*
Biol Sci **276**:4155–4162. doi:10.1098/rspb.2009.1397

Mathis A, Mamidanna P, Cury KM, Abe T, Murthy VN, Mathis MW, Bethge M. 2018.
DeepLabCut: markerless pose estimation of user-defined body parts with deep learning.
Nat Neurosci **21**:1281–1289. doi:10.1038/s41593-018-0209-y

Mitchell RW, Russell WH, Elliott WR. 1977. Mexican eyeless characin fishes, genus Astyanax:
Environment, distribution, and evolution., Special pu. ed, Special publications the museum
Texas Tech University. Texas: Texas Tech Press.

Montgomery JC, Macdonald JA. 1987. Sensory Tuning of Lateral Line Receptors in Antarctic
Fish to the Movements of Planktonic Prey Published by : American Association for the
Advancement of Science Stable URL : https://www.jstor.org/stable/1698962 **235**:195–196.

Nath T, Mathis A, Chen AC, Patel A, Bethge M, Mathis MW. 2019. Using {DeepLabCut} for

{3D} markerless pose estimation across species and behaviors. *Nat Protoc* **14**:2152–
2176. doi:10.1038/s41596-019-0176-0
Neumann S, Doubell TP, Leslie T, Woolf CJ. 1996. Inflammatory pain hypersensitivity
mediated by phenotypic switch in myelinated primary sensory neurons. *Nature* **384**:360–
364. doi:10.1038/384360a0
Nottebohm F. 1977. Asymmetries in neural control of vocalization in the canary In"
Lateralization of the Nervous System"(S. Harnard, ed.).
Nottebohm F. 1971. Neural lateralization of vocal control in a passerine bird. I. Song. *J Exp*
Zool **177**:229–261. doi:10.1002/jez.1401770210
Nottebohm F, Nottebohm ME. 1976. Left hypoglossal dominance in the control of canary and
white-crowned sparrow song. *J Comp Physiol ■ A* **108**:171–192. doi:10.1007/BF02169047
Ornelas-García C, Domínguez-Domínguez O, Doadrio I. 2008. Evolutionary history of the fish
genus Astyanax Baird & Girard (1854) (Actinopterygii, Characidae) in Mesoamerica
reveals multiple morphological homoplasies. *BMC Evol Biol* **8**:340. doi:10.1186/1471-
2148-8-340
Popper AN, Fay RR. 2011. Rethinking sound detection by fishes. *Hear Res* **273**:25–36.
doi:10.1016/j.heares.2009.12.023
Powers AK, Davis EM, Kaplan SA, Gross JB. 2017. Cranial asymmetry arises later in the life
history of the blind Mexican cavefish, Astyanax mexicanus. *PLoS One* **12**:e0177419.
doi:10.1371/journal.pone.0177419
Protas M, Jeffery WR. 2012. Evolution and development in cave animals: from fish to
crustaceans. *Wiley Interdiscip Rev Dev Biol* **1**:823–845. doi:10.1002/wdev.61
Rétaux S, Casane D, Rétaux S, Casane D. 2013. Evolution of eye development in the
darkness of caves: adaptation, drift, or both? *Evodevo* **4**:26.
Rogers LJ. 1989. eScholarship International Journal of Comparative Psychology Title Laterality
in Animals Permalink Publication Date. *Int J Comp Psychol* **3**.
Rogers LJ. 1981. Environmental influences on brain lateralization. *Behav Brain Sci* **4**:35–36.
Rogers LJ, Workman L. 1989. Light exposure during incubation affects competitive behaviour
in domestic chicks. *Appl Anim Behav Sci* **23**:187–198. doi:10.1016/0168-1591(89)90109-3
Schwartz G, Davidson R, Maer F. 1975. Right hemisphere lateralization for emotion in the
human brain: interactions with cognition. *Science (80-)* **190**:286–288.
doi:10.1126/science.1179210
Sovrano VA, Dadda M, Bisazza A. 2005. Lateralized fish perform better than nonlateralized
fish in spatial reorientation tasks. *Behav Brain Res* **163**:122–7.
doi:10.1016/j.bbr.2005.04.012
Strecker U, Hausdorf B, Wilkens H. 2012. Parallel speciation in Astyanax cave fish (Teleostei)
in Northern Mexico. *Mol Phylogenet Evol* **62**:62–70.
Tommasi L. 2009. Mechanisms and functions of brain and behavioural asymmetries. *Philos*
Trans R Soc Lond B Biol Sci **364**:855–9. doi:10.1098/rstb.2008.0293
Tricas TCTC. 2020. Acoustic Ecology, Communication and Peripheral Signal Processing in
Fishes. In: Fritzsch B, Grothe B, editors. The Senses: A Comprehensive Reference, Vol.
2. Cambridge: Academic Press. pp. 114–137. doi:10.1016/B978-0-12-809324-5.24220-X
Wang H, Dai Y, Fukuoka T, Yamanaka H, Obata K, Tokunaga A, Noguchi K. 2004.
Enhancement of stimulation-induced ERK activation in the spinal dorsal horn and gracile
nucleus neurons in rats with peripheral nerve injury. *Eur J Neurosci* **19**:884–890.
doi:10.1111/j.0953-816X.2004.03203.x
Wernicke C. 1874. Der aphasische Symptomencomplex: eine psychologische Studie auf
anatomischer Basis. Cohn.
Wilkens H, Hecht MK, Wallace B, Wilkens H. 1988. Evolution and genetics of epigean and

cave Astyanax-fasciatus (Characidae, Pisces) - Support for the neutral mutation theory In:
Hecht M K, Wallace B, editors. Evolutionary Biology. New York: Plenum Publishing
Corporation. pp. 271–367. doi:10.1007/978-1-4613-1043-3_8
Wilson EV V, Thomas RQQ, Evans LMM. 2004. Macrobrachium as a possible determinant of
Astyanax fasciatus distribution in a neotropical lowland stream. *Dartmouth Stud Trop Ecol*
2004:99–102.
Worsham M, Lima Fernandes VF, Settle A, Balaan C, Lactaoen K, Tuttle LJ, Iwashita M,
Yoshizawa M. 2019. Behavioral tracking and neuromast imaging of Mexican cavefish. *J*
Vis Exp **2019**. doi:10.3791/59099
Yoshizawa M. 2015. Behaviors of cavefish offer insight into developmental evolution. *Mol*
Reprod Dev **82**:268–280. doi:10.1002/mrd.22471
Yoshizawa M, Ashida G, Jeffery WR. 2012a. Parental genetic effects in a cavefish adaptive
behavior explain disparity between nuclear and mitochondrial DNA. *Evolution (N Y)*
66:2975–2982. doi:10.1111/j.1558-5646.2012.01651.x
Yoshizawa M, Gorički S, Soares D, Jeffery WR. 2010. Evolution of a behavioral shift mediated
by superficial neuromasts helps cavefish find food in darkness. *Curr Biol* **20**:1631–1636.
doi:10.1016/j.cub.2010.07.017
Yoshizawa M, Jeffery WR. 2011. Evolutionary tuning of an adaptive behavior requires
enhancement of the neuromast sensory system. *Commun Integr Biol* **4**:89–91.
doi:10.1016/j.cub.2010.0
Yoshizawa M, Robinson BG, Duboué ER, Masek P, Jaggard JB, O'Quin KE, Borowsky RL,
Jeffery WR, Keene AC. 2015. Distinct genetic architecture underlies the emergence of
sleep loss and prey-seeking behavior in the Mexican cavefish. *BMC Biol* **13**:15.
doi:10.1186/s12915-015-0119-3
Yoshizawa M, Yamamoto Y, O'Quin KE, Jeffery WR. 2012b. Evolution of an adaptive behavior
and its sensory receptors promotes eye regression in blind cavefish. *BMC Biol* **10**:108.
doi:10.1186/1741-7007-11-82
Zegers M, de Bruijne MC, Wagner C, Groenewegen PP, van der Wal G, de Vet HCW. 2010.
The inter-rater agreement of retrospective assessments of adverse events does not
improve with two reviewers per patient record. *J Clin Epidemiol* **63**:94–102.
doi:10.1016/j.jclinepi.2009.03.004

Figures

Figure 1. Sensory laterality in surface fish and cavefish populations of *Astyanax mexicanus*. VAB videos were analyzed using a home-developed Python script and a DeepLabCut deep learning algorithm. This method allowed us to detect the number of times fish made left or right approaches toward the vibrating rod (NOA) and the associated duration they spent within a 1.3-cm radius from the vibrating rod (DIR). (**A**) Video image (left) and schematic representation (right) of fish and the vibrating rod labeled with three markers on the head (left, center, and right) and one marker on the caudal fin. (**B**) Superficial neuromasts (SNs) in the infraorbital region (IO3 region delimited by a yellow line) were stained with 4-Di-1-ASP vital dye. (**C**) NOA data for the lab-raised populations indicated that Pachón cavefish showed the highest NOA values. The bars in the pirate plots represent means ± standard errors of the means (each population, n = 20). *, $P < 0.05$; **, $P < 0.01$; and ***, $P < 0.001$. Statistical scores are shown in Supplementary Table 1. (**D–G**) The number of SNs in the left or right IO3 area plotted against the total NOA in surface fish (**D**) and Los Sabinos (**E**), Tinaja (**F**), and Pachón cavefish (**G**). Statistical scores are available in Supplementary Table 2.

Figure 2. Ablation of superficial neuromasts (SNs) confirmed that the left-side SN promotes the level of VAB in Pachón cavefish. (**A**) Left and right sides of SNs in Pachón cavefish stained with 4-Di-1-ASP (green dots) before and after ablation of SNs in the infraorbital region (IO3 shown by a yellow line) of the same fish. (**B**) Proportion of ablated SNs on the left-side and right-side plotted against the total NOA in both cases. Regression lines with 95% confidence intervals (shaded gray or orange) are shown in each panel. See Supplementary Table 3 for additional details. **, P < 0.01.

807
Figure 2—figure supplement 1. Ablation of superficial neuromasts (SNs) did not induce
detectable effect on the level of VAB in surface fish. (**A**) Left and right sides of SNs in
Pachón cavefish stained with 4-Di-1-ASP (green dots) before and after ablation of SNs in the
infraorbital region (IO3 shown by a yellow line) of the same fish. (**B**) Proportion of ablated SNs
on the left and right sides plotted against the total NOA in both cases. Regression lines with
95% confidence intervals (shaded gray or orange) are shown in each panel. See
Supplementary Table 3 for additional details.
815

Figure 2--figure supplement 2

 Figure 2—figure supplement 2. Overall results of the SN ablation experiments in Pachón
cavefish. (**A**) Number of approaches (NOA) in Pachón cavefish. Ablation of the left SN
resulted in an increase in NOA (V = 74.5, P = 0.032); however, ablation of the right SN
changed NOA at an underdetection level (V = 121, P = 0.063). (**B**) Duratin in the radius (DIR)
in Pachón cavefish. Left SN or right SN ablation did not show detectable changes in DIR.
Pirate plot bars represent means ± standard errors of the means. Data were analyzed using a
generalized linear model followed by a post-hoc test (Wilcoxon test, adjusted by Holm's
correction): *, P < 0.05; **, P < 0.01; and ***, P < 0.001. Statistical scores are available in
Supplementary Table 4.

Figure 3

827
Figure 3. Laterality in behavioral output among surface fish and cavefish populations of
Astyanax mexicanus. (**A**) Schemes representing left or right approaches expressed as NOA
(top) and left or right adherence expressed as DIR (bottom). (**B**) Histograms for the L–R NOA
ratio (the right-side NOA divided by the total NOA: $\frac{NOA(right)}{NOA(right)+NOA(left)}$) across all four
populations. (**C**) Histograms for the L–R DIR ratio ($\frac{DIR(right)}{DIR(right)+DIR(left)}$) across all four
populations. Ratios of <0.33 indicate a strong left preference, ratios of 0.33–0.67 indicate
balanced approaches, and ratios of >0.67 indicate a strong right preference. Los Sabinos
cavefish showed a gaussian-like distribution, whereas many surface fish and the other
cavefish individulas exhibited strong left or right bias. Notably, all four populations included a
few individuals that showed strong left or right preferences (near 0.0 or 1.0, respectively).
838
839

Figure 3--figure supplement 1

840
Figure 3—figure supplement 1. *Astyanax mexicanus* populations did not exhibit left- or
right-biased approaches (NOA) or adhesion (DIR) at the population level.

(**A, B**) Pirate plots showing no detectable differences between the number of left- and right-
side approaches (NOA) (**A**) or the duration in the radius (DIR) (**B**) among the four tested
populations (2–3-year-old individuals). Data represent means ± standard errors of the means
(each population, n = 20). Further details are presented in Supplementary Table 1.
847

Figure 4. Plasticities of surface fish and Tinaja cavefish in total NOA and/or DIR, and the plastic left SN–NOA association in Pachón cavefish in response to fasting. (**A**) Pirate plots showing total NOA. Surface fish showed significantly reduced NOA after fasting, but neither Tinaja nor Pachón cavefish showed significant changes in NOA. (**B**) Pirate plots showing total DIR. Surface fish showed significantly reduced DIR after fasting, but Tinaja's DIR increased after fasting. Pachón cavefish showed no significant changes in DIR. (**C**) The number of left or right SNs plotted against changes in NOA after fasting (NOA_{after}–NOA_{before}) in surface fish, Tinaja cavefish, and Pachón cavefish. In Pachón cavefish, individuals with fewer left SNs showed a higher increase in total NOA, suggesting that a plastic response to fasting existed in individuals with few left SNs and a low NOA level. There was no detectable correlation between the number of right SNs and the change in NOA in Pachón cavefish or the other populations. All statistical scores are available in Supplementary Table 5 and 7.

Figure 5. Right-biased approaches (NOA) in Tinaja cavefish were reduced after fasting.
(**A, B**) Histograms and density plots showing the L–R NOA ratio (**A**) and L–R DIR ratio (**B**)
before and after fasting in the indicated populations. In **A**, more Tinaja cavefish individuals
showed L–R balanced approaches (0.33–0.67) after fasting; Pachón cavefish maintained L–R
balanced approaches (0.33–0.67) before and after fasting, whereas surface fish showed a
slight right bias. L–R NOA ratio = $\frac{NOA(right)}{NOA(right)+NOA(left)}$. In **B**, the trends for DIR were similar to
those in (A). L–R DIR ratio = $\frac{DIR(right)}{DIR(right)+DIR(left)}$. (**C, D**) Pirate plots showing L–R bias index for
NOA (**C**) and DIR (**D**). In **C**, Right bias index = $\frac{\max\{NOA(right),NOA(left)\}}{NOA(right)+NOA(left)}$. Tinaja cavefish showed a
significantly reduced bias index for NOA after fasting, indicating balanced L–R approaches,
whereas the other populations showed no detectable changes. In **D**, L–R bias index =

873 $\frac{\max\{DIR(right),DIR(left)\}}{DIR(right)+DIR(left)}$. None of the populations showed detectable changes in the bias index
874 after fasting, although Tinaja cavefish showed a nonsignificant trend for reduced bias after
875 fasting. All statistical scores are available in Supplementary Table 6. **, P < 0.01.
876

877
Figure 5—figure supplement 1. In response to starvation, lateralities in NOA and DIR are
plastic in Tinaja cavefish but not in Pachón cavefish. (**A, B**) Pirate plots showing the
number of left- and right-side approaches (NOA) (**A**) and left- and right-side-associated
durations within a 1.3-cm radius (DIR) (**B**) before and after a 6-day fasting period. In **A**, only
surface fish showed a significantly reduced NOA after fasting among the three tested
populations, and no L–R bias (laterality) was detectable. In **B**, surface fish showed significantly
reduced DIR after fasting, whereas Tinaja cavefish showed increased DIR after fasting. Tinaja
cavefish also showed right bias in DIR before and after fasting, but they showed more
significant right bias than left bias in DIR before fasting. Pachón cavefish did not exhibit any
changes in DIR or L–R bias in DIR. All statistical scores are available in Supplementary Table
5. *, P < 0.05; **, P < 0.01.

889
Figure 6. Relevance of sensory and behavior laterality. (**A**) General schematic of the model
showing food availability and the number of fish. Each cell represents the sensing range of a
fish. Red dots represent diet and black shapes represent fish. (**B**) In a given cell, the food
detection probability p_d is 1/4 for lateral sensing (top) and 1/2 for bilateral sensing (bottom); red
and green shaded areas represent overlapping areas in the food and sensing fields,
respectively. (**C**) Energy consumption (E_c) as a function of the number of food-acquiring fish
(N_c). (**D**) Energy gain from food (E_g) as a function of the number of food-acquiring fish (N_c). In
C and **D**, red arrows and lines indicate lateral sensing fish and green arrows and lines indicate

bilateral sensing fish. Gray and black lines are the functions E_g and depend on food-rich and -

limited conditions, respectively. (**E, F**) Evaluation of $\frac{dE}{dt}$. **E** shows E_g and E_c under food-rich

conditions, whereas **F** shows E_g and E_c under food-limited conditions. $\frac{dE}{dt}$ is based on the

subtraction of E_c from E_g and is shown as solid vertical lines (green and red). In **E** (food-rich

condition), the bilateral population is optimized because $\frac{dE}{dt}$ of the bilateral sensing population is

larger than that of the unilateral sensing population. Furthermore, there is no unconsumed food

in the bilateral sensing population, whereas unconsumed food is present in the unilateral

sensing population (red dotted line). In **F** (food-limited condition), the unilateral population is

optimized because $\frac{dE}{dt}$ of the unilateral sensing population has a positive value, whereas $\frac{dE}{dt}$ in

the bilateral sensing population is negative. (**G**) Food-approaching behavior. The gray circle

shows the 1.3-cm radius from the food (red circle). Blue and pink arrows represent steeper and

less steep turns, respectively. Fish with balanced approaches take either left or right

approaches, whereas fish with biased approaches (laterality) always take one-sided

approaches that frequently require steeper turns. (**H, I**) Evaluation of $\frac{dE}{dt}$ in food-rich (**H**) and

food-limited (**I**) environments. The E_c values for unilateral (steeper turn) fish and balanced fish

are shown as blue and pink dotted lines, respectively. $\frac{dE}{dt}$ is based on the subtraction of E_c

(dotted lines) from E_g (gray solid line in H or black in I); therefore, it is expressed as the blue

and pink vertical solid lines. Under food-rich conditions, $\frac{dE}{dt}$ for both unilateral and balanced fish

is positive; thus, both populations can gain energy. In contrast, under food-limited conditions,

the L–R balanced population is optimal because it shows a positive value for $\frac{dE}{dt}$. See

Supplementary file 1 for further information.