1

Dimerization of iLID Optogenetic Proteins Observed Using 3D Single-Molecule Tracking

in Live Bacterial Cells

Alecia M. Achimovich[a], Ting Yan[b] and Andreas Gahlmann[a,b#]

[a]Department of Molecular Physiology & Biological Physics, University of
Virginia School of Medicine, Charlottesville, Virginia, USA

[b]Department of Chemistry, University of Virginia, Charlottesville, Virginia

#Address correspondence to Andreas Gahlmann, agahlmann@virginia.edu

ABSTRACT

3D single molecule tracking microscopy has enabled measurements of protein diffusion in living cells, offering information about protein dynamics and the cellular environment. For example, different diffusive states can be resolved and assigned to protein complexes of different size and composition. However, substantial statistical power and biological validation, often through genetic deletion of binding partners, are required to support diffusive state assignments. When investigating some cellular processes, transient perturbation to protein spatial distributions is preferable to permanent genetic deletion of an essential protein. Optogenetic dimerization systems can be used to manipulate protein spatial distributions which could offer a means to deplete specific diffusive states observed in single molecule tracking experiments. Here, we evaluate the performance of the iLID optogenetic system in living *E. coli* cells using diffraction-limited microscopy and 3D single molecule tracking. We observed a robust optogenetic response in protein spatial distribution after 488 nm laser activation. Surprisingly, 3D single molecule tracking results indicate activation of the optogenetic response at high intensity wavelengths for which there is evidence of minimal photon absorbance by the LOV2 domain. However, the preactivation response was minimized through the use of iLID system mutants, and titration of protein expression levels.

INTRODUCTION

Many fluorescence microscopy-based techniques have been developed over the past decades to detect and quantify interactions between biomolecules *in vitro* and *in vivo*. Improvements of fluorescent probes and biomolecular labeling technologies in conjunction with instrumental improvements have enabled measurements that provide critical insights into cellular organization and the biochemical interactions occurring within them.[1] Spatial co-localization of emitters through multi-color imaging has been widely utilized to gauge whether biomolecules are close in space and are thus able to interact. The power of such measurements depends critically on the achievable spatial resolution of the instrument used. Diffraction-limited imaging provides 200-300 nm resolution, which is orders of magnitude larger than the size of a typical protein (~2 nm) or the size of small protein complexes (~20 nm). Diffraction-limited resolution is thus too low to conclusively determine whether two proteins interact directly or whether their interaction is mediated by a third protein[2]. Super-resolution microscopy approaches such as PALM/STORM and MINFLUX, have been successful in addressing this challenge by enabling precise single-molecule localization with tens of nanometers of precision. For example, Symborska et al. determined radial positions of protein subunits of the nuclear pore complex (NPC) with subnanometer precision using PALM/STORM in conjunction with particle averaging[3]. This approach was later extended to achieve a 3D reconstruction of the nuclear pore complex through iterative multi-color imaging of each NPC subunit relative to a reference protein also within the NPC.[4] More recently, Ries and co-workers used MINFLUX microscopy to pinpoint the position of subunits within the NPC with single nanometer precision without the need for radial averaging[5]. Thus, for relatively large, immobile, and highly symmetric structures like the nuclear pore

complex, super-resolution fluorescence microscopy can provide accurate co-localization data that

provides information on how individual subunits interact with each other.[6-8]

Detecting protein interactions within freely-diffusing complexes cannot be accomplished

by fluorescence co-localization and thus requires different approaches. Live cell measurements of

protein oligomerization is possible using fluorescence microscopy techniques such as

Fluorescence recovery after photobleaching (FRAP), and Fluorescence correlation spectroscopy

(FCS)[9-11]. These methods have gained popularity due to the increased availability of commercial

confocal laser scanning microscopes over the past twenty years. Because they provide

complimentary measurements, FCS and FRAP can be used together to probe binding kinetics of

protein-protein interactions[12,13]. However, it is difficult to identify transient binding events from

these data because these methods provide ensemble-averaged data. Alternatively, Forster

resonance energy transfer (FRET), measures the efficiency of non-radiative energy transfer from

an excited donor fluorophore to an acceptor fluorophore in the ground state over a distance of a

few nanometers[14,15]. Thus, FRET efficiency serves as a measurement of spatial proximity of

fluorescently labeled proteins and can be used to infer protein-protein interactions.

Detecting interactions of diffusing proteins using dual-color single-molecule microscopy

is not practically feasible because both fluorophores used to label the putative interacting proteins

must be in a fluorescent ON state at the same time within the same complex. The probability of

observing such co-diffusion directly in separate color-channels is negligibly small, because

fluorophore photo-activation or blinking in PALM/STORM occurs stochastically. To avoid this

problem, the diffusive behaviors of putative interacting proteins can be measured in separate

experiments. If the two proteins diffuse at the same rate, then it is possible that they exist within

the same protein complex. Because single-molecule trajectories in living cells are limited in

duration by fluorophore photobleaching, information from thousands of trajectories are often

pooled and statistical data analysis approaches are employed to resolve different diffusive states[16-

23]. Indeed, different diffusive states can be resolved in living cells due to homo- and

heterooligomeric complex formation among interacting proteins[23-25] or proteins binding to quasi-

stationary structures, such as DNA[26-36]. However, to assign diffusive states to specific

biomolecular complexes of different protein compositions, additional experimental perturbation is

needed. Single-point mutations or genetic deletion mutants of potential interacting partners can be

used as a tool to disrupt putative complexes and observe changes in diffusive behavior. A drawback

of such genetic approaches is that the permanent disruption of an interaction interface or absence

of a binding partner can interfere with cellular processes in ways that are difficult to predict or

control, or yield the cells inviable[37,38]. Therefore, controllable, transient perturbations of cellular

processes on experimentally-relevant time scales is preferable.

Optically and chemically induced dimerization provides a favorable alternative to

permanent genetic perturbations. Such optogenetic and chemogenetic dimerization systems enable

manipulation of protein spatial localization in so-called knocksideways assays[39,40]. In a

knocksideways experiment, one of the dimerizing molecules is attached to a cytosolic protein of

interest and the other dimerization partner is targeted to a cellular location that differs from the

native localization of the tagged protein of interest (e.g. the cell membrane). Upon optical or

chemical stimulation of dimerization, the protein of interest is sequestered away from its native

localization, and interacting proteins can be sequestered along with it. Optogenetic knocksideways

experiments can thus be used to detect molecular interactions in living cells.

We reasoned that manipulating the spatial distributions of otherwise cytosolic proteins

would aid the assignment of diffusive states (observed in single-molecule tracking experiments)

to protein complexes of specific protein composition. In the simplest scenario, depletion of the

cytosolic diffusive state corresponding to the suspected oligomer would be expected when either

interacting partner is sequestered to the membrane. Chemogenetic systems rely on dimerization

being induced by a diffusing molecule itself, whereas optogenetic systems can be activated by

photon absorption. Optogenetic systems thus allows for more precise spatial and temporal control

over dimerization than chemogenetic systems.

Here, we test the possibility of combining optogenetic manipulation with 3D single-

molecule tracking microscopy in live *Esherichia coli*. The improved Light Induced Dimerization

(iLID) system[4] was selected because it has been extensively characterized and engineered for

optimized performance in different model systems. Thus, a considerably large toolbox of iLID

variants with different affinities, and reversion times are available for implementation[41-43]. The

iLID protein contains the light sensing light oxygen voltage (LOV2) domain derived from *Avena*

sativa which incorporates a flavin cofactor during folding. The flavin cofactor acts as a

chromophore, and forms a cysteine adduct with the LOV2 domain after illumination with blue

light. As a result, the iLID protein changes conformation and exposes a binding site for the

interacting partner, SspB[44,45]. We demonstrate that the iLID system exhibits a robust response after

488 nm illumination in low-intensity excitation conditions used for diffraction-limited imaging.

Using single-molecule tracking, we found that $SspB_{micro}$ exhibited the most substantial 488 nm-

induced response among three tested SspB mutants. Surprisingly, our results also reveal that

activation of the optogenetic response also occurs under high intensity illumination conditions,

even at wavelengths for which iLID shows minimal absorbance. Titrating the iLID expression

level with respect to SspB reduced imaging laser-induced iLID:SspB interaction in 3D single-

molecule tracking experiments Together our results establish the need for careful calibration of the

optogenetic system prior to its application to imaging experiments.

RESULTS AND DISCUSSION

Diffraction-limited imaging indicates a robust redistribution of otherwise cytosolic

fluorescence to the membrane upon blue-light activation.

Optogenetic tools have been widely applied in eukaryotic cells in conjunction with

diffraction-limited fluorescence microscopy to obtain population-averaged, phenotypic

readouts[46,47]. Therefore, we sought to quantify the optogenetic response of the iLID system in K12

E.coli using diffraction-limited imaging. We expressed the iLID protein from an inducible

arabinose promoter and targeted it to the inner membrane using an N-terminal genetic fusion

containing a modified single transmembrane spanning helix derived from the *E.coli*

transmembrane protein, TatA[48,49]. This construct, herein referred to as MA-iLID, was previously

used for stable insertion into the inner membrane of gram negative bacteria and has been shown

to enable optogenetic disruption of type 3 protein secretion in *Yersinia enterocolitica.*[50]. We

genetically fused the strongest reported iLID binding partner, $SspB_{nano}$, with a C-terminal Halo

tag, and expressed this construct from a second, constitutively expressing plasmid. $SspB_{nano}$-Halo

was labeled with JFX549 for visualization in fluorescence imaging experiments[51] (**Figure 1a**). For

simplicity, the JFX549-labeled $SspB_{nano}$-Halo is referred to as $SspB_{nano}$ in this report.

Prior to optical activation of MA-iLID with 488 nm laser light, we observed a uniform distribution

of cytosolic $SspB_{nano}$. Redistribution of $SspB_{nano}$ to the membrane was dependent on both

expression of MA-iLID and 488 nm laser illumination (~1mW/cm^2) (**Figure 1b-d**). We quantified

$SspB_{nano}$ redistribution upon 488 nm laser illumination in each cell using the structural similarity

index measure (SSIM), which quantifies similarity between two images[52]. Cells expressing MA-

iLID were more dissimilar (1-SSIM) after 488 nm laser illumination compared to controls (**Figure**

1e). To estimate the fraction of molecules that reside at the membrane in each condition, we

simulated 2D diffraction-limited images that matched our imaging parameters and contained

realistic noise contributions (see methods) (**Figure 1f, SI Figure 1a**). Fluorescence line profiles

across the midsection of cells showed a Gaussian-like line shape for cytosolic fluorescence, while

membrane-associated fluorescence produced double-peaked line shapes. The experimental line

shape obtained with a fluorescently labeled, membrane-anchored iLID protein (MA-mCherry-

iLID) agreed well with line shape obtained based on simulated images of 100% membrane

localized molecules (**SI Figure 1b**). Based on qualitative comparison of simulated and measured

line shapes, we estimate that the fraction of molecules at the membrane in living cells is close to

0% pre-activation and greater than ~80% post-activation. These data suggest that the iLID system

can be used to efficiently sequester otherwise cytosolic molecules to the membrane.

Figure 1. Diffraction-limited imaging indicates a substantial redistribution of SspB$_{nano}$ to the membrane after blue-light activation. (a) Schematic depiction of a knocksideways experiment using the iLID system. The iLID protein undergoes a conformational change after 488 nm-light illumination and uncages a binding site for SspB. Halo-tagged (red star) cytosolic SspB binds to membrane-anchored iLID (MA-iLID) after 488 nm illumination. SspB-Halo is labeled with JFX549 for fluorescence imaging (hereafter referred to as SspB). **(b)** Diffraction-limited images of SspB$_{nano}$. Illumination with 488 nm light has no effect on the spatial distribution of SspB$_{nano}$ in the absence of MA-iLID. Fluorescence intensity line profiles across the midsection of the cell (normalized to the integrated line intensity) also exhibit no apparent shift in SspB$_{nano}$ spatial distribution. **(c)** Low intensity (~1 W/cm^2) 561 nm laser excitation over the entire exposure (80 seconds) is not sufficient to induce SspB$_{nano}$ sequestration to the membrane. **(d)** In the presence of MA-iLID, low intensity (~4 mW/ cm^2) 488 nm laser illumination shifts the spatial distribution of SspB$_{nano}$ to the cell membrane. The fluorescence intensity line profile across the midsection of the cell changes from a Gaussian-like line shape to a characteristic double-peaked line shape. **(e)** Left panel: Image dissimilarity (1-SSIM) of experimentally-acquired individual cell images before and after 488 nm illumination (n=200-350 cells in each condition). A population-level response in image dissimilarity is dependent on MA-iLID expression and 488 nm illumination. Right panel: Calibration of the image dissimilarity metric as a function of the fraction of membrane-associated fluorophores (relative to a simulated cell with 0% membrane-associated fluorophores). **(f)**

189 Simulated diffraction limited images of cells with 0% or 100% membrane-associated fluorophores,
190 respectively. Normalized line profiles of simulated diffraction-limited images (**SI Figure 1**)
191 indicate a progressive broadening, and redistribution of fluorescence as the fraction of membrane-
192 associated fluorophores increases.

3D single-molecule tracking data show increased SspB:iLID interaction prior to 488 nm illumination.

Single-molecule tracking microscopy has been used to resolve different diffusive states of

proteins that manifest in living cells due to protein complex formation[53]. Because optical activation

of the MA-iLID:SspB interaction allows for transient spatial redistribution of protein localization,

the light-driven interaction could provide a means to non-invasively perturb the diffusive states

that manifest in the cytosol of living cells. We therefore sought to measure SspB diffusive behavior

in living cells and determine the extent of MA-iLID association using 3D single-molecule tracking.

Based on the clear membrane localization of MA-iLID in diffraction limited images (SI Figure 1),

we reasoned that $SspB_{nano}$ interacting with MA-iLID should diffuse along the membrane at a much

slower rate than cytosolic $SspB_{nano}$. In 488 nm illuminated cells, the cumulative distribution

function of apparent diffusion coefficients indeed shows a bimodal curve with a transition at $D^* =$

0.15 μm^2/s (**Figure 2a**). When plotting the spatial trajectories of slowly diffusing molecules (D^*

< 0.15 μm^2/s) it is clear that they localize near the bacterial membrane. On the other hand, the

spatial trajectories of fast diffusing molecules ($D^* > 0.15$ um^2/s) localize in the cytosol (**Figure**

2b). We conclude that it is possible to clearly distinguish freely-diffusing, cytosolic $SspB_{nano}$ from

iLID-associated $SspB_{nano}$ molecules in living cells based on their diffusion rate and sub-cellular

localization.

The $D* = 0.15$ um^2/s threshold allowed us to estimate the relative population fractions of

iLID-associated and cytosolic SspB$_{nano}$. In cells that do not express MA-iLID, the vast majority

(95%) of SspB$_{nano}$ proteins diffuse fast and localize to the bacterial cytosol, as expected. The

remaining 5% of tracked proteins were classified as slowly diffusing, but the corresponding

trajectories showed no discernable preference for the membrane. The diffusive behavior and

spatial distribution of SspB$_{nano}$ remained unchanged upon 488 nm illumination in these cells

(**Figure 2a** and **c**). Surprisingly, and in contrast to the diffraction-limited images (**Figure 1**), a

large fraction of SspB$_{nano}$ proteins exhibited slow diffusion (48%) in cells that express MA-iLID,

even in the absence of 488 nm light. This shift towards slow diffusion is accompanied with spatial

redistribution of SspB$_{nano}$ to the membrane, whereas the fast diffusing population of proteins

remained uniformly localized to the cytosol (**Figure 2d**). When cells expressing both SspB$_{nano}$ and

MA-iLID were exposed to 488 nm light, we observed an additional shift towards slower diffusion

and membrane-proximal localization (56%) (**Figure 2e**). The increase in iLID-associated SspB

observed is marginal compared to the shift observed in diffraction-limited images (**Figure 1**).

However, we note that the line profile quantitation of our experimental images represents an

example of a single cell displaying an obvious change in SspB$_{nano}$ spatial distribution. Across a

population the degree of response observed in line profiles varies. Further, the population level

analysis of image dissimilarity performed on the diffraction limited images indicates a varied

response to 488 nm illumination. This is likely due to expression level heterogeneity which is

evident in whole cell fluorescence comparisons of both arabinose-promoter driven expression of

MA-iLID, and constitutive expression of SspB (**Figure 1, SI Figure 1**).

We hypothesize that the discrepancy of MA-iLID associated SspB seen in diffraction-

limited image quantitation compared to single-molecule trajectory data is due to the higher

excitation intensity required for single-molecule localization microscopy. The absorption spectrum

of iLID proteins shows very minimal absorption at wavelengths larger than 500 nm[54]. However,

optical activation of the iLID:SspB$_{nano}$ interaction is possible with 514 nm light at illumination

intensities typical for diffraction-limited imaging (~1 W/cm^2). (**SI Figure 2**). The intensity used

for JFX549 fluorescence excitation by 561 nm laser light was three orders of magnitude higher in

single-molecule imaging experiments (2 kW/cm^2). These data suggest that an increase in 561 nm

photon flux is able to activate the iLID optogenetic response.

Figure 2. 3D single-molecule tracking data show SspB:iLID interaction prior to 488 nm illumination (a) Apparent diffusion coefficient distributions of SspB$_{nano}$ in the absence (purple, solid line, N = 11,523 tracks, 120 cells) and presence (blue, solid line, N = 8,542 tracks, 78 cells) of MA-iLID. Illumination with 488 nm light marginally increases the population of slowly diffusing SspB$_{nano}$ in MA-iLID expressing cells (blue, dashed line, N = 6,360 tracks, 88 cells), but not in cells which do not express MA-iLID (purple, dashed line, N = 9,922 tracks, 94 cells).

Vertical dashed line indicates threshold used to distinguish fast diffusers ($D* > 0.15$ µm^2/s) from
slow diffusers ($D* < 0.15$ µm^2/s) **(b)** Single-molecule trajectories in a representative cell
expressing MA-iLID, after illumination with 488 nm light. Left: top view. Right: *en face* cross
sectional view. Slowly diffusing molecules (blue) clearly localize to the periphery of the cell and
diffuse laterally along the membrane, while fast diffusing molecules (red) localize predominately
to the cytosol and diffuse in 3D. **(c-f)** 2D histograms of localizations from slow and fast trajectories
within the population of cells imaged. Histograms recapitulate trends observed in panel b. Bin size
20 nm x 20 nm. Each histogram is normalized to the total number of molecules observed in that
experiment. Inset: rescaled (unnormalized) histograms, shown for clarity.

Single-molecule tracking of SspB mutants further indicate activation of iLID optogenetic

response by high intensity 561 nm light.

To decrease the amount of iLID-bound SspB prior to 488 nm illumination, we made a series of

previously characterized mutations to SspB[41,42]. These mutant versions of SspB, termed SspB$_{micro}$

and SspB$_{milli}$, have decreased binding affinity for the iLID protein. When we tracked single

SspB$_{micro}$ and SspB$_{milli}$ proteins in living cells, we indeed observed a decrease in the fraction of

slow diffusers near the cell membrane prior to 488 nm light illumination (**Figure 3a-c**). We found

that 31% of SspB$_{micro}$ proteins were iLID-associated, while iLID-associated SspB$_{milli}$ comprised

only 6% of the population. Upon 488 nm light illumination, the membrane-proximal, slow-

diffusing fractions increased to 45% and 7%, respectively (**Figure 3b** and **c**). The magnitude of

the optogenetic response thus differs between the mutants, with SspB$_{micro}$ showing the largest

response, under otherwise identical experimental conditions. Diffraction-limited images of

SspB$_{micro}$ and SspB$_{milli}$ recapitulate this trend (**SI Figure 3**). Previous work has also shown that

the SspB$_{milli}$:iLID interaction is only minimally affected by blue light, as judged based on SspB$_{milli}$

localization to iLID-labeled organelles in HeLa cells[55].

The reported dissociation constants for the three abovementioned SspB mutants differ by

six orders of magnitude. The data reported above was collected at the same expression levels for

each interacting protein components. $SspB_{nano}$ may thus saturate the available binding sites at the

membrane at the expression level used. Consequently, a large increase in binding upon 488 nm

illumination is not observed. At the other extreme, $SspB_{milli}$ may require a much higher number of

binding sites to observe binding both before and after 488 nm illumination due to its inherently

low affinity for the iLID protein. The comparatively larger optogenetic response observed for

$SspB_{micro}$ suggests that the concentration of each interacting parter is better-matched to the

dissociation constant for this mutant.

To test this model, we titrated the amount of MA-iLID, while keeping the expression level

of $SspB_{micro}$ constant. To estimate the concentration of MA-iLID expressed at each L-arabinose

induction concentration, we performed a calibration experiment in which we expressed MA-

mCherry-iLID under the same promoter. The number of MA-mCherry-iLID proteins was

estimated by dividing the intial fluorescence intensity of each cell by the average intensity of a

single mCherry protein. Uninduced cells still expressed MA-mCherry-iLID at a concentration of

~90 nM due to leaky expression from the arabinose promoter (**SI Figure 4**). At the maxium inducer

concetration used (13.3 mM), the MA-mCherry-iLID reached a concentration of ~510 nM. For the

following analyses, we assume that MA-mCherry-iLID and MA-iLID are expressed at the same

level for a given L-arabinose concentration.

Decreasing the concentration of MA-iLID, decreases the fraction of iLID-associated

$SspB_{micro}$. At the lowest MA-iLID concentration, $SspB_{micro}$ does not show an appreciable

population fraction of slowly diffusing, membrane-proximal molecules both before and after 488

nm illumination (**Figure 3d**). At the highest concentration of MA-iLID , we estimate that 48% of

$SspB_{micro}$ is iLID-associated. It is possible that this fraction could be augmented by further

increasing the expression level of MA-iLID. However, we found that high expression of MA-

mCherry-iLID leads to an increase in fluorescent foci formation, most often at the cell pole (**SI**

Figure 5a). Fluorescent foci formation can be indicative of protein overexpression and subsequent

aggregation and/or formation of inclusion bodies[56,57]. These processes do not necessarily interfere

with protein folding and function[56,57]. Indeed, we observe fluorescent foci formation in the MA-

mCherry-iLID data, but not to the same extent in the $SspB_{micro}$-Halo data (**SI Figure 5b**). These

observations suggest that mCherry remains correctly folded and functional within the MA-

mCherry-iLID construct, though the iLID protein within the fluorescent foci may not be functional

or available for interactions with SspB. Due to the significant increase in fluorescent foci

formation, we did not explore higher expression levels of MA-iLID.

Titration of MA-iLID expression allowed us to estimate the dissociation constant K_D of the

$SspB_{micro}$:MA-iLID interaction in living cells. Because we observe 48% association of $SspB_{micro}$

at 507 nM MA-iLID, we estimate that the K_D is approximately 510 nM before 488 nm illumination.

This value is about 100x smaller than K_D reported in *in vitro* fluorescence polarization assays in

unactivated conditions[41]. However, as we argued above, the high intensity 561 nm excitation light

activates the iLID optogentic response. Therefore, it is more appropriate to compare results under

activated conditions to the fluorescence polarization data. Indeed, the reported *in vitro* K_D is about

800 nM,[41] which is within a factor of two the K_D value we estimate in living cells. Further, we

observe ~45% association of $SspB_{micro}$ at ~260 nM MA-iLID in activated conditions, which

represents an approximately 2-fold increase in affinity upon 488 nm illumination. The reported

dynamic range derived from *in vitro* fluorescence polarization assays is reported to be a 58 fold

change in binding affinity due to activation of the system. However, we note that proteins

reconstituted in lipid bilayers show a shortened dynamic range of binding affinity, observing a 2.7-

fold change in affinity upon iLID-activation[58,59]. Together, these results suggest that proximity to

the membrane, and the steric hindrance imposed by it may affect the dynamics of the SspB:iLID

interaction.

Figure 3. Affinity of the SspB:iLID interaction determines the fraction of iLID-associated
SspB observed in single-molecule tracking experiments. **(a)** Apparent diffusion coefficient
distributions for three SspB mutants. The lowest affinity mutant, $SspB_{milli}$ (solid, yellow line, $N =$
4246 tracks, 65 cells) displays the highest population fraction of fast, cytosolic diffusion prior to
488 nm illumination, followed by $SspB_{micro}$ (solid, orange line, $N = 7{,}738$ tracks, 97 cells) and
$SspB_{nano}$ (solid, blue line, $N = 8{,}542$ tracks, 78 cells). Activation of these systems with 488 nm
illumination (dashed line distributions) shifts each mutant proteins's distribution towards larger
fractoins of slow diffusing molecules due to MA-iLID association. **(b)** 2D cross-sectional
histograms of $SspB_{micro}$ trajectories indicate MA-iLID association at the membrane increases in
frequency upon 488 nm illumination ($N = 6{,}491$ tracks, 77 cells). Each histogram is normalized to
the total number of molecules observed in that experiment. Inset: rescaled (unnormalized)
histograms, shown for clarity. **(c)** 2D cross-sectional histograms of $SspB_{milli}$ trajectories indicates

small, but qualitatively discernable MA-iLID association after 488 nm illumination (N = 5243
tracks, 99 cells). **(d)** Apparent diffusion coefficient distribution of SspB$_{micro}$ as a function of MA-
iLID expression level. Higher MA-iLID expression levels lead to increased population fractions
of slow, membrane-proximal diffusion of SspB$_{micro}$ (see also **SI Figure 4** and **6**).

Cumulative displacement analysis identifies state-switching events in full-length trajectories

Having quantified the dissociation constant of the iLID:SspB interaction, we next sought

to determine the kinetics of this interaction. Previous analyses were performed on trajectories

which were shortened to contain at most 10 displacements, to prevent averaging over multiple

states. Because our data sets contained many long trajectories, we reasoned that some of these

trajectories may contain information about the length of time a single SspB protein remains bound

to an MA-iLID protein at the membrane. To assess the diffusive behavior over the course of a

trajectory, we first calculated the cumulative displacement of each single-molecule over time

(**Figure 4a** and **4b**, see methods). These data show a clear separation for fast cytosolic diffusion

and slow, membrane-associated diffusion. Because of the clear distinction, we decided that

analysis cumulative displacements could be used to differentiate states within single trajectories.

To estimate the diffusion coefficients that describe motion in each state, we simulated cytosolic

and membrane-bound diffusion at different diffusion rates. We found that the cumulative

displacement over time of single-molecule trajectories simulated at 5.5 µm^2/s in the cytosol and

0.2 µm^2/s at the membrane best matched the experimentally measured cumulative displacement

data. These values were then used as references for subsequent analysis.

To identify state switching events within trajectories, we employed the workflow outlined

in Figure 4a and c. Briefly, the cumulative displacements of small segments within trajectories

were designated as fast diffusion or slow diffusion using a threshold corresponding to D = 1.75

µm^2/s. This threshold was chosen to match the upper-limit of the displacement sizes used in the

membrane-bound simulation at 0.2 µm^2/s. Trajectories which exhibited subsequent fast-slow-fast

diffusive states were ranked by evaluating the similarity of experimental slope of cumulative

displacement data to the abovementioned simulated slopes of cumulative displacement data

corresponding to $D = 5.5$ µm^2/s and 0.2 µm^2/s (**Figure 4d**). We further verified that the best scoring

trajectories indeed exhibited slow-diffusing segments at the membrane, and fast-diffusing

segments in the cytosol. The resulting subset of trajectories was then used to calculate t_{bound} (**Figure**

4e). As a control, we performed the same analysis for SspB trajectories in the absence of MA-

iLID, and, as expected, did not identify clear state switching signatures.

In the presence of 256 nM MA-iLID, SspB$_{micro}$ and SspB$_{nano}$ exhibit similar bound times

independent of 488 nm illumination (median = 300 and 350 ms, respectively) while SspB$_{milli}$

displayed comparatively shorter binding times (median = 175 ms) (**Figure 4e**). Thus, the

measurements of t_{bound} made in this manner do not show a trend that clearly correlates with each

SspB mutant's affinity for the iLID protein or with photoactivation of iLID. Trajectories which

were identified as acceptable for residence time measurements were rare (<1% or less of the total

number of trajectories analyzed). This low number stands in stark contrast to the much larger

fraction of molecules which remained iLID-bound for the duration of the trajectory which matched

the trend observed in the analyses of apparent diffusion coefficients (**SI Figure 7**, **Figures 2 and**

3). The length of the trajectories of fully-bound molecules ranges from 0.3 seconds up to 3 seconds

(longest observed trajectory – **Figure 4b**). The long lengths of these trajectories (which are

terminated by fluorophore photobleaching) suggest that SspB$_{micro}$ proteins can remain iLID-bound

for much longer than the median time scales estimated for t_{bound}. Previously published data

estimates the dissociation half-life of SspB$_{nano}$ and SspB$_{micro}$ with iLID to occur on the order of

minutes[58,59]. Thus, single-molecule trajectories are not long enough to capture both binding and

unbinding events of stable SspB:iLID interactions. These results suggest that short time scale

interactions that we capture are representative of a transient binding mode(s) which is(are) not

reflective of stable SspB:iLID binding.

Figure 4. Cumulative displacement analysis identifies state-switching events in single-molecule trajectories. (**a**) Workflow of cumulative displacement analysis to distinguish fast, cytosolic diffusing molecules from slow, membrane-associated molecules. (**b**) Top panel: Schematic of displacement data transformed into a 1D trajectory to illustrate displacement size over time. The cumulative displacement is the sum of all displacements up to a given time point, t. Bottom panel: Cumulative displacement over time for SspB$_{micro}$ trajectories collected in MA-iLID expressing cells (256 nM) in unactivated conditions. A threshold slope corresponding to $D = 1.75$ µm^2/s was used to distinguish fast diffusing molecules (blue) from slow diffusing molecules (red). Simulated trajectories in the cytosol at 5.5 µm^2/s and in the membrane at 0.2 µm^2/s recapitulated experimental cumulative displacement slopes and were used as reference slopes in the following steps. (**c**) Workflow used to identify trajectories with state-switching (see text for details). (**d**) Example of a SspB$_{micro}$ partial trajectory which exhibits state-switching. Three consecutive cumulative displacements are fit to a line, and the slope is compared to the threshold slope to classify the trajectory segment as cytosolic or membrane-bound. The slope of the segment is then compared to the reference slope for cytosolic diffusion or membrane diffusion depending on the state assignment, and scored by how closely they match. Trajectories with at least 2 consecutive segments in cytosolic and then membrane-associated states (or vice versa) are categorized as state-switching and used for later analysis. (**e**) Cumulative displacements over time of trajectories which show state transitions of fast-slow-fast. Trajectory localization of example fast(blue)-slow(red)-fast(blue) state-switching cumulative displacement plotted to highlight state transition at the membrane of the cell. Time spent in bound state (red portion of trajectory) is integrated to calculate the residence time t_{bound}. (**f**) Histogram of residence times of SspB affinity mutants (bin size = 250 ms) indicate that transient binding modes to MA-iLID do not exhibit residence times that are dependent on the affinity of the SspB:iLID interaction.

CONCLUSIONS

Here, we provide a quantitative analysis of the iLID optogenetic system in live K12 *E. coli* cells using 2D diffraction-limited imaging and 3D single-molecule tracking. We show that the iLID system enables efficient optogenetic manipulation of protein spatial distributions when used in conjunction with diffraction-limited imaging. Single-molecule tracking data acquired using high intensity 561 nm laser excitation leads to a substantial amount of SspB:iLID interaction even in the absence of optogenetic activation at 488 nm. This suggests that increased photon flux is somehow able to activate the iLID optogenetic response, even though the LOV2 domain absorption spectra suggest that wavelengths greater than 500 nm should not be absorbed[54]. This conclusion is consistent with our observation that even low intensity 514 nm illumination is able

to increase the SspB:iLID interaction. However, published LOV2 domain absorption spectra were

collected using purified proteins[54]. It is possible that cellular contents, proximity to the membrane

and imposed steric hindrance results in a shift in iLID conformational dynamics and thus a change

in interaction with SspB. A similar effect has been seen in other model systems utilizing the

iLID:SspB interaction[58,59]. This may explain the rare, transient binding events that we measured

in single- molecule trajectory analyses. Our results highlight the importance of calibrating the

effects of illumination conditions and cellular environment when the iLID system, or any

optogenetic system, is used to modulate an SspB-tagged protein's spatial distribution.

A notable advantage of optogenetic systems, such as iLIDs, is that dimerization can be

induced in real time using light. Optically-induced dimerization can be enabled much faster than

chemically induced dimerization, which are diffusion-limited. We aim to combine optogenetic

perturbation with single-molecule tracking to deplete cytosolic diffusive states that have been

assigned to hetero-oligomeric protein complexes based on control experiments in genetically

perturbed backgrounds. Although a pre-activated system dampens the magnitude of the

optogenetic response that can be observed, highly sampled distributions obtained from tens of

thousands of single-molecule trajectories, may allow for the observation of cytosolic diffusive

states depletion with sufficient statistical power. Additionally, titration of protein expression can

be used to incrementally sequester proteins of interest. Another alternative might be to use far-red

excitable fluorophores for single-molecule tracking. Based on the trends observed here we

speculate that high intensity red and far-red light illumination will not activate the iLID system.

Thus, single-molecule localization and tracking microscopy with red and far-red fluorophores

could enable observations of large optogenetic responses, similar to those observed in diffraction-

limited imaging.

METHODS:

Bacterial strains and plasmids

Escherichia coli K12 (MG1655) strains were generated by introducing pACYC and pBAD vectors

containing genes encoding the cytosolic prey protein, SspB, and the membrane-anchored bait

protein, iLID, respectively. pACYC $SspB_{nano}$-Halo was derived from pACYC $SspB_{nano}$-mCherry

(pFL109) which was a gift from Andreas Diepold. The construct was modified by removing

mCherry from the plasmid via restriction digest and the *halo* coding sequence was ligated in its

place, using XhoI and SalI cut sites. The plasmid containing the bait protein, pBad FLAG-iLID

(pFL108), also a gift from the Diepold lab, was transformed into bacteria without any further

modification.

Cell culture

E. coli cultures were inoculated from a freezer stock and grown overnight in LB media at 37° C

and shaking at 225 rpm. Strains expressing from the pACYC and pBAD plasmids were grown in

30 µg/mL chloramphenicol and 100 µg/mL ampicillin, respectively. Overnight cultures were

diluted to an OD_{600} of 0.05 in M9 minimal media and incubated at 37° C. Strains containing the

pBAD plasmid were induced with 5.33 mM L-arabinose after 2 hours unless indicated otherwise.

After an additional 1.5 hours of growth, 1.5 mL of cell culture was aliquoted and stained with 1.5

µL of 500 µM JFX-549 dye. The cell suspension mixture was incubated at 37° C for 30 minutes

and then washed four times with M9 minimal media. Cell pellets were resuspended in a final

volume of 10 µL for use in imaging experiments.

E. coli cells were transformed with plasmids via heat shock of chemically competent cells.

Transformed cells were selected for by growing on LB agar plates containing chloramphenicol,

ampicillin (100 µg/mL), or a combination of both for co-expressing strains. Transformants were

screened by performing a Western blot for FLAG or Halo proteins in pBAD and pACYC-

containing strains, respectively. Strains which showed a clear band in Western blot experiments

were used to make freezer stocks containing 15% glycerol.

Optical setup

Imaging of cells was performed on a home-built inverted fluorescence microscope as described

previously[24]. The microscope was modified for use in optogenetics experiments by incorporation

of an additional 488nm laser line (Coherent Genesis) for activation of the iLID system.

Fluorescence images were collected in the "red" channel of the imaging system.

Epifluorescence imaging

Stained cells were mounted on a #1.5 coverslips (VWR) and immobilized using 1.5% agarose pad

prepared in M9 minimal media. Halo-conjugated JFX549 was excited using 561 nm laser light at

~1 W/cm^2. Images were acquired at 5 frames/sec for 40 seconds pre- and post-activation of the

iLID optogenetics system. The iLID optogenetic system was activated using 488 nm laser light at

~4 mW/cm^2 continuously for 2 minutes prior to and then also during acquisition of post-activation

images. Epifluorescence images shown in the figures represent averages of these image sequences.

Brightfield images were acquired of the same field-of-view by illuminating cells with an LED, and

taking a single 25 ms exposure.

SSIM analysis of Diffraction-limited images

Pre- and post- activation images were background subtracted. Outlines of cells were derived by

using OUFTI[60] on inverted color to create a binary mask such that each cell was compared to itself

before and after activation. The built-in Matlab (The MathWorks, Inc, Natick, Massachusetts)

function ssim() was applied to the masked images with equal weighting of the comparison metrics,

luminance, structure, and contrast, to assess the degree of image similarity. Cells with abnormal

morphology (based on the associated brightfield image) were excluded from the analysis. Multiple

fields of view were acquired for each condition, and the experiments were replicated twice.

Super-resolution fluorescence imaging

Fluorescent fiducial markers (Invitrogen) were added to cell suspensions and the cell suspension

was then mounted on #1.5 glass coverslips. Cells were immobilized using solidified pads of 1.5%

agarose in M9 minimal media. Halo-conjugated JFX-549 was excited into the blinking state by

561 nm laser light at ~2 kW/cm^2 at the sample. Images were acquired at 40 frames/sec in the

presence or absence optogenetic activating 488 nm light. Cells were activated using 488 nm laser

light at ~4 mW/cm^2 continuously for 2 minutes prior to and during acquisition of post-activation

images.

Data Processing

Raw data was processed in Matlab (The MathWorks, Inc, Natick, Massachusetts) using a modified

version of easy-DHPSF software[24,61]. Fluorescent fiducial markers were used for sample drift

correction. Single-molecule localizations were registered to phase contrast images using a two-

step affine transformation in MATLAB as previously described[24]. Localizations outside of cells

were discarded from further analysis using axial bounds and OUFTI-derived cell outlines which

were generated from phase contrast images, as described above.

Single-molecule tracking analysis

To derive single-molecule displacements, localizations in subsequent frames were linked into

trajectories. A maximum linking distance of 2.5 um was used for linking analysis, and multiple

localizations which were present at the same time within a single cell were discarded to prevent

misassignment of molecules.

The Mean Squared Displacement (MSD) for each trajectory was calculated using

518
$$\text{MSD} = \frac{1}{(N-1)} \sum_{n=2}^{N} (x_n - x_{n-1})^2$$

519   where x is the 3D position at timepoint n, including up to 11 timepoints for calculating the mean

520   over 10 displacements. The remainder of the trajectory was not used in the MSD analysis to ensure

521   multiple diffusive states were not averaged over. The MSD measurement was then used to

522   calculate the apparent diffusion coefficient ($D*$) according to:

524
$$D^* = \frac{\text{MSD}}{2 \cdot m \cdot \Delta t}$$

Where m is the dimensionality of the measurement ($m=3$ for the 3D trajectories reported here),

and Δt is the camera exposure time used for imaging ($\Delta t=25$ ms under our conditions).

2D-cross section projection analysis

The vector describing the central axis of each cell was determined using the cell outline generated

from OUFTI. The outline was segmented into sections along the cell length, and localizations from

trajectories in each section were projected onto a 2D plane. The position of the central cell axis

was adjusted to match the centroid of all localizations within the section. Positions of localizations

from each cell were scaled to match the mean cell radius and mean cell length which was calculated

from OUFTI outlines. The trajectories were classified as slow or fast diffusing using the threshold

$D* = 0.15\,\mu\text{m}^2/\text{s}$ which was determined by identifying a transition point from the first major change

in slope in the CDF plot of apparent diffusion coefficients of data containing both iLID-associated

and cytosolic diffusion (**Figure 1**).

Determination of residence times - t_{bound}

To identify diffusive state transitions in single-molecule trajectories, we developed a

workflow to analyze displacement data over time (**Figure 4a and c**). First, we plotted the additive,

or cumulative displacement (CD) as a function of time elapsed according to:

$$\text{CD}(t) = \sum_{t=1}^{N} \sqrt{(x_t - x_{t-1})^2}$$

where x is the 3D position at timepoint t. The CD was calculated for each displacement, for the

full trajectory length (N).

The instantaneous rate of displacement was estimated by finding the slope within a small,

sliding window containing 3 displacements for the full length of the trajectory (**Figure 4d**). The

size of the sliding window was chosen to be small enough such that switching events were not

excessively averaged over, while being large enough to be somewhat insensitive to displacement

size fluctuations. Each segment of the trajectory was then classified as bound or free by calculating

its slope and comparing it to a threshold, which was chosen to match the upper-limit of the

displacement sizes used in the membrane-bound simulation at $0.2\,\mu\text{m}^2/\text{s}$. Trajectories which

contained consecutive segments of fast-slow-fast state assignments were used for further analysis

because of the presence of an internal bound state. To be considered for analysis, the molecule

must reside in each state for at least 2 segments (4 displacements) of the trajectory to ensure that

the identified state change was a true diffusive state change and not due to random fluctuation.

Each switching trajectory was ascribed a score (TS) by how well the slope of trajectory segments

matched simulated diffusion coefficient slopes derived for freely diffusing, cytosolic and

membrane-bound molecules:

558
$$TS = \frac{1}{N_{seg} - 2} \sum_{i=1}^{N_{seg}} e^{-|m_i - m_s|}$$

where m_i is the slope fit to the experimental trajectory segment, and m_s is the slope derived from

free cytosolic or membrane-bound simulated data, depending on the state assigned to the trajectory

segment. N_{seg} refers to the number of segments in the trajectory. Switching trajectories were ranked

by their TS, and the top-scoring trajectories (TS>~0.16) were inspected individually to verify that

switching events occurred at the membrane of the cell. The residence time *(t_{bound})* of bound

molecules in these switching trajectories was calculated by determine the number of displacements

the molecule was assigned to the bound state.

3D single-molecule trajectory simulation

Cytosolic trajectories were simulated as described previously.[23,24] Briefly, Monte Carlo

simulations of Brownian motion at 5.5 μm^2/s were confined to the volume of a cylinder matching

the dimensions of K12 *E. coli* (radius = 0.5 μm, length = 2.5 μm). Initial positions of molecules

were randomly selected to uniformly fill the volume of the cell, and subsequent positions in the

trajectory were selected at 100 ns intervals for a total of 125 ms. We simulated noisy, motion-

blurred images of the single-molecule trajectories, integrating images over subsequent 25 ms time

intervals, to match experimental camera exposure times for a total of 6 images per single-molecule

track. We simulated 1000 such trajectories. To best match data derived from experimental

conditions, we analyzed the simulated images of trajectories in the same way described for

experimental localization and tracking. Easy-DHPSF software was used to analyze the images and

detect single-molecule localizations.

Membrane-bound trajectories were simulated for molecules diffusing at 0.2 μm^2/s. Initial positions

were randomly chosen on the surface of a spherocylinder matching the dimensions of K12 *E. coli*

(radius = 0.5 µm, length = 2.5 µm). Monte Carlo simulations of molecular motion were confined

to the surface of the spherocylinder by translating displacements to changes in lateral position and

elevation angle on the cylindrical surface or elevation and azimuthal angle on the hemispherical

surface, such that the distance from the cell axis is always equal to the radius of the cell. Positions

were updated at 100 ns intervals for a total of 125 ms. Images were generated and analyzed as

described in the previous paragraph.

2D diffraction-limited image simulation

Trajectories of cytosolic and membrane-associated diffusing molecules were simulated as

described above with the following modifications to match diffraction-limited imaging conditions.

Positions were sampled every 1 µs for a total of 1.2 seconds. Molecule positions were sampled

ever 200 ms, resulting in 6 localization positions (macro-positions) per track.

The three dimensional molecule macro-positions were used to generate conventional diffraction-

limited 2D point-spread functions of isotropic emitters using a vectorial light propagation model.[62]

We modified the algorithm to omit the double helix phase mask and optical aberrations. In the

simulation, the nominal focal plane was positioned at the center of the cell. A total of 10,000

trajectories were simulated for both cytosolic and membrane-bound molecules, providing 60,000

total emitter positions in each condition. Images obtained for each emitter were added to attain the

total image. To attain mixed population images, trajectories were chosen at random from the total

simulated trajectory population to reflect the specified membrane-associated molecule fraction.

REFERENCES:

1   Achimovich, A. M., Ai, H. & Gahlmann, A. Enabling technologies in super-resolution

fluorescence microscopy: reporters, labeling, and methods of measurement. *Current*

Opinion in Structural Biology **58**, 224-232,

doi:https://doi.org/10.1016/j.sbi.2019.05.001 (2019).

2   Aaron, J. S., Taylor, A. B. & Chew, T.-L. Image co-localization – co-occurrence versus

correlation. *Journal of Cell Science* **131**, jcs211847, doi:10.1242/jcs.211847 (2018).

3   Szymborska, A. *et al.* Nuclear Pore Scaffold Structure Analyzed by Super-Resolution

Microscopy and Particle Averaging. *Science* **341**, 655-658,

doi:10.1126/science.1240672 (2013).

4   Sieben, C., Banterle, N., Douglass, K. M., Gönczy, P. & Manley, S. Multicolor single-

particle reconstruction of protein complexes. *Nature Methods* **15**, 777-780,

doi:10.1038/s41592-018-0140-x (2018).

5   Thevathasan, J. V. *et al.* Nuclear pores as versatile reference standards for

quantitative superresolution microscopy. *Nature Methods* **16**, 1045-1053,

doi:10.1038/s41592-019-0574-9 (2019).

6   Malkusch, S. *et al.* Coordinate-based colocalization analysis of single-molecule

localization microscopy data. *Histochemistry and cell biology*, doi:10.1007/s00418-

011-0880-5 (2011).

7   Larson, J. D., Rodgers, M. L. & Hoskins, A. A. Visualizing cellular machines with

colocalization single molecule microscopy. *Chem. Soc. Rev.* **43**, 1189-1200,

doi:10.1039/c3cs60208g (2014).

8   Levet, F. *et al.* A tessellation-based colocalization analysis approach for single-

molecule localization microscopy. *Nature communications* **10**, doi:10.1038/s41467-

019-10007-4 (2019).

9   Magde, D., Elson, E. & Webb, W. W. Thermodynamic Fluctuations in a Reacting

System—Measurement by Fluorescence Correlation Spectroscopy. *Phys. Rev. Lett.*

29, 705-708, doi:10.1103/physrevlett.29.705 (1972).

10   Axelrod, D. K., D. E.; Schlessinger J.; Elson, E.; Webb, W.W. Mobility measurement by

analysis of fluorescence photobleaching recovery kinetics. *Biophysical journal* **16**,

1055-1069, doi:https://doi.org/10.1016/S0006-3495(76)85755-4 (1976).

11   Koppel, D. E. A., D.; Schlessinger, J.; Elson, E. L.; Webb, W. W. . Dynamics of

fluorescence marker concentration as a probe of mobility. *Biophysical journal* **16**,

1315-1329 (1976).

12   Wachsmuth, M. Molecular diffusion and binding analyzed with FRAP. *Protoplasma*

251, 373-382, doi:10.1007/s00709-013-0604-x (2014).

13   Muller, K. P. *et al.* Multiscale analysis of dynamics and interactions of

heterochromatin protein 1 by fluorescence fluctuation microscopy. *Biophysical*

journal **97**, 2876-2885, doi:10.1016/j.bpj.2009.08.057 (2009).

14   Zeug, A., Woehler, A., Neher, E. & Evgeni. Quantitative Intensity-Based FRET

Approaches—A Comparative Snapshot. *Biophysical journal* **103**, 1821-1827,

doi:10.1016/j.bpj.2012.09.031 (2012).

15 Forster, T. Energiewanderung und fluoreszenz. *Naturwissenschaften* **33**, 166-175

(1946).

16 Elf, J. & Barkefors, I. Single-Molecule Kinetics in Living Cells. *Annual Review of*

Biochemistry **88**, 635-659, doi:10.1146/annurev-biochem-013118-110801 (2019).

17 Karslake, J. D. *et al.* SMAUG: Analyzing single-molecule tracks with nonparametric

Bayesian statistics. *Methods* **193**, 16-26, doi:10.1016/j.ymeth.2020.03.008 (2021).

18 Hansen, A. S. *et al.* Robust model-based analysis of single-particle tracking

experiments with Spot-On. *Elife* **7**, doi:10.7554/eLife.33125 (2018).

19 Persson, F., Lindén, M., Unoson, C. & Elf, J. Extracting intracellular diffusive states

and transition rates from single-molecule tracking data. *Nature Methods* **10**, 265-

269, doi:10.1038/nmeth.2367 (2013).

20 Monnier, N. *et al.* Inferring transient particle transport dynamics in live cells. *Nat*

Methods **12**, 838-840, doi:10.1038/nmeth.3483 (2015).

21 Chen, T. Y. *et al.* Quantifying Multistate Cytoplasmic Molecular Diffusion in Bacterial

Cells via Inverse Transform of Confined Displacement Distribution. *The journal of*

physical chemistry. B **119**, 14451-14459, doi:10.1021/acs.jpcb.5b08654 (2015).

22 Michalet, X. & Berglund, A. J. Optimal diffusion coefficient estimation in single-

particle tracking. *Physical Review E* **85**, doi:10.1103/PhysRevE.85.061916 (2012).

23 Rocha, J., Corbitt, J., Yan, T., Richardson, C. & Gahlmann, A. Resolving Cytosolic

Diffusive States in Bacteria by Single-Molecule Tracking. *Biophysical journal* **116**,

1970-1983, doi:https://doi.org/10.1016/j.bpj.2019.03.039 (2019).

24   Rocha, J. M. *et al.* Single-molecule tracking in live Yersinia enterocolitica reveals

distinct cytosolic complexes of injectisome subunits. *Integrative Biology* **10**, 502-

515, doi:10.1039/c8ib00075a (2018).

25   Prindle, J. R., Rocha, J., Wang, Y., Diepold, A. & Gahlmann, A. Distinct complexes

containing the cytosolic type III secretion system ATPase resolved by 3D single-

molecule tracking in live Yersinia enterocolitica. *bioRxiv*,

2022.2004.2025.488798, doi:10.1101/2022.04.25.488798 (2022).

26   Gahlmann, A. & Moerner, W. E. Exploring bacterial cell biology with single-molecule

tracking and super-resolution imaging. *Nature Reviews Microbiology* **12**, 9-22,

doi:10.1038/nrmicro3154 (2014).

27   Au - Uphoff, S., Au - Sherratt, D. J. & Au - Kapanidis, A. N. Visualizing Protein-DNA

Interactions in Live Bacterial Cells Using Photoactivated Single-molecule Tracking.

JoVE, e51177, doi:doi:10.3791/51177 (2014).

28   Badrinarayanan, A., Reyes-Lamothe, R., Uphoff, S., Leake, M. C. & Sherratt, D. J. *In*

Vivo Architecture and Action of Bacterial Structural Maintenance of Chromosome

Proteins. *Science* **338**, 528-531, doi:10.1126/science.1227126 (2012).

29   Stracy, M. *et al.* Single-molecule imaging of UvrA and UvrB recruitment to DNA

lesions in living Escherichia coli. *Nature communications* **7**, 12568,

doi:10.1038/ncomms12568 (2016).

30   Uphoff, S. *et al.* Stochastic activation of a DNA damage response causes cell-to-cell

mutation rate variation. *Science* **351**, 1094-1097, doi:10.1126/science.aac9786

(2016).

31   Uphoff, S., Reyes-Lamothe, R., Garza de Leon, F., Sherratt, D. J. & Kapanidis, A. N.

Single-molecule DNA repair in live bacteria. *Proceedings of the National Academy of*

Sciences of the United States of America **110**, 8063-8068,

doi:10.1073/pnas.1301804110 (2013).

32   Liao, Y., Schroeder, J. W., Gao, B., Simmons, L. A. & Biteen, J. S. Single-molecule

motions and interactions in live cells reveal target search dynamics in mismatch

repair. *Proc Natl Acad Sci U S A* **112**, E6898-6906, doi:10.1073/pnas.1507386112

(2015).

33   Liao, Y., Li, Y., Schroeder, J. W., Simmons, L. A. & Biteen, J. S. Single-Molecule DNA

Polymerase Dynamics at a Bacterial Replisome in Live Cells. *Biophysical journal* **111**,

2562-2569, doi:10.1016/j.bpj.2016.11.006 (2016).

34   Calkins, A. L. *et al.* Independent Promoter Recognition by TcpP Precedes

Cooperative Promoter Activation by TcpP and ToxR. *mBio* **12**, e0221321,

doi:10.1128/mBio.02213-21 (2021).

35   Li, Y., Schroeder, J. W., Simmons, L. A. & Biteen, J. S. Visualizing bacterial DNA

replication and repair with molecular resolution. *Curr Opin Microbiol* **43**, 38-45,

doi:10.1016/j.mib.2017.11.009 (2017).

36   Li, Y., Chen, Z., Matthews, L. A., Simmons, L. A. & Biteen, J. S. Dynamic Exchange of

Two Essential DNA Polymerases during Replication and after Fork Arrest.

Biophysical journal **116**, 684-693, doi:10.1016/j.bpj.2019.01.008 (2019).

37   Peters, J. M. *et al.* A Comprehensive, CRISPR-based Functional Analysis of Essential

Genes in Bacteria. *Cell* **165**, 1493-1506, doi:10.1016/j.cell.2016.05.003 (2016).

38   Ye, Y.-N., Hua, Z.-G., Huang, J., Rao, N. & Guo, F.-B. CEG: a database of essential gene

708   clusters. *BMC Genomics* **14**, 769, doi:10.1186/1471-2164-14-769 (2013).

709   39   Robinson, M. S. & Hirst, J. Rapid inactivation of proteins by knocksideways. *Curr*

710   *Protoc Cell Biol* **61**, 15 20 11-17, doi:10.1002/0471143030.cb1520s61 (2013).

711   40   Benedetti, L. *et al.* Light-activated protein interaction with high spatial subcellular

712   confinement. *Proc Natl Acad Sci U S A* **115**, E2238-E2245,

713   doi:10.1073/pnas.1713845115 (2018).

714   41   Guntas, G. *et al.* Engineering an improved light-induced dimer (iLID) for controlling

715   the localization and activity of signaling proteins. *Proc Natl Acad Sci U S A* **112**, 112-

716   117, doi:10.1073/pnas.1417910112 (2015).

717   42   Zimmerman, S. P. *et al.* Tuning the Binding Affinities and Reversion Kinetics of a

718   Light Inducible Dimer Allows Control of Transmembrane Protein Localization.

719   *Biochemistry* **55**, 5264-5271, doi:10.1021/acs.biochem.6b00529 (2016).

720   43   Lindner, F., Gahlmann, A. & Diepold, A. Optogenetic control of protein translocation:

721   Protein secretion and translocation into eukaryotic cells with high spatial and

722   temporal resolution by light-controlled activation of the bacterial type III secretion

723   system. European Patent Application 19166308 patent (2019).

724   44   Harper, S. M. Structural Basis of a Phototropin Light Switch. *Science (American*

725   *Association for the Advancement of Science)* **301**, 1541-1544.

726   45   Zayner, J. P., Antoniou, C. & Sosnick, T. R. The Amino-Terminal Helix Modulates

727   Light-Activated Conformational Changes in AsLOV2. *J. Mol. Biol.* **419**, 61-74,

728   doi:10.1016/j.jmb.2012.02.037 (2012).

46 Kramer, M. M., Lataster, L., Weber, W. & Radziwill, G. Optogenetic Approaches for

the Spatiotemporal Control of Signal Transduction Pathways. *International Journal*

of Molecular Sciences **22**, 5300, doi:10.3390/ijms22105300 (2021).

47 Wittmann, T., Dema, A. & Van Haren, J. Lights, cytoskeleton, action: Optogenetic

control of cell dynamics. *Current Opinion in Cell Biology* **66**, 1-10,

doi:10.1016/j.ceb.2020.03.003 (2020).

48 De Geyter, J., Smets, D., Karamanou, S. & Economou, A. 337-366 (Springer

International Publishing, 2019).

49 Cline, K. Mechanistic Aspects of Folded Protein Transport by the Twin Arginine

Translocase (Tat). *Journal of Biological Chemistry* **290**, 16530-16538,

doi:10.1074/jbc.r114.626820 (2015).

50 Lindner, F., Milne-Davies, B., Langenfeld, K., Stiewe, T. & Diepold, A. LITESEC-T3SS -

Light-controlled protein delivery into eukaryotic cells with high spatial and

temporal resolution. *Nature communications* **11**, 2381, doi:10.1038/s41467-020-

16169-w (2020).

51 Grimm, J. B. *et al.* A General Method to Improve Fluorophores Using Deuterated

Auxochromes. *JACS Au* **1**, 690-696, doi:10.1021/jacsau.1c00006 (2021).

52 Wang, Z., Simoncelli, E. P. & Bovik, A. C. (IEEE, 2003).

53 Elf, J. & Barkefors, I. Single-Molecule Kinetics in Living Cells. *Annu Rev Biochem* **88**,

635-659, doi:10.1146/annurev-biochem-013118-110801 (2019).

54 Swartz, T. E. *et al.* The Photocycle of a Flavin-binding Domain of the Blue Light

Photoreceptor Phototropin. *Journal of Biological Chemistry* **276**, 36493-36500,

doi:10.1074/jbc.m103114200 (2001).

55   Nijenhuis, W., Van Grinsven, M. M. P. & Kapitein, L. C. An optimized toolbox for the

753   optogenetic control of intracellular transport. *J. Cell Biol.* **219**,

754   doi:10.1083/jcb.201907149 (2020).

755   56   García-Fruitós, E. *et al.* Aggregation as bacterial inclusion bodies does not imply

756   inactivation of enzymes and fluorescent proteins. *Microbial Cell Factories* **4**, 27,

757   doi:10.1186/1475-2859-4-27 (2005).

758   57   Rinas, U. *et al.* Bacterial Inclusion Bodies: Discovering Their Better Half. *Trends in*

759   *Biochemical Sciences* **42**, 726-737, doi:10.1016/j.tibs.2017.01.005 (2017).

760   58   Bartelt, S. M. *et al.* Dynamic blue light-switchable protein patterns on giant

761   unilamellar vesicles. *Chemical communications* **54**, 948-951,

762   doi:10.1039/c7cc08758f (2018).

763   59   Bartelt, S. M., Steinkühler, J., Dimova, R. & Wegner, S. V. Light-Guided Motility of a

764   Minimal Synthetic Cell. *Nano Letters* **18**, 7268-7274,

765   doi:10.1021/acs.nanolett.8b03469 (2018).

766   60   Paintdakhi, A. *et al.* Oufti: An integrated software package for high-accuracy, high-

767   throughput quantitative microscopy analysis. *Molecular microbiology* **99**, 767-777,

768   doi:10.1111/mmi.13264 (2016).

769   61   Lew, M., Von Diezmann, A. R. S. & Moerner, W. E. Easy-DHPSF open-source software

770   for three-dimensional localization of single molecules with precision beyond the

771   optical diffraction limit. *Protoc. Exch.* (2013).

772   62   Yan, T., Richardson, C. J., Zhang, M. & Gahlmann, A. Computational correction of

773   spatially variant optical aberrations in 3D single-molecule localization microscopy.

774   *Optics express* **27**, 12582-12599 (2019).