Sex-bias in utero alters ovarian reserve but not uterine capacity in female offspring

Annika V Geijer-Simpson[1, 2], Haidee Tinning[1], Tiago H C de Bem[1], Ioannis Tsagakis[1], Alysha S

Taylor[1], Laura Hume[1], Lisa M Collins[2] and Niamh Forde[1*].

[1]Discovery and Translational Sciences Department, Leeds Institute of Cardiovascular and

Metabolic Medicine, Faculty of Medicine and Health, University of Leeds, LS2 9JT, United

Kingdom.

[2] School of Biology, Faculty of Biological Sciences, University of Leeds, Leeds, LS2 9JT.

* Correspondence should be addressed to: n.forde@leeds.ac.uk

ABSTRACT

Environmental stressors to which a foetus is exposed, affect a range of physiological functions in post-natal offspring. Such stressors include disproportionate steroid hormone concentrations in the uterine environment. We aimed to determine the *in-utero* effect of steroid hormones on reproductive potential of female offspring using a porcine model. Hypothesising that an *in-utero* sex bias will influence ovarian reserve and endometrial morphology in the breeding gilt. Reproductive tracts of pigs from female-biased litters (>65% female, n=15), non-biased litters (45-54.9% female, n=15), and male-biased litters (<35% females, n=9) were collected at slaughter (95-115 kg). Ovaries and uterine horns were processed for histological approaches and stained using H&E or IHC techniques. All measurements were conducted in QuPath (Bankhead et al, 2017). Variability of data within groups was analysed with a Levenes test, whilst data was analysed using linear models in R. In the ovarian reserve, there was a significant interaction between the birth weight and the sex ratio of a litter from which a pig originated (p=.015), with low-birth-weight pigs from male-biased litters having a higher number of primordial follicles and the opposite trend seen in pigs from female-biased litters. This was not reflected in recruited, nor atretic follicles. In the uterine horn sex bias held no effect on development as seen in this study. Birth weight held more effects on the gilts. A lower BW decreased the proportion of glands found in the endometrium (p=.045). BW was found to be far more variable in both male-biased and female-biased litters (p=.026). The variability of primordial follicles from male-biased litters was greater than non- and female-biased litters (p=.014). Similarly, endometrial stromal nuclei had a greater range in male- and female-biased litters than non-biased litters (p=.028). There was a greater effect on both ovarian reserve and uterine development of piglet BW than the litter bias. There seems a benefit of being androgenised on ovarian reserve whilst no effects were found for the morphology or endometrial gland proliferation of the uterine horns. However, a crucial finding was in the variability of the data. Both primordial follicles in the male-biased ovary, and stromal nuclei in the male- and female- biased uterine horns had a wider spread in numbers than non-biased litters. This could be inflating the variability of reproductive success seen in animals form male-biased litters by two means. Firstly, by a higher likelihood of insufficient primordial pools. Secondly, through a potential impact on stromal-derived growth factors or insufficient support of the underlying implantation

structures, leading to an increased variability in uterine implantation capabilities, and thus

survival of the embryo.

INTRODUCTION

The Developmental Origins of Health and Disease (DOHaD) describes how the developmental

plasticity and *in-utero* programming of offspring could contribute to susceptibility of a range

of adult disease with the maternal *intra-uterine* hormonal environment has been shown to

influence many aspects of offspring development, some of which are sexually dimorphic in

nature [1]. Studies specifically investigating the influence of foetal hormones have indicated

effects such as endocrine alterations [2], altered physiological development such as delayed

lung maturation [3], behavioural changes such as increased aggression [4], and reduced

maternal and paternal behaviour [5]. These effects are observed following an

androgenisation of the uterine environment in litter bearing species, often caused by

disproportionate numbers of males *in-utero*. This may be a consequence of either; females

positioned between males (which is proportionately more likely to happen in male-biased

litters) or; the overall proportion of males excreting testosterone from the point of sexual

differentiation [6]. A biased litter, one that skews toward a predominantly androgenised or

oestrogenised environment (those consisting of >60% of one sex), occur frequently both

within commercial production systems and among wild pig populations, where proportions

of 1.3:1 males to females per litter have been reported [7]. In the commercial pig a sex-biased

in utero environment hormonal environment has been shown to affect several aspects of

reproductive function in the offspring. Specifically, females that originate from male-biased

litters have fewer teats [8], a lower conception rate at first mating [9], increased sensitivity to

gonadotropins [10], and altered LH surge profiles [2], compared to offspring originating from

female-biased litters. The cause of these changes in reproductive efficacy remains unclear.

Arguably, the most vital organisational event in female reproductive development during

gestation is the developing Primordial Germ Cells (PGC). PGCs are originally derived from the

proximal epiblast cells of pre-gastrulating embryos [11], i.e. prior to generation of the three

primary germ layers. In pigs these have been identified in the dorsal mesentery at embryonic

day (ED)18-20 which then migrate to colonise forming the genital ridge at ED23-24 [12]. It is

established that the PGC's undergo epigenetic reprogramming over a period of several weeks,

beginning at ED12 [13]. In the pig, meiosis, initiation, and formation of primordial follicles

begins at E48 and continues until 25 days post parturition [14]. Differentiation of the Wolffian

duct in the male begins at ED26, at which point secretion of testosterone begins [15]. Hence,

laying down and formation of the primordial follicles in female offspring occurs once

testosterone from male littermates is present in the uterine environment. The point at which

the number of PGCs peaks is determined by morphogenesis and germ cell dynamics, and it is

still not understood how the ovarian reserve is maintained and what triggers activation of

folliculogenesis occurs [16]. It is clear in the pig that there is an effect on ovulation patterns.

However, it is currently unclear whether an androgenised uterine environment affects PGC

formation or follicular recruitment of the PGCs in the pig.

Data from other species indicates that an androgenised uterine environment effects the

development PGC establishment and subsequent follicular recruitment, but also gross

morphology of reproductive tracts. The ovulation rate of an individual does impose a clear

and direct limitation on potential litter sizes in the pig. However, despite ovulation rates

having improved in pigs via selective pressure this hasn't been reflected in any significant

increases to litter sizes [17]. With ovulation and conception rates in the breeding sow being

greater than 95% [18], the major limiting factor to maximized litter sizes is embryonic death.

Only 30-50% of fertilised ova survive through gestation [17]. This loss is predominantly seen

during the pre- and peri- implantation period defined as ED12-18 [17]–[19]. This coincides

with conceptus elongation, synthesis and release of oestrogen for maternal pregnancy

recognition, and trophectoderm differentiation, which is followed by foetal and epithelial

attachment [20]. There is then a secondary wave of embryonic loss at ED 30-40 due to

crowded *in-utero* conditions [21]. It is therefore clear that litter sizes, and conceptus survival,

is contingent on uterine capacity. This is determined by three main factors; uterine length,

uterine blood flow, and uterine gland development [21]. The uterine capacity of a pig is critical

it as it contributes to the reproductive potential of the female i.e. even if there are large

numbers of developmentally competent embryos present there is a physical limit on the

numbers that can implant. It is vital that the uterine tissues can recognise and respond to

maternal and conceptus signals crucial for a successful and established pregnancy [21]–[23].

Not only is this vital for pregnancy recognition, but the uterine capacity will define the

environment in which foetal development occurs [22]. The functional layer of the uterus that
is especially crucial for successful pregnancy outcomes, the endometrium, may be influenced
by an androgenised litter. The endometrium is a heterogenous tissue comprised of several
different cell types including various secretory cells such as the luminal and glandular
epithelium [19]. The endometrial surface of the pig is folded, which at conceptus attachment
(ED14) has a conceptus localised increase in endometrial surface folding [24]. The opposite
folds interlock, reducing the luminal space throughout the pregnancy, maximising luminal
epithelial and foetal contact [19], this is crucial to facilitate maternal and foetal
communication. Endometrial glands along with the luminal epithelium secret uterine luminal
fluid, a complex array of proteins and related substances [25]. The uterine luminal fluid is
critical for endometrial function and conceptus survival as they contain enzymes, growth
factors, cytokines, nutrients, transport proteins, and other regulatory molecules (See review
by Bazer and Johnson 2014 – "Pig blastocyst-uterine interactions") [26]. Maternal
endometrial gland hyperplasia and hypertrophy is extensive during gestation [27], [28] with
large amounts of granular, acid phosphate-positive material within the glands, indicative of a
high level of secretory activity [23]. Sheep with blocked uterine horn gland development
(UGKO – Sheep Uterine Gland Knock Out) indicate a failure of conceptus elongation at ED14
and are rendered infertile (11). Evidencing the fact that uterine glands are critical mediators
for the uterine ability to support a successful pregnancy. Formation of the uterine glands
occur in the neonatal piglet by branching and budding of the luminal epithelium [23], reaching
histoarchitectural maturity by 120 days of age [27]. Although the glands form neonatally, the
histogenesis of the initial uterine horn development, and luminal epithelium, takes place *in-*
utero [23], [30] making this process vulnerable to environmental effects of which the foetus
is exposed to. The understanding of the impact disruptions of uterine development may hold
on endometrial structure and function in adult mammals is crucial for unravelling the high
rates of peri-implantation embryonic loss in both livestock and humans [23], as this may
render the uterus unable to support especially small for gestational age individuals. For
example, adult cows exposed to progesterone and oestradiol benzoate *in-utero* demonstrate
a reduced number of endometrial glands [31]–[33]. Furthermore, in pigs, neonatal
progesterone treatment initially accelerated gland development, but reduced adult glandular
development [34]. Impairment, as described above, has been indicative of a reduced fertile
capacity [30].

This study aims to investigate how a sex biased *in-utero* environment may influence the

development of both the follicular pool, and endometrial glands; both critical components for

successful pregnancy. To do this we investigated the influence of *in-utero* sex ratio a female

was gestated in on the (i) follicular patterns in the pig, (ii) uterine morphology and

endometrial gland proliferation. This was achieved through investigation of the two following

objectives: (i) Investigation of how the established primordial germ cell pool is affected by

different *in-utero* hormonal biases and identification of any differences in follicular

recruitment, or follicular atresia profiles dependent on litter sex bias, (ii) effects the

development of gross uterine structures and level of proliferation within the uterus.

MATERIAL AND METHODS

After consultation with the University of Leeds Animal Welfare and Ethical Review Committee

no ethical approval was sought. It was deemed that due to no interventions taking place, and

the pigs remaining fully under the commercial management, ethics was not necessary. This

complies with the 3Rs strategy of utilising animals which are already part of a process.

ANIMAL MODEL

Our model recruited individual female offspring at birth that were retrospectively assigned to

one of three experimental groups. Females from 1) female-biased *in utero* litter (>65%

females), 2) non-biased (45.9-55% females), or 3) male-biased (<35% females) litters. Only

litters with 10 or more live-born piglets were included in the study and mummified piglets

were not included as accurate identification of sex was unreliable.

All animals were reared in a closed indoor system at the National Pig Centre on partially

slatted floor systems. The pigs used were either not part of any other trials or used as control

pigs in other dietary studies. These female pigs (JSR Large White x Landrace females JT dam-

line x JSR Pietrain-based Geneconverter 900 sire-line) were tracked through the production

system using RFID ear-tags. The pigs were weighed weekly in the month leading up to

slaughter in order to optimise an accurate estimate of slaughter weight. At commercial

slaughter weight (95-115 kg) pigs were delivered to the abattoir weekly over three collection

time points per batch, a total of 10 time points.

TISSUE COLLECTION AND DISSECTION

To control for within-litter variability, two females were selected at random (using
randomizer.org) for reproductive tract collection at slaughter from each litter. Reproductive
tract collections occurred over three periods in November 2018, June-August 2019, and July-
August 2020. The tracts were collected on the abattoir line and transported to the laboratory
on ice within 1.5 hours for tissue processing. The ovaries were dissected from the
reproductive tracts and individually weighed. The right ovary was divided in half and placed
in 10% neutral buffered formalin. The uterine horns were dissected away from the connective
tissue and whole cross sections from 1/3 of the way down from the tip of the uterine horn
(UH; Figure 1a) were also fixed in formalin for forty-eight hours. Ovaries and UH were
removed from formalin and ovaries further divided in half lengthways through the cortex,
resulting in a quartered ovary (**Error! Reference source not found.**b). UH were and ovary were
subsequently dehydrated through a series of ethanol washes and imbedded in paraffin for
histological examination. In total the reproductive tracts from 39 piglets were assessed; male-
biased = 9, non-biased = 15, and female-biased = 15. Unless otherwise stated all chemical
were obtained from Thermo Fisher Scientific, Waltham, US.

HISTOMORPHOMETRY

HEMATOXYLIN AND EOSIN STAIN

For histomorphometry of the ovary, 8 µm serial sections of the embedded ovary from an
individual were mounted on polysine-coated Microslides (VWR International, Radnor, US).
For UH two individual slides containing whole cross sections of UH per animal were assessed.
Slides were de-paraffinised three times in Histo-clear for 10 min. Slides were then hydrated
decreasing concentrations of ethanol (3 x 100% EtOH, 2 x 95% EtOH, and once in 70% EtOH).
All slides were finally rinsed in three times in tap water. Slides were incubated in hematoxylin
(Sigma Aldrich, St. Louis, US - 7g/L) for 5 min, slides rinsed in 3 x in tap water, dipped in 0.25%
acid alcohol for 5 sec and immersed in cold water. At this stage stain intensity was checked
under a microscope before proceeding. Stain was blued in hot tap water for 1 minute and
then rinsed 3 x in cold tap water. Slides were then placed in a Mordant of 95% EtOH. Slides
were then stained in Eosin for 10 min, and dehydrated through 2 changes of 95% EtOH, and
3 changes of 100% EtOH. Finally, slides were places in Histo-clear for 10 min x 3. Ten serial
sections per ovary per animal, (each 160 µm apart), were selected with the initial section

being closest to the ovarian cortex, (**Error! Reference source not found.**). As the distance

between sections a full cross section of the oocyte, <110 µm [35], could not be fully present

in two analysed sections, ensuring that no included follicle would be counted twice. For the

one section per UH of each individual were also stained.

IMMUNOHISTOCHEMISTRY – PROLIFERATING CELL NUCLEAR ANTIGEN

For investigating uterine gland proliferation one section per uterine horn were stained using

an immunohistochemistry (IHC) technique using the Vector Lab VectaStain Elite ABC-HRP kit.

Overnight incubations (4°C) of the sections were carried out with mouse monoclonal anti-

proliferating cell nuclear antigen (PCNA) (1:200, Invitrogen). Control sections were incubated

with a mouse IgG Isotype Control (1:200, Invitrogen). The sections were then incubated with

the secondary antibody and ABC reagent previously described for 60 min in a humidity

chamber. Development of sections was carried out using diaminobenzidine substrate

(ThermoFisher).

IMAGE ANALYSES

OVARIAN RESERVE

To assess the ovarian reserve the number of primordial, pre-antral, antral and atretic follicles

were counted and classified according to Almeida and colleagues [36] (Figure 1). Primordial

follicles (Figure 1– A) were identified as an intact oocyte surrounded by a single layer of

squamous (pre) granulosa cells. An enlarged oocyte that is surrounded by a single, or

multiple, layers of cuboidal granulosa cells was classified as pre-antral (Figure 1- B). Once the

follicle had developed a clear antral cavity that was the same size as, or larger than, the oocyte

it was classified as an antral follicle (Figure 1– C). The antral follicles have several layers of

granulosa cells and have a well-developed thecal layer. All the above follicle types had intact

oocytes with no signs of apoptosis or degradation as described by Almeida and colleagues

[36]/ If degenerative changes had occurred including reduction of the oocyte or condensation

of the nuclear chromatin, or changes to the antral cavity such as scattered granulosa cells, the

follicle was identified as atretic (Figure 1- D).

GROSS UTERINE MORPHOLOGY AND CELL PROLIFERATION

To assess endometrial capacity, we measured gross morphological structures. As a proxy

measure of uterine support we analysed the ratio of secretory cells (luminal and glandular

epithelia) to stromal cell area. The surface area of the endometriumper section was measured

(cm^2) along with the perimeter of the luminal gap (cm^2). Manual counts were made of the

glands in the endometrium of total sections, with the number of larger glands also counted.

An automated cell nuclei detection within QuPath was used to count the stromal cells within

an area of 20,000μm^2. These structures are visualised in Figure 3. Proliferative capacity of the

endometrium was measured using the PCNA IHC stain. An automated DAB stain analysis was

used for the entire sections of the uterine horns to measure stain intensity. This allowed for

identification of cells with a negative or positive stain, along with division of positively stained

cells into a low, moderate, or high stain intensity. Parameters used for the detection were

refined using three sections previously stained in the stain optimisation process.   The

parameters used for stain detection were as follows; pixel size, 0.5μm with thresholds being

Low; 0.1-0.3, Moderate; 0.3-0.5, High; 0.8-1.

DATA PROCESSING AND ANALYSIS

All statistical analyses were performed in RStudio [37] using *lme4* [38], and data was plotted

using PRISM (GraphPad Prism version 9.0.0 for MacOS, GraphPad Software, San Diego,

California USA, www.graphpad.com). Normality was assessed using appropriate tests as

detailed below, alongside histograms, and QQ-plots. Data considered to fit a Gamma

distribution were tested using **"gamma_test"** in package "**goft**". Collinearity between

predictor values was checked using the "**vif**" function in R package "**car**". Ovary weight (g)

and slaughter age (days) was removed due to collinearity (>3). Critical alpha level was applied

as p=0.05. Akaike Information Criterion (AIC) model selection was used to distinguish

between a set of possible models, each describing the relationship between the predictor

variables. The AIC of the final model used for each analyses is detailed in table 1 and 2.

Transformations were required for certain variables to fit their distributions. This was with

either a square root transformation, function "sqrt", or a log transformation, function "log"

and is detailed in table 1 and 2.

The models used were one of the following (specified per response variable in tables 1 and

2);

Models 1- *<Response variable>* per cm^2 with predictor variables being *litter sex ratio* and *birth*
weight as multiplicative, *slaughter weight* as additive, and *litter (ovary) or animal ID*
(UH) as random effect.

Models 2 - *<Response variable>* per cm^2 with predictor variables *litter* as random effect, *litter*
sex ratio, slaughter weight as additive, and *litter (ovary) or animal ID (UH)* as random
effect.

To investigate whether there was more variability within our investigated response variables
between pigs of different *in utero* sex ratios, variance of data points was measured using
"leveneTest" package.

OVARIAN RESERVE

Data was analysed in two ways. Firstly, looking at follicle numbers in the ovary as a whole;
and secondly controlling for variations in the manual dissection of the ovary by investigating
the number of follicles per cm^2 of observed tissue. As results were reflective of each other,
we have only reported the results per cm^2 for reader ease.

Gamma regression models were used to test the effect of predictor values on the following
response variables; (I) primordial follicle count, (II) recruited follicle count, (III) atretic follicle
count, and (IV) total follicular count. All predictor and response variables are described in
Table 1.

GROSS UTERINE MORPHOLOGY AND PROLIFERATION

Analyses of data was performed to investigate difference of <u>rudimentary morphology</u>.
<u>Proportional analyses</u> were conducted to investigate whether there was a difference in the
proportion of secretory structures between bias litters. The following comparisons were
made; luminal perimeter, stromal cells and endometrial glands in relation to endometrial
area. It was then important to compare whether the proportion of these secretory cells
differed in proportion to each other between individuals from different *in-utero* sex ratios.
These proportional comparisons are all detailed in Table 2. The IHC stained cells were
analysed in the same manner as for the morphological analyses. However, low stain intensity

was found to hold a normal distribution and was analysed using model 2, and high stain

intensity required a square root transformation. This is detailed in table 2.

RESULTS

In total 39 pigs were used for analysis, 15 females from female-biased litters, 15 from non-

biased litters, and 9 from male-biased litters. Pigs were excluded from the trial if they failed

to reach commercial slaughter, one pig was excluded once reproductive tracts were collected

due to an active infection of the tract. The mean birth weight and slaughter weight ($\pm$ SD) of

the pigs were: 1) female-biased - 1.65 kg ($\pm$ 0.446) and 107 ($\pm$ 8.140) kg, 2) non-biased - 1.47

($\pm$ 0.207) and 110 ($\pm$ 5.275) kg, and 3) male-biased - 1.51 ($\pm$ 0.303) and 113 ($\pm$ 5.915) kg. No

significant difference was observed.

The ovarian tissue analysed was on average 16.6 ($\pm$ 3.197) cm^2 in female-biased pigs, 17.8 ($\pm$

4.386) cm^2 in non-biased pigs, and 15.0 ($\pm$ 3.198) cm^2 in male-biased pigs.

VARIABILITY OF REPRODUCTIVE PARAMETERS IN INDIVIDUALS.

The variability of data points between the sex ratio groups were analysed as visualised in

Figure 5. This revealed a bigger range in birth weights (F-value=4.073, df=35, p=.026) of

females from both female- and male-biased litters compared to non-biased litters. There was

a significantly larger variance of both primordial (F-value=4.801, df=36, p=.014) and total (F-

value=5.381, df=36, p=.009) follicle numbers in females from male-biased compared to those

that originated from non- or female-biased litters. This also revealed that there was an

increased variability in the number of stromal cells in endometria recovered from females

from sex-biased litters compared to non-biased (F-value=3.8134, df=71, p=.027). The

variability did not differ between litter sex bias in endometrial area (F-value=.2472, df=71,

p=.7817), luminal perimeter (F-value=2.3527, df=71, p=.1025), nor total gland (F-

value=2.4973, df=71, p=.08951) or large endometrial gland counts (F-value=2.0419, df=71,

p=.1373).

INTERACTION OF BIRTHWEIGHT AND SLAUGHTER WEIGHT

The following results from the mixed models is summarised in Table 3.

FOLLICULAR COUNTS

For an individual there was no significant interaction with *in-utero* sex ratio of that piglet's

litter and their birthweight for the number of recruited follicles (GLMM; t-value=.789, n=34,

p=.436). There was a significant interaction in primordial (Gamma GLM; t-value=-2.637, n=34,

p=.008) and total (Gamma GLM; t-value=-2.754, n=34, p=.006) follicle numbers per cm^2

between these two predictor variables i.e. sex ratio *in-utero*. As seen in Figure 6, increasing

birth weight was negatively correlated with primordial and total follicle numbers in females

from male biased litters. Contrary to this, female-biased and non-biased litters showed no

difference in follicle numbers between different weights.

UTERINE MORPHOLOGY AND PROLIFERATING CELLS

There was no interaction observed between the sex ratio and birth weight for endometrial

area (Gamma GLM; z-value=-1.416, n64, p=.157), nor luminal perimeter (Gamma GLM; z

value=-1.595, n=64, p=.1107). No significant interaction was found between birthweight nor

sex ratio for the proportion of positively PCNA-stained nuclei (Gamma GLM; z-value=-1.266,

n=63, p=.206), nuclei with low stain intensity (GLMM; z-value=.713, n=63, p=.482), nuclei with

a moderate stain intensity (Gamma GLM; z-value=-.853, n=63, p=.394), nor the nuclei with a

high stain intensity (Gamma GLM; z-value=-1.533, n=63, p=.1252).

NON-INTERACTIVE EFFECTS OF SEX RATIO, BIRTH WEIGHT, AND SLAUGHTER WEIGHT

FOLLICULAR COUNTS

No significant difference in numbers of recruited (GLMM; t-value=.442, n=34, p=.679), nor

atretic (Gamma GLM; t-value=.817, n=34, p=.414) follicles were observed between sex ratios.

Similarly, both recruited (GLMM; t-value=669, n=34, p=.509) and atretic (Gamma GLM; t-

value=1.775, n=34, p=.0759) follicle numbers were not affected by the birthweight of an

individual. The slaughter weight had no effect on any aspect of follicle count measured

(primordial - Gamma GLM; t-value=860, n=34, p=.589; recruited - GLMM; t-value=.995, n=34,

p=.330; atretic - Gamma GLM; t-value=1.468, n=34, p=.142);, nor total - Gamma GLM; t-

value=2.754, n=34, p=.611).

UTERINE MORPHOLOGY

The sex ratio of the litter from which a pig originated, birth weight, or slaughter weight did

not significantly affect the total cross section of endometrial area (Gamma GLM, n=64; z-

value=.022, p=.825; z-value=1.376, p=.169; and z-value=.787, p=.431; respectively), size of the

luminal area as measured by luminal perimeter (Gamma GLM, n=64; z-value=1.096, p=.273;

z-value=1,915, p=.055; z-value=-.151, p=.880; respectively), nor number of stromal cells as

measured by nuclear staining (Poisson GLM, n=64: z-value=1.096, p=.2731; z-value=-1.915,

p=.0555; z-value=-.151, p=.8799; respectively).

The total endometrial gland numbers (Poisson GLM, n=64; z-value=.117, p=.907; z-

value=.058, p=.954; z-value=.054, p=.957; respectively) and larger glands alone (Poisson GLM,

n=64; z-value=-1.033, p=.302; z-value=.432, p=.666; z-value=-.908, p=.364; respectively) were

not significantly different between individuals of different *in-utero* sex ratios, birth weights,

or slaughter weights.

PROPORTIONAL ANALYSES

The proportion of the secretory structures were then analysed in proportion to the

endometrial area – this was to investigate whether the structures differed between sex ratio

litter individuals in regard to the size of their reproductive tracts. The luminal perimeter (cm^2),

glandular count, and stromal nuclei in relation to the endometrial area was analysed and no

effects were seen for the ratio of luminal perimeter nor stromal cell number nuclei to

endometrial area for females from different sex ratio litters (Gamma GLM, n=64; z-

value=.874, p=.382; z-value=.355, p=.723; respectively), birthweight (Gamma GLM, n=64; z-

value=-.142, p=.887; z-value=.941, p=.347; respectively), nor slaughter weight (Gamma GLM,

n=64; z-value=-1.397, p=.162; z-value=-1.315, p=.188; respectively). However, although the

ratio of the total number of glands in the endometrial area for females from different sex

ratios (Gamma GLM; z-value=-0.423, n=64, p=.67259), and slaughter weights (Gamma GLM;

z-value=-1.427, n=64, p=.15349) were not significantly different, there was a significant

difference in effect of the birth weight (Gamma GLM; z-value=2.005, n=64, p=.04492), Figure

7.

The analyses investigating the ratio of different secretory tissues against each other within

the uterine horn demonstrated no significant differences in ratios between stromal nuclei and

luminal perimeter (Gamma GLM, n=64; Sex ratio - z-value=-0.416, p=.6674; Birthweight - z-

value=1.919, p=.0550; Slaughter weight - z-value=-.912, p=.3619), gland counts and luminal

perimeter (Gamma GLM, n=64; Sex ratio - z-value=-1.315, p=.1884; Birthweight - z-

value=1.709, p=.0874; Slaughter weight - z-value=.045, p=.9642), nor gland counts and

stromal nuclei (Gamma GLM, n=64; Sex ratio - z-value=-.234, p=.815; Birthweight - z-value=-

.634, p=.526; Slaughter weight - z-value=.767, p=.443).

The sex ratio of the litter from which a pig originated, birth weight and slaughter weight did
not show a significant effect on the proportion of PCNA-positive stained cells (Gamma GLM,
n=63; z-value=.010, p=.992; z-value=.902, p=.367; z-value=.965, p=.335; respectively), nuclei
with a low stain intensity (GLMM, n=63; z-value=.225, p=.800; z-value=-.469, p=.619; z-
value=.020, p=.984; respectively), moderate stain intensity (Gamma GLM, n=63; z-value=-
.497, p=.619; z-value=1.056, p=.291; z-value=-.629, p=.529; respectively), and high stain
intensity (Gamma GLM, n=63; z-value=,204, p=.839; z-value=,737, p=.461; z-value=-.453,
p=.651; respectively).

DISCUSSION

The aim of this study was to investigate how gestation of individuals in a sex biased *in utero*
environment altered development and reproductive potential of the female offspring. We
hypothesised that gestation of a female in a predominantly male or female in utero
environment would alter primordial follicle pool and the development of the uterine capacity.

Irrespective of whether the follicle count was assessed on a per ovary or tissue size basis the
outcome was the same. We found that the primordial follicle pool and total follicle count in
the ovary was significantly more variable when females originated from a male biased uterine
environment. We also found that the follicle counts were affected by the uterine environment
differently by those from a male biased vs a female biased uterine environment, based on
their birth weight. Previous research in species including mice and pigs suggest an important
role for androgens on follicular and CL development. In gilts this is illustrated by increased
ovulation rates following dihydrotestosterone treatment [39], and CL dysfunction that is
marked by decreased progesterone production when flutamide was used to block androgenic
actions [40]. Overall, this study found no effects of sex ratio on the non-ovulatory recruitment,
nor atresia of follicles, suggesting that an androgenised uterine environment does not
interfere with non-cyclic folliculogenesis, nor the breakdown of follicles in the pre-pubertal
commercial pig. Androgens in sheep were found to increase follicular recruitment when
offspring were in a hyper-androgenised maternal circulation, resulting in detrimental effects
such as multi-follicular ovaries (similar to PCOS) and early cessation of cyclicity exposed to
hyper-androgenised maternal circulation in sheep [41]. However, it is important to note that

sheep normally only bear 1-3 offspring per gestation and hence aren't litter bearing. Thereby,

they are less likely to be exposed to a biased uterine environment than in the pig, potentially

leading to a higher sensitivity to hyper-androgenisation of the uterine environment. This may

contribute to the species-specificity observed on androgenisation….

Despite no effects seen in the number of follicles recruited females from male-biased litters

had a higher count of primordial and total follicles both per cm^2 when the individual pig held

a low birth weight. However, females with a higher birth weight were found to hold a higher

number of primordial and total follicles if from female-biased litters. This suggests that the

effect of an androgenised environment and an oestrogenised uterine environment seem to

have opposite influences on the development of the primordial follicle pool. This is further

affected by the birthweight of the particular individual. An androgenised uterine environment

may increase primordial germ cell proliferation, resulting in a larger TOR. A larger TOR would

be in accordance with Seyfang and colleagues [42] who found that androgenised female pigs

were more likely to ovulate and had higher CL counts when from a male-biased compared to

female-biased litters when treated with gonadotrophins at 18 weeks of age. This could be due

to in a higher TOR. Research in non-litter bearing species, who will hold different timings of

PGC establishment, conflicts with that in pigs; findings in sheep in which individuals from a

hyper-androgenised uterine environment displayed lower TOR's than their control

counterparts [41]. This may be due to the pre-mentioned species differences, or due to the

study design mimicking maternal testosterone via intramuscular testosterone propionate in

pregnant ewes, rather than uterine hyper-androgenisation. This was an unexpected finding,

as piglets that are below 1.3 kg at birth have been found to exhibit less competent post-natal

development and reduced survivability [43], suggesting a detrimental effect of low birth

weight on offspring. Compensatory growth following low birth weights has been found to

lead to delayed puberty onset in mice [44]. Low birthweight piglets have also been found to

grow less than normal and high birthweight piglets throughout their life course [45], and do

not display the "catch-up", that piglets that have been fed restrictively do [46]. Although our

findings suggest a benefit to the TOR in low birthweight piglets, with no effect on the

recruitment nor atresia of follicles, there may be long-term implications as indicated in for

example mice.

Despite there being, to our knowledge, no previous research specifically investigating the

effect of an *in-utero* sex bias on the uterine morphology or efficiency, post-natal uterine gland

development is influenced by lactocrine aspects [47] and oestrogenic influences leading to

reduced uterine responsiveness to embryotropic signals [30]. However, we did not find any

evidence suggesting that gestation of a female in a male-biased uterine environment is

detrimental to uterine development in our study. We did not demonstrate any effect of litter

sex ratio on any of the defined uterine measures, their ratios, or the cell proliferation.

However, it was found that the birth weight of a female pig had a significant effect on the

proportion of glands in the uterus, relative to the uterine size. In this instance the higher the

birth weight, the lower the proportion of glands. This may be due to a larger uterus rather

than specifically lower developed uterine glands. We propose that synchronised cyclic

animals should be used to further investigate whether this effect is having a real effect on the

reproducing gilts and sows.

The most important outcome that we found in this study is the increased variability in

reproductive measures in females originating from male biased litters. In mice, the strain has

shown to account for major variation with regard to follicular profiles [48]. The TOR has also

been found to be greatly variable within individuals of the same species, with reports of up

to 20-fold differences in individuals through the neonatal period and puberty [49], [50]. This

would partially clarify the high variation seen in follicular numbers of pigs from male-biased

litters in our data, however, it is interesting to note that the variation is considerably smaller

in both non- and female-biased litters. There may be an underlying cause of the increased

variation of TOR in male biased litters that hold functional consequences not yet understood.

What is also evident is that the number of stromal cell nuclei within the endometrium was

significantly more variable in females that came from either extreme, male- and female-

biased litters. This variability in females from male- and female- biased litters was also seen

in the birth weight of pigs. This could potentially impact on stromal-derived growth factors

that enhance secretion from the luminal and glandular epithelium [51]. The stromal cells are

also key in supporting the underlying implantation structures and hence [51], variability

within their numbers could lead to variability in uterine implantation capabilities, and thus

survival of the embryo. Variability is known to be a major issue for pig producers for the many

reasons and is commonly found in many different aspects of production. Placental efficiency

was found to be highly variable in the large white at an approximate level of three-fold [52],

with two-fold variation within litters. As indicated in these findings, higher levels of variation

was seen in both male-, and female- biased litters, compared to the non-biased litters. This

suggests that a biased litter may be a contributor to the variability of reproductive output

commonly reported in female pigs. Furthermore, such biased litters were found to lead to

more variation in the birth weight of offspring. Selection for larger litter sizes over time has

resulted in litters of higher numbers but with low and greatly variable birth weights [53]. Low

birthweight piglets from large litters are often cross fostered or euthanized as they will not

be able to compete with their larger siblings for teats and have poor pre-weaning survival

rates [21].

Under the premise of this study pigs were expected to, at slaughter, not yet have become

cyclic. Some pigs may however have reached puberty earlier than the norm, skewing results.

Therefore, this suggests that to truly understand the effects of a biased litter on the

recruitment and atresia of follicles, future work should utilise synchronised gilts, and these

current findings should be cautiously interpreted. There may be an increase in the depletion

of TOR in older pigs, which would normally see a 70% decrease from E50 to 300 days after

birth [54]. Finally, albeit currently a very novel avenue, there may be an effect of an

androgenised uterine environment on the development of oogonial stem cells, if they

functionally exist within the pig. Potentially inhibiting a maintenance of the TOR. However,

the subjective nature of manual quantification of the TOR must be taken into consideration

when interpreting the results from this study. Future research should investigate the TOR of

synchronised, pre-pubertal pigs to further understand the effects of a sex biased litter on the

follicular profile of commercial pigs. Synchronisation would allow for a detailed understanding

of deviating recruitment patterns observed in previous studies, which this experimental

design doesn't hold sensitivity to fully investigate. Further to this study, investigating the

depletion of the ovarian reserve over time would help understand the long-term effects that

a bias may hold on reproductive longevity.

In conclusion, females that originate from a predominantly male-biased litter i.e. an

androgenised uterine environment, and have low birth weight, display increased primordial

ovarian reserve along with increased variability of primordial follicle numbers as compared to

pigs that originate from non- and female-biased litters. Conversely, a higher birth weight

resulted in a greater primordial ovarian reserve if the female pig originated from an oestrogenised uterine environment. Pigs from either litter bias (male or female) were found to have a significantly higher variation in birth weight than if a pig originated from a non-biased litter. These data have implications for reproductive potential of females gestated in sex-biased in utero environments.

ACKNOWLEDGEMENTS

Microscopy was facilitated by the University of Leeds Bioimaging Facility, and we would also like to thank the National Pig Research Centre for their work in relation to this publication. Work in NF's group is supported by N8 agri-food pump priming, QR GCRF, UN, LTHT, as well as BBSRC grant number BB/R017522/1.

REFERENCES

[1]   S. Pérez-Cerezales, P. Ramos-Ibeas, D. Rizos, P. Lonergan, P. Bermejo-Alvarez, and A. Gutiérrez-Adán, "Early sex-dependent differences in response to environmental stress," *Reproduction*, vol. 155, no. 1, pp. R39–R51, 2018.

[2]   J. Seyfang, R. N. Kirkwood, A. J. Tilbrook, and C. R. Ralph, "Sex bias of the birth litter affects surge but not tonic LH secretion in gilts1," *Journal of Animal Science*, vol. 96, no. 6, pp. 2195–2203, Jun. 2018, doi: 10.1093/jas/sky151.

[3]   A. Rodriguez, R. M. Viscardi, and J. S. Torday, "Fetal androgen exposure inhibits fetal rat lung fibroblast lipid uptake and release," *EXPERIMENTAL LUNG RESEARCH*, vol. 27, no. 1, pp. 13–24, 2001, doi: 10.1080/019021401459743.

[4]   J. Seyfang, K. J. Plush, R. N. Kirkwood, A. J. Tilbrook, and C. R. Ralph, "The sex ratio of a litter affects the behaviour of its female pigs until at least 16 weeks of age," *APPLIED ANIMAL BEHAVIOUR SCIENCE*, vol. 200, pp. 45–50, Mar. 2018, doi: 10.1016/j.applanim.2017.12.001.

[5]   S. Ichikawa and Y. Fujii, "Effect of prenatal androgen treatment on maternal behavior in the female rat," *Horm Behav*, vol. 16, no. 2, pp. 224–233, 1982.

[6]   J. J. Ford, R. K. Christenson, and R. R. Maurer, "Serum Testosterone Concentrations in Embryonic and Fetal Pigs During Sexual Differentiation," *Biology of Reproduction*, vol. 23, no. 3, pp. 583–587, Oct. 1980, doi: 10.1095/biolreprod23.3.583.

[7] C. Fonseca, A. A. da Silva, J. Alves, J. Vingada, and A. M. V. M. Soares, "Reproductive
performance of wild boar females in Portugal," *EUROPEAN JOURNAL OF WILDLIFE*
RESEARCH, vol. 57, no. 2, pp. 363–371, Apr. 2011, doi: 10.1007/s10344-010-0441-6.

[8] L. C. Drickamer, T. L. Rosenthal, and R. D. Arthur, "Factors affecting the number of teats
in pigs," *Reproduction*, vol. 115, no. 1, pp. 97–100, 1999.

[9] L. C. Drickamer, R. D. Arthur, and T. L. Rosenthal, "Conception failure in swine:
importance of the sex ratio of a female's birth litter and tests of other factors," *J Anim*
Sci, vol. 75, no. 8, pp. 2192–2196, 1997.

[10] J. Seyfang, P. Langendijk, T. Y. Chen, E. Bouwman, and R. N. Kirkwood, "Human
chorionic gonadotrophin in early gestation induces growth of estrogenic ovarian
follicles and improves primiparous sow fertility during summer," *ANIMAL*
REPRODUCTION SCIENCE, vol. 172, pp. 21–25, Sep. 2016, doi:
10.1016/j.anireprosci.2016.06.009.

[11] Q. LEI *et al.*, "Reprogramming of the pig primordial germ cells into pluripotent stem
cells: a brief review," *Frontiers (Boulder)*, vol. 1, 2019.

[12] Y. Takagi, N. C. Talbot, C. E. Rexroad Jr, and V. G. Pursel, "Identification of pig primordial
germ cells by immunocytochemistry and lectin binding," *Molecular Reproduction and*
Development: Incorporating Gamete Research, vol. 46, no. 4, pp. 567–580, 1997.

[13] Q. Zhu *et al.*, "Specification and epigenomic resetting of the pig germline exhibit
conservation with the human lineage," *Cell Rep*, vol. 34, no. 6, p. 108735, 2021.

[14] D. Monniaux *et al.*, "The Ovarian Reserve of Primordial Follicles and the Dynamic
Reserve of Antral Growing Follicles: What Is the Link?1," *Biology of Reproduction*, vol.
90, no. 4, Apr. 2014, doi: 10.1095/biolreprod.113.117077.

[15] L. L. Anderson, "Reproductive biology of pigs," *Iowa State University Animal Industry*
Report, vol. 6, no. 1, 2009.

[16] J. B. Kerr, M. Myers, and R. A. Anderson, "The dynamics of the primordial follicle
reserve," *Reproduction*, vol. 146, no. 6, pp. R205–R215, 2013.

[17] W. F. Pope, "Embryonic mortality in swine," *Embryonic mortality in domestic species*,
vol. 53, p. 77, 1994.

[18] R. D. Geisert and R. A. M. Schmitt, "Early embryonic survival in the pig: Can it be
improved?," *Journal of Animal Science*, vol. 80, no. E-suppl_1, pp. E54–E65, Jan. 2002,
doi: 10.2527/animalsci2002.0021881200800ES10009x.

[19]   H. W. Stroband and T. van der Lende, "Embryonic and uterine development during early pregnancy in pigs," *J Reprod Fertil Suppl*, vol. 40, pp. 261–277, 1990.

[20]   J. W. Ross, M. D. Ashworth, D. R. Stein, O. P. Couture, C. K. Tuggle, and R. D. Geisert, "Identification of differential gene expression during porcine conceptus rapid trophoblastic elongation and attachment to uterine luminal epithelium," *Physiological Genomics*, vol. 36, no. 3, pp. 140–148, 2009.

[21]   J. L. Vallet, A. K. McNeel, G. Johnson, and F. W. Bazer, "TRIENNIAL REPRODUCTION SYMPOSIUM: Limitations in uterine and conceptus physiology that lead to fetal losses," *JOURNAL OF ANIMAL SCIENCE*, vol. 91, no. 7, pp. 3030–3040, Jul. 2013, doi: 10.2527/jas.2012-6138.

[22]   F. F. Bartol, A. A. Wiley, T. E. Spencer, J. L. Vallet, and R. K. Christenson, "Early uterine development in pigs," *JOURNAL OF REPRODUCTION AND FERTILITY-SUPPLEMENT-*, p. 99, 1993.

[23]   C. A. Gray *et al.*, "Developmental Biology of Uterine Glands1," *Biology of Reproduction*, vol. 65, no. 5, pp. 1311–1323, Nov. 2001, doi: 10.1095/biolreprod65.5.1311.

[24]   R. Geisert, M. Lucy, J. Whyte, J. Ross, and D. Mathew, "Cytokines from the pig conceptus: roles in conceptus development in pigs," *J Anim Sci Biotechnol*, vol. 5, p. 51, Nov. 2014, doi: 10.1186/2049-1891-5-51.

[25]   F. W. Bazer, "Uterine protein secretions: relationship to development of the conceptus," *J Anim Sci*, vol. 41, no. 5, pp. 1376–1382, 1975.

[26]   F. W. Bazer and G. A. Johnson, "Pig blastocyst–uterine interactions," *Differentiation*, vol. 87, no. 1–2, pp. 52–65, 2014.

[27]   J. S. Perry and P. R. Crombie, "Ultrastructure of the uterine glands of the pig.," *J Anat*, vol. 134, no. Pt 2, p. 339, 1982.

[28]   F. Sinowatz and A. E. Friess, "Uterine glands of the pig during pregnancy," *Anat Embryol (Berl)*, vol. 166, no. 1, pp. 121–134, 1983.

[29]   C. A. Gray *et al.*, "Endometrial glands are required for preimplantation conceptus elongation and survival," *Biol Reprod*, vol. 64, no. 6, pp. 1608–1613, 2001.

[30]   B. J. Tarleton, T. D. Braden, A. A. Wiley, and F. F. Bartol, "Estrogen-induced disruption of neonatal porcine uterine development alters adult uterine function," *Biology of Reproduction*, vol. 68, no. 4, pp. 1387–1393, 2003.

[31]   F. F. Bartol *et al.*, "Uterine differentiation as a foundation for subsequent fertility.," *J*
595   *Reprod Fertil Suppl*, vol. 54, pp. 287–302, 1999.

596   [32]   J. G. Floyd *et al.*, "Fertility of beef heifers treated from birth with growth promoting
597   implants," in *ASAS Southern Section Abstracts*, 2001, vol. 5.

598   [33]   F. F. Bartol *et al.*, "Neonatal exposure to progesterone and estradiol alters uterine
599   morphology and luminal protein content in adult beef heifers," *Theriogenology*, vol.
600   43, no. 5, pp. 835–844, 1995.

601   [34]   J. L. Vallet, R. K. Christenson, F. F. Bartol, and A. A. Wiley, "Effect of treatment with
602   retinyl palmitate, progesterone, oestradiol and tamoxifen on secretion of a protein
603   similar to retinol-binding protein during uterine gland development in neonatal pigs,"
604   *Reproduction*, vol. 103, no. 1, pp. 189–197, 1995.

605   [35]   D. E. Morbeck, K. L. Esbenshade, W. L. Flowers, and J. H. Britt, "Kinetics of Follicle
606   Growth in the Prepubertal Gilt," *Biology of Reproduction*, vol. 47, no. 3, pp. 485–491,
607   Sep. 1992, doi: 10.1095/biolreprod47.3.485.

608   [36]   F. Almeida, A. L. N. Alvarenga Dias, L. P. Moreira, A. T. L. Fiúza, and H. Chiarini-Garcia,
609   "Ovarian follicle development and genital tract characteristics in different birthweight
610   gilts at 150 days of age," *Reproduction in Domestic Animals*, vol. 52, no. 5, pp. 756–762,
611   2017.

612   [37]   R. C. Team, "R: A language and environment for statistical computing," 2013.

613   [38]   D. Bates, M. Mächler, B. Bolker, and S. Walker, "Fitting linear mixed-effects models
614   using lme4," *arXiv preprint arXiv:1406.5823*, 2014.

615   [39]   H. Cardenas, J. R. Herrick, and W. F. Pope, "Increased ovulation rate in gilts treated with
616   dihydrotestosterone," *REPRODUCTION-CAMBRIDGE-*, vol. 123, no. 4, pp. 527–533,
617   2002.

618   [40]   M. Grzesiak, K. Knapczyk-Stwora, R. E. Ciereszko, A. Golas, I. Wieciech, and M.
619   Slomczynska, "Androgen deficiency during mid-and late pregnancy alters progesterone
620   production and metabolism in the porcine corpus luteum," *Reproductive Sciences*, vol.
621   21, no. 6, pp. 778–790, 2014.

622   [41]   T. Steckler, J. Wang, F. F. Bartol, S. K. Roy, and V. Padmanabhan, "Fetal Programming:
623   Prenatal Testosterone Treatment Causes Intrauterine Growth Retardation, Reduces
624   Ovarian Reserve and Increases Ovarian Follicular Recruitment," *Endocrinology*, vol.
625   146, no. 7, pp. 3185–3193, Jul. 2005, doi: 10.1210/en.2004-1444.

[42] J. Seyfang, C. R. Ralph, A. J. Tilbrook, and R. N. Kirkwood, "Response to gonadotrophins
differs for gilts from female- and male-biased litters," *ANIMAL REPRODUCTION*
SCIENCE, vol. 182, pp. 134–137, Jul. 2017, doi: 10.1016/j.anireprosci.2017.05.012.
[43] C. Rehfeldt *et al.*, "Limited and excess protein intake of pregnant gilts differently affects
body composition and cellularity of skeletal muscle and subcutaneous adipose tissue
of newborn and weanling piglets," *EUROPEAN JOURNAL OF NUTRITION*, vol. 51, no. 2,
pp. 151–165, Mar. 2012, doi: 10.1007/s00394-011-0201-8.
[44] L. S. Monteiro and D. S. Falconer, "Compensatory growth and sexual maturity in mice,"
Animal Science, vol. 8, no. 2, pp. 179–192, 1966, doi: DOI:
10.1017/S0003356100034565.
[45] E. Václavková, P. Daněk, and M. Rozkot, "The influence of piglet birth weight on growth
performance," *Research in pig breeding*, vol. 6, no. 1, pp. 1–3, 2012.
[46] A. M. S. Huting, P. Sakkas, I. Wellock, K. Almond, and I. Kyriazakis, "Once small always
small? To what extent morphometric characteristics and post-weaning starter regime
affect pig lifetime growth performance," *Porcine Health Manag*, vol. 4, no. 1, pp. 1–14,
2018.
[47] D. J. Miller, A. A. Wiley, J. C. Chen, C. A. Bagnell, and F. F. Bartol, "Nursing for 48 Hours
from Birth Supports Porcine Uterine Gland Development and Endometrial Cell
Compartment-Specific Gene Expression1," *Biology of Reproduction*, vol. 88, no. 1, Jan.
2013, doi: 10.1095/biolreprod.112.105056.
[48] J. B. Kerr, R. Duckett, M. Myers, K. L. Britt, T. Mladenovska, and J. K. Findlay,
"Quantification of healthy follicles in the neonatal and adult mouse ovary: evidence for
maintenance of primordial follicle supply," *Reproduction*, vol. 132, no. 1, pp. 95–109,
2006.
[49] M. J. Faddy, "Follicle dynamics during ovarian ageing," *Mol Cell Endocrinol*, vol. 163,
no. 1–2, pp. 43–48, 2000.
[50] W. H. B. Wallace and T. W. Kelsey, "Human ovarian reserve from conception to the
menopause," *PLoS One*, vol. 5, no. 1, p. e8772, 2010.
[51] C. Chen, T. E. Spencer, and F. W. Bazer, "Fibroblast Growth Factor-10: A Stromal
Mediator of Epithelial Functionin the Ovine Uterus," *Biology of Reproduction*, vol. 63,
no. 3, pp. 959–966, Sep. 2000, doi: 10.1095/biolreprod63.3.959.

[52] M. E. Wilson, N. J. Biensen, and S. P. Ford, "Novel insight into the control of litter size in pigs, using placental efficiency as a selection tool," *Journal of Animal Science*, vol. 77, no. 7, pp. 1654–1658, Jul. 1999, doi: 10.2527/1999.7771654x.

[53] E. Zotti *et al.*, "Impact of piglet birthweight and sow parity on mortality rates, growth performance, and carcass traits in pigs," *Revista Brasileira de Zootecnia*, vol. 46, pp. 856–862, 2017.

[54] H. D. Guthrie and W. M. Garrett, "Apoptosis during folliculogenesis in pigs.," *Reprod Suppl*, vol. 58, pp. 17–29, 2001.

Table 1

Table 1. The response variables analysed for the ovary and the specific model, model type,

distribution of the variable, and AIC value of the most fitting model. All analyses were

performed in RStudio and were carried out using *lme4*. Follicle numbers manually counted on

H&E stained histological sections and reproductive tracts were collected from pigs originating

from either female biased (n = 15), non biased (n = 15), or male biased groups (n = 9).

Repsonse variable	Model	Model Type	Distribution	AIC value
Primordial follicles	1	GLM	Gamma	369.0617
Recruited follicles	1	GLMM	Gamma	186.1362
Atretic follicles	2	GLM	Gamma	190.557
Total follicles	1	GLM	Gamma	371.1647

Table 2

Table 2. The response variables analysed for the uterine horn and the specific model, model

type, distribution of the variable, and AIC value of the most fitting model. All analyses were

performed in RStudio and were carried out using *lme4*. Reproductive tracts were collected

from pigs originating from either female biased (n = 15), non biased (n = 15), or male biased

groups (n = 9). All observations were made from histological sections using either H&E stains

or an IHC stain for cell proliferation (PCNA).

Response variable	Model	Model Type	Distribution	Transformation	AIC value
Rudimentary analysis					
Endometrial area (EA)	1	GLM	Gamma		-127.258
Luminal perimeter (LP)	1	GLM	Gamma	Log	-12.6513
Stromal nuclei (SN)	2	GLM	Poisson		575.5295
Gland count (GC)	2	GLM	Poisson		1398.412
Large gland count (LGC)	2	GLM	Poisson		789.2035
IHC analysis					
Positively stained cells	1	GLM	Gamma		-70.7763
Low stain intensity	1	GLMM	Gaussian		-121.477
Moderate stain intensity	1	GLM	Gamma		-218.274
High stain intensity	1	GLM	Gamma	Square root	-149.745
Proportional analysis					

LP: EA	2	GLM	Gamma		1476.293
SN: EA	2	GLM	Gamma		531.1063
GC: EA	2	GLM	Gamma		1013.363
SN: LP	2	GLM	Gamma	Square root	117.1338
GC: LP	2	GLM	Gamma		749.575
GC:SN	2	GLM	Gamma		749.575

695 **Table 3**

696 **Table 3. Mixed model outputs for each response variable.** Outputs from the models run for each response variable including the number of

697 animals in the analyses, specific test statistic, and p-value. Signif. Codes: 0 '***'; 0.001 '**'; 0.01 '*'; 0.05 '.'.

Response variable	N of animals / df	Interaction		Litter sex bias		Birth weight		Slaughter weight	
		Test statistic	p-value	Test statistic	p-value	Test statistic	p-value	Test statistic	p-value
Primordial follicles	34	2.637	**0.008***	NA	NA	NA	NA	0.541	0.589
Recruited follicles	34	0.789	0.436	0.442	0.679	-0.669	0.509	0.995	0.330
Atretic follicles	34	NA	NA	0.817	0.414	-1.775	0.076	1.468	0.142
Total follicles	34	-2.754	**0.006***	NA	NA	NA	NA	-2.754	0.611
Endometrial area (EA)	64	-1.416	0.157	0.022	0.825	-1.376	0.169	0.787	0.431
Luminal perimeter (LP)	64	-1.595	0.1107	1.096	0.2731	-1.915	0.055	-0.151	0.880
Stromal nuclei (SN)	64	NA	NA	-0.287	0.774	-0.916	0.359	1.237	0.216
Gland count (GC)	64	NA	NA	0.117	0.907	0.058	0.954	0.54	0.957

Large gland count (LGC)	64	NA	NA	-1.033	0302	0.432	0.666	-0.908	0.364
LP:EA	64	NA	NA	0.874	0.382	-0.142	0.887	-1.397	0.162
SN:EA	64	NA	NA	0.355	0.723	0.941	0.347	-1.315	0.188
GC:EA	64	NA	NA	-0.423	0.673	2.001	**0.045.**	-1.427	0.153
SN:LP	64	NA	NA	-0.416	0.667	1.919	0.055	-0.912	0.362
GC:LP	64	NA	NA	-1.315	0.188	1.709	0.087	0.045	0.964
GC:SN	64	NA	NA	-0.234	0.815	-0.634	0.526	0.767	0.443
Positively stained cells	63	-1.266	0.206	0.010	0.992	0.902	0.367	-0.965	0.335
Low stain intensity	63	0.713	0.482	0.225	0.800	-0.496	0.624	0.020	0.984
Moderate stain intensity	63	-0.853	0.394	-0.497	0.619	1.056	0.291	-0.629	0.529
High stain intensity	63	-1.533	0.125	0.204	0.839	0.737	0.461	-0.453	0.651

Figure 1

Figure 1. Schematic of how the reproductive tracts were dissected. A) uterine horn sections

(1cm) were taken at the marked points, and B) ovaries were dissected, resulting in the final

ovary sections. The place at which the first section was taken is marked out under 3. Quartered

ovary. Created with Biorender.com.

Figure 2

Figure 1. Images taken of follicles representing the different stages of development. A) A

cluster of primordial follicles. B) Pre-antral follicle. C) Antral follicle. D) Atretic follicle.

Arrowheads indicate location of the oocyte.

Figure 3

Figure 3. Images taken of uterine horn cross-sections. Example of uterine cross section with

an H&E stain and indicators of the morphological structures investigated.

Figure 4

Figure 4. Images of IHC stained sections (8um) using a proliferating cell nuclear antigen

antibody, including the identified stain intensity and the control section. The automated

stain detection output can be seen including visual representations of cells considered to have

no stain, low stain, moderate stain, and high stain intensities.

Figure 5

Figure 5. Weight and reproductive parameters from females gestated in different in utero

environments. Variability of data represented by a violin plot of the birth weight (kg), stromal

nuclei count, primordial follicles, and total follicles between male-biased (n=9), non-biased

(n=15), and female-biased (n=15) groups. Data was analysed using a Levenes test. Birth

weights of piglets were measured on their first day post parturition post their first suckling

event. Stromal nuclei count, primordial follicle counts and total follicle counts were all

analysed on H&E stained sections. Manual counts were made for follicles, and an automated

nuclei count used in QuPath 0.2.0.

Figure 6

Figure 6. Birth weight of piglets for both primordial and total follicle numbers per cm2.

Scatter plots hold fitted regression lines, individual points have been grouped according to the

bias of the litter as either female-biased (>65% females), non-biased (45-54.9% females), and

male-biased (<35% females), n=34. Pigs from male biased litters held a higher follicle count

(both primordial and total) when they held a lower birth weight.

785 **Figure 7**

787 **Figure 7. The ratio of glands to endometrial area in female pigs in relation to their birth**

788 **weight (kg).** Ratio given for those individuals gestated in either female-biased (>65% females;

789 Blue dots), non-biased (45-54.9% females; Yellow dots), and male-biased (<35% females;

790 Green dots), n=34 in total.

792