Delivery of loaded MR1 monomer results in efficient ligand exchange to host MR1
and subsequent MR1T cell activation

[Corinna A. Kulicke[1], Gwendolyn M. Swarbrick[2]], Nicole A. Ladd[3], Meghan Cansler[2], Megan Null[2],

Aneta Worley[1], Chance Lemon[1], Tania Ahmed[2], Joshua Bennett[2], Taylor N. Lust[1], Chelsea M.

Heisler[1], Megan E. Huber[5], Jason R. Krawic[6], Laurisa M. Ankley[7], Savannah K. McBride[5], Fikadu G.

Tafesse[5], Andrew J. Olive[7], William H. Hildebrand[6], Deborah A. Lewinsohn[2], Erin J. Adams[3], David

M. Lewinsohn[1,4], Melanie J. Harriff*[1,4, 5]

Affiliations

[1]Division of Pulmonary, Allergy, and Critical Care Medicine, Oregon Health & Science University,

Portland, OR 97239, USA

[2]Division of Infectious Diseases, Department of Pediatrics, Oregon Health & Science University,

Portland, OR 97239, USA

[3]Department of Biochemistry and Molecular Biology, University of Chicago, Chicago, IL 60637,

USA

[4]VA Portland Health Care System, Portland, OR 97239, USA

[5]Department of Molecular Microbiology and Immunology, Oregon Health & Science University,

Portland, OR 97239, USA

[6]Department of Microbiology and Immunology, University of Oklahoma Health Sciences Center,

Oklahoma City, OK 73104, USA

[7]Department of Microbiology and Molecular Genetics, College of Osteopathic Medicine,

Michigan State University, East Lansing, MI 48824 , USA

* corresponding author: harriffm@ohsu.edu

Keywords

Abstract

MR1-restricted T cells have been implicated in microbial infections, sterile inflammation, wound healing and cancer. Similar to other antigen presentation molecules, evidence supports multiple, complementary MR1 antigen presentation pathways. To investigate ligand exchange pathways for MR1, we used MR1 monomers and tetramers loaded with 5-(2-oxopropylideneamino)-6-d-ribitylaminouracil (5-OP-RU) to deliver the antigen. Using MR1-deficient cells reconstituted with wild-type MR1 or MR1 molecules that cannot bind 5-OP-RU, we show that presentation of monomer-delivered 5-OP-RU is dependent on cellular MR1 and requires the transfer of ligand from the soluble molecule onto MR1 expressed by the antigen presenting cell. This new mode of antigen delivery strengthens the evidence for post-ER ligand exchange pathways for MR1, which could represent an important avenue by which MR1 acquires antigens derived from endocytosed pathogens.

Introduction

MR1-restricted T (MR1T) cells respond to small molecule antigens presented in the context of MHC I-related protein 1 (MR1) and include mucosal associated invariant T (MAIT) cells, which express the canonical TRAV1-2 TCR alpha chain [1]. They are increasingly being recognized for their role in early responses to microbial infection, but also in non-infectious conditions including a variety of autoimmune disorders, wound healing and cancer (reviewed e.g. in [1, 2]). Despite the heightened interest in these donor-unrestricted T cells, the mechanisms governing the presentation of MR1T cell antigens remain incompletely defined.

Successful antigen presentation requires the antigen presenting molecule to leave the endoplasmic reticulum (ER), bind an antigen, and display the bound antigen at the cell surface for T cell receptor (TCR) recognition. How these three features are achieved mechanistically, what compartments they take place in, and in what sequence they occur differs substantially between antigen presentation pathways. For major histocompatibility complex (MHC) I/peptide complexes, antigen binding generally precedes, and is indeed required for, ER egress and followed by translocation to the cell surface (reviewed e.g. in [3]). MHC II and CD1 molecules, by

contrast, exit the ER bound to a surrogate ligand which is exchanged for antigenic ligand in the

endosomal compartment after internalization from the cell surface (reviewed e.g. in [4, 5]).

For MR1, an MHC I-like mode of antigen acquisition has been described in which more MR1

associates with β_2-microglobulin (β_2m) and acquires endoglycosidase H (EndoH) resistance upon

addition of ligand to the culture medium [6, 7]. In line with this, McWilliam, Mak, et al. have used

a fluorescently labelled MR1 antigen analogue to directly demonstrate loading of exogenous

ligand in the ER [7]. Studies have linked the mechanism for MR1 ER egress to the charge

neutralization of a lysine residue within the MR1 ligand binding groove, which has been

suggested to result in a conformational change conducive to ER egress [6, 7]. The most potent

MR1 ligand described to date, 5-OP-RU, forms a Schiff base with this lysine residue, resulting in

its neutralization [6, 8].

In addition, recent findings support the hypothesis that there are complementary, non-exclusive

presentation pathways for MR1, allowing the molecule to survey diverse intracellular

compartments [9-14]. Multiple lines of evidence suggest that ligand exchange can occur on pre-

loaded MR1 molecules and that this may take place in an endosomal compartment. Most

notably, releasing MR1 from the ER prior to antigen exposure by pre-incubation with the MR1

ligand 6-formylpterin (6-FP) enhances presentation of exogenous antigenic ligand to MR1T cells

[10]. In line with this, siRNA knock-down of proteins involved in the trafficking and recycling of

endosomal vesicles such as VAMP4, Rab6 [9], Stx4, Stx16, and VAMP2 [10] impairs MR1-

mediated antigen presentation. Based on these observations, we and others have proposed

ligand exchange in a post-ER compartment as a complementary mechanism for MR1 ligand

acquisition [6, 10, 11, 14]. In a mechanism akin to MHC II and CD1 molecules, we hypothesize

that MR1 molecules internalized from the cell surface can be re-loaded with microbial antigens

in the endosomal compartment and recycle to the cell surface to activate MR1T cells. Here we

investigate the mechanisms underlying MR1 ligand exchange using a novel delivery method for

MR1T cell antigens. We find that MR1 ligands are efficiently presented when delivered in the

context of soluble MR1 molecules. This presentation is dependent on host MR1 and involves the

transfer of antigen from the recombinant molecule onto MR1 protein expressed by the cell.

Building upon our previously described tetraSPOT assay [15], this new method, the exSPOT,

extends the tool box available to investigate MR1 biology and provides important insights into

the mechanisms of MR1-mediated antigen presentation.

Results

MR1 ligands delivered to antigen presenting cells on soluble MR1 molecules are efficiently

recognized by an MR1T cell clone

The MR1 ligand 5-OP-RU is inherently unstable in aqueous environments [8, 16], which

presumably includes the intracellular milieu. However, the formation of a covalent bond with the

lysine residue at position 43 of MR1 stabilizes the antigen [8]. We have successfully used such

MR1/5-OP-RU complexes in their tetramerized form in plate-bound assays to activate MR1T cells

[15]. Similarly, Priya and Brutkiewicz have used latex beads coated with MR1/5-OP-RU complexes

to expand and activate human MAIT cells [17]. While these studies used recombinant MR1

molecules immobilized on a scaffold, here we sought to improve the efficiency of ligand delivery

into physiologically relevant antigen presenting cells (APCs) using soluble MR1.

To test this system, we incubated human dendritic cells (DCs) with MR1 tetramers loaded with

the canonical ligand 5-OP-RU (exSPOT), or applied the same concentration of tetramer directly

to the ELISPOT plate prior to adding an MR1T cell clone (tetraSPOT). Intriguingly, the MR1T cells

readily produced IFN-γ in response to the DCs incubated with soluble MR1/5-OP-RU in a dose-

dependent manner (Figure 1a). As expected based on previous work [15], they also produced

IFN-γ in the tetraSPOT. A quantitative comparison between the two assays was not possible due

to unknown coating efficiency in the tetraSPOT. MR1T cells did not respond to the MR1/6-FP

tetramer in the tetraSPOT, but we observed MR1T responses to the MR1/6-FP tetramers

incubated with DCs at the highest concentrations (Figure 1a). Similar to the tetraSPOT [15], MR1T

cell activation by DCs incubated with soluble MR1 tetramers was dependent on MR1, as blocking

with the anti-MR1 antibody clone 26.5 markedly reduced the response (Figure 1b, EC_{50} of 0.95

pM *vs* 31.62 pM, p<0.0001). Thus, 5-OP-RU delivered in the context of soluble, recombinant MR1

is efficiently presented by DCs and activates a human MR1T cell clone in an MR1-dependent

manner.

As professional APCs, DCs are particularly well equipped for phagocytosis of particulates of

various sizes and cross-presentation [18]. To investigate whether such specific functionality was

required in the exSPOT, we incubated the monocytic cell line THP-1, a B cell line (LCL), or the

airway epithelial cell line BEAS-2B with MR1/5-OP-RU tetramers and quantified IFN-γ responses.

As with DCs, the MR1T cell clone produced IFN-γ in response to all of the tetramer-incubated cell

types (Figure 1c). However, DCs were nearly three orders of magnitude more efficient at

presenting the antigen compared to the other cell types (EC$_{50}$ of 0.55 pM *vs* 142 − 265 pM,

p<0.0001).

Ligand delivered in the context of MR1 monomer is presented more efficiently than free ligand

Next, we compared the presentation of MR1/5-OP-RU tetramers and MR1/5-OP-RU monomers

to reduce the variability associated with tetramerization (e.g. biotinylation efficiency,

streptavidin binding, and ligand saturation). Using the MR1 monomer further circumvented

potential variability introduced by instability of the tetramer complex over time.

To our surprise, presentation of ligand from monomer-incubated DCs to MR1T cells was even

more efficient than presentation of ligand from tetramer-incubated DCs, when accounting for

valency (Figure 2a). Although the EC$_{50}$ values of the complexes were comparable (Tetramer EC$_{50}$:

0.204 pM *vs* Monomer EC$_{50}$: 0.22 pM, p=0.7517), there was a marked difference in the estimated

EC$_{50}$ for 5-OP-RU which takes into account that each tetramer carries up to four 5-OP-RU

molecules whereas each monomer carries one at most (Tetramer 5-OP-RU EC$_{50}$: 0.816 pM *vs*

Monomer 5-OP-RU EC$_{50}$: 0.22 pM, p<0.0001). Comparing monomer-delivered 5-OP-RU to free

ligand further revealed improved efficiency of the monomer over the unbound antigen (Figure

2a; Monomer EC$_{50}$: 0.22 pM *vs* Free EC$_{50}$: 4.97 pM, p<0.0001). Of note, these calculations assume

full occupancy of monomers and tetramers.

To further investigate the relative efficiency of loading free *vs* monomer-bound 5-OP-RU, we

wanted to specifically detect MR1 molecules bearing 5-OP-RU at the cell surface by flow

cytometry. To this end, we generated a TCR tetramer based on the TCR sequence of the same

previously characterized MR1T cell clone that was used in the functional assays (TRAV1-2[+] clone

D426-G11 – see references [15, 19]). We confirmed that this reagent specifically stained MR1

molecules loaded with 5-OP-RU but not 6-FP (Supplementary Figure 1). When we incubated

BEAS-2B cells overexpressing GFP-tagged MR1 (BEAS-2B.MR1-GFP) with either MR1/5-OP-RU

monomer or free 5-OP-RU overnight and stained with the TCR tetramer to track the cell surface

translocation of MR1 bound to the antigen, almost 100-fold higher concentrations of free 5-OP-

RU were needed to reach cell surface levels comparable to monomer-delivered ligand (Figure

2b). Together, these results demonstrate that soluble, recombinant MR1 is an efficient

mechanism for delivery of MR1T cell antigens in a variety of cell lines *in vitro.*

Presentation of MR1 ligands delivered on soluble MR1 molecules requires cellular MR1

In principle, the MR1/5-OP-RU complex that interacts with the TCR in our assay could be either

endogenously expressed MR1 bound to 5-OP-RU from the monomer or the recombinant MR1/5-

OP-RU monomer itself. Thus, we tested whether endogenous MR1 was required. We incubated

WT BEAS-2B cells, MR1[-/-] BEAS-2B cells lacking all isoforms of MR1, and MR1[-/-] BEAS-2B cells

reconstituted with tetracycline-inducible MR1A [20, 21] with MR1/5-OP-RU monomer and used

them as APCs. As expected, we detected responses to WT BEAS-2B cells incubated with MR1/5-

OP-RU monomers (Figure 3a, left), but when these monomers were incubated with the MR1[-/-]

BEAS-2B cell line, the response was completely abrogated (Figure 3a, middle). Reconstitution of

MR1 expression in MR1-deficient cells resulted in complete recovery of MR1T cell responses to

the monomer-delivered ligand (Figure 3a, right). These observations were confirmed in THP-1

cells (Figure 3b).

Ligands are transferred from soluble MR1 molecules onto cellular MR1

Having established that MR1 expression by the APC itself was required for presentation of

monomer-delivered ligand, we considered two possible scenarios: (1) dimerization or indirect

binding of the monomer to host MR1 at the cell surface and (2) uptake of the MR1/5-OP-RU

complex and transfer of the antigen from the monomer onto cellular MR1.

Since the MR1 molecules used to generate the ligand-loaded monomers and tetramers do not

contain a transmembrane domain [8], their presence on the cell surface and, hence, ability to

activate MR1T cells, would require lateral interactions with other membrane-bound proteins. Of

note, Lion and colleagues have demonstrated that a GFP-tagged version of the MR1B isoform

weakly associated with the MR1A isoform although that observation was considered a potential

technical artifact [22]. To directly investigate the presence of the monomer on the cell surface,

we incubated BEAS-2B.MR1-GFP cells with the biotinylated MR1/5-OP-RU monomer and stained

the cells with fluorophore-coupled streptavidin. We did not detect any signal from the

biotinylated monomer although our controls confirmed that we added sufficient monomer to

induce MR1 surface translocation and sufficient streptavidin to detect biotinylated molecules at

the cell surface (Figure 4a). Interestingly, we detected a small but notable shift in total MR1

surface levels after incubation with the monomer (Figure 4a, right). This suggests that enough 5-

OP-RU ligand is delivered to increase not just TCR-reactive, but overall MR1 surface levels.

Together, these data do not support MR1 dimer formation at the cell surface.

Since we could not find evidence for direct presentation by the monomer, we next tested the

hypothesis that the 5-OP-RU is transferred from the soluble MR1/5-OP-RU complex onto the MR1

expressed by the APC. First, we sought to establish whether actin-dependent internalization of

soluble MR1 molecules was required for the presentation of monomer-delivered 5-OP-RU. Thus,

we treated BEAS-2B.MR1-GFP cells with latrunculin B (LatB), a well-characterized inhibitor of

actin polymerization. Surface levels of 5-OP-RU-loaded MR1 were reduced in cells treated with

LatB as compared to the DMSO control (Figure 4b). Of note, LatB treatment also reduced surface

expression of TCR-reactive MR1 in response to free 5-OP-RU (Supplementary Figure 4a). Total

MR1 as measured by the GFP expression was not reduced by LatB treatment (Supplementary

Figure 4b). We concluded that an active actin cytoskeleton is required for presentation of the

monomer-derived ligand. This is consistent with previous results demonstrating a requirement

for actin for an efficient MR1 response to *Escherichia coli* (*E.coli*) [13, 23] and could indicate a

requirement for either internalization or intracellular trafficking.

189 Additional evidence that intracellular processing is required came from the observation that the

190 kinetics of presentation differ between monomer-delivered ligand compared to free 5-OP-RU.

191 Here, we incubated BEAS-2B.MR1-GFP cells with concentrations of MR1/5-OP-RU monomer (50

192 nM) and free 5-OP-RU (6.4 µM) that resulted in intermediate surface expression (Figure 2b) and

193 followed MR1 surface translocation over time. In agreement with McWilliam et al., who detected

194 surface expression of 5-OP-RU-loaded MR1 after 2 h [6], we saw loaded MR1 as soon as 1 h after

195 addition of the free ligand (Figure 4c). By contrast, it took 6 h for the MR1 surface expression to

196 reach comparable levels after addition of the MR1/5-OP-RU monomer (Figure 4c). These kinetics

197 are consistent with a longer time needed for internalization, processing, ligand exchange, and

198 recycling of monomer-delivered ligand compared to free 5-OP-RU.

199 To directly investigate transfer onto cellular MR1, we tested whether the MR1 expressed by the

200 APC needed to be capable of presenting 5-OP-RU or whether its presence was sufficient. We

201 reconstituted the MR1$^{-/-}$ BEAS-2B cell line with a CRISPR-resistant version of MR1A that carried

202 the R9H mutation described by Howson et al. [24]. This mutation allows stable MR1 protein

203 expression and binding of 6-FP but renders the protein incapable of presenting 5-OP-RU ([24]

204 and Supplementary Figures 2 and 3). Indeed, reconstituting the MR1-deficient cells with this

205 mutant did not rescue the presentation of either monomer-delivered 5-OP-RU or *Mycobacterium*

206 *smegmatis (M. smegmatis)* supernatant (Figure 4d), further supporting the idea that it is the

207 cellular MR1 that presents the 5-OP-RU ligand.

208 *Ligand exchange occurs in post-ER compartments and does not require acidification*

209 Our data demonstrate the transfer of monomer-delivered 5-OP-RU onto MR1 expressed by the

210 APC but the mechanism of this exchange is unclear. Our lab has previously shown that pre-

211 incubation with the non-stimulatory ligand 6-FP can increase MR1-mediated antigen

212 presentation, presumably by increasing the pool of MR1 molecules available for ligand exchange

213 [10]. When we applied this concept to the monomer delivery assay, we found a similar effect:

214 releasing MR1 from the ER by pre-incubation with 6-FP decreased the EC$_{50}$ for presentation of

215 monomer-delivered ligand from 1.027 pM to 0.1365 pM (p<0.0001; Figure 5a). Thus, MR1

216 molecules pre-loaded with a non-stimulating ligand appear to contribute to the activation of

MR1T cells after delivery of antigenic ligand on monomers, supporting the notion of ligand

exchange. In this context, it is worth noting that 6-FP concentrations multiple orders of

magnitude higher were required to inhibit 5-OP-RU presentation in a functional assay [25], which

suggests stronger binding of 5-OP-RU [6] and may help push the equilibrium towards binding of

the activating ligand.

To probe to what extent the monomer delivery could be extended to other antigen presentation

pathways and potentially gain mechanistic insights from any parallels, we repeated the assay

with peptide-loaded MHC I and II tetramers. We first used a fluorophore-conjugated HLA-

B*08:01/CFP10$_{3-11}$ tetramer to stain a CD8$^+$ T cell clone restricted by this MHC I/peptide complex

(D480-F6 – see references [26] and [27]) for flow cytometry, confirming the T cell clone reacted

with the tetramer (Figure 5b). However, using the same tetramer to load autologous DCs in an

ELISPOT with the same T cell clone did not result in the production of IFN-γ in an APC-dependent

manner (Figure 5c). As expected, the T cells responded to DCs incubated with free peptide (Figure

5c). Next, we tested an HLA-DRB1*14:03/ESAT6$_{73-78}$ tetramer in combination with a T cell clone

that was specific to the antigen but restricted by a different MHC II allele (D481-D1, from an HLA-

DRB1*14:03 negative individual). As expected, this tetramer did not stain the T cell clone by flow

cytometry (Figure 5d). Nevertheless, feeding the non-matched tetramer to DCs autologous to the

same MHC II-restricted T cell clone elicited an IFN-γ response (Figure 5e). The only source of

matched MHC II molecules in this case are the DCs, further supporting a requirement for transfer

of the antigen from the soluble protein onto antigen presenting molecules expressed by the APC

itself. In line with this, the T cells did not respond when using HLA-mismatched DCs or the A549

cell line which express MHC II alleles different from Donor 481 ([28], Cellosaurus cell line A-549

(CVCL_0023) (expasy.org)) (Figure 5f). Thus, like with MR1, presentation of the MHC II antigen

delivered in the context of recombinant antigen presenting molecules requires transfer onto

proteins expressed by the APC.

Since soluble proteins could efficiently deliver peptide for MHC II presentation, we reasoned that

the MR1 loading in our assay was similar to the MHC II pathway, which involves exchange of a

surrogate ligand in an endosomal compartment (reviewed e.g. in [4]). To test whether MR1 ligand

transfer from the monomer could be mediated in an analogous fashion, we investigated the

effect of brefeldin A (BFA) on the presentation of monomer-delivered 5-OP-RU. BFA disrupts the

Golgi apparatus and effectively traps secretory proteins in the ER. Indeed, incubation with BFA

diminished surface translocation of 5-OP-RU-loaded MR1 in response to the MR1/5-OP-RU

monomer (Figure 5g, left). However, this experiment is unable to distinguish between two

different scenarios: 1) ligand exchange occurs in the ER and MR1 needs to subsequently traverse

the Golgi for surface presentation, or 2) MR1 needs to leave the ER to be available for exchange

in a post-ER compartment. To distinguish between these two hypotheses, we pre-incubated the

cells with 6-FP to induce ER egress of a pool of MR1 molecules before adding the BFA. Under

these conditions, BFA had a minimal effect on presentation of monomer-delivered ligand (Figure

5g, second from left). As expected, presentation of free 5-OP-RU was similarly impaired in the

presence of BFA and improved after pre-incubation with 6-FP (Supplementary Figure 5). It is

important to note that BFA treatment also reduced total MR1 levels as measured by GFP

expression (Figure 5g, right panels). These data clearly show that 5-OP-RU delivered on soluble,

recombinant MR1 molecules can be transferred to cellular MR1 in a post-ER compartment.

Exchange of exogenous ligand onto MHC II [4] and CD1 [5] molecules is potentiated by the low

pH in the endolysosomal compartment. Pertinently, endosomal acidification is required for

efficient presentation of *E.coli* by MR1 [13, 29] and McWilliam et al. have proposed that the Schiff

base that links 5-OP-RU to the MR1 ligand binding groove is labile under acidic conditions [6, 14].

Therefore, we investigated whether presentation of monomer-delivered ligand was dependent

on endosomal acidification using both a cell-free and a functional approach. First, we tested the

stability of the MR1/5-OP-RU and MR1/6-FP monomers under increasingly acidic conditions in

an ELISA. Fully folded, recombinant MR1 loaded with either 6-FP or 5-OP-RU was readily

detectable using both a conformation-dependent anti-MR1 antibody [30] and an anti-β_2m

antibody at neutral pH (Figure 6a). As expected, at neutral pH, a control MHC I/peptide complex

was detected by the anti-β_2m antibody but not by the anti-MR1 reagent and the recombinant $\gamma\delta$

TCR used as a control was detected by neither antibody. Since MR1 requires a ligand to efficiently

refold *in vitro* [8, 31], both read-outs specifically report on fully folded trimeric complexes.

Surprisingly, binding of neither the anti-MR1 nor the anti-β_2m antibody was affected by overnight

treatment of the plate-bound monomer with buffers of increasing acidity, covering the range of

those observed in the endomembrane compartments (pH 4.5-7.5). Acidic conditions, therefore,

did not destabilize the antibody-reactive conformation of MR1 in this assay. Hence, it appears

unlikely that loaded MR1 ligands are released simply by virtue of the acidic conditions in the

endosomal compartment.

To confirm these results in a cellular assay, we pre-incubated BEAS-2B WT cells with bafilomycin

before performing the exSPOT. Bafilomycin inhibits vacuolar ATPases [32] and diminishes the

presentation of an endosomal processing-dependent 5-A-RU pro-drug to MR1T cells [12].

Whereas the bafilomycin concentrations we used here reduced the MR1T cell response to this

pro-drug in a dose-dependent manner, presentation of the monomer-delivered 5-OP-RU was not

affected by the inhibitor (Figure 6b). Together, these data support a model in which the transfer

of 5-OP-RU delivered in the context of a soluble MR1 monomer onto the MR1 expressed by the

APC can displace pre-bound 6-FP and is independent of endosomal acidification.

Transfer of monomer-bound 5-OP-RU requires breaking a covalent bond. Since acidification was

not needed for ligand transfer (Figure 6b), we hypothesized that this process is catalyzed by an

exchange chaperone akin to HLA-DM for MHC II [4, 33]. TAPBPR, tapasin, and the adapter protein

AP2A1 have been shown to bind MR1 and modulate its antigen presentation [7, 34, 35]. However,

knock down (KD) of these candidate chaperones alone or together did not affect how efficiently

monomer-delivered 5-OP-RU was presented to our MR1T cell clone (Supplementary Figure 7).

To further characterize the compartment where ligand exchange in the exSPOT occurs, we

performed immunofluorescence microscopy. We observed AF647-labeled MR1/5-OP-RU

tetramer in compartments labeled with Rab5 (23.51%), an early endosome marker, or LAMP1

(33.1%), which localizes to late endosomes and lysosomes in these cells [36] (Figure 6c). These

observations support the hypothesis that ligand transfer occurs in post-ER compartments,

including the early and late endosome.

Vesicular trafficking between the cell surface and the endosomal compartments is mediated by

soluble N-ethylmaleimide-sensitive factor attachment protein receptors (SNAREs) [37]. The

SNARE syntaxin-4 (STX4) is required for exocytosis in various cellular contexts [38-41] including

the fusion of recycling endosomes to the plasma membrane in cytotoxic T cells [38] and post-

synaptic membranes on dendritic spines [39]. Our lab has previously shown that siRNA-mediated

knock down of STX4 reduces MR1-mediated antigen presentation of *M. smegmatis* supernatant

but not live *Mycobacterium tuberculosis* [10]. To investigate the involvement of this mediator of

vesicular trafficking in the exSPOT, we generated BEAS-2B cells deficient for STX-4 using

CRISPR/Cas9 technology. Consistent with previous results in the knock down system [10], STX-4

KO cells presented *M. smegmatis* supernatant less efficiently than their Cas9-expressing parent

cell line (Figure 6d, right, p<0.0001). Similarly, presentation of monomer-delivered antigen was

reduced in the absence of the SNARE (Figure 6d, left, p=0.0032). These results implicate STX4 in

the mechanism underlying the exSPOT and suggest that it may share mechanistic parallels with

presentation of microbial ligand.

Discussion

Distinct and complementary cellular pathways have been proposed for MR1 loading and

trafficking [11, 14, 42, 43]. While most research efforts have been focused on the ER pathway,

increasing evidence supports exchange in a post-ER compartment as a mechanism by which MR1

acquires antigenic ligand(s). For example, both Karamooz et al. and McWilliam et al. have shown

that MR1 molecules in cells pre-incubated with the MAIT cell antagonist 6-FP can present

activating MR1 ligands upon subsequent incubation with agonistic ligand [6, 10]. This exchange

requires internalization of the molecules and partially counters the ability of Brefeldin A (BFA) to

block surface expression of MR1/5-OP-RU complexes, indicating re-loading in a post-ER

compartment [6]. Indeed, pre-incubation with 6-FP enhances MR1-mediated presentation of

bacterial supernatant, presumably by increasing the pool of MR1 molecules available for re-

loading [10]. Using a fluorescence internalization probe (FIP), McWilliam et al. have directly

demonstrated recycling of a small proportion of MR1 molecules in C1R cells [6]. In our model of

mycobacterial infection of airway epithelial cells, this pathway contributes substantially to

presentation of antigen derived from intracellular infection [9, 10].

A recently developed pro-drug requires endosomal processing to release the 5-OP-RU precursor

5-A-RU [12]. Since the antigen is generated in endosomes in this scenario, it appears likely that

this compound is loaded onto MR1 in this compartment as opposed to trafficking to another

intracellular organelle, which would require it to cross multiple membranes. Interestingly, while

Ac-6-FP inhibits T cell activation after delivery of free 5-A-RU, it does not interfere with the

presentation of this pro-drug [12]. This is consistent with the earlier observation that 6-FP blocks

presentation of bacterial supernatant but not of intracellular infection [9]. Together, these results

suggest that there are distinct pools of MR1 in the ER and in a post-ER compartment [11, 12, 14,

43], that these compartments are differentially accessed by exogenous ligands, and/or that

conditions are more favorable for ligand exchange in the endosomal compartment than in the

ER. This notion is not without precedent as multiple, complementary pathways including both

direct loading in the ER and ligand exchange in the endosomal compartment have been shown

for MHC I (reviewed e.g. in [44]).

Of note, a synthetic MR1 ligand modified to be able to track its location by microscopy resides in

small vesicles also occupied by the micropinocytosis marker dextran [7]. This supports the idea

that both membrane-bound internalized MR1 and soluble MR1 ligands could occupy this

compartment at the same time, facilitating loading [14]. Furthermore, MR1-mediated antigen

presentation of *M. smegmatis* supernatant is reduced upon knock-down of endosomal trafficking

molecules [9, 10], further supporting involvement of this pathway in efficient loading of some

antigens.

In this study, we report a novel delivery method for MR1 ligands which allowed us to directly

demonstrate transfer of the MR1T cell-activating ligand 5-OP-RU from recombinant, soluble MR1

onto cellular MR1. Intriguingly, monomer-delivered antigen was presented more efficiently than

tetramer-delivered ligand. We speculate that the bulky structure of the tetramer may result in

steric hindrance of interactions with chaperones required for ligand transfer. The smaller, more

compact monomer, on the other hand, may be better suited to adopt the correct orientation.

Moreover, DCs were much more efficient at presenting the tetramer-derived ligand than other

cell lines tested. This could be due to their ability to cross-present internalized antigen, which

allows their MHC I molecules to sample diverse sources of antigen and may enable MR1 to do
likewise [18].

Monomer-delivered 5-OP-RU was also presented to MR1T cells more efficiently than free 5-OP-
RU. One explanation for this observation is that binding to the monomer protects the inherently
unstable antigenic molecule from dehydrative cyclization [8]. However, the analogous results
with MHC II tetramers argue that the enhanced antigen presentation associated with monomer
delivery is not simply attributable to increased stability as peptide antigens are more stable.
Therefore, we think it more likely that the monomer delivery enhances the efficiency of
presentation by improving another aspect of antigen processing or presentation.

Our efforts to identify the molecular mechanism that enables transfer of the antigenic ligand
from the soluble MR1 to the cellular MR1 show that this process is dependent on actin dynamics
and enhanced if cellular MR1 is pre-loaded with 6-FP. Peptide editing on MHC I and MHC II
depends on the relative affinities of the available ligands and their competition for transiently
ligand-receptive antigen presenting molecules [3, 4, 33]. In line with this concept, we propose
that the endosomal compartment is a uniquely suited place for MR1 ligand exchange by virtue
of its high local concentration of antigens derived from internalized pathogens relative to the cell
as a whole. This enrichment of otherwise low abundance ligands may drive the equilibrium from
pre-bound ligands to internalized antigens specifically in endosomal compartments.

This analogy breaks down, however, in that peptides are not covalently bound to their antigen
presenting molecules. By contrast, a covalent bond needs to be broken in our exchange assay to
release the antigenic ligand from the soluble MR1. The Schiff base that anchors both 6-FP and 5-
OP-RU to the MR1 binding groove is considered labile under acidic conditions [6, 14].
Nevertheless, our results suggest that ligand exchange on MR1 does not require endosomal
acidification. Indeed, plate-bound MR1 was resistant to conformational changes associated with
ligand release under acidic conditions. This is in striking contrast to MR1-mediated presentation
of fixed whole *E.coli* [13] and the 5-A-RU pro-drug [12]. While the latter was designed to require
acidic conditions for release, the first suggests that the molecular mechanisms for MR1 loading
differ between *E.coli* and 5-OP-RU-loaded monomer. Importantly, *Mycobacterium tuberculosis* is

known to prevent endosomal acidification to evade immune recognition [45], making antigen

loading mechanisms that function at higher pH desirable. Alternatively, ligand transfer after

monomer delivery may be insensitive to bafilomycin because it occurs in the ER. Previous studies

have demonstrated that whole proteins from the extracellular environment can be imported into

this compartment [46-48], so this may be true for the MR1 monomer, too. In that case, we would,

however, expect monomer delivery to efficiently supply ligand to MHC I molecules as well, which

was not the case (Figure 5c). Our BFA experiments also show that MR1 ligand exchange can take

place after MR1 is released from the ER bound to 6-FP (Figure 5g).

While the molecular mechanism and the subcellular location of ligand exchange remain to be

determined, our data lead us to propose that it is a facilitated process catalyzed by a yet

unidentified exchange chaperone (Supplementary Figure 9). Monomer delivery may increase the

efficiency of exchange by enhancing the interaction between pre-loaded MR1 and this

chaperone. Indeed, specific chaperones facilitate the release of pre-loaded ligands, stabilize the

transiently unoccupied antigen presenting molecule, and mediate the re-loading with antigenic

cargo in other examples of ligand exchange. For MHC I, for instance, this function can be fulfilled

by TAPBPR in the Golgi (reviewed e.g. in [49] and [50]). For MHC II, HLA-DM chaperones

endosomal ligand exchange and exchange kinetics are further modulated by its regulator HLA-

DO (recently reviewed in [33] and [4]). Recent screens for modulators of MR1 loading and

trafficking have identified numerous interaction partners through both genetic techniques and

protein co-immunoprecipitation [7, 51]. These studies found MR1-mediated antigen

presentation to be regulated by tapasin and TAPBPR [7, 34] and the P5-type ATPase ATP13A1

[51], but have largely focused on ER and Golgi-resident proteins. Discovery approaches more

specifically targeted at interaction partners during ligand exchange in endosomal compartments

may be needed to identify the chaperone(s) involved in this process.

In this manuscript, we explored the mechanisms by which MR1 ligands can be delivered

exogenously using MR1 itself to stabilize an inherently unstable ligand. This mode of ligand

delivery is a powerful tool to identify chaperones involved in the re-loading of pre-loaded MR1

molecules which can then be tested for their involvement in physiological settings such as

mycobacterial infection. 5-OP-RU was also recently successfully used to boost mucosal immunity

in the context of vaccinations against respiratory pathogens [52]. Thus, in addition to its use in

exploring the molecular underpinnings of ligand exchange, we also postulate MR1/ligand

complexes could be used as an adjuvant or antigen delivery strategy in novel vaccination

approaches.

Methods

Human subjects

This study was conducted according to the principles expressed in the Declaration of Helsinki.

Study participants, protocols, and consent forms were approved by the Institutional Review

Board at Oregon Health & Science University (OHSU) (IRB00000186).

Cells and bacteria

M. smegmatis (strain mc^{2}155) supernatant was obtained by growing *M. smegmatis* and filtering

the supernatant as described previously [10]. PBMCs were isolated from whole blood obtained

by leukapheresis with informed consent, as previously described [53]. Human monocyte-derived

dendritic cells (DCs) were isolated from PBMCs by plate adherence and cultured in RPMI+10%

heat-inactivated human serum (HuS) with 30 ng/ml GM-CSF (Immunex) and 10 ng/ml IL-4 (R&D

Systems) as previously described [36]. BEAS-2B cells were obtained from the American Type

Culture Collection (ATCC) and cultured as recommended. BEAS-2B MR1$^{-/-}$, BEAS-2B MR1$^{-/-}$: MR1A

[20, 21] and the BEAS-2B cell line overexpressing GFP-tagged MR1A (BEAS-2B.MR1-GFP) [9] were

described previously. THP-1 cells were obtained from ATCC. THP-1 MR1$^{-/-}$ and THP-1 MR1$^{-/-}$: MR1

cell lines were described previously [21, 54]. The THP-1 MR1$^{-/-}$ from the Sewell lab constitutively

express Cas9 protein whereas the THP-1 MR1$^{-/-}$: MR1 cells from the Cerundolo lab do not. EBV-

transformed B cell lines (LCL) were generated in our laboratory using supernatants from the cell

line 9B5–8 (ATCC). A549 cells were obtained from ATCC. The MR1 R9H res construct was based

on [24]. MR1 res and MR1 R9H res contain silent mutations to make the sequence resistant to

CRISPR/Cas9 editing in the MR1$^{-/-}$ background. The coding sequence with a downstream IRES GFP

[55] was ordered from GeneArt (ThermoFisher) and cloned into our previously described pCI AscI

vector [10]. Plasmids were sequenced at the OHSU sequencing core and transfected into BEAS-

2B MR1$^{-/-}$ cells using the Amaxa nucleofection system with Kit T solution (Lonza). The transfection

efficiency was evaluated via GFP expression by flow cytometry (Supplementary Figure 3).

Trichoplusia ni (Hi-5) and *Spodoptera fugiperda* (Sf9) cells for baculovirus production and

recombinant protein expression were maintained as described previously [56].

CRISPR/Cas9-mediated knock out of STX4

Early passage BEAS-2B WT cells were stably transduced with lentiviral particles encoding Cas9 by

spinoculation as previously described [57]. Transduced cells were selected by culturing with 5

µg/ml Blasticidin (Life Tech) for 12 days. Three sgRNA constructs targeting STX4 were generated

and packaged into lentiviral particles as described [58]. Early passage BEAS-2B.Cas9 cells were

then transduced with supernatant containing these lentiviral particles in the presence of 200 µg

Polybrene (Sigma). Transduced cells were selected by culturing with 8 µg/ml Puromycin (Sigma)

for 6 days. Monoclonal populations were created by limiting dilution and a functional screen was

performed by ELISPOT. Genomic editing efficiency and clonality were verified by Sanger

sequencing. Briefly, DNA was isolated from STX4 KO clones and BEAS-2B.Cas9 control cells using

the QIAamp DNA Micro Kit (Qiagen) per manufacturer's protocols. DNA surrounding the sgRNA

target site was amplified by PCR and purified using the QIAquick PCR Purification Kit (Qiagen) per

manufacturer's protocols. Sanger sequencing was performed by capillary electrophoresis at the

OHSU Vollum Institute DNA Sequencing Core and sequences were analyzed by TIDE [59] and ICE

[60]. Sequence of the sgRNA used to generate the STX4 KO clonal cell line used in this manuscript:

AGAACGTGGAGCGGATTCGG.

T cell clones

All T cell clones were expanded and maintained as previously described [19, 26]. Three T cell

clones were used in this work. Two clones have been published previously (D426-G11 [15, 19]

and D480-F6 [26, 27]). D481-D1 is a CD4$^+$ T cell clone that recognizes Esat6$_{73-87}$. This clone was

generated in a limiting dilution assay by adding either 100, 200 or 600 CD4$^+$ T cells (CD4

microbeads, Miltenyi) isolated from PBMC from D481, a participant who had active tuberculosis,

to wells of a 96 well plate containing 1e5 irradiated (3,500 rad using a Cs source) allogeneic

PBMCs, 1e4 irradiated (3,500 rad using a Cs source) autologous dendritic cells (prepared as

described above), 2 µg/ml Esat6 protein (BEI Resources), 5 ng/ml interleukin-2 (Proleukin) and

incubated at 37°C for two weeks. Wells with growth were assessed for response to an Esat6

peptide pool consisting of 15mers overlapping by 11 amino acids in an IFN-γ ELISPOT. T cell

clones that responded to the Esat6 peptide pool were expanded as described previously [19, 26]

and further screened against the individual 15mers. The restricting allele is not known for this

clone, but D481 expresses these class II alleles: DRB1*08:01, DRB1*13:01, DPB1*01:01,

DPB1*04:01, DQB1*04:02 and DQB1*06:03 (HLA typing by Kashi Clinical Laboratories, Portland,

OR).

ELISPOT assays

For the plate-bound tetramer ELISPOT (tetraSPOT) assay, ELISPOT plates were coated with an

anti–IFN-γ antibody, as previously described [61]. At the time of coating, MR1/5-OP-RU and

MR1/6-FP tetramers conjugated to PE obtained from the NIH tetramer core [8] were added at a

range of concentrations. After overnight incubation at 4°C, ELISPOT plates were washed three

times with phosphate-buffered saline and then blocked with RPMI 1640 + 10% human serum

(RPMI+10% HuS) for at least 1 h. T cell clones (1e3-2e4) were added to wells overnight. IFN-γ

ELISPOTs were enumerated following development as previously described [61]. For all other

ELISPOT assays, ELISPOT plates were coated with anti-IFN-γ antibody and blocked as described

above. 1e4 DC, LCL, THP-1 or BEAS-2B cells were plated in the ELISPOT wells in RPMI+10% HuS.

Tetramer, monomer, ligand or peptide were added to the wells at concentrations indicated for

each experiment. PHA was used as positive control. After at least 1 h, 1e3-2e4 T cell clones were

added and the plate was incubated overnight at 37°C. 6-FP pre-treatment assays were performed

as described [10]. Briefly, BEAS-2B.MR1-GFP cells were plated in 6-well tissue culture plates and

treated with 6-FP *vs* NaOH control over night. On the next day, the cells were harvested and used

as APCs in an IFN-γ ELISPOT assay as described above. For experiments with Bafilomycin A1

(BAFA1, MedChemExpress), BEAS-2B cells were incubated with BAFA1 for 2 h at indicated

concentrations prior to the addition of 100 pM MR1/5-OP-RU monomer or 50 nM 5-OP-RU

prodrug (MedChemExpress) in an IFN-γ ELISPOT assay as described above.

5-OP-RU Ligand

5-(2-oxopropylideneamino)-6-d-ribitylaminouracil (5-OP-RU) was made from two sources of 5-

amino-6-D-ribitylaminouracil (5-A-RU). For the first method, 5-nitro-6-d-ribitylaminouracil (5-N-

RU) was prepared by the OHSU Medicinal Chemistry Core. 5-A-RU was freshly prepared from 5-

N-RU following a previously described procedure [8]. In brief, 4.8 mg of 5-N-RU was dissolved in

400 µl of 50 mM MES, pH 5.8. 8-fold molar excess of sodium dithionite powder was added and

incubated in the dark at 90°C for five minutes under argon. The solution was chilled in an ice

water bath. To prepare 5-OP-RU, at least 15 molar equivalents of methylglyoxal was added to 5-

A-RU in 100 mM HEPES, pH 7.2, 150 mM NaCl to give a stock solution at 20 mM. For the second

method, 5-A-RU*HCl was prepared by the OHSU Medicinal Chemistry Core following a previously

described procedure [62]. To prepare 5-OP-RU, equal volumes of circa 650 mM methylglyoxal

(Sigma) and 32 mM 5-A-RU*HCl dissolved in water were combined immediately before addition

to the assay to give a stock solution of 16 mM of 5-OP-RU. Mass spectrometry analysis confirmed

near-complete conversion of 5-A-RU into 5-OP-RU under similar conditions (Supplementary

Figure 8). For this, samples were directly injected into an AB Sciex TripleTOF 5600 quadrupole

time of flight mass spectrometer using the built-in syringe pump through a Hamilton Model 1710

N 100 µl syringe. Samples passing through the syringe were ionized by a PicoView ion source by

New Objective before passing through a Co-Ann technologies 11cm 20 µm inner diameter pulled

glass emitter which tapered to a 10 µm inner diameter at the tip. A stock solution of 32 mM 5-A-

RU was diluted 1:10 before injection into the mass spectrometer. 5-OP-RU was produced by

combining a 5 µl stock solution of 32 mM of 5-A-RU with a 5 µl 1:10 dilution from a 6.5 M stock

solution of methylglyoxal (Sigma). This 5-OP-RU was then diluted 1:10 for injection into the mass

spectrometer. Samples were examined in negative ion mode with the source voltage set at -2600

V. MS1 and MS2 data was acquired using Analyst 1.8 in manual tuning mode, with a TOF mass

range of 30 to 500 Da after 1 minute. MS2 data was collected in high sensitivity mode with a

declustering potential of -40 and a collision energy of -30 for 5-A-RU and -38 for 5-OP-RU. Data

was analyzed using PeakView 1.2.

Flow Cytometry assays

The following reagents were obtained through the NIH Tetramer Core Facility: MR1/5-OP-RU

tetramer and monomer and MR1/6-FP tetramer and monomer, HLA*B08:01/CFP10$_{3-11}$ tetramer,

HLA-DRB1*14:03/Esat6$_{73-87}$ tetramer, HLA-DRB1*14:03/control tetramer. Identification of MAIT

cells with MR1 tetramers was described in [8]. The D426-G11 TCR tetramer was made as specified

below. To confirm the specificity of the TCR tetramer as a staining reagent, BEAS-2B.MR1-GFP

cells were incubated with the indicated concentrations of 5-OP-RU or 6-FP, or the equivalent

volume of 0.01 M NaOH overnight. Cells were harvested with Trypsin and split into three wells

of a U-bottom 96-well plate each for staining with TCR tetramer, anti-MR1 antibody 26.5

(BioLegend, 361108), or isotype control antibody (BioLegend, 400222) for at least 30 min on ice.

BEAS-2B MR1-/- cells transfected with the MR1 res or MR1 R9H res plasmids were analyzed

similarly. For ligand titration experiments, 5e4 BEAS-2B.MR1-GFP cells per well were incubated

with the indicated concentrations of free 5-OP-RU or MR1/5-OP-RU monomer in a 24-well plate

overnight. Cells were harvested with Trypsin, transferred to a 96-well U-bottom plate, washed,

and stained with TCR tetramer for at least 30 min at 4° C. For time course experiments, 5e4 BEAS-

2B.MR1-GFP cells per well were plated in 24-well plates. Free 5-OP-RU or MR1/5-OP-RU were

added to the indicated final concentrations for the indicated amounts of time. Samples from all

time points were harvested and stained at the same time as above. Free ligand and monomer

titrations were performed in parallel in both sets of experiments. For the actin blockade, 5e4

BEAS-2B.MR1-GFP cells were plated in 24-well plates and left to adhere overnight. Cells were

then treated with 10 μM Latrunculin B (Sigma) or the equivalent volume of DMSO (Sigma). After

1 h, MR1/5-OP-RU or MR1/6-FP monomer was added to a final concentration of 100 nM and the

cells were incubated for another 4 h before staining as above. For Brefeldin A blocking

experiments, BEAS-2B.MR1-GFP cells were incubated with 94 μM 6-formylpterin (6-FP, Schircks)

or the equivalent volume of 0.01 M NaOH (Sigma) overnight at 37°C at 5e4/well in a 24-well plate.

6-FP was washed off and the cells treated with 50 μg/ml Brefeldin A (BFA, Sigma) or the

equivalent volume of DMSO (Sigma) for 1 h, followed by the addition of MR1/6-FP or MR1/5-OP-

RU monomer (100 nM) or free 5-OPRU (5 μM) or 6-FP (100 μM) and incubation at 37°C for 4 h.

Cells were then stained with TCR tetramer as above. The gating strategy is shown in

Supplementary Figure 10. TCR tetramer was used at 150 or 300 nM. BEAS-2B WT cells were used

for unstained and single color controls. Anti-MHC I antibody W6/32 (BioLegend, 311410) was

used for the APC single color control due to low MR1 expression on BEAS-2B WT cells. For surface

staining of BEAS-2B.MR1-GFP cells with streptavidin, 3e5 cells were incubated with 50 nM of

MR1/5-OP-RU monomer or medium overnight. These cells were split into multiple wells and

either stained with the TCR tetramer as described above or blocked with PBS + goat, human, and

bovine serum for at least 30 min on ice and then stained with Streptavidin-APC (Agilent) for 40

min on ice or with biotinylated anti-MR1 antibody 26.5 for 40 minutes on ice, followed by

Steptavidin-APC for 40 minutes on ice. For tetramer staining of the CD8 T cell clone D480-F6, 1e5

T cell clones were incubated with the HLA*B08:01/CFP10$_{3-11}$ tetramer at 1:500 (0.127 µg/well)

for 45 minutes at 37°C. For tetramer staining of the CD4 T cell clone D481-D1, 1e5 T cell clones

were incubated with the HLA-DRB1*14:03/Esat6$_{73-87}$ or HLA-DRB1*14:03/control at 6 µg/ml for

1 h at 37°C. Cells were subsequently stained with an antibody cocktail containing LIVE/DEAD

Fixable Dead Cell Stain Kit (Thermo Fisher), and surface stained with the antibodies to CD4 (BD

Biosciences, clone SK3 348789), CD8 (Biolegend, clone SK1344714), and CD3 (BD Biosciences,

clone SK7340663), for 30 minutes at 4°C. Samples were washed and fixed with 1% PFA.

Acquisition was performed using a BD FACSymphony, LSRII, or Canto cytometer. The gating

strategy is shown in Supplementary Figure 11. Flow cytometric analysis was performed using

FlowJo version 10.6.1 or 10.7.1 (Treestar, Ashland, OR, USA). Plots show GFP+ gates to ensure

only cells over-expressing MR1 are included in the analysis.

Knock down of candidate chaperones

For siRNA-mediated knock down of candidate chaperones, siRNAs targeting TAPBPR (Assay ID

s229969), tapasin (Assay ID 41850), or AP2A1 (Assay ID s183) or a missense control (4390844)

(all Thermo Fisher Scientific) were nucleofected into 1e6 BEAS-2B WT cells at 300 nM according

to the manufacturer's instructions (Lonza, Nucleofector Kit T). Cells were allowed to rest

overnight (TAPBPR and tapasin) or two days (AP2A1) before use in an ELISPOT assay as

described above. RNA levels of targeted transcripts were measured by qRT PCR at the time of

ELISPOT using Taqman probes Hs00794155_m1, Hs00917451_g1, and Hs00900330_m1 with

GAPDH (Hs02758991_g1) as the housekeeping gene (all Thermo Fisher Scientific). For AP2A1,

equal cell numbers were also lysed in Lysis Buffer (Chromotek) with Protease Inhibitor cocktail

(Roche) at the time of ELISPOT and lysates of all three experiments were analyzed together by

Western blot. Briefly, lysates were combined with Loading Buffer and Reducing Agent (both

Invitrogen), separated on 4-20% polyacrylamide gels (Bio-Rad), and transferred onto PVDF

membrane (Millipore). The membrane was blocked with Blocking Buffer (Odyssey) for 1 h at

room temperature before incubation with mouse anti-AP2A1 (BD, 610501) and rabbit anti-

ACTB (Abcam, ab8227) primary antibodies overnight at 4° C. The blot was developed with LI-

COR IRDye secondary antibodies (goat anti-mouse 800CW and goat anti-rabbit 680RD) and

imaged simultaneously on an Odyssey imager.

Recombinant protein expression and purification

All experiments using MR1 monomer and tetramer were performed with the reagent(s)

provided by the NIH Tetramer Core Facility (NTCF). Recombinant DP10.7 γδ TCR and HLA-

A2/MART-1 peptide used as controls in the ELISA were produced as previously described [63,

64]. The D426-G11 TCR was produced as previously described for other MAIT TCRs [65]. Briefly,

the D426-G11 TCR V(D)J region of each TCR chain was cloned separately into the pAcGP67a

vector after the GP67 signal sequence and before either a "cysteine-optimized" TRAC or TRBC

sequence [66]. To ensure heterodimer formation, the chains were tagged with cleavable C-

terminal acidic and basic zippers [67] each followed with a 6xHis tag. The alpha chain construct

contained a C-terminal AviTag before the zipper cleavage site. The P1 and P2 baculoviruses

were produced in Sf9 cells [56], then the baculoviruses were added in tandem to Hi-5 cells and

the cells were allowed to express the recombinant protein for 72 h. The supernatant was

harvested and the protein purified by Ni-NTA chromatography. Recombinant TCR was then

treated with recombinant HRV 3C protease overnight and the cleaved zippers were removed

with Ni-NTA. The protein was biotinylated *in vitro* using recombinant BirA ligase and the

biotinylation efficiency checked by a streptavidin gel-shift assay. Finally, it was purified by size

exclusion chromatography and tetramerized with streptavidin-APC (Agilent). This reagent is

available upon request after the completion of an MTA that includes a clear description of how

the material would be used.

Microscopy

BEAS-2B cells (1e5) were seeded on a chamber slide and incubated for 18 h at 37C. MR1/5-OP-

RU tetramer conjugated to AF647 (200 nM) was added to the experimental wells for 2 h at RT.

Control wells received no tetramer. To determine whether the tetramer was observed in the

cells, CellMask Orange Plasma (1:5000 dilution, Invitrogen) was added for 10 min. To examine

whether the tetramer localized to endosomal compartments, CellLight reagents (Thermofisher)

for early endosomes (Rab5) or lysosomes (LAMP1) were added for 18 h prior to addition of the

tetramer. For all imaging, DAPI was added during PBS washes, and cells were fixed prior to data

acquisition. Images were acquired on a Keyence BZ-X810 Fluorescence microscope or high-

resolution wide-field CoreDV microscope (Applied Precision) with CoolSNAP ES2 HQ (Nikon),

using nuclear staining with DAPI to identify un-biased fields of view. Localization of the tetramer

to endosomal compartments was quantified using Imaris image analysis software as previously

described [36].

Cell-free acidification assay

In addition to MR1/5-OP-RU and MR1/6FP from the NIH Tetramer Core, recombinant,

biotinylated γδ TCR ("DP10.7") [63] and HLA-A2:MART-1 [64] were made in-house as previously

described and used as controls for this ELISA assay. Extensive washing with PBS pH 7.4 + 0.5%

Tween-20 was performed between each step/treatment outlined below. First, traptavidin, a

variant of streptavidin with a slower biotin off-rate [68], was immobilized on an ELISA plate in 0.1

M Na_2CO_3, pH 9.6 overnight. The plate was blocked with 0.5% BSA in 1xPBS, then biotinylated

MR1, TCR, and HLA-A2 were immobilized as appropriate. Eight working buffers of 1xPBS were

made from a house-made 10xPBS stock spanning pH 4.6-7.4. Protein was treated with either PBS

pH 7.4 (controls) or PBS of variable pH overnight at 4°C. (This experiment was also performed

with a citrate-phosphate buffer system, which unlike PBS, has a buffering range spanning the

experimental conditions, and the results did not differ from those using PBS, which has a

buffering range of ~5.8-7.4 pH units.) After removal of the acidic buffer and washing with PBS pH

7.2, either anti-human MR1 (unlabeled, clone 26.5, BioLegend 361102) or anti-human β_2m

(unlabeled, clone B2M-01, Novus Biologicals NB500-317) antibody was applied to the plate,

followed by a rabbit anti-mouse IgG H+L, HRP-conjugated secondary antibody (Abcam ab6728).

The ELISA was developed with a TMB substrate kit (Thermo Scientific 34021) and the absorbance

was detected at 450 nm using a Tecan Safire 2 plate reader.

Data analysis and Statistics

Data were analyzed with Prism 9 GraphPad Software. For measurements of functional avidity,

curves were determined by non-linear regression and best-fit EC_{50} was calculated. EC_{50} values

were compared with the extra sum-of-squares F test.

Data Availability

645 The authors declare that the data supporting the findings of this study are available within the

646 paper and its supplementary information files.

Acknowledgements

We would like to thank the participants who gave time and dedication to this health research as

well as Erin Merrifield, Department of Pediatrics, OHSU, for her contributions to this study. We

also thank Sobhan Roy and Mayuri Viswanathan for kindly providing recombinant γδ TCR and

HLA-A2 for use in this study.

This project has been funded in whole or in part with Federal funds from the National Institutes

of Allergy and Infectious Diseases, National Institutes of Health, Department of Health and

Human Services, under grant no AI147954 (EA, DML, WHH), AI134790 (DML), GM007183 (NAL),

AI129976 (MH), 1R01AI141549 (FT) and AI165618 (AO). This work was also supported in part by

Merit Award #I01 CX001562 from the U.S. Department of Veterans Affairs Clinical Sciences

Research and Development Program (MJH), Merit Award #I01 BX000533 from the U.S.

Department of Veterans Affairs Biomedical Laboratory (DML).

Author Contributions

CK, GS, NAL, DAL, EJA, DML and MH contributed to the conception and/or design of the work. CK and GS are joint first authors. DML and MH jointly supervised the work. MC, MN, DAL and DML contributed to clinical activities. AO, WHH, FT, EJA, DML and MH raised grants to fund the research. CK, GS, NAL, MC, MN, AW, CL, TA, JB, TNL, CMH, MEH, JK, LA, AO, WHH, SM, FT, DAL, EJA, DML and MH substantially contributed to the acquisition, analysis, or interpretation of data and drafting of the manuscript. All authors substantially contributed to revising and critically reviewing the manuscript for important intellectual content. All authors approved the final version of this manuscript to be published and agree to be accountable for all aspects of the work.

References

1. Crowther, M.D. and A.K. Sewell, *The burgeoning role of MR1-restricted T-cells in infection, cancer and autoimmune disease.* Curr Opin Immunol, 2021. **69**: p. 10-17.
2. Hinks, T.S.C. and X.W. Zhang, *MAIT Cell Activation and Functions.* Front Immunol, 2020. **11**: p. 1014.
3. Trowitzsch, S. and R. Tampé, *Multifunctional Chaperone and Quality Control Complexes in Adaptive Immunity.* Annual Review of Biophysics, 2020. **49**(1): p. 135-161.
4. Alvaro-Benito, M. and C. Freund, *Revisiting nonclassical HLA II functions in antigen presentation: Peptide editing and its modulation.* HLA, 2020. **96**(4): p. 415-429.
5. Moody, D.B. and R.N. Cotton, *Four pathways of CD1 antigen presentation to T cells.* Curr Opin Immunol, 2017. **46**: p. 127-133.
6. McWilliam, H.E., et al., *The intracellular pathway for the presentation of vitamin B-related antigens by the antigen-presenting molecule MR1.* Nat Immunol, 2016. **17**(5): p. 531-7.
7. McWilliam, H.E.G., et al., *Endoplasmic reticulum chaperones stabilize ligand-receptive MR1 molecules for efficient presentation of metabolite antigens.* Proc Natl Acad Sci U S A, 2020. **117**(40): p. 24974-24985.
8. Corbett, A.J., et al., *T-cell activation by transitory neo-antigens derived from distinct microbial pathways.* Nature, 2014. **509**(7500): p. 361-5.
9. Harriff, M.J., et al., *Endosomal MR1 Trafficking Plays a Key Role in Presentation of Mycobacterium tuberculosis Ligands to MAIT Cells.* PLoS Pathog, 2016. **12**(3): p. e1005524.
10. Karamooz, E., et al., *MR1 recycling and blockade of endosomal trafficking reveal distinguishable antigen presentation pathways between Mycobacterium tuberculosis infection and exogenously delivered antigens.* Scientific Reports, 2019. **9**(1): p. 4797.
11. Kulicke, C., et al., *Covering All the Bases: Complementary MR1 Antigen Presentation Pathways Sample Diverse Antigens and Intracellular Compartments.* Frontiers in Immunology, 2020. **11**.

12. Lange, J., et al., *The Chemical Synthesis, Stability, and Activity of MAIT Cell Prodrug Agonists*
That Access MR1 in Recycling Endosomes. ACS Chemical Biology, 2020.

13. Ussher, J.E., et al., *TLR signaling in human antigen-presenting cells regulates MR1-*
dependent activation of MAIT cells. European Journal of Immunology, 2016. **46**(7): p. 1600-
1614.

14. McWilliam, H.E.G. and M. Salio, *Understanding and modulating the MR1 metabolite*
antigen presentation pathway. Molecular Immunology, 2021. **129**: p. 121-126.

15. Harriff, M.J., et al., *MR1 displays the microbial metabolome driving selective MR1-restricted*
T cell receptor usage. Sci Immunol, 2018. **3**(25).

16. Mak, J.Y., et al., *Stabilizing short-lived Schiff base derivatives of 5-aminouracils that activate*
mucosal-associated invariant T cells. Nat Commun, 2017. **8**: p. 14599.

17. Priya, R. and R.R. Brutkiewicz, *MR1 Tetramer-Based Artificial APCs Expand MAIT Cells from*
Human Peripheral Blood That Effectively Kill Glioblastoma Cells. Immunohorizons, 2021.
5(6): p. 500-511.

18. Cruz, F.M., A. Chan, and K.L. Rock, *Pathways of MHC I cross-presentation of exogenous*
antigens. Semin Immunol, 2023. **66**: p. 101729.

19. Gold, M.C., et al., *Human mucosal associated invariant T cells detect bacterially infected*
cells. PLoS Biol, 2010. **8**(6): p. e1000407.

20. Narayanan, G.A., et al., *Alternative splicing of MR1 regulates antigen presentation to MAIT*
cells. Sci Rep, 2020. **10**(1): p. 15429.

21. Salio, M., et al., *Ligand-dependent downregulation of MR1 cell surface expression.* Proc Natl
Acad Sci U S A, 2020. **117**(19): p. 10465-10475.

22. Lion, J., et al., *MR1B, a natural spliced isoform of the MHC-related 1 protein, is expressed*
as homodimers at the cell surface and activates MAIT cells. Eur J Immunol, 2013. **43**(5): p.
1363-73.

23. Salerno-Goncalves, R., T. Rezwan, and M.B. Sztein, *B cells modulate mucosal associated*
invariant T cell immune responses. Front Immunol, 2014. **4**: p. 511.

24. Howson, L.J., et al., *Absence of mucosal-associated invariant T cells in a person with a*
homozygous point mutation in MR1. Sci Immunol, 2020. **5**(49).

25. Eckle, S.B., et al., *A molecular basis underpinning the T cell receptor heterogeneity of*
mucosal-associated invariant T cells. J Exp Med, 2014. **211**(8): p. 1585-600.

26. Lewinsohn, D.A., et al., *Immunodominant tuberculosis CD8 antigens preferentially*
restricted by HLA-B. PLoS Pathog, 2007. **3**(9): p. 1240-9.

27. Lancioni, C., et al., *Recognition of CD8(+) T-cell epitopes to identify adults with pulmonary*
tuberculosis. Eur Respir J, 2019. **53**(5).

28. Adams, S., et al., *HLA class I and II genotype of the NCI-60 cell lines.* J Transl Med, 2005.
3(1): p. 11.

29. Le Bourhis, L., et al., *Antimicrobial activity of mucosal-associated invariant T cells.* Nat
Immunol, 2010. **11**(8): p. 701-8.

30. Huang, S., et al., *Evidence for MR1 antigen presentation to mucosal-associated invariant T*
cells. J Biol Chem, 2005. **280**(22): p. 21183-93.

31. Kjer-Nielsen, L., et al., *MR1 presents microbial vitamin B metabolites to MAIT cells.* Nature,
2012. **491**(7426): p. 717-23.

32.   Bowman, E.J., A. Siebers, and K. Altendorf, *Bafilomycins: a class of inhibitors of membrane*
739   *ATPases from microorganisms, animal cells, and plant cells.* Proc Natl Acad Sci U S A, 1988.
740   **85**(21): p. 7972-6.

741   33.   Sadegh-Nasseri, S., *Partnering for the major histocompatibility complex class II and*
742   *antigenic determinant requires flexibility and chaperons.* Curr Opin Immunol, 2021. **70**: p.
743   112-121.

744   34.   McShan, A.C., et al., *TAPBPR employs a ligand-independent docking mechanism to*
745   *chaperone MR1 molecules.* Nat Chem Biol, 2022. **18**(8): p. 859-868.

746   35.   Lim, H.J., et al., *A specialized tyrosine-based endocytosis signal in MR1 controls antigen*
747   *presentation to MAIT cells.* J Cell Biol, 2022. **221**(12).

748   36.   Harriff, M.J., et al., *Human Lung Epithelial Cells Contain Mycobacterium tuberculosis in a*
749   *Late Endosomal Vacuole and Are Efficiently Recognized by CD8+ T Cells.* PLOS ONE, 2014.
750   **9**(5): p. e97515.

751   37.   Scheller, R.H., *In search of the molecular mechanism of intracellular membrane fusion and*
752   *neurotransmitter release.* Nat Med, 2013. **19**(10): p. 1232-5.

753   38.   Spessott, W.A., et al., *Syntaxin 4 mediates endosome recycling for lytic granule exocytosis*
754   *in cytotoxic T-lymphocytes.* Traffic, 2017. **18**(7): p. 442-452.

755   39.   Kennedy, M.J., et al., *Syntaxin-4 defines a domain for activity-dependent exocytosis in*
756   *dendritic spines.* Cell, 2010. **141**(3): p. 524-35.

757   40.   Chatterjee Bhowmick, D., et al., *Conventional and Unconventional Mechanisms by which*
758   *Exocytosis Proteins Oversee beta-cell Function and Protection.* Int J Mol Sci, 2021. **22**(4).

759   41.   Black, H.L., et al., *Knockout of syntaxin-4 in 3T3-L1 adipocytes reveals new insight into*
760   *GLUT4 trafficking and adiponectin secretion.* J Cell Sci, 2022. **135**(1).

761   42.   Lamichhane, R. and J.E. Ussher, *Expression and trafficking of MR1.* Immunology, 2017.
762   **151**(3): p. 270-279.

763   43.   Karamooz, E., M.J. Harriff, and D.M. Lewinsohn, *MR1-dependent antigen presentation.*
764   Seminars in Cell & Developmental Biology, 2018. **84**: p. 58-64.

765   44.   Blander, J.M., *Regulation of the Cell Biology of Antigen Cross-Presentation.* Annu Rev
766   Immunol, 2018. **36**: p. 717-753.

767   45.   Kinsella, R.L., et al., *Perspectives and Advances in the Understanding of Tuberculosis.* Annu
768   Rev Pathol, 2021. **16**: p. 377-408.

769   46.   Ackerman, A.L., et al., *Access of soluble antigens to the endoplasmic reticulum can explain*
770   *cross-presentation by dendritic cells.* Nat Immunol, 2005. **6**(1): p. 107-13.

771   47.   Spooner, R.A., et al., *Retrograde transport pathways utilised by viruses and protein toxins.*
772   Virol J, 2006. **3**: p. 26.

773   48.   Mukhopadhyay, S. and A.D. Linstedt, *Retrograde trafficking of AB_5 toxins: mechanisms to*
774   *therapeutics.* J Mol Med (Berl), 2013. **91**(10): p. 1131-41.

775   49.   Hafstrand, I., A. Aflalo, and L.H. Boyle, *Why TAPBPR? Implications of an additional player in*
776   *MHC class I peptide presentation.* Current Opinion in Immunology, 2021. **70**: p. 90-94.

777   50.   Thomas, C. and R. Tampe, *MHC I assembly and peptide editing - chaperones, clients, and*
778   *molecular plasticity in immunity.* Curr Opin Immunol, 2021. **70**: p. 48-56.

779   51.   Kulicke, C.A., et al., *The P5-type ATPase ATP13A1 modulates major histocompatibility*
780   *complex I-related protein 1 (MR1)-mediated antigen presentation.* J Biol Chem, 2022.
781   **298**(2): p. 101542.

782   52.   Pankhurst, T.E., et al., *MAIT cells activate dendritic cells to promote T(FH) cell differentiation*
783   *and induce humoral immunity.* Cell Rep, 2023. **42**(4): p. 112310.

784   53.   Lewinsohn, D.M., et al., *Mycobacterium tuberculosis-reactive CD8+ T lymphocytes: the*
785   *relative contribution of classical versus nonclassical HLA restriction.* J Immunol, 2000.
786   **165**(2): p. 925-30.

787   54.   Laugel, B., et al., *Engineering of Isogenic Cells Deficient for MR1 with a CRISPR/Cas9*
788   *Lentiviral System: Tools To Study Microbial Antigen Processing and Presentation to Human*
789   *MR1-Restricted T Cells.* J Immunol, 2016. **197**(3): p. 971-82.

790   55.   Olive, A.J., et al., *Chlamydia trachomatis-induced alterations in the host cell proteome are*
791   *required for intracellular growth.* Cell Host Microbe, 2014. **15**(1): p. 113-24.

792   56.   Murhammer, D.W.e., *Baculovirus and insect cell expression protocols.* Methods in
793   Molecular Biology. Vol. 388. 2007, Totowa, NJ: Humana Press.

794   57.   Tafesse, F.G., et al., *Disruption of Sphingolipid Biosynthesis Blocks Phagocytosis of Candida*
795   *albicans.* PLoS Pathog, 2015. **11**(10): p. e1005188.

796   58.   Thomas, S.T., et al., *Fetal Liver-Derived Alveolar-like Macrophages: A Self-Replicating Ex*
797   *Vivo Model of Alveolar Macrophages for Functional Genetic Studies.* Immunohorizons,
798   2022. **6**(2): p. 156-169.

799   59.   Brinkman, E.K., et al., *Easy quantitative assessment of genome editing by sequence trace*
800   *decomposition.* Nucleic Acids Res, 2014. **42**(22): p. e168.

801   60.   Conant, D., et al., *Inference of CRISPR Edits from Sanger Trace Data.* Crispr j, 2022. **5**(1): p.
802   123-130.

803   61.   Heinzel, A.S., et al., *HLA-E-dependent presentation of Mtb-derived antigen to human CD8+*
804   *T cells.* J Exp Med, 2002. **196**(11): p. 1473-81.

805   62.   Li, K., et al., *Synthesis, stabilization, and characterization of the MR1 ligand precursor 5-*
806   *amino-6-D-ribitylaminouracil (5-A-RU).* PLoS One, 2018. **13**(2): p. e0191837.

807   63.   Luoma, A.M., et al., *Crystal structure of Vδ1 T cell receptor in complex with CD1d-sulfatide*
808   *shows MHC-like recognition of a self-lipid by human γδ T cells.* Immunity, 2013. **39**(6): p.
809   1032-42.

810   64.   Benveniste, P.M., et al., *Generation and molecular recognition of melanoma-associated*
811   *antigen-specific human γδ T cells.* Sci Immunol, 2018. **3**(30).

812   65.   López-Sagaseta, J., et al., *The molecular basis for Mucosal-Associated Invariant T cell*
813   *recognition of MR1 proteins.* Proc Natl Acad Sci U S A, 2013. **110**(19): p. E1771-8.

814   66.   Boulter, J.M., et al., *Stable, soluble T-cell receptor molecules for crystallization and*
815   *therapeutics.* Protein Eng, 2003. **16**(9): p. 707-11.

816   67.   Chang, H.C., et al., *A general method for facilitating heterodimeric pairing between two*
817   *proteins: application to expression of alpha and beta T-cell receptor extracellular segments.*
818   Proc Natl Acad Sci U S A, 1994. **91**(24): p. 11408-12.

819   68.   Chivers, C.E., et al., *A streptavidin variant with slower biotin dissociation and increased*
820   *mechanostability.* Nat Methods, 2010. **7**(5): p. 391-3.

 Figures

Fig 1. MR1 ligands delivered to antigen presenting cells on soluble MR1 molecules are efficiently recognized by MR1T cell clones. A. MR1T cell clone (1e4) IFN-γ responses to MR1/5-OP-RU or MR1/6-FP tetramer, either added to DCs (1e4) (exSPOT) or bound to ELISPOT plate (tetraSPOT). **B.** MR1T cell clone (1e4) IFN-γ response to MR1/5-OP-RU is inhibited by anti-MR1 antibody blockade (2 μg/ml). **C**. MR1T cell clone (1e4) IFN-γ response to MR1/5-OP-RU tetramer, added to either DCs, THP-1 cells, BEAS-2B (B2B) cells or B-LCL (1e4). EC_{50} calculated using non-linear regression and the difference in the EC_{50} was statistically significant between the curves (p<0.0001) by the extra sum-of-squares F test. Data are representative of three independent experiments each. Error bars denote SD between technical duplicates within the representative experiment shown. For EC_{50} values of replicate experiments see Supplementary Tables 1+2.

Fig 2. Ligand delivered in the context of MR1 monomer is presented more efficiently than free
ligand. A. MR1T cell clone (1e4) IFN-γ response to DCs incubated with MR1/5-OP-RU tetramer,
MR1/5-OP-RU monomer, and soluble 5-OP-RU. EC_{50} calculated using non-linear regression. The
difference in the EC_{50} was statistically significant between the MR1/5-OP-RU tetramer and 5-OP-
RU and between MR1/5-OP-RU monomer and 5-OP-RU curves (p<0.0001), but there was no
significant difference in the EC_{50} of MR1/5-OP-RU tetramer and monomer by the extra sum-of-
squares F test (p=0.7486). Data are representative of three independent experiments. Error bars
denote SD between technical duplicates within the representative experiment shown. For EC_{50}
values of replicate experiments see Supplementary Table 3. **B.** BEAS-2B.MR1-GFP cells were
incubated with the indicated concentrations of MR1/5-OP-RU monomer or free 5-OP-RU
overnight and surface stained for flow cytometry with APC-conjugated TCR tetramer.
Representative histograms are shown on the left with pooled geometric mean fluorescence
intensity from four experiments shown on the right. The same unstained and untreated control
are shown in both histogram plots for easy comparison. Error bars denote SD.

873
874
875
Fig 3. Presentation of MR1 ligands delivered on soluble MR1 molecules requires cellular MR1.
A. MR1T cell clone (1e3) IFN-γ response to BEAS-2B, BEAS-2B MR1[-/-] and BEAS-2B MR1[-/-] : MR1A
(1e4) incubated with MR1/5-OP-RU or MR1/6-FP monomers. Also included MR1T clone IFN-γ
response to MR1/5-OP-RU monomer only with no antigen presenting cells. **B.** Same as **A.**, except
the antigen presenting cells are THP-1, THP-1 MR1[-/-] and THP-1 MR1[-/-] : MR1A (1e4). Data are
representative of two independent experiments. Error bars denote SD of technical duplicates
within the representative experiments shown.
883

884
885
Fig 4. Ligands are transferred from the soluble MR1 molecules onto cellular MR1.
A. BEAS-2B.MR1-GFP cells (3e5) were pre-incubated with 50 nM of MR1/5-OP-RU monomer
overnight before staining with either Streptavidin (left), TCR tetramer (middle) or biotinylated
26.5 anti-MR1 antibody followed by Streptavidin (right). The same unstained sample is shown in
all panels and the same primary antibody only controls are shown in the left and right panels for
easy comparison. Data are representative of three independent experiments. **B.** BEAS-2B.MR1-
GFP cells were pre-incubated with 10 µM of latrunculin B (LatB) for 1 h before addition of MR1/5-
OP-RU or MR1/6-FP monomer for an additional 4 h. Cells were surface stained with TCR tetramer.
Representative histograms are shown on the left with pooled geometric mean fluorescence
intensity from three experiments shown on the right. The same unstained and untreated control
are shown in both histogram plots for easy comparison. Error bars denote SD. **C.** BEAS-2B.MR1-
GFP cells were incubated with either 50 nM MR1/5-OP-RU monomer or 6.4 µM of free 5-OP-RU
for indicated amounts of time before surface staining for flow cytometry with TCR tetramer.
Representative histograms are shown on the left with pooled geometric mean fluorescence
intensity from three experiments shown on the right. Error bars denote SD. **D.** MR1T cell clone
(1e3) IFN-γ response to BEAS-2B MR1$^{-/-}$ cells (1e4) that had been transfected with the MR1 or
MR1 R9H plasmid, followed by addition of the MR1/5-OP-RU monomer, or 1 µL of *M. smegmatis*
supernatant. Data are representative of three independent experiments. Error bars denote SD
between technical duplicates within the representative experiment shown.
905
906
907

908
909
Fig 5. Ligand exchange can occur in post-ER compartments. A. MR1T cell clone (1e4) IFN-γ
response to BEAS-2B.MR1-GFP (1e4) that were pretreated with 6-FP *vs* NaOH using MR1/5-OP-
RU monomer as the antigen. EC_{50} calculated using non-linear regression and the difference in the
EC_{50} was statistically significant between the curves (p<0.0001) by the extra sum-of-squares F
test. For EC_{50} values of replicate experiments see Supplementary Table 4. **B.** HLA*B08:01-
restricted T cell clone D480-F6 was stained with the HLA-B*08:01 tetramer to $CFP10_{3-11}$, along
with antibodies to CD3, CD4, CD8 and a live/dead discriminator. Live, $CD3^+CD4^-CD8^+$ T cells are
shown.　**C.** HLA*B08:01-restricted T cell clone D480-F6 (5e3) IFN-γ response to DCs (1e4) that
were incubated with either the $CFP10_{3-11}$ tetramer or $CFP10_{3-11}$ peptide. 9.5e4 irrelevant T cells
(MR1-restricted D426-G11) were also added to the well to decrease T:T presentation of the
classical T cell clone. **D.** MHC II-restricted T cell clone D481-D1 was stained with the HLA-
DRB1*14:03 tetramer to $Esat6_{73-87}$ or a control tetramer along with antibodies to CD3, CD4, CD8
and a live/dead discriminator. Live, $CD3^+CD8^-CD4^+$ T cells are shown.　**E.** MHC II-restricted T cell
clone D481-D1 (2e4) IFN-γ response to D481 (autologous) dendritic cells (2e4) that were
incubated with either the $Esat6_{73-87}$ tetramer, control tetramer or $Esat6_{73-87}$ peptide. **F.** MHC II-
restricted D481-D1 T cell clone (2e4) IFN-γ response to D481 (autologous) dendritic cells, D520
mismatched dendritic cells or A549 cells (2e4) that were incubated with the $Esat6_{73-87}$ tetramer.
G. BEAS-2B.MR1-GFP cells were pre-incubated with 6-FP or NaOH overnight. 6-FP was washed
off and cells were incubated with 50 ug/ml brefeldin A (BFA) for 1h before addition of MR1/5-
OP-RU or MR1/6-FP monomer for an additional 4 h. Cells were surface stained for flow cytometry
with TCR tetramer. Representative histograms are shown on the left with pooled geometric mean

fluorescence intensity from three experiments shown on the right. The same unstained control is shown in all plots. Error bars denote SD. Data in A, C, E and F are representative of three independent experiments. Error bars denote SD between technical duplicates within the representative experiment shown.

954
955

Fig. 6. Ligand exchange does not require acidification.
A. Recombinant, biotinylated MR1 monomer, γδ TCR, or HLA-A2 were immobilized on a
traptavidin-coated plate before incubation with PBS at the indicated pH overnight at 4°C. Binding
of either anti-MR1 (clone 26.5) or anti-β_2m (clone B2M-01) antibody was assessed with an HRP-
conjugated secondary antibody and TMB substrate. A recombinant γδ TCR was included as a
negative control and HLA-A2 was included as a positive control for the anti-β_2m antibody and a
negative control for the anti-MR1 antibody. Error bars denote SD of 3 replicates. **B.** MR1T cell
clone (5e3) IFN-γ response to BEAS-2B (1e4) pre-treated with Bafilomycin A1 for 2 h, followed by
addition of the MR1/5-OP-RU monomer or 5-OP-RU prodrug. Data in **B** are representative of
three independent experiments. Error bars denote SD between technical duplicates within the
representative experiment shown. **C**. BEAS-2B cells (1e5 or 1e4) were seeded on chamber slide
and CellLight RFP reagents for Rab5 and LAMP1 were added and incubated for 18 h at 37C.
Tetramer was then added for 2 h at RT, followed by washing 2X in PBS. All images acquired were
using the same microscope settings. Data are representative of three independent experiments.
Error bars denote SD from a representative experiment. **D**. MR1T cell clone (1e3) IFN-γ response
to BEAS-2B.Cas9 or BEAS-2B.Cas9 STX4 KO cells (1e4) in the presence of the indicated
concentrations of *M. smegmatis* supernatant or MR1/5-OP-RU monomer. Data were normalized
to the maximum response and pooled from three experiments. Error bars denote SD. EC_{50}
calculated using non-linear regression and the difference in the EC_{50} was statistically significant
between the curves (p=0.0032) by the extra sum-of-squares F test.