Optimised DNA isolation from marine sponges for natural sampler DNA (nsDNA) metabarcoding

Lynsey R. Harper[1,2*], Erika F. Neave[1,4], Graham S. Sellers[3], Alice V. Cunnington[1], María Belén Arias[4], Jamie Craggs[4], Barry MacDonald[5], Ana Riesgo[4,6], Stefano Mariani[1]

[1] School of Biological and Environmental Sciences, Liverpool John Moores University, Liverpool, L3 3AF, UK

[2] NatureMetrics Ltd, 1 Occam Court, Surrey Research Park, Guildford, GU2 7HJ, UK

[3] Department of Biological and Marine Sciences, University of Hull, Kingston upon Hull, HU6 7RX, UK

[4] Department of Life Sciences, The Natural History Museum, London, SW7 5BD, UK

[5] Fisheries and Oceans Canada, 1 Challenger Drive, Dartmouth, Nova Scotia, B2Y 4A2, Canada

[6] Department of Biodiversity and Evolutionary Biology, Museo Nacional de Ciencias Naturales (CSIC), C/José Gutiérrez Abascal 2, 28006, Madrid, Spain

*Corresponding author:

Lynsey R. Harper

lynsey.harper2@gmail.com

Running title: DNA extraction for sponge nsDNA metabarcoding

Abstract

Marine sponges have recently been recognised as natural samplers of environmental DNA (eDNA) due to their effective water filtration and their ubiquitous, sessile and regenerative nature. However, laboratory workflows for metabarcoding of sponge tissue have not been optimised to ensure that these natural samplers achieve their full potential for community survey. We used a phased approach to investigate the influence of DNA extraction procedures on the biodiversity information recovered from sponges. In Phase 1, we compared three treatments of residual ethanol preservative in sponge tissue alongside five DNA extraction protocols. The results of Phase 1 informed which ethanol treatment and DNA extraction protocol should be used in Phase 2, where we assessed the effect of starting tissue mass on extraction success and whether homogenisation of sponge tissue is required. Phase 1 results indicated that ethanol preservative may contain unique and/or additional biodiversity information to that present in sponge tissue, but blotting tissue dry generally recovered more taxa and generated more sequence reads from the wild sponge species. Tissue extraction protocols performed best in terms of DNA concentration, taxon richness and proportional read counts, but the non-commercial tissue protocol was selected for Phase 2 due to cost-efficiency and greater recovery of target taxa. In Phase 2 overall, we found that homogenisation may not be required for sponge tissue and more starting material does not necessarily improve taxon detection. These results combined provide an optimised DNA isolation procedure for sponges to enhance marine biodiversity assessment using natural sampler DNA metabarcoding.

Keywords: biodiversity, DNA extraction, environmental DNA, fish, marine, natural samplers, Porifera, vertebrates

1. Introduction

The emergence of environmental DNA (eDNA) analysis signalled a new age of biodiversity monitoring, facilitating faster, non-invasive, cost-efficient and comprehensive surveys for single species and whole communities than was previously thought possible (Lawson Handley 2015; Deiner et al., 2017; Miya 2021). Such has been the rapid, exponential growth of this field in the last decade (Tsuji et al., 2019; Thalinger et al., 2021) that it is now beginning to branch. In addition to the now relatively standard analyses of aquatic eDNA and soil/sediment eDNA, studies have been published on species detection using eDNA extracted from snow (Franklin et al., 2019; Kinoshita et al., 2019) and air (Clare et al., 2021; Lynggaard et al., 2022; Roger et al., 2022). Innovative research has also demonstrated that water can be used to aggregate eDNA deposited on plants (Valentin et al., 2020) or fishing nets (Russo et al., 2020) for species detection. Less closely intertwined but stemming from the same roots is natural sampler DNA (nsDNA) (Siegenthaler et al., 2019), which refers to the analysis of DNA collected by live organisms. This is not a novel concept as researchers have been analysing the DNA present in faeces and gut contents of consumers (Pompanon et al., 2012; Nørgaard et al., 2021) and hematophagous insects (Schnell et al., 2012; Massey et al., 2021) for years, but more creative applications are now being explored, including flowers and cow dung as samplers of insect assemblages (Thomsen & Sigsgaard, 2019; Sigsgaard et al., 2020; Harper et al., 2021), dung beetles as samplers of mammal communities (Drinkwater et al., 2021), and sponges as samplers of marine biodiversity (Mariani et al., 2019; Turon et al., 2020; Jeunen et al., 2021).

Marine sponges can filter thousands of litres of water in a single day (Gökalp et al., 2020), vastly exceeding the volume that could be collected with artificial devices for aquatic eDNA metabarcoding. As water is filtered, particles are trapped and concentrated in sponge tissue until digestion or excretion, meaning that a biopsy can be taken to provide DNA for metabarcoding analysis (Mariani et al., 2019; Turon et al., 2020; Jeunen et al., 2021). Sponge nsDNA metabarcoding offers several advantages over aquatic eDNA metabarcoding. The high-tech robotics being used to sample large volumes of water for aquatic eDNA (e.g. automated underwater vehicles) are expensive

to build and operate, are ineffective for inaccessible and/or complex habitats, and often
target a limited set of taxa (Mariani et al., 2019). Although passive filtration has been
explored as an alternative to active filtration using peristaltic or vacuum pumps, they
require long submergence times and have saturation limits (Kirtane et al., 2020; Bessey
et al., 2021b; a). Sponges are present in nearly every aquatic habitat, and sample
collection is inexpensive and non-destructive with minimal environmental impact.
Provided that biopsies are not destructive and conducted carefully, sponges will
regenerate quickly. Therefore, sponges offer a low-tech, affordable and standardised
method of surveying marine biodiversity, and could serve as long-term monitoring
stations (Mariani et al., 2019). However, there is much work to be done before sponges
can be reliably and routinely used in community surveys, specifically sampling and
laboratory protocols (Jeunen et al., 2021).

Two sponge nsDNA studies to date (Mariani et al., 2019; Turon et al., 2020)
utilised samples from existing collections and extracted DNA from tissue with
commercial kits. The other study (Jeunen et al., 2021) compared dive surveys and
aquatic eDNA to sponge tissue and the ethanol used to preserve sponge tissue, and
also extracted DNA using a commercial kit. It remains undetermined whether certain
sponge species are better samplers than others and if this is related to phenotype. The
spatial extent of biodiversity information recovered by a single sponge is yet to be
ascertained along with the optimal mass/volume for biopsies to maximise data
generation without being lethal to the sponge sampled (Mariani et al., 2019). The
influence of DNA extraction method on biodiversity detection from sponge tissues is
unknown (Jeunen et al., 2021), but this has been found to affect detection success with
aquatic eDNA (Deiner et al., 2018; Jeunen et al., 2019), bulk invertebrates (Hermans et
al., 2018; Majaneva et al., 2018) and faecal samples (Kaunisto et al., 2017).
Furthermore, it has not been investigated if samples should be homogenised to ensure
DNA is evenly distributed throughout the sample prior to extraction, similar to bulk
invertebrate and faecal samples (Gosselin et al., 2017; Pereira-da-Conceicoa et al.,
2020). The optimal amount of starting material to maximise species recovery without
introducing PCR inhibitors is also unknown, but using larger amounts of soil was found
to improve invertebrate detection (Kirse et al., 2021).

Here, we focus on the influence of laboratory workflows on the performance of
sponge nsDNA metabarcoding, specifically DNA isolation procedures. We used a
phased approach with three sponge species to investigate whether residual ethanol
preservative, DNA extraction protocol, type of starting material, and amount of starting
material influence sponge nsDNA metabarcoding performance using the following
extraction success criteria: 1) total DNA yield, 2) target DNA concentration, 3) taxon
richness, 4) proportional read counts, and 5) community composition. In Phase 1, we
compared three treatments of residual ethanol preservative in sponge tissue (no
removal, removal via blotting, removal via centrifugation) alongside five DNA extraction
protocols, including commercial kits and non-commercial protocols. Using the optimal
ethanol treatment and DNA extraction protocol from Phase 1, we compared different
volumes of homogenised sponge tissue to different weights of dried sponge tissue. We
synthesise the results from each experimental phase to provide an optimised DNA
isolation procedure for sponge tissue to enhance marine biodiversity assessment using
sponge nsDNA metabarcoding.

135   2. Methods

137   2.1 Workspace and decontamination procedures

A unidirectional workflow was used for sample processing. Sponge tissue was stored
and handled in a laboratory dedicated to the processing of environmental samples with
low DNA concentrations. Prepared PCR reactions were transported to separate
laboratories for: addition of PCR positive controls and PCR amplification; gel
electrophoresis, library preparation, and storage; and library quality checks,
quantification, and sequencing. Bench space and non-immersible equipment in all
laboratories were sterilised before and after use by wiping with fresh paper towel and
10% v/v bleach solution (made using Cleanline thin bleach containing 4.53% sodium
hypochlorite), followed by 70% v/v ethanol solution. Immersible equipment was
sterilised in a 10% v/v bleach bath for at least 10 minutes, then immersed in a 5% v/v
Lipsol detergent bath and rinsed with deionised water. Bleach and Lipsol detergent

baths were changed weekly or daily during periods of heavy use. Stainless steel 5 mm

beads (Qiagen, Germany) were additionally sterilised by heating at 220°C for 3 hours.

Plastics and hoods were sterilised with UV light for at least 30 minutes. Full details of

decontamination procedures are provided in Supporting Information.

2.2 Sponge selection

For this experiment, we used samples from three marine sponges which are the focus

of ongoing research employing nsDNA metabarcoding: 1) an invasive sponge present in

the aquarium facility of the Horniman Museum and Gardens, London, 2) *Vazella*

pourtalesii, which is a siliceous sponge that colonises hydroacoustic moorings in Nova

Scotia, Canada, and 3) *Phakellia ventilabrum*, which is an axinnellid that forms relatively

dense aggregations on rock-sand habitats and is broadly distributed across the North

Atlantic. The Horniman invasive sponge provided an opportunity to groundtruth nsDNA

metabarcoding in a controlled tank environment with 25 species of known abundance,

whereas *V. pourtalesii* and *P. ventilabrum* were collected from natural marine

environments with more diverse species assemblages and differed in phenotype as well

as filtration efficiency. The Horniman invasive sponge and *V. pourtalesii* were used in

Phase 1, whereas *V. pourtalesii* and *P. ventilabrum* were used in Phase 2 due to the

greater amounts of starting material required for DNA extraction.

2.3 Sponge collection

All sponges were detached from their substrate using a sterile diving knife or sterile

disposable surgical scalpel (Swann-Morton No. 21, Fisher Scientific, UK) whilst wearing

disposable gloves. Individuals of the Horniman invasive sponge were small thus an ~1

cm^3 biopsy was taken. Each biopsy was placed in a 2 ml DNA LoBind tube (Eppendorf,

Fisher Scientific, UK) containing 1 ml of molecular grade 100% ethanol (Fisher

Scientific, UK) and stored at -20°C for transport to Liverpool John Moores University

(LJMU). At LJMU, forceps were used to transfer each biopsy to a new 2 ml DNA LoBind

tube containing 1 ml of molecular grade 100% ethanol as water present in tissue may

have diluted the original ethanol. Samples were stored at -20°C for three months until

DNA extraction. As *V. pourtalesii* colonises hydroacoustic moorings, samples were

collected during mooring retrieval undertaken by Ocean Tracking Network and Fisheries

and Oceans Canada in July 2021. Whole individuals were removed from the moorings,

placed in sterile pots containing 500 ml of molecular grade 100% ethanol, and stored at

-20°C once research vessels returned to shore. The ethanol in each pot was changed

twice before samples were shipped to LJMU and stored at -20°C for three months until

DNA extraction. For *P. ventilabrum,* a ~5 cm^3 biopsy was taken from each individual

sampled from Rockall Bank (coordinates: 57.1705, -13.638) at 185-189 m depth in May

2018, then placed in a 50 ml falcon tube containing 25 ml of 100% ethanol and stored at

4°C. Again, the ethanol was changed twice before samples were shipped to LJMU in

January 2021 where they were stored at -20°C for three months until DNA extraction.

2.4 DNA extraction

2.4.1 Phase 1

Three Horniman invasive sponge samples and three *V. pourtalesii* samples were used

in Phase 1 (Fig. 1). Each sample from each sponge was subjected to a different ethanol

treatment as ethanol is known to interact with chemicals used during DNA extraction

and inhibit PCR amplification, but residual ethanol preservative may be present in

porous sponge tissue. The ethanol treatments were: 1) no removal of residual ethanol

(wet treatment), 2) removal of residual ethanol by blotting sponge tissue against filter

paper inside a petri dish (blot treatment), and 3) biopsies were transferred to a new 2 ml

DNA LoBind tube and centrifuged for 1 minute at maximum speed, following which the

ethanol was removed with a pipette and centrifugation repeated until no more ethanol

was observed (spin treatment). After each ethanol treatment, samples were cut into

small pieces within a petri dish using dissection scissors. Pieces were added to a weigh

boat until a maximum weight of 500 mg was observed. Each subsample was transferred

to a high-impact 2 ml screw cap microtube (Starlab, UK) containing a 5 mm stainless

steel bead and twice the amount of 1x TE buffer to tissue weight was added to each

microtube (e.g. 500 µl to 250 mg). This resulted in 20-30 subsamples per *V. pourtalesii*

sample. Each Horniman invasive sponge sample weighed ~500 mg individually thus no

subsamples were required. The (sub)samples were homogenised on a Qiagen

TissueLyser II for 2 minutes at maximum speed. Using a 1000 µl pipette tip, the

homogenate from each subsample was pooled in a 15 ml falcon tube corresponding to each original sample. Twelve 200 µl aliquots of each *V. pourtalesii* sample and twelve 50 µl aliquots of the Horniman invasive sponge (Fig. 1) were transferred to individual 2 ml DNA LoBind tubes and frozen at -20°C alongside the homogenised samples.

The aliquots were used as the starting material for extraction with three commercial and two non-commercial DNA extraction protocols: 1) Qiagen DNeasy PowerSoil Kit (QPS), 2) Qiagen DNeasy Blood and Tissue Kit (QBT), 3) Qiagen DNeasy Blood and Tissue Kit and post-extraction inhibitor removal with the Zymo Research OneStep™ PCR Inhibitor Removal Kit (QBT-ZYMO), 4) Mu-DNA soil protocol (MUS), and 5) Mu-DNA tissue protocol (MUT) with an added inhibitor removal step (Sellers et al*.,* 2018) (Fig. 1). The QPS, QBT, MUS and MUT protocols were used on three aliquots of each sample (*n* = 72) following the manufacturer's instructions or steps outlined in Sellers et al. (2018) with one modification. Elution buffer (100 µl) was added to each spin column, incubated for 5 minutes at room temperature before centrifugation for 1 minute at 10,000 x g, then the eluate was collected and added to the same spin column again for incubation for 5 minutes at room temperature before centrifugation for 1 minute at 10,000 x g. An extraction blank (extraction buffers only) was processed alongside samples for each extraction protocol (*n* = 4).

The extracts resulting from QBT extraction were halved (Fig. 1), and one half was processed with the Zymo Research OneStep™ PCR Inhibitor Removal Kit (Cambridge Bioscience, UK) for the QBT-ZYMO protocol (*n* = 19). Notably, three preps from the OneStep™ PCR Inhibitor Removal Kit failed, resulting in only 16 extracts for the QBT-ZYMO protocol (Fig. 1). A 25 µl and 5 µl aliquot of each extract (*n* = 92) were transferred to PCR strip tubes (Starlab, UK). All original DNA extracts and 25 µl aliquots were frozen at -20°C until PCR preparation. A Nanodrop was used to measure DNA purity and a Qubit™ 4 Fluorometer (Thermo Scientific, UK) with Qubit™ dsDNA HS Assay Kit (Fisher Scientific, UK) was used to measure total DNA yield from the 5 µl aliquots, which were subsequently discarded.

2.4.2 Phase 2

Three *V. pourtalesii* samples and three *P. ventilabrum* samples were used in Phase 2

(Fig. 1). Based on the findings of Phase 1 (see Results), all samples were blotted

against filter paper in a petri dish to remove residual ethanol preservative and extracted

with the MUT protocol. Samples from each sponge species were halved (n = 12), with

each half being used to provide either dry tissue or homogenate as starting material

(Fig. 1). Homogenisation of each sample was performed as described in Phase 1.

Three dry tissue weights and three homogenate volumes were compared in triplicate for

each sample (n = 108), and the volume of Lysis Master Mix (1 ml contains 730 µl of

Lysis Solution, 250 µl of Tissue Lysis Additive and 20 µl of Proteinase K) for the MUT

protocol was adjusted proportionately: 1) 50 mg and 100 µl (≈50 mg) to 300 µl of Lysis

Master Mix, 2) 100 mg and 200 µl (≈100 mg) to 600 µl of Lysis Master Mix, and 3) 500

mg and 1000 µl (≈500 mg) to 1 ml of Lysis Master Mix. Extraction followed the steps

outlined in Sellers et al. (2018) with the same modification for elution as Phase 1.

Notably, there was not enough tissue for some samples to obtain triplicate 500 mg

weights or 1000 µl volumes, even when additional samples collected from the same

location were used as replicates (PHAK4, PHAK5), resulting in 105 DNA extracts. An

extraction blank (extraction buffers only) was included for each round of extractions (n =

4) (Fig. 1). A 25 µl and 5 µl aliquot of each extract (n = 109) were transferred to PCR

strip tubes (Starlab, UK). All original DNA extracts and 25 µl aliquots were frozen at -

20°C until PCR preparation. A Nanodrop was used to measure DNA purity and a

Qubit™ 4 Fluorometer was used to measure total DNA yield.

2.5 PCR amplification and library preparation

DNA extracts were PCR amplified using the Tele02 primers (F: 5'-

AAACTCGTGCCAGCCACC-3', R: 5'-GGGTATCTAATCCCAGTTTG-3') which target a

~167 bp fragment of the mitochondrial 12S ribosomal RNA (rRNA) gene and achieved

>98% teleost species detection when tested *in silico* (Taberlet *et al.,* 2018). Although

the primers were designed to target teleost fish, mammal and bird DNA may also

amplify (Mariani *et al.,* 2021). To ease sample demultiplexing and mitigate cross-

contamination and/or tag switching during sequencing, each sample was PCR amplified using the locus primers attached to a unique 8 bp tag shared by the forward and reverse primer. Each tag (*n* = 96) differed by at least three base pairs from other tags and included 2-4 degenerate bases (Ns) at the beginning of the tag sequence to improve clustering during initial sequencing. For Phase 2, samples were processed in two batches to facilitate use of the same tags in different pools for adapter ligation.

Each sample was PCR amplified in triplicate using 20 µl reactions consisting of 10 µl of MyFi™ Mix (Meridian Bioscience, UK), 1 µl of each forward and reverse primer (10 µM, Eurofins), 0.16 µl of Bovine Serum Albumin (20 mg/ml, Thermo Scientific, UK), 5.84 µl of UltraPure™ Distilled Water (Invitrogen, UK), and 2 µl of DNA template. Two PCR positive controls (iridescent catfish, *Pangasianodon hypophthalmus*, tissue DNA at 0.05 ng/µl) and two PCR negative controls (molecular grade water) were included in each PCR run (up to 96 reactions). PCR was performed on a T100 Thermal Cycler (Bio-Rad Laboratories Ltd, UK) with the following profile: 95°C for 10 minutes, followed by 40 cycles of 95°C for 30 s, 60°C for 45 s, and 72°C for 30 s, and final elongation at 72°C for 5 minutes. PCR products were run on a 2% agarose gel, made with 150 ml of 1x TBE buffer (Invitrogen, UK), 3 g of agarose powder (Fisher Scientific, UK), and 1.5 µl of SYBR™ Safe DNA Gel Stain (Invitrogen, UK), at 100 V for 40 minutes using 1 µl of PCR product to 1 µl of 10x BlueJuiceTM Gel Loading Buffer (Invitrogen, UK). Gels were visualised with a Bio-Rad Gel Doc™ EZ Imaging System.

PCR replicates for each sample were pooled prior to purification with Mag-Bind[®] Total Pure NGS Beads (Omega Bio-tek Inc., GA, USA), following the double size selection protocol established by Bronner et al. (2009). Ratios of 1x and 0.6x magnetic beads to 30 µl of PCR product were used. Eluted DNA (20 µl) was stored at -20°C until quantification using a Qubit™ 4 Fluorometer. Purified samples were normalised and pooled in equimolar concentration. Phase 1 consisted of one pool of 96 samples/controls. Phase 2 required two pools - the first consisted of 65 samples/controls from Phase 2 and the second consisted of 40 samples/controls from Phase 2 and 25 samples/controls from other projects. Sample pools were purified using the aforementioned bead ratios and elution in 25 µl, then concentrated using a 1x bead ratio and elution in 45 µl.

End repair, adapter ligation and PCR were performed using the NEXTFLEX®

Rapid DNA-Seq Kit 2.0 for Illumina® Platforms (PerkinElmer, UK) according to the

manufacturer's protocol. An Agilent 2200 TapeStation and High Sensitivity D1000

ScreenTape (Agilent Technologies, UK) indicated secondary product remained, thus gel

extraction was performed on each pool using the GeneJET Gel Extraction Kit (Thermo

Scientific, UK) with elution in 20 µl. Each pool was quantified using quantitative PCR

(qPCR) on a Rotor-Gene Q (Qiagen, Germany) with the NEBNext® Library Quant Kit for

Illumina® (New England Biolabs, UK) and diluted to 1 nM. For Phase 2, 6 µl of each

pool were combined into one library. The final libraries and PhiX Control were quantified

using qPCR before the Phase 1 library was sequenced at 50 pM with 10% PhiX Control

and the Phase 2 library was sequenced at 60 pM with 10% PhiX Control on an Illumina®

iSeq™ 100 using iSeq™ 100 i1 Reagent v2 (300-cycle) (Illumina Inc., CA, USA).

2.6 Bioinformatics

2.6.1 Reference databases

Sequence data were processed in two batches to facilitate taxonomic assignment

against different reference databases. Sequences for Horniman invasive sponge

samples and associated controls were compared against a reference database for the

aquarium fish species, the PCR positive control, and common domestic species (human

Homo sapiens, cow *Bos taurus*, pig *Sus scrofa*, chicken *Gallus gallus*, sheep *Ovis aries*,

dog *Canis lupus*, and cat *Felis catus*). Sequences for *V. pourtalesii* and *P. ventilabrum*

samples and associated controls were compared against a marine vertebrate reference

database, the PCR positive control, and common domestic species. Full details of

database creation can be found in the Supporting Information.

2.6.2 Data processing and taxonomic assignment

Sequence data were automatically demultiplexed to separate (forward and reverse) fastq files

per library using the onboard Illumina MiSeq Reporter software. Library sequence reads were

further demultiplexed to sample using a custom Python script. *Tapirs*, a reproducible workflow

for the analysis of DNA metabarcoding data (https://github.com/EvoHull/Tapirs), was used for

processing and taxonomic assignment of demultiplexed sequence reads. *Tapirs* uses the

Snakemake workflow manager (Köster & Rahmann, 2012) and a *conda* virtual environment to
ensure software compatibility.

Raw reads were quality trimmed from the tail with a 5 bp sliding window (qualifying
phred score of Q30 and an average window phred score of Q30) using fastp (Chen et al., 2018),
allowing no more than 40% of the final trimmed read bases to be below Q30. Poly X/G tail
trimming (≥ 5 bp) removed any remaining Illumina iSeq™ 100 sequencing artefacts. Primers
were removed by trimming the first 18 bp and 20 bp of the forward and reverse reads
respectively. Reads were then tail cropped to a maximum length of 150 bp and reads shorter
than 90 bp were discarded. Sequence read pairs were merged into single reads using fastp,
provided there was a minimum overlap of 20 bp, no more than 5% mismatches and no more
than 5 mismatched bases between pairs. Only forward reads were kept from read pairs that
failed to be merged. A final length filter removed any reads longer than 200 bp to ensure
sequence lengths approximated the expected fragment size (~167 bp). Redundant sequences
were removed by clustering at 100% read identity and length (--derep_fulllength) in *VSEARCH*
(Rognes et al., 2016). Clusters represented by less than three sequences were omitted from
further processing. Reads were further clustered (--cluster_unoise) to remove redundancies due
to sequencing errors (retaining all cluster sizes). Retained sequences were screened for
chimeric sequences with *VSEARCH* (--uchime3_denovo).

The final clustered, non-redundant query sequences were then compared
against the corresponding reference database using *BLAST* (Zhang et al., 2000).
Taxonomic identity was assigned using a custom majority lowest common ancestor
(MLCA) approach based on the top 2% query *BLAST* hit bit-scores, with at least 90%
query coverage. A minimum *BLAST* hit identity filter of 80% was used for the Horniman
invasive sponge samples and the aquarium fish reference database, whereas a
minimum identity of 98% was used for *V. pourtalesii* and *P. ventilabrum* samples and
the marine vertebrate reference database. Of these filtered hits for both batches, 80% of
unique taxonomic lineages therein had to agree at descending taxonomic rank (domain,
phylum, class, order, family, genus, species) for it to be assigned a taxonomic identity. If
a query had a single *BLAST* hit, it was assigned directly to this taxon only if it met all
previous MLCA criteria. Read counts assigned to each taxonomic identity were
calculated from query cluster sizes. Lowest taxonomic rank was to species and
assignments higher than order were classed as unassigned.

2.7 Data analysis

2.7.1 Dataset refinement

Taxonomically assigned data was combined for downstream processing in R v3.6.3 (R Core Team, 2020). All plots were produced using the packages ggplot2 v3.3.1 (Wickham, 2016) and ggpubr v0.3.0 (Kassambara, 2020). We split the data by Phase 1 (comparison of ethanol treatments and DNA extraction protocols) and Phase 2 (comparison of types and amounts of starting material). Assignments were corrected: family and genera containing a single species were reassigned to that species, species were reassigned to domestic subspecies, and misassignments were corrected. Manual reassignment duplicated some assignments thus the read count data for these assignments were merged. Proportional read counts for each taxon were calculated from the total read counts per sample. Taxon-specific sequence thresholds (i.e. maximum sequence frequency for each taxon detected in process controls) were applied to the sponge samples to mitigate cross-contamination and false positives (Fig. S1). A second sequence threshold (0.01%) was applied to the sponge samples to mitigate low noise detections. Higher taxonomic assignments, iridescent catfish (PCR positive control), human, domestic species, and terrestrial species were then removed. Taxonomic assignments remaining in the refined dataset were predominantly of species resolution and considered true positives (Fig. S2).

2.7.2 Data summaries

We examined overall fish and mammal taxon richness and read counts across samples processed with each ethanol treatment and DNA extraction protocol in Phase 1 (Fig. S3) and samples of each type and amount of starting material in Phase 2 (Fig. S4) for each sponge species. We also examined detection consistency across extraction replicates, in terms of taxon richness and number of replicates each taxon was detected in, for samples processed with each ethanol treatment and DNA extraction protocol in Phase 1 (Fig. S5) and samples of each type and amount of starting material in Phase 2 (Fig. S6) for each sponge species.

2.7.2 Statistical analysis

We compared total DNA yield, post-PCR DNA concentration, taxon richness and

proportional read counts of individual samples according to ethanol treatment and DNA

extraction protocol for Phase 1, and according to type and amount of starting material

for Phase 2. Only fish detections were used to compare extraction protocols and

ethanol treatments in terms of taxon richness, proportional read counts, and community

similarity as the metabarcoding primers used were designed to target fish. The data

were not normally distributed and the assumptions of a two-way Analysis of Variance

(ANOVA) were violated, thus statistical treatment followed Generalised Linear Models

(GLMs) with Quasi-Poisson error family or Gaussian error family (for total DNA yield

and post-PCR concentration), Poisson error family and the log-link function or negative

binomial error family (for taxon richness), or binomial error family (for proportional read

counts). If a significant effect was found using the *drop1* function from the package stats

v3.6.3, multiple pairwise comparisons were performed using the *pairs* function from the

package emmeans v1.7.2 (Lenth, 2022). Where assumptions of GLMs were violated,

the non-parametric Kruskal-Wallis Test from the package stats was used, and multiple

pairwise comparisons were performed using the non-parametric Dunn's Test (p-values

adjusted with the Benjamini-Hochberg method) from the package FSA v0.8.30 (Ogle et

al., 2020). In Phase 1, data were split by ethanol treatment to compare extraction

protocols and/or vice versa where significant effects were found using GLMs. In Phase

2, data were split by type of starting material to compare amounts of starting material

and/or vice versa where significant effects were found using GLMs.

The package betapart v1.5.1 (Baselga & Orme, 2012) was used to estimate total

beta diversity, partitioned by nestedness-resultant (i.e. community dissimilarity due to

taxon subsets) and turnover (i.e. community dissimilarity due to taxon replacement),

across samples in Phases 1 and 2 with the *beta.multi* function. Total beta diversity and

its two components (Jaccard dissimilarity) were estimated for sponge samples in both

phases using the *beta.pair* function. For total beta diversity and both partitions, we

calculated the average distance of group members to the group centroid in multivariate

space using the *betadisper* function in the package vegan v2.5-6 (Oksanen et al.,

2019). This was done for each group of samples within ethanol treatments and DNA

extraction protocols in Phase 1 and within types and amounts of starting material in
Phase 2. The distances of group members to the group centroid were subjected to an
ANOVA to test for homogeneity of multivariate dispersions (MVDISP), that is whether
the dispersions of one or more groups were different, prior to permutational multivariate
analysis of variance (PERMANOVA). Community dissimilarity for total beta diversity and
both partitions was visualised with Non-metric Multidimensional Scaling (NMDS) using
the *metaMDS* function with 1000 permutations, and tested statistically with
PERMANOVA using the function *adonis* in the package vegan. The PERMANOVA
included an interaction term between ethanol treatment and DNA extraction protocol for
Phase 1 and an interaction term between type and amount of starting material for
Phase 2. Pre-defined cut-off values were used for effect size, where PERMANOVA
results were interpreted as moderate and strong effects if $R^2 > 0.09$ and $R^2 > 0.25$
respectively. These values are broadly equivalent to correlation coefficients of $r = 0.3$
and 0.5 which represent moderate and strong effects accordingly (Nakagawa & Cuthill,
2007; Macher et al., 2018).
442
443

3. Results

3.1 Sequencing performance
The Phase 1 library generated 3,757,770 raw sequence reads, of which 3,532,041
remained after quality control and merging. Following dereplication, denoising, and
chimaera detection, 2,995,356 reads (average read count 31,201 per sample including
controls) remained for taxonomic assignment. The Phase 2 library generated 3,005,701
raw sequence reads, of which 2,879,596 remained after quality control and merging.
Following dereplication, denoising, and chimaera detection, 2,170,282 reads (average
read count 16,823 per sample including controls) remained for taxonomic assignment.
For the Horniman invasive sponge samples, 85.05% of remaining reads were
taxonomically assigned while 15.95% could not be assigned to any taxonomic rank. For
P. ventilabrum and *V. pourtalesii* samples, 78.87% of remaining reads were
taxonomically assigned, while 21.13% could not be assigned to any taxonomic rank.

Contamination was restricted to human DNA in one extraction blank and one PCR

negative control in Phase 1, but naturally occurring marine species were also detected

in extraction blanks and PCR positive controls in Phase 2. Before dataset refinement,

29 taxa were identified in Phase 1 and 94 taxa were identified in Phase 2. After dataset

refinement, 18 taxa were identified in Phase 1 and 49 taxa were identified in Phase 2

(Fig. S2).

For simplicity, we only report significant results from the final statistical analyses

applied to the data. Full results from all statistical tests, including non-significant results

and violations of model assumptions, are detailed in the Supporting Information.

Statistical comparisons were not performed for DNA purity due to lack of variation in

A260/A280 and A260/A230 ratios between ethanol treatments, extraction protocols,

types of starting material, and amounts of starting material.

3.2 Phase 1: ethanol treatment and DNA extraction protocol

For the Horniman invasive sponge, total DNA yield significantly differed between

extraction protocols for the spin ethanol treatment (Kruskal-Wallis Test: H_4 = 11.314, P

= 0.023) and between ethanol treatments for the QBT (Kruskal-Wallis Test: H_2 = 7.200,

P = 0.027) and MUT (Kruskal-Wallis Test: H_2 = 7.261, P = 0.027) protocols. For the spin

treatment, none of the Dunn's Test adjusted p-values were significant, but the largest

differences in effect size were observed between the QBT and MUS protocols (Z = -

2.749, P = 0.060), the QBT and MUT protocols (Z = -2.107, P = 0.117), and the QBT

and QPS protocols (Z = 2.565, P = 0.052). For both the QBT and MUT protocols, the

spin treatment produced lower total DNA yields than the wet treatment (QBT: Z = -

2.683, P = 0.022; MUT: Z = -2.695, P = 0.021) (Fig. 2a). With *V. pourtalesii*, total DNA

yield significantly differed between extraction protocols for the wet (Kruskal-Wallis Test:

H_4 = 13.233, P = 0.010) and blot (Kruskal-Wallis Test: H_4 = 11.833, P = 0.019) ethanol

treatments. The QBT protocol produced higher total DNA yields than the QPS protocol

with the wet ethanol treatment (Dunn's Test: Z = 3.286, P = 0.010) and the MUT

protocol with the blot treatment (Dunn's Test: Z = -2.921, P = 0.035) (Fig. 2a).

For the Horniman invasive sponge, ethanol treatment influenced post-PCR DNA

concentration with the MUS protocol (Kruskal-Wallis Test: H_2 = 6.006, P = 0.050), where

the spin treatment produced lower post-PCR DNA concentrations than the wet

treatment (Dunn's Test: Z = -2.395, P = 0.050) (Fig. 2b). For *V. pourtalesii*, post-PCR

DNA concentration was positively influenced by the MUT (GLM: 0.669 ± 0.327, t =

2.044, P = 0.048) and QBT (GLM: 1.005 ± 0.311, t = 3.229, P = 0.003) protocols, but

the MUT protocol (12.13 ± 2.31 ng/µl) produced lower post-PCR DNA concentrations

than the QBT protocol (16.97 2.74 ng/µl; pairwise comparison: Z = -3.229, P = 0.011)

(Fig. 2b).

Only the MUS and QBT extraction protocols produced incidental mammal

detections (Fig. 3). Using a GLM, neither extraction protocol or ethanol treatment were

found to influence fish taxon richness for the Horniman invasive sponge, but model

assumptions were met. For *V. pourtalesii*, no effect of extraction protocol on fish taxon

richness was found using Kruskal-Wallis Test (Fig. 4a).

Proportional read counts significantly differed between ethanol treatments with

the MUS (Kruskal-Wallis Test: H_2 = 6.161, P = 0.046) and QPS (Kruskal-Wallis Test: H_2

= 6.006, P = 0.050) protocols for the Horniman invasive sponge. The wet treatment

produced higher proportional read counts than the spin treatment with the MUS protocol

(Dunn's Test: Z = -2.426, P = 0.046), and than the blot treatment with the QPS protocol

(Dunn's Test: Z = -2.395, P = 0.050) (Fig. 4b). For *V. pourtalesii*, no effect of extraction

protocol on proportional read counts was found, but proportional reads counts

significantly differed between ethanol treatments with the QBT-ZYMO protocol (Kruskal-

Wallis Test: H_2 = 7.624, P = 0.022), where the blot treatment produced higher

proportional read counts than the spin (Dunn's Test: Z = 2.391, P = 0.025) or wet

(Dunn's Test: Z = 2.391, P = 0.050) treatments (Fig. 4b).

MVDISP differed between DNA extraction protocols for total beta diversity for *V.*

pourtalesii, but not either partition for *V. pourtalesii* or the Horniman invasive sponge

(Table 1). MVDISP differed between ethanol treatments for total beta diversity and the

turnover component for the Horniman invasive sponge, and for the turnover and

nestedness-resultant components of beta diversity for *V. pourtalesii*. DNA extraction

protocol alone did not influence beta diversity for either sponge species (Table 2; Figs.

4ci-iii). Ethanol treatment had a strong positive influence on turnover and total beta

diversity of communities detected from the Horniman invasive sponge, but did not

influence beta diversity for *V. pourtalesii* (Table 2; Figs. 4ci-iii). DNA extraction protocol

combined with ethanol treatment also had a moderate positive effect on turnover for the

Horniman invasive sponge (Table 2; Figs. 4ci-iii). Therefore, taxa detected from the

Horniman invasive sponge with one ethanol treatment were replaced by different taxa

detected with other ethanol treatments, although this finding should be interpreted with

caution due to MVDISP heterogeneity.

3.3 Phase 2: treatment and amount of sponge tissue

Using GLMs, amount of starting material influenced total DNA yield for both *P.*

ventilabrum (F_2 = 16.485, P < 0.001) and *V. pourtalesii* (F_2 = 3.795, P = 0.030), but type

of starting material influenced total DNA yield for *P. ventilabrum* only (F_1 = 14.641, P <

0.001). Model assumptions were met for *P. ventilabrum*. Total DNA yield was higher

using homogenised tissue (0.460 ± 0.120, t = 3.840, P < 0.001) and increased as the

amount of starting material increased (amount 2: 0.451 ± 0.136, t = 3.321, P = 0.002;

amount 3: 0.981 ± 0.165, t = 5.931, P < 0.001). Dried tissue (30.30 ± 2.81 ng/µl)

produced lower total DNA yields than homogenate (48.00 ± 3.82 ng/µl; Z = -3.840, P <

0.001). Amount 3 (63.10 ± 7.96 ng/µl) produced higher total DNA yields than amount 2

(37.10 ± 3.18 ng/µl; Z = -3.474, P = 0.002) or amount 1 (23.70 ± 2.53 ng/µl; Z = -5.931,

P < 0.001). Amount 1 also produced lower total DNA yields than amount 2 (Z = -3.321,

P = 0.003) (Fig. 5a). Model assumptions were not met for *V. pourtalesii* thus Kruskal-

Wallis Test was applied, but amount of starting material was not found to influence total

DNA yield.

Using GLMs, amount of starting material did not influence post-PCR DNA

concentration for either *P. ventilabrum* or *V. pourtalesii*, and type of starting material

influenced post-PCR DNA concentration for *P. ventilabrum* only (F_1 = 5.054, P = 0.031).

Model assumptions for *P. ventilabrum* were violated thus Kruskal-Wallis Test was

applied, but no effect of type of starting material was found (Fig. 5b).

V. pourtalesii produced more incidental mammal detections than *P. ventilabrum*

(Fig. 6). Using GLMs, neither type or amount of starting material were found to influence

fish taxon richness for *P. ventilabrum*, and type of starting material influenced fish taxon

richness for *V. pourtalesii* only (χ_1 = 6.451, P = 0.011). Type of starting material had a

positive influence on taxon richness (0.476 ± 0.190, t = 2.514, P = 0.012), where taxon

richness was higher using homogenate (2.780 ± 0.334) than dried tissue (1.720 ±

0.253; Z = -2.514, P = 0.012) (Fig. 6a).

Using GLMs, neither type or amount of starting material were found to influence

fish proportional read counts for *P. ventilabrum*, and only amount of starting material

influenced fish proportional read counts for *V. pourtalesii* (F_2 = 3.266, P = 0.047). Model

assumptions for *V. pourtalesii* were violated thus Kruskal-Wallis Test was applied, but

no effect of amount of starting material was found (Fig. 6b).

MVDISP did not differ between types of starting material for total beta diversity or

its partitions for either sponge. MVDISP did not differ between amounts of starting

material for total beta diversity or its partitions for *V. pourtalesii*, but did differ between

amounts for total beta diversity and turnover for *P. ventilabrum* (Table 3). For *P.*

ventilabrum, amount of starting material influenced turnover and total beta diversity, but

type of starting material did not influence beta diversity (Table 4; Figs. 7ci-iii). For *V.*

pourtalesii, type of starting material influenced turnover and total beta diversity, but

amount of starting material did not influence beta diversity (Table 4; Figs. 7ci-iii).

Therefore, taxa detected with one type or amount of starting material were replaced by

different taxa detected with the other type (*V. pourtalesii*) or amounts (*P. ventilabrum*) of

starting material, although the latter should be interpreted with caution due to MVDISP

heterogeneity.

4. Discussion

We have illustrated how the performance of sponge nsDNA metabarcoding can be

affected by DNA isolation procedures, specifically treatment to remove residual ethanol

prior to DNA extraction, DNA extraction protocol, type of starting material, and amount

of starting material. These effects can also be sponge-specific. Our experiments have

provided an optimised DNA isolation procedure that will inform the development and

application of sponge nsDNA metabarcoding for aquatic biodiversity assessment.

4.1 Phase 1: ethanol treatment and DNA extraction protocol

582
The QBT and MUT extraction protocols outperformed others for different ethanol
treatments of sponge tissue in terms of total DNA yield and post-PCR DNA
concentration. Lower DNA purity was observed with the MUT and QBT-ZYMO protocols
for the wet ethanol treatments of *V. pourtalesii* tissue and Horniman invasive sponge
tissue respectively, but otherwise extraction protocols produced consistent purity ratios.
Extraction protocol was not found to influence fish taxon richness, proportional read
counts or community composition (arguably the most important extraction success
criteria) for either sponge species.

The spin ethanol treatment produced lower total DNA yields and post-PCR DNA
concentrations with certain extraction protocols for the Horniman invasive sponge, but
no differences were observed between ethanol treatments for *V. pourtalesii*. Ethanol
treatment did not influence fish taxon richness for either sponge species, but did
influence fish proportional read counts with certain extraction protocols, where the wet
or blot treatments produced higher fish proportional read counts than the spin treatment.
Ethanol treatment influenced community composition of Horniman invasive sponge
tissue but not *V. pourtalesii* tissue, where different taxa were detected in Horniman
invasive sponge tissue with different ethanol treatments.

DNA extraction protocols have been found to influence the performance of
metabarcoding across sample types and habitats (Deiner et al., 2018; Jeunen et al.,
2019; Johnson et al., 2019; Geraldi et al., 2019; Sanches & Schreier, 2020), thus it was
not surprising that this also turned out to be the case for sponge tissue. All extraction
protocols produced some biodiversity information, but some protocols were more
effective than others, which has implications for detection of rare and/or low density
species in diverse and complex marine systems using sponge nsDNA metabarcoding.
Our findings echo those of Sellers et al. (2018), where the Mu-DNA protocols were
comparable to or outperformed commercial kits, and other studies which found the QBT
protocol produced higher DNA yield than other commercial kits and unbranded
extraction protocols (Hinlo et al., 2017; Deiner et al., 2018; Jeunen et al., 2019). Mu-
DNA is advantageous due to adaptability and cost-efficiency. Although an inhibitor
removal step is not included in the tissue protocol published in Sellers et al. (2018), it

can be added from the soil or water protocols, which is desirable for obtaining high DNA yields whilst removing inhibitors (Jeunen et al., 2019). A QBT extraction costs four times as much as a MUT extraction (Sellers et al., 2018), and requires a commercial post-extraction inhibitor removal kit (e.g. Zymo Research) to remove potential PCR inhibitors. Reduced cost means that the Mu-DNA protocols are more scalable for large projects and buffer volumes can be adjusted proportionately to the amount of starting material without quickly depleting budgets.

Residual ethanol in tissue can retain unique DNA that may leech from tissue into ethanol (Zizka et al., 2018; Marquina et al., 2019; Jeunen et al., 2019; Derycke et al., 2021; Persaud et al., 2021; Wang et al., 2021), but also react with buffers used for DNA extraction and/or inhibit PCR amplification downstream (Schrader et al., 2012). Our DNA concentration results seem to indicate that residual ethanol contained DNA unique and/or additional to that present in sponge tissue (at least for the Horniman invasive sponge) as higher total DNA yields and post-PCR DNA concentrations were observed with the wet ethanol treatment. However, this did not translate to higher fish taxon richness, and conflicting effects were found for fish proportional read counts from each sponge species, where the wet ethanol treatment generated higher proportional read counts for the Horniman invasive sponge and the blot ethanol treatment produced higher proportional read counts for *V. pourtalesii*. Beta diversity analyses also indicated that different taxa were detected using different ethanol treatments for the Horniman invasive sponge. This reaffirms findings of studies on bulk invertebrate samples (Zizka et al., 2018; Derycke et al., 2021; Persaud et al., 2021; Wang et al., 2021) and marine sponges (Jeunen et al., 2019) where eDNA from ethanol preservative detected different taxa to DNA from tissue, although eDNA from ethanol preservative generally detected more taxa than bulk invertebrate tissue but less taxa than marine sponge tissue.

Sponge species, extraction protocol and ethanol treatments appear to have specific interactions and effects on total DNA yield, DNA purity, target DNA concentration, taxon recovery, and sequence generation. This has been observed for aquatic eDNA in terms of filter material and extraction protocol (Deiner et al., 2017), although other studies have suggested that these aspects of the eDNA workflow do not interact with each other (Sanches & Schreier, 2020). A more elaborate experiment

incorporating numerous individuals of different sponge species would be required to test
this hypothesis. Based on our results, we recommend that these future studies focus on
comparing tissue-based extraction protocols and DNA extraction from ethanol alone
versus sponge tissue alone versus ethanol and sponge tissue combined. For the
purposes of experimental Phase 2 in the present study, blotting sponge tissue dry after
ethanol preservation followed by DNA extraction with the MUT protocol appeared to be
the best combination for extracting DNA from tissue of naturally occurring marine
sponges like *V. pourtalesii*.

4.2 Phase 2: treatment and amount of sponge tissue
Homogenised tissue produced higher total DNA yields for *P. ventilabrum* tissue but not
V. pourtalesii tissue. DNA from homogenised tissue from both sponge species was of
lower purity. Type of starting material did not influence post-PCR DNA concentration for
either sponge species. Homogenised tissue resulted in higher fish taxon richness for *V.*
pourtalesii, although not *P. ventilabrum*, and type of starting material did not influence
fish proportional read counts for either sponge species. Different communities were
observed using different types of starting material for *V. pourtalesii*, but community
composition from homogenate and dried tissue was highly consistent for *P. ventilabrum*.

Larger amounts of starting material produced higher total DNA yields for *P.*
ventilabrum tissue, but not *V. pourtalesii* tissue. DNA purity was consistent across
different amounts of starting material for both sponge species. Amount of starting
material did not influence post-PCR DNA concentration, fish taxon richness or fish
proportional read counts for either sponge species. However, different communities
were observed using different amounts of starting material for *P. ventilabrum*, but not for
V. pourtalesii.

Homogenisation is often used with complex sample types to account for uneven
distribution of organisms/DNA throughout the sample by facilitating the release of DNA
via bead beating, blending or vortexing, but introduces contamination risk via more
handling of samples and reusable equipment (Gosselin et al., 2017; Marquina et al.,
2019; Pereira-da-Conceicoa et al., 2020; Hestetun et al., 2021; Zizka et al., 2022). Until
now, the distribution of eDNA in marine sponge tissue and whether homogenisation was

required was unknown. Our results suggest that eDNA has a relatively even distribution
in sponge tissue. Type of starting material did not influence fish taxon richness for *P.*
ventilabrum or fish proportional read counts for either sponge species. Although higher
taxon richness was observed with homogenised *V. pourtalesii* tissue, the gains were
marginal compared to dried *V. pourtalesii* tissue. A homogenous distribution of eDNA in
sponge tissue offers additional advantages to using these natural samplers for marine
biodiversity assessment over aquatic eDNA which can exhibit substantial spatial
heterogeneity (Bruce et al., 2021). Dried tissue appears to be sufficient for taxon
recovery and would avoid the contamination risk associated with homogenisation.
However, sponge nsDNA accumulation and degradation rates still require investigation
to quantify temporal variation in eDNA signals from these natural samplers (Jeunen et
al., 2021) compared to aquatic eDNA accumulation and degradation rates which differ
by target taxa and habitat (Harrison et al., 2019). This has been done in an artificial
capacity (Cai et al., 2022), but must also be investigated in natural settings.

689    Total DNA yield increased as volume of homogenate and weight of dry tissue
increased. This relationship was expected (bar the risk of introducing inhibitors by using
too much tissue for DNA extraction) as using more starting material was found to
improve detection success for metabarcoding of bulk invertebrate (Zizka et al., 2022)
and soil (Kirse et al., 2021) samples. However, more starting material did not result in
higher fish taxon recovery or fish proportional read counts, which were consistent
across volumes of homogenate and weights of dried tissue. Different amounts of
starting material also produced similar communities for V. pourtalesii, albeit different
communities were observed with different amounts of P. ventilabrum tissue. Therefore,
there seems to be little benefit to using more starting material in sponge nsDNA
metabarcoding as more complex extraction procedures are required to handle larger
quantities of tissue. Instead, it may be more advantageous to invest effort in performing
more extraction replicates with smaller amounts of starting material (Lanzén et al.,
2017; Hestetun et al., 2021).

4.3 Future directions

Our findings have advanced the field of sponge nsDNA metabarcoding and emphasise the importance of sequencing samples used in comparisons of DNA extraction procedures rather than relying only on DNA concentration or purity values and gel images. Nevertheless, a number of unanswered questions remain. The sample size used in our study for each sponge, ethanol treatment, and DNA extraction protocol as well as type and amount of starting material was small due to the various permutations on DNA extraction methods being investigated and the available budget. Thousands of sponges occupy marine systems at every depth and latitude across the globe (Morganti et al., 2021). They can vary in shape, size, and hardness giving rise to variable characteristics, including height, diameter, depth, growth form (free-standing, encrusting, tubular, massive), substrate (hard rock, soft sediment, animal, floating debris), spicules (spongin fibre, calcium carbonate, silica), colour, symmetry (radial symmetry or asymmetrical), body plan (Asconoid, Syconoid, Leuconoid), and number and cross-sectional area of oscula. They also differ in terms of reproduction (sexual or asexual), feeding (carnivorous or herbivorous), defence (toxicity or boring activity), disease (fungi, viruses, cyanobacteria), and symbionts (fish, shrimps, plants, bacteria, algae) (Kahn et al., 2015; Kandler et al., 2019; Morganti et al., 2021). Only by pooling empirical nsDNA-inferred taxon inventories from a variety of wild sponges in natural settings will it be possible to gauge a more comprehensive understanding of the robustness and reliability of this approach across habitats and geographical regions.

Different sponges may not have the same pumping and filtration efficiency or even have viable tissue for DNA extraction due to microbial loads and PCR inhibitors (Cai et al., 2022). Therefore, future research must identify the best natural samplers in both benthic and pelagic habitats from those sponges that are accessible and can withstand repeat sampling as well as understand how different sponge characteristics influence eDNA recovery (Mariani et al., 2019; Jeunen et al., 2021). The spatial and temporal resolution of sponge nsDNA under different environmental conditions and across seasons must be investigated as we do not know the spatial extent or recency of eDNA trapped in sponge tissue (Turon et al., 2020) and whether eDNA in sponge tissue is affected by the same factors that hasten degradation of aquatic eDNA (e.g.

temperature, pH, salinity, microbial activity), but these factors are known to influence the
pumping and filtration action of sponges (Gökalp et al., 2020; Morganti et al., 2021). It is
also undetermined whether sponges exhibit selectivity for eDNA particles of different
sizes akin to filter membranes used for aquatic eDNA (Turner et al., 2014) and the
sampling effort (whether replicates from the same individual or replicates from different
individuals) required for sponges to achieve set species detection probabilities (Turon et
al., 2020).

In addition to these biological considerations, there are further technical
considerations for sponge nsDNA metabarcoding. Many species were only detected in
one extraction replicate performed on tissue from wild *V. pourtalesii* specimens,
emphasising the potential stochasticity of sponge tissue and need for extraction
replicates, similar to sediment (Lanzén et al., 2017; Hestetun et al., 2021) and bulk
invertebrate samples (Elbrecht & Steinke, 2018). Marker and primer choice is crucial to
metabarcoding success. Existing sponge nsDNA studies have focused on 12S and 16S
rRNA primers for the detection of fish (Mariani et al., 2019; Turon et al., 2020; Jeunen et
al., 2021), but as future studies diverge and other markers are explored, primers that
minimise non-target amplification (Collins et al., 2019) and blocking primers
(Boessenkool et al., 2012) to prevent amplification of DNA from the sponge itself could
become essential (Jeunen et al., 2021).

Finally, sequencing depth may play an important role in detection of low
concentration eDNA signals in sponge tissue, whether these represent species that are
rare, small or release less DNA into the environment. Very few studies have compared
sequencing platforms for metabarcoding, but comparisons have been made for the
popular Illumina MiSeq with the larger Illumina NovaSeq (Singer et al., 2019) and with
the smaller Illumina iSeq 100 (Nakao et al., 2021). The Illumina NovaSeq detected more
species from seawater than the Illumina MiSeq, and higher sequencing depth was also
found to improve species detection from sediment (Lanzén et al., 2017) and faecal
(Alberdi et al., 2018) samples with metabarcoding. However, species detection rates
and community composition were highly similar between the Illumina MiSeq and
Illumina iSeq 100 (Nakao et al., 2021). Given these disparities, future studies should
carefully consider sequencing depth for nsDNA metabarcoding when testing the

performance of new natural samplers and addressing the outstanding questions
identified here.

These biological and technical considerations are well-exemplified by the
Horniman invasive sponge in our study which detected 4 of 25 (after dataset
refinement) or 11 of 25 (before dataset refinement) species present in the aquarium
tank. The sponge individuals were relatively young, having regrown over a few weeks
after attempts to remove the sponge from the aquarium tank. This should have been
enough time for eDNA to accumulate in the tissues of these sponges (Cai et al., 2022),
but they may not have reached a large enough size to adequately pump and filter eDNA
from the volume of water held in the tank. The BLAST identity for taxonomic assignment
also had to be reduced to 80% in order to recover more species known to be present in
the aquarium tank. This suggests that the primers used may have insufficient taxonomic
resolution to discriminate between tropical fish species, namely members of the families
Acanthuridae, Chaetodontidae, Labridae, Pomacanthidae, Serranidae and Siganidae.
780
781

5. Conclusions

Our study represents another step forward in the development of sponge nsDNA
metabarcoding as a means to assess aquatic biodiversity, and highlights how
methodological choices (specifically DNA isolation procedures) can influence the taxa
and subsequent communities recovered using this approach. Based on our results, we
recommend blotting sponge tissue dry to remove ethanol preservative prior to extraction
with a tissue-based protocol, whether commercial or non-commercial. However, we
note that more extensive comparisons of eDNA in ethanol preservative only, sponge
tissue only, and ethanol and sponge tissue combined extracted with tissue-based
protocols would be worthwhile. Homogenisation does not seem to be necessary for
sponges, eliminating a time-consuming step that presents a contamination risk. Amount
of starting material did not substantially influence taxon recovery or community
composition, but further comparisons of multiple extraction replicates with small
amounts of sponge tissue versus a single extraction with a large amount of tissue are

needed to determine whether time is better invested in simpler but more numerous extractions or more complex but fewer extractions. With these additional comparative studies, we will have a comprehensive understanding of the optimal DNA isolation procedure for sponge nsDNA metabarcoding which future investigations, whether of a biological or technical nature, can draw upon to advance this growing field.

Acknowledgements

This study was funded by the UK Natural Environment Research Council grant NE/T007028/1. A.R. was also supported by the Spanish Ministry of Science and Innovation grant (RYC2018-024247-I) and CSIC intramural grant (202030E006). M.B.A. was supported by the National Agency for Research and Development (ANID), fellowship program Postdoctorado en el extranjero/2019 - 74200143.

Author Contributions

S.M. and L.R.H. designed the study; J.C., A.R., and B.M. facilitated sample collection; L.R.H, E.F.N. and A.C. performed the laboratory work with guidance from G.S., M.B.A. and A.R.; G.S. performed the bioinformatic analyses; S.M. and A.R. obtained the funding; L.R.H. analysed the data and wrote the manuscript, which all authors revised.

Data Accessibility

Raw sequence reads have been archived on the NCBI Sequence Read Archive (BioProject: PRJNA854174; BioSamples: SAMN29421799 - SAMN29421894 [Phase 1] and SAMN29444657 - SAMN29444784 [Phase 2]; SRA accessions: SRR19906974 - SRR19907069 [Phase 1] and SRR19912485 - SRR19912613 [Phase 2]). Scripts and

corresponding data have been deposited in a dedicated Zenodo repository (https://doi.org/10.5281/zenodo.6815267).

References

Alberdi, A., Aizpurua, O., Gilbert, M.T.P., & Bohmann, K (2018) Scrutinizing key steps for reliable metabarcoding of environmental samples. *Methods in Ecology and Evolution*, **9**, 134–147. https://doi.org/10.1111/2041-210X.12849

Baselga, A., & Orme, C.D.L. (2012) betapart: an R package for the study of beta diversity: Betapart package. *Methods in Ecology and Evolution*, **3**, 808–812. https://doi.org/10.1111/j.2041-210X.2012.00224.x

Bessey, C., Gao, Y., Truong, Y.B., Miller, H., Jarman, S.N., & Berry, O. (2021) Comparison of materials for rapid passive collection of environmental DNA. *bioRxiv*, 2021.11.08.467842. https://doi.org/10.1111/1755-0998.13640

Bessey, C., Jarman, S.N., Simpson, T., Miller, H., Stewart, T., Keesing, J.K., & Berry, O. (2021) Passive eDNA collection enhances aquatic biodiversity analysis. *Communications Biology*, **4**, 236. https://doi.org/10.1038/s42003-021-01760-8

Boessenkool, S., Epp, L.S., Haile, J., Bellemain, E., Edwards, M., Coissac, E., Willerslev, E., & Brochmann, C. (2012) Blocking human contaminant DNA during PCR allows amplification of rare mammal species from sedimentary ancient DNA. *Molecular Ecology*, **21**, 1806–1815. https://doi.org/10.1111/j.1365-294X.2011.05306.x

Bronner, I.F., Quail, M.A., Turner, D.J., & Swerdlow, H. (2009) Improved Protocols for Illumina Sequencing. *Current Protocols in Human Genetics*, **18**, 18.2.1–18.2.42. https://doi.org/10.1002/0471142905.hg1802s62

Bruce, K., Blackman, R., Bourlat, S.J., Hellström, A.M., Bakker, J., Bista, I., Bohmann, K., Bouchez, A., Brys, R., Clark, K., Elbrecht, V., Fazi, S., Fonseca, V., Hänfling, B., Leese, F., Mächler, E., Mahon, A.R., Meissner, K., Panksep, K., Pawlowski, J., Schmidt Yáñez, P., Seymour, M., Thalinger, B., Valentini, A., Woodcock, P., Traugott, M., Vasselon, V., & Deiner, K. (2021) A practical guide to DNA-based methods for biodiversity assessment. *Advanced Books*. https://doi.org/10.3897/ab.e68634

Cai, W., Harper, L.R., Neave, E.F., Shum, P., Craggs, J., Arias, M.B., Riesgo, A., & Mariani, S. (2022) Environmental DNA persistence and fish detection in captive sponges. *Molecular Ecology Resources*. https://doi.org/10.1111/1755-0998.13677

Chen, S., Zhou, Y., Chen, Y., & Gu, J. (2018) fastp: an ultra-fast all-in-one FASTQ preprocessor. *Bioinformatics*, **34**, i884–i890. https://doi.org/10.1093/bioinformatics/bty560

Clare, E.L., Economou, C.K., Faulkes, C.G., Gilbert, J.D., Bennett, F., Drinkwater, R., & Littlefair, J.E. (2021) eDNAir: proof of concept that animal DNA can be collected from air sampling. *PeerJ*, **9**, e11030. https://doi.org/10.7717/peerj.11030

Collins, R.A., Bakker, J., Wangensteen, O.S., Soto, A.Z., Corrigan, L., Sims, D.W., Genner, M.J., & Mariani, S. (2019) Non-specific amplification compromises

environmental DNA metabarcoding with COI. *Methods in Ecology and Evolution*, **10**, 1985–2001. https://doi.org/10.1111/2041-210X.13276

Deiner, K., Bik, H.M., Mächler, E., Seymour, M., Lacoursière-Roussel, A., Altermatt, F., Creer, S., Bista, I., Lodge, D.M., de Vere, N., Pfrender, M.E., & Bernatchez, L. (2017) Environmental DNA metabarcoding: Transforming how we survey animal and plant communities. *Molecular Ecology*, **26**, 5872–5895. https://doi.org/10.1111/mec.14350

Deiner, K., Lopez, J., Bourne, S., Holman, L., Seymour, M., Grey, E.K., Lacoursière, A., Li, Y., Renshaw, M.A., Pfrender, M.E., Rius, M., Bernatchez, L., & Lodge, D.M. (2018) Optimising the detection of marine taxonomic richness using environmental DNA metabarcoding: the effects of filter material, pore size and extraction method. *Metabarcoding and Metagenomics*, **2**, e28963. https://doi.org/10.3897/mbmg.2.28963

Derycke, S., Maes, S., Van den Bulcke, L., Vanhollebeke, J., Wittoeck, J., Hillewaert, H., Ampe, B., Haegeman, A., Hostens, K., & De Backer, A. (2021) Detection of Macrobenthos Species With Metabarcoding Is Consistent in Bulk DNA but Dependent on Body Size and Sclerotization in eDNA From the Ethanol Preservative. *Frontiers in Marine Science*, **8**, 637858. https://doi.org/10.3389/fmars.2021.637858

Drinkwater, R., Williamson, J., Clare, E.L., Chung, A.Y.C., Rossiter, S.J., & Slade, E. (2021) Dung beetles as samplers of mammals in Malaysian Borneo—a test of high throughput metabarcoding of iDNA. *PeerJ*, **9**, e11897. https://doi.org/10.7717/peerj.11897

Elbrecht V., & Steinke, D. (2018) Scaling up DNA metabarcoding for freshwater macrozoobenthos monitoring. *Freshwater Biology*, **64**, 380–387. https://doi.org/10.1111/fwb.13220

Franklin, T.W., McKelvey, K.S., Golding, J.D., Mason, D.H., Dysthe, J.C., Pilgrim, K.L., Squires, J.R., Aubry, K.B., Long, R.A., Greaves, S.E., Raley, C.M., Jackson, S., MacKay, P., Lisbon, J., Sauder, J.D., Pruss, M.T., Heffington, D., & Schwartz, M.K. (2019) Using environmental DNA methods to improve winter surveys for rare carnivores: DNA from snow and improved noninvasive techniques. *Biological Conservation*, **229**, 50–58. https://doi.org/10.1016/j.biocon.2018.11.006

Geraldi, N.R., Díaz-Rúa, R., Shea, L.A., & Duarte, C.M. (2019) Performance of extraction methods for extracellular DNA from sediments across marine habitats. *Environmental DNA*, **4**, e6372. https://doi.org/10.1002/edn3.48

Gökalp, M., Kooistra, T., Rocha, M.S., Silva, T.H., Osinga, R., Murk, A.J., & Wijgerde, T. (2020) The Effect of Depth on the Morphology, Bacterial Clearance, and Respiration of the Mediterranean Sponge *Chondrosia reniformis* (Nardo, 1847). *Marine Drugs*, **18**. https://doi.org/10.3390/md18070358

Gosselin, E.N., Lonsinger, R.C., & Waits, L.P. (2017) Comparing morphological and molecular diet analyses and fecal DNA sampling protocols for a terrestrial carnivore: Noninvasive Diet Analysis Methodologies. *Wildlife Society Bulletin*, **41**, 362–369. https://doi.org/10.1002/wsb.749

Harper, L.R., Niemiller, M.L., Benito, J., Paddock, L.E., Knittle, E., Molano-Flores, B., & Davis, M.A. (2021) BeeDNA: microfluidic environmental DNA metabarcoding as a tool for connecting plant and pollinator communities. *bioRxiv*, 2021.11.11.468290. https://doi.org/10.1101/2021.11.11.468290

Harrison, J.B., Sunday, J.M., & Rogers, S.M. (2019) Predicting the fate of eDNA in the
environment and implications for studying biodiversity. *Proceedings of The Royal*
Society B, **286**, 20191409. https://doi.org/10.1098/rspb.2019.1409
Hermans, S.M., Buckley, H.L., & Lear, G. (2018) Optimal extraction methods for the
simultaneous analysis of DNA from diverse organisms and sample types. *Molecular*
Ecology Resources, **18**, 557–569. https://doi.org/10.1111/1755-0998.12762
Hestetun, J.T., Lanzén, A., Skaar, K.S., & Dahlgren, T.G. (2021) The impact of DNA
extract homogenization and replication on marine sediment metabarcoding diversity
and heterogeneity. *Environmental DNA*, **3**, 997–1006.
https://doi.org/10.1002/edn3.223
Hinlo, R., Gleeson, D., Lintermans, M., & Furlan, E. (2017) Methods to maximise
recovery of environmental DNA from water samples. *PLoS ONE*, **12**, e0179251.
https://doi.org/10.1371/journal.pone.0179251
Jeunen, G.-J., Cane, J.S., Ferreira, S., Strano, F., von Ammon, U., Cross, H., Day, R.,
Hesseltine, S., Ellis, K., Urban, L., Pearson, N., Olmedo-Rojas, P., Kardailsky, A.,
Gemmell, N.J., & Lamare, M. (2021) Assessing the utility of marine filter feeders for
environmental DNA (eDNA) biodiversity monitoring. *bioRxiv*, 2021.12.21.473722.
https://doi.org/10.1101/2021.12.21.473722
Jeunen, G.-J., Knapp, M., Spencer, H.G., Taylor, H.R., Lamare, M.D., Stat, M., Bunce,
M., & Gemmell, N.J. (2019) Species-level biodiversity assessment using marine
environmental DNA metabarcoding requires protocol optimization and
standardization. *Ecology and Evolution*, **9**, 1323–1335.
https://doi.org/10.1002/ece3.4843
Johnson, M.D., Cox, R.D., & Barnes, M.A. (2019) Analyzing airborne environmental
DNA: A comparison of extraction methods, primer type, and trap type on the ability to
detect airborne eDNA from terrestrial plant communities. *Environmental DNA*, **1**,
176–185. https://doi.org/10.1002/edn3.19
Kahn, A.S., Yahel, G., Chu, J.W.F., Tunnicliffe, V., & Leys, S.P. (2015) Benthic grazing
and carbon sequestration by deep-water glass sponge reefs. *Limnology and*
Oceanography, **60**, 78–88. https://doi.org/10.1002/lno.10002
Kandler, N.M., Wooster, M.K., Leray, M., Knowlton, N., de Voogd, N.J., Paulay, G., &
Berumen, M.L. (2019) Hyperdiverse Macrofauna Communities Associated with a
Common Sponge, Stylissa carteri, Shift across Ecological Gradients in the Central
Red Sea. *Diversity*, **11**, 18. https://doi.org/10.3390/d11020018
Kassambara A (2020) ggpubr: 'ggplot2' Based Publication Ready Plots. R package
version 0.4.0. https://CRAN.R-project.org/package=ggpubr
Kaunisto, K.M., Roslin, T., Sääksjärvi, I.E., & Vesterinen, E.J. (2017) Pellets of proof:
First glimpse of the dietary composition of adult odonates as revealed by
metabarcoding of feces. *Ecology and Evolution*, **7**, 8588–8598.
https://doi.org/10.1002/ece3.3404
Kinoshita, G., Yonezawa, S., Murakami, S., & Isagi, Y. (2019) Environmental DNA
Collected from Snow Tracks is Useful for Identification of Mammalian Species.
Zoological Science, **36**, 198–207. https://doi.org/10.2108/zs180172
Kirse, A., Bourlat, S.J., Langen, K., & Fonseca, V.G. (2021) Unearthing the potential of
soil eDNA metabarcoding—towards best practice advice for invertebrate biodiversity

assessment. *Frontiers in Ecology and Evolution*, **9**, 630560. https://doi.org/10.3389/fevo.2021.630560

Kirtane, A., Atkinson, J.D., & Sassoubre, L.M. (2020) Design and Validation of Passive Environmental DNA Samplers (PEDS) using Granular Activated Carbon (GAC) and Montmorillonite Clay (MC). *Environmental Science & Technology*, **54**, 11961–11970. https://doi.org/10.1021/acs.est.0c01863

Köster, J., & Rahmann, S. (2012) Snakemake—a scalable bioinformatics workflow engine. *Bioinformatics*, **28**, 2520–2522. https://doi.org/10.1093/bioinformatics/bts480

Lanzén, A., Lekang, K., Jonassen, I., Thompson, E.M., & Troedsson, C. (2017) DNA extraction replicates improve diversity and compositional dissimilarity in metabarcoding of eukaryotes in marine sediments. *PLoS ONE*, **12**, e0179443. https://doi.org/10.1371/journal.pone.0179443

Lawson Handley, L. (2015) How will the "molecular revolution" contribute to biological recording? *Biological Journal of the Linnean Society*, **115**, 750–766. https://doi.org/10.1111/bij.12516

Lenth, R.V. (2022). emmeans: Estimated Marginal Means, aka Least-Squares Means. R package version 1.7.2. https://CRAN.R-project.org/package=emmeans

Lynggaard, C., Bertelsen, M.F., Jensen, C.V., Johnson, M.S., Frøslev, T.G., Olsen, M.T., & Bohmann, K. (2022) Airborne environmental DNA for terrestrial vertebrate community monitoring. *Current Biology*, **32**, 701–707.e5. https://doi.org/10.1016/j.cub.2021.12.014

Macher, J.-N., Vivancos, A., Piggott, J.J., Centeno, F.C., Matthaei, C.D., & Leese, F. (2018) Comparison of environmental DNA and bulk-sample metabarcoding using highly degenerate cytochrome c oxidase I primers. *Molecular Ecology Resources*, **18**, 1456–1468. https://doi.org/10.1111/1755-0998.12940

Majaneva, M., Diserud, O.H., Eagle, S.H.C., Hajibabaei, M., & Ekrem, T. (2018) Choice of DNA extraction method affects DNA metabarcoding of unsorted invertebrate bulk samples. *Metabarcoding and Metagenomics*, **2**, e26664. https://doi.org/10.3897/mbmg.2.26664

Mariani, S., Baillie, C., Colosimo, G., & Riesgo, A. (2019) Sponges as natural environmental DNA samplers. *Current Biology*, **29**, PR401–PR402. https://doi.org/10.1016/j.cub.2019.04.031

Mariani, S., Harper, L.R., Collins, R.A., Baillie, C., Wangensteen, O.S., McDevitt, A.D., Heddell-Cowie, M., & Genner, M.J. (2021) Estuarine molecular bycatch as a landscape-wide biomonitoring tool. *Biological Conservation*, **261**, 109287. https://doi.org/10.1016/j.biocon.2021.109287

Marquina, D., Esparza-Salas, R., Roslin, T., & Ronquist, F. (2019) Establishing arthropod community composition using metabarcoding: Surprising inconsistencies between soil samples and preservative ethanol and homogenate from Malaise trap catches. *Molecular Ecology Resources*, **19**, 1516–1530. https://doi.org/10.1111/1755-0998.13071

Massey, A.L., Bronzoni, R.V. de M., da Silva, D.J.F., Allen, J., de Lazari, P.R., dos Santos Filho, M., Canale, G.R., Bernardo, C.S.S., Peres, C.A., & Levi, T. (2021) Invertebrates for vertebrate biodiversity monitoring: comparisons using three insect taxa as iDNA samplers. *Molecular Ecology Resources*, **22**, 962–977. https://doi.org/10.1111/1755-0998.13525

Miya, M. (2021) Environmental DNA Metabarcoding: A Novel Method for Biodiversity Monitoring of Marine Fish Communities. *Annual Review of Marine Science*, **14**, 161–185. https://doi.org/10.1146/annurev-marine-041421-082251

Morganti, T.M., Ribes, M., Moskovich, R., Weisz, J.B., Yahel, G., & Coma, R. (2021) In situ Pumping Rate of 20 Marine Demosponges Is a Function of Osculum Area. *Frontiers in Marine Science*, **8**, 583188. https://doi.org/10.3389/fmars.2021.583188

Nakagawa, S., & Cuthill, I.C. (2007) Effect size, confidence interval and statistical significance: a practical guide for biologists. *Biological Reviews*, **82**, 591–605. https://doi.org/10.1111/j.1469-185X.2007.00027.x

Nakao, R., Inui, R., Akamatsu, Y., Goto, M., Doi, H., & Matsuoka, S. (2021) Illumina iSeq 100 and MiSeq exhibit similar performance in freshwater fish environmental DNA metabarcoding. *Scientific Reports*, **11**, 15763. https://doi.org/10.1038/s41598-021-95360-5

Nørgaard, L., Olesen, C.R., Trøjelsgaard, K., Pertoldi, C., Nielsen, J.L., Taberlet, P., Ruiz-González, A., De Barba, M., & Iacolina, L. (2021) eDNA metabarcoding for biodiversity assessment, generalist predators as sampling assistants. *Scientific Reports*, **11**, 6820. https://doi.org/10.1038/s41598-021-85488-9

Ogle, D.H., Doll, J.C., Wheeler, P., & Dinno, A. (2021) FSA: Fisheries Stock Analysis. R package version 0.9.1. https://github.com/droglenc/FSA

Oksanen, J., Blanchet, F.G., Friendly, M., Kindt, R., Legendre, P., McGlinn, D., Minchin, P.R., O'Hara, R.B., Simpson, G.L., Solymos, P., Stevens, M.H.H., Szoecs E., & Wagner, H. (2020). vegan: Community Ecology Package. R package version 2.5-7. https://CRAN.R-project.org/package=vegan

Pereira-da-Conceicoa, L., Elbrecht, V., Hall, A., Briscoe, A., Barber-James, H., & Price, B. (2020) Metabarcoding unsorted kick-samples facilitates macroinvertebrate based biomonitoring with increased taxonomic resolution, while outperforming environmental DNA. *Environmental DNA*, **3**, 353–371. https://doi.org/10.1002/edn3.116

Persaud, S.F., Cottenie, K., & Gleason, J.E. (2021) Ethanol eDNA Reveals Unique Community Composition of Aquatic Macroinvertebrates Compared to Bulk Tissue Metabarcoding in a Biomonitoring Sampling Scheme. *Diversity*, **13**, 34. https://doi.org/10.3390/d13010034

Pompanon, F., Deagle, B.E., Symondson, W.O.C., Brown, D.S., Jarman, S.N., & Taberlet, P. (2012) Who is eating what: diet assessment using next generation sequencing. *Molecular Ecology*, **21**, 1931–1950. https://doi.org/10.1111/j.1365-294X.2011.05403.x

Roger, F., Ghanavi, H.R., Danielsson, N., Wahlberg, N., Löndahl, J., Pettersson, L.B., Andersson, G.K.S., Boke Olén, N., & Clough, Y. (2022) Airborne environmental DNA metabarcoding for the monitoring of terrestrial insects—A proof of concept from the field. *Environmental DNA*. https://doi.org/10.1002/edn3.290

Rognes, T., Flouri, T., Nichols, B., Quince, C., & Mahé, F. (2016) VSEARCH: a versatile open source tool for metagenomics. *PeerJ*, **4**, e2584. https://doi.org/10.7717/peerj.2584

Russo, T., Maiello, G., Talarico, L., Baillie, C., Colosimo, G., D'Andrea, L., Di Maio, F., Fiorentino, F., Franceschini, S., Garofalo, G., Scannella, D., Cataudella, S., & Mariani, S. (2020) All is fish that comes to the net: metabarcoding for rapid fisheries

catch assessment. *Ecological Applications,* **31**, e02273. https://doi.org/10.1002/eap.2273

Sanches, T.M., & Schreier, A.D. (2020) Optimizing an eDNA protocol for estuarine environments: Balancing sensitivity, cost and time. *PLoS ONE*, **15**, e0233522. https://doi.org/10.1371/journal.pone.0233522

Schnell, I.B., Thomsen, P.F., Wilkinson, N., Rasmussen, M., Jensen, L.R.D., Willerslev, E., Bertelsen, M.F., & Gilbert, M.T.P. (2012) Screening mammal biodiversity using DNA from leeches. *Current Biology*, **22**, R262–R263. https://doi.org/10.1016/j.cub.2012.02.058

Schrader, C., Schielke, A., Ellerbroek, L. & Johne, R. (2012) PCR inhibitors - occurrence, properties and removal. *Journal of Applied Microbiology*, **113**, 1014–1026. https://doi.org/10.1111/j.1365-2672.2012.05384.x

Sellers, G.S., Di Muri, C., Gómez, A., & Hänfling, B. (2018) Mu-DNA: a modular universal DNA extraction method adaptable for a wide range of sample types. *Metabarcoding and Metagenomics*, **2**, e24556. https://doi.org/10.3897/mbmg.2.24556

Siegenthaler, A., Wangensteen, O.S., Soto, A.Z., Benvenuto, C., Corrigan, L., & Mariani, S. (2019) Metabarcoding of shrimp stomach content: Harnessing a natural sampler for fish biodiversity monitoring. *Molecular Ecology Resources*, **19**, 206–220. https://doi.org/10.1111/1755-0998.12956

Sigsgaard, E.E., Olsen, K., Hansen, M.D.D., Hansen, O.L.P., Høye, T.T., Svenning, J.-C., & Thomsen, P.F. (2020) Environmental DNA metabarcoding of cow dung reveals taxonomic and functional diversity of invertebrate assemblages. *Molecular Ecology*, **30**, 3374-3389. https://doi.org/10.1111/mec.15734

Singer, G.A.C., Fahner, N.A., Barnes, J.G., McCarthy, A., & Hajibabaei, M. (2019) Comprehensive biodiversity analysis via ultra-deep patterned flow cell technology: a case study of eDNA metabarcoding seawater. *Scientific Reports*, **9**, 5991. https://doi.org/10.1038/s41598-019-42455-9

Taberlet, P., Bonin, A., Zinger, L., & Coissac, E. (2018) *Environmental DNA: For Biodiversity Research and Monitoring*. Oxford University Press.

Thalinger, B., Deiner, K., Harper, L.R., Rees, H.C., Blackman, R.C., Sint, D., Traugott, M., Goldberg, C.S., & Bruce, K. (2021) A validation scale to determine the readiness of environmental DNA assays for routine species monitoring. *Environmental DNA*, **3**, 823–836. https://doi.org/10.1002/edn3.189

Thomsen, P.F., & Sigsgaard, E.E. (2019) Environmental DNA metabarcoding of wild flowers reveals diverse communities of terrestrial arthropods. *Ecology and Evolution*, **9**, 1665–1679. https://doi.org/10.1002/ece3.4809

Tsuji, S., Takahara, T., Doi, H., Shibata, N., & Yamanaka, H. (2019) The detection of aquatic macroorganisms using environmental DNA analysis—A review of methods for collection, extraction, and detection. *Environmental DNA*, **1**, 99–108. https://doi.org/10.1002/edn3.21

Turner, C.R., Barnes, M.A., Xu, C.C.Y., Jones, S.E., Jerde, C.L., & Lodge, D.M. (2014) Particle size distribution and optimal capture of aqueous macrobial eDNA. *Methods in Ecology and Evolution*, **5**, 676–684. https://doi.org/10.1111/2041-210X.12206

Turon, M., Angulo-Preckler, C., Antich, A., Præbel, K., & Wangensteen, O.S. (2020) More Than Expected From Old Sponge Samples: A Natural Sampler DNA

Metabarcoding Assessment of Marine Fish Diversity in Nha Trang Bay (Vietnam).
Frontiers in Marine Science, **7**, 605148. https://doi.org/10.3389/fmars.2020.605148

Valentin, R.E., Fonseca, D.M., Gable, S., Kyle, K.E., Hamilton, G.C., Nielsen, A.L., &
Lockwood, J.L. (2020) Moving eDNA surveys onto land: Strategies for active eDNA
aggregation to detect invasive forest insects. *Molecular Ecology Resources*, **20**, 746–
755. https://doi.org/10.1111/1755-0998.13151

Wang, Y., Chen, K., Gao, J., Wang, M., Dong, J., Xie, Y., Giesy, J.P., Jin, X. & Wang,
B. (2021) Environmental DNA of preservative ethanol performed better than water
samples in detecting macroinvertebrate diversity using metabarcoding. *Diversity &*
Distributions, **27**, 1989–2002. https://doi.org/10.1111/ddi.13284

Wickham H (2016) *ggplot2: Elegant Graphics for Data Analysis*. Springer-Verlag New
York.

Zhang, Z., Schwartz, S., Wagner, L., & Miller, W. (2000) A greedy algorithm for aligning
DNA sequences. *Journal of Computational Biology*, **7**, 203–214.
https://doi.org/10.1089/10665270050081478

Zizka, V.M.A., Geiger, M.F., Hörren, T., Kirse, A., Noll, N.W., Schäffler, L., Scherges,
A.M., & Sorg, M. (2022) Recommendations for tissue homogenisation and extraction
in DNA metabarcoding of Malaise trap samples. *bioRxiv*, 2022.01.25.477667.
https://doi.org/10.1101/2022.01.25.477667

Zizka, V.M.A., Leese, F., Peinert, B., & Geiger, M.F. (2018) DNA metabarcoding from
sample fixative as a quick and voucher-preserving biodiversity assessment method.
Genome, **62**, 122–136. https://doi.org/10.1139/gen-2018-0048

1119

1120

1121 **Table 1.** Summary of analyses statistically comparing homogeneity of multivariate
1122 dispersions (MVDISP) between the communities (ANOVA) produced by different
1123 extraction protocols and ethanol treatments. Relative contributions of taxon turnover
1124 and nestedness-resultant to total beta diversity (Jaccard dissimilarity) for each protocol
1125 and treatment are given in brackets.
1126

	Mean distance to centroid ± SE	df	F	*P*
Horniman invasive sponge				
DNA extraction protocol				
Turnover		4	0.059	0.993
MUT (100.00%)	0.375 ± 0.268			
MUS (100.00%)	0.333 ± 0.267			
QBT (96.92%)	0.446 ± 0.276			
QBT-ZYMO (88.89%)	0.413 ± 0.054			
QPS (96.55%)	0.429 ± 0.286			
Nestedness-resultant		4	2.013	0.118
MUT (0.00%)	$4.939 \times 10^{-9} \pm 4.647 \times 10^{-17}$			
MUS (0.00%)	$9.697 \times 10^{-9} \pm 2.341 \times 10^{-16}$			
QBT (3.08%)	0.091 ± 0.015			
QBT-ZYMO (11.11%)	0.110 ± 0.046			
QPS (3.45%)	$8.700 \times 10^{-9} \pm 1.361 \times 10^{-16}$			
Total beta diversity		4	0.132	0.969
MUT (100.00%)	0.375 ± 0.268			
MUS (100.00%)	0.333 ± 0.267			
QBT (100.00%)	0.500 ± 0.199			
QBT-ZYMO (100.00%)	0.413 ± 0.053			
QPS (100.00%)	0.429 ± 0.286			
Ethanol treatment				
Turnover		2	12.343	**<0.001**
Blot (96.73%)	0.500 ± 0.286			
Spin (0.00%	$2.363 \times 10^{-9} \pm 7.259 \times 10^{-17}$			
Wet (0.00%)	$1.129 \times 10^{-9} \pm 1.784 \times 10^{-17}$			
Nestedness-resultant		2	0.294	0.747
Blot (3.27%)	$3.833 \times 10^{-6} \pm 2.959 \times 10^{-11}$			
Spin (100.00%)	0.039 ± 0.019			
Wet (100.00%)	0.036 ± 0.018			
Total beta diversity		2	8.593	**0.001**
Blot (100.00%)	0.500 ± 0.286			
Spin (100.00%)	0.039 ± 0.019			
Wet (100.00%)	0.036 ± 0.018			
Vazella pourtalesii				

DNA extraction protocol

Turnover		4	1.902	0.204
MUT (98.52%)	0.572 ± 0.077			
MUS (100.00%)	$0.500 \pm 2.465 \times 10^{-32}$			
QBT (100.00%)	0.500 ± 0.000			
QBT-ZYMO (100.00%)	$0.575 \pm 2.115 \times 10^{-5}$			
QPS	0.000			
Nestedness-resultant		4	1.620	0.260
MUT (1.48%)	0.078 ± 0.006			
MUS (0.00%)	0.000 ± 0.000			
QBT (0.00%)	$1.273 \times 10^{-8} \pm 1.819 \times 10^{-18}$			
QBT-ZYMO (0.00%)	$6.332 \times 10^{-9} \pm 1.203 \times 10^{-16}$			
QPS	0.000			
Total beta diversity		4	5.107	**0.024**
MUT (100.00%)	0.593 ± 0.030			
MUS (100.00%)	0.500 ± 0.000			
QBT (100.00%)	$0.500 \pm 1.233 \times 10^{-32}$			
QBT-ZYMO (100.00%)	$0.577 \pm 3.513 \times 10^{-9}$			
QPS	0.000			

Ethanol treatment

Turnover		2	10.920	**0.003**
Blot (99.36%)	0.651 ± 0.009			
Spin (98.08%	0.521 ± 0.079			
Wet (0.00%)	$1.343 \times 10^{-8} \pm 0.000$			
Nestedness-resultant		2	11.748	**0.002**
Blot (0.64%)	$2.915 \times 10^{-9} \pm 5.948 \times 10^{-17}$			
Spin (1.92%)	0.137 ± 0.026			
Wet (100.00%)	0.333 ± 0.000			
Total beta diversity		2	4.055	0.051
Blot (100.00%)	0.635 ± 0.009			
Spin (100.00%)	0.559 ± 0.041			
Wet (100.00%)	0.333 ± 0.000			

1127

1128

1129 **Table 2.** Results of PERMANOVA used to compare variation in community composition
1130 produced by different extraction protocols and ethanol treatments, individually and
1131 combined.
1132

	df	F	R^2	*P*	Stress
Horniman invasive sponge					0
Turnover					
DNA extraction protocol	4	2.782	0.060	0.057	
Ethanol treatment	2	67.309	0.720	**0.001**	
DNA extraction protocol : Ethanol treatment	7	2.883	0.108	**0.025**	
Nestedness-resultant					0
DNA extraction protocol	4	0.670	0.122	0.569	
Ethanol treatment	2	-2.124	-0.193	0.947	
DNA extraction protocol : Ethanol treatment	7	0.361	0.115	0.704	
Total beta diversity					0
DNA extraction protocol	4	2.152	0.063	0.087	
Ethanol treatment	2	46.617	0.677	**0.001**	
DNA extraction protocol : Ethanol treatment	7	2.132	0.108	0.054	
Vazella pourtalesii					
Turnover					0
DNA extraction protocol	4	0.677	0.263	0.950	
Ethanol treatment	2	1.117	0.217	0.365	
DNA extraction protocol : Ethanol treatment	2	0.683	0.133	0.830	
Nestedness-resultant					5.113×10^{-5}
DNA extraction protocol	4	1.984×10^{-15}	0.895	0.112	
Ethanol treatment	2	-2.317×10^{-15}	-0.523	0.877	
DNA extraction protocol : Ethanol treatment	2	2.783×10^{-15}	0.628	0.098	
Total beta diversity					0
DNA extraction protocol	4	0.772	0.288	0.886	
Ethanol treatment	2	1.007	0.188	0.427	
DNA extraction protocol : Ethanol treatment	2	0.816	0.152	0.723	

1133
1134

Table 3. Results of ANOVA used to compare homogeneity of multivariate dispersions (MVDISP) between the communities produced by different types and amounts of starting material. Relative contributions of taxon turnover and nestedness-resultant to total beta diversity (Jaccard dissimilarity) for each type and amount of starting material are given in brackets.

	Mean distance to centroid ± SE	df	F	P
Phakellia ventilabrum				
Type of starting material				
Turnover		1	0.172	0.681
Dried tissue (85.02%)	0.115 ± 0.021			
Homogenate (88.16%)	0.131 ± 0.010			
Nestedness-resultant		1	1.729	0.196
Dried tissue (14.98%)	0.074 ± 0.004			
Homogenate (11.84%)	0.051 ± 0.002			
Total beta diversity		1	0.059	0.809
Dried tissue (100.00%)	0.165 ± 0.026			
Homogenate (100.00%)	0.175 ± 0.011			
Amount of starting material				
Turnover		2	6.141	**0.005**
1 (79.35%)	0.106 ± 0.005			
2 (79.46%)	0.096 ± 0.004			
3 (92.15%)	0.247 ± 0.038			
Nestedness-resultant		2	0.425	0.657
1 (20.65%)	0.073 ± 0.005			
2 (20.54%)	0.059 ± 0.002			
3 (7.85%)	0.049 ± 0.002			
Total beta diversity		2	4.898	**0.013**
1 (100.00%)	0.153 ± 0.009			
2 (100.00%)	0.137 ± 0.005			
3 (100.00%)	0.287 ± 0.030			
Vazella pourtalesii				
Type of starting material				
Turnover		1	0.327	0.570
Dried tissue (97.03%)	0.489 ± 0.059			
Homogenate (96.63%)	0.451 ± 0.046			

Nestedness-resultant		1	0.696	0.409
Dried tissue (2.97%)	0.146 ± 0.019			
Homogenate (3.37%)	0.186 ± 0.035			
Total beta diversity		1	0.143	0.707
Dried tissue (100.00%)	0.569 ± 0.024			
Homogenate (100.00%)	0.554 ± 0.014			
Amount of starting material				
Turnover		2	0.165	0.849
1 (96.29%)	0.513 ± 0.025			
2 (97.05%)	0.477 ± 0.055			
3 (95.02%)	0.480 ± 0.044			
Nestedness-resultant		2	0.645	0.530
1 (3.71%)	0.132 ± 0.010			
2 (2.95%)	0.177 ± 0.030			
3 (4.98%)	0.115 ± 0.037			
Total beta diversity		2	0.022	0.978
1 (100.00%)	0.569 ± 0.015			
2 (100.00%)	0.563 ± 0.021			
3 (100.00%)	0.559 ± 0.012			

1141

1142

Table 4. Results of PERMANOVA used to compare variation in community composition according to type and amount of starting material, individually and combined.

	df	F	R^2	*P*	Stress
Phakellia ventilabrum					
Turnover					0.065
Type of starting material	1	1.917	0.034	0.194	
Amount of starting material	2	7.513	0.269	**0.004**	
Type of starting material : Amount of starting material	2	1.981	0.071	0.141	
Nestedness-resultant					0.045
Type of starting material	1	1.941	0.054	0.194	
Amount of starting material	2	0.385	0.021	0.659	
Type of starting material : Amount of starting material	2	-0.756	-0.042	0.994	
Total beta diversity					0.077
Type of starting material	1	1.323	0.026	0.265	
Amount of starting material	2	5.571	0.220	**0.001**	
Type of starting material : Amount of starting material	2	1.638	0.065	0.135	
Vazella pourtalesii					
Turnover					0.119
Type of starting material	1	3.058	0.067	**0.012**	
Amount of starting material	2	0.664	0.030	0.757	
Type of starting material : Amount of starting material	2	0.507	0.022	0.863	
Nestedness-resultant					0.190
Type of starting material	1	-6.207	-0.145	1.000	
Amount of starting material	2	2.410	0.113	0.196	
Type of starting material : Amount of starting material	2	2.105	0.098	0.249	
Total beta diversity					0.113
Type of starting material	1	2.055	0.045	**0.019**	
Amount of starting material	2	1.026	0.045	0.396	
Type of starting material : Amount of starting material	2	0.751	0.033	0.849	

1148
1149
1150 **Figure 1.** Flow diagram illustrating the laboratory workflow for the phased experiment.

1153 **Figure 2.** Box plots summarising **(a)** total DNA yield, **(b)** post-PCR DNA concentration,
1154 **(c)** A260/280 ratios, and **(d)** A260/A230 ratios for each sample from each sponge
1155 species (Horniman invasive sponge (H-INV) and *Vazella pourtalesii* (VAZ)) processed
1156 with each ethanol treatment and each DNA extraction protocol.

1157
1158
1159 **Figure 3.** Bubble plot summarising proportional read counts for each taxon and
1160 detection consistency across extraction replicates for samples from each sponge
1161 species (Horniman invasive sponge (H-INV) and *Vazella pourtalesii* (VAZ)) that were
1162 processed with each ethanol treatment (blot, spin, wet) and DNA extraction protocol
1163 (MUS, MUT, QBT, QBT-ZYMO and QPS).
1164
1165

1166
1167

Figure 4. Differences in fish taxon richness **(a)**, proportional read counts **(b)** and community similarity **(c)**, including turnover **(i)** and nestedness-resultant **(ii)** components of total beta diversity **(iii)**, for samples processed with each ethanol treatment (blot, spin, wet) and each DNA extraction protocol (MUT, QBT and QBT-ZYMO). Abbreviations: Horniman invasive sponge (H-INV), *Vazella pourtalesii* (VAZ).

1173
1174

1175

1176

Figure 5. Box plots summarising **(a)** total DNA yield, **(b)** post-PCR DNA concentration, **(c)** A260/280 ratios, and **(d)** A260/A230 ratios for each sample from each sponge species (*Phakellia ventilabrum* (PHAK) *Vazella pourtalesii* (VAZ)) processed using different types and amounts of starting material.

Figure 6. Bubble plots summarising proportional read counts for each taxon and detection consistency across extraction replicates for different types and amounts of starting material from **(a)** *Vazella pourtalesii* (VAZ) and **(b)** *Phakellia ventilabrum* (PHAK).

1189
1190
1191 **Figure 7.** Differences in taxon richness **(a)**, proportional read counts **(b)** and community
1192 similarity **(c)**, including turnover **(i)** and nestedness-resultant **(ii)** components of total
1193 beta diversity **(iii)**, for fish detected in samples of each type and amount of starting
1194 material from each sponge species (*Phakellia ventilabrum* (PHAK), *Vazella pourtalesii*
1195 (VAZ)).
1196
1197