Nationwide genomic biobank in Mexico unravels demographic history and complex trait architecture from 6,057 individuals

Mashaal Sohail[1, 2,3,#], Amanda Y. Chong[4,*], Consuelo D. Quinto-Cortes[1,*], María J. Palma-Martínez[1,*], Aaron Ragsdale[1], Santiago G. Medina-Muñoz[1], Carmina Barberena-Jonas[1], Guadalupe Delgado-Sánchez[5], Luis Pablo Cruz-Hervert[5,6], Leticia Ferreyra-Reyes[5], Elizabeth Ferreira-Guerrero[5], Norma Mongua-Rodríguez[5], Andrés Jimenez-Kaufmann[1], Hortensia Moreno-Macías[7,8], Carlos A. Aguilar-Salinas[9], Kathryn Auckland[4], Adrián Cortés[10], Víctor Acuña-Alonzo[11], Alexander G. Ioannidis[12], Christopher R. Gignoux[13], Genevieve L. Wojcik[14], Selene L. Fernández-Valverde[1], Adrian V.S. Hill[4,15], María Teresa Tusié-Luna[7], Alexander J. Mentzer[4, 10], John Novembre[2,16], Lourdes García-García[5,#], Andrés Moreno-Estrada[1,#]

[1]Laboratorio Nacional de Genómica para la Biodiversidad (LANGEBIO), Unidad de Genómica Avanzada (UGA), CINVESTAV, Irapuato, Guanajuato, México.
[2]Department of Human Genetics, University of Chicago, Chicago, IL, USA.
[3]Centro de Ciencias Genómicas (CCG), Universidad Nacional Autónoma de México (UNAM), Cuernavaca, Morelos, México.
[4]The Wellcome Centre for Human Genetics, University of Oxford, Oxford, UK.
[5]Instituto Nacional de Salud Pública (INSP), Cuernavaca, Morelos, México.
[6]Facultad de Odontología, Universidad Nacional Autónoma de México (UNAM), Ciudad de México, México.
[7]Unidad de Biología Molecular y Medicina Genómica, UNAM-INCMNSZ, México City, México.
[8]Universidad Autónoma Metropolitana, México City, México.
[9]Division de Nutrición, Instituto Nacional de Ciencias Médicas y Nutrición Salvador Zubirán, Mexico City, México.
[10]Big Data Institute, Li Ka Shing Centre for Health Information and Discovery, University of Oxford, Oxford, UK.
[11]Escuela Nacional de Antropología e Historia (ENAH), Mexico City, México.
[12]Department of Biomedical Data Science, Stanford University, Stanford, CA, USA.
[13]Colorado Center for Personalized Medicine, University of Colorado Anschutz Medical Campus, Aurora, CO, USA.
[14]Department of Epidemiology, Johns Hopkins Bloomberg School of Public Health, Baltimore, MD, USA.
[15]The Jenner Institute, University of Oxford, Oxford, UK.
[16]Department of Ecology and Evolution, University of Chicago, Chicago, IL, USA.

*co-authorship
#co-corresponding

Emails for correspondence: andres.moreno@cinvestav.mx, garcigarml@gmail.com, mashaal@ccg.unam.mx

Abstract:

Latin America continues to be severely underrepresented in genomics research, and fine-scale genetic histories as well as complex trait architectures remain hidden due to the lack of Big Data. To fill this gap, the Mexican Biobank project genotyped 1.8 million markers in 6,057 individuals from 32 states and 898 sampling localities across Mexico with linked complex trait and disease information creating a valuable nationwide genotype-phenotype database. Through a suite of state-of-the-art methods for ancestry deconvolution and inference of identity-by-descent (IBD) segments, we inferred detailed ancestral histories for the last 200 generations in different Mesoamerican regions, unraveling native and colonial/post-colonial demographic dynamics. We observed large variations in runs of homozygosity (ROH) among genomic regions with different ancestral origins reflecting their demographic histories, which also affect the distribution of rare deleterious variants across Mexico. We analyzed a range of biomedical complex traits and identified significant genetic and environmental factors explaining their variation, such as ROH found to be significant predictors for trait variation in BMI and triglycerides.

The genetic architecture of complex traits in admixed genomes cannot be understood outside the
context of their underlying histories. Present-day Mexico covers seven Mesoamerican regions
with rich civilizational histories[1]. Archaeology and anthropology regionalize Mexico into the
north of Mexico, the north of Mesoamerica, the center, occident and gulf of Mexico, Oaxaca and
the Mayan region[2](Fig. 1a). These are based on noting specific pre-Hispanic civilizations and
cultures, which began flourishing very early in the Mayan region, in Oaxaca, in the occident and
in the gulf of Mexico[3], and later in the center and north of Mesoamerica. Such histories have also
been used to classify Mesoamerican chronology into preclassical, classical, postclassical,
colonial, and postcolonial periods.
74
In the last five hundred years, Spanish colonization has left an indelible mark on this native
tapestry. In a colonial context, ancestries that trace to European, African and Asian sources can
be identified in living Mexicans, however they vary in structure and timing between
Mesoamerican regions[4,5-8,9-11]. The heterogeneity of such a mixture at a genetic level has been
characterized, revealing extensive fine-scale population substructure and ancestry sources across
Mexico[12-16]. These studies have also identified genes potentially under selection for some traits in
different native groups[12,15].
82
Further, such varying genetic histories, as captured by ancestry distributions, have been shown to
impact variation in complex traits in Mexicans in traits such as lung force capacity[12], and a
number of other complex traits and diseases[17]. Nevertheless, a large gap remains in the
representation of Mexicans from across Mexico in cohorts with linked genotypes and
phenotypes, which could enable finer-scale studies of genetic history and a better understanding
of complex trait architecture among individuals with diverse ancestries from the Americas and
those living in rural areas[18]. Past efforts have been limited to studying individuals from the United
States and Mexico City and have not simultaneously modelled the influence on complex trait
variation of a rich array of genetic and environmental factors as is possible with a nationwide
Biobank.
93
To bridge this gap, we launched the Mexican Biobank (MXB) project, densely genotyping 6,057
individuals from all 32 states across Mexico (Fig. S1-S2) recruited as part of the National Health
Survey in 2000 (ENSA2000), which sampled more than 40,000 participants nationwide. To
select the samples for genomic and biochemical characterization, we enriched for those
individuals that can speak an indigenous language in each state while maximizing the
representation of rural localities (~70% of the MXB out of a total of 898 localities, Fig. S2-S5) to
increase the representation of indigenous ancestries. The MXB is 70% female and comprised of
individuals born between 1910 and 1980, all sampled in the year 2000[18](Table S1). These
individuals were genotyped at ~1.8 million SNPs and have linked information for traits such as
height, BMI, triglycerides, glucose, cholesterol, blood pressure and various socioeconomic and
biogeographical markers (Table S2).
105
Here, we leverage rich archaeological and anthropological information to guide a regionalized
analysis of Mexico, and harness the power of local ancestry estimation genome-wide and
segments of identity-by-descent (IBD) to decipher fine-scale genetic histories using ancestry-
specific approaches to denote origins and historical population size changes[19,20]. We reveal a very
heterogeneous landscape of both, painting a genetically informed picture of varying demographic

trajectories of Mesoamerican civilizations, as well as colonial migrations and dynamics in
different regions of Mexico. We further investigate the role of these evolutionary histories as
captured by proxies of genetic ancestries in shaping genetic variation and complex traits patterns
in Mexico today. We show that these histories result in marked geographic and ancestry-specific
patterns in the distributions of runs of homozygosity (ROH) and of the genomic burden of rare
deleterious mutations.
117
Lastly, we study the impact of these histories which could associate certain trait-relevant
genotypes with certain genetic backgrounds, along with portions of the genome in ROH and
other sociocultural and biogeographical factors capturing environmental context, on creating trait
variation in complex and medically-relevant traits such as height, BMI, triglycerides, glucose
levels, and others in Mexico. Our results can help guide sampling and design for future genetic
mapping efforts by determining which environmental and genetic axes maximize trait variation,
to help increase power in genome-wide association studies. They can also help determine cases
where environmental interventions are more likely to bring a desired improvement in public
health.
127
Genetic structure across Mexico is shaped by native diversity and historical migrations.
We begin by excavating the population structure in the MXB at different geographic resolutions
and time-scales. Principal components analysis (PCA)[21] captures predominant axes of genetic
similarity. Further, a proxy for genetic ancestries from different regions can be quantified using
ADMIXTURE[22] (see note on genetic ancestries in methods). When we visualize the Mexican
biobank samples using PCA with individuals from around the world (1000 Genomes[23], HGDP[24],
and PAGE[25]), we find that most Mexican individuals lie on a cline between living Europeans and
indigenous Americans, which we interpret as reflecting the history of admixture in Mexico since
Spanish colonization (Fig. 1B, Figs. S6). We also observe a "pull" towards present-day Africans,
likely reflecting the genetic impact of the trans-Atlantic slave trade during the colonial period
and subsequent migrations that brought many Africans to Mexico. When analyzed alone, the
MXB individuals show a striking population substructure delineation between the Mayan region
and the rest of the country (Fig. 1C, Fig. 1E, Figs. S7-S17). In the rest of the regions, only a
subtle genetic substructure mirroring Mesoamerican geography is visible in the MXB, likely
reflecting the effects of movement and mating among the different regions sampled in the year
2000.
144
We infer ancestry proxies at different geographic resolutions reflecting mating dynamics in the
colonial and post-colonial periods by analyzing MXB with global individuals as well as with
only native individuals using the software ADMIXTURE (Fig. 1D, Table S3, Fig. S11). We
observe that individuals in the Mexican Biobank are inferred to be admixed with varying degrees
of ancestries that are found most abundantly in individuals of the Americas ("American
ancestries") and Europe ("European ancestries"). Higher levels of ancestries from the Americas
were inferred in the central and southern states of Mexico, compared to the northern states. We
observe the largest genetic differentiation as measured using Fst along a north to southeast cline
(Fig. S14-17). We observe some ancestries from Africa in individuals found in every single state
(Table S3). We observe that only 3 states in the north (Chihuahua, Nuevo Leon and Sinaloa)
have more ancestries on average from Europe than ancestries from the Americas, with ancestries
from the Americas being the majority ancestries in every other state (Table S3). Lastly, we note

the presence of a small but significant proportion of ancestries from East Asia in almost every state (0-2.3%), the highest in the state of Guerrero (2.3%), and an even more modest but significant amount of ancestries from South Asia in the majority of states as well (0-0.8%).

We use an ancestry-specific PCA or MDS approach to pinpoint the origins of the ancestries from the Americas, Africa, and Asia observed in the Mexican Biobank within those regions. For ancestries from the Americas, we observe that such ancestries tend to originate from indigenous cultures predominant in the region an individual is from (Fig. S18, Table S4). For example, such ancestries in the Yucatan peninsula originate from the Maya and Tzotzil. We observe that most ancestries from Africa in Mexico originate from West Africa[26], in agreement with historical records of shipping voyages from the trans-Atlantic slave trade (Fig. S19)[4]. For the ancestries from East Asia, we find for individuals in Guerrero, such segments projecting to East Asian regions that were linked to the Manila Galleon trade, as reported in preliminary findings[16]. In contrast, for individuals from northern states such as Chihuahua, such segments project to China and Japan, likely reflecting later migrations from East Asia to Mexico (Fig. S20). Similarly, for ancestries from South Asia, as illustrated for individuals from Guerrero, the landing state of the Manila Galleon, we observe diverse roots in South Asia (Fig. S21-S22).

The identification of individuals with genetic ancestries from East Asia in Mexico dating to the Manila Galleon trade agrees with one other recent study[16]. These genetic observations are plausibly explained by the poorly appreciated history of Mexico's Asian population[5-8]. Using voyage records, slaving documents, and other sources, historians have documented arrivals from the Manila port in the Spanish Philippines to the Acapulco port in Mexico through the 16[th] and 17[th] centuries, with origins as diverse as the Philippines, Indonesia, Malaysia, India, Bengal, and Sri Lanka (though they were collectively referred to as "Chinos" by the colonists). They entered from the Acapulco port, but moved through most of Mexico, with even what was called the "China Road" existing between Acapulco and Mexico City. There have also been later 19[th] and 20[th] century migrations from China and Japan, especially to the north of Mexico, and inheritance from these ancestors likely explains part of the ancestries from East Asia we observe in the northern states of the MXB today[9-11].

Ancestry-specific IBD tracts recover 200 generations of genetic history within Mexico
Apart from the small amount of recent ancestry from Asia discussed above, the ancestry of contemporary Mexicans arises predominantly from lineages that would have been found in Central America, Western Europe, and West Africa prior to 15[th] century. Each of these sources had different demographic histories prior to and after their arrival in present-day Mexico. To reveal the more recent history of population sizes of these three ancestries, we analyze identity-by-descent segments[27] overlapped with their corresponding local ancestry inference[19]. We use this approach to estimate effective population size (N_e) trajectories 200 generations into the past for these ancestries in the Mexican Biobank as a whole, as well as in specific Mesoamerican regions (Figure 2). This analysis helps us reveal genetic histories of present-day Mexicans, which is of anthropological interest, as well as relevant for patterns of genetic and complex trait variation as shown in later sections.

In the entire MXB, contextualized using Mesoamerican chronology (Fig. 2A), we find that, for indigenous lineages, (i.e., those present in the area before the arrival of the Spaniards), the

effective population size went through a slow and steady decline in the classical period (250 – 900 CE). This decline was followed by an increase in the postclassical period (900 – 1521 CE), right before the arrival of the Spaniards, and then a decline later in the colonial period (1521 – 1821 CE) and in the post-colonial period (1821 – present) (Fig. 2B).

Further, we observe fine-scale structure in N_e trajectories for indigenous lineages which we interpret in the context of the different cultural histories of Mesoamerican regions (Fig. 2C)[1]. Starting chronologically, archaeologists document that Mesoamerican civilizations flourished very early in the Mayan region, in Oaxaca, in the occident and in the gulf of Mexico, where we also observe large N_e already in the classical period[3]. For example, in the gulf, where we observe high N_e since the pre-classical period (2500 BCE – 250 CE), there is archaeological evidence, among a myriad of other groups, of the Olmecs in the pre-classical period, the Totonacs in the classical period, and the Huastecs in the post-classical period[28]. In Oaxaca, we observe N_e rapidly growing in the pre-classical to the classical period, in line with archaeological inferences that the Zapotecs were already starting to create sedentary settlements in the pre-classical period followed by a rise in social and political structures in the classical period. This was followed by a more militaristic period in the post-classical causing warfare[29], and our genetic evidence suggests a significant population decline toward the end of the post-classical period. In the Yucatan peninsula, the Maya had prominent civilizational spread in the classical period (peak N_e observed), and started going through a slow decline only in the post-classical period due to what archaeologists have inferred as a combination of different political and ecological factors, and this trajectory is supported in the N_e trend[3]. We further observe that native groups in both Oaxaca and the Mayan peninsula started to increase in population size again through the colonial and post-colonial (1821 – present) periods, after the arrival of the Spaniards.

This is in contrast with the center and north of Mesoamerica, where the Aztec empire had a strong-hold most recently and where we see increasing N_e in the post-classical right before the arrival of the Spaniards and into part of the colonial period, after which we start to see a population decline in N_e. Thus, the decline in N_e after the arrival of the Spaniards is most prominent in the center and north of Mesoamerica, and is actually followed by an increase in Oaxaca and the Mayan region, where native ancestries from Central America are most prevalent today as evidenced by the Admixture analysis (Table S3). As generational time can vary, we present our analysis at two extremes of 20 and 30 years per generation[30](Fig. S23 and 2C, respectively).

We observe that ancestries from Western Europe that entered the contemporary Mexican gene pool went through a sharp decline in effective population size during the colonial period. The extent of the founder effect varied by region, with the strongest effect seen in Oaxaca and the Mayan region (Fig. S24, S25). Similarly, ancestries from Western Africa in Mexico revealed stronger founder effects that varied by region with N_e ranging between 10^3 and 10^4 in the colonial period. The population size in the post-colonial period continued to grow in some regions such as the occident and north of Mexico and the Mayan region, compared to others (Fig. S26, S27).

Demographic histories impact patterns of genetic variation in Mexico
Small ROH prevalence is correlated with ancestry proxies

We next analyze the patterns of ROH in the MXB and their relationship with geography and
ancestry proxies. ROH patterns help further illuminate demographic and mating histories of
Mexicans[31], and are relevant for variation in complex traits if trait-relevant variation is partially
recessive[32]. We identify ROH ($\geq$ 1 Mb) in the Mexican Biobank and observe that both the
number of ROHs and the total length of ROH per individual increases as we move from north to
southeast in the country (Fig. S28-29). We confirm that this is primarily due to individuals with
more genetic ancestries from Central America also having more ROH in their genomes (Fig. 3A,
Figs. S30-31, Table S5).
256
Next, we asked whether this signal of higher ROH associated with higher ancestry from Central
America is due to historical bottlenecks and small population sizes, or due to consanguinity. A
bottleneck event or a long-term small population size will result in a large number of small
ROH[33]. Consanguinity or marriage between relatives would instead result in fewer but longer
ROH[33]. To answer this question, we analyze the total length of ROH per individual in each state,
after first partitioning ROH by size (Fig. 3B). We observe that there are more ROH per
individual moving southwards in Mexico in large part due to small ROH (smaller than those
expected from recent consanguinity e.g., < 8 Mb), implying that bottlenecks and small
population sizes rather than consanguinity have been largely responsible for more ROH in
individuals with higher ancestries from Central America.
267
Lastly, we observe that ROH that are found on segments of the genome from Central America
are more frequently found in younger individuals compared to older individuals (Spearman's rho
= 0.31, p = 0.016) (Fig. 3C). We also verified that this correlation with birth year primarily
derives from small ROH (rho = 0.35, p = 0.006), and small ROH found on genomic segments
from the Americas (rho = 0.39, p = 0.002) (Fig. 3C). This result is at least partly due to younger
individuals carrying more ancestries from Central America compared to older individuals,
especially in the rural localities (Fig. S51), and agrees with recent observations about ancestry
and ROH made in Mexican-Americans[17]. The observation of higher ancestries from Central
America in younger individuals in rural areas may be due to either individuals in rural areas pro-
creating at a higher rate, or individuals with other ancestries moving out from rural to urban
areas. The observation of a larger number of small ROH in younger individuals in the MXB is
relevant for parsing the genetic architecture of complex traits and diseases, especially those with
a recessive component.
281
Rare deleterious variant burden is correlated with ancestry proxies
We also investigated the effects of demographic history on the frequency distribution of genetic
variants. If such an effect exists for variants that contribute to trait variation, it would imply
varying genetic architectures for some traits that may be captured by ancestry proxies. This
analysis is motivated by previous theoretical and empirical work showing that undergoing a
bottleneck changes the allele frequency distribution in the group that experienced the
bottleneck[23,34,35]. In particular, rare variants are lost or increase in frequency after the bottleneck.
289
We can evaluate this effect within and between genes in the genome by calculating the genome
load of genetic variants, or mutation burden. We calculated the genome-wide mutation burden by
summing the derived alleles at each SNP in the genome for different types of variants
(intergenic, synonymous, putatively deleterious). When considering only rare variants (DAF<=

5%), we observe that the total number of rare variants is negatively correlated with ancestries
from Central America, while it is positively correlated with ancestries from Western Europe and
West Africa (Fig. 4). Thus, individuals with higher ancestries from Central America carry fewer
rare variants likely due to a history of more bottlenecking events compared to individuals with
higher ancestries from Western Europe and West Africa. We observed the same general pattern
for different types of variants. However, this effect of varying demographic histories on variant
frequencies is strongest for putatively neutral variants (Fig. 4). We have verified these
observations for rare variants with whole-genome sequences from the subset of Mexicans living
in Los Angeles (MXL) from the 1000 Genomes Project to rule out ascertainment biases due to
the array genotyping as the source of this effect (Fig. S50).
304
As shown in previous studies, the effect of demographic history on the total mutation burden is
minimal in MXB[35-37]. When we consider all frequencies (DAF <= 100%), we see only a small
correlation between mutation burden and ancestries from different regions likely due to
ascertainment bias as this correlation does not persist in whole genome sequence data from 1000
Genomes (Fig. 4, Fig. S50). This is because while some rare variants are lost, some increase in
frequency, compensating the total mutation burden, which overall remains unchanged.
311
Complex traits display varying roles of genetics and environment
Lastly, we assessed the contribution of genetic variation towards impacting variation in complex
traits or disease in Mexico (Fig. S32). For example, ROH have been previously shown to have
associations with a broad range of complex traits, and estimated to be negatively associated with
height, weight, and cholesterol[32]. Such associations, if due to genetic factors, point towards a
recessive architecture of the traits. Further, genetic ancestry proxies can also be associated with
complex traits due to genetic factors, or due to differential experience of discrimination and other
socioeconomic factors (Fig. S32). Such genetic factors can be different distributions of ROH or
other differential patterns of genetic variation caused by demographic and environmental
histories that vary among ancestries, and that can lead to the association of particular causal
genotypes with ancestry proxies. Indeed, genetic ancestry proxies in Mexico are correlated with
the number and length of ROH (Fig. 3A). We therefore model the association of genetic factors
such as ancestry proxies and ROH with trait variation in the same model to disentangle these
effects. As genetic ancestry proxies can also reflect differential environmental exposures, we
consider in our model several environmental factors, to further disentangle the role of genetic
factors reflected in ancestry proxies compared to environmental factors. Our model therefore
also includes variables available in the MXB related to discrimination, socioeconomic
opportunities, and living environment (collectively called sociocultural and biogeographical
factors). We also account for cryptic relatedness and potential unmodelled environmental factors
using a genetic relationship matrix and city or town of origin as random effects in a mixed model
framework. Significant associations would give insight into the architecture, including the
genetic architecture, of the trait in Mexico to help guide future efforts in genetic mapping, and in
considering other interventions towards improving public health.
335
Aiming to first understand how the traits are distributed geographically and relative to single
model covariates, we first visualize average trait values by units of our biogeographical and
sociocultural factors to understand the dimensions of trait variation (Figs. 5A, S33-41). Next, we
use a mixed model to estimate the contribution of genetic factors to trait variation jointly

modelled with the environmental factors (Figs. 5B, D-F, S42-49). Finally, we visualize trait
values by birth year to assess their changes over time. We focus on several quantitative traits:
height, BMI, triglycerides, cholesterol, glucose, blood pressure, and others. Our test predictive
variables in the full model include genetic factors (genetic ancestry proxies from ADMIXTURE
and ancestry-specific MDS analyses, and ROH in kb in each genome), life history factors (age,
sex), sociocultural factors (educational attainment as a proxy for income levels (Fig. S42),
whether they speak an indigenous language or not as a proxy for differential experience of
discrimination/other cultural factors such as diet, whether they live in an urban or rural
environment), and biogeographical factors (altitude, latitude and longitude). For height and other
traits analyzed, a significant association with ancestry proxies could reflect the association of
particular causal genotypes with those ancestries or associated unmodelled environmental factors
such as nutrition. If the association is due to genetic factors, it should still not be interpreted as
deterministic of a trait value, but makes a case for inclusion of diverse individuals in genetic
studies of complex traits.
354
Height. When viewed as averages per state, height values show a clear increasing pattern from
southeast to northwest in the MXB (Fig. 5A). Even though every state shows a large variance
(Fig. 5A), height shows a significant correlation with both latitude and longitude univariately
(Figs. S34-35). With our explanatory mixed model, we can explain 66.33% of the variance for
height. We find that individuals with higher ancestries from Central America are significantly
shorter ($\beta = -0.42$, $p < 2.2 \times 10^{-16}$) (Figure 5b). Further, considering ancestries at a finer
resolution, we observe decreased height with a change in ancestries from the North of Mexico
(Huichol, Tarahumara) to the Mayan region (Tojolabal, Maya) (Fig. S48). Notably, individuals
having higher educational attainment are also estimated to be taller. After Bonferroni correction
across traits and predictors, the relationship between ROH and height is not statistically
significant ($\beta = -0.07, p = 0.03$). Nevertheless, younger individuals with any range of
ancestries from Central America are taller than older individuals with the same ancestries (Fig.
5C). As the positive correlation between birth year and height for all individuals regardless of
their ancestries demonstrates, height can also vary due to environmental factors or aging.
369
370 *Body mass index (BMI).* BMI similarly shows a significant correlation with latitude and
371 longitude univariately (Figs. S34-35). Our full mixed model explains 30.10% of the variation in
372 BMI. In the full model, ROH remain significantly associated with lower BMI ($\beta = -0.18, p = 3.95 \times 10^{-5}$), while ancestry does not (Fig. 5D). BMI is also significantly correlated with birth
373 3.95×10^{-5}), while ancestry does not (Fig. 5D). BMI is also significantly correlated with birth
374 year, increasing with older age, as well as with being female (Fig. 5D).
375
Triglycerides. There are clear differences in levels by region that are more striking for
cholesterol (see below) than for triglycerides (Fig. S33). Our full model explains 37.23% of the
variation in triglyceride levels. Ancestries from Central America are significantly associated with
higher triglyceride levels ($\beta = 0.19, p = 3.17 \times 10^{-4}$) while the ROH carried by an individual
are significantly associated with lower triglyceride levels ($\beta = -0.18, p = 1.4 \times 10^{-4}$) (Fig.
5D). Notably, age, being male, lower educational attainment and high altitude are also associated
with higher triglyceride levels (Fig. 5D).
383
Cholesterol. Cholesterol levels show a significant correlation with latitude and longitude
univariately (Fig. S33-35). In the full model (explaining 29.49% of trait variation), we do not see

any correlation with genetic ancestry proxies or ROH, but we estimate significantly lower
cholesterol in individuals who speak an indigenous language ($\beta = -0.26, p = 6.93 \times 10^{-7}$).
We also estimate higher cholesterol in those living in an urban environment, at high altitude or of
a higher age (Fig. S43). For HDL and LDL levels, we similarly find a significantly lower
cholesterol in those who speak an indigenous language but not related to ancestry (Fig. S43)
likely indicating that cultural/diet factors are stronger than the genetic factors tested here. Living
in in an urban environment is also significantly associated with high HDL and LDL levels (Fig.
S43).
394
Glucose. Glucose levels show a significant correlation with latitude in univariate analysis (Fig.
S34). In the full model (explaining 28.62% of the trait variation), we estimate ancestries from
Central America to be significantly correlated with higher glucose levels ($\beta = 0.23, p =$
2.452×10^{-5}). Glucose also remains significantly associated with latitude (increasing
northwards) and higher age (Figure 5d). For fasting glucose, where we reduce sample size by
about four-fifths, ancestry from Central America still has a positive estimated coefficient (Fig.
S45), but it is not significant with the smaller sample size.
402
403 *Other traits.* We also analyze creatinine, and systolic and diastolic blood pressure
404 (Fig. S44). Individuals that speak an indigenous language have significantly lower creatinine
405 ($\beta = -0.18, p = 1.32 \times 10^{-3}$). Living in an urban environment and altitude are significantly
406 associated with higher creatinine. Only age and sex are significantly associated with diastolic and
407 systolic blood pressure (adjusted by medication status) (Fig. S44).
408
Overall, ancestries from Central America are significantly associated with trait variation for
height, triglycerides, and glucose levels (Fig. S46), the ROH present in a genome with BMI and
triglycerides (Fig. S47) levels, and native ancestry variation within Central America with height
variation (Fig. S48-49). In contrast, cholesterol levels, creatinine levels, and blood pressure are
significantly associated with environmental, but not genome-wide genetic factors. This does not
rule out the effect that specific gene variants may have in the variation of these traits, as
illustrated before by candidate gene approaches discovering functional variants exclusive to the
Americas[38].
417
Conclusion.
Our work is a demonstration of the value of generating genotype-phenotype data on
underrepresented populations to reveal lesser-known genetic histories and generate findings of
biomedical relevance. It is also an illustration of the joint modeling of genetic and environmental
effects to reveal the etiology of complex traits and disease. In this project, we ensured diverse
indigenous and rural presence in our sampling strategy, considered the fluidity of ancestries from
different local and global regions in our analyses and evaluated their reflection in genetic and
disease-relevant complex trait variation. By leveraging the largest nationwide genomic biobank
in Mexico, we find diverse sources of ancestries in Mexico in light of its unique history, infer
population size changes and runs of homozygosity using ancestry-specific haplotype identity that
reveal an elaborate fine-scale structure in the country. We also show that demographic history
affects the frequency distribution of genetic variants thus changing how many rare variants
individuals with different ancestries carry. We observe a significant impact of genetic ancestries,
ROH, as well as socioeconomic and biogeographic variables on a variety of complex traits

implicating the importance of both genetic and environmental factors in explaining complex trait
variation and in considerations of potential public health interventions. Our results can inform
the design of future studies in other admixed populations and the MXB will hopefully motivate
additional efforts to strengthen local research capacity across Latin America and benefit
underserved populations globally.

Funding

This work was supported by "The Mexican Biobank Project: Building Capacity for Big Data
Science in Medical Genomics in Admixed Populations", a binational initiative between Mexico
and the UK co-funded equally by CONACYT (Grant number FONCICYT/50/2016), and The
Newton Fund through The Medical Research Council (Grant number MR/N028937/1) awarded
to AME and AH. MS was also supported by the Chicago Fellows program of the University of
Chicago. Training activities in Mexico were hosted by CINVESTAV and supported in part by
CABANA, a capacity strengthening project for bioinformatics in Latin America, funded by the
Global Challenges Research Fund (GCRF) of the UK.

Data availability

449 The dataset for the 6,057 newly genotyped individuals from the MX biobank project are
450 available at the European Genome-phenome Archive (EGA) through a Data Access Agreement
451 with the Data Access Committee (EGA accession number in process).

Code availability

454 All custom scripts/approaches described in Methods will be made available from
455 https://github.com/msohail88/MXB_popstruct_complextraits on publication and all existing
456 software packages and versions used are noted in Methods

Acknowledgments

We thank the participants of the *Encuesta Nacional de Salud, 2000* (2000 National Health
Survey, ENSA 2000), conducted in Mexico nationwide by the *Secretaría de Salud (*Health
Secretariat) and the *Instituto Nacional de Salud Pública* (National Institute of Public Health,
INSP). We are grateful to Mauricio Hernández and Celia Alpuche-Aranda for Institutional
support from INSP, Mitzi Flores, Rocío Nájera, and Adriana Garmendia for project management
support, and to Carlos Conde, Victor Guerrero Lemus, Armando Mendez Herrera, Cruz Portugal
García, Rosario Rodriguez, and Manuel Velazquez Mesa for biobank maintenance and sample
preparation. We thank Mary Ortega, Cecilia Gutiérrez, and Sara García for technical assistance,

Jacob Cervantes for IT support, Harald Ringbauer for useful advice with the ROH analysis, Juan
Esteban Rodriguez for helpful conversations about population structure in Mexico, and Arslan
Zaidi for useful comments on an earlier draft of this manuscript.

Inclusion and Ethics statement

References

1. Michael D. Coe, R. K. *Mexico: From the Olmecs to the Aztecs*. (Thames & Hudson, 2013).

2. Vela, E. Áreas culturales: Oasisamérica, Aridamérica y Mesoamérica. *Arqueología*

Mexicana, edición especial 28–29 (2018).

3. Gugliotta, G. The Maya: Glory and Ruin. *The National Geographic Magazine* 68–109

(2007).

4. Trans-atlantic slave trade database. *Slave Voyages.* https://www.slavevoyages.org/ (2018).

5. Seijas, T. *Asian Slaves in Colonial Mexico: From Chinos to Indians*. (Cambridge University

Press, 2014).

6. Chávez, C. P. M. El alcalde de los chinos en la Provincia de Colima durante el siglo XVII:

Un sistema de representación en torno a un oficio. *Letras Históricas E-ISSN: 2448-8372*

(2009).

7. Keresey, D. O. LA ESCLAVITUD ASIÁTICA EN EL VIRREINATO DE LA NUEVA ESPAÑA,

1565-1673. *Hist. Mex.* **61**, 5–57 (2011).

8.   Carrillo, R. Asia llega a América. Migración e influencia cultural asiática en Nueva España

(1565-1815). *Asiadémica* **3**, 81–98 (2014).

9.   Mishima, M. E. O. *Siete migraciones japonesas en México: 1890-1978*. (El Colegio de

Mexico, 1982).

10. Augustine-Adams, K. Prohibir el mestizaje con chinos: solicitudes de amparo, Sonora,

1921-1935. *Rev. Indias* **72**, 409–432 (2012).

11. Guillén, M. L. Vivir para trabajar. La inserción laboral de los inmigrantes chinos en Chiapas,

siglos XIX y XX. *Studium: Revista de humanidades* 113–140 (2013).

12. Moreno-Estrada, A. *et al.* The genetics of Mexico recapitulates Native American

substructure and affects biomedical traits. *Science* **344**, 1280–1285 (2014).

13. García-Ortiz, H. *et al.* The genomic landscape of Mexican Indigenous populations brings

insights into the peopling of the Americas. *Nat. Commun.* **12**, 5942 (2021).

14. Romero-Hidalgo, S. *et al.* Demographic history and biologically relevant genetic variation of

Native Mexicans inferred from whole-genome sequencing. *Nat. Commun.* **8**, (2017).

15. Ávila-Arcos, M. C. *et al.* Population History and Gene Divergence in Native Mexicans

Inferred from 76 Human Exomes. *Mol. Biol. Evol.* **37**, 994–1006 (2020).

16. Rodríguez-Rodríguez, J. E. *et al.* The genetic legacy of the Manila galleon trade in Mexico.

Philos. Trans. R. Soc. Lond. B Biol. Sci. **377**, 20200419 (2022).

17. Spear, M. L. *et al.* Recent shifts in the genomic ancestry of Mexican Americans may alter

the genetic architecture of biomedical traits. *Elife* **9**, (2020).

18. Popejoy, A. B. & Fullerton, S. M. Genomics is failing on diversity. *Nature* **538**, 161–164

(2016).

19. Browning, S. R. *et al.* Ancestry-specific recent effective population size in the Americas.

PLoS Genet. **14**, 1–22 (2018).

20. Moreno-Estrada, A. *et al.* Reconstructing the Population Genetic History of the Caribbean.

PLoS Genet. **9**, (2013).

21. Novembre, J. *et al.* Genes mirror geography within Europe. *Nature* **456**, 98–101 (2008).

22. Alexander, D. H., Novembre, J. & Lange, K. Fast Model-Based Estimation of Ancestry in

Unrelated Individuals. *Genome Res.* **19**, 1655–1664 (2009).

23. 1000 Genomes Project Consortium *et al.* A global reference for human genetic variation.

Nature **526**, 68–74 (2015).

24. Li, J. Z. *et al.* Worldwide human relationships inferred from genome-wide patterns of

variation. *Science* **319**, 1100–1104 (2008).

25. Wojcik, G. L. *et al.* Genetic analyses of diverse populations improves discovery for complex

traits. *Nature* **570**, 514–518 (2019).

26. Patin, E. *et al.* Dispersals and genetic adaptation of Bantu-speaking populations in Africa

and North America. *Science* **356**, 543–546 (2017).

27. Browning, S. R. & Browning, B. L. Accurate Non-parametric Estimation of Recent Effective

Population Size from Segments of Identity by Descent. *Am. J. Hum. Genet.* **97**, 404–418

(2015).

28. Diehl, R. A. *The Olmecs : America's first civilization*. (London : Thames & Hudson, 2004).

29. Marcus, J. & Flannery, K. Cultural Evolution in Oaxaca: the Origins of the Zapotec and

Mixtec Civilizations. in *The Cambridge History of the Native Peoples of the Americas* 358–

406 (Cambridge University Press, 2000). doi:10.1017/CHOL9780521351652.009.

30. Wang, R. J., Al-Saffar, S. I., Rogers, J. & Hahn, M. W. Human generation times across the

past 250,000 years. (2021).

31. Ceballos, F. C., Joshi, P. K., Clark, D. W., Ramsay, M. & Wilson, J. F. Runs of

homozygosity: windows into population history and trait architecture. *Nat. Rev. Genet.* **19**,

220–234 (2018).

32. Clark, D. W. *et al.* Associations of autozygosity with a broad range of human phenotypes.

Nat. Commun. **10**, 4957 (2019).

33. Ringbauer, H., Novembre, J. & Steinrücken, M. Parental relatedness through time revealed

by runs of homozygosity in ancient DNA. *Nat. Commun.* **12**, 5425 (2021).

34. Henn, B. M. *et al.* Distance from sub-Saharan Africa predicts mutational load in diverse

human genomes. *Proceedings of the National Academy of Sciences* 201510805 (2015)

doi:10.1073/pnas.1510805112.

35. Henn, B. M., Botigué, L. R., Bustamante, C. D., Clark, A. G. & Gravel, S. Estimating

Mutation Load in Human Genomes. *Nat. Rev. Genet.* **16**, 333–343 (2015).

36. Simons, Y. B., Turchin, M. C., Pritchard, J. K. & Sella, G. The deleterious mutation load is

insensitive to recent population history. *Nat. Genet.* **46**, 220–224 (2014).

37. Do, R. *et al.* No evidence that selection has been less effective at removing deleterious

mutations in Europeans than in Africans. *Nat. Genet.* **47**, 126–131 (2015).

38. Acuña-Alonzo, V. *et al.* A functional ABCA1 gene variant is associated with low HDL-
cholesterol levels and shows evidence of positive selection in Native Americans. *Hum. Mol.*
Genet. **19**, 2877–2885 (2010).

563
Figure 1. Visualizing genetic structure across the geography of the Mexican Biobank as inferred
by dimensionality reduction and unsupervised clustering. A) Mexico regionalized into
Mesoamerican regions according to anthropological and archaeological context. B) Principal
components analysis (PCA) of MXB with reference global data from the 1000G project, HGDP
and PAGE. Some specific sampled cohorts are labelled across the plot to orient the reader. C)
PCA with only MXB colored and regionalized into Mesoamerican regions. D) Unsupervised
clustering using Admixture and global reference panels (same as in B). E) UMAP analysis of
MXB colored by Mesoamerican regions.
572

573
Figure 2. Effective population size (N_e) changes inferred using identity-by-descent (IBD) tracts
across ancestries and geographies reveal the different histories present within Mexico. A)
Mesoamerican chronology coloring different periods in Mesoamerican history using
anthropological and archaeological context B) Ancestry-specific effective population size
changes over past 200 generations across Mexico (colored by chronology from A assuming 30
years per generation (see Figs. S23-27 for other generation intervals and ancestries). C)
Ancestry-specific effective population changes over time for ancestries from Central America in
different Mesoamerican regions of Mexico.
582

Figure 3. Analysis of runs of homozygosity (ROH) in each individual in MXB across ancestries, geographies and birth year reveals the role of ancient and recent demographic movements to and within Mexico. A) ROH are correlated with ancestries from global region in each individual reflecting the impact of varied and shared demographic histories and bottlenecks. B) Distribution of ROH segments of different sizes for each Northern, Central, Southern and Southeastern state. The y-axis was truncated to aid visualization, truncating the first bar for some states. C) ROH as a function of birth year. Solid lines show ROH overall, and dashed lines indicate ROH overlapping ancestries from Central America. ROH are divided into small, medium and large ROH same as in (B).

Figure 4. Mutation burden in different ancestries show effects of bottleneck in causing loss of
rare variants. Rare variants are correlated with levels of ancestries from Central America,
Western Europe or West Africa for rare variants (DAF < 5%), and common variants. Analysis of
WGS from 1000 genomes MXL shows that the rare mutation burden result is robust while the
full mutation burden correlation is caused by ascertainment bias of the MEGA array (Fig. S50).
Variants were annotated using VEP, and deleterious variants are a combined set of missense
variants predicted to be damaging by polyphen2 along with splice, stoploss and stopgain
variants.

Figure 5. Height variation over space and birth year, and analysis of the factors influencing
height and other complex trait variation. A) Map of average height in Mexico (inset shows
boxplots of height variation in each state from North-west to South-east). B) Explanatory model
for height variation implicates the role of genetics and environment. The plot shows effect size
estimates and confidence intervals from a mixed model analysis. All quantitative predictors are
centered and scaled by 2 standard deviations. Asterisks indicate significance of the effect of a
predictor after Bonferroni correction (** $P < 10^{-5}$, *** $P < 10^{-6}$) across traits and predictors
analyzed. C) Height as a function of birth year in quantiles of ancestries from Central America.
D) Trait profiles for BMI, Triglycerides, and Glucose. Results of mixed model analysis same as
in (B). Educational attainment is on a scale from 0-8 (low to high educational attainment), and
altitude is measured in meters (low to high).