Unraveling the relationship between cancer and life history traits in vertebrates

Short title: **Cancer prevalence across vertebrates**

Stephanie E. Bulls[1,2], Laura Platner[3,4], Wania Ayub[1], Nickolas Moreno[1], Jean-Pierre Arditi[5], Saskia Dreyer[3],

Stephanie McCain[6], Philipp Wagner[4], Silvia Burgstaller[7], Leyla R. Davis[7], Linda GR. Bruins - van

Sonsbeek[8], Dominik Fischer[3,9], Vincent J. Lynch[10], Julien Claude[11,12], Scott Glaberman[2,13], Ylenia

Chiari[1,14*]

[1] Department of Biology, George Mason University, Fairfax, VA, USA

[2] Department of Environmental Science & Policy, George Mason University, Fairfax, VA, USA

[3] Der Grüner Zoo, Wuppertal, Germany

[4] Allwetterzoo, Münster, Germany

[5] Department of Biology, University of South Alabama, Mobile, AL, USA

[6] Birmingham Zoo, Birmingham, AL, USA

[7] Zoo Zurich, Zürich, Switzerland

[8] Diergaarde Blijdorp, Rotterdam Zoo, Rotterdam, The Netherlands

[9] Wildlife Zoo, Hellental, Germany

[10] Department of Biological Sciences, University at Buffalo, SUNY, USA

[11] Institut des Sciences de l'Evolution, Université de Montpellier/CNRS/IRD, Montpellier, France [12] Faculty

of Science, Chulalongkorn University, Bangkok, Thailand

[13] Centre for Environmental Research and Justice, School of Biosciences, University of Birmingham, UK

[14] School of Life Sciences, University of Nottingham, Nottingham, UK

*Author(s) for correspondence:

Ylenia Chiari

School of Life Sciences

University of Nottingham

Nottingham NG7 2TQ, UK

Email: ylenia.chiari@nottingham.ac.uk

ABSTRACT

Identifying species with unusually low cancer prevalence can provide new insights into cancer resistance.

Most studies have focused on mammals, but the genetic, physiological, and ecological diversity among

vertebrates can influence cancer susceptibility. We used necropsies from over a thousand species of

amphibians, birds, crocodilians, mammals, squamates, and turtles to investigate relationships between

cancer prevalence, intrinsic cancer risk, body mass, and lifespan. Previous studies often relied on species

averages, leading to inaccurate interpretations. Our innovative statistical approach uses raw cancer data and

resampling to improve accuracy. We found remarkably low cancer prevalence in turtles, high prevalence

in squamates and mammals, and lower-than-expected prevalence based on lifespan and body mass in

multiple groups. Our results show lifespan influences neoplasia and malignancy transformation rates in

mammals, while body mass affects neoplasia prevalence in amphibians and squamates. These data reveal

a complex relationship between life history traits and cancer risk, identifying vertebrates with potential

novel cancer resistance mechanisms.

KEYWORDS: Body mass, cancer prevalence, comparative oncology, intrinsic cancer risk, lifespan,

neoplasia, phylogenetic comparative method, Peto's paradox, vertebrates, zoo.

STATEMENT OF SIGNIFICANCE: Biodiversity is an untapped natural resource for understanding

cancer. Our study reveals a wide divergence in cancer prevalence among vertebrate groups, with notably

low rates in turtles and high rates in squamates and mammals. These findings can lead to new breakthroughs

in understanding the biology of cancer.

INTRODUCTION

Understanding cancer prevalence across animal species can identify novel cancer resistance mechanisms. Theoretically, larger, longer-lived species should have higher cancer susceptibility due to increased cell numbers and more extensive lifetime cellular turnover [1]. However, empirical studies have not observed this pattern in mammals [2,3,4] or other vertebrates [5,6]. This suggests that species with extreme body sizes and lifespans must have evolved mechanisms to offset their elevated cancer risk [1,7,8]. Our study seeks to bridge this gap by investigating cancer prevalence within and among vertebrate groups, including underrepresented ones such as reptiles and amphibians, to identify species with novel anti-cancer mechanisms [9].

Prior research has documented marked differences in cancer prevalence among vertebrates, with higher rates in mammals compared to other groups [10, 11, 12,13,14]. But studies have largely overlooked diversity within certain groups, especially reptiles and amphibians. There haven't been any phylogenetic comparative studies of cancer that discriminate among major lineages of non-avian reptiles including crocodilians, squamates (lizards and snakes), and turtles. Previous data on cancer occurrence hint at low cancer prevalence in turtles and possibly crocodilians [15,16], underscoring the need for more in-depth analysis in these groups.

In this study, we assess cancer prevalence across all major vertebrate groups except fish—including amphibians, birds, crocodilians, mammals, squamates, and turtles—using zoo necropsy data. Our goal is to uncover the relationship between cancer prevalence and species-specific traits such as body mass, lifespan, and intrinsic cancer risk. We introduce an innovative statistical approach to overcome the significant limitations of previous research, employing a phylogenetic comparative method specifically designed for binomial distributions [17]. This is crucial because the rarity of cancer within species results in prevalence data that are heavily skewed towards zero, fundamentally violating normality assumptions that many statistical models, including phylogenetic generalized least squares (PGLS) regression, depend on. Indeed, PGLS regression—commonly used in past studies [3,6,] (see [10] for analysis of data using modified PGLS and the approach presented here) —relies on species averages, making it less appropriate for handling non-

normal data characteristic of cancer prevalence.

In contrast, our innovative method leverages individual data points, offering a more precise account

of intraspecific sampling effort and avoiding the common issue of treating prevalence as a continuous

variable. This strategy establishes stronger correlations between cancer incidence and key biological traits

across various vertebrate groups. Additionally, we incorporate a jackknife resampling technique to evaluate

the robustness of our findings, particularly the impact of species sampling effort. This approach not only

addresses the shortcomings of previous methodologies, but also provides a more reliable foundation for

future research into cancer prevalence across vertebrates. Finally, we contextualize our findings within

existing research by reanalyzing publicly available mammalian data using our improved statistical

methodology [3,4] (see also [10] for a similar approach in comparing results obtained following the innovative

approach used in this paper and a modified PGLS method).

Our analysis shows that neoplasia and cancer prevalence exhibit significant variation within and

among vertebrate groups. Unlike previous studies [2,3,4,5,6], our results indicate that body mass and lifespan

are influential factors for cancer prevalence in amphibians, squamates, and mammals. We further show that

cancer prevalence is extremely low in turtles. This highlights the importance of employing precise statistical

methods to reveal complex biological relationships and provides new perspectives on cancer biology across

different vertebrate lineages.

RESULTS

Necropsy Data

Our dataset included 9,631 necropsy reports from 1,030 species (1,060 species, including additional

squamate data from [18]). Out of the total described species in each vertebrate group, our sampling represents

0.7% of all amphibians, 4.5% of all birds, 34.7% of all crocodilians, 4.0% of all mammals, 1.8% of all

squamates, and 21.0% of all turtles (**Table 1**). Birds were highly represented with 507 species and 5,121

necropsy reports, while turtles and crocodilians had fewer necropsies (207 and 26, respectively). The

Shannon Equitability Index indicated that the allocation of necropsies among species was similar within

each group (**Supplementary Materials Table 2**).

Table 1. Total number of necropsies and species for each tetrapod group. The total recognized number

of species for each group was obtained from IUCN (*The IUCN Red List of Threatened Species*, [19]). "Total

neoplasia" includes all benign and malignant tumor counts whereas "Total cancer" includes only those

tumors that were diagnosed as malignant by a veterinary pathologist and confirmed by histology.

Malignancy transformation rates is indicated as "malignancy prevalence" and was calculated as neoplasia

count divided by cancer count.

	Amphibians	Birds	Crocodilians	Mammals	Squamates	Squamates*	Turtles
Species in dataset	49	507	8	230	179	209	57
Recognized species in group	7296	11162	23	5968	9855	9855	269
Percentage of recognized species represented	0.7	4.5	34.7	4.0	1.8	2.1	21.0
Total necropsies	383	5121	26	2841	1053	2608	207
Total neoplasia	15	96	2	191	84	198	1
Total cancer	10	61	1	111	57	164	0
Neoplasia prevalence	0.04	0.02	0.08	0.07	0.08	0.08	0.005
Cancer prevalence	0.03	0.01	0.04	0.04	0.05	0.06	0
Malignancy prevalence	0.7	0.6	0.5	0.6	0.7	0.8	0

114
*Includes data from [18]

Prevalence of Neoplasia, Cancer, and Malignancy Transformation Rate

We found clear differences among tetrapod groups (**Figures 1, 2,** and **3**), with some groups having

notably low neoplasia, cancer prevalence, and/or malignancy transformation rates. Turtles had no instances

of cancer and only one case of neoplasia. Crocodilians had only two neoplasms, one of which was malignant

(**Table 1**). Neoplasia and cancer prevalence values were also particularly low in birds (2% and 1%,

respectively), while the malignancy transformation rate was 60% (**Table 1**). These results indicate that

while cancer is relatively rare in birds, it is frequently malignant when it does occur.

Amphibians, birds, mammals, and squamates had outlier species with prevalence values (neoplasia,

cancer, or malignancy rate) significantly different ($p < 0.05$) from the group mean (**Supplementary**

Materials Table 3). Notably, among the species that were different from the group mean and a sample size

of at least 20 necropsies per species, some outliers for neoplasia prevalence were the domestic chicken

(7/157, 5.6%), the common hamster (24/114, 22%), and the diamondback rattlesnake (6/21, 29%).

Additionally, three mammal outliers had 0% neoplasia prevalence despite a large number of necropsies per

species: the artiodactyl collard peccary (0/89), domestic goat (0/91), and the rodent Patagonian mara

(0/103).

Within our dataset, neoplasia prevalence ranked from highest to lowest as follows: 8% in both

squamates and crocodilians, 7% in mammals, 4% in amphibians, 2% in birds, and 0.5% in turtles (**Table 1**

and **Figure 2**). Cancer prevalence ranked from highest to lowest as follows: 5% in squamates, 4% in both

mammals and crocodilians, 3% in amphibians, 1% in birds, and 0% in turtles. Malignancy transformation

rate (i.e., cancer divided by neoplasia) was highest in squamates and amphibians with both groups having

70% malignancy transformation rate (80% in squamates when including data from [18]), followed by

mammals and birds with 60%, and crocodilians with 50% (**Table 1** and **Figure 2**). Squamates continue to

have the highest neoplasia prevalence and malignancy transformation rate at ≥10 and ≥20 necropsies per

species (**Supplementary Materials Table 4**). The observed prevalence values and their 95% confidence

intervals among tetrapod groups indicated that birds and turtles had much lower neoplasia and cancer

prevalence than other groups, but birds were similar to other groups for malignancy transformation rate

(**Figure 2**). Except for crocodilian and amphibian cancer and malignancy transformation rates, 95%

confidence intervals for prevalence values were narrow (**Figure 2**), indicating that estimates were precise.

Pairwise comparisons of neoplasia, cancer, and malignancy transformation rate calculated using

prop.test and *p.adjust* with the BH method to account for multiple testing further supported differences

among tetrapod groups (**Table 2**). Birds and turtles, with the lowest neoplasia and cancer prevalence, were

not significantly different from each other but differed from all other groups (except crocodilians, **Table**

2). For neoplasia, amphibians differed from squamates, which had the highest neoplasia prevalence.

Amphibians and squamates also differed from each other for cancer prevalence, but not for malignancy

transformation rate (**Table 2**). Finally, mammals and squamates did not differ from each other in terms of

neoplasia prevalence, but they were found to be different for cancer and malignancy transformation rate

(**Table 2**). These trends were generally supported when the data was curated to only species with ≥10

necropsies (**Supplementary Materials Tables 5**). Birds differed from mammals and squamates for

neoplasia and cancer for ≥20 necropsies, while only squamates differed from the other two groups for

malignancy transformation rate (**Supplementary Materials Tables 6**).

Within each studied vertebrate group, we found wide variation in neoplasia and cancer prevalence

among families and orders (**Figure 3, Supplementary Materials Table 7,** full comparisons on Dryad *after*

manuscript acceptance).

Figure 1. Summary of (a) neoplasia prevalence and (b) cancer prevalence across tetrapod groups. Each

point represents the prevalence of a single species. Pink lines represent the mean prevalence of each group.

Groups ordered left to right by descending mean prevalence. Prevalence is based on data as in Table 1.

*Indicates squamate data from [18] included.

(a)

(b)

Figure 2. Prevalence of neoplasia, cancer, and malignancy transformation rate across tetrapod groups using

the full dataset as in **Table 1**. Malignancy transformation rates is indicated as "malignancy prevalence" and

was calculated as neoplasia count divided by cancer count. Points represent observed prevalence values and

lines represent the 95% confidence intervals surrounding the observed prevalence value. *Indicates

squamate dataset with data from [18] included. Please notice that due to the large variation in percentage

among neoplasia, cancer, and malignancy transformation rate, the scale of the y-axis is different in the three

graphs.

Figure 3. Interspecific variation in neoplasia (N) and cancer (C) prevalence within a) mammals including

all species, b) mammals including only species with 20 or more necropsies, c) birds including all species,

d) birds including only species with 20 or more necropsies, e) squamates, including all species, f) squamates

including only species with 20 or more necropsies, and g) amphibians including all species. Taxa are color

coded either by order (mammals and birds) or family (squamates and amphibians).

Table 2: Pairwise comparison results for (a) neoplasia, (b) cancer, and (c) malignancy prevalence proportions run on full dataset (as in **Table 1) using *prop.test* a**nd ***p.adjust*** with the Benjamini & Hochberg ("BH") method (R package {stats}). Numbers in parentheses are proportion numbers per group calculated as # group neoplasia / # group total necropsy or # group cancer/ # group necropsy total, and # group cancer / # group total neoplasia for malignancy. Malignancy transformation rates is indicated as "malignancy prevalence" and was calculated as neoplasia count divided by cancer count. Significant values are in bold. Malignancy transformation rate could not be calculated for turtles since there were no instances of cancer in these groups. The asterisk denotes squamates data included from [18].

(a)

GROUP NEOPLASIA PREVALENCE							
	Amphibians (15/383)	Birds (96/5124)	Crocodilians (2/25)	Mammals (191/2841)	Squamates (84/1053)	Squamates* (198/2608)	Turtles (1/207)
Turtles (1/207)	0.07	0.3	0.07	**2.70E-03**	**1.50E-15**	**1.10E-03**	1
Squamates* (198/2608)	**0.03**	**1.50E-15**	1	0.3	1	1	
Squamates (84/1053)	**0.03**	**1.50E-15**	1	0.3	1		
Mammals (191/2841)	0.1	**1.50E-15**	1	1			
Crocodilians (2/25)	1	0.2	1				
Birds (96/5124)	**0.03**	1					

(b)

GROUP CANCER PREVALENCE							
	Amphibians (10/383)	Birds (61/5124)	Crocodilians (1/25)	Mammals (111/2841)	Squamates (57/1053)	Squamates* (164/2608)	Turtles (0/207)

Turtles (0/207)	**1.50E-15**	0.4	0.4	**0.02**	**0.004**	**0.002**	1
Squamates* (164/2608)	**0.02**	**1.50E-15**	1	**4.0E-4**	0.7	1	
Squamates (57/1053)	0.1	**1.50E-15**	1	0.1	1		
Mammals (111/2841)	0.5	**1.50E-15**	1	1			
Crocodilians (1/25)	1	1	1				
Birds (61/5124)	0.08	1					

199
200
201 (c)

GROUP MALIGNANCY PREVALENCE						
	Amphibians (10/15)	Birds (55/82)	Crocodilians (1/2)	Mammals (106/185)	Squamates (50/75)	Squamates* (140/173)
Squamates* (140/173)	0.8	**0.004**	1	**3.20E-06**	**0.05**	1
Squamates (50/75)	1	1	1	0.8	1	
Mammals (106/185)	1	1	1	1		
Crocodilians (1/2)	1	1	1			
Birds (55/82)	1	1				

202
203
204

Intrinsic Cancer Risk Versus Neoplasia, Cancer, and Malignancy Transformation Rate

Intrinsic cancer risk corresponds to the predicted risk of developing cancer based on a combination

of species lifespan and body mass [20,21]. Across tetrapods, we found ICR to be highly variable (**Figure 4**).

Certain species showed extraordinarily high ICR values compared to other species in their groups, including

Chinese giant salamanders (*Andrias davidianus*), common ostriches (*Struthio camelus*), Asian elephants

(*Elaphas maximus*), Komodo dragons (*Varanus komodoensis*), and Galapagos tortoises (*Geochelone nigra*

complex). Overall, based on the species in our dataset, amphibians and turtles had lower average ICR values

than other tetrapod groups (**Figure 4**). Note that the ICR calculation (ICR = lifespan^6 x body mass) is more

strongly affected by variation in lifespan than body mass.

Using phylogenetic comparative methods (*gee*), we found significant relationships between ICR

and neoplasia prevalence in squamates (positive, p=0.007) and mammals (negative, p=0.0002), and

between ICR and cancer prevalence in amphibians (positive, p=0.002; **Table 3**). ICR was a highly

correlated to neoplasia or cancer prevalence for squamates, mammals, and amphibians, but not for other

groups, and was not related to malignancy transformation rates in any group.

Table 3. Influence of intrinsic cancer risk (lifespan^6 x body mass) on neoplasia, cancer or malignancy

transformation rate. Malignancy transformation rate is indicated as "Malignancy" in the table. Analyses

were run using the generalized estimating equations (*gee*; see Materials and Methods for additional

information). Species included in analysis were curated to those for which both lifespan and body mass life

history traits were available. P-values <0.05 are in bold with slope coefficients in parentheses to show

direction effect and jackknife 95% confidence intervals italicized in brackets. Dashed lines represent

models that did not run due to non-converging. NA in turtles indicates that analyses could not be run

because cancer was not observed in these groups.

Vertebrate Group	N Species	Neoplasia	Cancer	Malignancy
		ICR p-value (slope) [*jackknife*]	ICR p-value (slope) [*jackknife*]	ICR p-value (slope) [*jackknife*]
Amphibians	40	0.2 (0.2) [*0.09 , 0.2*]	**0.002** **(0.3)** [***0.2 , 0.4***]	–
Birds	263	0.2 (0.05) [*0.04 , 0.08*]	0.7 (0.02) [*0.009 , 0.02*]	0.4 (-0.06 , [*0.05 , 0.05*]
Mammals	218	**2.0e-4** **(-0.1)** [***-0.1 , -0.07***]	0.45 (-0.02) [*-0.03 , -0.02*]	0.2 (0.04) [*-0.1 , -0.1*]
Squamates	99	**0.007** **(0.1)** [***0.08 , 0.1***]	0.12 0.07 [*0.06 , 0.1*]	0.05 (-0.2) [*0.1 , 0.1*]
Squamates *	113	**3.0e-7** **(0.2)** [***0.17 , 0.2***]	**8.0e-8** **(0.25)** [***0.2 , 0.3***]	0.4 (0.06) [*0.2 , 0.2*]
Turtles	38	0.9 (-0.02) [*-0.03 , -0.02*]	NA	NA

Figure 4. Relative estimated intrinsic cancer rates (ICR) based on body mass and lifespan data for all tetrapod species examined in this study (see Materials and Methods for ICR calculation). Gray lines in each violin represent the mean ICR for the tetrapod group. For visual representation, original ICR values were log scaled and transformed using a min-max normalization (ICR – min(ICR))/(max(ICR) – min(ICR)) to bring them to a 0-1 range within each group, where 0 and 1 represent the species with the lowest and highest intrinsic cancer risk, respectively, within each lineage. Please note that species with 1 in the graph do not necessarily have a 100% intrinsic cancer rate, but they just correspond to the higher intrinsic rate in the group.

Neoplasia, Cancer, and Malignancy Transformation Rate as a Function of Body Mass and Lifespan

We also tested the influence of body mass or lifespan separately on the prevalence of neoplasia, cancer, and malignancy transformation rates in each tetrapod group, while accounting for phylogenetic relationships among species within each group. We found that body mass was positively correlated with neoplasia and cancer in amphibians, neoplasia in squamates, and negatively correlated to neoplasia and malignancy transformation rate in mammals (**Table 4**). Lifespan was positively correlated with cancer in amphibians, neoplasia and cancer in squamates, and malignancy transformation rates in mammals, and

negatively correlated with neoplasia in mammals (**Table 4**). Thus, in amphibians and no other studied

group, both body mass and lifespan influence cancer, while in mammals and no other group, both body

mass and lifespan influence neoplasia and malignancy transformation rate (**Table 4**). A positive correlation

means that when taking into account phylogenetic relationships among the studied species, an increase in

that life history trait correspond to higher prevalence, which supports our predictions using ICR (**Table 3**).

On the other hand, in mammals, we found a negative relationship between body mass and lifespan and

neoplasia, as well as body mass and malignancy transformation rates, meaning that increases in that life

history trait correspond to a decrease in prevalence (**Table 4**). For groups with ≥20 species and ≥10 or 20

necropsies per species (birds, mammals, and squamates), lifespan influenced neoplasia and malignancy

transformation rate in mammals, and neoplasia and cancer in squamates (**Supplementary Materials Table**

8). To note that while a threshold of 10 or 20 or more necropsies per species has been used in previous

studies, due to the reduced species representation and prevalence distribution on each of the group

phylogenies (**Figure 3**), phylogenetic heterogeneity in the data is lost and analyses fail to converge due to

lack of variation in the data across the reduced phylogenetic representation. The fact that less tests reach

convergence or are significant when species with few individuals are dropped from the analyses it is further

illustrated by the jackknife confidence intervals. 95% confidence intervals in fact increase when the number

of species included in the analysis is reduced (see confidence intervals of **Table 4 and Supplementary**

Materials Table 8) The meaning of this is a reduced confidence in the results obtained using less species

with more necropsies than the entire dataset.

Table 4. Influence of body mass and lifespan on cancer, neoplasia, and malignancy prevalence.

Analyses run using the full dataset on cancer, neoplasia, and malignancy transformation rates using either

lifespan or body mass as the predicting variable and taking into account group's phylogenetic relationships.

Analyses were run using the generalized estimating equations (*gee*; see Materials and Methods for

additional information). Malignancy transformation rates is indicated as "malignancy prevalence" and was

calculated as neoplasia count divided by cancer count. P-values <0.05 are in bold with slope parameters in

parentheses to show direction effect and jackknife 95% confidence intervals italicized in brackets. Lines

represent models that did not run due to non-converging (see Materials and Methods for additional

information). Numbers in the column "N Species" refers to the number of species per trait being analyzed.

If a species mass or lifespan history trait was unknown it was removed from that analysis. Crocodilians

could not be analyzed due to an insufficient number of species. Turtles could not be analyzed for cancer

and malignancy transformation rate as no instances of cancer were found.

Vertebrate Group	Total Neoplasia				Cancer				Malignancy			
	Mass		Lifespan		Mass		Lifespan		Mass		Lifespan	
	N Species	p-value (slope) [jack knife]	N Species	p-value (slope) [jack knife]	N Species	p-value (slope) [jack knife]	N Species	p-value (slope) [jack knife]	N Species	p-value (slope) [jack knife]	N Species	p-value (slope) [jack knife]
Amphibians	47	**0.003 (1.6)** *[1.3 , 2.1]*	42	0.31 (0.7) *[0.2 , 1.5]*	47	**0.0014 (1.5)** *[1.3 , 2.3]*	42	**0.009 (1.8)** *[1.1 , 2.2]*	8	0.85 (0.3) *[0.12 , 1.4]*	8	—
Birds	299	0.4 (0.07) *[0.07 , 0.1]*	267	0.3 (0.3) *[0.3 , 0.4]*	299	0.6 (0.07) *[0.06 , 0.08]*	267	0.96 (-0.02) *[0.03 , 0.08]*	57	0.12 (0.3) *[0.2 , 0.3]*	48	0.3 (-0.5) *[-0.7 , -0.4]*
Mammals	220	**0.002 (-0.2)** *[-0.2 , -0.15]*	218	**0.001 (-0.8)** *[-0.8 , -0.4]*	220	0.6 (0.03) *[-9e-4 , 0.05]*	218	0.3 (-0.3) *[-0.4 , -0.3]*	77	**0.005 (-0.2)** *[-0.3 , -0.2]*	77	**0.001 (1.1)** *[1.0 , 1.1]*
Squamates	102	**0.02 (0.2)** *[0.16 , 0.2]*	122	0.2 (0.4) *[0.3 , 0.6]*	102	0.14 (0.1) *[0.09 , 0.2]*	122	0.8 (0.08) *[-0.02 , 0.16]*	32	0.07 (-0.4) *[0.4 , -0.2]*	32	0.2 (-0.9) *[-1.1 , -0.7]*
Squamates *	116	**0.0003 (0.24)** *[0.2 , 0.3]*	142	**0.0003 (0.9)** *[0.8 , 1.1]*	116	**0.0004 (0.3)** *[0.2 , 0.3]*	142	**0.0004 (1.07)** *[1.0 , 1.2]*	56	0.6 (0.1) *[0.05 , 0.2]*	56	0.5 (0.4) *[0.07 , 0.6]*

Vertebrate Group	Total Neoplasia				Cancer				Malignancy			
	Mass		Lifespan		Mass		Lifespan		Mass		Lifespan	
	N Species	p-value (slope) *[jack knife]*	N Species	p-value (slope) *[jack knife]*	N Species	p-value (slope) *[jack knife]*	N Species	p-value (slope) *[jack knife]*	N Species	p-value (slope) *[jack knife]*	N Species	p-value (slope) *[jack knife]*
Turtles	46	0.8 (0.07) *[0.05 , 0.07]*	42	0.8 (-0.3) *[-0.6 , 0.13]*	NA	NA	NA	NA	NA	NA	NA	NA

288
289   *Includes data from [18]

Re-analysis of Publicly Available Data from Previous Studies on Mammals

Previous studies in mammals [3,4] have found that change of body mass and lifespan is not associated with a change of neoplasia and cancer prevalence (but see [10] on the influence of body weight after correcting for gestational period). To assess the influence of our statistical approach on the results obtained from prior studies, we re-analyzed their data using *gee*. Based on the data from Boddy et al. [3], which included 36 species (28 of them with ≥10 necropsies per species), we found no influence of body mass or lifespan change on neoplasia or cancer prevalence evolution, supporting their results (**Supplementary Materials Table 9**). However, as their interspecific sampling is lower than ours and as parameter estimates were close to 0 (**Supplementary Materials Table 9**, see jackknife values), it could also mean that their sample size is too low to detect any relationships. However, when we used the data from Vincze et al. [4], including 189 species with 20 or more necropsies per species, our statistical approach indicated that both changes in body mass and life expectancy were significantly (p<0.001) related to changes in neoplasia prevalence, in contrast to what was found by the authors. Note that in Vincze et al. [4], the neoplasm was defined as the cause of death, so in this sense it is cancer mortality data and does not exactly correspond to any of the categories – neoplasia, cancer, or malignancy transformation rate – used in our study. The fact that we

found a significant relationship between life history traits and cancer mortality data when using their dataset

supports the relationship in mammals between cancer mortality and life history traits (**Supplementary**

Materials Table 9).

DISCUSSION

We estimated neoplasia and cancer prevalence and malignancy transformation rates across

vertebrates, including amphibians, birds, crocodilians, mammals, squamates, and turtles, focusing on

variation both within and among groups. Previous studies have explored cancer prevalence in birds and

reptiles as a single taxon (sauropsids) [5,10] or focusing on birds alone [6,13], which limits understanding of

cancer diversity among these groups. Our study addresses this gap by examining each vertebrate group

individually. We found that neoplasia and cancer prevalence, as well as malignancy transformation rates,

were higher in squamates, especially in large-bodied lizards like Helodermatidae (e.g., gila monster). This

could be due to the greater number of cells associated with larger body sizes, increasing cancer risk. We

also found high malignancy transformation rates almost uniformly across snake families (67-100%). In

contrast, we found no cancer and only one case of neoplasia (*Chelus fimbriata*) in turtles. In crocodilians,

only two out of eight species had neoplasia, one of which was malignant (*Osteolaemus tetraspis*).

Despite the remarkably low prevalence values, neoplasia and cancer do occur in turtles and

crocodilians [12,15,16,22,23]. These low rates may be due to slower metabolic rates, reduced oxidative stress, and

other unique physiological traits. Previous work, which did not distinguish between benign and malignant

tumors, found 1.2-2.7% neoplasia prevalence in turtles [15,16] and 2.2% in crocodilians [15]. Turtles have lower

germline mutation rates in mitochondrial and nuclear DNA than mammals and birds [24], possibly related to

lower somatic mutation rates and lower cancer incidence. In addition, turtles were found to have slower

aging rates and derived cellular phenotypes associated with delayed aging and cancer resistance [25,26,27].

We identified low rates of neoplasia and cancer in birds compared to mammals and reptiles,

consistent with previous studies [11,12]. This low prevalence may be related to the bursa of Fabricius, a

specialized organ producing B-cells in juvenile birds, and enhanced immune system development[13]. Higher

prevalence of neoplasia and malignancy transformation rate in some avian species has been correlated with

larger clutch sizes [6]. Our data confirmed that birds have very low neoplasia rates, but malignancy

transformation rates are comparable to mammals and squamates. This might be due to the efficiency of

their immune system in preventing neoplasia, but once tumors develop, they rapidly progress to

malignancy. Although this effect could be due to bias in the types of tissues sent for pathology analysis, if

confirmed, these results indicate that when neoplasia occurs in birds, it is usually malignant (>50% of the

time).

Previous studies have found conflicting results regarding the influence of life history traits, such as

body mass and lifespan, on neoplasia and cancer prevalence across tetrapod groups. Our work is the first

to compare expected rates of intrinsic cancer risk (ICR) based on body mass and lifespan with observed

rates across each studied vertebrate group. We found higher ICR to be correlated with higher neoplasia

prevalence in squamates. Conversely, in mammals, higher ICR values correlate with lower neoplasia

prevalence, possibly due to evolved tumor suppression mechanisms in larger, longer-lived species. ICR is

also a good positive predictor of cancer in amphibians. However, ICR does not correlates to neoplasia or

cancer in birds, turtles, or crocodilians, nor malignancy transformation rates across any studied vertebrate

groups. These results highlight the complexity of cancer evolution across different vertebrate groups and

underscores the need for taxon-specific approaches in future comparative oncology research.

When analyzing the influence of body mass and lifespan separately, at least one of these two life

history traits has an influence on neoplasia or cancer in several vertebrate groups including amphibians,

squamates, and mammals. Our results contradict a previous study that observed a positive association

between body mass and neoplasia prevalence in birds [13]. In mammals, change in cancer prevalence was

previously found to be unrelated to body mass or lifespan evolution (but see [10]), but instead potentially

influenced by diet, with higher prevalence values for cancer observed in carnivores [4,12] or by gestation

period and litter size [3,10]. In contrast, we found a positive relationship between lifespan and malignancy

transformation rate in mammals, but inverse relationships between body mass and both neoplasia and

malignancy transformation rates as well as an inverse relationship between lifespan and neoplasia. This

suggests that animals with larger body mass have lower neoplasia and malignancy transformation rates, and

as such, larger animals have effectively evolved mechanisms to reduce mortality due to cancer by

experiencing a decreased occurrence of neoplasia than except for their body size. At the same time, our

results indicate a more complex relationship existing between lifespan, neoplasia, and malignancy

transformation. Longer-lived species may have evolved mechanisms protecting them from developing

neoplasia, but once it occurs, it may often become malignant, hence the positive relationship with

malignancy transformation.

In this work, we have established the novel findings that squamates and amphibians both have a

positive relationship between body mass and neoplasia prevalence, and amphibians also have a positive

relationship between body mass, lifespan, and cancer. It has been proposed that regenerative capacities, as

observed in some amphibian species, may be associated with tumor suppression mechanisms [28,29], and that

regeneration in amphibians, unlike mammals, does not necessarily result in increased cancer prevalence

[30,31]. This hypothesis needs further exploration since we found that amphibians had lower neoplasia and

cancer prevalence than mammals and squamates, but higher than birds. Regenerative capacities are not

uniform across species, orders, and life-stages of amphibians [32], possibly explaining why neoplasia and

cancer prevalence is relatively high in this group. Finally, we found that neoplasia, cancer, and malignancy

transformation rates are remarkably frequent in snakes and occurred in almost all snake families included

in this study, which is similar to [18].

Our study pioneers an approach for analyzing the relationship between life history traits and cancer

prevalence data, marking a significant advancement in the field. Unlike previous studies that treated cancer

prevalence as a continuous variable using Phylogenetic Generalized Least Squares (PGLS) for correlation

analysis [3,4], we recognize that prevalence data are proportional, adhering to a binomial rather than a normal

distribution [17]. The skewness towards zero in cancer occurrence across species challenges the PGLS

method's assumption of a normal distribution. We employ Generalized Estimating Equations (*gee*) to

encompass the binomial distribution and account for the skewed nature of cancer prevalence data. This

method allows for the inclusion of all collected data, improving representation even for species with sparse

samples. *Gee* utilizes individual neoplasia and cancer counts instead of averaging prevalence per species,

enabling a more detailed analysis that considers variations in sample size and distribution across the

phylogeny. Our analysis, conducted without imposing a minimum threshold for necropsies per species, and

then with subsets of species having at least 10 or 20 necropsies, underscores the limitations of data

censoring. Excluding species based on arbitrary necropsy thresholds reduces phylogenetic representation

and variation in prevalence across clades, potentially leading to non-converging analyses or insignificantly

large confidence errors. Thus, we advocate for using statistical methods suited for proportional data and

individual rather than mean species data.

Finally, to assess whether our analytical approach may explain why our findings differ from

previously published work, we reanalyzed the mammalian data from both Boddy et al. [3] and Vincze et al. [4]

using our statistical methodology. We found that when the Vincze et al.'s [4] data were reanalyzed with our

approach, their data did support a relationship between cancer mortality and life history traits. This suggests

that a relationship between life history traits and neoplasia or cancer mortality may previously not have

been recovered due to the use of less suitable statistical approaches.

Caveats And Future Directions

Our study has identified a significant influence of body mass and lifespan on neoplasia and cancer

prevalence across various tetrapod groups. The additional jackknife analysis we conducted suggests that

limitations such as sparse data, long evolutionary branches within the phylogeny of focal species, and the

uneven distribution of neoplasia, cancer, or rates of malignancy transformation across the phylogeny can

affect the outcomes of comparative phylogenetic methods. Addressing these limitations by expanding the

dataset both in terms of species diversity and the number of necropsies per species could mitigate some of

these issues. However, the acquisition of more data is challenged by the fact that certain species, especially

those in zoos like turtles, are endangered or critically endangered [33], limiting the availability of individuals

for study. Analyzing data from all available individuals ensures a more comprehensive phylogenetic

representation and strengthens the robustness of the results.

Acknowledging potential biases is crucial in studies of this nature. The frequency and type of tissue

samples submitted for pathological analysis can vary significantly across vertebrate groups, zoos, and over

time, influenced by evolving necropsy practices. A strategy to minimize such biases involves

comprehensive sampling across a wide range of groups, zoological institutions, and geographical areas.

Furthermore, the zoo environment, predominantly housing captive-bred species often at risk of inbreeding,

may influence cancer patterns differently than in wild populations. Although zoo data provide invaluable

insights for answering broad comparative questions, as demonstrated in our study, future research should

aim to address these considerations to enhance the reliability and applicability of findings. Finally, since

animals housed in zoos may have different lifespans and aging processes compared to their wild

counterparts [34], obtaining data on cancer prevalence in wild animals is essential for a complete

understanding of cancer occurrence in these organisms. However, accurately measuring cancer prevalence

in wild animals is challenging, especially for elusive species. Sick or dead animals are likely to be eaten or

difficult to find, leading to significant underestimates of cancer prevalence.

Conclusions

Our findings reveal that vertebrate groups offer distinct insights into the evolution of cancer

suppression mechanisms. In mammals, our data support Peto's Paradox, showing that evolutionary

pressures have favored the suppression of neoplasia in larger species. We also observed a positive

correlation between malignancy transformation rates and lifespan in mammals, suggesting that neoplasia,

once formed, are prone to becoming malignant. This pattern implies that evolutionary strategies may have

prioritized preventing neoplasia over cancer, guiding future research toward uncovering specific cancer

suppression mechanisms in mammals, including humans.

In amphibians and squamates, body mass and lifespan emerged as reliable predictors of neoplasia

prevalence, although this did not extend to malignancy transformation rates. In contrast, birds exhibited

relatively low rates of both neoplasia and cancer, hinting at evolved resistance mechanisms to neoplasia

irrespective of their life history traits. However, the high rate of malignancy transformation in birds suggests

that neoplasia that do develop are likely to become malignant.

Our study also highlights the urgent need for more comprehensive data on amphibians, a group

where the dual influence of lifespan and body mass on cancer prevalence was noted, yet remains

underrepresented in comparative oncology research. The limited species representation in our dataset

indicates a gap in understanding cancer across vertebrates.

Turtles, and possibly crocodilians, stand out as valuable models for cancer resistance research due

to their exceptionally low neoplasia rates and even rarer malignancy transformation. These findings

underscore the diversity of cancer prevalence and resistance mechanisms across vertebrate groups and

emphasize the importance of broadening the scope of comparative oncology to include a wider array of

species.

MATERIALS AND METHODS

Cancer Data Collection and Curation

We searched paper reports or digital databases of six European zoos (Allwetterzoo, Münster,

Germany; ZOOM Erlebniswelt Gelsenkirchen, Gelsenkirchen, Germany; Der Grüner Zoo Wuppertal,

Wuppertal, Germany; Wildlife Zoo Hellenthal, Hellenthal, Germany; Rotterdam Zoo, Rotterdam, The

Netherlands; Zoo Zürich, Zürich, Switzerland) and one in the United States (Birmingham Zoo, Alabama)

for necropsies recorded between 1998 and 2019 (**Supplementary Material Table 1**). The databases for

Allwetterzoo, ZOOM Erlebniswelt, Zoo Wuppertal, Zoo Hellenthal, Zürich Zoo, and Birmingham Zoo

were all local databases, while Rotterdam Zoo data was obtained through local paper reports and the

Species360 Zoological Information Management System (ZIMS [35]) with Rotterdam Zoo's authorization.

We also used a dataset of neoplasia (benign or malignant) in snakes from Duke et al. [18], which is based on

cancer prevalence calculated from the number of biopsies and necropsies of the total number of individuals

(live and dead) of each species housed at a given time at participating institutions in the study. Since this

prevalence was partly based on live individuals instead of strictly necropsy reports, like the rest of our data,

we performed subsequent analyses both with and without the data from Duke et al. [18].

The list of keywords used to search for neoplasia within the necropsy reports can be found in the

file "Neoplasia_Terms" (**Supplementary Materials**). To build the dataset for this study, we only included

tumor data confirmed by histological reports from veterinary pathologists. Any neoplastic individual that

was too autolytic to diagnose potential cancer incidence was removed from the study altogether. To be

conservative in our dataset, any individual that died during the first month of life was removed from the

dataset since they may have a low risk of developing cancer, which would bias the data toward lower cancer

prevalence. To note that although the data in this study all come from necropsy reports (except from [18]), the

cause of death in these reports was not necessarily recognized to be the tumor(s).

We tallied the total number of necropsies, independently of whether they had a tumor or not, for

each species represented by a necropsy report during the time frame considered in this study (1998-2019).

The data were organized into cases of 'neoplasia' (any benign or malignant neoplasia confirmed by

histology) and 'cancer' (malignant neoplasia diagnosed by abnormal cellular nuclei, tissue invasion, and/or

metastasis). Finally, two species, *Rousettus aegyptiacus* (mammal) and *Thamnophis radix* (squamate from

[18]) had more than 500 necropsies and were removed from the dataset as outliers, defined as having

necropsies more than three standard deviations above the group mean.

"Neoplasia prevalence" per group was calculated as the total number of neoplasias divided by the

total number of necropsies for each group following the rate of tumorigenesis calculation in [36]. "Cancer

prevalence" per group was calculated as the total number of cancers divided by the total number of

necropsies for each group. "Malignancy transformation rate" per group was calculated as the total number

of cancers divided by the total number of neoplasias for each group also following [36]. Finally, although

malignancy requires neoplasia to occur first [36], our data are based only on deceased individuals (i.e.,

necropsies). Thus, the number of neoplasias could be skewed if the individual did not die from it or other

causes. For that reason, we report neoplasia, cancer, and malignancy transformation rate values throughout

our results. The full dataset can be found on Dryad (*after manuscript acceptance*).

Life History Data Collection

Maximum body mass and lifespan were collected for each species when available using the Animal

Aging and Longevity Database (AnAge) [37]. If no species data was found in AnAge, primary sources were

used when available. Data origin for life history trait information is listed in the full dataset file under

"Sources LHT" for each group (*Dryad after manuscript acceptance*). If no source for a verified maximum

lifespan or body mass could be found, then that species was removed from life history trait analyses. Most

amphibian life history trait data were collected from the AmphiBIO database [38]. As body mass information

was lacking for most amphibians, snout-vent length was used as a proxy for body mass. This is the most

common measurement for amphibian body size since mass can be highly variable within the same species

and within individuals, even over a short time period, due to factors such as hydration, bladder size,

physiological state, and reproductive state [39,40].

Intrinsic Cancer Risk Analysis

Intrinsic cancer risk (ICR) due to body size and lifespan was estimated for each species in the

dataset following Peto's model [20], where ICR = $\text{lifespan}^6 \times \text{body mass}$. Only species for which both body

mass and lifespan data are both available could be included in this analysis. To test the relationship between

expected cancer risk and observed neoplasia, cancer, and malignancy transformation rate for each group,

while taking phylogenetic relationships into account, we used generalized estimating equations

(*compar.gee*, R package {ape}) [17,41] between the log scaled intrinsic cancer risk and neoplasia, cancer, or

malignancy transformation rate for each group using R v4.1.1 [42].

Statistical Analyses

The Shannon-Wiener Diversity Index and the Shannon Equitability Index were used to assess

species diversity and the evenness of species distribution in our dataset [43,44]. Additional information on

Methods and Results for this can be found in the Supplementary Materials (**Supplementary Materials**

Table 2).

To determine if the prevalence of neoplasia, cancer, and malignancy transformation rates differ

significantly among groups, we calculated 95% confidence intervals for each group's prevalence values

separately. Prevalence of neoplasia, cancer and malignancy transformation rate among tetrapod groups

were compared in pairwise fashion using the test of equal proportion using the R function *prop.test* and

p.adjust with the Benjamini & Hochberg ("BH") method to account for multiple testing (R package {stats}).

To our knowledge, this is the first work that considers pairwise comparisons of cancer prevalence within

and between each group, thus assessing which prevalence values are significantly different from one

another. Between species, prevalence, using neoplasia/cancer incidence and sample size, were tested

against the mean of their respective group using *prop.test* (R package {stats}) to identify outliers with

significantly (p = <0.05) different prevalence from the group mean (**Supplementary Materials Table 3**).

To assess the influence of lifespan or body mass on neoplasia, cancer, and malignancy

transformation rate, statistical analyses were run for neoplasia prevalence, cancer prevalence, and

malignancy transformation rate. Analyses were run for the entire dataset and also considering species with

≥10 and 20 necropsies per species – both thresholds used in previously published papers on this topic [2,3,4,6,10]

(**Supplementary Materials Table 4** contains necropsy sample sizes for each group for which 20 or more

species were available with 10 and 20 or more necropsies). We repeated analyses with the full dataset and

with species having more necropsies to ensure higher phylogenetic resolution and avoid bias from

overrepresented clades (**Figure 3**). Furthermore, as in the statistical approach used in this paper, individual

data and not species prevalence are used for the analyses and the relative weight of the number of

individuals per species is taken into account by this approach, we want to assess the influence of using a

different number of necropsies per species on the results. Finally, in order to obtain an idea of the influence

of interspecific sampling within groups, analyses were run by jackknifing species of the dataset and a 95%

confidence interval was obtained for the estimated parameters.

For the statistical analyses, mass and lifespan data were log-transformed to linearize potential

allometric effects. We used generalized estimating equations (*compar.gee*) to estimate the influence of body

mass or lifespan on neoplasia, cancer, and malignancy transformation rate while considering the

phylogenetic relationships within each group [17,41]. As cancer prevalence is the response variable and is a

proportion, the binomial family was used for the model and the number of positive (occurrence of neoplasia,

cancer) and negative (non-occurrence) individuals was entered as raw data for each species. For these

comparative analyses in which phylogenetic relationships and each life history trait was analyzed, we only

used species that could be matched with a published phylogeny for their group and for which at least one

life history trait (lifespan and/or body mass) could be found online or in literature (see Life History Data

Collection). The novelty of this approach is that instead of using proportions for neoplasia, cancer, or

malignancy transformation rate per species, as commonly done [3,4,6,10], we directly used individual count for

each species. This ensures that per species sample size is considered directly in the model, without the need

of giving different weights according to sampling; this also helps to directly model the prevalence as a

logistic response. Numbers of species per group for all analyses can be found in **Table 1** and in the

Supplementary Materials (**Table 4**). These analyses were not run for the crocodilians as there were too few

species to allow convergence of the model.

Phylogenetic trees for each studied group were obtained from the following sources: amphibians

[45], birds [46], crocodilians [47], mammals [48], squamates [49], and turtles [50]. All phylogenetic trees were transformed

to ultrametric trees using the *chronos* function in R package [ape], using default parameters (no fossil

calibration). All the statistical analyses were run in R v4.1.1.

Re-analysis of Previously Published Results on Mammals

To assess the influence of our statistical approach on the results obtained from previous studied,

we reanalyzed data from Boddy et al. [3] and Vincze et al. [4] separately using *gee*. Body mass and lifespan

data were log transformed as the authors did in their papers. For the Vincze et al. [4] data, we used the

estimates of life expectancy data calculated by the authors instead of maximum lifespan for species,

following what the authors did in their paper. Vincze et al. [4] also had only one category for neoplastic

conditions represented by any neoplasm that was diagnosed as cause of death while Boddy et al. [3] separated

total neoplasms and cancer as in our work. As such, the data of Vincze et al. [4] represent cancer mortality

prevalence, while the data of Boddy et al. [3] were analyzed for neoplasia, cancer, and malignancy

transformation rate. Note that while Boddy et al. [3] did not use their data to calculate the malignancy

transformation rate, their data could be used to calculate all the categories used in this paper.

ACKNOWLEDGMENTS

This project was supported by NSF IOS joint collaborative awards (No. 2028458 to VJL and No. 2028459

to SG and YC). We are grateful to Harald Schmidt for helping us to obtain the data at the Rotterdam Zoo,

to Isabella Capellini for providing the avian phylogeny used in our study, and to Beth Reinke for allowing

us access to reptile and amphibian life history trait data. We are also grateful to Robert Ossiboff for our

discussion on veterinary medicine and pathology reports on different vertebrate groups. Finally, we are

thankful to the comments received by two anonymous reviewers, which strongly improved the clarity and

potential impact of our paper.

AUTHOR CONTRIBUTIONS

Conceptualization: YC, SG; Investigation: SB, LP, WA, NM, J-PA, SD, SMcC, PW, SB, LRD, LGRB -

vanS, DF, SG, YC; Data curation: SB, SG, YC; Formal Analysis: SB, JC, SG, YC; Writing – Original

Draft: SB, SG, YC; Writing – Review and Editing: SMcC, PW, LGRB - vanS, DF, VJL, JC.

COMPETING INTERESTS

Authors claim no competing interests.

DATA AVAILABILITY

All data and R codes used for this paper will be available on Dryad after manuscript acceptance.

REFERENCES

1. Caulin, A. F., & Maley, C. C. (2011). Peto's Paradox: Evolution's prescription for cancer prevention. *Trends in Ecology & Evolution*, *26*(4), 175–182. https://doi.org/10.1016/j.tree.2011.01.002

2. Abegglen, L.M., Caulin, A.F., Chan, A., Lee, K., Robinson, R., Campbell, M., Kiso, W.K., Schmitt, D.L., Waddell, P.J., Bhaskara, S., Jensen, S.T., Maley, C.C., Schiffman, J.D. (2015). Potential mechanisms for cancer resistance in elephants and comparative cellular response to DNA damage in humans. JAMA. 2015;314(17):1850-1860. https://doi.org/10.1001/jama.2015.13134

3. Boddy, A. M., Abegglen, L. M., Pessier, A. P., Aktipis, A., Schiffman, J. D., Maley, C. C., & Witte, C. (2020a). Lifetime cancer prevalence and life history traits in mammals. *Evolution, Medicine, and Public Health*, *2020*(1), 187–195. https://doi.org/10.1093/emph/eoaa015

4. Vincze, O., Colchero, F., Lemaître, J.-F., Conde, D. A., Pavard, S., Bieuville, M., Urrutia, A. O., Ujvari, B., Boddy, A. M., Maley, C. C., Thomas, F., & Giraudeau, M. (2022). Cancer risk across mammals. *Nature*, *601*(7892), 263–267. https://doi.org/10.1038/s41586-021-04224-5

5. Harris, V. (2022). *An Evolutionary Perspective on Neoplastic Disease Across the Tree of Life* [Arizona State University]. ProQuest. https://www.proquest.com/docview/2671962967?fromopenview=true

6. Kapsetaki, S., Compton, Z., Dolan, J., Harris, V., Rupp, S., Duke, E., Harrison, T., Aksoy, A., Giraudeau, M., Vincze, O., McGraw, K., Aktipis, A., Tollis, M., Boddy, A., & Maley, C. (2024). Life history traits and cancer in birds. *Evolution, Medicine, and Public Health* eoae011, https://doi.org/10.1093/emph/eoae011

7. Nunney, L. (1999). Lineage selection and the evolution of multistage carcinogenesis. *Proceedings of the Royal Society of London. Series B: Biological Sciences*, *266*(1418), 493–498. https://doi.org/10.1098/rspb.1999.0664

8. Nunney, L. (2015) Size matters: height, cell number, and a person's risk of cancer. Proc. R. Soc. B 285: 20181743. http://dx.doi.org/10.1098/rspb.2018.1743

9. Chiari, Y., Glaberman, S., & Lynch, V. J. (2018). Insights on cancer resistance in vertebrates: Reptiles

as a parallel system to mammals. *Nature Reviews Cancer*, *18*, 525. https://doi.org/10.1038/s41568-018-0033-4

10. Compton, Z., Mellon, W., Harris, V., Mellon, W., Rupp, S., Mallo, D., Kapsetaki, S., Wilmot, M., Kennington, R., Noble, K., Baciu, C., Ramirez, L., Peraza, A., Martins, B., Sudhakar, S., Aksoy, S., Furukawa, G., Vincze, O., Giraudeau, M., Duke, E., Spiro, S., Flach, E., Davidson, H., Zehnder, A., Graham, T., Troan, B., Harrison, T., Tollis, M., Schiffman, J., Aktipis, A., Abegglen, L., Maley, C., & Boddy, A. (2024). Cancer Prevalence Across Vertebrates. *BioRxiv.* Preprint. https://www.biorxiv.org/content/10.1101/2023.02.15.527881v2

11. Effron, M., Griner, L., & Benirschke, K. (1977). Nature and Rate of Neoplasia Found in Captive Wild Mammals, Birds, and Reptiles at Necropsy. *JNCI: Journal of the National Cancer Institute*, *59*(1), 185–198. https://doi.org/10.1093/jnci/59.1.185

12. Madsen, T., Arnal, A., Vittecoq, M., Bernex, F., Abadie, J., Labrut, S., Garcia, D., Faugère, D., Lemberger, K., Beckmann, C., Roche, B., Thomas, F., & Ujvari, B. (2017). Cancer Prevalence and Etiology in Wild and Captive Animals. In *Ecology and Evolution of Cancer* (pp. 11–46). Elsevier. https://doi.org/10.1016/B978-0-12-804310-3.00002-8

13. Møller, A. P., Erritzøe, J., & Soler, J. J. (2017). Life history, immunity, Peto's paradox and tumours in birds. *Journal of Evolutionary Biology*, *30*(5), 960–967. https://doi.org/10.1111/jeb.13060

14. Pesavento, P. A., Agnew, D., Keel, M. K., & Woolard, K. D. (2018). Cancer in wildlife: Patterns of emergence. *Nature Reviews Cancer*, *18*(10), 646–661. https://doi.org/10.1038/s41568-018-0045-0

15. Garner, M. M., Hernandez-Divers, S. M., & Raymond, J. T. (2004). Reptile neoplasia: A retrospective study of case submissions to a specialty diagnostic service. *Veterinary Clinics: Exotic Animal Practice*, *7*(3), 653–671. https://doi.org/10.1016/j.cvex.2004.04.002

16. Sykes, J. M., & Trupkiewicz, J. G. (2006). Reptile neoplasia at the Philadelphia Zoological Garden, 1901–2002. *Journal of Zoo and Wildlife Medicine*, *37*(1), 11–19. https://doi.org/10.1638/04-112.1

17. Paradis, E., & Claude, J. (2002). Analysis of Comparative Data Using Generalized Estimating Equations. *Journal of Theoretical Biology*, *218*(2), 175–185.

https://doi.org/10.1006/jtbi.2002.3066

18. Duke, E. G., Harrison, S. H., Moresco, A., Trout, T., Troan, B. V., Garner, M. M., Smith, M., Smith,

S., & Harrison, T. M. (2022). A Multi-Institutional Collaboration to Understand Neoplasia,

Treatment and Survival of Snakes. *Animals*, *12*(3), 258. https://doi.org/10.3390/ani12030258

19. *The IUCN Red List of Threatened Species*. (n.d.). IUCN Red List of Threatened Species. Retrieved June

1, 2022, from https://www.iucnredlist.org/en

20. Peto, R. (2015). Quantitative implications of the approximate irrelevance of mammalian body size and

lifespan to lifelong cancer risk. *Philosophical Transactions of the Royal Society B: Biological*

Sciences, *370*(1673), 20150198. https://doi.org/10.1098/rstb.2015.0198

21. Vazquez, J. M., & Lynch, V. J. (2021). Pervasive duplication of tumor suppressors in Afrotherians

during the evolution of large bodies and reduced cancer risk. *ELife*, *10*, e65041.

https://doi.org/10.7554/eLife.65041

22. Chang, P., Wang, F., Chin, S., Pang, V. F., Jeng, C., Liu, C., Chu, P., & Zhuo, Y. (2012). Spontaneous

neoplasms in zoo mammals, birds, and reptiles in Taiwan – a 10-year survey. *Animal Biology*,

62(1), 95-110. https://doi.org/10.1163/157075611X616941

23. Kitsoulis, C.V., Baxevanis, A.D. & Abatzopoulos, T.J. (2020) The occurrence of cancer in vertebrates:

a mini review. *Journal of Biological Research-Thessaloniki*, 27, 9. https://doi.org/10.1186/s40709-

020-00119-0

24. Lourenço, J. M., Glémin, S., Chiari, Y., & Galtier, N. (2013). The determinants of the molecular

substitution process in turtles. *Journal of Evolutionary Biology*, *26*(1), 38–50.

https://doi.org/10.1111/jeb.12031

25. da Silva, R., Conde, D. A., Baudisch, A., & Colchero, F. (2022). Slow and negligible senescence among

testudines challenges evolutionary theories of senescence. *Science*, 376(6600), 1466–1470.

26. Glaberman, S., Bulls, S. E., Vazquez, J. M., Chiari, Y., & Lynch, V. J. (2021). Concurrent evolution of

anti-aging gene duplications and cellular phenotypes in long-lived turtles *Genome Biology and*

Evolution, 13 (12), evab244. https://doi.org/10.1093/gbe/evab244

27. Reinke, B. A., Cayuela, H., Janzen, F. J., Lemaître, J.-F., Gaillard, J.-M., Lawing, A. M., Iverson, J. B.,

Christiansen, D. G., Martínez-Solano, I., Sánchez-Montes, G., Gutiérrez-Rodríguez, J., Rose, F. L.,

Nelson, N., Keall, S., Crivelli, A. J., Nazirides, T., Grimm-Seyfarth, A., Henle, K., Mori, E., …

Miller, D. A. W. (2022). Diverse aging rates in ectothermic tetrapods provide insights for the

evolution of aging and longevity. *Science*, 376(6600), 1459–1466.

https://doi.org/10.1126/science.abm0151

28. Corradetti, B., Dogra, P., Pisano, S., Wang, Z., Ferrari, M., Chen, S.-H.., Sidman, R. L., Pasqualini, R.,

Arap, W., Cristini, V. (2021). Amphibian regeneration and mammalian cancer: Similarities and

contrasts from an evolutionary biology perspective. *BioEssays,* 43, e2000339.

https://doi.org/10.1002/bies.202000339.

29. Pomerantz, J. H., Blau, H. M. (2013) Tumor suppressors: enhancers or suppressors of regeneration?

Development, 140 (12), 2502–2512. https://doi.org/10.1242/dev.084210

30. Sarig, R., Tzahor, E. (2017) The cancer paradigms of mammalian regeneration: can mammals

regenerate as amphibians? *Carcinogenesis*, 38 (4), 359–366.

https://doi.org/10.1093/carcin/bgw103

31. Torres-Dimas, E., Cruz-Ramírez, A., Bermúdez-Cruz, R. M. (2022). Cancer in Amphibia, a rare

phenomenon? *Cell Biology International*, 46(12), 1992-1998. doi: 10.1002/cbin.11888.

32. Vieira WA, Anderson K, Glass Campbell L, McCusker CD. Characterizing the regenerative capacity

and growth patterns of the Texas blind salamander (*Eurycea rathbuni*). *Developmental*

Dynamics. 2021; 250: 880–895. https://doi.org/10.1002/dvdy.245

33. Ginal, P., Stahlberg, J., Rauhaus, A., Wagner, P., Rödder, D., Ziegler, T. (2023). Threatened turtles and

tortoises (Testudines) in zoos: A ZIMS database analysis for improved One Plan Approach to

Conservation actions. *Salamandra* 59(3): 262–274.

34. Tidière, M., Gaillard, JM., Berger, V. *et al.* Comparative analyses of longevity and senescence reveal

variable survival benefits of living in zoos across mammals. *Scientific Reports* 6, 36361 (2016).

https://doi.org/10.1038/srep36361

35. ZIMS. Species360 Zoological Information Management System. https://zims.species360.org.

36. Wagner, G. P., Kshitiz, & Levchenko, A. (2020). Comments on Boddy et al. 2020: Available data

suggest positive relationship between placental invasion and malignancy. *Evolution, Medicine, and*

Public Health, *2020*(1), 211–214. https://doi.org/10.1093/emph/eoaa024

37. Tacutu, R., Thornton, D., Johnson, E., Budovsky, A., Barardo, D., Craig, T., Diana, E., Lehmann, G.,

Toren, D., Wang, J., Fraifeld, V. E., & de Magalhães, J. P. (2018). Human Ageing Genomic

Resources: New and updated databases. *Nucleic Acids Research*, *46*(D1), D1083–D1090.

https://doi.org/10.1093/nar/gkx1042

38. Oliveira, B. F., São-Pedro, V. A., Santos-Barrera, G., Penone, C., & Costa, G. C. (2017). AmphiBIO, a

global database for amphibian ecological traits. *Scientific Data*, *4*(1), 170123.

https://doi.org/10.1038/sdata.2017.123

39. Dodd, CK (2010). Amphibian Ecology and Conservation. A Handbook of Techniques. Oxford

University Press, Oxford

40. Vitt, LJ., Caldwell, JP. (2013). Herpetology: An Introductory Biology of Amphibians and Reptiles.

Academic Press, USA

41. Paradis, E., Claude, J., & Strimmer, K. (2004). APE: Analyses of Phylogenetics and Evolution in R

language. *Bioinformatics*, *20*(2), 289–290. https://doi.org/10.1093/bioinformatics/btg412

42. R Core Team. (2018). *R: A language and environment for statistical computing.* R Foundation for

Statistical Computing, Vienna, Austria. https://www.R-project.org/

43. Shannon, C. E. (1948). A mathematical theory of communication. *The Bell System Technical Journal*,

27(3), 379–423. https://doi.org/10.1002/j.1538-7305.1948.tb01338.

44. Tuomisto, H. (2012). An updated consumer's guide to evenness and related indices. *Oikos*, 121(8),

1203–1218. https://doi.org/10.1111/j.1600-0706.2011.19897.x

45. Pyron, A. R., & Wiens, J. J. (2011). A large-scale phylogeny of Amphibia including over 2800 species,

and a revised classification of extant frogs, salamanders, and caecilians. *Molecular Phylogenetics*

and Evolution, *61*(2), 543–583. https://doi.org/10.1016/j.ympev.2011.06.012

46. Jetz, W., Thomas, G. H., Joy, J. B., Hartmann, K., & Mooers, A. O. (2012). The global diversity of

721   birds in space and time. *Nature*, 491(7424), 444–448. https://doi.org/10.1038/nature11631

722   47. Groh, S. S., Upchurch, P., Barrett, P. M., & Day, J. J. (2020). The phylogenetic relationships of

723   neosuchian crocodiles and their implications for the convergent evolution of the longirostrine

724   condition. *Zoological Journal of the Linnean Society*, *188*(2), 473–506.

725   https://doi.org/10.1093/zoolinnean/zlz117

726   48. Upham, N. S., Esselstyn, J. A., & Jetz, W. (2019). Inferring the mammal tree: Species-level sets of

727   phylogenies for questions in ecology, evolution, and conservation. *PLOS Biology*, *17*(12),

728   e3000494. https://doi.org/10.1371/journal.pbio.3000494

729   49. Pyron, R. A., Burbrink, F. T., & Wiens, J. J. (2013). A phylogeny and revised classification of Squamata,

730   including 4161 species of lizards and snakes. *BMC Evolutionary Biology*, *13*(1), 93.

731   https://doi.org/10.1186/1471-2148-13-93

732   50. Thomson, R. C., & Shaffer, H. B. (2010). Sparse Supermatrices for Phylogenetic Inference: Taxonomy,

733   Alignment, Rogue Taxa, and the Phylogeny of Living Turtles. *Systematic Biology*, *59*(1), 42–58.

734   https://doi.org/10.1093/sysbio/syp075