Including Phylogenetic Conservatism of Shortgrass Prairie Restoration Species Does Not

Improve Species Germinability Prediction

Yanni Chen[1], Dylan W. Schwilk[1], Robert D. Cox[2], Matthew G. Johnson[1]

[1]Department of Biological Sciences, Texas Tech University, 2901 Main Street, Lubbock, Texas 79409, USA

[2]Department of Natural Resources Management, Texas Tech University, Lubbock, TX 79410, USA

ABSTRACT

PREMISE

We investigated whether phylogenetic conservatism can improve the performance of seed

germinability prediction models. Previous studies in tallgrass prairie and alpine meadow revealed

that seed morphological traits demonstrate phylogenetic conservatism. We hypothesized that

phylogenetic conservatism in seed traits could help predict the seed germinability, under the

assumption that seed traits contain phylogenetic signals.

METHODS

We measured seed germination percentage and seed morphological traits (seed mass,

seed height, and seed surface area) on 34 native species from shortgrass prairie in North

America. We supplemented these data with similar data from the literature on 11 more species.

We calculated the robustness of the phylogenetic signal of each trait to the number of species

sampled. We also compressed the phylogenetic distance matrix to a two-dimensional space, and

applied the Akaike information criterion to evaluate the effects of phylogeny on seed

germinability prediction models.

KEY RESULTS

We found weak but significant phylogenetic signals in seed mass and seed height in the

full data set. These phylogenetic signals were not able to improve seed germinability prediction

model performance among shortgrass prairie species. Our robustness tests of phylogenetic

signals using random sub-sampling showed that the detection rate of phylogenetic signals in seed

mass was increased along with the expansion of species pool, and nearly 100% at 40 species.

However, the detection rate of phylogenetic signals in seed height was constantly low, around

20%.

CONCLUSIONS

When the phylogenetic signals are weak, the phylogenetic position does not improve

germinability prediction model performance. Therefore, phylogenetic signals detected during a

single species pool calculation may not accurately reflect the phylogenetic conservatism of the

trait in a plant community. We suggest testing for robustness of phylogenetic signals using

random sub-sampling tests.

Keywords: ecological restoration, seed germinability prediction, phylogenetic comparative

method, phylogenetic conservatism, phylogenetic signal

Introduction

The need for ecological restoration is constantly increasing. For example, the September

2014 United Nations Climate Summit suggested the need for 350 million hectares to be restored

worldwide by 2030 (Bonn Challenge, https://www.bonnchallenge.org/). Tremendous numbers of

native species will be needed to meet this need. Most ecological restoration projects select only a

small number of species out of the community species list to conduct ecological restoration

(Kiehl *et al.* 2010). Given the low numbers of species selected for any specific restoration

project, maximizing the benefit from selected species is key. Thus, ensuring that the selected

species have high final germination percentages is a high priority because seed germination

ranks as one of the top restoration challenges (Larson *et al.* 2015). Therefore, lab assessment

formulas to narrow down the restoration species list could aid species selection in many

restoration projects.

Seed dormancy regulates seed germination but is complicated and hard to predict. In over

90% of species, seeds dry and start primary dormancy by the time of harvest (Finch-Savage and

Leubner-Metzger 2006; Subbiah *et al.* 2019). After dispersal, seeds can have secondary

dormancy, a shallow physiological dormancy which is broken by responses to environmental

cues (Finch-Savage and Leubner-Metzger 2006). Multiple categorical seed dormancy types are

widely represented in plant species, including morphological dormancy (MD), physical

dormancy (PY), physiological dormancy (PD), and morphophysiological dormancy (MPD;

(Baskin and Baskin 1998). Physiological dormancy is thought to be the ancestral state of seed

dormancy and also serves as the diversification hub for different dormancy types (Willis *et al.*

2014). Considering the complexity of dormancy stages and the lengthy experiments needed to

distinguish these types (Finch-Savage and Leubner-Metzger 2006), it is desirable to predict seed

germinability success through other related traits.

Low germination rate hinders restoration and, given limited resources, managers desire to

only include species with predictably high germination rates. Several seed traits are related to

seed germination and might serve as more easily measured predictors of final germination

percentage. In general, mass is a good indicator of seed germination, as small seeds tend to

germinate faster (Westoby *et al.* 2002a; Barak *et al.* 2018), while large seeds can stay dormant

longer and produce stronger seedlings after germination (Leishman *et al.* 2000; Westoby *et al.*

2002a). The rationale behind this phenomenon is related to nutrition stored in the seed under

either a "larger-seed-later-deployment" interpretation (Ganade and Westoby 1999; Leishman *et*

al. 2000; Kidson and Westoby 2000) or "cotyledon functional morphology" hypothesis (Hladik

and Miquel 1990; Kitajima 1996a, b). Furthermore, seed size and seed shape are also traits

influencing seed germination by stimulating or delaying seed germination through wind, water,

or animal dispersal (Howe and Smallwood 1982). Large seeds generally have advantages for

dispersal related to entrapment strategies, such as net trapping, surface tension, and wake

trapping (Jager *et al.* 2019), especially for wind-dispersed species (Zhu *et al.* 2019). Specifically,

seed morphological traits influence both seed primary dispersal (seed departure from parent

plants) and secondary wind dispersal (seed lifting off the ground by wind power) (Zhu *et al.*

2019). Primary dispersal is mainly driven by dispersal height and terminal falling velocity, which

are influenced by seed morphology (Sheldon and Burrows 1973; Jongejans and Telenius 2001).

Secondary dispersal distance strongly depends on the lift-off velocity, which is influenced by

seed height and seed surface area (van Tooren 1988; Schurr *et al.* 2005; Zhu *et al.* 2022). There

are many other seed physiological traits associated with seed germination that are not commonly

tested, such as base water potential, cardinal temperature, thermal time and hydrothermal time

for germination (Bradford 2002; Hardegree *et al.* 2013).

Seed germination trials are time consuming, therefore, predicting germinability for

species without conducting such trials could benefit restoration. Seed morphology traits are

potential predictors of germination rate. If dormancy and lack thereof are evolutionarily

conserved, then it may be possible to predict seed germination rate of unmeasured species based

on the rates of closely related taxa. A phylogenetic tree models the inferred evolutionary

branching history of a group of taxa (Baum and Smith 2013). A phylogenetically conserved trait

will tend to be most similar among species close together on the phylogenetic tree. The common

test for such phylogenetic signals is Blomberg's K (Blomberg *et al.* 2003, Revell 2008), but it is

also possible to include all pairwise phylogenetic distances among taxa in linear models through

the method of phylogenetic residuals (Revell 2010). Phylogenetic trait conservatism is common

across many traits and clades (Bu *et al.* 2016, Barak *et al.* 2018, Duncan *et al.* 2019). Adding

phylogenetic residuals to the generalized least square model can take the evolution of

unmeasured traits into account and improve the prediction model's accuracy. This work has two

major goals. The first goal is to test whether adding phylogenetic information among species

(presented by x-y coordinates transferred from phylogenetic tree topology) can improve

predictions of germination rate based on seed morphology. Adding phylogenetic information

might improve predictions if the germination rate shows a phylogenetic signal or if the seed

morphology effect on germination rate interacts with phylogeny. There is precedent for using

phylogeny for this purpose: In a study of species native to tallgrass prairie, Barak et al. (2018)

confirmed that adding the phylogenetic residual improved the accuracy of the seed germinability

prediction model due to the phylogenetic conservatism in both seed germination and

morphological traits. However, because phylogenetic tools are unfamiliar and inaccessible to

restoration practitioners and due to a historical separation between evolutionary biology and

applied ecology, phylogenetic methods have not been broadly applied to restoration practice

(Hipp *et al.* 2015).

The second major goal of this work is to determine how the size of a sample of taxa from

an ecological community influences the power to detect phylogenetic signals in traits. The

sample size and combination of given species influence tree topology and branch length during

phylogenetic signal calculation. In empirical examples, the detection of phylogenetic signals is

strongly related with the number of species included, with twenty or more species usually

considered sufficient for estimation of Blomberg's K (Blomberg *et al.* 2003) . However, in

phylogenetic comparative analysis aimed at answering evolutionary questions, the combination

of species is commonly fixed. For applied restoration use, the practitioner will need to measure

traits on some sample of species from a particular community. By examining how this sample of

taxa influences the calculation of Blomberg's K, we aim to provide guidelines for estimating the

robustness of this calculation.

To address these two major goals and test the potential role of phylogeny for improving

restoration practice, we asked four research questions: 1) Do seed traits and seed final

germination percentages exhibit phylogenetic signals? 2) Among seed traits, which one is the

best predictor of seed final germination percentage? 3) Does including phylogenetic residuals

improve the seed germinability prediction? 4) Do the sampling size and species composition

influence phylogenetic conservatism detection in shortgrass prairie species?

Material and Methods:

To determine the relationship between seed germinability and seed morphological traits,

we measured seed germination percentage, seed mass, seed height, and seed surface area in 45

species which are native to the shortgrass prairie of North America (Table 1). All of our raw data

and calculations were demonstrated in our interactive Shiny Application (Figure 5,

https://chenyanniii.shinyapps.io/Phylo_Compar_Traits/).

<u>Seed Germination Percentage and Morphological Traits Measurements</u>

Seed germination percentage was obtained from two sources: our own germination trials

and previous publications. In all cases, we defined "germination percentage" as the maximum

final germination percentage obtained. The germination trials followed a simple germination

protocol without cold stratification or other attempts to break dormancy, which simulated

minimum requirements for restoration projects. This simple protocol is essentially a

measurement of lack of dormancy assuming the tested seeds were full viable. For 34 of the 45

species, we conducted new germination trials. Our new germination trials were trying to simulate

the scenario that practitioners want to find some easy to use native species. Because the

experiment is trying to simulate the scenario in which practitioners are attempting to find easy to

use native species, we bought seeds from a local restoration seed vendor (Native American

Seed), and chose species for which they offered local seed sources (and recorded the seed

source), with seeds that were harvested less than 6 years ago. When seeds arrived, we stored the

seeds in a dry and dark place at room temperature (20 °C) until experiments started. Although

it's possible that some species may exhibit dormancy, we didn't use any dormancy breaking

treatment, in order to simulate simple restoration practice. For the germination experiment, we

used triple replicated germination trials: disposable petri dishes with lids were placed in

germination chambers (20 °C day and night, with 15 hours and 9 hours day night shift). Inside a

petri dish a piece of filter paper was placed to observe auto-claved water to keep the seeds moist.

We checked the water sufficiency every day. In each germination trial we split a total of 50 seeds

of each species into 5 petri dishes. Since our study used commercial seeds and focused on

species dormancy status, we assumed our seeds will either be dormant or start germination

within a month. Our observations during experiments proved this assumption. The seeds

generally started germinating within 10 days or stayed dormant through the whole germination

trial. Our germination trials ran until one week after the last seed germinated. Most of the seed

germination trials finished within a month, and all the trials finished within two months. Three

independent trials happened in July 2019, September 2019, and November 2019. For the

remaining 11 species, we used final germination percentages reported in two published studies

(Schwilk and Zavala 2012; Chou *et al.* 2012). These two studies were originally designed for

detecting smoke effects on shortgrass prairie species, but we used the control treatment data only

which provided conditions similar to those in our trials (20-25 °C, 12-16 hours illumination).

We measured seed mass using an electronic balance (Sartorius Analytical Balance LA

230P, 0.1mg readability) in lab conditions with 10 replicates of 100 seeds each per species. For

species in which we could not obtain 100 seeds, we used 30 seeds per replicate.

We measured seed surface area and seed height through digital image processing with 10

replicates. The seed surface was defined by the two largest orthogonal axes, the height was

defined as the third axis. We calculated the seed surface area by digital image of the maximum

surface area of seeds and imaged under a stereomicroscope at 400 magnification. We

transformed the images to 8-bit (black and white) and calculated the surface area using the

"analyze particle" function in ImageJ (Fiji, Schindelin et al. 2019). We also recorded the seeds'

heights calculated by the z-stack image function and NIS-Element BR 4.60.00 software.

<u>Species Phylogenetic Information</u>

We generated a phylogenetic tree of all study species using two methods: pruning

existing phylogeny (Zanne et al. 2014) and binding non-existing tips to the phylogeny based on

their taxonomic information. The phylogeny (Zanne et al. 2014) we used in this study was a

time-calibrated maximum-likelihood-based phylogenetic tree, built with seven genes (18S

rDNA, 26S rDNA, ITS, matK, rbcL, atpB, and trnL-F) downloaded from GenBank. First, we

confirmed that every genus in our study was on the Zanne phylogeny. Second, we created a

function to prune species which were not on the tree, and we also swapped the species under the

same genus if the exact species was not on the tree (see the function of func_prun_replac on

https://github.com/chenyanniii/Traits4 repo for more detail). The results showed that 30 species

on the tree and 15 missing species (*Argemone albiflora, Asclepias asperula, Astragalus*

crassicarpus, Callirhoe leiocarpa, Centaurea americana, Chasmanthium latifolium, Corydalis

curvisiliqua, Digitaria californica, Eragrostis trichodes, Herbertia lahue, Liatris mucronata,

Linum rigidum, Pavonia lasiopetala, Polytaenia nuttallii, Tradescantia occidentalis). After

applying func_prun_replac, 13 of 15 species were placed based in their genus and only two

species (*Callirhoe leiocarpa* and *Digitaria californica*) were missing. Thus, we added the

missing species (*Callirhoe leiocarpa* and *Digitaria californica*) as sister tips to *Callirhoe*

involucrate and *Digitaria ciliaris* under the same genus assuming that phylogenetic relationships

were consistent with their taxonomic grouping. Our final tree contained all species was a

dichotomous tree (Figure 1).

To incorporate phylogenetic relatedness in the general linear models, we represented the

phylogeny by all pairwise phylogenetic distances across taxa. We converted the pairwise

distance matrix to points distributed in a two-dimensional coordinate system, using nonmetric

multidimensional scaling (NMDS) (isoMDS function in the package MASS, Venables and Ripley

2002). We evaluated phylogenetic signals for individual traits as Blomberg's K (Blomberg *et al.*

2003) using the phylosig function in the phytools R package (Revell 2012). We tested for

phylogenetic signal using a randomization test (phylosig function) that compared the measured

value of Blomberg's K against a distribution of K calculated when trait values were randomized

across the tips of the phylogeny.

<u>Germinability Prediction Model Selection</u>

To generate and evaluate generalized linear models, we applied backward stepwise model

comparison based on the Akaike information criterion (Akaike 1998) using the AICc function in

the AICcmodavg package (Mazerolle 2020). We also used seed germination percentage, three

seed morphological traits (seed mass, seed height and seed surface area) and phylogenetic

positions to generate a global general linear model. Then, we used AIC to correct for small

sample sizes (AICc) and evaluate the fitness of models. We standardized all input parameters to

the mean of zero to produce standardized coefficients between parameters for numeric reasons in

fitting. We also tested correlation among morphological traits (seed mass, seed height and seed

surface area). All original data and scripts that we used to calculate phylogenetic signals,

phylogenetic residuals, and seed germinability prediction models are available on GitHub

website (https://github.com/chenyanniii/Traits4, DOI: 10.5281/zenodo.6609175).

<u>Random Sub-sampling of Different Species Pool Size</u>

To estimate the minimum species pool size for obtaining a stable phylogenetic signal, we

created 31 different species pool subsets, from 10 species to 40 species. For each pool size, we

randomly withdrew 100 times at each pool size species from the whole species pool, thus

generating 100 sub-pools of each species pool size by random sub-sampling. The phylogenetic

signals of each sub-pool were calculated for their Blomberg's K and related p value. We

analyzed the relationship between sample size and detection rate of phylogenetic signals was

analyzed to evaluate the effect of sample size to estimated phylogenetic signals in traits.

<u>Shiny Application</u>

Shiny is a web framework for displaying data. Shiny is a good data processing

demonstration tool, an interactive way for users to experience how different input and procedure

affect output. We designed our shiny application to import with our full dataset and display data

analysis and results. Users can see our full dataset result (as default), or interactively calculate all

parameters for any sub-pools using checkboxes of species (Figure 5).

Results

In this study, we used 45 commonly selected restoration species to explore the

phylogenetic distance among shortgrass prairie species by pruning unnecessary species and

adding desired species to the existing phylogenetic tree of flowering plants (Figure 1).

<u>Seed Final Germination Percentage and Morphological Traits Measurements</u>

When examining species' trait value with the phylogenetic tree (Figure 1), we found the

phylogenetic patterns in seed mass, seed height, seed surface area and seed germination rate were

varied. We were not able to germinate eight species (Figure 1, *Argemone albiflora*, *Callirhoe*

leiocarpa, *Corydalis curvisiliqua*, *Herbertia lahue*, *Oenothera rhombipetala*, *Pavonia*

lasiopetala, *Phytolacca americana*, and *Polytaenia nuttallii*). *Eragrostis trichodes* had the

highest final germination percentage, 82%. For seed mass, *Sporobolus airoides* had the lightest

weight per seed, 0.0945 ± 0.0083 mg per seed; the heaviest seed was *Pavonia lasiopetala*, 18.75

± 0.3487 mg per seed. The seed height measurement ranged from 0.658 ± 0.1051 (*Coreopsis*

tinctoria) to 2.995 ± 0.1334 mm (*Pavonia lasiopetala*); and the seed surface areas ranged from

0.361 ± 0.0083 (*Sporobolus cryptandrus*) to 25.258 ± 1.322 (*Polytaenia nuttallii*) mm^2 (Figure

1).

Species Phylogenetic Information

We used nonmetric multidimensional scaling (NMDS) to compress the phylogenetic

distance matrix to a two-dimensional space, with a pressure of 17.86. Our results showed that 45

species were grouped into three clusters: Monocot, Asteraceae and eudicots-except Asteraceae

(Figure 2). NMDS compressed phy1 (x-axis) corresponded to separating monocot and eudicots,

while the phy2 (y-axis) separated Asteraceae from other families.

Our measurements of phylogenetic signals, Blomberg's K (using species shuffling

method), were low for all four seed traits, indicating a departure from signal under strict

Brownian motion and suggesting that these traits are evolutionarily labile. Although Blomberg's

K were low, indicating a weak phylogenetic signal, we found significant phylogenetic signals for

seed mass ($K = 0.07$, $p = 0.01$) and seed height ($K = 0.05$, $p = 0.05$).

Germinability Prediction Model Selection

    The full set of models built from morphological traits and phylogenetic information were

evaluated using adjusted AIC (AICc). The AICc values range from 129.9 to 139.4. The best

prediction model is using seed height to predict seed germination (AICc = 129.9), slightly better

than the model using seed mass to predict germination (AICc = 130.5). The models with low

AICc values were clustered by using one morphological trait as a predictor or the combination of

two morphological traits. This indicated that morphological traits out-perform phylogenetic

distance in predicting seed germination. Pearson correlation coefficient analysis revealed a

strong correlation between seed mass and seed height (r = 0.66, p < 0.01); a medium correlation

between seed mass and seed surface area (r = 0.49, p <0.01); no correlation was detected

between seed height and seed surface area.

Random Sub-sampling of Different Species Pool Size

    We calculated phylogenetic signals of morphological traits (seed mass, seed height, and

seed surface area) and seed germination rate of all 3,100 sub-pools. All Blomberg's K values

were between 0 and 1 in all phylogenetic signal calculations, except 9 of them were larger than

1. In general, phylogenetic signals distributed widely at small species pool sizes, and became less

varied while increasing species pool sizes (Figure 3). For seed height, seed surface area, and seed

germination, the probability of detecting phylogenetic signals were consistently low regardless of

the species pool size. This was true even for seed height, for which we detected a significant

phylogenetic signal in our full dataset. In contrast, the probability of detecting the phylogenetic

signal of seed mass increased with species pool size (Figure 4).

Discussion:

295   Aiming to verify the usefulness of trait conservatism in restoration seed selection, we

measured seed traits, ran seed germination tests, calculated phylogenetic signals in seed traits,

and presented the phylogenetic residual in seed germinability prediction models. We quantified

weak phylogenetic signals in seed mass and seed height, but we found no phylogenetic signal in

seed surface area nor in seed final germination percentage. In those traits that did exhibit

phylogenetic signals, the signals were weak: closely related species were more similar than

expected under species shuffling, but more different in their trait values than expected under

Brownian motion.

Phylogenetic Tree

305   The phylogenetic tree of 45 commonly selected species in shortgrass prairie ecological

restoration was clustered in Poaceae within monocots and were relatively clustered in Asteraceae

and Lamiaceae within eudicots (Figure 1), which reflects that the species composition may be

clustered in shortgrass prairie. The phylogenetic comparative methods displayed trait values

indicated that the closely related species had similar trait values in seed mass and seed height,

but not in seed germination (Figure 1). The NMDS compressing phylogenetic distance into two-

dimensions shows three distinct clusters (Figure 2). The results showed that shortgrass prairie

families were grouped into 3 clusters: one monocot group and two eudicot groups (Asteraceae

and others, Figure 2). Meanwhile the tallgrass prairie species (Barak *et al.* 2018) were grouped

into 4 clusters: one monocot group, three eudicot groups (Asteraceae, Fabaceae, and others).

Our development of the Shiny application demonstrated: (1) the procedure of pruning the

synthetic phylogenetic tree (Zanne et al. 2014) to the desired species tree (Figure 5). (2) the

calculation of compressing phylogenetic distance into two-dimensions. The interactive

demonstration allows users to select all or a portion of desired species and understand the effect

of species selection on phylogenetic calculation.

Phylogenetic Signal in Traits

Phylogenetic signal indicates that closely related species have more similar trait value

than expected under species shuffling across tips of a phylogeny. We found significant

phylogenetic signals in seed mass and seed height, but no such signals in seed surface area nor in

seed final germination percentage. Although germination traits are not specific or constant in

each species (but vary in space and time), since we chose seeds from the same eco-region, our

results are able to represent our region and still allow generalization when considering

germinability predictions. Generally, seed mass is phylogenetically conserved in sample taxas

from different ecosystems (tallgrass prairie, Barak et al. 2018; alpine grassland, Bu *et al.* 2016

and globally Westoby 2002). In our set of taxa, we found a weak but significant pattern. Seed

mass often predicts energy and nutrient provisioning (Westoby *et al.* 2002b), which increases

seed germination rates and stress tolerance (Leishman 2000, Moles 2018). This assumes,

however, that mass is primarily the embryo and nutrients. It is possible for a large portion of the

seed mass to be seed defense structures (i.e. seed coat).

We used seed height and seed surface area as proxies for seed dispersal syndrome,

because these dimensions influence primary wind dispersal (seed departure from mother plants,

Sheldon and Burrows 1973; Jongejans and Telenius 2001) and secondary wind dispersal (seed

lifting off ground by wind power, van Tooren 1988; Schurr *et al.* 2005; Zhu *et al.* 2022). Primary

dispersal is mainly related to dispersal height and terminal falling velocity, which is influenced

by seed morphology (Sheldon and Burrows 1973; Jongejans and Telenius 2001). Secondary

dispersal distance strongly depends on the lift-off velocity, which is influenced by seed height

and the planform area of a seed exposed to airflow (van Tooren 1988; Schurr *et al.* 2005; Zhu *et*

al. 2022). Classically, seed shape was measured by the roundness or closeness of a seed to

specific shape, such as ellipse or cardioid (Cervantes et al. 2016) and linked with seed

persistence in soil seed bank (Moles *et al.* 2000; Laughlin 2014). Some recent studies link seed

morphological shape with evolutionary constraint and selective pressure of seeds and its

potential relationship with seed germination (Barak et al. 2018, Bu *et al.* 2016). In our study,

seed mass and seed height were positively correlated. We found a weak pattern of phylogenetic

trait conservatism in two traits, but this signal did not aid in improving seed germinability

prediction models.

Seed germination is a complex phenomenon. Our measure of total germination was, in

effect, a dormancy proxy: high germination rates indicated a lack of dormancy in our research.

Our experiment didn't include any dormancy breaking retreatments, only supplying light and

water during experiments to simulate practitioners' low effort practices. Seed germination can be

influenced by abiotic factors, such as wetland species germination impacted by water level

(Keddy 1992); or arid zone woody species developing rapid germination in response to

unpredictable rainfall (Duncan *et al.* 2019). Seed germination can also be influenced by biotic

factors, such as small- and large-seeded species diverging in the species they associate with,

regarding seed mass and understory light preference (Umaña *et al.* 2020). We didn't detect a

phylogenetic signal in germination rate indicating this trait is highly labile. This result was

different from a similar study of tallgrass prairie species (Barak *et al.* 2018), where the authors

found significant phylogenetic trait conservatism in germination percentage under control and

gibberellic acid treatment, and including phylogeny improve time-to-germination (survival)

model. However, the survival model (Barak *et al.* 2018) includes both germination time and

pretreatment for germination rate and doesn't measure dormancy. Differing patterns in

phylogenetic signal in germination rate of two prairie studies are reasonable, in consideration of

environmental differences between two different ecosystems, and the germination experiment

setting in two studies.

<u>Germinability Prediction Model Selection</u>

The germinability predictive models with morphological data did not improve when

adding phylogenetic information using the full dataset. This means adding phylogenetic

information to morphological measurements increased the complexity of models but did not

increase the fitness of models. This is not surprising given that we found no phylogenetic signal

in seed germination rate and only weak signals in two other traits.

<u>Random Sub-sampling of Different Species Pool Size</u>

From the distribution of Blomberg's K, we can tell the species sample size will greatly

influence phylogenetic signal calculation (Blomberg *et al.* 2003). Our shortgrass prairie

restoration species results showed that the phylogenetic signal would be less impacted by the

species composition, and less varied with sufficient species, around 35 to 40 (Figure 3). This also

indicates the 45 species we have in our study is sufficient.

In the full dataset (45 species), we were able to detect phylogenetic signals for both seed

mass and seed height. However, the subsampling exploration method demonstrates that detecting

a phylogenetic signal in seed height is a low probability event. On the other hand, our sub-

sampling in seed mass showed that the probability of detecting a phylogenetic signal increased

along with the increase in the number of species in the species pool. The Blomberg's K value is

stable at 40 species, which could indicate that if researchers or practitioners have over 40 species

sub-sampling of shortgrass prairie restoration species, their studies should be able to detect

phylogenetic signals. The random sampling methods to verify sample size method could apply in

sampling species to estimate phylogenetic conservatism in plant communities.

Shiny Application

From the Shiny application, restoration practitioners could use interactive methods to

explore our data and statistical analysis and results visualization. For readers who are first

exposed to phylogenetic comparative methods, the interactive graphic user interface can lower

the bar for exploring our data, as well as increase engagement. Our checkbox of species list

allows users to design their composition of species, and to investigate the impact of species

choice on phylogenetic signal and germinability prediction. Our Shiny application was published

on GitHub website (https://github.com/chenyanniii/Traits_Shiny, DOI:

10.5281/zenodo.6609191) and on shinyapps.io

(https://chenyanniii.shinyapps.io/Phylo_Compar_Traits/).

Comparison Between Tallgrass Prairie and Shortgrass Prairie Studies

Seed germination is a complex physiological phenomenon that could be studied for its

optimization using dormancy breaking treatments (Barak et al. 2018), as well as could be a

dormancy proxy, such as high germination rates indicated a lack of dormancy in our research.

Our research can be contrasted with a was similar tallgrass prairie study (Barak et al. 2018), in

which: (1) the phylogenetic signals of germination were detected in morphological traits and

seed germination percentage; (2) phylogenetic information improves the seed germinability

prediction model. We saw the potential of applying phylogenetic information in ecological

restoration, so we tested the phylogenetic application in simply restoration setting: (1) We

selected regional appropriated seed sources from a local restoration vendor. (2) We proxy

dormancy in seed sources by running germination trails without any dormancy breaking

treatment to approximate the conditions preferred by restoration practitioners. (3) We tested our

results against null models: confirming our confidence in sample size, examining the robustness

of our conclusion while ensuring we can generalize results for the whole shortgrass prairie plant

community. Our unique restoration scenario of shortgrass prairie showed a few advancements of

knowledge. First, only seed mass and seed height detected phylogenetic signals in 45 species.

The phylogenetic signal in seed mass is well preserved and can be generalized to estimate the

phylogenetic signal for the shortgrass prairie plant community. On the opposite, detecting a

phylogenetic signal in seed height is a low chance event that the phylogenetic signal in 45

species should not be generalized to estimate the phylogenetic signal for the shortgrass prairie

plant community. Second, estimating phylogenetic signals for a plant community needs a larger

sample size than a single fixed group. The shortgrass prairie plant community needs at around 40

species for detecting a general pattern (Figure 5), which is twice of the twenty species

assumption in a fixed species comparative study (Blomberg et al. 2003).

Conclusion and Future Studies

Overall, we have demonstrated that the phylogenetic signal calculation can be influenced

by size and composition of seed pool. We recommend running a sub-sampling test to verify the

sufficiency of species and phylogenetic conservatism in traits for a community study. Our Shiny

application is on GitHub website (https://github.com/chenyanniii/Traits_Shiny, DOI:

10.5281/zenodo.6609191) and on shinyapps.io

(https://chenyanniii.shinyapps.io/Phylo_Compar_Traits/), using an interactive way to

demonstrate how species composition directly impacts the phylogenetic signal calculation.

Our work demonstrated that morphological traits (seed mass and seed height) are highly

conserved traits in shortgrass prairie, North America. Yet our study could not detect the benefit

of adding phylogenetic information using morphological traits to predict seed germination. The

inconsistent role of phylogeny in different ecosystems needs further exploration, especially

taking advantage of large standard databases of seed traits and the tree of life.

References

Agrawal AA. 2007. Macroevolution of plant defense strategies. *Trends Ecol Evol* **22**: 103–9.

Akaike H. 1998. Information Theory and an Extension of the Maximum Likelihood Principle. In: Parzen
E, Tanabe K, Kitagawa G (Eds). Selected Papers of Hirotugu Akaike. New York, NY: Springer New
York.

Barak RS, Lichtenberger TM, Wellman-Houde A, *et al.* 2018. Cracking the case: Seed traits and
phylogeny predict time to germination in prairie restoration species. *Ecol Evol* **8**: 5551–62.

Baskin CC and Baskin JM. 1998. Seeds: ecology, biogeography, and evolution of dormancy and
germination. Academic press.

Baum DA and Smith SD. 2013. Tree thinking: an introduction to phylogenetic biology. Roberts
Greenwood Village (CO).

Blomberg SP and Garland T. 2002. Tempo and mode in evolution: phylogenetic inertia, adaptation and
comparative methods. *J Evol Biol* **15**: 899–910.

Blomberg SP, Garland T Jr, and Ives AR. 2003. Testing for phylogenetic signal in comparative data:
behavioral traits are more labile. *Evolution* **57**: 717–45.

Bradford KJ. 2002. Applications of hydrothermal time to quantifying and modeling seed germination and
dormancy. *Weed Sci* **50**: 248–60.

Bu H-Y, Wang X-J, Zhou X-H, *et al.* 2016. The ecological and evolutionary significance of seed shape
and volume for the germination of 383 species on the eastern Qinghai-Tibet plateau. *Folia Geobot*
51: 333–41.

Chou Y-F, Cox RD, and Wester DB. 2012. Smoke Water and Heat Shock Influence Germination of

Shortgrass Prairie Species. *Rangeland Ecol Manage* **65**: 260–7.

Duncan C, Schultz N, Lewandrowski W, *et al.* 2019. Lower dormancy with rapid germination is an

important strategy for seeds in an arid zone with unpredictable rainfall. *PLoS One* **14**: e0218421.

Finch-Savage WE and Leubner-Metzger G. 2006. Seed dormancy and the control of germination. *New*

Phytol **171**: 501–23.

Finch-Savage WE and Leubner-Metzger G. 2006. Seed dormancy and the control of germination. *New*

Phytol **171**: 501–23.

Ganade G and Westoby M. 1999. Seed Mass and the Evolution of Early-Seedling Etiolation. *Am Nat* **154**:

469–80.

Garamszegi LZ. 2014. Uncertainties Due to Within-Species Variation in Comparative Studies:

Measurement Errors and Statistical Weights. In: Garamszegi LZ (Ed). Modern Phylogenetic

Comparative Methods and Their Application in Evolutionary Biology: Concepts and Practice.

Berlin, Heidelberg: Springer Berlin Heidelberg.

Hardegree SP, Moffet CA, Flerchinger GN, *et al.* 2013. Hydrothermal Assessment of Temporal

Variability in Seedbed Microclimate. *Rangeland Ecol Manage* **66**: 127–35.

Hipp AL, Larkin DJ, Barak RS, *et al.* 2015. Phylogeny in the service of ecological restoration. *Am J Bot*

102: 647–8.

Hladik A and Miquel S. 1990. Seedling types and plant establishment in an African rain forest.

Reproductive ecology of tropical forest plants **7**: 1990.

Howe HF and Smallwood J. 1982. Ecology of Seed Dispersal. *Annu Rev Ecol Syst* **13**: 201–28.

Jager M, Kaphingst B, Janse EL, *et al.* 2019. Seed size regulates plant dispersal distances in flowing

water (C Nilsson, Ed). *J Ecol* **107**: 307–17.

Jongejans E and Telenius A. 2001. Field experiments on seed dispersal by wind in ten umbelliferous

species (Apiaceae). *Plant Ecol* **152**: 67–78.

Keddy PA. 1992. A Pragmatic Approach to Functional Ecology. *Funct Ecol* **6**: 621–6.

Kidson R and Westoby M. 2000. Seed mass and seedling dimensions in relation to seedling

establishment. *Oecologia* **125**: 11–7.

Kiehl K, Kirmer A, Donath TW, *et al.* 2010. Species introduction in restoration projects – Evaluation of

different techniques for the establishment of semi-natural grasslands in Central and Northwestern

Europe. *Basic Appl Ecol* **11**: 285–99.

Kitajima K. 1996a. Ecophysiology of Tropical Tree Seedlings. In: Mulkey SS, Chazdon RL, Smith AP

(Eds). Tropical Forest Plant Ecophysiology. Boston, MA: Springer US.

Kitajima K. 1996b. Cotyledon functional morphology, patterns of seed reserve utilization and

regeneration niches of tropical tree seedlings. In "The ecology of tropical forest tree seedlings".(Ed.

MD Swaine) pp. 193--210.

Larson JE, Sheley RL, Hardegree SP, *et al.* 2015. Seed and seedling traits affecting critical life stage

transitions and recruitment outcomes in dryland grasses (M Cadotte, Ed). *J Appl Ecol* **52**: 199–209.

Laughlin DC. 2014. The intrinsic dimensionality of plant traits and its relevance to community assembly.

J Ecol **102**: 186–93.

Leishman MR, Wright IJ, Moles AT, *et al.* 2000. The evolutionary ecology of seed size. *Seeds: the*

ecology of regeneration in plant communities **2**: 31–57.

Martins EP and Hansen TF. 1997. Phylogenies and the Comparative Method: A General Approach to

Incorporating Phylogenetic Information into the Analysis of Interspecific Data. *Am Nat* **149**: 646–67.

Mazerolle MJ. 2020. AICcmodavg-package: Model Selection and Multimodel Inference Based on

(Q)AIC(c) in AICcmodavg: Model Selection and Multimodel Inference Based on

(Q)AIC(c)https://rdrr.io/cran/AICcmodavg/man/AICcmodavg-package.html. Viewed 30 Apr 2022.

Moles AT. 2018. Being John Harper: Using evolutionary ideas to improve understanding of global

patterns in plant traits. *J Ecol* **106**: 1–18.

Moles AT, Hodson DW, and Webb CJ. 2000. Seed Size and Shape and Persistence in the Soil in the New

Zealand Flora. *Oikos* **89**: 541–5.

Paradis E and Schliep K. 2019. ape 5.0: an environment for modern phylogenetics and evolutionary

analyses in R. *Bioinformatics* **35**: 526–8.

Revell LJ. 2010. Phylogenetic signal and linear regression on species data. *Methods in Ecology and*

Evolution **1**: 319–29.

Revell LJ. 2012. phytools: an R package for phylogenetic comparative biology (and other things).

Methods Ecol Evol **3**: 217–23.

Revell LJ, Harmon LJ, and Collar DC. 2008. Phylogenetic signal, evolutionary process, and rate. *Syst*

Biol **57**: 591–601.

Schurr FM, Bond WJ, Midgley GF, and Higgins SI. 2005. A mechanistic model for secondary seed

dispersal by wind and its experimental validation. *J Ecol* **93**: 1017–28.

Schwilk DW and Zavala N. 2012. Germination response of grassland species to plant-derived smoke. *J*

Arid Environ **79**: 111–5.

Sheldon JC and Burrows FM. 1973. The dispersal effectiveness of the achene-Pappus units of selected

compositae in steady winds with convection. *New Phytol* **72**: 665–75.

Subbiah A, Ramdhani S, Pammenter NW, *et al.* 2019. Towards understanding the incidence and

evolutionary history of seed recalcitrance: An analytical review. *Perspect Plant Ecol Evol Syst* **37**:

11–9.

Tooren BF van. 1988. The Fate of Seeds after Dispersal in Chalk Grassland: The Role of the Bryophyte

Layer. *Oikos* **53**: 41–8.

Umaña MN, Arellano G, Forero-Montaña J, *et al.* 2020. Large- and small-seeded species have contrasting

functional neighborhoods in a subtropical forest. *Ecosphere* **11**: 314.

Venables WN and Ripley BD. 2002. Modern Applied Statistics with S. Springer, New York, NY.

Westoby M, Falster DS, Moles AT, *et al.* 2002a. Plant ecological strategies: some leading dimensions of

variation between species. *Annu Rev Ecol Syst*: 125–59.

Westoby M, Falster DS, Moles AT, *et al.* 2002b. Plant Ecological Strategies: Some Leading Dimensions

of Variation between Species. *Annu Rev Ecol Syst* **33**: 125–59.

Willis CG, Baskin CC, Baskin JM, *et al.* 2014. The evolution of seed dormancy: environmental cues,

evolutionary hubs, and diversification of the seed plants. *New Phytol* **203**: 300–9.

Zanne AE, Tank DC, Cornwell WK, *et al.* 2014. Three keys to the radiation of angiosperms into freezing

environments. *Nature* **506**: 89–92.

Zhu J, Buchmann CM, and Schurr FM. 2022. Effects of seed morphology and orientation on secondary

seed dispersal by wind. *J Plant Ecol*.

Zhu J, Liu M, Xin Z, *et al.* 2019. A trade-off between primary and secondary seed dispersal by wind.

Plant Ecol **220**: 541–52.

Table 1 Forty-five native species were selected in this study, which are commonly involved in

restoration practice and range management in shortgrass prairie. Most of the species were bought

from Native American Seed, tested in controlled environments, 6 species were cited from (Chou

et al. 2012)[1] and 5 species were cited from (Schwilk and Zavala 2012)[2].

Species	Family
Andropogon gerardii Vitman	Poaceae
Argemone albiflora Hornem.	Papaveraceae
Aristida purpurea Hutt.	Poaceae
Asclepias asperula (Decne.) Woodson	Asclepiadaceae
Astragalus crassicarpus Nutt.[1]	Fabaceae
Bouteloua curtipendula (Michx.) Torr.	Poaceae
Bouteloua gracilis (Willd. ex Kunth) Lag. ex Griffiths	Poaceae
Callirhoe involucrata (Torr. & A. Gray) A. Gray	Malvaceae
Callirhoe leiocarpa R.F. Martin	Malvaceae
Centaurea americana Nutt.	Asteraceae
Chasmanthium latifolium (Michx.) Yates	Poaceae
Chloris cucullata Bisch.	Poaceae
Coreopsis lanceolata L.[2]	Asteraceae
Coreopsis tinctoria Nutt.	Asteraceae
Corydalis curvisiliqua Engelm.	Fumariaceae
Desmanthus illinoensis (Michx.) MacMill. ex B.L. Rob. & Fernald	Fabaceae
Digitaria californica (Benth.) Henr.	Poaceae
Digitaria ciliaris (Retz.) Koeler	Poaceae
Echinacea angustifolia DC.[2]	Asteraceae
Eragrostis trichodes (Nutt.) Alph. Wood	Poaceae
Eryngium leavenworthii Torr. & A. Gray	Apiaceae

Gutierrezia sarothrae (DC.) A. Gray	Asteraceae
Helianthus annuus L.	Asteraceae
Herbertia lahue (Molina) Goldblatt	Iridaceae
Ipomopsis rubra (L.) Wherry	Polemoniaceae
Liatris mucronata Hook. var. *mucronata* (DC.) B.L. Turner[2]	Asteraceae
Linum rigidum Pursh	Linaceae
Monarda citriodora Cerv. ex Lag.	Lamiaceae
Oenothera rhombipetala Nutt. ex Torr. & A. Gray	Onagraceae
Pavonia lasiopetala Scheele	Malvaceae
Penstemon cobaea Nutt.[2]	Scrophulariaceae
Phacelia congesta Hook.	Hydrophyllaceae
Phytolacca americana L.	Phytolaccaceae
Polytaenia nuttallii DC.	Apiaceae
Ratibida columnifera (Nutt.) Wooton & Standl.	Asteraceae
Rivina humilis L.	Phytolaccaceae
Salvia azurea Michx. ex Lam.	Lamiaceae
Salvia coccinea P.J. Buchoz ex Etlinger[2]	Lamiaceae
Salvia farinacea Benth.	Lamiaceae
Salvia lyrata L.	Lamiaceae
Schizachyrium scoparium (Michx.) Nash	Poaceae
Sorghastrum nutans (L.) Nash	Poaceae
Sporobolus airoides (Torr.) Torr.	Poaceae
Sporobolus cryptandrus (Torr.) A. Gray	Poaceae
Tradescantia occidentalis (Britton) Smyth	Commelinaceae

Figure 1 Phylogenetic tree of species and species seed traits values (seed mass, seed height)

distribution along the phylogenetic tree. Phylogenetic tree was generated from the pruned Zanne

et al. tree (Zanne *et al.* 2014), including 15 species (*Astragalus crassicarpus*, *Argemone*

albiflora, *Asclepias Asperula*, *Callirhoe leiocarpa*, *Centaurea americana*, *Chasmanthium*

tifolium, *Corydalis curvisiliqua*, *Digitaria californica*, *Eragrostis trichodes*, *Herbertia lahue*,

Linum rigidum, *Pavonia lasiopetala*, *Polytaenia nuttallii*, *Tradescantia occidentalis*, *Liatris*

mucronata) were placed within under the same genus/family. The center of each plot is the mean

value, the other two lines are -/+ standard errors. The colors were coded corresponding to the

grouping of phylogenetic positions (Figure 2).

Figure 2 Phylogenetic position of 45 species, represented by family, were clustered in three

groups. The phylogenetic positions were generated from paired-wise distances of species on the

phylogenetic tree (see Figure 1). The nonmetric multidimensional scale (NMDS) was applied, at

the stress of 17.86, displayed in two axes. For the convenience of display the phylogenetic

positions were grouped and color coded by vision.

571   Table 2 Phylogenetic signal was tested in seed morphological traits and overall seed final

572   germination percentage. Blomberg's K was used to evaluate phylogenetic signals (Blomberg et

573   al. 2003).

Trait	Blomberg's K	P-value
Seed Mass*	**0.07**	**0.01**
Seed Height *	**0.05**	**0.05**
Seed Surface Area	0.03	0.14
Seed Final Germination Percentage	0.02	0.20

575   K = 1, the traits is perfectly fit with Brownian motion model

576   K > 1, the traits is more conserved than expected comparing to Brownian motion model

577   K < 1, the traits is less conserved than expected comparing to Brownian motion model

578   * indicate traits containing phylogenetic signal (P =< 0.05)

Figure 3. The distribution of Blomberg's K along the size of the species pool in random

subsampling tests. The species were resampled 100 times from 10 species to 40 species, and

phylogenetic signal (Blomberg's K) was calculated for each trait, 3100 times for each trait. The

dots represent the Blomberg's K value of each resampling pool. The color of dots indicates the

p-value of Blomberg's K ($p < = 0.05$, black; $p > 0.05$, grey).

Figure 4 The proportion of subsamples with significant phylogenetic signal along the change of

number of species in species pools. The species were resampled 100 times from 10 species to 40

species. The dots represent the proportion of Blomberg's K value in each resampling pool.

Figure 5 Shiny application of Interactive Learning of Phylogenetic Comparative Methods. This

is a screenshot of the shiny application. The checkbox of species could be used to choose

different combinations of species and explore its impact on phylogenetic signals.

Figure 6 General protocol for generating a germinability prediction model with phylogenetic

information for a plant community. This model needs a pool species with phylogenetic

information, morphological data and germination data to build. It will be able to explore the

germination pattern of the community.

Glossary

Phylogeny / Phylogenetic tree: branching evolutionary histories / to graphs that represent these evolutionary histories. Phylogenetic tree including gene tree and species tree. In this paper, we only refer to species' tree (Baum and Smith 2013).

Phylogenetic conservatism: the hypothesis that closely related species share more traits than distantly related species. (Agrawal 2007).

Phylogenetic position: the relative position between species commonly used nearest neighbor and paired-wise distance. We used paired-wise distance in our calculation.

Phylogenetic signal: to describe a tendency for evolutionarily related organisms, under assumption of following a certain evolutionary model, to resemble each other. (Blomberg *et al.* 2003).

Phylogenetic residual: incorporate the phylogeny through error structure, such as estimating ancestral states, rates of evolution, phylogenetic effects. (Garamszegi 2014).

629   **Supplemental Material**: Model Evaluation for all seed germinability prediction models. AICc is

630   the adjusted AIC value due to the small sample size in biological tests.

Interactive Learning of Phylogenetic Comparative Methods

Which species do you want to using for building phylogeny?

- ☑ Eryngium_leavenworthii
- ☑ Polytaenia_nuttalli
- ☑ Asclepias_asperula
- ☑ Centaurea_americana
- ☑ Coreopsis_lanceolata
- ☑ Coreopsis_tinctoria
- ☑ Echinacea_angustifolia
- ☑ Gutierrezia_sarothrae
- ☑ Helianthus_annuus
- ☑ Liatris_mucronata
- ☑ Ratibida_columnifera
- ☑ Tradescantia_occidentalis
- ☑ Astragalus_crassicarpus
- ☑ Desmanthus_illinoensis
- ☑ Corydalis_curvisiliqua
- ☑ Phacelia_congesta
- ☑ Herbertia_lahue
- ☑ Monarda_citriodora
- ☑ Salvia_azurea
- ☑ Salvia_coccinea
- ☑ Salvia_farinacea
- ☑ Salvia_lyrata
- ☑ Linum_rigidum

This is a shiny application to help user to experience the phylogenetic comparative method. The left hand side allow user to select species. Based on species selection, the display tab will have display related phylogenetic tree, phylogenetic position, phylogenetic signal and model selection results

Please try multiple combination of species and see differences between results of phylogenetic signal calculation, the user could understand the sampling of species is critical for phylogenetic signal estimation for the community

| Phylogenetic_Tree | Phylogenetic_Position | Phylogenetic_Signal | Model_Selection |

Phylogenetic position of selected species, which are calculated based on phylogenetic tree.

Figure 6 General protocol for generation a germination prediction model with phylogenetic information for a plant community. This model needs a pool species with phylogenetic information, morphological data and germination data to build. It will be able to explore the germination pattern of the community.