Evolutionary innovation through transcription factor promiscuity in microbes is constrained by pre-existing gene regulatory network architecture

Authors: Matthew J. Shepherd[1], Aidan P. Pierce[1], Tiffany B. Taylor[1]*

1 - Milner Centre for Evolution, Department of Life Sciences, University of Bath, Claverton Down, Bath BA2 7AY, UK

* - Corresponding author: Tiffany B. Taylor, Milner Centre for Evolution, Department of Life Sciences, University of Bath, Claverton Down, Bath BA2 7AY, UK, +44 (0)1225 384398, t.b.taylor@bath.ac.uk

The contributions to this work were as follows:

Experimental work performed by MJS. Rhamnose expression plasmids built by APP. Manuscript written by MJS. Manuscript editing by all authors. Project supervised by TBT. Funding secured by TBT.

Acknowledgements:

This work was funded by a Royal Society Research Fellows Grant (RG160491; awarded to TBT) supporting MJS; Royal Society Research Fellows Enhancement Award (RGF\EA\201057; awarded to TBT) supporting APP; a Royal Society Dorothy Hodgkin Research Fellowship (DH150169; awarded to and supporting TBT). Bioinformatics analyses for the article was carried out using the Medical Research Council's (MRC) Cloud Infrastructure for Microbial Bioinformatics (CLIMB), Illumina Whole-Genome Sequencing conducted by MicrobesNG, Birmingham, UK (http://www.microbesng.com) which was supported by the BBSRC (grant number BB/L024209/1), and Illumina RNA sequencing with ribodepletion was conducted by Oxford Genomics Centre, Oxford UK. We would like to thank Prof. Laurence Hurst for insightful discussion of earlier manuscript versions, Dr. Jonathan Nzakizwanayo and Dr. Iain MacArthur for providing us with *E.coli* strains SP50 pRK2073, S17 pTNS2 and ST18, Dr. Ellie Harrison for providing the R script that Fig.5A is based upon, and Dr. James Horton, Louise Flanagan and Josie Elliott for helpful discussion during the course of the study.

Data availability:

All raw data for this study is available on the Open Science Framework (OSF), and can be accessed at https://osf.io/pcdhx/

Abstract

The survival of a population during environmental shifts depends on whether the rate of phenotypic adaptation keeps up with the rate of changing conditions. A common way to achieve this is via change to gene regulatory network (GRN) connections – known as rewiring – that facilitate novel interactions and innovation of transcription factors. To understand the success of rapidly adapting organisms, we therefore need to determine the rules that create and constrain opportunities for GRN rewiring. Here, using an experimental microbial model system we reveal a hierarchy among transcription factors that are rewired to rescue lost function, with alternative rewiring pathways only unmasked after the preferred pathway is eliminated. We identify three key properties - high activation, high expression, and pre-existing low-level affinity for novel target genes – that facilitate transcription factor innovation via gain of functional promiscuity. Ease of acquiring these properties is constrained by pre-existing GRN architecture, which was overcome in our experimental system by both targeted and global network alterations. This work reveals the key properties that determine transcription factor evolvability, and as such, the evolution of GRNs.

Introduction

During conditions of environmental upheaval and niche transition events, rapid phenotypic adaptation is essential for evolutionary success[1–3]. To understand patterns of diversification in novel environments we need to understand why some evolutionary transitions occur more rapidly than others, and what allows some organisms to succeed where others fail. Key to this is understanding the evolution of gene regulatory networks (GRNs), control circuits common throughout the domains of life that determine the magnitude and timing of gene expression[4] in response to environmental and internal signals[5,6]. GRNs are frequent sites of adaptive mutation driving phenotypic evolution[7–9], and often underscore adaptation to - and survival in - new and changeable environments[10–13]. Determining how these networks evolve and identifying rules governing when and how a regulatory circuit adapts will allow us to better understand their role in determining the evolutionary success of an organism.

Within GRNs, a key mechanism of innovation is transcription factor rewiring, in which transcription factors can gain or lose regulatory connections to target genes creating new network architectures and opportunity for phenotypic innovation[14]. Rewiring

events can have dramatic effects on the transcriptome[15] that can drive phenotypic

diversification. However, the majority of past studies on transcription factor rewiring

involve retrospective experimental dissection of networks that have already

diverged[16–21], which allows inference of past rewiring events but does not address

the evolutionary factors driving the rewiring process. For rewiring to occur, a

transcription factor must first have the potential for promiscuous activity (interaction

with non-cognate regulatory targets) allowing regulation of a new gene[22], as is the

case for innovation in other proteins[23,24]. Promiscuity is also referred to as cross-talk,

however this term can describe cognate interactions where transcription factors have

overlapping regulons, so we avoid it in this study. These non-cognate interactions

can become meaningful and drive adaptation if favoured by natural selection[25,26].

However, promiscuity cannot be commonplace within GRNs as this will likely result

in dysregulation of gene expression and fitness costs for an organism in the

environment to which it is adapted[27,28].

Whilst there exists some empirical evidence of the importance of transcription factor

promiscuity in driving evolutionary innovation[21,29,30], in each case it is unknown why

the transcription factor in question became promiscuous as opposed to any of the

other regulators within their protein family; what role – if any – GRN structure played

in the process; and whether the rewired transcription factors were unique in their

ability to become promiscuous within each study system. To address these

questions, we set out to investigate the evolution of transcription factor promiscuity

using rescue of flagellar motility via rewiring of NtrC documented by Taylor et al.

(2015b). In this model system, *Pseudomonas fluorescens* SBW25 is engineered to

be non-motile via deletion of the master regulator for flagellar synthesis (*fleQ*) and

abolishment of biosurfactant production. Under strong selection for motility, the

bacteria rapidly and reliably evolve new regulatory network wiring to rescue flagellar

motility (a schematic of these experiments is laid out in Fig.1A)[31] with the same

transcription factor (*ntrC*) repurposed to rescue flagellar driven motility each time, to

the exclusion of all other homologous transcription factors within the protein family[32].

However, we do not know the factors that determine this evolutionary preference. To

test this, we constructed a double *fleQ ntrC* knockout and placed this mutant under

strong selection for swimming motility. This forces evolutionary utilisation of an

alternative transcription factor in order to rescue motility, and is an established

method for unveiling hidden evolutionary pathways[33]. We reveal a hierarchy among
transcription factors that are rewired to rescue lost flagellar function, with alternative
rewiring pathways only revealed after the preferred co-opted transcription factor,
NtrC, is eliminated. Identification of additional transcription factors capable of gaining
promiscuity within the same GRN background allows us to investigate rules
governing when and where a transcription factor becomes promiscuous within its
GRN, and is important for understanding how these regulatory systems innovate
during adaptation to environmental challenges.
111

Results
Evolutionary rescue of flagellar motility can occur in the absence of _ntrC_,
through _de-novo_ mutation to an alternative two-component system
Is the reason we only see NtrC co-opted to rescue FleQ function because it is the
only transcription factor capable of doing this? It is known that within families of
homologous transcription factors there is variation in the ability to bind non-cognate
sites[34], so it is possible that NtrC is unique in its ability to become promiscuous and
rewire. FleQ and NtrC are part of a family of 22 structurally related transcription
factors called R̲poN-dependent e̲nhancer b̲inding p̲roteins (RpoN-EBPs), many of
which are predicted to be more structurally similar to FleQ than NtrC[32]. To identify if
any other RpoN-EBPs were capable of promiscuity, the double knockout non-motile
P. fluorescens (ΔfleQΔntrC) was challenged to rescue flagellar motility in 0.25% agar
LB plates. Motile zones re-emerged in a two-step manner (a slow-swimming variant,
then a faster-swimming variant) as typical of previous studies[31]. Motile isolates were
sampled and whole genome re-sequenced at each step. Motility-granting mutations
were identified in the gene _PFLU1131_ for all first-step motile isolates (n=15), and in
13/15 cases this was the only mutant gene (Supplementary excel file E1). _PFLU1131_
is unstudied and encodes a putative sensor histidine kinase. The gene is situated in
an operon between two other genes, _PFLU1130_ and _PFLU1132_ which encode a
hypothetical GNAT-acetyltransferase, and a putative RpoN-dependent transcriptional
regulator respectively. PFLU1132 is a known FleQ-homolog[32] and together with
PFLU1131 forms a putative two-component system. The most frequent mutation in
PFLU1131 was an identical 15bp deletion (Fig.2A) in the histidine-kinase phospho-
acceptor domain (73%). This mutation results in loss of 5 amino acids (368-GEVAM-
372) in the protein product (henceforth referred to as PFLU1131-del15). Other

mutations included a highly similar 15bp deletion resulting in loss of 5 amino acids a

few amino acids downstream (369-EVAMG-373), as well as a SNP resulting in

A375V. All of these mutations (86% of the first-step mutations) cluster to the same

26bp of the 1770bp *PFLU1131* ORF, and result in amino acid changes in a site

directly adjacent to the catalytically active H-box[35] (amino acids 376-382, Fig.2A)

suggesting a significant effect on the catalytic function of the putative kinase. We

additionally constructed a *PFLU1132* knockout in the *ΔfleQΔntrC* background to

identify any further RpoN-EBPs capable of promiscuity. *ΔfleQΔntrCΔPFLU1132* failed

to rescue flagellar motility within the 6-week assay cut-off (Supplementary excel file

E2, n= 192), suggesting evolutionary gain of promiscuity in other transcription factors

may not be easily achieved under the conditions we tested.

Together, these results suggest that in the absence of FleQ and NtrC, flagellar

motility can be rescued via rewiring of an alternative transcription factor in the same

protein family, PFLU1132. To confirm that observed motility phenotypes were

dependent on the PFLU1131/2 two-component system, the *PFLU1132* gene was

deleted and complemented with and without the presence of the most common first-

step kinase mutation (*PFLU1131*-del15). Knockout of *PFLU1132* abolished flagellar

motility, and complementation restored motility only in the presence of the kinase

mutation (Fig.2B). This experiment was repeated for *ntrBC* and produced the same

result (Supplementary Fig.S1). The *PFLU1131*-del15 mutation also grants flagellar

motility when *ntrC* is present (i.e., in a *ΔfleQ* background), so does not depend on

the absence of *ntrC* to function (Supplementary Fig.S2). Transcriptomic analysis of

the *PFLU1131*-del15 mutant by RNA sequencing indicates an alteration to the

activity of the PFLU1131/2 two-component system that results in a net up-regulatory

effect on the transcriptome (Fig.2C). These regulatory changes include upregulation

of the flagellar genes (Supplementary Fig.S4) and the *PFLU1130/1/2* operon. In

sum, in the "commonly rewired route", repeatable mutations in the NtrBC (Fig.1B)

two-component system were found to rescue flagellar motility. In the absence of both

FleQ and NtrC, we identified the "unmasked rewiring route", via PFLU1131/2

(Fig.1C), that was also capable of rescuing flagellar motility.

First-step mutations in the unmasked rewiring pathway grant promiscuous

activity to a FleQ homologous transcription factor

In the common rewiring pathway, NtrC, a homolog of FleQ, was recruited to recover

lost FleQ function. This occurred as a two-step process; an initial motility-granting

mutation to the gene *ntrB* encoding NtrC's cognate kinase and a secondary motility-

enhancing mutation to the helix-turn-helix (HTH) DNA binding domain of the NtrC

transcription factor. Similarly, In the Δ*fleQ*Δ*ntrC* background strain, first-step

mutations restored slow motility, showing that alternative rewiring routes were

possible, but not utilised in the presence of *ntrC*. To determine the mechanism of this

unmasked rewiring route we performed whole genome sequencing and

transcriptome analysis on ancestral and evolved isolates. In 13/15 first-step mutants,

only a single mutation to *PFLU1131* was identified. To confirm that no other

mutations were needed for flagellar motility, the *PFLU1131*-del15 mutation was

introduced into the ancestor (Δ*fleQ*Δ*ntrC*). The resulting engineered strain was motile

(Supplementary Fig.S2). As no mutational change to the DNA binding domain of

PFLU1132 is needed, this suggests a non-specific mechanism through which this

transcription factor induces the flagellar genes.

To investigate this, our transcriptomic data was used to assess the impact of the

PFLU1131-del15 mutation on the expression of a list of genes[36] controlled by all 22

homologous transcription factors (part of the RpoN-EBP family) present in *P.*

fluorescens SBW25[37]. Regulatory changes predictably include upregulation of the

flagellar genes (Supplementary Fig.S3) and the *PFLU1130/1/2* operon, but also

include upregulation of many other genes. In particular, we saw significant

upregulation for 54% of all RpoN-EBP controlled genes in the unmasked rewiring

pathway (Fig.3). For comparison, 70% of RpoN-EBP controlled genes were

upregulated in the common rewiring pathway (via NtrC)[34]. Flagellar genes account

for 12% of all RpoN-EBP controlled genes, and whilst the native regulatory targets of

PFLU1132 are unknown it is unlikely that it natively regulates 54% of these.

Similarly, the first step motile mutant reported in Taylor et al., (2015b) leads to

upregulation of genes known to be involved in nitrogen assimilation (as expected as

mutations are located in the Ntr pathway), however it also results in upregulation of

many genes with no known regulatory link to NtrC. The large percentages of the

RpoN regulon induced by these specific kinase mutations, in both the common and

unmasked rewiring pathways, suggest first-step mutations confer a state of

promiscuous regulatory activity to PFLU1132 resulting in non-cognate interactions

across many RpoN-EBP controlled genes including the flagellar genes.

Evolutionary rescue of motility through unmasked rewiring pathway is

significantly constrained

If alternative rewiring routes are available to natural selection – why have we not seen

them utilised in the presence of the commonly rewired route? In the unmasked rewiring

pathway, we considered the time taken for swimming mutants to evolve and the

strength of the evolved phenotype in comparison to the common rewiring route. Time

to emergence (the length of time taken for motility to evolve on the soft agar plate) was

recorded for all replicate experimental lines of the Δ*fleQ*Δ*ntrC* ancestor, as well as

Δ*fleQ* as a comparison. Strikingly, only 9.4% of replicate experimental lines (n=160) of

Δ*fleQ*Δ*ntrC* evolved within the 6-week assay cut-off compared to 100% for the Δ*fleQ*

background (n=22). Initial first-step motile Δ*fleQ*Δ*ntrC* strains also evolved within an

average of 18.7 days from the assay start (Fig.4A), significantly longer than Δ*fleQ* took

to rescue motility (4.2 days, P = 0.002, Dunn test). After first-step PFLU1131

mutations, second-step mutants evolved rapidly in Δ*fleQ*Δ*ntrC* lines within an average

of 2.2 days (Fig.4A). This was faster than the second-step mutants in the Δ*fleQ*

background which took 3.3 days to emerge (*P* = 0.0017, Dunn test).

The greatly increased time, and low frequency for rescue motility in an Δ*fleQ*Δ*ntrC*

background may be reflective of a small pool of accessible PFLU1131 mutations that

can grant promiscuity to PFLU1132, perhaps due to a small mutational target size.

One replicate line gained a *de novo* mutation in the DNA mismatch-repair gene *mutS*

(Supplementary excel file E1), resulting in a frame shift and probable loss of

function[38]. This strain, along with strains derived from it possessed large numbers of

additional SNPs (>80) including a SNP in PFLU1131. Loss of function to *mutS* is

known to enhance mutability in *Pseudomonads*[39]*,* and its occurrence may increase

access to motility-rescuing mutations in PFLU1131.

From a phenotypic perspective, the first-step mutations in the *PFLU1131* pathway in

a Δ*fleQ*Δ*ntrC* strain, provide a far poorer motility phenotype than the analogous first-

step mutations in a Δ*fleQ* background (Fig.4B). In a race assay, *PFLU1131*-del15

mutants swam 0.31 mm for every 1 mm swam by the most common first step mutant

in the common rewiring route (P=0.000356, Wilcox test). One possible explanation

for this poor motility is a lack of significant upregulation for the flagellar filament

subunit FliC (Supplementary Fig.S4) in the *PFLU1131*-del15 mutant.

The *PFLU1131*-del15 mutant additionally does not confer a significant defect to

growth in shaking LB broth compared to its ancestral strain (Fig.4C, average relative

area under the growth curve (rAUGC) of 1.55 and 1.48 respectively, P=0.1132, Dunn

test), and grew significantly better than the most common first step mutant in the

common rewiring route (rAUGC of 1, P=0.0081, Dunn test). The poor motility

phenotype and the lack of significant fitness cost associated with *PFLU1131*-del15

likely indicates that the PFLU1131/2 system is a weaker activator of the flagellar

motility that incurs less sever pleiotropy compared to NtrBC.

These findings indicate that the unmasked pathway is the far poorer option for

gaining promiscuous regulation of the flagellar genes compared to the common

NtrBC pathway - motility rescue in the Δ*fleQ*Δ*ntrC* background took significantly

longer to occur, was significantly less frequent, and provides a significantly poorer

flagellar motility phenotype with lower pleiotropic fitness cost. However, it is not clear

why this should be the case – both NtrC and PFLU1132 are FleQ-homologs, and are

gaining mutations to their cognate kinases that unlock promiscuity. This suggests

that there are other factors that constrain the evolution of promiscuity in PFLU1132.

Motility enhancing second-step mutations suggest alterations to global gene

regulatory network can facilitate promiscuity in unmasked rewiring pathway

Following the first-step mutations in PFLU1131 that unlock flagellar motility, our

motile isolates develop second-step mutations that act to boost motility speed

(Supplementary Fig.S5A), often accompanied by significant pleiotropic fitness costs

for growth in LB broth (Fig. S5B). These second-step mutations offer clues to the

nature of constraining factors limiting innovation through promiscuity of PFLU1132,

as they represent evolutionary solutions to the poor motility provided to the first-step

mutations.

Whole genome resequencing identified a diverse set of motility enhancing second-

step mutations occurring at both a local (PFLU1131/2 locus) and global regulatory

scale (Fig.5A, full details of all mutations provided in supplementary excel file E1,

second step n = 18 (several first step lineages generates multiple second step

zones). In contrast to the frequent second-step mutations observed in the HTH DNA-

binding domain of NtrC[31], no analogous mutations were observed in the same

domain of PFLU1132. Mutations that did occur in the *PFLU1132* gene were parallel

and identified in only 2 lines, impacting the receiver domain of the transcription

factor, which may further increase activity of the PFLU1132 regulator by modifying

the interactions with its kinase. One line gained a secondary *PFLU1131* mutation, a

SNP in the HATPase domain along with the first-step SNP A547D already present

(Fig.2B), which may further boost kinase interactions. A mutation to the promoter

region of the PFLU1131/2 operon was also observed.

Other secondary mutations can be grouped into 3 other broad categories. (i) The first

are mutations in the operon *PFLU1583/4*, accounting for 24% of secondary

mutations (Fig.5A, Purple). This gene pair encodes a putative anti-sigma factor

PP2C-like phosphatase and a putative STAS family anti-sigma factor antagonist

(*PFLU1583* and *PFLU1584* respectively). Mutations in *PFLU1583* were generally

loss of function suggesting loss of repression on an unknown sigma factor. This may

act on RpoN the partner sigma factor of PFLU1132 in regulating gene expression,

however this cannot be tested easily through *rpoN* knockout, and *P. fluorescens*

SBW25 encodes ~31 other putative sigma factors[37] which may instead be the

mechanistic targets of this mutation. Transcriptomic analysis of a *PFLU1583* mutant

(PFLU1583Δ48-74) identified sigma factor RpoE as upregulated, however *rpoE*

knockout did not negate the motility enhancing effect of the mutation (Supplementary

Fig.S6A). To test that PFLU1583/4 are not acting to enhance motility through

another RpoN-EBP, we engineered *PFLU1583*Δ48-74 in a Δ*fleQ*Δ*ntrC* background in

the absence of a *PFLU1131* mutation. This strain was immotile, indicating that these

anti-sigma factor mutations depend on the function of the PFLU1131/2 system to

confer a swimming phenotype (Supplementary Fig.S6B). (ii) The second category of

mutations impact core gene expression components that will affect global GRN

function (Fig.5A, Green). These included mutations to *rho*, *rpsK* and *rpoC* that

encode core gene expression machinery and will impact expression of most genes in

the genome. (iii) The final category of motility enhancing mutations fit a metabolic

theme (Fig.5A, blue). These mutations occur in genes involved in central carbon

metabolism, which will likely have a significant impact on global gene expression
through alterations to central carbon flux and may act on RpoN indirectly – for
example via core carbon metabolic regulator CbrB (an RpoN-EBP) – or signal to the
PFLU1131 sensor-kinase via internal metabolic flux.
The diversity and nature of these second-step mutations suggest global regulatory
strategies to facilitate flagellar gene expression through the unmasked rewiring
pathway. To understand their regulatory impact, transcriptome analysis was
performed on a pair of representative mutants: the most common second-step
mutant (anti-sigma factor mutant, *PFLU1583* Δ48-74) which represented a mutation
with global regulatory effects, and the *PFLU1131/2* promoter mutation as a
representative of mutation with predicted "local" regulatory effects. Together these
mutations represent the two loci that constitute 56% of identified second-step
mutations.
Principle component analysis indicates that the transcriptomic profile of the anti-
sigma factor mutant differs significantly from the profile of the *PFLU1131/2* promoter
mutation. Both mutations resulted in similar variation across PC1 relative to the first-
step *PFLU1131*-del15 mutant, but opposite variation in PC2 (Fig.5B). Both the
PFLU1131/2 promoter and anti-sigma factor mutations result in net up-regulatory
effects on the transcriptome reflected by positive skewed volcano plots (Fig.5C and
Fig.5D) albeit with differing expression patterns. Both mutations have the effect of
further upregulating RpoN-EBP controlled genes with 22% and 60% being
upregulated for the anti-sigma factor mutant and the *PFLU1131/2* promoter mutant
respectively (Fig.5E).
In general, for the unmasked rewiring pathway, genomic and transcriptomic analysis
of second-step mutations indicate that second-step mutations are increasing
promiscuous activity of the rewired transcription factor (PFLU1132), albeit through
differing (local and global) mechanisms. These results highlight that diverse
mutations with both targeted and global regulatory effects can influence promiscuity
to facilitate rewiring and innovation in a transcription factor.
330
Second-step promoter capture mutation suggests importance of transcription
factor expression in gaining promiscuity

Although second-step mutations were diverse in the unmasked rewiring pathway,

one promoter capture event in particular provides evidence for the role of increased

expression of the rewired transcription factor in strengthening the motility phenotype.

For promiscuous activation of non-cognate genes, the PFLU1132 transcription factor

will need to first saturate its native regulatory interactions before it can engage in

low-affinity promiscuous interactions with FleQ-controlled genes. High expression of

active transcription factor can provide these conditions, elevating the concentration

of transcription factor in the cell to permit more frequent promiscuous action[40,41].

Whilst many of the second-step mutations detailed above may impact *PFLU1131/2*

expression indirectly (RpoC, Rho), the *PFLU1131/2* promoter mutation directly

effects expression of the two-component system. This mutation is a 1.59Kbp deletion

resulting in total loss of the *PFLU1130* gene, and in *PFLU1131/2* becoming part of

the *PFLU1127/8/9* operon (Fig.6A). Upstream of this new combined operon sits a

predicted RpoN binding site[36], so this deletion positions *PFLU1132* in a new operon

under the control of RpoN which may create a positive feedback loop where this

RpoN-dependent regulator drives its own expression – either through promiscuous

regulation or native control of this promoter. This genetic rearrangement significantly

upregulates both *PFLU1131* and *PFLU1132* with 4.26 and 3.79 Log2Fold increases

in expression compared to the non-motile Δ*fleQ* strain respectively (Fig.6B, raw

values for this and other tested strains provided in supplementary excel file E5).

Upregulation of this two-component system corresponds with the 60% increase in

expression of RpoN-EBP controlled genes discussed above (Fig.5E), indicating an

increase in promiscuous transcription factor activity. The 1.59Kbp deletion results in

loss of an intergenic region that typically separates the two combined operons. This

region is predicted to contain a rho-dependent terminator (Supplementary Excel file

E3) which may typically prevent transcriptional readthrough into *PFLU1130/1/2*.

Such a terminator will restrict *PFLU1131/2* expression via readthrough and constrain

its ability to achieve high concentrations that may facilitate promiscuity. The *rho*

mutation observed in another second-step motile strain may also act to increase

readthrough at this site. In comparison to this promoter mutation, the other second-

step mutation tested (anti-sigma factor *PFLU1583* Δ48-74) does not significantly

impact *PFLU1131/2* expression (Fig.6B), so this mutation likely influences the

promiscuity of the *PFLU1132* transcription factor through a separate mechanism.

In summary, expression of *PFLU113/2* appears to be a major factor constraining promiscuous regulatory activity and motility rescue through this regulatory system, and overcoming this constraint facilitates evolutionary innovation through this unmasked rewiring route.

Increasing transcription factor expression in motile mutants results in faster motility phenotypes

To test the importance of transcription factor gene expression in gain of promiscuous activity, we made use of the complementation strains presented in Fig.2C and supplementary Fig.S1. These experiments produced strains where the common (*ntrC*) and unmasked (*PFLU1132)* rewired transcription factor genes are deleted from their native loci and reintroduced on an L-rhamnose titratable promoter system. This was done in the presence of their respective kinase mutations that conferred first-step slow spreading motility in each rewiring pathway. Concentration of L-rhamnose added to the media positively correlated with distance moved (Fig.7A) for both the common rewiring pathway (i.e., NtrC expression system; $\rho = 0.984$, Spearman's test) and the unmasked rewiring pathway (i.e., PFLU1132 expression system; $\rho = 0.865$, Spearman's test). We confirmed increasing L-rhamnose concentration increased expression from the L-rhamnose titratable system (RhaSR-PrhaBAD) in our bacterial strain backgrounds by testing LacZ activity with increasing L-rhamnose concentration for *lacZ* under control of the L-rhamnose titratable promoter construct in the Δ*fleQ* and Δ*fleQ*Δ*ntrC* backgrounds. In both cases LacZ activity positively correlated with L-rhamnose concentration ($\rho = 0.981$ and $\rho = 0.955$ respectively, Spearman's tests), indicating that increasing L-rhamnose concentration results in increasing expression of the gene introduced in the *rhaSR-PrhaBAD* expression system (Supplementary Fig.S7).

This allows us to conclude that increasing expression of *ntrC* and *PFLU1132* in the presence of their first-step kinase mutations granted stronger motility phenotypes, likely through stronger promiscuous action. The expression level of a transcription factor can therefore significantly impact its propensity for promiscuous regulatory activity, and more highly expressed transcription factor may be expected to engage in non-canonical regulatory interactions more readily.

Discussion

Essential to understanding why some organisms adapt faster than others in the face of environmental challenges is knowledge of how GRNs and regulatory systems evolve. Evolutionary innovation of transcription factors within these networks depends on gain of functional promiscuity as a first step, however the rules of when and where such properties develop within the context of the GRN is currently unknown. Using our microbial model system, we have identified two available pathways for rescuing flagellar motility in a maladapted Δ*fleQ* network; through mutations to the NtrBC and PFLU1131/2 two component systems. However, one pathway – the NtrBC pathway – is always used to the exclusion of the other when both are present. By comparing these available evolutionary pathways, we can resolve why NtrC is more accessible for gaining promiscuity and rewiring due to its position within its GRN. In addition, we can identify three key properties that facilitate transcription factor promiscuity and opportunities for evolutionary innovation. **Firstly, our results confirm the importance of shared structural homology between the absent and recruited transcription factor.** In our system, the shared structural homology of NtrC and PFLU1132 with FleQ grants a basal affinity for regulating flagellar genes that evolution can act upon. This likely explains why structurally unrelated transcription factors do not rewire in our model, and indicates that similarity to FleQ facilitates the gain of promiscuity observed in NtrC and PFLU1132. However it is not the case that the regulators that are most structurally homologous to FleQ are the first to be repurposed[32], so homology is not the deciding factor in which RpoN-EBPs become promiscuous to rescue flagellar motility. It is possible that the RpoN-EBPs represent a hyper-evolvable family of transcription factors through the nature of their connectivity and mechanism of transcriptional regulation via RpoN. To investigate this evolutionary rescue of a non-RpoN dependent trait would need to be tested using a different maladapted GRN rescue model. **Secondly, hyperactivation of the co-opted transcription factor benefits promiscuous regulatory activity**. In our model system, first step mutations that granted a slow motile phenotype were always achieved through mutation of a kinase – resulting in hyper-activation of their cognate transcription factors. Most transcription factors are subject to some form of post-translational control, most commonly through phosphorylation of a REC receiver domain in bacteria[42].

Promiscuous binding may occur transiently or only briefly, so ensuring that the

transcription factor is constantly activated increases the likelihood of these non-

cognate regulatory events leading to altered gene expression. **Finally - and**

perhaps most importantly - high gene expression of the transcription factor

boosts promiscuous activity. High expression of a transcription factor can lead to

saturation of native binding sites. Any unbound, active transcription factor is then

free to engage in promiscuous activity. These low affinity interactions will occur more

readily if there is a higher concentration of transcription factor present, following laws

of mass action kinetics[43]. Typically, transcription factors are expressed at low levels

in bacteria[44], due to only a small protein copy number being needed for a small

number of binding sites in most cases[45]. Transcription factor expression level may

well be optimised at a low level to prevent mis-regulation. Indeed, highly expressed

and connected transcription factors were found to result in greater network

perturbations when synthetically rewired[15], which may reflect a greater ability to act

promiscuously. Many molecular biology studies use overexpression/overactivation of

proteins of interest as a means to determining their biological functions[46]. Our

findings also suggest that this approach should perhaps be used more cautiously,

certainly when studying transcription factor function.

Aside from mutations directly targeting activation and expression of *PFLU1131/2*, we

observed several distinct strategies that enhance PFLU1132 promiscuity through

global regulator changes. These strategies likely impact the wider GRN and

transcriptome significantly, with mutations to RpoC, RpsK and Rho being particularly

striking. RpoC and RpsK mutations may act to increase expression of most genes

including *PFLU1131/2* and the flagellar genes. Mutations to the *PFLU1583/4* anti-

sigma factor system also indicated a strategy to boost promiscuity that would have a

significant impact of the wider GRN. Global regulators are often sites of mutation and

innovation[47,48], with mutations affecting sigma factor function in particular having

large impacts on global gene expression which can drive phenotypic innovation[49].

Changes to sigma factors have been shown to facilitate GRN remodelling, as was

the case for a study of *P. aeruginosa* during adaptation to the CF lung[13]. Our study

highlights for the first time that such mutational strategies can also achieve increases

in promiscuity for specific transcription factors to facilitate evolutionary innovation.

Each of these mutational strategies that enhanced functional promiscuity in

PFLU1132 acted to alter the pre-existing GRN architecture around the transcription

factor, whether at a local level by altering the *PFLU1131/2* promoter or at a global

regulatory level.

Our findings suggest that pre-existing GRN architecture can be a key constraining

factor in gaining functional promiscuity and evolutionary innovation by a transcription

factor. For example, the ease by which a transcription factor can gain high

expression depends on the GRN structure it sits within. Many transcription factors

promote their own expression[50] (positive autoregulation) and can generate high self-

expression through runaway positive feedback. Such positive feedback loops are

typically contained by a negative repressor[51–53], including in the case of NtrBC

(Fig.1B). Mutational loss of repression can therefore be an evolutionarily accessible

mechanism to achieve promiscuity-enhancing high expression levels of a

transcription factor. In our model system, GlnK prevents runaway *ntrBC*

autoregulation by inhibiting NtrB[54] (Fig.1B). All previously observed NtrBC pathway

mutations were single *de-novo* mutations that act to remove repression through

GlnK, both directly or indirectly[31,55,56]. These have a dual effect, both hyperactivating

the NtrBC system, and strongly upregulating *ntrBC* expression through their positive

autoregulatory loop. In contrast, we have identified in this study that to achieve the

same effect of high activity and high expression in PFLU1131/2, multiple mutations

were required (Fig.1C). This highlights the clear evolutionary advantage NtrBC has

in becoming promiscuous compared to PFLU1131/2.

From the evidence outlined, pre-existing GRN motifs can be predicted to significantly

bias the evolvability of the transcription factors under their control. This is strongly

supported by the literature: mutational loss of a negative repressor leading to

overactivity and overexpression of a positively autoregulated transcription factor has

been observed to drive adaptation in several experimental and clinical settings[57–59].

Considering the alternative scenario, a transcription factor that represses its own

expression (negative autoregulation) will hinder the gain of a high expression level,

due to the nature of self-repression. Interestingly, the hierarchy favouring NtrBC over

PFLU1132 matches previous studies of evolutionary pathway heirarchies[33], where

mutational target size predicts that loss of repressors will be favoured over other

precise mutations. Our study highlights that in the context of GRN structures,

connectivity to other network components can constrain innovation through gain of

promiscuity, alongside other effects including mutational target size.

Our study provides empirical evidence for mechanisms and evolutionary drivers of

regulatory evolution. GRNs are key to the adaptability of organisms, and our results

reveal the genetic regulatory factors that generate evolvable control systems. We

find that feedback loops that can generate high expression of regulators after a small

number of mutations can facilitate rapid innovation and adaptation. Previously, we

have hypothesised that some transcription factors may be 'primed' for evolutionary

innovation by virtue of the GRN architecture they exist within[26]. Our data support this

by suggesting that promiscuity can be more easily evolved in transcription factors

with positive feedback structures. These findings can also be utilised in the design of

genetic regulatory systems to have reduced evolvability, a key future goal for

synthetic biology projects where the evolutionary change of a constructed system is

detrimental[60,61]. Designing systems that avoid easily accessible routes to

hyperactivation and hyperexpression of their components could help prevent rewiring

of engineered GRNs. Pre-existing GRN structure can be expected to significantly

bias patterns of diversification in regulatory systems, and subsequently shape

genomic evolution by influencing the speed in which genetic circuitry can adapt to

changing conditions. This is highly important for populations adapting during sudden

changes such as niche transitions including the emergence of new pathogens. Our

data moves us closer to the eventual goal of empirically building a set of principles

by which GRNs adapt and offers unique insights into how these systems can

function and influence the evolution of their components.

Materials and Methods

Strains and culture conditions

All ancestral lines in this study are derived from *Pseudomonas fluorescens*

SBW25Δ*fleQ* IS-ΩKm-hah: PFLU2552 (referred to throughout as Δ*fleQ*). Removal of

the flagellum master regulator FleQ and transposon-insertional disruption of the

gene *viscB* (PFLU2552), rendering this strain immotile as detailed previously[31,62]. All

strains were cultured on lysogeny broth (LB; Miller) media at 27°C. *Escherichia coli*

strains for cloning were cultured on LB media at 37°C. Strains, plasmids, media

supplements and primers are detailed in Supplementary Excel file E4.

Knockout mutant construction by two-step allelic exchange

Knockouts of ntrC and PFLU1132 were achieved using two-step allelic exchange

following the protocol of Hmelo et al., 2015 with some alterations. In brief, 400bp

flanking regions up- and down-stream of the target gene were amplified and joined

by strand overlap extension (SOE) PCR to create a knockout allele. This was

inserted into the allelic-exchange suicide vector pTS1 (Containing *sacB* and *tetR*) by

SOE-cloning[64] to create the knockout plasmid, and transformed into the conjugal *E.*

coli strain ST18 by chemical-competence heat-shock. Knockout plasmids were

transferred from *E. coli* ST18 to Δ*fleQ* by two-parent puddle-mating and merodiploids

selected for on LB supplemented with kanamycin sulphate and tetracycline

hydrochloride but lacking 5-ALA supplement required for growth of the *E. coli* ST18

auxotrophic Δ*hemA* mutant. Merodiploids were cultured overnight in LB broth lacking

tetracycline selection and diluted before spread plating onto NSLB media

supplemented with 15% w/v sucrose for *sacB* levansucrase-mediated

counterselection of the pTS1 plasmid backbone. Sucrose-resistant colonies were

isolated and screened for tetracycline sensitivity. Chromosomal presence of the

knockout allele and absence of target gene coding sequences was confirmed by

colony PCR. Knockout of ntrC left an ORF encoding an 11 amino acid truncated

protein, and knockout of PFLU1132 left an ORF encoding an 8 amino acid truncated

protein. All other genomic features including operon structure and terminator regions

were left intact.

Motility evolution experiments

Δ*fleQ*Δ*ntrC* and Δ*fleQ*Δ*ntrC*Δ*PFLU1132* were challenged to rescue motility in the

absence of the FleQ master flagellar regulator on soft agar, as described

previously[31,56]. Pure colonies were picked and inoculated into 0.25% agar LB plates

made as described in Alsohim et al., 2014, and incubated at 27°C. Plates were

checked a minimum of twice daily for motility, recording time to emergence. Motile

zones were sampled immediately and always from the leading edge. Motile isolates

were streaked on LB agar, and a pure colony picked and stored at -80°C as glycerol

stocks of LB overnight cultures. All subsequent analysis was conducted on these

pure motile isolates. Experiment was run for six weeks and any replicates without

motility after this cut-off recorded as having not evolved.

Bacterial growth and motility fitness assays

The fitness of the motility phenotype of evolved Δ*fleQ*Δ*ntrC* strains was tested by

measuring distance moved over 24 hrs of incubation in 0.25% agar LB plates. Six

biological replicates of each strain were grown as separate overnight cultures.

Cultures were adjusted to an OD595 = 1 and resuspended in PBS. Soft agar plates

were inoculated with 1 µL of these suspensions, by piercing the surface of the plate

with the pipette tip, and then effusing the sample into the gap left by the tip. Plates

were incubated for 24 h at 27°C, and photographs taken of motile zones. Surface

area moved was then calculated from the radius of the concentric motile zone

measured from these images ($A = \pi r^2$). Values were square root transformed before

plotting.

Growth in shaking LB broth was measured by inoculating 99 µL of sterile LB broth

with 1 µL of the OD595=1 PBS cell suspensions for each replicate in a 96-well plate.

Plates were incubated at 27°C with 180 rpm shaking in a plate reader, recording

OD595 every hour for 24 hours. Area under the bacterial growth curve was

calculated using the growthcurver package in R and plotted[65]. Area under the

bacterial growth curve provides a metric for fitness, as it accounts for the

characteristics of lag- and log-phase growth, as well as the final carrying capacity of

the population. All growth curves were plotted and inspected prior to calculation of

AUGC to ensure similar AUGC values corresponded to similar curve shapes

(Supplementary Fig. S3)

Mutation identification by whole genome resequencing and PCR amplicon

sanger sequencing

To identify motility rescuing mutations, genomic DNA was extracted from motile

strains and their ancestral strain using the Thermo Scientific GeneJET Genomic

DNA Purification Kit. Genomic DNA was quality checked using BR dsDNA Qubit

spectrophotometry to determine concentration and nanodrop spectrophotometry to

determine purity. Illumina HiSeq sequencing was provided by MicrobesNG

(Birmingham, UK), with a minimum 30x coverage and a sample average of 114x

coverage. Returned paired-end reads were aligned to the *P. fluorescens* SBW25

reference genome[37] using the Galaxy platform[66]. INDELs were identified using the

integrated genomics viewer[67]. For SNP identification, the variant calling software

SNIPPY was used with default parameters (Seemann, 2015). Protein domains

affected by mutations detailed in this article were predicted from amino acid

sequences using pfam, SMART and BLASTp. For a subset of additional motile

mutants, PCR amplification and subsequent sanger sequencing of *PFLU1131* was

performed as this gene was mutated in all previously sequenced strains. This was

done using the service provided by Eurofins Genomics. PCR amplicons were

purified using the Monarch® PCR & DNA Cleanup Kit (New England Biolabs).

Mutations were identified by alignment of the returned PFLU1131 sequence against

the *P. fluorescens* SBW25 reference genome using NCBI BLAST.

Three second-step isolates derived from line 24 were sampled due to observed

colony morphology variation. After whole genome re-sequencing it became apparent

that this variation was due to the presence of a *mutS* mutation and not due to diverse

motility strategies. These isolates all have the mutation *PFLU1132*-D111G

(Supplementary excel file E1, isolate ID's 24-S1, 24-LCV, and 24-SCV), however this

is likely not due to parallel evolution, but instead a single instance of a *PFLU1132*-

D111G mutant that then diversified due to *mutS* mutation. For all subsequent

analysis these three isolates were treated as a single instance of *PFLU1132*-D111G.

RNA-sequencing

Whole-cell RNA was extracted from 20 OD units of *P. fluorescens* cultures in mid-log

phase growth (OD_{595} ~1.5). Cultures were incubated in LB broth at 27°C and 180

rpm shaking. Extractions were performed for biological triplicates of each strain of

interest. Upon reaching the desired OD, growth and RNA expression was halted by

addition of a ½ culture volume of ice-cold killing buffer (20 mM NaN_3, 20 mM Tris-

HCl, 5 mM $MgCl_2$). Cells were pelleted, and the killing buffer removed. A lysis buffer

of β-mercaptoethanol in buffer RLT from the Qiagen RNeasy extraction kit was used

to resuspend pellets, which were then lysed by bead-milling at 4500 rpm for 45 s

with lysing matrix B. Lysates were spun through columns from the Qiagen RNeasy

extraction kit, and the extraction completed following the RNeasy kit protocol. A

DNase I treatment step was included between washes with buffer RW1, by adding

DNase I directly to the column from the RNase-free DNase kit (Qiagen) following kit

protocol. Samples were eluted in nuclease-free water, and subsequently treated with

TURBO DNase from the Turbo DNA-free kit (Invitrogen) following kit protocols.

Purified RNA concentration was measured by Qubit RNA BR assay (Thermo-
scientific), RNA quality by nanodrop spectrophotometry, and RNA integrity by
agarose gel electrophoresis. RNA sequencing was provided by Oxford Genomics
(Oxford UK). Samples were ribodepleted, and the mRNA fraction converted to cDNA
with dUTP incorporated during second strand synthesis. The cDNA was end-
repaired, A-tailed and adapter-ligated. Prior to amplification, samples underwent
uridine digestion. The prepared libraries were size selected, multiplexed and QC'ed
before paired end sequencing over one unit of a flow cell in a NovaSeq 6000.
Returned data was quality checked before it was returned in fastq format.
Reference-based transcript assembly was performed using the Rockhopper RNA
sequencing analysis software[68], to generate transcript read counts. Differential gene
expression analysis was performed using the DESeq2 package of the R statistical
coding language to create lists of differentially expressed genes and corresponding
Log2Fold changes in gene expression.

Rhamnose-inducible expression system constructs

Complementation of *ntrC* and *PFLU1132* knockouts, as well as inducible expression
experiments were performed by single-copy chromosomal re-introduction of the
RpoN-EBP genes under control of a *rhaSR-PrhaBAD* L-rhamnose inducible
expression system[69]. The RpoN-EBP ORF was amplified and a strong ribosome
binding site (stRBS) introduced upstream after a 7bp short spacer between the site
and the start codon. This construct was placed downstream of the *PrhaBAD*
promoter on the miniTn7 suicide-vector pJM220 (obtained from the Addgene plasmid
repository, plasmid #110559) by restriction-ligation and transferred to *E. coli* DH5α
by chemical-competence heat-shock. The miniTn7 transposon containing the *rhaSR*
genes and the PrhaBAD-stRBS-RpoN-EBP construct were transferred to the *P.*
fluorescens chromosome by transposonal insertion downstream of the *glmS* gene
via four-parent puddle-mating conjugation[70]. The relevant *E. coli* DH5α pJM220-
derived plasmid donor was combined with recipient *P. fluorescens* Δ*fleQ* strains,
transposition helper *E. coli* SM10 λpir pTNS2 and conjugation helper *E. coli* SP50
pRK2073, and Gentamicin resistant *Pseudomonas* selected for on LB supplemented
with Gentamicin sulphate and Kanamycin sulphate. Chromosomal insertion of the
correct miniTn7 transposon was confirmed by colony PCR.

β-galactosidase activity assay

To validate the expression dynamics of the rhaSR-PrhaBAD system in the genetic

backgrounds used in this work, the same system described above was introduced

into *P. fluorescens ΔfleQ* and *ΔfleQΔntrC* with *lacZ* under the control of PrhaBAD.

This was obtained from the pJM230 plasmid[69] (obtained from the Addgene plasmid

repository, plasmid #110560). Overnight cultures of each strain were set up in

biological triplicate, and used to inoculate 9 mL cultures of LB broth supplemented

with a range of L-rhamnose concentrations at a bacterial OD_{600} = 0.05. Cultures

were incubated shaking 180 r.p.m. at 27°C until they reached and OD_{600} = 0.5, 2 mL

was spun down and frozen at -20°C until needed. Cell pellets were resuspended in

Z-buffer (60mM $Na_2HPO_4*7H_2O$, 40mM NaH_2PO4*H_2O, 10mM KCl, 1mM

$MgSO_4*7H_2O$), OD_{600} recorded, and 400μL cells added to 600μL fresh Z-buffer. Cell

suspension was lysed with addition of 40 μL Chloroform, 20 μL 0.1% SDS and

vortexing followed by 10 minutes incubation at 30°C. β-galactosidase activity assay

was begun with addition of 200μL o-nitrophenyl-β-D-galactopyraniside (ONPG) and

incubated at 30°C for 20 minutes. If reactions become visibly yellow they were halted

immediately by addition of 500 μL Na_2CO_3 and the time recorded. All other reactions

were halted in this manner after 20 minutes. Reactions were centrifuged to remove

debris and 900 μL added to a cuvette and A_{420} measured. LacZ reporter activity

(MU) was calculated by: MU = (A_{420} * 1000) / (Time(mins) * 0.4 * OD_{600} of initial Z-

buffer cell suspension).

Statistical analyses and data handling

All statistical analysis and data handling was performed using R core statistical

packages and the Dunn.test package. Shapiro-Wilks normality tests were performed

to confirm non-normality of datasets. To compare group medians for more than two

groups, a Kruskal-Wallis test with post-hoc Dunn test and Benjamini-Hochberg

correction was performed, with a $P ≤ 0.025$ taken to indicate significance. For

comparisons of two groups only, a two-sample Wilcox test was used. Spearman's

correlation was used to test significance of correlations between motility rate and

rhamnose concentration, using the R core statistical function cor.test with the

spearman method specified.

For analysis of transcriptomic data, RNA sequencing output was run through

Rockhopper for reference-based transcript assembly. Transcript count data were

then analysed using DESeq2 package in R, for differential gene expression analysis,

as well as generation of PCA and Volcano plots. A gene was taken as being differentially expressed if it differed by a Log2Fold change of ≥1 or ≤ -1 between two samples (representing a doubling or halving of expression level respectively) with an adjusted P-value of ≤0.001.

To identify whether Rho dependent terminators were present downstream of genes of interest, the RhoTermPredict algorithm[71] was run on the *P. fluorescens* SBW25 reference genome in Python.

<u>References</u>

1. Sheppard SK, Guttman DS, Fitzgerald JR. Population genomics of bacterial host adaptation. *Nat Rev Genet*. 2018;19(9):549-565. doi:10.1038/s41576-018-0032-z

2. Cohan FM, Koeppel AF. The Origins of Ecological Diversity in Prokaryotes. *Curr Biol*. 2008;18(21):1024-1034. doi:10.1016/j.cub.2008.09.014

3. Martin CH, Richards EJ. The Paradox behind the Pattern of Rapid Adaptive Radiation: How Can the Speciation Process Sustain Itself Through an Early Burst? *Annu Rev Ecol Evol Syst*. 2019;50:569-593. doi:10.1146/annurev-ecolsys-110617-062443

4. Fang X, Sastry A, Mih N, et al. Global transcriptional regulatory network for Escherichia coli robustly connects gene expression to transcription factor activities. *Proc Natl Acad Sci U S A*. 2017;114(38):10286-10291. doi:10.1073/pnas.1702581114

5. López-Maury L, Marguerat S, Bähler J. Tuning gene expression to changing environments: From rapid responses to evolutionary adaptation. *Nat Rev Genet*. 2009;10(1):68. doi:10.1038/nrg2500

6. Martínez-Antonio A, Janga SC, Salgado H, Collado-Vides J. Internal-sensing machinery directs the activity of the regulatory network in Escherichia coli. *Trends Microbiol*. 2006;14(1):22-27. doi:10.1016/j.tim.2005.11.002

7. Mcdonald JMC, Reed RD. Patterns of selection across gene regulatory networks. *Semin Cell Dev Biol*. 2022;(xxxx). doi:10.1016/j.semcdb.2022.03.029

8. Isalan M, Lemerle C, Michalodimitrakis K, et al. Evolvability and hierarchy in rewired bacterial gene networks. *Nature*. 2008;452(7189):840-845. doi:10.1038/nature06847

9. Mack KL, Ballinger MA, Phifer-Rixey M, Nachman MW. Gene regulation

underlies environmental adaptation in house mice. *Genome Res*.

2018;28(11):1636-1645. doi:10.1101/gr.238998.118

10. Luo X, Song R, Acar M. Multi-component gene network design as a survival

strategy in diverse environments. *BMC Syst Biol*. 2018;12(1):85.

doi:10.1186/s12918-018-0609-3

11. Ali F, Seshasayee ASN. Dynamics of genetic variation in transcription factors

and its implications for the evolution of regulatory networks in Bacteria. *Nucleic*

Acids Res. 2021;48(8):4100-4114. doi:10.1093/NAR/GKAA162

12. Turkarslan S, Reiss DJ, Gibbins G, et al. Niche adaptation by expansion and

reprogramming of general transcription factors. *Mol Syst Biol*. 2011;7(554):1-

16. doi:10.1038/msb.2011.87

13. Damkiær S, Yang L, Molin S, Jelsbak L. Evolutionary remodeling of global

regulatory networks during long-term bacterial adaptation to human hosts.

Proc Natl Acad Sci U S A. 2013;110(19):7766-7771.

doi:10.1073/pnas.1221466110

14. Johnson AD. The rewiring of transcription circuits in evolution. *Curr Opin*

Genet Dev. 2017;47:121-127. doi:10.1016/j.gde.2017.09.004

15. Baumstark R, Hänzelmann S, Tsuru S, et al. The propagation of perturbations

in rewired bacterial gene networks. *Nat Commun*. 2015;6:1-5.

doi:10.1038/ncomms10105

16. Kruzel EK, Giles SS, Hull CM. Analysis of cryptococcus neoformans sexual

development reveals rewiring of the pheromone-response network by a

change in transcription factor identity. *Genetics*. 2012;191(2):435-449.

doi:10.1534/genetics.112.138958

17. Reece-Hoyes JS, Pons C, Diallo A, et al. Extensive rewiring and complex

evolutionary dynamics in a C.elegans multiparameter transcription factor

network. *Mol Cell*. 2013;51(1):116-127. doi:10.1016/j.molcel.2013.05.018

18. Tebung WA, Choudhury BI, Tebbji F, Morschhäuser J. Rewiring of the Ppr1

Zinc Cluster Transcription Factor from Purine Catabolism to Pyrimidine

Biogenesis in the Saccharomycetaceae. *Curr Biol*. 2016;26(13):1677-1687.

doi:10.1016/j.cub.2016.04.064

19. Adhikari S, Erill I, Curtis PD. Transcriptional rewiring of the GcrA/CcrM

bacterial epigenetic regulatory system in closely related bacteria. *PLoS Genet*.

2021;17(3):1-30. doi:10.1371/journal.pgen.1009433

20.   Martchenko M, Levitin A, Hogues H, Nantel A, Whiteway M. Transcriptional
765   Rewiring of Fungal Galactose-Metabolism Circuitry. *Curr Biol*.
766   2007;17(12):1007-1013. doi:10.1016/j.cub.2007.05.017

767   21.   Hsu PC, Lu TC, Hung PH, et al. Plastic Rewiring of Sef1 Transcriptional
768   Networks and the Potential of Nonfunctional Transcription Factor Binding in
769   Facilitating Adaptive Evolution. *Mol Biol Evol*. 2021;38(11):4732-4747.
770   doi:10.1093/molbev/msab192

771   22.   Voordeckers K, Pougach K, Verstrepen KJ. How do regulatory networks
772   evolve and expand throughout evolution? *Curr Opin Biotechnol*. 2015;34:180-
773   188. doi:10.1016/j.copbio.2015.02.001

774   23.   Copley SD. An evolutionary biochemist's perspective on promiscuity. *Trends
775   Biochem Sci*. 2015;40(2):72-78. doi:10.1016/j.tibs.2014.12.004

776   24.   Guzmán GI, Sandberg TE, LaCroix RA, et al. Enzyme promiscuity shapes
777   adaptation to novel growth substrates. *Mol Syst Biol*. 2019;15(4):1-14.
778   doi:10.15252/msb.20188462

779   25.   Aharoni A, Gaidukov L, Khersonsky O, Gould SM, Roodveldt C, Tawfik DS.
780   The "evolvability" of promiscuous protein functions. *Nat Genet*. 2005;37(1):73-
781   76. doi:10.1038/ng1482

782   26.   Taylor TB, Shepherd MJ, Jackson RW, Silby MW. Natural selection on
783   crosstalk between gene regulatory networks facilitates bacterial adaptation to
784   novel environments. *Curr Opin Microbiol*. 2022;67:102140.
785   doi:10.1016/j.mib.2022.02.002

786   27.   Capra EJ, Perchuk BS, Skerker JM, Laub MT. Adaptive mutations that prevent
787   crosstalk enable the expansion of paralogous signaling protein families. *Cell*.
788   2012;150(1):222-232. doi:10.1016/j.cell.2012.05.033

789   28.   Zarrinpar A, Park SH, Lim WA. Optimization of specificity in a cellular protein
790   interaction network by negative selection. *Nature*. 2003;426(6967):676-680.
791   doi:10.1038/nature02178

792   29.   Pougach K, Voet A, Kondrashov FA, et al. Duplication of a promiscuous
793   transcription factor drives the emergence of a new regulatory network. *Nat
794   Commun*. 2014;5:1-11. doi:10.1038/ncomms5868

795   30.   Brunkard JO, Runkel AM, Zambryski PC. A promiscuous intermediate
796   underlies the evolution of LEAFY DNA binding specificity. *Science (80-)*.
797   2015;347(6222):645-649. doi:10.1126/science.1255437

798   31.   Taylor TB, Mulley G, Dills AH, et al. Evolutionary resurrection of flagellar
799   motility via rewiring of the nitrogen regulation system. *Science (80-)*.
800   2015;347(6225):1014-1017. doi:10.1126/science.1259145

801   32.   Taylor T, Mulley G, McGuffin L, et al. Evolutionary rewiring of bacterial
802   regulatory networks. *Microb Cell*. 2015;2(7):256-258.
803   doi:10.15698/mic2015.07.215

804   33.   Lind PA, Farr AD, Rainey PB. Experimental evolution reveals hidden diversity
805   in evolutionary pathways. *Elife*. 2015;4:1-17. doi:10.7554/elife.07074

806   34.   Igler C, Lagator M, Tkačik G, Bollback JP, Guet CC. Evolutionary potential of
807   transcription factors for gene regulatory rewiring. *Nat Ecol Evol*.
808   2018;2(10):1633-1643. doi:10.1038/s41559-018-0651-y

809   35.   Kim DJ, Forst S. Genomic analysis of the histidine kinase family in bacteria
810   and archaea. *Microbiology*. 2001;147(5):1197-1212. doi:10.1099/00221287-
811   147-5-1197

812   36.   Jones J, Studholme DJ, Knight CG, Preston GM. Integrated bioinformatic and
813   phenotypic analysis of RpoN-dependent traits in the plant growth-promoting
814   bacterium Pseudomonas fluorescens SBW25. *Environ Microbiol*.
815   2007;9(12):3046-3064. doi:10.1111/j.1462-2920.2007.01416.x

816   37.   Silby MW, Cerdeño-Tárraga AM, Vernikos GS, et al. Genomic and genetic
817   analyses of diversity and plant interactions of Pseudomonas fluorescens.
818   *Genome Biol*. 2009;10(5):1-16. doi:10.1186/gb-2009-10-5-r51

819   38.   Oliver A, Baquero F, Blázquez J. The mismatch repair system (mutS, mutL
820   and uvrD genes) in Pseudomonas aeruginosa: Molecular characterization of
821   naturally occurring mutants. *Mol Microbiol*. 2002;43(6):1641-1650.
822   doi:10.1046/j.1365-2958.2002.02855.x

823   39.   Luján AM, Maciá MD, Yang L, Molin S, Oliver A, Smania AM. Evolution and
824   adaptation in pseudomonas aeruginosa biofilms driven by mismatch repair
825   system-deficient mutators. *PLoS One*. 2011;6(11).
826   doi:10.1371/journal.pone.0027842

827   40.   Gutierrez PS, Monteoliva D, Diambra L. Cooperative Binding of Transcription
828   Factors Promotes Bimodal Gene Expression Response. *PLoS One*. 2012;7(9).
829   doi:10.1371/journal.pone.0044812

830   41.   Kribelbauer JF, Rastogi C, Bussemaker HJ, Mann RS. Low-affinity binding
831   sites and the transcription factor specificity paradox in eukaryotes. *Annu Rev*

Cell Dev Biol. 2019;35:357-379. doi:10.1146/annurev-cellbio-100617-062719

42. Galperin MY. Structural classification of bacterial response regulators: Diversity of output domains and domain combinations. *J Bacteriol*. 2006;188(12):4169-4182. doi:10.1128/JB.01887-05

43. Kim JK, Tyson JJ. Misuse of the Michaelis–Menten rate law for protein interaction networks and its remedy. *PLoS Comput Biol*. 2020;16(10):1-21. doi:10.1371/journal.pcbi.1008258

44. Ishihama Y, Schmidt T, Rappsilber J, et al. Protein abundance profiling of the Escherichia coli cytosol. *BMC Genomics*. 2008;9:1-17. doi:10.1186/1471-2164-9-102

45. Alkema WBL, Lenhard B, Wasserman WW. Regulog analysis: Detection of conserved regulatory networks across bacteria: Application to Staphylococcus aureus. *Genome Res*. 2004;14(7):1362-1373. doi:10.1101/gr.2242604

46. Prelich G. Gene overexpression: Uses, mechanisms, and interpretation. *Genetics*. 2012;190(3):841-854. doi:10.1534/genetics.111.136911

47. Philippe N, Crozat E, Lenski RE, Schneider D. Evolution of global regulatory networks during a long-term experiment with Escherichia coli. *BioEssays*. 2007;29(9):846-860. doi:10.1002/bies.20629

48. Harrison E, Guymer D, Spiers AJ, Paterson S, Brockhurst MA. Parallel Compensatory Evolution Stabilizes Plasmids across the Parasitism-Mutualism Continuum. *Curr Biol*. 2015;25(15):2034-2039. doi:10.1016/j.cub.2015.06.024

49. Pletnev P, Pupov D, Pshanichnaya L, et al. Rewiring of growth-dependent transcription regulation by a point mutation in region 1.1 of the housekeeping σ factor. *Nucleic Acids Res*. 2020;48(19):10802-10819. doi:10.1093/nar/gkaa798

50. Macneil LT, Walhout AJM. Gene regulatory networks and the role of robustness and stochasticity in the control of gene expression. *Genome Res*. 2011;21:645-657. doi:10.1101/gr.097378.109.21

51. Rao SD, Igoshin OA. Overlaid positive and negative feedback loops shape dynamical properties of PhoPQ two-component system. *PLoS Comput Biol*. 2021;17(1 December):1-18. doi:10.1371/journal.pcbi.1008130

52. Weyder M, Prudhomme M, Bergé M, Polard P, Fichant G. Dynamic modeling of Streptococcus pneumoniae competence provides regulatory mechanistic insights into its tight temporal regulation. *Front Microbiol*. 2018;9(JUL):1-25. doi:10.3389/fmicb.2018.01637

53. diCenzo GC, Sharthiya H, Nanda A, Zamani M, Finan TM. PhoU allows rapid adaptation to high phosphate concentrations by modulating PstSCAB transport rate in Sinorhizobium meliloti. *J Bacteriol*. 2017;199(18):1-20. doi:10.1128/JB.00143-17

54. Hervás AB, Canosa I, Santero E. Regulation of glutamate dehydrogenase expression in Pseudomonas putida results from its direct repression by NtrC under nitrogen-limiting conditions. *Mol Microbiol*. 2010;78(2):305-319. doi:10.1111/j.1365-2958.2010.07329.x

55. Shepherd MJ, Horton JS, Taylor TB. A near-deterministic mutational hotspot in Pseudomonas fluorescens is constructed by multiple interacting genomic features. *Mol Biol Evol*. 2022;39(6):1-7. doi:10.1093/molbev/msac132

56. Horton JS, Flanagan LM, Jackson RW, Priest NK, Taylor TB. A mutational hotspot that determines highly repeatable evolution can be built and broken by silent genetic changes. *Nat Commun*. 2021;12(1). doi:10.1038/s41467-021-26286-9

57. Bretl DJ, Ladd KM, Atkinson SN, Müller S, Kirby JR. Suppressor mutations reveal an NtrC-like response regulator, NmpR, for modulation of Type-IV Pili-dependent motility in Myxococcus xanthus. *PLoS Genet*. 2018;14(10):1-28. doi:10.1371/journal.pgen.1007714

58. Patel V, Matange N. Adaptation and compensation in a bacterial gene regulatory network evolving under antibiotic selection. *Elife*. 2021;10:1-27. doi:10.7554/eLife.70931

59. Schofield MC, Rodriguez DQ, Kidman AA, et al. The anti-sigma factor MucA is required for viability in Pseudomonas aeruginosa. *Mol Microbiol*. 2021;116(2):550-563. doi:10.1111/mmi.14732

60. Gallup O, Ming H, Ellis T. Ten future challenges for synthetic biology. *Eng Biol*. 2021;5(3):51-59. doi:10.1049/enb2.12011

61. Chalovich JM, Eisenberg E. Engineering reduced evolutionary potential for synthetic biology. *Biophys Chem*. 2005;257(5):2432-2437. doi:10.1039/c3mb70606k.Engineering

62. Alsohim AS, Taylor TB, Barrett GA, et al. The biosurfactant viscosin produced by Pseudomonas fluorescensSBW25 aids spreading motility and plant growth promotion. *Environ Microbiol*. 2014;16(7):2267-2281. doi:10.1111/1462-2920.12469

63. Hmelo LR, Borlee BR, Almblad H, et al. Precision-engineering the
Pseudomonas aeruginosa genome with two-step allelic exchange. *Nat Protoc*.
2015;10(11):1820-1841. doi:10.1038/nprot.2015.115

64. Bryksin A V., Matsumura I. Overlap extension PCR cloning: a simple and
reliable way to create recombinant plasmids. *Biotechniques*. 2010;48(6):463-
465. doi:10.2144/000113418.Overlap

65. Sprouffske K, Wagner A. Growthcurver: An R package for obtaining
interpretable metrics from microbial growth curves. *BMC Bioinformatics*.
2016;17(1):17-20. doi:10.1186/s12859-016-1016-7

66. Afgan E, Baker D, Batut B, et al. The Galaxy platform for accessible,
reproducible and collaborative biomedical analyses: 2018 update. *Nucleic*
Acids Res. 2018;46(W1):W537-W544. doi:10.1093/nar/gky379

67. Robinson JT, Thorvaldsdóttir H, Winckler W, et al. Integrative genomics
viewer. *Nat Biotechnol*. 2011;29(1):24-26. doi:10.1038/nbt.1754

68. Brian Tjaden. A Computational System for Identifying Operons Based on RNA-
Seq Data. *Methods*. 2020;176:60-72. doi:10.1016/j.ymeth.2019.03.026.A

69. Meisner J, Goldberg JB. The Escherichia coli rhaSR-PrhaBAD inducible
promoter system allows tightly controlled gene expression over a wide range in
Pseudomonas aeruginosa. *Appl Environ Microbiol*. 2016;82(22):6715-6727.
doi:10.1128/AEM.02041-16

70. Choi KH, Schweizer HP. mini-Tn7 insertion in bacteria with single attTn7 sites:
Example Pseudomonas aeruginosa. *Nat Protoc*. 2006;1(1):153-161.
doi:10.1038/nprot.2006.24

71. Di Salvo M, Puccio S, Peano C, Lacour S, Alifano P. RhoTermPredict: An
algorithm for predicting Rho-dependent transcription terminators based on
Escherichia coli, Bacillus subtilis and Salmonella enterica databases. *BMC*
Bioinformatics. 2019;20(1):1-11. doi:10.1186/s12859-019-2704-x

950   **Figures**
951

952
Figure 1: Rewiring of RpoN-EBP transcription factors to rescue flagellar

motility through promiscuous control of FleQ-controlled genes. A) Flagellar

motility rescue experiment routmap outlines typical progression of motility evolution.

Pathway diagrams display key components of the common (B, Taylor et al., 2015)

and unmasked (C, this study) rewiring pathways for rescue of flagellar gene

expression. Genes are coloured in white, protein components in green. Mutational

targets and their effects are shown in red, and whether a mutation occurs in the first

or second evolutionary step indicated.

966
967
Figure 2: First-step motility rescue mutations in *PFLU1131* A: Diagram of
PFLU1131 protein, indicated positions of first-step motility-rescuing mutations.
Predicted protein domains (pfam) are indicated in lilac, amino acid positions are
indicated as numbers below each domain. Mutations are indicated in blue boxes,
with the number of replicate lines gaining each mutation indicated in brackets. **B)**
FleQ-homolog encoding gene *PFLU1132* is essential for rescued flagellar motility in
the *PFLU1131*-del15 mutant strain and depends on the presence of the kinase
mutation. Scale bars (white) = 12.6mm. Transcription factor gene *PFLU1132* is
deleted and reintroduced as a single-copy chromosomal insertion expressed from an
L-rhamnose inducible promoter system (*rhaSR-PrhaBAD*). The same
complementation lacking the *PFLU1131*-del15 mutation, as well as an empty
expression system transposon were included as further controls. Photographs of
motility after 1 day incubation in 0.25% agar LB plates supplemented with or without
0.15% L-rhamnose for induction of transcription factor expression. **C)** Volcano plots
indicating impact of *PFLU1131*-del15 mutation on the transcriptome relative to the

Δ*fleQ* ancestor. Red points indicate significantly differentially expressed genes.
Triangles indicate flagellar genes, Squares indicate *PFLU1131/2* and adjacent
genes, and Circles indicate all other genes.
986

Figure 3: Log2Fold changes in gene expression for RpoN-EBP controlled genes after first-step motility mutations, relative to Δ*fleQ* ancestor. Δ*fleQ*Δ*ntrC* and Δ*fleQ*Δ*ntrC* first-step differentially expressed genes were determined by RNA sequencing (this work). Δ*fleQ* first-step differentially expressed genes were identified by RNA microarray and originally reported by Taylor *et al.* 2015. List of RpoN-EBP controlled genes was obtained from Jones *et al.,* 2007. Groups of genes by function are indicated by the coloured bars above the plot: Red – Flagellum, Blue – Nitrogen, Teal – Other/unknown.

996
997

Figure 4: Phenotypes of motility rescue mutants. A) Time to emergence of motile
zone (Days) for first-and second-step mutants in the Δ*fleQ*Δ*ntrC* and Δ*fleQ*
backgrounds. Percentages above each plot indicate proportion of replicate evolving
that step within 6 week assay cutoff. **** indicates significant difference in mean time
to emergence from all other groups (Dunn test). **B)** Race assay as measure of
motility fitness. Distance moved over 24hrs in 0.25% Agar LB plates measured
relative to the Δ*fleQ ntrB*-T97P mutant. P-value indicated is generated from a two-
sample Wilcox test. **C)** Fitness in LB broth measured as area under the 24hr growth
curve (AUGC) relative to Δ*fleQ ntrB*-T97P. P-values above plots generated from
Dunn tests. AUGC values and growth curve plots are provided in supplementary
Fig.S3. For all boxplots – box represents first to third quartile range, middle line
represents median value, whiskers range from quartiles to maxima and minima.

1010
1011

Figure 5: Diverse second-step mutations indicate multiple global regulatory strategies for enhancing promiscuity through PFLU1132. A) Diagram of mutational distribution across the SBW25 chromosome for first and second step motility mutants. Each ring represents the chromosome, with black dots indicating non-synonymous mutations. The size of each dot is proportional to the number of mutations occurring in that locus in independent replicates. Mutational target names are coloured by functional category: Red – *PFLU1131/2*, Purple – *PFLU1583/4*, Blue – Metabolic, Green – Global regulatory, Grey – other. **B)** Principle component analysis of transcriptomic data for ancestral, first-step and second-step motility mutants. **C and D)** Volcano plots of differentially expressed genes for second-step mutations relative to Δ*fleQ* ancestor*.* Red points are significantly differentially expressed. Triangles indicate flagellar genes, Squares indicate *PFLU1131/2* and adjacent genes, and Circles indicate all other genes. **E)** Log2Fold change in gene expression for RpoN-EBP controlled genes after *PFLU1583* Δ48-74 mutation – relative to the first-step *PFLU1131*-del15 mutant. RpoN-EBP controlled genes grouped by function are indicated by the coloured bars above the plot: Red – Flagellum, Blue – Nitrogen, Teal – Other/unknown

1031
1032

Figure 6: Second-step *PFLU1131/2* promoter mutation increases *PFLU1131/2*

expression. A) *PFLU1130/1/2* genetic locus in the SBW25 chromosome, and

impact of the 1.59kbp deletion identified in one second-step motile mutant. Location

of the RpoN binding site upstream of PFLU1127/8/9 was predicted by Jones et al.,

(2007). The presence of the rho-dependent terminator was determined by use of

RhoTermPredict[71] **B)** Log2Fold change in gene expression (relative to Δ*fleQ*

ancestor) heatmap of *PFLU1130/1/2* and nearby RpoN-EBP controlled genes,

including *PFLU1127/8/9* in the ancestral Δ*fleQ*Δ*ntrC* strain, first and second step

motility mutants**.**

1042
1043

Figure 7: Impact of RpoN-EBP gene expression on motility speed. A) Mean
motility speed (mm relative to Δ*fleQ ntrB*-T97P) plotted against increasing L-
Rhamnose concentration (%w/v). Mean of six biological replicates for each point.
first-step motility mutants Δ*fleQ ntrB*-T97P and Δ*fleQΔntrC PFLU1131*-del15 had
their respective RpoN-EBPs knockout out, and reintroduced on a miniTn7
transposon with the RpoN-EBP under control of the PrhaBAD and *rhaSR* rhamnose
inducible expression system. In full, these were – Δ*fleQ* ntrB-T97P Δ*ntrC*
miniTn7[*rhaSR*-PrhaBAD-stRBS-ntrC] (rha-*ntrC*) and Δ*fleQΔntrC PFLU1131*-del15
ΔPFLU1132 miniTn7[*rhaSR*-PrhaBAD-stRBS -PFLU1132] (rha-1132). **B)** Motility
speed (mm relative to Δ*fleQ ntrB*-T97P) of common pathway ntrB-T97P first step
mutant, *PFLU1131*-del15 first-step and PFLU1131/2 promoter second-step mutants.
Whiskers represent standard deviation above and below the mean value.
1056

1057

1058

1059

<u>Supplementary figures</u>

Supplementary FigS1*:* FleQ-homolog encoding gene *ntrC* is essential for rescued
flagellar motility in *ntrB* mutant strain and depends on the presence of a kinase
mutation. Transcription factor gene NtrC was deleted, and then reintroduced as a
single-copy chromosomal insertion expressed from an L-rhamnose inducible
promoter system (*rhaSR-PrhaBAD*). The same complementation lacking the ntrB-
T97P mutation, as well as an empty expression system transposon were included as
further controls. Photographs of motility after 1 day incubation in 0.25% agar LB
plates supplemented with or without 0.15% L-rhamnose for induction of transcription
factor expression.

1070
1071

1072 **Supplementary Figure S2:** PFLU1131-del15 mutation engineered into the ΔfleQ
1073 and ΔfleQΔntrC ancestral lines provides motility in both conditions. Indicates that
1074 rescue of motility by this mutation does not depend on ΔntrC and is the sole mutation
1075 required to do so.
1076

Strains (Area under growth curve - AUGC)			
ΔfleQΔntrC	ntrB-T97P	PFLU1131-del15	Replicate
11.658605	6.393025	11.3905	Rep1
11.513825	6.358134	10.943426	Rep2
11.596943	6.674162	10.755457	Rep3
11.385187	6.434133	10.448776	Rep4
11.830055	6.381795	10.696076	Rep5
12.16704	5.922087	10.351345	Rep6

Supplementary Figure S3: Individual microbial growth curves that provide area under the growth curve (AUGC) values that are plotted in Fig.4C. Six biological replicates for each of the three strains were assayed for change in OD_{595} over 24 hours of incubation in shaking LB broth. AUGC values corresponding to each curve are displayed in the table below the graphs.

1120   ***Supplementary Figure S4:*** Impact of motility rescue mutations of flagellar and
1121   chemotaxis gene expression. Log2Fold changes in gene expression relative to the
1122   Δ*fleQ* ancestor are shown for all genes associated with flagellar motility in *P.*
1123   *fluorescens* SBW25. Functional groups of genes are indicated by the coloured bars
1124   and labels to the left of the plot.

1125
1126

Supplementary Figure S5: Phenotypes of motility rescue second-step mutants. **A)**
Race assay as measure of motility fitness. Distance moved over 24hrs in 0.25%
Agar LB plates measured relative to the Δ*fleQ* ancestor *ntrB*-T97P mutant. **B)**
Fitness in LB broth measured as area under the 24hr growth curve (AUGC) relative
to Δ*fleQ* ancestor *ntrB*-T97P. In both plots, mutations are coloured by functional
category: Red – PFLU1131/2, Purple – PFLU1583/4, Blue – Metabolic, Yellow –
Global regulatory, Grey – other. For all boxplots – box represents first to third quartile
range, middle line represents median value, whiskers range from quartiles to
maxima and minima.

1136
1137
1138

1139   **Supplementary Figure S6:** PFLU1583 Δ48-74 mutation depends to presence of
1140   PFLU1131 mutation, and does not act through sigma factor RpoE. **A)** Motility of
1141   PFLU PFLU1583 Δ48-74 with and without accompanying *PFLU1131*-del15 mutation.
1142   **B)** *rpoE* knockout does not revert PFLU1583 mutant to first-step motility phenotype.

1143
1144
1145   **Supplementary Figure S7:** Activity of LacZ expression reporter (Miller units – MU)
1146   with increasing L-rhamnose concentration (%w/v) in both the Δ*fleQ* and Δ*fleQ*Δ*ntrC*
1147   ancestral genetic backgrounds with miniTn7[rhaSR-PrhaBAD-stRBS-LacZ].
1148   Whiskers represent standard deviation above and below the mean value.

1149