1 # Single-nucleus ATAC-seq elucidates major modules of gene
2 # regulation in the development of non-alcoholic fatty liver disease
3

Fumihiko Takeuchi [1,2] * , Yi-Qiang Liang [1] , Hana Shimizu-Furusawa [3] , Masato

Isono [1] , Mia Yang Ang [1,4] , Kotaro Mori [2] , Taizo Mori [5] , Eiji Kakazu [5] , Sachiyo

Yoshio [5] , Norihiro Kato [1,2,4]

[1] Department of Gene Diagnostics and Therapeutics, Research Institute,

National Center for Global Health and Medicine, Japan.

[2] Medical Genomics Center, Research Institute, National Center for Global

Health and Medicine, Japan.

[3] Department of Hygiene and Public Health, School of Medicine, Teikyo

University, Japan.

[4] Department of Clinical Genome Informatics, Graduate School of Medicine,

The University of Tokyo, Japan.

[5] Department of Liver Diseases, The Research Center for Hepatitis and

Immunology, National Center for Global Health and Medicine, Japan.

* Correspondence: fumihiko@takeuchi.name

Abstract

Background: Non-alcoholic fatty liver disease (NAFLD) develops from fatty liver to steatohepatitis during which multiple cell types may play different roles. Aiming to understand the tissue composition of cell types, their gene expression and global gene regulation in the development of NAFLD, we performed single-nucleus and bulk ATAC-seq on the liver of rats fed with a high-fat diet (HFD).

Methods: Male Spontaneously Hypertensive Rats were fed a normal diet or a HFD. Rats fed HFD for 4 weeks developed fatty liver, and those fed HFD for 8 weeks further developed steatohepatitis. Under the washout condition, where 4 weeks of HFD is followed by 4 weeks of a normal diet, fatty liver was partially ameliorated. For each dietary condition, we performed single-nucleus ATAC-seq on one animal and bulk ATAC-seq on four animals.

Results: In accordance with the pathological progression from fatty liver to steatohepatitis, the proportion of inflammatory macrophages dramatically increased. By machine learning, we divided the global gene expression into modules, such that transcription factors in a module regulate the genes in the same module. Consequently, many of the modules rediscovered known regulatory relationship between the transcription factors and biological processes. For the discovered biological processes, we searched core genes, which were defined as genes central regarding co-expression and protein-protein interaction. A large part of the core genes overlapped with previously implicated NAFLD genes.

Conclusions: Single-nucleus ATAC-seq combined with data-driven statistical analysis help elucidate the global gene regulation *in vivo* as a combination of modules and discover core genes of the relevant biological processes.

48 **Keywords**

49 Non-alcoholic fatty liver disease, single-nucleus ATAC-seq, transcription factor,

50 gene regulation

Background

Non-alcoholic fatty liver disease (NAFLD) is a continuum of liver abnormalities ranging from non-alcoholic fatty liver (NAFL) to non-alcoholic steatohepatitis (NASH). It has a global prevalence of ~25% [1]. NAFLD can gradually develop from fatty liver, steatohepatitis and cirrhosis to hepatic cancer. No approved drugs are available for NAFLD, and biomarkers are currently insufficient. The liver is composed of multiple cell types, including hepatocytes, cholangiocytes, stellate cells, endothelial cells, and white blood cells, each of which plays a different role in disease development. Recent single-cell RNA-seq studies have uncovered cell type-specific gene expression changes in NAFLD [2–4].

A key upstream event in gene expression regulation is transcription factors (TFs) binding to DNA. The conventional ChIP-seq assay has a methodological weak point since it is limited to measuring the binding of one TF in bulk of cells. In contrast, single-nucleus ATAC-seq (snATAC-seq) realized simultaneously measuring the binding of all TFs in single cells. In ATAC-seq, open chromatin sites are observable and indicate 1) TF binding at the nearby motifs on the genome and 2) expression of a gene located close to the site. Since previous single-cell analyses of NAFLD were not ATAC-seq but all RNA-seq, TFs binding to chromatin could not be measured.

The tissue composition of cell types changes during disease development in various diseases and affects the outcome. Cell type counting of solid tissue became feasible through the single-cell assay, which is still expensive. Instead, deconvolution, the statistical inference of the cell type composition of tissue samples assayed in bulk, has been used for transcriptome and DNA methylation assays [5, 6]. For ATAC-seq, Li and associates [7] previously proposed an algorithm for deconvolution, but no experimental validation has been done with single-cell data.

Aiming to understand the cell type composition and global TF regulation in NAFLD development, we performed single-nucleus and bulk

ATAC-seq on the liver of rats fed or withdrawn from a high-fat diet (HFD). We

first analyzed the composition of cell types in the liver and the gene

expression separately by cell type. Then, to explore global TF regulation in

each cell type, we developed a new statistical method that decomposes TF

regulation into major modules, each being characterized by biological

processes. We also developed a statistical method to detect core genes for

the individual biological processes. Taking snATAC-seq data of an *in vivo*

experimental model as the input, our data-driven reductionist workflow

proceeds from global TF regulation, TF regulation modules, biological

processes to core genes. The core genes thus identified overlapped well with

the previously known NAFLD genes, providing experimental validation to our

newly developed statistical methods.

Results

Cell type composition of the liver

Spontaneously Hypertensive Rats (SHR) fed HFD for 4 weeks developed fatty

liver, and those fed HFD for 8 weeks further developed inflammation and

premature fibrosis (steatohepatitis) (Figure 1A; Table S1). Under the washout

condition, where 4 weeks of HFD is followed by 4 weeks of a normal diet,

fatty liver was partially ameliorated. Along with control rats only fed a normal

diet, five dietary conditions were examined.

In order to catalog cell types in the liver, we performed snATAC-seq

on one animal per dietary condition, five animals in total. We obtained a total

of 14,486 nuclei (~3000 per sample), which were grouped into 16 clusters

by the similarity of chromatin opening. Hepatocytes (seven clusters),

endothelial cells (three clusters), stellate cells (two clusters) and white blood

cells (four clusters) separated clearly (Figures 1B and 1C). The clusters for

white blood cells corresponded to inflammatory macrophages, non-

inflammatory macrophages, T cells and NK cells combined, and B cells.

Cholangiocytes were included in the hepatocyte clusters, and neutrophils and
Kupffer cells were included in the macrophage clusters.

To measure the changes in cell type population through dietary
intervention, we performed bulk ATAC-seq on four animals per condition, 20
animals in total. Using snATAC-seq as reference, we inferred the cell type
composition in the bulk ATAC-seq samples by deconvolution. Beforehand,
using the five samples assayed in both single-nucleus and bulk ATAC-seq, the
accuracy of deconvolution for ATAC-seq was validated (Figure 1D). The
difference of the inferred proportion from the true proportion counted by
snATAC-seq had average of zero and standard deviation of 0.035, indicating
high accuracy. By using four animals per condition, the standard error would
become 0.018. Compared with the normal diet condition, hepatocytes
significantly decreased after 8 weeks of HFD (Figure 1E). Inflammatory
macrophages largely increased after 4 weeks of HFD and further increased
after 8 weeks of HFD. Non-inflammatory macrophages also increased after 8
weeks of HFD. Additionally, B cells increased under HFD conditions. In
accordance, previous studies of HFD-induced NAFLD animal models reported
an increase in liver-resident macrophages [8], recruitment of monocyte-
derived macrophages [9], and an increase in intrahepatic B cells [10, 11].
Under the washout condition, the cell type composition returned to the same
as the normal diet condition. Here, we elucidated how the quantity of cell
types changed through dietary intervention, and we next investigate the
qualitative change.
133

Differential gene expression under HFD in each cell type

We analyzed cell type-specific differential gene expression by comparing the
nuclei of the same cell type between dietary conditions (Table S2; Dataset
S1). Hereafter, clusters for inflammatory and non-inflammatory macrophages
are combined and regarded as the macrophage cell type. T cells, NK cells,

and B cells were not analyzed because the number of nuclei was limited. The

biological processes characteristic of the differentially expressed genes were

cell type-dependent (Figure 2). Steroid metabolism and fatty acid metabolism

genes were expressed in hepatocytes differently under HFD and washout

compared with the normal diet. As of inflammation genes, the expression was

altered in macrophages after 4 weeks of HFD and washout. The inflammation

genes and apoptosis genes were expressed in hepatocytes differently after 8

weeks of HFD. Genes for the actin filament-based process were expressed in

stellate cells differently after 8 weeks of HFD.

Conventional differential gene expression analysis using bulk tissue

samples was limited in detecting 1) changes in minor cell types, 2) changes

in a cell type with a low basal expression of the gene relative to other cell

types, and 3) concurrent changes in gene expression and cell population that

mutually cancel, for example, downregulated genes in a cell type that is

increasing in population. For the comparison between the 4 weeks of HFD and

the normal diet, cell type-specific differential gene expression was cross-

checked with bulk tissue transcriptome analysis, and the results were found

to replicate under the abovementioned limitations (Figure S1). The

differential gene expression was replicated in hepatocytes, but to a lesser

degree in minor cell types. For macrophages, whose proportion increased

after 4 weeks of HFD, the upregulated genes were replicated in bulk

experiments but the downregulated genes were not.

Major modules of TF regulation and associated biological processes

To elucidate the global picture of TF regulation, we inferred the regulatory

impact of all TFs on all genes and extracted major modules of the regulatory

relation. By snATAC-seq, we measured in all nuclei of each cell type (from

rats of all dietary conditions) the genome-wide binding of TFs and the

expression of genes, and inferred the regulation of the latter by the former

by applying machine learning. The regulatory relation was divided into

modules. A module comprises a subset of TFs and a subset of genes, such

that TFs belonging to a module regulate genes belonging to the same module

but do not regulate genes belonging to other modules. In hepatocytes, we

obtained four modules (Figure 3A). Modules are simply a grouping of

TFs/genes detected by machine learning and have no biological meaning by

themselves. By applying gene set enrichment analysis to the genes belonging

to a module, but not using the information on TFs, we searched for biological

processes that characterize the module (Figure 3B). In the following, we

describe modules for hepatocytes, endothelial cells (Figure 4), stellate cells

(Figure 5), and macrophages (Figure 6). If a biological process was assigned

to a module, we sought the literature that supports the regulation of the

biological process by the TFs belonging to the module. In five of the seven

discovered modules, there was support for the discovered linkage between

the TFs and biological processes, suggesting the validity of the module

discovery algorithm.

Module 1 of hepatocytes was characterized by the binding of STAT

family TFs (namely, STAT5B, STAT4 and STAT3) and by genes for steroid

metabolism.

Module 2 of hepatocytes, Module 1 of endothelial cells, and Module 1

of macrophages were characterized by the binding of AP-1 family TFs (namely,

FOS, JUND, BACH1 and BACH2), SMARCC1 and NFKB1, and by genes for

TNFα signaling via NF-κB. AP-1 TFs respond to cytokine stimuli [12].

SMARCC1 is a member of the SWI/SNF family of proteins, which function as

a helicase and interacts with NF-κB [13].

Module 3 of hepatocytes was characterized by the binding of TCF/LEF

family TFs (namely, LEF1 and TCF7L1) and by genes in the farnesoid X

receptor pathway for bile acid synthesis and genes for the glutamine family

amino acid biosynthesis. Within the liver lobule, a unit that assembles to form

the liver, a central vein is located in the center, the portal triad (portal vein,

hepatic artery, and bile duct) is located in the periphery, and the metabolism

of hepatocytes in between differs between the pericentral and periportal

zones. The glutamine family amino acid biosynthesis genes include *Glul* and

Ass1, which are highly expressed in the pericentral and periportal zones,

respectively [14]. Additionally, the LEF1 TF binds to β-catenin protein and

activates the Wnt signaling pathway [15], and liver-specific β-catenin KO mice

exhibited defective cholesterol and bile acid metabolism in the liver [16].

Moreover, Wnt-activated genes are highly expressed in the pericentral zone.

Thus, LEF1 TF presumably regulates bile acid metabolism and zonated gene

expression in the liver.

Module 2 of endothelial cells was characterized by the DNA binding of

AHCTF1 (aka ELYS) and ZNF740 and by genes for angiogenesis. AHCTF1 is a

DNA-binding protein that is required in mitosis [17]. ZNF740 activates

angiogenesis in pulmonary artery endothelial cells of rats [18].

Module 1 of stellate cells was characterized by SOX9 TF binding and

genes in the semaphorin-plexin signaling pathway.

Module 2 of macrophages was characterized by the binding of Maf

family TFs (namely, MAFA, MAFB and MAFG) and by genes for complement

system activation. In *Mafb*-deficient macrophages of mice, C1q production

decreased [19].

Module 3 of macrophages was characterized by the binding of IRF

family TFs (namely, IRF1, IRF2 and IRF7), which regulate immune responses,

and by genes for angiogenesis. IRF1 contributes to the commitment of pro-

inflammatory M1 macrophages, which produce angiogenic stimulators [20].

Core genes in specific biological processes

In the previous section, we discovered major modules of TF regulation and

then searched across databases, such as Gene Ontology, for biological

processes that characterize the modules. We equated a biological process

with its comprising genes, and defined them as a gene set (GS). A GS

characterizes a module if it significantly overlaps with the genes of the module.

Here, we aim to find "core genes" of a biological process. In preparation, we

compute the activity of a GS in single nuclei as the "GS activity score" by

taking the average binding/expression level of TFs/genes in the set or, more

precisely, the first principal component. In hepatocytes, regarding the

biological process of TNFα signaling via NF-κB, its GS activity score was the

highest among nuclei from 8 weeks of HFD (Figure 7A, top panel).

Core genes are the genes that are central regarding two connections

of the GS, co-expression and protein-protein interaction. For co-expression,

we first compute the correlation coefficient between the GS activity score and

the expression level of a gene (or the binding of a TF). The absolute value of

the correlation coefficient is required to be large for a core gene. Moreover,

the total amount of protein-protein interactions between the gene and the GS

is required to be large. Therefore, by plotting the correlation coefficient on

the horizontal axis and the total protein-protein interactions on the vertical

axis, core genes are expected to appear in the top left or top right corner.

This approach can be used to prioritize the known genes in the target GS

(Figure 7A, middle panel) and to discover novel genes not included in the GS

(Figure 7A, bottom panel). In the following, we inspect some of the biological

processes that appeared in the previous section.

In hepatocytes, the GS activity score for TNFα signaling via NF-κB

(Figure 7A) was the highest after 8 weeks of HFD. TF binding that was

positively correlated and had abundant protein-protein interactions included

FOS/JUN and NFKB1/RELA. The positively correlated core genes were *Tnf*,

Nfkb1, *Bcl2l1* and *Il1r1*. The negatively correlated core gene was *Cxcl12*.

In endothelial cells, the GS activity score for TNFα signaling via NF-

κB (Figure 7B) was higher under HFD conditions. The positively correlated

core genes were *Socs3* and *Pecam1*. The negatively correlated core genes

were *Tlr2*, *Irs2*, *Tlr4*, *Il15*, *Ccr5* and *Cxcr4* (receptor for the chemoattractant

CXCL12 [21]).

In macrophages, the GS activity score for TNFα signaling via NF-κB

(Figure 7C) was higher after 4 weeks of HFD. TF binding that was positively

correlated and had abundant protein-protein interactions included JUN and

KLF6. The positively correlated core genes were *Cd44*, *Fn1* and *Cxcr4*. KLF6

regulates the switching of macrophages between pro- and anti-inflammatory

[22]. In accordance, *Lyz2* and *Marco*, the marker genes for inflammatory and

non-inflammatory macrophages, showed a correlation of 0.39 and −0.49,

respectively, with KLF6 binding in our single-nucleus data.

In hepatocytes, the GS activity score for steroid metabolism (Figure

S2A) was the highest after 8 weeks of HFD and to a lesser degree after 4

weeks of HFD. The positively correlated core genes included *Abcg1*, *Abca1*

and *Scd*. The negatively correlated core genes included *Sqle*, *Hmgcr*, *Apoa1*,

Sult2a1, *Acox2* and *Alb*. Among the 15 cytochrome P450 genes showing a

strong negative correlation (correlation smaller than −0.3 and protein-protein

interactions within the top 5%), 13 were oxidoreductase of EC 1.14.14.1. For

reference, we labeled the binding of steroid sensing TFs, NR1H2/3 (aka

LXRα/β), SREBF1 and SREBF2, which were negatively correlated with the GS

expression.

To investigate lobular zonation genes in hepatocytes, the GS was

taken from the list of zonation genes in normal mice (q-value < 10^{-6} in Table

S3 in [14]). The GS activity score showed a modest difference between

dietary conditions (Figure S2B), which agrees with zonation being a normal

phenomenon unrelated to NAFLD. When classified by the correlation with the

GS activity score, the genes with correlation coefficient > 0.3 were all

pericentral and those with correlation coefficient < −0.3 were all periportal,

showing that the GS activity score rediscovered zonation.

In stellate cells, the GS activity score for semaphorin-plexin signaling

was higher at 8 weeks of HFD (Figure S2C). The positively correlated core

gene was *Nrp2*. The negatively correlated core genes were *Nrp1*, *Plxnd1*,

semaphorin genes (*Sema3d*, *Sema3e* and *Sema6d*) and *Ntn1*.

Discussion

By performing single-nucleus and bulk ATAC-seq on the liver of rats fed or

withdrawn from the HFD, we studied the transition in tissue composition of

cell types and cell type-specific gene regulation in a rat model of NAFLD. The

accuracy of ATAC-seq-based deconvolution was validated and used to analyze

cell type composition. In accordance with the pathological progression from

fatty liver to steatohepatitis that could have occurred between 4 and 8 weeks

of the HFD intervention in our rat model, the proportion of inflammatory

macrophages dramatically increased. In contrast, after 8 weeks of HFD, the

proportion of hepatocytes largely decreased. In the washout condition, where

4 weeks of HFD was followed by 4 weeks of normal diet, the cell type

composition returned similar to rats only fed with the normal diet.

From the global analysis of TF binding and gene expression in single

nuclei, we identified major modules of TF regulation (Figure 8), which could

be shared between different cell types or be specific to certain cell types. One

module shared among hepatocytes, endothelial cells, and macrophages was

characterized by the binding of AP-1 TFs and a biological process of TNFα

signaling via NF-κB associated with core genes that appear to be specific to

cell types. For many of the discovered modules, TFs in a given module were

known to regulate biological processes assignable to the genes in the same

module, suggesting the validity of our module discovery algorithm. Finally,

for the biological processes that emerged in TF regulation analysis, by

incorporating known protein-protein interactions, we could identify core

genes, many of which overlap with previously implicated NAFLD genes (see

below).

We consider that the gene regulatory machinery of a cell is modular
and some of the modules would become active when perturbed by HFD. As
an analogy, a car consists of many machinery parts, and on a cold sunny day
the engine and air conditioning are active but the wiper is not. To discover
gene regulatory modules that respond to HFD stimulus, we integratively
analyzed the TF binding and gene expression in nuclei derived from rats under
various dietary conditions. In each cell type, we used machine learning to
decompose the global variation of TF regulation into major modules, where
the TFs belonging to a given module appear to regulate the genes that belong
to the same module. In the modules identified as such, the TFs and the
function of regulated genes agree with the literature: e.g., the AP-1 TFs that
regulate inflammation genes (Module 2 of hepatocytes, Module 1 of
endothelial cells and Module 1 of macrophages) are known to respond to
cytokine stimuli [12], and LEF1 that regulates zonation (Module 3 of
hepatocytes) is known to activate Wnt signaling [15]. Previous snATAC-seq
studies have characterized the difference between cell types [23] using TF
binding. In contrast, we illustrated how TFs control the states of cells
belonging to one cell type (e.g., the hepatocytes with various degrees of
inflammation) possibly in a cell-type specific manner (e.g., a distinct set of
core genes and other target genes associated with AP-1 TFs in different cell
types).

One novel method developed in this article is the computation of the
GS activity score in single cells to indicate the activity of a GS, as if it were
the expression of a representative gene. In experiments, gene expression
can be directly measured. The activity of a pathway is of interest but not
directly measurable. One way would be to take the average expression of
genes in that pathway, but this is not optimal for the following reasons. A
simple average would give equal weight to both core and periphery genes in

the pathway, increasing noise from the latter. Moreover, activator and

suppressor genes contribute to the pathway in opposite directions. Instead of

a simple average, we used the first principal component to deal with these

issues. The principal component approach was adapted from the PLAGE

program [24], which is used to compute pathway activity scores in bulk tissue

specimens. The GS activity score could improve the understanding of

biological processes. For example, *Abcg1* and *Sqle* correlated positively and

negatively, respectively, with the activity score for the steroid metabolism

pathway (Figure S2A). As this score became higher under the HFD, the GS

activity represented by the score presumably works in response to the

overdose of cholesterol contained in the diet. Thus, a positive correlation of

Abcg1 (for lipid efflux) and a negative correlation of *Sqle* (for sterol

biosynthesis) are reasonable. Moreover, the cell type-specific nature of the

GS activity score could unravel cell type-specific dynamics; the inflammation

process peaked under 4 weeks of HFD in macrophages and under 8 weeks of

HFD in hepatocytes (Figure 7). The GS activity score can also be applied to

single-cell and single-nucleus RNA-seq data.

Another novel method in this article is finding the core genes of

biological processes. Given a GS comprising known genes for the process, we

identified core genes as those that show a strong correlation with the GS

activity score as well as abundant protein-protein interactions with the GS.

We obtained evidence supporting the former from single-nucleus experiment

data and the latter from a public database. We sought literature indicating

that core genes cause NAFLD or serve as biomarkers for NAFLD and found

indications for a large part of the core genes (see below): for the TNFα

signaling via NF-κB process, 4 of 5 in hepatocytes, 4 of 8 in endothelial cells,

and 1 of 3 in macrophages; for steroid metabolism 3 of 9 in hepatocytes; for

semaphorin-plexin signaling 3 of 7 in stellate cells. The presence of

substantial overlap between the current findings and known NAFLD genes

provides a proof-of-concept for our data-driven approach, using snATAC-seq

data from a rat model of dietary intervention as input, extracting modules,

biological processes, and then core genes. Regarding TNFα signaling via NF-

κB in hepatocytes, in animal studies, *Tnf* KO mouse was protected from

NAFLD [25], *Nfkb1* KO mouse exacerbated NAFLD [26], and *Il1r1* KO mouse

had less hepatic triacylglycerol [27]. CXCL12 (aka SDF1) induced migration

of stellate cells [28], endothelial cells [21] and CD4+ T cells [29]. In human

biomarker studies, higher circulating TNFα level was associated with NAFLD

[30] and future NAFLD [31]. In NAFLD patients, soluble IL1R1 was higher

(Ajmera et al., 2017). Regarding TNFα signaling via NF-κB in endothelial cells,

in animal studies, *Pecam1* KO mouse developed NASH [32], whereas *Tlr4* KO

mouse [33] and *Il15* KO mouse [34] were protected from NAFLD. *Ccr5*

antagonist eased NAFLD in mice [35]. Regarding TNFα signaling via NF-κB in

macrophages, *Cd44* KO mouse had less liver injury under NAFLD inducing

diet, and NAFLD patients had elevated serum soluble CD44 [36]. Regarding

steroid metabolism in hepatocytes, *Scd1* KO mouse exacerbated NAFLD [37],

and *Acox2* KO mouse developed NAFLD [38]. NAFLD patients had lower

serum APOA1 [39, 40]. Regarding semaphorin-plexin signaling in stellate

cells, in animal studies, *Nrp2* conditional KO reduced liver fibrosis in mice

[41]. *Nrp1* KO attenuated cell motility and overexpression increased motility

[42]. *Sema3e* KO eased liver fibrosis in mice [43].

In search of drug targets or biomarkers of NAFLD, we looked up

proteins whose subcellular localization is extracellular or plasma membrane

proteins [44] among the genes strongly upregulated under a dietary

intervention (Table S2) or the core genes positively correlated with a NAFLD-

related activity. Previous reports of NAFLD biomarkers in humans are cited.

For hepatocytes, we found *Il1r1* [45], *Il1r2* and *Tnf* [30]. For endothelial cells,

there was *Hgf* [46–48]. *Pecam1* was listed for both endothelial and stellate

cells. For stellate cells, there were *Faslg* [49] and *Nrp2*. For macrophages,

there were RT1-Db1, *Cd44* [36], *Cxcr4* and *Fn1*. We also searched DGIdb

(2022-Feb) for existing drugs that can potentially be repurposed and found

antagonist/antibody/inhibitor of upregulated or positively correlated genes,

Bcl2l1, *Cxcr4*, *Hgf*, *Il1r1* and *Tnf*, and agonist of downregulated or negatively

correlated genes, *Ccr5*, *Cxcr4*, *Tlr2* and *Tlr4* (Table S3). *Cxcr4* was positively

correlated with TNFα signaling via NF-κB in macrophages but negatively

correlated in endothelial cells.

This study has several limitations. Firstly, it was difficult to distinguish

TFs with very similar motifs. A specific TF was regarded to bind a specific site

if the chromatin was open near the site and if the DNA sequence of the site

matched with the known TF-binding motif. However, instead, a different TF

with a similar motif could have bound there. Secondly, the interpretation of

gene expression requires caution. Gene expression can be quantified by the

amount of either A) mRNA transcription per time, B) mRNA in the nucleus,

C) mRNA in the cell, D) protein translation per time, or E) protein.

Measurement for bulk tissue is conducted by ATAC-seq (A), RNA-seq (C),

Ribo-seq (D), or Western blotting (E), and measurement for single

nucleus/cell is performed by single-nucleus ATAC-seq (A), single-nucleus

RNA-seq (B), or single-cell RNA-seq (C). While our measurement, A, could

reflect the state of gene regulation, it could differ from E, which reflects the

state of metabolism, and the fold-change value would differ among the types

of gene expression. Thirdly, the TF regulation could differ between rat, mouse

and human.

To discover TF modules, we used machine learning whose

performance can be affected by the choice of hyperparameters. Under a

hyperparameter that allows the machine learning model to be too complex

and flexible, compared to the amount of data, the model can pick up noise

as signal and overfit. We chose hyperparameters based on previous empirical

validations (for the number of candidate regulators at each node in GENIE3)

or residual error (for the number of factors in NMF) but did not explore

hyperparameters extensively. We assessed the adopted hyperparameter post

hoc. Under a range of hyperparameters, we randomly divided the hepatocytes

into two sets of halves, performed TF module discovery in each set and

assessed the consistency. We did not observe overfitting for the

hyperparameters adopted in this study (Figure S3). Nevertheless, an

assessment using data from an independent experiment is needed.

Conclusions

By performing single-nucleus ATAC-seq in a rat model of NAFLD and using

novel statistical methods, we elucidated a global picture of *in vivo* TF

regulation in each cell type as a set of modules and discovered core genes

for NAFLD-relevant biological processes. The core genes can serve as novel

candidates responsible for NAFLD and our statistical methods can be applied

to elucidate the TF regulation in biological systems other than NAFLD.

Methods

Animal experiment

We used male rats of SHR/Izm strain, which is an inbred genetic model of

essential hypertension. The strain was supplied by Japan SLC. Rats were

weaned at 4 weeks after birth and placed on a normal diet (SP diet, Funabashi

Farm, Japan). Adult rats were fed the normal diet or a high-fat atherogenic

diet (24% fat, 15% protein, 5% cholesterol, 2% cholic acid) according to the

study design. Compared to models based on nutritional deficiency (such as

methionine and choline-deficient diet) or chemical-induced models (such as

carbon tetrachloride), HFD models resemble more human NAFLD, both

pathophysiologically and phenotypically [50]. SHR can mimic NAFLD under a

hypertension background. The rats were killed under pentobarbital

anesthesia (200 mg/kg via intraperitoneal infusion), and the organs were

excised and immediately frozen at −70 C. We used the liver's largest lobe (left

lateral lobe) for pathology imaging and the right segment of the second

largest lobe (left median lobe) for genomics analysis. The measured

outcomes were genomic and biochemical assays of the liver, pathology

imaging of the liver, biochemical assays of the serum, organ and body weight,

and blood pressure. The experimental unit was the individual animal. No

animal was excluded during the experiment. The allocation of animals to

dietary conditions was not randomized. We controlled the dietary conditions

but did not control for other possible confounders. The group allocation was

known to all experimenters except for the pathologist, who inspected

unlabeled preparations. All animal experiments conformed to the Guidelines

for Animal Experiments of the National Center for Global Health and Medicine

(NCGM), and were performed in the animal facility of NCGM. The animal

experiments and the protocol were registered and approved by the Animal

Research Committee of NCGM (permission number 21009).

Library preparation and sequencing for snATAC-seq

Nuclei were isolated from flash-frozen liver tissue according to the 10x Genomics protocol CG000212 (samples m154207 and m154211) or using the Minute Single Nucleus Isolation Kit for Tissues/Cells (Invent Biotechnologies, SN-047) (samples m167108, m167203 and m168101). For each sample, the sequencing library was prepared using the Chromium Next GEM Single Cell ATAC Reagent Kits v1.1 (10x Genomics, CG000209) and sequenced using the Novaseq 6000 sequencer (Illumina).

Library preparation and sequencing for bulk tissue ATAC-seq

Bulk tissue ATAC-seq was performed according to the Omni-ATAC protocol [51] using flash-frozen liver tissue. The library was sequenced using the NextSeq sequencer (Illumina).

Measuring triglycerides in liver tissue

Triglyceride in the liver tissue was measured enzymatically using the Adipogenesis Colorimetric/Fluorometric Assay Kit (BioVision, K610-100). Tissue homogenate was obtained by dispersing 10 mg tissue in 1 ml lipid extraction solution under 2200 rpm for 1 min, using a handheld disperser (KINEMATICA, PT 1300 D). For fatty liver tissues, we prepared 1/3 dilution of the homogenate by adding lipid extraction solution and examined both the original and the dilution. The triglyceride was measured in a colorimetric assay according to the manual. Triglyceride quantity in mol unit was converted to g unit by multiplying 885.7.

Measuring total cholesterol in liver tissue

Total cholesterol in the liver tissue was measured enzymatically using the Total Cholesterol and Cholesteryl Ester Colorimetric Assay Kit II (BioVision, K623-100). Tissue homogenate was obtained by dispersing 15 mg tissue in

600 ul CHCl$_3$:IPA:NP-40 (7:11:0.1) at 2200 rpm for 1 min, using a handheld

disperser (KINEMATICA, PT 1300 D). The homogenate was then centrifuged

for 5 min at 15,000 g. Thereafter, we transferred 200 ul of the liquid phase

to a new tube, dried with air-flow-in at 40 C for 10 min using a centrifugal

concentrator (TOMY Digital Biology, CC-105), and dissolved the dried lipids

with 400 ul Cholesterol Assay Buffer. Total cholesterol was measured in a

colorimetric assay according to the manual.

Data cleaning for snATAC-seq

Data cleaning was conducted separately by samples. ATAC-seq fragments

were assigned to single nuclei, and nuclei of good quality were retained as

described below. In principle, one nucleus and one barcoded bead are

contained in one droplet of the Chromium assay; thus, ATAC-seq fragments

from the same nucleus are identifiable by the shared unique barcode. Using

the Cell Ranger ATAC software (version 2.0.0), we mapped the paired-end

reads to the mRatBN7.2/rn7 rat reference genome and identified ATAC-seq

fragments, which were assigned to single barcodes.

Occasionally in the Chromium assay, more than one barcoded beads

are contained in one droplet, in which case, excessive ATAC-seq fragments

would be assigned in common to the barcodes. Such cases were detected,

and the barcodes were unified and renamed using the bap2 software (version

0.6.7) [52]. In the downstream analysis, we can assume that one barcode

corresponds to one droplet.

We managed the snATAC-seq data using the ArchR package (version

1.0.1) [53] of the R software (version 4.1.1). ATAC-seq fragments from the

autosomes and chromosome X were retained. We retained nuclei that had

abundant and enriched fragments around transcription start sites (TSS) as

follows: ≥1000 fragments within 1 kb to TSS, the proportion of fragments

within 1 kb to TSS being ≥7.5% (samples m154207, m154211 and m167203)

or ≥15% (samples m167108 and m168101), and Tn5 insertions (which occur
at both ends of a fragment) within 25 bp to TSS occurring ≥4 times frequently
than the insertions at 1900–2000 bp from TSS.

For each nucleus, the proportion of fragment counts from a
chromosome (e.g., chromosome 1) is approximately constant, being
proportional to the chromosome length, but could deviate if a droplet is
contaminated by free DNA from broken nuclei [54]. Therefore, we
represented the proportion by a matrix indexed by nuclei (rows) and
chromosomes (columns). To standardize the matrix, we subtracted column-
wise averages and divided by the standard deviation. By taking the sum of
the absolute values of a row, we defined the deviation score of a nucleus. In
order to omit contaminated droplets, we discarded nuclei with a score ≥40.

Since Tn5 inserts less efficiently into nucleosome-associated DNA,
ATAC-seq fragment size distribution shows periodicity [23]. Using the
program provided by Cusanovich et al.
(https://atlas.gs.washington.edu/mouse-atac/docs/), for each nucleus, we
computed the spectral densities for the 100–300 bp range, which is the pitch
of nucleosomes. In order to omit nuclei with an aberrant chromatin state, we
discarded nuclei below the 1st or above the 99th percentile in this score.

When two or more nuclei are contained in one droplet, called a
multiplet, those nuclei share the same barcode, violating the one-nucleus-to-
one-barcode assumption. We removed the barcodes for multiplets in two
ways. First, for the set of ATAC-seq fragments assigned to a barcode, we
inferred the number of homologous chromosomes from which the fragments
were derived. The number is analogous to ploidy, but differs because the
container is not a cell, but a droplet. By counting overlaps among the
fragments, the number of homologous chromosomes was estimated to be
either $2n$, $4n$ or $\geq 6n$ using the R package scPloidy (version 0.3.0). Since the
majority of hepatocytes in rat liver are diploids or tetraploids [55] and other

cell types in the liver are diploids, ≥$6n$ droplets were regarded as possible

multiplets and discarded. Second, we detected doublets formed by two

heterogeneous cell types using ArchR. Droplets with doublet enrichment

score >10 were discarded.

Chromatin accessibility peak discovery and nucleus clustering for

snATAC-seq

The Tn5 insertions observed in ATAC-seq occur at open chromatin regions,

such as promoters and enhancers, and the genome-wide profile of Tn5

insertions characterizes the types of cells. We clustered the nuclei based on

the similarity of Tn5 insertions. Firstly, we represented the Tn5 insertions in

snATAC-seq as a "tile matrix": a binary matrix indexed by tiles of 500 bp

width across the genome (rows) and nuclei (columns), coding the occurrence

(1) or non-occurrence (0) of Tn5 insertions. Using ArchR, we obtained a

matrix with 5,230,329 rows and 14,615 columns. We then grouped the nuclei

according to the similarity of corresponding columns (see next section). For

each group, pseudo-bulk ATAC-seq datasets were generated, and Tn5

insertion peaks were computed using ArchR and the MACS2 program (version

2.1.4) [56]. We obtained 599,035 chromatin accessibility peaks of width 501

bp.

Secondly, we represented the Tn5 insertions as a "peak matrix": a

binary matrix indexed by peaks (rows) and nuclei (columns), coding the

occurrence or non-occurrence of Tn5 insertions. Again, we grouped the nuclei

according to the similarity of corresponding columns. The nuclei clustering

based on the peak matrix was used in downstream analysis, as it would be

more accurate than the clustering based on the tile matrix.

Algorithm for dimensionality reduction and nucleus clustering

The tile and peak matrices are large binary matrices indexed by genomic

positions (rows) and nuclei (columns). Since the column vectors are

extremely high dimensional, we need to reduce the dimensions in order to

properly analyze the similarity among nuclei. For the dimensionality reduction,

we used latent semantic indexing [57] and singular value decomposition

(SVD). The dimensionality reduction was performed for all samples combined.

Since experimental conditions, including Tn5:DNA ratio, could differ among

the samples, we introduced a sample batch correction step. We applied the

same algorithm to the tile and peak matrices.

The input binary matrix (f_{ij}) with n columns was treated as the

term-document matrix in the latent semantic indexing literature. We defined

the inverse document frequency (IDF) as

$$g_i = \log_2 \left(\frac{n}{\sum_j f_{ij}} \right).$$

For the i-th row of the matrix weighted by IDF, the variance becomes

$$g_i{}^2 \left(\frac{\sum_j f_{ij}}{n} \right) \left(1 - \frac{\sum_j f_{ij}}{n} \right)$$

and ranges between zero and one. In order to restrict noise, we retained rows

with IDF-weighted variance >0.4 of the tile matrix and >0.1 of the peak

matrix. We defined the document normalization factor as

$$d_j = \frac{1}{\sqrt{\sum_i f_{ij}{}^2}}.$$

We applied sample batch correction independently to the rows of the

matrix. For row i, the normalized elements $d_j \cdot f_{ij}$ vary among nuclei j, but

the average across many nuclei in a sample is anticipated to be approximately

constant. The difference in the averages among samples was regarded as a

batch effect and was corrected by scaling. For the nucleus j belonging to the

sample s, the correction factor is

$$\gamma_{ij} = \frac{\text{Mean}\left[\{ d_{j'} \cdot f_{ij'} : 1 \leq j' \leq n \} \right]}{\text{Mean}\left[\{ d_{j'} \cdot f_{ij'} : j' \text{ belongs to sample } s \} \right]}. \quad (1)$$

When the denominator equals zero, γ_{ij} is set as one. The above correction

requires no knowledge of cell types. If cell type classification of nuclei is

available, the correction can be refined. For the nucleus j of cell type c

belonging to sample s, the correction factor becomes

$$\gamma_{ij} = \frac{\text{Mean}\big[\{d_{j'} \cdot f_{ij'} : j' \text{ is cell type } c\}\big]}{\text{Mean}\big[\{d_{j'} \cdot f_{ij'} : j' \text{ is cell type } c \text{ and belongs to sample } s\}\big]}. \quad (2)$$

Since we had no knowledge of cell type in advance, we first applied the batch

correction in (1) and then inferred coarse cell type classification. Using the

coarse classification, we applied the batch correction in (2). To avoid the

inflation of γ_{ij} under small denominator, for each nucleus j, we computed the

square of the third quartile of γ_{ij} divided by the second quartile, and imposed

this upper limit.

Dimensionality reduction and nucleus clustering were performed in

the following steps:

1. Compute the sample batch correction factor γ_{ij} in (1).

2. Compute the matrix $(g_i \cdot d_j \cdot f_{ij} \cdot \gamma_{ij})$ and centralize each row by

subtracting the average.

3. Compute truncated SVD, UDV^T, for the 30 largest singular values using

the R package irlba (version 2.3.3) [58].

4. Take the truncated SVD for the two largest singular values, project the

nuclei to the VD coordinates, and detect and discard outlier nuclei.

Outliers were detected based on the sum of the squared distance to the

five nearest neighbors.

5. Take the truncated SVD for the non-noise singular values, and project the

nuclei to the VD coordinates. As the 11th and later singular values

bottomed out and the corresponding columns of V appeared randomly

distributed in our data, we took the ten largest singular values.

6. Identify clusters of nuclei by computing the shared nearest-neighbor

graph with the R package Seurat (version 4.0.0) [59] and optimizing the

modularity with the Leiden algorithm (version 0.8.4) [60]. By merging

the clusters, define coarse clustering that clearly separates in the VD

coordinates. In our data, the coarse clusters corresponded to hepatocytes,

stellate cells, endothelial cells and white blood cells.

7. Compute the sample batch correction factor γ_{ij} in (2) by incorporating

the coarse clustering of nuclei.

8. Compute the matrix $\left(g_i \cdot d_j \cdot f_{ij} \cdot \gamma_{ij}\right)$ and centralize each row by

subtracting the average.

9. Compute truncated SVD, UDV^T, for the 30 largest singular values.

10. Take the truncated SVD for the non-noise singular values, and project the

nuclei to the VD coordinates. We took the eight largest singular values.

11. Identify clusters of nuclei by the shared nearest-neighbor modularity

optimization.

Cell type assignment for snATAC-seq

For each nuclei cluster obtained above, we assigned a cell type based on

chromatin accessibility at known marker genes. For each cluster, chromatin

accessibility at the TSS and gene body of the genes were visually inspected

using the browser of ArchR. We used the following marker genes in the rat:

• *Alb*, *Apoc3*, *Cyp2e1* and *Cyp2f4* for hepatocyte

• *Epcam* and *Krt19* for cholangiocyte

• *Stab2* for endothelial cell

• *Dcn*, *Des*, *Lrat* and *Acta2* for stellate cell

• *Csf3r* for granulocyte

• *Elane*, *Ly6g6c*, *Ly6g6d*, *Ly6g6e* and *Mpo* for neutrophil

• *Csf1r* for macrophage (including Kupffer cell)

• *Lyz2* for inflammatory macrophage

• *Marco* for non-inflammatory macrophage

• *Cd3g* and *Gata3* for T cell

• *Gzma* and *Prf1* for cytotoxic T cell and NK cell

• *Ebf1* for B cell

Data cleaning for bulk ATAC-seq

According to the ATAC-seq data processing workflow [61], we trimmed reads,

mapped reads to the reference genome, removed mitochondrial DNA reads,

subsampled to equate the library complexity, and removed PCR duplicates.

Inferring cell type composition of bulk tissue samples

We inferred the cell type composition of liver tissue samples by comparing

their bulk ATAC-seq data with the reference data generated by snATAC-seq.

As references, pseudo-bulk ATAC-seq data of pure cell types were computed

from snATAC-seq results. We combined several published deconvolution

methods. Let the indices be i for a chromatin accessibility peak, j for a cell

type, and k for a sample, either assayed in bulk, in single nuclei or both.

By adding Tn5 insertion counts of snATAC-seq over a_{jk} nuclei of cell

type j from sample k, we obtain Tn5 insertion count u_{ijk} at peak i. The

counts across all peaks are represented as a vector $\boldsymbol{u}_{jk}$ in bold font, in which

the index i is dropped. We quantile normalize the vectors $\boldsymbol{u}_{jk}$ and obtain

vectors $\boldsymbol{v}_{jk}$. Although quantile normalization is not a linear transformation,

we can regard the total number of nuclei in $\boldsymbol{v}_{jk}$ as

$$b_{jk} = \frac{\sum_i v_{ijk}}{\sum_i u_{ijk}} \cdot a_{jk}.$$

We factorize the count as

$$b_{jk} \approx c_j \cdot d_k,$$

where c_j represents how many nuclei are included in the normalized ATAC-

seq library for pure cell type j and d_k adjusts for experimental batch effect

between samples. Specifically, we solve a linear regression whose dependent

variable is the logarithm of b_{jk} and independent variables are the indicator

functions of j and k, and take the exponential of the obtained coefficients.

For the solution to be unique, we set $d_1 = 1$. We represent by C the diagonal

matrix with c_j in the diagonal. The values of c_j for hepatocytes were half the

values for other cell types due to higher ploidy of hepatocytes.

We used the DeconPeaker software (version 1.0) [7] to select the

peaks (rows) that discriminate cell types. DeconPeaker selects discriminating

peaks based on t-tests while minimizing the conditional number. Let $\tilde{v}_{jk}$ be

the vector truncated from v_{jk} accordingly. The vectors $\tilde{v}_{jk}$ across the

samples from the same cell type j are approximately equal, and their

average becomes the reference vector w_j for chromatin accessibility:

$$w_{ij} = \text{mean}_k\{\tilde{v}_{ijk}\}.$$

We represent by W the matrix whose columns are w_j. By regarding the

samples as replicates, the error variance for peak i was estimated as

$$e_i^2 = \text{mean}_j\left\{\text{var}_k\{\tilde{v}_{ijk}\}\right\}.$$

For samples assayed in snATAC-seq, we can generate pseudo-bulk

chromatin accessibility for mixtures based on W. For sample k, the

proportion of cell type j is $a_{jk}/(\sum_{j'} a_{j'k})$, and the mixture becomes

$$x_k = \sum_j \frac{a_{jk}}{\left(\sum_{j'} a_{j'k}\right)} \cdot \frac{1}{c_j} \cdot w_j.$$

Here, the vector is scaled to one nucleus (under the experimental condition

of sample $k = 1$). In other words, each element is the average count of Tn5

insertions at the site in one nucleus.

The bulk ATAC-seq results of mixture samples are quantile normalized

and represented as vector y_k for sample k. The elements y_{ik} are taken only

for the peaks i that were selected by DeconPeaker. Five samples were

assayed both in bulk and in single nuclei, for which we have vectors x_k and

y_k. Although, in principle, the two vectors should be proportional, the ratio

y_{ik}/x_{ik} actually varies by peaks, due to measurement differences between

different assay technologies. Correction of between-peak variability can

improve deconvolution [62]. We compute the correction factor as

728
$$f_i = \exp\left(\text{mean}_k\left\{\log\left(\frac{y_{ik}}{x_{ik}}\right)\right\}\right).$$

To avoid over-correction, we cap f_i to range between Q_1^2/Q_2 and Q_3^2/Q_2,

where Q_1, Q_2, Q_3 are the quartiles of the original value. We represent by F the

diagonal matrix with f_i in the diagonal. A bulk ATAC-seq experiment y_k after

correction becomes $F^{-1}y_k$. Notice that the vector is scaled to one nucleus.

For deconvolution of a bulk sample k, we first compute $\boldsymbol{\theta}_k$ that

approximates

735
$$F^{-1}y_k \approx W\boldsymbol{\theta}_k$$

736   under $\theta_{jk} \geq 0$. Using robust linear regression [63], we solve

737
$$F^{-1}y_k = W\boldsymbol{\theta}_k + \boldsymbol{\varepsilon},$$

738
$$\varepsilon_i \sim N\left(0, e_i^2\sigma^2\right),$$

where σ^2 is the parameter for error variance. Peaks with large variability e_i^2

are allowed to have a larger variance [64]. To avoid over-correction, we cap

e_i^2 to range between Q_1^2/Q_2 and Q_3^2/Q_2, where Q_1, Q_2, Q_3 are the quartiles of

the original value. The negative elements of the estimate $\widehat{\boldsymbol{\theta}_k}$ are set to zero.

When computing $\boldsymbol{\theta}_k$, the independent variable $F^{-1}y_k$ was scaled to one

nucleus, and the regressors w_j were scale to c_j nuclei. We multiply $\widehat{\boldsymbol{\theta}_k}$ by C

and obtain $C\widehat{\boldsymbol{\theta}_k}$, which is the fractional number of nuclei from each cell types

included per nucleus of sample k, which in spirit is the cell type composition.

Although the sum of the elements in $C\widehat{\boldsymbol{\theta}_k}$ was approximately equal to one, in

order to be precise, we rescale to obtain the cell type composition for sample

k as

750
$$\frac{C\widehat{\boldsymbol{\theta}_k}}{\sum_j c_j\widehat{\theta_{jk}}}.$$

Comparing cell type composition among dietary conditions

For each cell type, the imputed composition in bulk samples was compared

among dietary conditions. We performed multiple regression where the

dependent variable was the composition and independent variables were

indicators of dietary condition, with the normal diet being the reference. Using

Bonferroni's correction for multiple testing of seven cell types and three

conditions (apart from normal diet), the significance level was set to

$0.05/[7 \times 3]$.

Quantifying gene expression in snATAC-seq

For each nucleus, the expression level of a gene was inferred from the number

of Tn5 insertions in the gene body and vicinity. The targeted biological

phenomenon is the chromatin opening of the gene rather than the amount of

mRNA in a nucleus or cell. Using ArchR, we computed the Gene Score Matrix,

which is indexed by genes (rows) and nuclei (columns) and represents the

insertion counts. As the matrix is analogous to the count matrix obtained in

single-cell RNA-seq, we applied library size normalization of the R package

Linnorm (version 2.16.0) [65] and imputation of the R package SAVER

(version 1.1.2) [66]. Before imputation, 47% of the matrix elements were

zero, and after imputation, 0.2% were set to the lowest value (0.001). We

excluded low expression genes, where the lowest value occurred in ≥1% of

nuclei. We took log2 of the matrix and applied quantile normalization. The

obtained matrix was used for inspecting cell type marker genes. We applied

SVD and subtracted the first two components from the matrix, which captured

the sample batch effect. Although the two components also captured major

cell type differences, we anticipated that the removal would not harm

downstream analysis, which compares nuclei of the same cell type between

samples.

Quantifying TF binding in snATAC-seq

If the DNA sequence of a chromatin accessibility peak includes the motif of a

TF, Tn5 insertion in the peak suggests the binding of the corresponding TF.

For each nucleus, by aggregating over all peaks, we can infer the genome-

wide binding of the TF. This is obtained as the chromVAR [67] deviation z-

score in ArchR. The CIS-BP database of TF binding motifs [68] was used. In

the same way as we did for the Gene Score Matrix, we applied quantile

normalization to the Motif Matrix and subtracted the first two components of

SVD.

Gene set enrichment analysis of differentially expressed genes under

HFD

To seek which biological processes and/or pathways are differentially

expressed under a particular dietary intervention, we performed gene set

enrichment analysis. The target GS's were obtained from the Molecular

Signatures Database (MSigDB) (version 7.4) [69] using the R package

msigdbr (version 7.4.1). Among the datasets in MSigDB, we used Gene

Ontology for Biological Process, Hallmark gene sets, KEGG pathways,

Reactome pathways, and WikiPathways. Using the sample batch-corrected

Gene Score Matrix, we compared nuclei from a pair of dietary conditions, for

example, the hepatocytes from 4 weeks of HFD *vs.* the normal diet. The

statistical significance of gene set enrichment was tested using the R package

PADOG (version 1.34.0) [70] and by performing permutation for 10,000

times or 20 times the number of GS's in a dataset, whichever was larger.

Within each dataset, the GS's that attained a false discover rate <0.05 were

considered statistically significant. Significant GS's were visually inspected

using the EnrichmentMap App (version 3.3.3) of the Cytoscape software

(version 3.8.2).

Discovering major modules of TF regulation and assigning biological

processes

To elucidate a global picture of TF regulation, we inferred the regulatory

impacts of all TFs on all genes and then extracted major modules of the

regulatory relationship between TFs and genes. For each gene, we computed

its regulatory TFs using the R package GENIE3 (version 1.14.0) [71], which

implements a random forest algorithm and thus can model nonlinear

regulation. The algorithm aims to predict the expression level of the gene

based on the binding of TFs and dietary conditions. As for the number of

candidate regulators randomly selected at each node of the random forests,

we chose the square root of the number of predictors, as suggested in the

GENIE3 publication. GENIE3 quantified the impacts of regulators by the

proportion of explained variance. By combining the results from all genes, we

obtained a regulator-regulatee matrix, in which the rows correspond to TFs

and dietary conditions, the columns correspond to genes, and an element

represents the importance of a TF to a gene.

Next, we applied the nonnegative matrix factorization implemented in

the R package NMF (version 0.23.0) [72] to the regulator-regulatee matrix.

We used the Brunet algorithm [73] with 100 random starts. The number of

factors was chosen based on data as the inflection point of residual error [74].

In the output of NMF, the membership of TFs (regulators) and genes

(regulatees) to a factor was defined by nonnegative weight. We called the

factors as "modules" of TF regulation.

We discovered biological processes that characterize a module by

applying gene set enrichment analysis to genes belonging to the module. For

each module of TF regulation, we tested if heavily weighted genes were

enriched in some GS. For each GS (see above section), we computed the P-

value and t-statistics to test if genes in the given GS had a larger weight than

other genes. To account for multiple testing by the GS's, we performed 100

trials of permuting gene names and computed family-wise error rates. The

GS's that attained a family-wise error rate of <0.05 were considered

statistically significant. The permutation test does not require the assumption

of normality.

GS activity score in single nuclei

To infer the activity of a GS in single nuclei, we computed the "average" level

of TF binding and gene expression, with regard to the TFs/genes included in

the GS. Formally, we defined the weight of TFs/genes as one if it was included

in the GS and as zero otherwise. We then performed SVD for the TF binding

and gene expression matrix incorporating the weight. In the Motif Matrix (see

above), each TF (row) was standardized by subtracting the mean and being

divided by the standard deviation and then multiplied by the weight of the TF.

Similarly, in the Gene Score Matrix (see above), each gene (row) was

standardized by subtracting the mean and being divided by the standard

deviation and then multiplied by the weight of the gene. By stacking the two

matrices, we obtained a large matrix, in which the rows correspond to TFs

and genes and the columns correspond to nuclei. Finally, SVD was applied

and the first right-singular vector was defined as the GS activity score in

single nuclei.

Although the abovementioned activity score can always be computed,

it would be meaningless if the TFs/genes in the GS were not co-expressed.

Therefore, we conducted a statistical test to exclude such meaningless GS.

We checked whether the TFs/genes in the GS were mutually correlated more

than by chance. We computed the null distribution of the first singular value

by performing 100 trials of the SVD procedure but with the weight of TFs

permutated and the weight of genes permutated and verified whether the

actual first singular value could attain an empirical P-value of <0.05. The

permutation test does not require the assumption of normality.

The principal component approach was adapted from the PLAGE

program [24] that was developed to compute the pathway activity score of a

GS in bulk tissue specimens. Instead, we applied to nuclei. Compared with

the number of specimens, the large number of nuclei enabled significance

testing by permutation.

Abbreviations

GS, gene set; IDF, inverse document frequency; NAFL, non-alcoholic fatty

liver; NAFLD, non-alcoholic fatty liver disease; NASH, non-alcoholic

steatohepatitis; snATAC-seq, single-nucleus ATAC-seq; SVD, singular value

decomposition; TF, transcription factor

Declarations

Ethics approval and consent to participate

Not applicable.

Consent for publication

Not applicable.

Availability of data and materials

The single-nucleus and bulk ATAC-seq raw data generated during this study

are available at DDBJ (DRA DRA014511 and DRA014458 within BioProject

PRJDB13870). Datasets after cleaning are available at figshare (DOI

10.6084/m9.figshare.20236509). The code generated during this study is

available at GitHub https://github.com/fumi-

github/rat_singlecell_liver_ArchR

Competing interests

The authors declare no competing interests.

Funding

This work was supported by the NCGM Intramural Research Fund (20A1013)

and by JSPS KAKENHI [grant number JP16K07218]. The funding body had no

role in the design of the study and collection, analysis and interpretation of

data and in writing the manuscript.

Authors' contributions

Conceptualization, F.T., Y-Q.L., H.S-F. and N.K.;

Software, F.T.;

Formal Analysis, F.T.;

Investigation, F.T., Y-Q.L., H.S-F., M.I., M.Y.A. and E.K.;

Writing – Original Draft, F.T.;

Writing – Review & Editing, all authors;

Supervision, N.K.

All authors have read and approved the manuscript.

Acknowledgments

The authors thank the Center for Omics and Bioinformatics, Graduate School

of Frontier Sciences, University of Tokyo and Azenta for their assistance in

snATAC-seq. The authors thank Enago (www.enago.jp) for the English

language review.

FIGURE LEGENDS

Figure 1

Cell types observed in the liver and their composition

(A) Experimental design.

(B) UMAP plot of the observed nuclei. The number of observed nuclei is shown

in parentheses.

(C) Expression level of cell type marker genes in nuclei (vertical axis) is

compared between nuclei clusters (horizonal axis) in violin plots.

(D) Validation of deconvolution based on ATAC-seq using the samples assayed

in both single-nucleus and bulk ATAC-seq. The true cell type proportion

counted in snATAC-seq (horizontal axis) is compared with the proportion

inferred based on bulk ATAC-seq (vertical axis).

(E) Proportion of nuclei from a specific cell type (vertical axis) is compared

between dietary conditions (horizontal axis). The proportion was measured

by bulk tissue ATAC-seq (n=4 samples per condition).

Figure 2

Biological processes that were differentially expressed under dietary

interventions

The heatmap shows whether a biological process (row) was differentially

expressed in a cell type under a dietary intervention compared with the

normal diet (column). The coloring indicates statistical significance of gene

set enrichment analysis: red if significant adjusting for multiple testing,

brown if nominally significant at P<0.01, light brown if P<0.05, and white

otherwise.

Figure 3

Major modules of TF regulation in hepatocytes

(A) The central heatmap shows the regulatory impact of TF binding (columns)

on genes (rows). Major modules are highlighted by red rectangles. The

heatmap at the bottom shows the membership of TFs (columns) to modules

(rows). The heatmap on the right shows the membership of genes (rows) to

modules (columns). The heatmaps only include the TFs/genes with

membership ranking in the top five and membership weight more than half

of the top.

(B) Gene set enrichment analysis for genes belonging to the modules.

Excessive overlap between genes belonging to a module and the GS for a

biological process (horizontal axis) was tested, and the P-value was plotted

(vertical axis). Bars are colored black if statistically significant, adjusting for

multiple testing, and gray otherwise. Biological processes that were

significant for at least one module are shown.

Figure 4

Major modules of TF regulation in endothelial cells

See legend in Figure 3.

Figure 5

Major modules of TF regulation in stellate cells

See legend in Figure 3.

Figure 6

Major modules of TF regulation in macrophages

See legend in Figure 3.

Figure 7

Core genes for the TNFα signaling via NF-κB processes are extracted based

on correlation with GS activity score and protein-protein interaction

(A) Analysis in hepatocytes using the GS for TNFα signaling via NF-κB. In the

top panel, the GS activity score in single nuclei (vertical axis) is compared

between dietary conditions (horizontal axis) in violin plots. Box plots within

the violins indicate quartiles. The score distribution under a diet intervention

was compared with that under normal diet condition by the area under the

receiver operating characteristic curve: * and ** indicate larger than 0.75

and 0.9, respectively. In the remaining panels, TFs and genes are plotted by

the correlation with the GS activity score (horizontal axis) and the total

amount of protein-protein interactions with the GS (vertical axis). The points

for TFs/genes having protein-protein interactions within the top 5% are

colored red, and the others are colored blue. The TFs/genes in the GS (middle

panel) and not in the GS (bottom panel) are plotted separately; the former

have more abundant protein-protein interactions. TFs are labeled in all capital

letters. The GS was taken from the MSigDB Hallmark database. Protein-

protein interactions were taken from the STRING database (version 11.0).

(B) Analysis in endothelial cells.

(C) Analysis in macrophages.

Figure 8

Modules of global TF regulation, biological processes, and core genes were

detected by the data-driven approach

See text for the full list of core genes.

REFERENCES

1. Friedman SL, Neuschwander-Tetri BA, Rinella M, Sanyal AJ. Mechanisms of
NAFLD development and therapeutic strategies. Nat Med. 2018;24:908-22.

2. Chu AL, Schilling JD, King KR, Feldstein AE. The Power of Single Cell Analysis for
the Study of Liver Pathobiology. Hepatology. 2020;:hep.31485-23.

3. Ramachandran P, Dobie R, Wilson-Kanamori JR, Dora EF, Henderson BEP, Luu
NT, et al. Resolving the fibrotic niche of human liver cirrhosis at single-cell level.
Nature. 2019;575:512-8.

4. Xiong X, Kuang H, Ansari S, Liu T, Gong J, Wang S, et al. Landscape of
Intercellular Crosstalk in Healthy and NASH Liver Revealed by Single-Cell
Secretome Gene Analysis. Molecular cell. 2019;75:644-660.e5.

5. Sturm G, Finotello F, Petitprez F, Zhang JD, Baumbach J, Fridman WH, et al.
Comprehensive evaluation of transcriptome-based cell-type quantification methods
for immuno-oncology. Bioinformatics (Oxford, England). 2019;35:i436-45.

6. Teschendorff AE, Zheng SC. Cell-type deconvolution in epigenome-wide
association studies: a review and recommendations. Epigenomics. 2017;9:757-68.

7. Li H, Sharma A, Luo K, Qin ZS, Sun X, Liu H. DeconPeaker, a Deconvolution Model
to Identify Cell Types Based on Chromatin Accessibility in ATAC-Seq Data of
Mixture Samples. Frontiers in Genetics. 2020;11:392.

8. Kiki I, Altunkaynak BZ, Altunkaynak ME, Vuraler O, Unal D, Kaplan S. Effect of
High Fat Diet on the Volume of Liver and Quantitative Feature of Kupffer Cells in
the Female Rat: A Stereological and Ultrastructural Study. Obes Surg. 2007;17:1381.

9. Zhong L, Huang L, Xue Q, Liu C, Xu K, Shen W, et al. Cell-specific elevation of
Runx2 promotes hepatic infiltration of macrophages by upregulating MCP-1 in high-
fat diet-induced mice NAFLD. J Cell Biochem. 2019;120:11761-74.

10. Zhang F, Jiang WW, Li X, Qiu XY, Wu Z, Chi YJ, et al. Role of intrahepatic B cells

in non-alcoholic fatty liver disease by secreting pro-inflammatory cytokines and

regulating intrahepatic T cells. J Digest Dis. 2016;17:464–74.

11. Barrow F, Khan S, Fredrickson G, Wang H, Dietsche K, Parthiban P, et al.

Microbiota-Driven Activation of Intrahepatic B Cells Aggravates NASH Through

Innate and Adaptive Signaling. Hepatology Baltim Md. 2021;74:704–22.

12. Hess J, Angel P, Schorpp-Kistner M. AP-1 subunits: quarrel and harmony among

siblings. J Cell Sci. 2004;117:5965–73.

13. Zinzalla G. A New Way Forward in Cancer Drug Discovery: Inhibiting the

SWI/SNF Chromatin Remodelling Complex. Chembiochem. 2016;17:677–82.

14. Halpern KB, Shenhav R, Matcovitch-Natan O, Tóth B, Lemze D, Golan M, et al.

Single-cell spatial reconstruction reveals global division of labour in the mammalian

liver. Nature. 2017;542:352–6.

15. Sun J, Weis WI. Biochemical and Structural Characterization of β-Catenin

Interactions with Nonphosphorylated and CK2-Phosphorylated Lef-1. J Mol Biol.

2011;405:519–30.

16. Behari J, Yeh T-H, Krauland L, Otruba W, Cieply B, Hauth B, et al. Liver-Specific

β-Catenin Knockout Mice Exhibit Defective Bile Acid and Cholesterol Homeostasis

and Increased Susceptibility to Diet-Induced Steatohepatitis. Am J Pathology.

2010;176:744–53.

17. Rasala BA, Orjalo AV, Shen Z, Briggs S, Forbes DJ. ELYS is a dual

nucleoporin/kinetochore protein required for nuclear pore assembly and proper cell

division. Proc National Acad Sci. 2006;103:17801–6.

18. Yu X, Chen X, Zheng XD, Zhang J, Zhao X, Liu Y, et al. Growth Differentiation

Factor 11 Promotes Abnormal Proliferation and Angiogenesis of Pulmonary Artery

Endothelial Cells. Hypertension. 2018;71:729–41.

19. Tran MTN, Hamada M, Jeon H, Shiraishi R, Asano K, Hattori M, et al. MafB is a

critical regulator of complement component C1q. Nat Commun. 2017;8:1700.

20. Chistiakov DA, Myasoedova VA, Revin VV, Orekhov AN, Bobryshev YV. The
impact of interferon-regulatory factors to macrophage differentiation and
polarization into M1 and M2. Immunobiology. 2018;223:101–11.

21. Salcedo R, Wasserman K, Young HA, Grimm MC, Howard OMZ, Anver MR, et al.
Vascular Endothelial Growth Factor and Basic Fibroblast Growth Factor Induce
Expression of CXCR4 on Human Endothelial Cells In Vivo Neovascularization
Induced by Stromal-Derived Factor-1α. Am J Pathology. 1999;154:1125–35.

22. Date D, Das R, Narla G, Simon DI, Jain MK, Mahabeleshwar GH. Kruppel-like
Transcription Factor 6 Regulates Inflammatory Macrophage Polarization. J Biol
Chem. 2014;289:10318–29.

23. Cusanovich DA, Hill AJ, Aghamirzaie D, Daza RM, Pliner HA, Berletch JB, et al. A
Single-Cell Atlas of In Vivo Mammalian Chromatin Accessibility. Cell.
2018;174:1309–1324.e18.

24. Tomfohr J, Lu J, Kepler TB. Pathway level analysis of gene expression using
singular value decomposition. Bmc Bioinformatics. 2005;6:225–225.

25. Kakino S, Ohki T, Nakayama H, Yuan X, Otabe S, Hashinaga T, et al. Pivotal Role
of TNF-a in the Development and Progression of Nonalcoholic Fatty Liver Disease
in a Murine Model. Horm Metab Res. 2017;50:80–7.

26. Locatelli I, Sutti S, Vacchiano M, Bozzola C, Albano E. NF-κB1 deficiency
stimulates the progression of non-alcoholic steatohepatitis (NASH) in mice by
promoting NKT-cell-mediated responses. Clin Sci. 2012;124:279–87.

27. Roos B de, Rungapamestry V, Ross K, Rucklidge G, Reid M, Duncan G, et al.
Attenuation of inflammation and cellular stress-related pathways maintains insulin
sensitivity in obese type I interleukin-1 receptor knockout mice on a high-fat diet.
Proteomics. 2009;9:3244–56.

28. Sawitza I, Kordes C, Reister S, Häussinger D. The niche of stellate cells within
rat liver. Hepatology. 2009;50:1617–24.

29. Boujedidi H, Robert O, Bignon A, Cassard-Doulcier A-M, Renoud M-L, Gary-
Gouy H, et al. CXCR4 dysfunction in non-alcoholic steatohepatitis in mice and
patients. Clin Sci. 2014;128:257-67.

30. Potoupni V, Georgiadou M, Chatzigriva E, Polychronidou G, Markou E, Gakis CZ,
et al. Circulating tumor necrosis factor-α levels in non-alcoholic fatty liver disease:
A systematic review and a meta-analysis. J Gastroen Hepatol. 2021;36:3002-14.

31. Seo YY, Cho YK, Bae J-C, Seo MH, Park SE, Rhee E-J, et al. Tumor Necrosis
Factor-α as a Predictor for the Development of Nonalcoholic Fatty Liver Disease:
A 4-Year Follow-Up Study. Endocrinol Metab. 2013;28:41-5.

32. Goel R, Boylan B, Gruman L, Newman PJ, North PE, Newman DK. The
proinflammatory phenotype of PECAM-1-deficient mice results in atherogenic diet-
induced steatohepatitis. Am J Physiol-gastr L. 2007;293:G1205-14.

33. Zhang C, Jia Y, Liu B, Wang G, Zhang Y. TLR4 knockout upregulates the
expression of Mfn2 and PGC-1α in a high-fat diet and ischemia-reperfusion mice
model of liver injury. Life Sci. 2020;254:117762.

34. Cepero-Donates Y, Lacraz G, Ghobadi F, Rakotoarivelo V, Orkhis S, Mayhue M,
et al. Interleukin-15-mediated inflammation promotes non-alcoholic fatty liver
disease. Cytokine. 2016;82:102-11.

35. Pérez-Martínez L, Pérez-Matute P, Aguilera-Lizarraga J, Rubio-Mediavilla S,
Narro J, Recio E, et al. Maraviroc, a CCR5 antagonist, ameliorates the development
of hepatic steatosis in a mouse model of non-alcoholic fatty liver disease (NAFLD).
J Antimicrob Chemoth. 2014;69:1903-10.

36. Patouraux S, Rousseau D, Bonnafous S, Lebeaupin C, Luci C, Canivet CM, et al.
CD44 is a key player in non-alcoholic steatohepatitis. J Hepatol. 2017;67:328-38.

37. Li ZZ, Berk M, McIntyre TM, Feldstein AE. Hepatic Lipid Partitioning and Liver
Damage in Nonalcoholic Fatty Liver Disease ROLE OF STEAROYL-CoA
DESATURASE. J Biol Chem. 2009;284:5637-44.

38. Zhang Y, Lu Z, Zeng W, Zhao J, Zhou X. Two sides of NNMT in alcoholic and
nonalcoholic fatty liver development. J Hepatol. 2020;74:1250-3.

39. Chen C, Li H, Song J, Zhang C, Li M, Mao Y, et al. Role of Apolipoprotein A1 in
PPAR Signaling Pathway for Nonalcoholic Fatty Liver Disease. Ppar Res.
2022;2022:4709300.

40. Yang MH, Sung J, Gwak G-Y. The associations between apolipoprotein B, A1,
and the B/A1 ratio and nonalcoholic fatty liver disease in both normal-weight and
overweight Korean population. J Clin Lipidol. 2016;10:289-98.

41. Rigotti FDA, Wiedmann L, Hubert MO, Vacca M, Hasan SS, Moll I, et al.
Semaphorin 3C exacerbates liver fibrosis. Biorxiv. 2022;:2021.07.29.454292.

42. Cao S, Yaqoob U, Das A, Shergill U, Jagavelu K, Huebert RC, et al. Neuropilin-1
promotes cirrhosis of the rodent and human liver by enhancing PDGF/TGF-β
signaling in hepatic stellate cells. J Clin Invest. 2010;120:2379-94.

43. Yagai T, Miyajima A, Tanaka M. Semaphorin 3E Secreted by Damaged
Hepatocytes Regulates the Sinusoidal Regeneration and Liver Fibrosis during Liver
Regeneration. Am J Pathology. 2014;184:2250-9.

44. Binder JX, Pletscher-Frankild S, Tsafou K, Stolte C, O'Donoghue SI, Schneider
R, et al. COMPARTMENTS: unification and visualization of protein subcellular
localization evidence. Database J Biological Databases Curation. 2014;2014:bau012.

45. Ajmera V, Perito ER, Bass NM, Terrault NA, Yates KP, Gill R, et al. Novel plasma
biomarkers associated with liver disease severity in adults with nonalcoholic fatty
liver disease. Hepatology. 2017;65:65-77.

46. Koutsogiannis D, Summers K, George B, Adams P, Marotta P, Chakrabarti S.
Identification of Serum Biomarkers in End Stage Liver Disease. Open Biomarkers J.
2010;3:1-6.

47. Krawczyk M, Zimmermann S, Hess G, Holz R, Dauer M, Raedle J, et al. Panel of
three novel serum markers predicts liver stiffness and fibrosis stages in patients
with chronic liver disease. Plos One. 2017;12:e0173506.

48. An X, Liu J, Zheng X, Dou Z, Li Y, Suo Y, et al. Serum HGF ,PCIII and PLT are
Noninvasive Markers for the diagnosis of nonalcoholic fatty liver disease. 2019.
https://doi.org/10.21203/rs.2.18035/v1.

49. Page S, Birerdinc A, Estep M, Stepanova M, Afendy A, Petricoin E, et al.
Knowledge-Based Identification of Soluble Biomarkers: Hepatic Fibrosis in NAFLD
as an Example. Plos One. 2013;8:e56009.

50. Herck MAV, Vonghia L, Francque SM. Animal Models of Nonalcoholic Fatty Liver
Disease—A Starter's Guide. Nutrients. 2017;9:1072.

51. Corces MR, Trevino AE, Hamilton EG, Greenside PG, Sinnott-Armstrong NA,
Vesuna S, et al. An improved ATAC-seq protocol reduces background and enables
interrogation of frozen tissues. Nature Methods. 2017;14:959-62.

52. Lareau CA, Ma S, Duarte FM, Buenrostro JD. Inference and effects of barcode
multiplets in droplet-based single-cell assays. Nature Communications. 2020;11:866.

53. Granja JM, Corces MR, Pierce SE, Bagdatli ST, Choudhry H, Chang HY, et al.
ArchR is a scalable software package for integrative single-cell chromatin
accessibility analysis. Nature Genetics. 2021;53:403 – 411.

54. Orchard P, Kyono Y, Hensley J, Kitzman JO, Parker SCJ. Quantification,
Dynamic Visualization, and Validation of Bias in ATAC-Seq Data with ataqv. Cell
Systems. 2020;10:298-306.e4.

55. Katsuda T, Hosaka K, Matsuzaki J, Usuba W, Prieto-Vila M, Yamaguchi T, et al.
Transcriptomic Dissection of Hepatocyte Heterogeneity: Linking Ploidy, Zonation,
and Stem/Progenitor Cell Characteristics. Cellular and Molecular Gastroenterology
and Hepatology. 2020;9:161-83.

56. Zhang Y, Liu T, Meyer CA, Eeckhoute J, Johnson DS, Bernstein BE, et al. Model-
based Analysis of ChIP-Seq (MACS). Genome Biol. 2008;9:R137-R137.

57. Berry MW, Browne M. Understanding Search Engines. 2nd edition. SIAM; 2005.

58. Baglama J, Reichel L. Augmented Implicitly Restarted Lanczos Bidiagonalization
Methods. Siam J Sci Comput. 2005;27:19-42.

59. Hao Y, Hao S, Andersen-Nissen E, Mauck WM, Zheng S, Butler A, et al.
Integrated analysis of multimodal single-cell data. Cell. 2021;184:3573-3587.e29.

60. Traag VA, Waltman L, Eck NJ van. From Louvain to Leiden: guaranteeing well-
connected communities. Sci Rep-uk. 2019;9:5233.

61. Reske JJ, Wilson MR, Chandler RL. ATAC-seq normalization method can
significantly affect differential accessibility analysis and interpretation. Epigenetics &
chromatin. 2020;13:22.

62. Jew B, Alvarez M, Rahmani E, Miao Z, Ko A, Garske KM, et al. Accurate
estimation of cell composition in bulk expression through robust integration of
single-cell information. Nature Communications. 2020;11:1971.

63. Teschendorff AE, Breeze CE, Zheng SC, Beck S. A comparison of reference-
based algorithms for correcting cell-type heterogeneity in Epigenome-Wide
Association Studies. BMC Bioinformatics. 2017;18:105.

64. Wilson DR, Jin C, Ibrahim JG, Sun W. ICeD-T Provides Accurate Estimates of
Immune Cell Abundance in Tumor Samples by Allowing for Aberrant Gene
Expression Patterns. J Am Stat Assoc. 2019;115:1-23.

65. Yip SH, Wang P, Kocher J-PA, Sham PC, Wang J. Linnorm: improved statistical
analysis for single cell RNA-seq expression data. Nucleic Acids Research.
2017;45:e179.

66. Huang M, Wang J, Torre E, Dueck H, Shaffer S, Bonasio R, et al. SAVER: gene
expression recovery for single-cell RNA sequencing. Nature Methods. 2018;15:539 –
542.

67. Schep AN, Wu B, Buenrostro JD, Greenleaf WJ. chromVAR: inferring
transcription-factor-associated accessibility from single-cell epigenomic data. Nat
Methods. 2017;14:975-8.

68. Weirauch MT, Yang A, Albu M, Cote AG, Montenegro-Montero A, Drewe P, et al.
Determination and Inference of Eukaryotic Transcription Factor Sequence
Specificity. Cell. 2014;158:1431-43.

69. Subramanian A, Tamayo P, Mootha VK, Mukherjee S, Ebert BL, Gillette MA, et al.
Gene set enrichment analysis: A knowledge-based approach for interpreting

genome-wide expression profiles. Proceedings of the National Academy of Sciences
of the United States of America. 2005;102:15545–50.

70. Tarca AL, Draghici S, Bhatti G, Romero R. Down-weighting overlapping genes
improves gene set analysis. Bmc Bioinformatics. 2012;13:136.

71. Huynh-Thu VA, Irrthum A, Wehenkel L, Geurts P. Inferring Regulatory Networks
from Expression Data Using Tree-Based Methods. Plos One. 2010;5:e12776.

72. Gaujoux R, Seoighe C. A flexible R package for nonnegative matrix factorization.
BMC Bioinformatics. 2010;11:367.

73. Brunet J-P, Tamayo P, Golub TR, Mesirov JP. Metagenes and molecular pattern
discovery using matrix factorization. Proc National Acad Sci. 2004;101:4164–9.

74. Hutchins LN, Murphy SM, Singh P, Graber JH. Position-dependent motif
characterization using non-negative matrix factorization. Bioinformatics.
2008;24:2684–90.

1195

Figure 1

Figure 2

Legend: 3 = FDR <0.05; 2 = P <0.01; 1 = P <0.05. Compared to normal diet.

Database	Gene set	Hepatocyte HFD 4w	Hepatocyte HFD 8w	Hepatocyte washout	Endothelial cell HFD 4w	Endothelial cell HFD 8w	Endothelial cell washout	Stellate cell HFD 4w	Stellate cell HFD 8w	Stellate cell washout	Macrophage HFD 4w	Macrophage HFD 8w	Macrophage washout
GOBP	STEROID_METABOLIC_PROCESS	3	3	3	2	2	2	2	0	2	0	2	0
REACTOME	CHOLESTEROL_BIOSYNTHESIS	2	0	3	2	0	1	3	0	0	0	1	0
REACTOME	REGULATION_OF_CHOLESTEROL_BIOSYNTHESIS_BY_SREBP_SREBF	2	0	3	0	0	0	1	0	0	0	0	0
GOBP	FATTY_ACID_METABOLIC_PROCESS	3	2	3	0	1	2	2	0	0	2	1	0
GOBP	LIPID_STORAGE	3	3	3	0	0	1	0	0	0	1	2	0
KEGG	PPAR_SIGNALING_PATHWAY	1	1	3	0	1	0	0	0	0	0	0	0
H	XENOBIOTIC_METABOLISM	2	0	3	0	1	0	0	0	0	0	0	0
H	TNFA_SIGNALING_VIA_NFKB	1	3	0	2	2	0	0	0	0	2	1	0
GOBP	CYTOKINE_PRODUCTION	1	3	0	2	1	0	0	0	0	2	0	3
GOBP	RESPONSE_TO_CYTOKINE	2	3	0	2	0	2	1	1	0	3	0	3
REACTOME	INTERLEUKIN_1_SIGNALING	2	3	0	1	0	0	0	0	0	0	0	0
REACTOME	INTERLEUKIN_10_SIGNALING	1	3	0	2	2	1	3	1	0	0	0	0
REACTOME	INFLAMMASOMES	1	3	0	0	0	0	0	0	0	0	0	0
GOBP	ADAPTIVE_IMMUNE_RESPONSE	0	2	0	1	0	1	0	0	0	3	2	3
KEGG	APOPTOSIS	2	3	0	0	0	0	0	0	0	0	0	0
GOBP	ACTIN_FILAMENT_BASED_PROCESS	1	2	0	0	0	0	0	3	0	1	0	2
GOBP	CELL_MIGRATION	3	3	1	2	1	2	0	3	2	2	2	3
GOBP	INSULIN_LIKE_GROWTH_FACTOR_RECEPTOR_SIGNALING_PATHWAY	1	0	3	0	0	0	0	2	0	0	0	0

Figure 3. Hepatocyte

Figure 4. Endothelial

Figure 5. Stellate

A

B

- log10(P)

SEMAPHORIN PLEXIN
SIGNALING PATHWAY

Module 1
Module 2
Module 3
Module 4

Figure 6. Macrophage

Figure 7

Figure 8

	Machine learning	Gene set enrichment analysis	Co-expression, protein-protein interaction
Global TF regulation in	→ Modules (TFs therein)	→ Biological processes	→ Core genes

Hepatocyte
- Module 1 (STAT family) — Steroid metabolism — *Abcg1*, *Sqle*
- Module 2 (AP-1 family) — TNFα signaling via NF-κB — FOS/JUN, *Tnf*
- — Apoptosis — FOS/JUN, *Bcl2l1*
- Module 3 (TCF/LEF family) — Lobular zonation — *Glul*, *Cyp7a1*
- Module 4 — Undetected

Endothelial cell
- Module 1 (AP-1 family) — TNFα signaling via NF-κB — *Socs3*, *Tlr2*
- Module 2 (AHCTF1, ZNF740) — Angiogenesis — *Pxn*
- Module 3 — Undetected

Stellate cell
- Module 1 (SOX9) — Semaphorin-plexin signaling — *Nrp1*, *Nrp2*
- Module 2 — Undetected
- Module 3 — Undetected
- Module 4 — Undetected

Macrophage
- Module 1 (AP-1 family) — TNFα signaling via NF-κB — JUN, *Cd44*
- Module 2 (Maf family) — Complement system — *C1qc*, *Fcnb*
- Module 3 (IRF family) — Angiogenesis — *Pak1*, *Kdr*
- Module 4 — Undetected