Mapping Pharmacologically-induced Functional Reorganisation onto the Brain's Neurotransmitter Landscape

Andrea I. Luppi[1,2,3,4]*, Justine Y. Hansen[5], Ram Adapa[1], Robin L. Carhart-Harris[6], Leor Roseman[7], Christopher Timmermann[7], Daniel Golkowski[8], Andreas Ranft[9], Rüdiger Ilg[8,10], Denis Jordan[11], Vincent Bonhomme[12,13], Audrey Vanhaudenhuyse[12], Athena Demertzi[14], Oceane Jaquet[12], Mohamed Ali Bahri[14], Naji L.N. Alnagger[12], Paolo Cardone[12], Alexander R. D. Peattie[1,2], Anne E. Manktelow[1], Draulio B. de Araujo[15], Stefano L. Sensi[16,17], Adrian M. Owen[18], Lorina Naci[19], David K. Menon[1,20], Bratislav Misic[5], Emmanuel A. Stamatakis[1,2]

[1]Division of Anaesthesia, University of Cambridge, Cambridge, UK.

[2]Department of Clinical Neurosciences, University of Cambridge, Cambridge, UK.

[3]Leverhulme Centre for the Future of Intelligence, University of Cambridge, Cambridge, UK.

[4]The Alan Turing Institute, London, UK.

[5]McConnell Brain Imaging Center, Montreal Neurological Institute, McGill University, Montreal, QC, Canada.

[6]Psychedelics Division - Neuroscape, Department of Neurology, University of California San Francisco, San Francisco, CA, USA.

[7]Center for Psychedelic Research, Department of Brain Sciences, Imperial College London, London, UK.

[8]Department of Neurology, Klinikum rechts der Isar, Technical University Munich, München, Germany.

[9]Technical University of Munich, School of Medicine, Department of Anesthesiology and Intensive Care, Munich, Germany.

[10]Asklepios Clinic, Department of Neurology, Bad Tölz, Germany.

[11]Department of Anaesthesiology and Intensive Care Medicine, Klinikum rechts der Isar, Technical University Munich, München, Germany.

[12]Department of Anesthesia and Intensive Care Medicine, Liege University Hospital, Liege, Belgium.

[13]Anesthesia and Perioperative Neuroscience Laboratory, GIGA-Consciousness Thematic Unit, GIGA-Research, Liege University, Liege, Belgium.

[14]GIGA-Cyclotron Research Centre-In Vivo Imaging, University of Liege, Liege, Belgium.

[15]Brain Institute, Federal University of Rio Grande do Norte, Natal, RN, Brazil.

[16]Department of Neuroscience and Imaging and Clinical Science, Center for Advanced Studies and Technology, Institute for Advanced Biomedical Technologies, University "G. d'Annunzio" Chieti-Pescara, Chieti, Italy.

[17]Institute for Memory Impairments and Neurological Disorders, University of California-Irvine, Irvine, CA, USA.

18 Department of Psychology and Department of Physiology and Pharmacology, Western
Institute for Neuroscience (WIN), Western University, London, ON, Canada.

19 Trinity College Institute of Neuroscience, School of Psychology, Trinity College Dublin,
Dublin, Ireland.

20 Wolfon Brain Imaging Centre, University of Cambridge, Cambridge, UK.

*Corresponding author: email: al857@cam.ac.uk

Abstract

To understand how pharmacological interventions can exert their powerful effects on brain function, we need to understand how they engage the brain's rich neurotransmitter landscape. Here, we bridge microscale molecular chemoarchitecture and pharmacologically-induced macroscale functional reorganisation, by relating the regional distribution of 19 neurotransmitter receptors and transporters obtained from Positron Emission Tomography, and the regional changes in functional MRI connectivity induced by 10 different mind-altering drugs: propofol, sevoflurane, ketamine, LSD, psilocybin, DMT, ayahuasca, MDMA, modafinil, and methylphenidate. Our results reveal that psychoactive drugs exert their effects on brain function by engaging multiple neurotransmitter systems. The effects of both anaesthetics and psychedelics on brain function are organised along hierarchical gradients of brain structure and function. Finally, we show that regional co-susceptibility to pharmacological interventions recapitulates co-susceptibility to disorder-induced structural alterations. Collectively, these results highlight rich statistical patterns relating molecular chemoarchitecture and drug-induced reorganisation of the brain's functional architecture.

Keywords: neurotransmitters; receptors; functional MRI; pharmacology; anaesthesia; psychedelic; cognitive enhancer; Positron Emission Tomography.

Introduction

Understanding how the brain orchestrates complex signals across spatial and temporal scales to support cognition and consciousness is a fundamental challenge of contemporary neuroscience. By inducing profound but reversible alterations of brain function, psychoactive compounds provide neuroscientists with the means to manipulate the brain without requiring surgical intervention. In combination with non-invasive brain imaging techniques such as functional MRI, acute pharmacological interventions have therefore emerged as a prominent tool for causal investigation of the relationship between brain and cognitive function in healthy humans[1].

Mind-altering pharmacological agents also play a fundamental role in modern clinical practice. The invention of anaesthesia was a major milestone in medical history, enabling millions of life-saving surgeries to take place every year[2]. Other drugs that influence the mind without suppressing consciousness, such as the cognitive enhancers modafinil and methylphenidate, have found useful applications in alleviating the cognitive symptoms of syndromes such as ADHD, narcolepsy, and traumatic brain injury (TBI)[3–11]. More recently, classic and "atypical" psychedelics are increasingly being investigated for their potential to provide breakthrough avenues to treat psychiatric conditions, with recent successes in clinical trials heralding a possible end to the current scarcity of therapies for treatment-resistant depression and other neuropsychiatric disorders[12–17]. For these convergent reasons, the effects of anaesthetics, psychedelics, and cognitive enhancers on brain function are becoming the focus of intense investigation, revealing both similarities and differences between them[18–26].

Pharmacological agents exert their mind-altering effects by tuning the brain's neurotransmitter landscape. Neurotransmitters engage receptors on neurons' membrane to mediate the transfer and propagation of signals between cells, modulate the functional configurations of neuronal circuits, and ultimately shape network-wide communication[27–31]. Several psychoactive drugs appear to exert their effects on the mind and brain primarily through one or few specific neurotransmitters: the main action of the general anaesthetic propofol is enhancement of synaptic transmission mediated by GABA-A receptors, a mechanism that is also shared by

sevoflurane, which in addition attenuates glutamatergic synaptic signalling (mediated by both AMPA and NMDA receptors)[2,32–39]. Ketamine (a dissociative anaesthetic at high doses, and atypical psychedelic at low doses) is an NMDA receptor antagonist[40–45]; the classic psychedelics LSD, psilocybin, and DMT are agonists of the serotonin 2A receptor, with a strong dependence between subjective efficacy and 2A receptor affinity[46–49].

However, in the words of Sleigh and colleagues, "Linking observed molecular actions for any particular drug with its clinical effects is an abiding pharmacological problem"[50]: knowing the primary molecular target is not sufficient to understand a drug's effects on brain function, for several reasons. First, given the brain's intricate, nested feedback loops and recurrent pathways of connectivity, even a relatively selective drug can end up influencing unrelated systems beyond what may be apparent from in vitro studies. Second, most mind-altering compounds are also known to have affinity for other receptors. Indeed, evidence has been accumulating that multiple neurotransmitter influences may be involved in both the neural and subjective experiences induced by many consciousness-altering drugs. In the last year, human neuroimaging studies identified the involvement of the dopaminergic system in both propofol-induced anaesthesia[51] and the subjective effects of LSD[52]. More broadly, a recent large-scale study, combining receptor expression from transcriptomic data with linguistic processing of several thousand subjective reports of psychedelic use, identified complex multivariate patterns of association between neurotransmitters and their effects on the mind elicited by a wide variety of psychedelics, even for putatively selective agents[53]. At the same time, molecularly different compounds can exert intriguingly similar effects on both the mind and brain: for instance, LSD and (sub-anaesthetic) ketamine can produce subjectively similar effects and changes in terms of structure-function coupling and the complexity of brain activity - despite acting on different pathways[21]. This suggests both divergent and convergent effects of different pharmacological agents on the brain's rich neurotransmitter landscape.

Finally, the human brain exhibits rich patterns of anatomical, functional, cytoarchitectonic, and molecular variations[54–59]. Such patterns also extend to the regional distribution of different neurotransmitter receptors and transporters, which

vary widely not only in terms of their affinity, time-scales, and downstream effects on neuronal excitability, but also their distribution across regions, layers and neuron types[27–30]. Therefore, our knowledge of how a drug influences neurotransmission must take into account the neuroanatomical distribution of its target neurotransmitters - an essential step towards explaining how different neurotransmitters mediate the capacity of different drugs to shape the functional and computational properties of the brain's architecture[27,31].

Here, we sought to address this question in a data-driven way, mapping the neurotransmitter landscape of drug-induced alterations in the brain's functional connectivity. To do so, we leveraged two unique datasets: (i) a recently assembled collection of in vivo maps of regional receptor expression from 19 different receptors, obtained from PET scanning of over 1200 total subjects, providing the most detailed information about neuromodulators and their spatial distribution available to date[31]; and (ii) resting-state functional MRI (rs-fMRI) data acquired under the effects of the serotonergic psychedelics LSD[60], psilocybin[61], DMT (Timmermann *et al.,* under review), ayahuasca[62], and MDMA[63]; sub-anaesthetic doses of ketamine (acting as an "atypical psychedelic")[64] as well as anaesthetic doses (acting as a "dissociative anaesthetic"); the cognitive enhancers modafinil[65] and methylphenidate[10]; and the anaesthetics sevoflurane[66] and propofol[67,68] (which we compared against pre-anaesthesic baseline as well as post-anaesthetic recovery); representing a total of 382 sessions of pharmacological-MRI from 224 distinct subjects and 10 distinct pharmacological agents. Through pharmacologically modulated rs-fMRI, we can characterise a drug's effects on the brain's spontaneous activity, without the interference of any specific task[1].

Thus, our goal was to obtain a comprehensive mapping between the cortical distributions of neurotransmitters and a set of diverse psychoactive pharmacological agents (covering the range from anaesthetics to psychedelics), in terms of their effects of functional connectivity. There have been other studies looking at the relationships between brain changes induced by one or few psychoactive drugs, and one or few neurotransmitter systems[51,52,69–74] – and a previous effort considering how changes in cerebral blood flow induced by different psychiatric medications depend on the distribution of receptors[75]. However, to our knowledge, this is the largest fMRI

184 study to date not only in terms of the number, variety, and potency of psychoactive
185 pharmacological agents included, but also the breadth and coverage of
186 neurotransmitter systems considered.

Results

To establish a relationship between neurotransmitter systems and
pharmacologically-induced reorganisation of the brain's functional architecture, we
combine two sets of neuroimaging data, each collected from across multiple studies.
On one hand, we characterise drug-induced functional reorganisation as the
changes in functional connectivity (FC) obtained by contrasting resting-state
functional MRI (rs-fMRI) at baseline and under the acute effect of a psychoactive
drug. We considered the general anaesthetics propofol (two independent datasets)
and sevoflurane; the cognitive enhancers modafinil and methylphenidate; ketamine,
acting as both atypical psychedelic (at sub-anaesthetic doses) and as dissociative
anaesthetic[24,75,76]; and the serotonergic psychedelics lysergic acid diethylamide
(LSD), psilocybin, DMT, ayahuasca, and MDMA (Figure 1). For sevoflurane and both
propofol datasets, we considered two contrasts: drug versus pre-induction baseline,
and drug versus post-anaesthetic recovery (recovery data were not available for
ketamine). We followed the same preprocessing and denoising procedure for each
dataset, to ensure comparability (see Methods).

On the other hand, we consider the cortical distribution of 15 neurotransmitter
receptors and 4 transporters, obtained from in vivo Positron Emission Tomography[31].
Overall, 9 neurotransmitter and neuromodulatory systems ("neurotransmitters" for
short) are covered: dopamine (D1[77], D2[78–81], DAT[82]), norepinephrine (NET[83–86]),
serotonin (5-HT1A[87], 5-HT1B[87–90,90–92], 5-HT2A[93], 5-HT4[93], 5-HT6[94,95], 5-HTT[93]),
acetylcholine ($\alpha 4\beta 2$[96,97], M1[98], VAChT[99,100]), glutamate (mGluR5[101,102], NMDA[103,104]),
GABA (GABA-A[105]), histamine (H3[106]), cannabinoid (CB1[107–110]), and opioid
(MOR[111]). (Figure 1A,B). Both rs-fMRI and PET maps were parcellated into 100
functionally defined regions according to the Schaefer atlas[112].

Figure 1. Overview of receptors and pharmacological rs-fMRI data. (A) For each psychoactive drug, its pattern of pharmacologically-induced functional reorganisation is quantified as the average (across subjects) of the within-subject difference in regional FC density between task-free fMRI scans at baseline and under the drug's effects. The result is a map of 100 cortical regions by 15 drug-related contrasts. (B) Neurotransmitter systems are mapped with Positron Emission Tomography with radioligands for 15 receptors and 4 transporters, resulting in a map of 100 cortical regions by 19 neurotransmitters. Correlating each of these sets of maps against itself yields two region-by-region matrices of pharmacological co-susceptibility (C) and neurotransmitter co-expression (D), respectively, which are significantly correlated even after removing the exponential relationship with Euclidean distance between regions (E).

Brain regions with shared chemoarchitecture also respond similarly across pharmacological perturbations

Receptors and transporters shape the way that neurons respond to neurotransmission and neuromodulatory influences. In turn, psychoactive drugs exert their effects (primarily) by acting on neurotransmitters and neuromodulators. Therefore, we reasoned that everything else being equal, regions that express similar patterns of receptors and transporters should exhibit similar patterns of susceptibility to drug-induced functional reorganisation.

To address this question, we computed matrices of pharmacological co-susceptibility and neurotransmitter co-expression between pairs of regions, by correlating respectively the regional patterns of drug-induced FC changes (across all subjects), and the regional patterns of neurotransmitter expression across all 19 receptor and transporter PET maps. To account for spatial autocorrelation in molecular and FC attributes, we regressed out from both matrices the exponential trend with Euclidean distance[113–116].

Supporting our hypothesis, we found that pharmacological co-susceptibility is significantly correlated with neurotransmitter profile similarity: the extent to which two regions' FC patterns are similarly affected by perturbations induced by different psychoactive drugs, is predicted by the extent to which they co-express neurotransmitter receptors and transporters: *rho* = 0.34, *p* < 0.001 after regressing out the effects of Euclidean distance (Figure 1C-E). In other words, regions that exhibit shared chemoarchitecture also respond similarly across pharmacological perturbations.

Multivariate Receptor-Drug Associations

The previous analysis revealed of a relationship between large-scale patterns of neurotransmitter expression, and large-scale patterns of functional susceptibility to pharmacological perturbations – complementing previous work that identified relationships between individual drugs and individual receptors. However, neither of these two approaches captures the full richness of the two datasets employed here. To obtain a synthesis between these two approaches, we employed a multivariate association technique, Partial Least Squares correlation (PLS, also known as Projection to Latent Structures[117,118]), which enabled us to identify multivariate patterns of maximum covariance between drug-induced effects on functional connectivity, and the cortical distributions of neurotransmitter expression[119,120].

This analysis indicated the presence of two statistically significant latent variables (linear weighted combinations of the original variables) relating pharmacologically-induced functional reorganisation to neurotransmitter profiles, together accounting for nearly 85% of covariance. Significance was assessed against autocorrelation-preserving spin-based null models, embodying the null hypothesis that drug effects and neurotransmitters are spatially correlated with each other purely because of inherent spatial autocorrelation[121–124] (Figure 2). We further cross-validated this result using a distance-dependent method; out-of-sample $r = 0.46$ for PLS1 and 0.54 for PLS2, both $p < 0.001$ from t-test against spin-based null distributions) (Figure S1).

Figure 2. PLS analysis reveals spatially covarying patterns of pharmacologically-induced functional reorganisation and neurotransmitter expression. (A) PLS analysis relates two data domains by correlating the variables across brain regions and subjecting this to singular value decomposition. This results in multiple latent variables: linear weighted combinations of the original variables (neurotransmitter weights and drug weights) that maximally covary with each other. (B) Latent variables are ordered according to effect size (the proportion of covariance explained between neurotransmitter expression and drug-induced functional reorganisation they account for) and shown as red dots. (C) The first two latent variables (PLS1 and PLS2) were statistically significant, with respect to the spatial autocorrelation-preserving null model shown in grey (10,000 permutations). The first latent variable accounted for 57% of covariance, and the second latent variable accounted for 28%. Neurotransmitter (drug) scores are defined as the projection of the original neurotransmitter density (drug-induced FC changes) matrix onto the neurotransmitter (drug) weights, such that each brain region is associated with a neurotransmitter and drug score. By design, neurotransmitter and drug scores correlate highly.

For each latent variable, each brain region is associated with a neurotransmitter and drug score. In turn, neurotransmitter (drug) loadings are defined as the correlation between the PLS-derived score pattern and each neurotransmitter's density of expression (resp., drug-induced FC changes) across brain regions. Taking into account the first latent variable (PLS1), drug loadings showed a distinction of

pharmacological effects into two groups, with all anaesthetics (except ketamine) on one side, and both ketamine datasets dominating the opposite side, together with LSD, ayahuasca, and modafinil (Figure 3A). Neurotransmitter loadings divided the receptors from transporters: at the positive end (orange), the noradrenaline, serotonin and acetylcholine transporters (with the dopamine transporter following closely, but narrowly including zero in its 95% CI); all receptors except NMDA were instead at the negative end (blue), although some included zero in their CI (Figure 3B).

Figure 3. PLS scores and loadings from significant latent variables. (A-B) Scores and loadings for PLS1. (C-D) scores and loadings for PLS2. Brain plots: Drug scores (top row) and neurotransmitter scores (bottom row) for each brain region are obtained by projecting the original neurotransmitter and drug data back onto the PLS analysis-defined drug/neurotransmitter weights, indexing the extent to which a brain region expresses covarying drug/ neurotransmitter patterns. In turn, neurotransmitter (drug) loadings are defined as the Pearson's correlation between each neurotransmitter's density of expression (drug-induced FC changes) across brain regions and the PLS analysis-derived score pattern. Error bars indicate 95% confidence interval, and colour indicates direction of the effect: positive (orange), negative (blue), or null (grey). Same-coloured loadings and scores co-vary positively, whereas opposite-coloured drugs and scores co-vary negatively. The label "ketamine 1" refers to the sub-anaesthetic dose, and "ketamine 2" is the anaesthetic dose.

Pertaining to the second latent variable (PLS2), neurotransmitter loadings mainly identified a monoamine-rich end (with dopamine and serotonin), although 5-HT1b occupied the opposite end. However, the drug loadings were less clearly discernible, with modafinil alone at one end, and a mixture of propofol, psychedelics, and both ketamine datasets at the other end. Both neurotransmitter and drug scores markedly separated dorsal and ventral aspects of the brain for this second latent variable (Figure 3).

Pharmacologically-induced alterations align with functional, anatomical and molecular hierarchies

Neurotransmitter and drug scores (whose spatial similarity PLS is designed to maximise) provide information about the regional distribution of neurotransmitter-drug associations. Neurotransmitters and drugs whose activity correlates positively with the score pattern covary with one another in the positively scored regions, and vice versa for negatively scored regions.

PLS1 scores correspond to the main axis of covariance between neurotransmitter expression and pharmacologically-induced functional reorganisation. For both drug and receptor scores, we observed that their regional distribution reflected the brain's organisation into intrinsic resting-state networks (RSNs)[125], setting apart visual and somatomotor cortices from association cortices (Figures 3,4). It is possible that the correspondence of PLS1 scores with RSNs may be in part driven by the fact that these networks are predicated in terms of functional neuroimaging, which we also used to characterise drug-induced functional reorganisation in our data. Therefore, we next sought to determine whether our data-driven topographic patterns reflect other cortical gradients of variation in terms of functional, anatomical, and molecular attributes. To this end, we considered intracortical myelination obtained from T1w/T2w MRI ratio[58]; evolutionary cortical expansion obtained by comparing human and macaque[126]; the principal component of variation in gene expression from the Allen Human Brain Atlas transcriptomic database ("AHBA PC1")[59,127]; the principal component of variation in task activation from the NeuroSynth database ("NeuroSynth PC1")[59,128]; and the principal gradient of functional connectivity[57].

Since pharmacological interventions exert their effects on the brain via the
bloodstream, we also included a map of cerebral blood flow[54]. Finally, we included a
recently derived gradient of regional prevalence of different kinds of information, from
redundancy to synergy[129].

We observed significant correlations (assessed against spin-based null models)
between each cortical hierarchy and both neurotransmitter and drug scores for PLS1
(except for PLS1 drug scores versus NeuroSynth PC1; $p_{spin} = 0.054$) (Figure 4). The
scores for PLS2 instead identified a ventral-dorsal pattern of regional variation
(Figure 3 and Figure S2), which did not significantly correlate with any of the
canonical gradients of hierarchical organisation (all $p > 0.05$ against spin-based null
models, except for PLS2 drug scores versus NeuroSynth PC1; $p_{spin} = 0.013$).

Figure 4. Correspondence between the principal axis of drug-neurotransmitter scores and functional, anatomical and molecular hierarchies. (A) Cortical distribution of drug scores for PLS1, and their association with intrinsic resting-state networks. (B) Radial plot represents the absolute value of the correlation between PLS1 drug and neurotransmitter scores, and each of seven cortical hierarchies obtained from different neuroimaging modalities (note that the myelin and AHBA PC1 maps are reversed with respect to the remaining hierarchies).

Neurotransmitter landscape of pharmacologically-induced functional reorganisation

Taking into account the first two PLS latent variables shows how each drug-specific pattern of pharmacologically-induced functional reorganisation can be interpreted in

terms of contributions from different receptors (note that sign is arbitrary) (Figure 5).

As already shown in Figure 3, the first latent variable revealed a stark division

between transporters and receptors, which discriminates between traditional

anaesthetics and other psychoactive substances. In terms of pharmacological

alterations, the non-monoaminergic end of the second latent variable loaded onto

drugs with relatively stronger effects on subjective experiences (the higher doses of

anaesthetic, including ketamine, and LSD and MDMA). However, methylphenidate

and sub-anaesthetic ketamine also loaded onto this end of the second latent

variable. Altogether, we find that the first latent variable captures a strong

relationship between drug interventions and receptor systems that is both biologically

relevant and aligns with the functional organisation of the brain.

406 Co-susceptibility to pharmacological and pathological
407 alterations

Finally, we wondered if the functional co-susceptibility of different regions to transient

pharmacological perturbations may provide a functional proxy for their co-

susceptibility to structural perturbations resulting from different neurological,

neurodevelopmental, and psychiatric disorders. To this end, we combined 11 spatial

maps of cortical thickness abnormalities made available by the Enhancing Neuro

Imaging Genetics Through Meta Analysis (ENIGMA) consortium[130]: 22q11.2 deletion

syndrome[131], attention-deficit/hyperactivity disorder[132], autism spectrum disorder[133],

idiopathic generalized epilepsy[134], right temporal lobe epilepsy[134], left temporal lobe

epilepsy[134], depression[135], obsessive-compulsive disorder[136], schizophrenia[137],

bipolar disorder[138], and Parkinson's disease[139]. For simplicity, we refer to diseases,

disorders, and conditions as "disorders" throughout the text. The cortical abnormality

maps summarise contrasts between over 21,000 adult patients and 26,000 controls,

collected following identical processing protocols to ensure maximal comparability[130].

Following the same procedure used to obtain the region x region matrices of

pharmacological co-susceptibility and neurotransmitter co-expression in Figure 1, we

obtained a region x region matrix of co-susceptibility to disorder-induced cortical

abnormality by correlating the regional patterns of cortical abnormality across all 11

disorders[113] (Figure 6A,B). Correlating this matrix of regional co-susceptibility to

disease-associated perturbations against the previously derived matrix of regional

co-susceptibility to pharmacological perturbations, we found a statistically significant

relationship (Spearman's *rho* = 0.31, *p* < 0.001 after regressing out the effect of

Euclidean distance) (Figure 6C). This result goes beyond a recent demonstration

that molecular similarity and disorder similarity are correlated[113], by showing that a

correlation also exists between different kinds of perturbations: anatomical and
pharmacological.

This observation suggests that there may be common patterns of regional co-
susceptibility to perturbations, whether structural or functional. To explore this
possibility explicitly, we resorted to a non-linear dimensionality reduction algorithm,
diffusion map embedding[57,140], to obtain joint gradients of variation from
pharmacological and disease-associated co-susceptibility using a recently developed
method for network fusion[141]. We found that the main axis of variation in regional
joint susceptibility to pharmacological and neuropsychiatric alterations corresponds
to the well-known principal gradient of functional connectivity[57], setting apart
unimodal from transmodal cortices, reminiscent of the PLS1 scores (Figure 6D). The
second gradient instead sets apart anterior dorsal and more ventral parts of the
brain, reminiscent of the PLS2 scores (Figure 6E). Together, these two gradients
account for nearly half of the variation in regional co-susceptibility (Figure 6F).

When applying diffusion map embedding to the matrix of pharmacological co-
susceptibility only, we found that the first two gradients of variation in regional
pharmacological susceptibility coincide with the two principal gradients of functional
connectivity of Margulies *et al*[57] (Figure S4): the first gradient sets apart unimodal
from transmodal cortices, coinciding with the first gradient of joint susceptibility,
whereas the second gradient is anchored in visual cortex at one end, and
somatomotor cortex at the other end. This observation suggests that co-susceptibility
to pharmacological perturbations recapitulates intrinsic functional architecture, as
well as the co-susceptibility to disorder-induced structural perturbations.

457
Figure 6. Co-susceptibility to pharmacological and pathological alterations. Brain regions that
are similarly affected by pharmacology, in terms of functional reorganisation (A) are also similarly
affected across disorders (B), in terms of cortical thickness abnormalities. This relationship persists
after regressing out the exponential trend with Euclidean distance (C). (D-E) First two principal
gradients of regional joint susceptibility to pharmacological and neuropsychiatric and neurological
alterations, obtained from diffusion map embedding. (F) Scree plot of the scaled eigenvalues from
diffusion map embedding versus number of components.

466 Validation and additional analyses

Results for a different cortical parcellation (Lausanne-114[142]) are provided as
Supplementary Information (Figures S5, S6); likewise, we show a qualitatively similar
mapping between drugs and neurotransmitters on the first two latent variables, even
when the methylphenidate dataset (obtained from patients rather than healthy
controls) is excluded (Figure S7).

We also performed a separate PLS analysis on the subcortex, which revealed a
monoamine- (dopamine- and serotonin-) dominated principal latent variable,
positively associated with the drugs having strongest effects, and separating
caudate, putamen, and anterior thalamus from posterior thalamus, hippocampus and
amygdala (Figure S8).

Discussion

Here, we characterised how mind-altering pharmacological agents engage the brain's rich neurotransmitter landscape to exert their effects on brain function. We mapped the functional chemoarchitecture of the human brain, by developing a computational framework to relate the regional reorganisation of fMRI functional connectivity induced by 10 different mind-altering drugs, and the cortical distribution of 19 neurotransmitter receptors and transporters obtained from PET[31]. This approach allowed us to discover large-scale spatial gradients relating pharmacologically-induced changes in functional connectivity to the underlying neurotransmitter systems. By relating microscale molecular chemoarchitecture and macroscale functional reorganisation induced by drugs with potent acute effects on the mind, our results provide a first step to bridge molecular mechanisms and their effects on subjective experience, cognition, and behaviour, via their effects on the brain's functional architecture.

Using our computational framework, we found that psychoactive drugs are best understood in terms of contributions from multiple neurotransmitter systems. We also found that anaesthetics and psychedelics/cognitive enhancers are largely opposite in terms of their association with neurotransmitters in the cortex – although not without exceptions. Remarkably, the effects of mind-altering drugs are topographically organised along multiple hierarchical gradients of brain function, anatomy, and neurobiology. Finally, we found that co-susceptibility to pharmacological perturbations recapitulates co-susceptibility to disorder-induced structural perturbations.

The diverse mapping between drug-induced functional reorganisation and neurotransmitters that we recovered (Figure 5) clearly shows the power of our multivariate approach to detect both expected and novel relationships between drugs and neurotransmitters. Many of the drugs considered here are known to have varied molecular targets, beyond the primary ones through which they exert their effects. The present results add another dimension to recent work employing a similar multivariate approach to relate gene expression of receptors with subjective reports

of psychedelic experiences, which also found widespread involvement of multiple receptors[53]. In addition, the drugs we considered here have profound effects on the mind after a single acute dose[143], from cognitive enhancement to hallucinations to the suppression of consciousness altogether. Such far-reaching effects are accompanied by sometimes drastic repercussions on brain function and dynamics: it stands to reason that such widespread reorganisation would not leave many neurotransmitter pathways unaffected - even those that are not directly involved in generating the altered state in question.

The broadly opposite characterisation of traditional anaesthetics and most psychedelics is aligned with their respective effects on the complexity of brain activity and connectivity, which is reduced by GABA-ergic anaesthesia but increased by LSD, ayahuasca and psychedelic doses of ketamine, as well as other psychedelics[18,21,23,62,144–154]. Similarly, psychedelics (including sub-anaesthetic ketamine) and anaesthetics were recently shown to exert opposite effects on structure-function coupling: whereas anaesthesia increases the dependence of brain activity on the underlying structural network, LSD, psilocybin, and sub-anaesthetic ketamine induce fMRI BOLD signals that are increasingly liberal with respect to the underlying structural network organisation[21]. Intriguingly, we found that anaesthetic doses of ketamine align more closely with psychedelic (sub-anaesthetic) doses of ketamine than with anaesthetics such as propofol and sevoflurane. Although the ketamine-anaesthetised volunteers were behaviourally unresponsive, as for the traditional anaesthetics, they subsequently reported a wide range of vivid hallucinatory experiences[75]. Thus, as per ketamine's characterisation as a dissociative anaesthetic, in subjective terms their consciousness was not suppressed but rather profoundly altered in a manner more similar to psychedelics than anaesthetics. Therefore, the neurotransmitter signatures of the two levels of ketamine align with the molecular effects and subjective effects, rather than with the behavioural effects.

The main division we observed in terms of neurotransmitters is between receptors and transporters, which displayed opposite associations with drug-induced effects. Specifically pertaining to PLS1, we found that transporters covary with cognitive enhancers and most psychedelics in primary sensory and motor regions, whereas

receptors covary with GABA-ergic anaesthetics in transmodal association cortices. Hierarchical organisation of pharmacologically-induced functional reorganisation stands to reason based on prior evidence: both psychedelics and anaesthetics have been shown to have potent effects on the activity and connectivity of higher-order association cortices, and the default mode network in particular[18,60,61,66,75,155–157]. In addition, serotonergic psychedelics also exert powerful influences on the visual cortex at the other end of the cortical hierarchy[60], and as a result they have been shown to induce a "flattening" of the principal gradient of functional connectivity[158].

Having established that the effects of mind-altering drugs are hierarchically organised, the question then becomes: why should mind-altering drugs exert their effects in such a hierarchically organised fashion? Multiple aspects of neuroanatomy may contribute to this effect. First, the principal component of variation of receptor expression is itself organised along the brain's sensory-to-association hierarchical axis[27] - and so is, for instance, the distribution of the serotonin 2A receptor, the main direct target of serotonergic psychedelics[31]. Second, transmodal cortices are characterised by increased excitability[159] and a predominance of feedback efferent connections[27]: combined with their high diversity of receptor expression across layers[27], these regions may be especially susceptible to receive and amplify multiple pharmacological influences.

Third, we observed that for the first latent variable, neurotransmitter and drug scores also correlate with the map of regional cerebral blood flow; since ultimately the bloodstream is how drugs reach their regional molecular targets, greater cerebral blood flow in transmodal cortices may facilitate especially high availability of the drug in these regions (although it should be noted that some drugs can also have effects on heart rate and neuro-vascular coupling). Finally, regions of transmodal cortex have high neuron and synapse density[129,160] and tend to have numerous, far-reaching, and diversely distributed anatomical connections[161], as well as the highest prevalence of synergistic (complementary) interactions with the rest of the brain[129], so that any effects that are exerted in these regions may quickly reverberate throughout the whole cortex.

To summarise, we conjecture that the hierarchical organisation of pharmacologically-induced changes in FC may be explained as follows: transmodal association cortices are especially diverse in their receptor profiles, and rich in some key receptors; in addition to being more susceptible to pharmacological intervention due to higher expression of receptors, blood flow is poised to bring greater amounts of drug to these very cortices; and once these cortices' activity is perturbed, the perturbation can reverberate widely, thanks to their widespread and diverse connectivity. Of course, the drugs we included were chosen precisely because of their powerful effects on cognition and subjective experience, so it stands to reason that their effects should align with the division between primary and higher-order cortices (which also aligns with the principal component of variation obtained from NeuroSynth term-based meta-analysis). In other words, drugs whose effects on functional connectivity are less selective for higher versus lower ends of the cortical hierarchy may simply be less likely to exert mind-altering effects of the kind that we chose to focus on in this work.

More broadly, we found that pairs of regions that are more similar in terms of their susceptibility to pharmacologically-induced FC changes, are also more similar in their susceptibility to cortical alterations associated with a variety of neuropsychiatric disorders. This observation suggests a broader pattern of both pharmacological (acute) and neuroanatomical (chronic) susceptibility across regions. We speculate that this joint susceptibility may be related to regional relevance for cognitive function: indeed, we found that this joint vulnerability can be understood in terms of two multimodal principal gradients of variation over the cortex: one of them resembling the principal gradient of functional connectivity (and principal latent variable of neurotransmitter-drug association), and the other anchored in dorsal prefrontal cortex at one end, and temporal cortex at the other. The association between disorder co-susceptibility and co-susceptibility to pharmacologically-induced functional reorganisation sheds new light on recent evidence that the principal gradient of neurotransmitter expression is particularly relevant for predicting a wide spectrum of disease-specific cortical morphology[113], by showing that this observation extends to the effects of engaging different receptors. This interpretation is further supported by our own evidence that pharmacological perturbations are shaped by neurotransmitter co-expression.

The results reported here open new possibilities for data-driven, multivariate mapping between the brain's high-dimensional neurotransmitter landscape and the effects of potent pharmacological interventions on the brain's functional architecture. Crucially, neuropsychiatric disorders and candidate pharmacological treatments for them ultimately need to exert their effects on cognition and behaviour by influencing brain function. In this light, it is intriguing that susceptibility to disorder-related cortical abnormalities correlates with susceptibility to pharmacological intervention. This observation suggests that regions that are structurally most vulnerable to disease (which presumably in turn shapes their functional architecture) may also be the ones that are most susceptible to re-balancing of their functional organisation by an appropriate choice of pharmacological intervention. This work represents the necessary first step towards identifying novel and perhaps unexpected associations between drugs and neurotransmitters, as well as elucidating the known ones in a data-driven manner.

Limitations and future directions

Although the main strength of our study is our extensive coverage of both neurotransmitters and pharmacological data, it is important to acknowledge that neither is complete: in particular, our sample did by no means exhaustively include all mind-altering drugs that have been studied: prominent additions for future work may include the psychedelic kappa opioid receptor agonist salvinorin-A[162], the sedative dexmedetomidine, an alpha-2 receptor agonist[163–165] – but also alcohol or caffeine, arguably the two most widely used psychoactive substances. We also acknowledge that the pharmacological datasets included here come from limited samples that have been studied before, and future replication in different datasets with the same drugs (as we have done here for propofol) would also be desirable.

The datasets included here come from different sources and locations, and were acquired under a variety of conditions. We endeavoured to mitigate scanner and acquisition differences by re-preprocessing all data with the same pipeline, and following uniform denoising procedures, rather than following the various pipelines originally employed by each group. Further mitigation of the acquisition differences

between datasets should come from our within-subject design in healthy individuals
(except for the methylphenidate dataset, which comes from TBI patients[10]; although
we showed that our results remain qualitatively the same if this dataset is excluded;
Fig. S6). Nevertheless, we cannot exclude some residual influence of such
differences on our results (e.g., eyes open versus closed; the ayahuasca data were
acquired at a lower field strength of 1.5T; the TRs varied from 1.671s for modafinil, to
3s for psilocybin). Similar considerations about the differences between datasets
apply for the PET data, as discussed in detail in the original publication collecting the
PET maps[31]. Likewise, the coverage of neurotransmitter receptors and transporters,
though the most extensive available to date and obtained *in vivo* rather than post-
mortem, is far from exhaustive. The same limitation also applies to the ENIGMA
disorder data[130]: many more disorders, diseases, and conditions exist than the ones
considered here. And although the ENIGMA consortium provides datasets from large
samples with standardised pipelines, ensuring robust results, the patient populations
may exhibit co-morbidities and/or be undergoing treatment. In addition, the available
maps do not directly reflect changes in tissue volume, but rather the effect size of
patient-control statistical comparisons, in terms of only one low resolution cortical-
only parcellation.

In addition to the inevitable limitations of analysing large-scale datasets from multiple
sites, there are also limitations of our analytic framework. Although we report a
macroscale spatial association between neurotransmitter expression and
pharmacologically-induced functional reorganisation that is statistically unexpected
based on autocorrelation alone, caution is warranted when drawing inferences from
statistical results to the underlying biology. We used linear models that assume
independence between observations – an assumption that mostly does not hold in
the brain, given the possibility of nonlinear effects in how drugs exert their effects on
the brain's intricately connected neurotransmitter systems. To mitigate this limitation,
throughout this work we triangulated towards a robust statistical mapping between
neurotransmitters and drugs by combining cross-validation and conservative null
models that account for the spatial dependencies between regions[59].

Another limitation is that, due to data availability and well documented differences in
PET radioligand uptake between cortical and subcortical structures[31,166,167], our work

was mainly restricted to the cortex, and we considered the subcortex separately. The

thalamus, brainstem, and other subcortical structures are prominently involved in

mediating cortico-cortical interactions and the effects of both psychedelics,

anaesthetics, and cognitive enhancers[16,18,51,66,163,168–173]. We expect that future work

suitable data for whole-brain coverage (ideally including cerebellum and brainstem)

may provide richer insights than the sum of their individual contributions.

More broadly, the other main limitation of this work is its correlational nature:

receptors and drugs were mapped in separate cohorts of individuals, and identifying

spatially correlated patterns does not guarantee the causal involvement of the

neurotransmitters in question. Experimental interventions will be required to

conclusively demonstrate causal involvement, and elucidate the underlying

neurobiological pathways. However, we emphasise that our results generate

empirically testable hypotheses about which neurotransmitters may be involved with

the macroscale effects of different drugs on brain function. Such hypotheses may be

tested experimentally, but also *in silico*: whole-brain computational modelling is

becoming increasingly prominent as a tool to investigate the causal mechanisms that

drive brain activity and organisation in healthy and pathological conditions [174–177].

Crucially, the more biologically-inspired models (e.g. dynamic mean-field) can also

be enriched with further information, such as regional myelination [159], or the regional

distribution of specific receptors and ion channels obtained from PET or

transcriptomics [71–73,178,179], to reflect neurotransmitter influences. This approach may

complement experimental manipulations, making it possible to systematically

evaluate the causal effects of combinations of different neuromodulators on the

brain's functional connectivity.

Conclusion
704

Here, we mapped the functional chemoarchitecture of the human brain, by relating

the regional changes in fMRI functional connectivity induced by 10 different mind-

altering drugs, and the regional distribution of 19 neurotransmitter receptors and

transporters obtained from PET. This work provides a computational framework to

characterise how mind-altering pharmacological agents engage the brain's rich

neurotransmitter landscape to exert their effects on brain function. Our analytic

workflow could find application across the breadth of human cognitive and clinical neuroscience, with the potential to shed light on alterations of neurotransmission underlying neuropsychiatric conditions, which are known to involve a combination of anatomical and neurochemical imbalances. More broadly, our framework could also find fruitful application for data-driven prediction of the effects of candidate drugs on the brain: the mapping between neurotransmitters and pharmacological effects on brain function offers an indispensable biological lens that can reveal neurotransmitter targets for therapeutic intervention. In summary, we demonstrate that diverse patterns of neurotransmitter expression are variously engaged by an array of potent pharmacological interventions, ultimately manifesting as a large-scale hierarchical axis. Collectively, these results highlight a statistical link between molecular dynamics and drug-induced reorganisation of functional architecture.

Materials and Methods

Description of datasets

Propofol

Propofol (2,6-diisopropylphenol) is perhaps the most common agent used for intravenous induction and maintenance of general anaesthesia[32]. One of the chief reasons for its widespread use, both in the operating room and for scientific studies, is propofol's rapid action, which allow for precise titration and therefore greater control over the induction and emergence process. Additionally, propofol has minimal effects on both regional cerebral blood flow[180], and the coupling between blood flow and metabolism[181], thereby reducing the number of potential confounding effects. Propofol is a potent agonist of inhibitory GABA-A receptors, directly activating them as well as increasing their affinity for agonists[33,34], leading to suppressed neuronal activity. Propofol also blocks Na+ channels, inhibiting glutamate release[182] and more broadly it inhibits neurotransmitter release at

presynaptic terminals[183]. There is also some evidence that it may affect the
dopaminergic system[51,184,185]. Here, we included two independent propofol datasets.

Western University dataset: Recruitment

The Western University ("Western") propofol data were collected between May and
November 2014 at the Robarts Research Institute, Western University, London,
Ontario (Canada), and have been published before[18,151,186,187]. The study received
ethical approval from the Health Sciences Research Ethics Board and Psychology
Research Ethics Board of Western University (Ontario, Canada). Healthy volunteers
(n=19) were recruited (18–40 years; 13 males). Volunteers were right-handed, native
English speakers, and had no history of neurological disorders. In accordance with
relevant ethical guidelines, each volunteer provided written informed consent, and
received monetary compensation for their time. Due to equipment malfunction or
physiological impediments to anaesthesia in the scanner, data from n=3 participants
(1 male) were excluded from analyses, leaving a total n=16 for analysis[18].

Western University dataset: Study protocol

Resting-state fMRI data were acquired at different propofol levels: no sedation
(Awake), Deep anaesthesia (corresponding to Ramsay score of 5) and also during
post-anaesthetic recovery. As previously reported[18], for each condition fMRI
acquisition began after two anaesthesiologists and one anaesthesia nurse
independently assessed Ramsay level in the scanning room. The anaesthesiologists
and the anaesthesia nurse could not be blinded to experimental condition, since part
of their role involved determining the participants' level of anaesthesia. Note that the
Ramsay score is designed for critical care patients, and therefore participants did not
receive a score during the Awake condition before propofol administration: rather,
they were required to be fully awake, alert and communicating appropriately. To
provide a further, independent evaluation of participants' level of responsiveness,
they were asked to perform two tasks: a test of verbal memory recall, and a
computer-based auditory target-detection task. Wakefulness was also monitored
using an infrared camera placed inside the scanner.

Propofol was administered intravenously using an AS50 auto syringe infusion pump
(Baxter Healthcare, Singapore); an effect-site/plasma steering algorithm combined
with the computer-controlled infusion pump was used to achieve step-wise sedation
increments, followed by manual adjustments as required to reach the desired target
concentrations of propofol according to the TIVA Trainer (European Society for
Intravenous Aneaesthesia, eurosiva.eu) pharmacokinetic simulation program. This
software also specified the blood concentrations of propofol, following the Marsh 3-
compartment model, which were used as targets for the pharmacokinetic model
providing target-controlled infusion. After an initial propofol target effect-site
concentration of 0.6 μg mL^{-1}, concentration was gradually increased by increments
of 0.3 μg mL^{1}, and Ramsay score was assessed after each increment: a further
increment occurred if the Ramsay score was lower than 5. The mean estimated
effect-site and plasma propofol concentrations were kept stable by the
pharmacokinetic model delivered via the TIVA Trainer infusion pump. Ramsay level
5 was achieved when participants stopped responding to verbal commands, were
unable to engage in conversation, and were rousable only to physical stimulation.
Once both anaesthesiologists and the anaesthesia nurse all agreed that Ramsay
sedation level 5 had been reached, and participants stopped responding to both
tasks, data acquisition was initiated. The mean estimated effect-site propofol
concentration was 2.48 (1.82- 3.14) μg mL^{-1}, and the mean estimated plasma
propofol concentration was 2.68 (1.92- 3.44) μg mL^{-1}. Mean total mass of propofol
administered was 486.58 (373.30- 599.86) mg. These values of variability are typical
for the pharmacokinetics and pharmacodynamics of propofol. Oxygen was titrated to
maintain SpO2 above 96%.

At Ramsay 5 level, participants remained capable of spontaneous cardiovascular
function and ventilation. However, the sedation procedure did not take place in a
hospital setting; therefore, intubation during scanning could not be used to ensure
airway security during scanning. Consequently, although two anaesthesiologists
closely monitored each participant, scanner time was minimised to ensure return to
normal breathing following deep sedation. No state changes or movement were
noted during the deep sedation scanning for any of the participants included in the
study[18]. Propofol was discontinued following the deep anaesthesia scan, and
participants reached level 2 of the Ramsey scale approximately 11 minutes

afterwards, as indicated by clear and rapid responses to verbal commands. This

corresponds to the "recovery" period.

As previously reported[18], once in the scanner participants were instructed to relax

with closed eyes, without falling asleep. Resting-state functional MRI in the absence

of any tasks was acquired for 8 minutes for each participant. A further scan was also

acquired during auditory presentation of a plot-driven story through headphones (5-

minute long). Participants were instructed to listen while keeping their eyes closed.

The present analysis focuses on the resting-state data only; the story scan data have

been published separately[188] and will not be discussed further here.

Western University dataset: MRI Data Acquisition

As previously reported[18], MRI scanning was performed using a 3-Tesla Siemens Tim

Trio scanner (32-channel coil), and 256 functional volumes (echo-planar images,

EPI) were collected from each participant, with the following parameters: slices = 33,

with 25% inter-slice gap; resolution = 3mm isotropic; TR = 2000ms; TE = 30ms; flip

angle = 75 degrees; matrix size = 64x64. The order of acquisition was interleaved,

bottom-up. Anatomical scanning was also performed, acquiring a high-resolution T1-

weighted volume (32-channel coil, 1mm isotropic voxel size) with a 3D MPRAGE

sequence, using the following parameters: TA = 5min, TE = 4.25ms, 240x256 matrix

size, 9 degrees flip angle[18].

Cambridge University dataset: Recruitment

The Cambridge University ("Cambridge") propofol dataset is described in detail in a

previous publication[67]. Sixteen healthy volunteer subjects were initially recruited for

scanning. In addition to the original 16 volunteers, data were acquired for nine

additional participants using the same procedures, bringing the total number of

participants in this dataset to 25 (11 males, 14 females; mean age 34.7 years, SD =

9.0 years). Ethical approval for these studies was obtained from the Cambridgeshire

2 Regional Ethics Committee, and all subjects gave informed consent to participate

in the study. Volunteers were informed of the risks of propofol administration, such

as loss of consciousness, respiratory and cardiovascular depression. They were also

informed about more minor effects of propofol such as pain on injection, sedation

and amnesia. In addition, standard information about intravenous cannulation, blood
sampling and MRI scanning was provided.

Cambridge University dataset: Study protocol

Three target plasma levels of propofol were used - no drug (Awake), 0.6 mg/ml (Mild
sedation) and 1.2 mg/ml (Moderate sedation). Scanning (rs-fMRI) was acquired at
each stage, and also at Recovery; anatomical images were also acquired. The level
of sedation was assessed verbally immediately before and after each of the
scanning runs. Propofol was administered intravenously as a "target controlled
infusion" (plasma concentration mode), using an Alaris PK infusion pump
(Carefusion, Basingstoke, UK). A period of 10 min was allowed for equilibration of
plasma and effect-site propofol concentrations. Blood samples were drawn towards
the end of each titration period and before the plasma target was altered, to assess
plasma propofol levels. In total, 6 blood samples were drawn during the study. The
mean (SD) measured plasma propofol concentration was 304.8 (141.1) ng/ml during
mild sedation, 723.3 (320.5) ng/ml during moderate sedation and 275.8 (75.42)
ng/ml during recovery. Mean (SD) total mass of propofol administered was 210.15
(33.17) mg, equivalent to 3.0 (0.47) mg/kg. Two senior anaesthetists were present
during scanning sessions and observed the subjects throughout the study from the
MRI control room and on a video link that showed the subject in the scanner.
Electrocardiography and pulse oximetry were performed continuously, and
measurements of heart rate, non-invasive blood pressure, and oxygen saturation
were recorded at regular intervals.

Cambridge University dataset: MRI Data Acquisition

The acquisition procedures are described in detail in the original study[37]. Briefly, MRI
data were acquired on a Siemens Trio 3T scanner (WBIC, Cambridge). For each
level of sedation, 150 rs-fMRI volumes (5 min scanning) were acquired. Each
functional BOLD volume consisted of 32 interleaved, descending, oblique axial
slices, 3 mm thick with interslice gap of 0.75 mm and in-plane resolution of 3 mm,

field of view = 192x192 mm, repetition time = 2000 ms, acquisition time = 2 s, time

echo = 30 ms, and flip angle 78. T1-weighted structural images at 1 mm isotropic

resolution were also acquired in the sagittal plane, using an MPRAGE sequence with

TR = 2250 ms, TI = 900 ms, TE = 2.99 ms and flip angle = 9 degrees, for localization

purposes. During scanning, volunteers were instructed to close their eyes and think

about nothing in particular throughout the acquisition of the resting state BOLD data.

Of the 25 healthy subjects, 15 were ultimately retained (7 males, 8 females): 10 were

excluded, either because of missing scans (n=2), or due of excessive motion in the

scanner (n=8, 5mm maximum motion threshold). For the analyses presented in this

paper, we only considered the Awake, Moderate (i.e., loss of behavioural

responsiveness) and Recovery resting-state scans.

Sevoflurane

Sevoflurane is an inhalational anaesthetic: specifically, a halogenated ether.

Although its exact molecular mechanisms of action are yet to be fully elucidated, in

vivo and in vitro evidence indicates that it acts primarily via GABA-A receptors[2,35–38],

but also interacts with NMDA, AMPA[39,189,190] and nicotinic ACh receptors[191,192].

Additionally, electrophysiologic investigation suggests possible affinity for as well as

Na+, K+ and hyperpolarization-activated cyclic nucleotide-gated (HCN) channels[193].

Sevoflurane dataset: Recruitment

The data included here have been published before [66,194–196], and we refer the reader

to the original publication for details[66]. The ethics committee of the medical school of

the Technische Universität München (München, Germany) approved the current

study, which was conducted in accordance with the Declaration of Helsinki. Written

informed consent was obtained from volunteers at least 48 h before the study

session. Twenty healthy adult men (20 to 36 years of age; mean, 26 years) were

recruited through campus notices and personal contact, and compensated for their

participation in the study.

Before inclusion in the study, detailed information was provided about the protocol
and risks, and medical history was reviewed to assess any previous neurologic or
psychiatric disorder. A focused physical examination was performed, and a resting
electrocardiogram was recorded. Further exclusion criteria were the following:
physical status other than American Society of Anesthesiologists physical status I,
chronic intake of medication or drugs, hardness of hearing or deafness, absence of
fluency in German, known or suspected disposition to malignant hyperthermia, acute
hepatic porphyria, history of halothane hepatitis, obesity with a body mass index
more than 30 kg/m2, gastrointestinal disorders with a disposition for
gastroesophageal regurgitation, known or suspected difficult airway, and presence of
metal implants. Data acquisition took place between June and December 2013.

Sevoflurane dataset: Study protocol

Sevoflurane concentrations were chosen so that subjects tolerated artificial
ventilation (reached at 2.0 vol%) and that burst-suppression (BS) was reached in all
participants (around 4.4 vol%). To make group comparisons feasible, an
intermediate concentration of 3.0 vol% was also used. In the MRI scanner,
volunteers were in a resting state with eyes closed for 700s. Since EEG data were
simultaneously acquired during MRI scanning [66] (though they are not analysed in the
present study), visual online inspection of the EEG was used to verify that
participants did not fall asleep during the pre-anaesthesia baseline scan.
Sevoflurane mixed with oxygen was administered via a tight-fitting facemask using
an fMRI-compatible anaesthesia machine (Fabius Tiro, Dräger, Germany). Standard
American Society of Anesthesiologists monitoring was performed: concentrations of
sevoflurane, oxygen and carbon dioxide, were monitored using a cardiorespiratory
monitor (DatexaS/3, General electric, USA). After administering an end-tidal
sevoflurane concentration (etSev) of 0.4 vol% for 5 min, sevoflurane concentration
was increased in a stepwise fashion by 0.2 vol% every 3 min until the participant
became unconscious, as judged by the loss of responsiveness (LOR) to the
repeatedly spoken command "squeeze my hand" two consecutive times.
Sevoflurane concentration was then increased to reach an end-tidal concentration of
approximately 3 vol%. When clinically indicated, ventilation was managed by the

physician and a laryngeal mask suitable for fMRI (I-gel, Intersurgical, United

Kingdom) was inserted. The fraction of inspired oxygen was then set at 0.8, and

mechanical ventilation was adjusted to maintain end-tidal carbon dioxide at steady

concentrations of 33 ± 1.71 mmHg during BS, 34 ± 1.12 mmHg during 3 vol%, and

33 ± 1.49 mmHg during 2 vol% (throughout this article, mean ± SD). Norepinephrine

was given by continuous infusion ($0.1 ± 0.01$ µg $\cdot$ kg^{-1} $\cdot$ min^{-1}) through an intravenous

catheter in a vein on the dorsum of the hand, to maintain the mean arterial blood

pressure close to baseline values (baseline, 96 ± 9.36 mmHg; BS, 88 ± 7.55 mmHg;

3 vol%, 88 ± 8.4 mmHg; 2 vol%, 89 ± 9.37 mmHg; follow-up, 98 ± 9.41 mmHg). After

insertion of the laryngeal mask airway, sevoflurane concentration was gradually

increased until the EEG showed burst-suppression with suppression periods of at

least 1,000 ms and about 50% suppression of electrical activity (reached at 4.34 ±

0.22 vol%), which is characteristic of deep anaesthesia. At that point, another 700s

of electroencephalogram and fMRI was recorded. Further 700s of data were

acquired at steady end-tidal sevoflurane concentrations of 3 and 2 vol%,

respectively, each after an equilibration time of 15 min. In a final step, etSev was

reduced to two times the concentration at LOR. However, most of the subjects

moved or did not tolerate the laryngeal mask any more under this condition:

therefore, this stage was not included in the analysis[66].

Sevoflurane administration was then terminated, and the scanner table was slid out

of the MRI scanner to monitor post-anaesthetic recovery. The volunteer was

manually ventilated until spontaneous ventilation returned. The laryngeal mask was

removed as soon as the patient opened his mouth on command. The physician

regularly asked the volunteer to squeeze their hand: recovery of responsiveness was

noted to occur as soon as the command was followed. Fifteen minutes after the time

of recovery of responsiveness, the Brice interview was administered to assess for

awareness during sevoflurane exposure; the interview was repeated on the phone

the next day. After a total of 45 min of recovery time, another resting-state combined

fMRI-EEG scan was acquired (with eyes closed, as for the baseline scan). When

participants were alert, oriented, cooperative, and physiologically stable, they were

taken home by a family member or a friend appointed in advance.

Sevoflurane dataset: MRI Data Acquisition

Although the original study acquired both functional MRI (fMRI) and
electroencephalographic (EEG) data, in the present work we only considered the
fMRI data. Data acquisition was carried out on a 3-Tesla magnetic resonance
imaging scanner (Achieva Quasar Dual 3.0T 16CH, The Netherlands) with an eight-
channel, phased-array head coil. The data were collected using a gradient echo
planar imaging sequence (echo time = 30 ms, repetition time (TR) = 1.838 s, flip
angle = 75°, field of view = 220 × 220 mm^2, matrix = 72 × 72, 32 slices, slice
thickness = 3 mm, and 1 mm interslice gap; 700-s acquisition time, resulting in 350
functional volumes). The anatomical scan was acquired before the functional scan
using a T1-weighted MPRAGE sequence with 240 × 240 × 170 voxels (1×1×1 mm
voxel size) covering the whole brain. A total of 16 volunteers completed the full
protocol and were included in our analyses; one subject was excluded due to high
motion, leaving N=15 for analysis. Here, we used fMRI data from the Awake, 3% vol,
and Recovery scans.

Ketamine

Ketamine is a multi-faceted drug, in terms of both neurophysiology and how it affects
subjective experience. Depending on dosage, it can act as a "dissociative"
anaesthetic (high dose)[22,24,75,76] or as an "atypical psychedelic" (at sub-anaesthetic
dose)[40–44]. At small doses, it has also found recent use as a fast-acting
antidepressant[197,198]. Both the anaesthetic and psychedelic effects of ketamine are in
some respect unusual; unlike widely used anaesthetics like propofol and
sevoflurane, ketamine does not exert its anaeshtetic function through agonism of
GABA receptors, nor does it recruit sleep-promoting hypothalamic nuclei, which it
appears to suppress instead[24,43]. Likewise, although ketamine does induce
psychedelic-like symptoms such as perceptual distortions, vivid imagery and
hallucinations, like classic psychedelics, it also induces prominent dissociative
symptoms of disembodiment [43,64,199–201]. Its psychedelic action is not mediated by the
serotonin 2A receptor, on which classic psychedelics operate[46,49]: although its
precise mechanisms of action are yet to be fully elucidated, ketamine appears to be

primarily an antagonist of NMDA and HCN1 receptors; however, evidence suggests
that cholinergic, aminergic, and opioid systems may also play modulatory
roles[44,45,50].

Anaesthetic ketamine dataset: Recruitment

The anaesthetic ketamine data used here have been published before [75], and we
refer the reader to the original publication for details. The original study was
approved by approval by the ethics committee of the Medical school of the university
of Liege (University Hospital, Liege, Belgium), registered at eudract 2010-023016-13.
14 right-handed volunteers were recruited via advertisements in an Internet forum (5
women; median age [range], 25 [19 to 31] years). Each participant provided written
informed consent to participation, and underwent medical interview and physical
examination before their participation.
1000

Anaesthetic ketamine dataset: Study protocol

The volunteers were requested to fast for at least 6 h from solids and 2 h from liquids
before the experimental session. After structural MR image acquisition, subjects
were removed from the MRI scanner, and 64 electroencephalogram (EEG) scalp
electrodes were placed to allow for simultaneous EEG-fMRI. Here, we only focus on
the functional MRI data.

1007

An 18-gauge intravenous catheter (BD Insyte-W; Becton Dickinson Infusion Therapy
Systems inc., USA) was then placed into a vein of the left forearm and infused using
normal saline at a rate of 20 ml/h. The intravenous line served for ketamine infusion
and eventual administration of rescue medications. a 20-gauge arterial catheter
(Arrow International Inc., USA) was also placed into the left radial artery, under strict
sterile conditions and after performing local anesthesia with 3 ml of 1% lidocaine.
This catheter was equipped with a monitoring set (TruWave, Edwards Lifesciences,
Dominican Republic) and served for arterial blood sampling and gas analysis.
standard MRI compatible anaesthesia monitoring (Magnitude 3150M; Invivo
Research, inc., USA) was also placed to allow continuous monitoring and recording
of the electrocardiogram, heart rate, blood pressure, pulse oxymetry (spO2), and

breathing frequency throughout the scanning and recovery periods. Through a loosely fitting plastic facemask, additional oxygen at a rate of 5 l/min was provided to volunteers, whose breathing always remained spontaneous. One certified anaesthesiologist and one neurologist were present throughout the experiment. After setting all needed equipment and monitoring, the volunteers were comfortably installed in the MRI tray. The most comfortable supine position attainable was sought to avoid painful stimulation related to position. all volunteers wore earplugs to attenuate noise and earphones to allow communication with investigators; one investigator remained in the MRI scan room at all times.

Ketamine was administered using a computer-controlled intravenous infusion device composed of a separate laptop computer. A 50-ml syringe was filled with normal saline containing racemic ketamine (Ketalar, Pfizer Ltd., Turkey) at a concentration of 10mg/ml. The pharmacokinetic model used to drive the pump was the domino model, which has been demonstrated to have acceptable predictive performance. This system commands infusion pump rates to allow targeting precise effect-site and plasma concentrations of ketamine, based on several biometric parameters. For each change in ketamine concentration, a 5-min equilibration period was allowed after reaching the target, to permit equilibration of ketamine concentration between body compartments. The depth of sedation was assessed using the Ramsay Scale and the University of Michigan Sedation Scale. Each evaluation took place immediately before and after each fMRi data acquisition sequence. Volunteers were asked to strongly squeeze the hand of the investigator, and the command was repeated twice. For that purpose, and for close watch of the volunteer, an investigator continuously stayed inside the MRI room.

A first fMRI data acquisition was performed in the absence of any infusion of ketamine. Ketamine infusion was then started, and its target concentration was increased by steps of 0.5 μg/ml until a level of sedation corresponding to RS 3 to 4 or UMSS 1 to 2 was reached (light sedation, S1). After the 5-min equilibration period, a novel sequence of data acquisition occurred, consisting of the same sequence of events as during W1. Ketamine target concentration was then further increased by steps of 0.5 μg/ml until RS 5 to 6 or UMSS 4 (deep sedation [s2]), and the same sequence of data acquisition was again performed. Because ketamine has a long

elimination half-life and to limit time spent in the fMRI scanner for the volunteer, the temporal order of those clinical states was not randomized. For the same reason, a recovery experimental condition could not be achieved. After those acquisitions, the infusion of ketamine was stopped, and the subject was removed from the fMRI scanner to allow for comfortable recovery. The presence of dreaming during ketamine infusion was checked through a phone call at distance from the experimental session. Here, we only consider the awake and deep sedation scans.

Anaesthetic ketamine dataset: MRI Data Acquisition

MRI data were acquired on a 3T Siemens Allegra scanner (Siemens AG, Germany; Echo Planar Imaging sequence using 32 slices; repetition time = 2,460 ms; echo time = 40 ms; field of view = 220 mm; voxel size = 3.45 × 3.45× 3 mm; and matrix size = 64 × 64× 32). A high-resolution structural T1 image was acquired in each volunteer at the beginning of the whole experiment for coregistration to the functional data.

A total of 6 participants had to be excluded from the study and further data analysis because of excessive agitation and movements (5 subjects) or voluntary withdrawal (1 subject), leaving 8 for analysis[75].

Psychedelic ketamine dataset: Recruitment

The sub-anaesthetic ("psychedelic") ketamine data included in this study have been published before[64], and we refer the reader to the original publication for details. Briefly, a total of 21 participants (10 males; mean age 28.7 years, SD = 3.2 years) were recruited via advertisements placed throughout central Cambridge, UK[64]. All participants underwent a screening interview in which they were asked whether they had previously been diagnosed or treated for any mental health problems and whether they had ever taken any psychotropic medications. Participants reporting a personal history of any mental health problems or a history of any treatment were excluded from the study. All participants were right-handed, were free of current of previous psychiatric or neurological disorder or substance abuse problems, and had

1083 no history of cardiovascular illness or family history of psychiatric disorder/substance
1084 abuse. The study was approved by the Cambridge Local Research and Ethics
1085 Committee, and all participants provided written informed consent in accordance with
1086 ethics committee guidelines.

1087

Psychedelic ketamine dataset: Study protocol

Participants were scanned (resting-state functional MRI and anatomical T1) on two
occasions, separated by at least 1 week. On one occasion, they received a
continuous computer-controlled intravenous infusion of a racemic ketamine solution
(2 mg/ml) until a targeted plasma concentration of 100 ng/ml was reached. This
concentration was sustained throughout the protocol. A saline infusion was
administered on the other occasion. Infusion order was randomly counterbalanced
across participants. The infusion was performed and monitored by a trained
anaesthetist (R.A.) who was unblinded for safety reasons, but who otherwise had
minimal contact with participants. At all other times, participants were supervised by
investigators blinded to the infusion protocol. The participants remained blinded until
both assessments were completed. Bilateral intravenous catheters were inserted into
volunteers' forearms, one for infusion, and the other for serial blood sampling. A
validated and previously implemented[202] three-compartment pharmacokinetic model
was used to achieve a constant plasma concentration of 100 ng/ml using a
computerized pump (Graseby 3500, Graseby Medical, UK). The infusion continued
for 15 min to allow stabilization of plasma levels. Blood samples were drawn before
and after the resting fMRI scan and then placed on ice. Plasma was obtained by
centrifugation and stored at −70 °C. Plasma ketamine concentrations were measured
by gas chromatography–mass spectrometry.

1108

1109 Psychedelic ketamine dataset: MRI Data Acquisition

1110 All MRI and assessment procedures were identical across assessment occasions.
1111 Scanning was performed using a 3.0 T MRI scanner (Siemens Magnetom, Trio Tim,
1112 Erlangen, Germany) equipped with a 12-channel array coil located at the Wolfson

Brain Imaging Centre, Addenbrooke's Hospital, Cambridge, UK. T2*-weighted echo-planar images were acquired under eyes-closed resting-state conditions. Participants were instructed to close their eyes and let the minds wander without going to sleep. Subsequent participant debriefing ensured that no participants fell asleep during the scan. Imaging parameters were: 3x3x3.75mm voxel size, with a time-to-repetition (TR) of 2000 ms, time-to-echo (TE) of 30 ms, flip angle of 781 in 64x64 matrix size, and 240mm field of view (FOV). A total of 300 volumes comprising 32 slices each were obtained. In addition, high- resolution anatomical T1 images were acquired using a three-dimensional magnetic-prepared rapid gradient echo (MPPRAGE) sequence. In all, 176 contiguous sagittal slices of 1.0mm thickness using a TR of 2300 ms, TE of 2.98 ms, flip angle of 91, and a FOV of 256mm in 240x256 matrix were acquired with a voxel size of $1.0mm^3$. One participant was excluded due to excessive movement, resulting in a final sample of N=20 subjects.

LSD

LSD (lysergic acid diethylamide) is perhaps the best-known among classic psychedelics, inducing a powerful state of altered consciousness with subjective experiences including hallucinations and "ego dissolution"[46,48,49]. Substantial work in humans and animals has demonstrated that LSD influences neuromodulation, having affinity for multiple receptors, primarily serotonergic (5-HT2A, 5-HT1A/B, 5-HT6, 5-HT7) and dopaminergic (D1 and D2 receptors)[46,48,52,203,204].

The main neural and subjective effects of LSD originate from its agonism of the 5-HT2A receptor: both effects are abolished by pre-treatment with the non-selective 5HT2 antagonist ketanserin, which has highest affinity for the 5HT2A receptor[171,205]. In humans, functional connectivity under LSD shows significant correspondence with the spatial distribution of the 5HT2A receptor[206]. Providing evidence for a mechanistic role, both PET maps and transcriptomic maps of the 5HT2A receptor (but not other serotonin receptors) have been shown to improve the ability of computational models to recapitulate the effects of LSD on brain activity and connectivity, as measured by fMRI[72,73,178]. Therefore, pharmacological and in silico

1144 evidence converge towards the central role of the 5HT2A receptor for LSD's ability to
1145 alter consciousness and its neural underpinnings – although other receptors have
1146 also been shown to play an auxiliary role[52].

1147

LSD dataset: Recruitment

The LSD data employed here have been extensively published, and we refer to the
original publication for details[60]. Briefly, collection of these data[60] was approved by
the National Research Ethics Service Committee London–West London and was
conducted in accordance with the revised declaration of Helsinki (2000), the
International Committee on Harmonization Good Clinical Practice guidelines and
National Health Service Research Governance Framework. Imperial College London
sponsored the research, which was conducted under a Home Office license for
research with schedule 1 drugs. All participants were recruited via word of mouth
and provided written informed consent to participate after study briefing and
screening for physical and mental health. The screening for physical health included
electrocardiogram (ECG), routine blood tests, and urine test for recent drug use and
pregnancy. A psychiatric interview was conducted and participants provided full
disclosure of their drug use history. Key exclusion criteria included: < 21 years of
age, personal history of diagnosed psychiatric illness, immediate family history of a
psychotic disorder, an absence of previous experience with a classic psychedelic
drug (e.g. LSD, mescaline, psilocybin/magic mushrooms or DMT/ayahuasca), any
psychedelic drug use within 6 weeks of the first scanning day, pregnancy,
problematic alcohol use (i.e. > 40 units consumed per week), or a medically
significant condition rendering the volunteer unsuitable for the study. Twenty healthy
volunteers with previous experience using psychedelic drugs were scanned.

1169

LSD dataset: Study protocol

Volunteers underwent two scans, 14 days apart. On one day they were given a
placebo (10-mL saline) and the other they were given an active dose of LSD (75 μg
of LSD in 10-mL saline). The order of the conditions was balanced across

participants, and participants were blind to this order but the researchers were not.
Participants carried out VAS-style ratings via button-press and a digital display
screen presented after each scan, and the 11-factor altered states of conscious-
ness (ASC) questionnaire was completed at the end of each dosing day[60]. All
participants reported marked alterations of consciousness under LSD.

The data acquisition protocols were described in detail in the original publication [60],
so we will only describe them in brief here. The infusion (drug/placebo) was
administered over 2 min and occurred 115 min before the resting-state scans were
initiated. After infusion, subjects had a brief acclimation period in a mock MRI
scanner to prepare them for the experience of being in the real machine. ASL and
BOLD scanning consisted of three seven-minute eyes closed resting state scans.
The ASL data were not analysed for this study, and will not be discussed further.

1186

LSD dataset: MRI Data Acquisition

The first and third scans were eyes-closed, resting state without stimulation, while
the second scan involved listening to music; however, this scan was not used in this
analysis. The precise length of each of the two BOLD scans included here was 7:20
minutes. For the present analysis, these two scans were concatenated together in
time. Imaging was performed on a 3T GE HDx system. High-resolution anatomical
images were acquired with 3D fast spoiled gradient echo scans in an axial
orientation, with field of view = 256x256x192 and matrix = 256x256x129 to yield
1mm isotropic voxel resolution. TR/TE = 7.9/3.0ms; inversion time = 450ms; flip
angle = 20. BOLD-weighted fMRI data were acquired using a gradient echo planer
imaging sequence, TR/TE = 2000/35ms, FoV = 220mm, 64x64 acquisition matrix,
parallel acceleration factor = 2, 90 flip angle. Thirty five oblique axial slices were
acquired in an interleaved fashion, each 3.4mm thick with zero slice gap (3.4mm
isotropic voxels). One subject aborted the experiment due to anxiety and four others
were excluded for excessive motion (measured in terms of frame-wise
displacement), leaving 15 subjects for analysis (11 males, 4 females; mean age 30.5
years, SD = 8.0 years)[60].

1204

Psilocybin

Psilocybin (4-phosphoryloxy-N,N-dimethyltryptamine), a prodrug of psilocin (4-OH-N,N-dimethyltrayptamine), is a classic psychedelic, the active compound of "magic mushrooms" of the *Psilocybe* family. Although its psychedelic effects are exerted via agonism of the serotonin 2A receptor[16,207–210], psilocin also has demonstrated affinity for additional receptors, in particular serotonin 1A and 2C receptors[211].

Psilocybin dataset: Recruitment

Data acquisition for this dataset is described in detail previously[61], and will only be summarised here. Volunteers were at least 21 years of age, with no personal or family history of a major psychiatric disorder, no substance dependence, no cardiovascular disease, and no history of adverse response to a psychedelic drug yielded datasets for nine participants. All subjects had used psilocybin at least once before, but not within 6 weeks of the study. The study was approved by a National Health Service research ethics committee and all participants gave informed consent to participate in the study.

Psilocybin dataset: Study protocol

In brief, fifteen health volunteers underwent two MRI scanning sessions at least 14 days apart. In each session, subjects were injected. with either psilocybin (2 mg dissolved in 10 mL of saline, 60-s intravenous injection) or a placebo (10 mL of saline, 60-s i.v. injection) in a counterbalanced design. The infusions began exactly 6 min after the start of the 12-min fMRI scans and lasted 60s. The subjective effects of psilocybin were felt almost immediately after injection and sustained for the remainder of the scanning session. At the end of each session, subjects were asked to rate the overall intensity of their subjective experience under the drug (or placebo) and to comment on their wakefulness level throughout the scan. As expected, all participants rated the subjective effects of psilocybin (mean intensity = 6.9/10 ± 2.6) as much stronger than placebo (mean intensity = 0.4/10 ± 0.6) and none of the

1234 subjects reported falling asleep during either scanning session. The 5 minutes of
1235 post-infusion data were used for the present analysis.

1236

Psilocybin dataset: MRI Data Acquisition

Neuroimaging data were acquired using a 3T GE HDx MRI system. Anatomical
scans were performed before each functional scan and thus prior to administering
either the drug or placebo. Structural scans were collected using a 3D fast spoiled
gradient echo scans in an axial orientation, with field of view = $256 \times 256 \times 192$ and
matrix = $256 \times 256 \times 192$ to yield 1 mm isotropic voxel resolution (repetition time/echo
time TR/TE = 7.9/3.0 ms; inversion time = 450 ms; flip angle = 20). BOLD-weighted
fMRI data were acquired at 3T using a gradient echo EPI sequence, TR/TE 3000/35
ms, field-of-view = 192 mm, 64×64 acquisition matrix, parallel acceleration factor =
2, 90° flip angle. Fifty-three oblique axial slices were acquired in an interleaved
fashion, each 3 mm thick with zero slice gap ($3 \times 3 \times 3$-mm voxels). Following the
same exclusion criteria for motion described above for the LSD dataset, N=9
subjects were kept for analysis (seven men; age, 32 ± 8.9 SD years of age).

1250

DMT

The tryptamine N,N-di-methyltryptamine (DMT) is a fast-acting mind-altering drug,
capable of inducing an immersive state of altered consciousness, with vivid and
detailed visual hallucinations. It is found endogenously in trace amounts in the
human body[46,212,213], but it is orally inert and therefore primarily studied as
intravenous injection, whereupon its effects have very rapid onset (2-5 minutes) and
offset, effectively fading within 30 minutes of administration[214–217]. Pharmacologically,
DMT binds to sigma-1 and serotonin (particularly 2A and 2C) receptors, but also
dopamine D1 and alpha-adrenergic receptors[218–221].

1260

DMT dataset: Recruitment

The original DMT study (Timmermann *et al.,* under review) was approved by the
National Research Ethics (NRES) Committee London – Brent and the Health
Research Authority and was conducted under the guidelines of the revised
Declaration of Helsinski (2000), the International Committee on Harmonisation Good
Clinical Practices guidelines, and the National Health Service Research Governance
Framework. Imperial College London sponsored the research, which was conducted
under a Home Office license for research with Schedule 1 drugs. An initial visit was
focused on assessing physical and mental health to ensure suitability, and
participants provided written informed consent. In total, 20 participants completed all
study visits (7 female, mean age = 33.5 years, SD = 7.9).

1272
1273

DMT dataset: Study protocol

This was a single-blind, placebo-controlled, counter-balanced design. Volunteers
participated in two testing days, 2 weeks apart. On each testing day, volunteers
(after testing for drugs of abuse) were involved in 2 separate scanning sessions. In
this initial session (task-free) they received intravenous (IV) administration of either
placebo (saline) or DMT (in fumarate form) in a counter-balanced order (half of the
participants received placebo and the other half received DMT). This first session
always consisted of continuous resting-state scans which lasted 28 minutes with
DMT/placebo administered at the end of the 8th minute and scanning was over 20
minutes after injection. Participants lay in the scanner with their eyes closed (an eye
mask was used to prevent eyes-opening). Following the scanning procedure,
participants were interviewed and completed questionnaires designed to assess the
subjective effects experienced during the scan. Here, we focused on the 8 minutes
post-infusion, corresponding to peak DMT experience.

1288

DMT dataset: MRI Data Acquisition

MR images were acquired in a 3T MR scanner (Siemens Magnetom Verio syngo MR
B17) using a 12-channel head coil for compatibility with EEG acquisition. Functional
imaging was performed using a T2*-weighted BOLD sensitive gradient echo planar
imaging sequence (repetition time (TR) = 2000ms, echo time (TE) = 30ms,

acquisition time (TA) = 28.06 mins, flip angle (FA) = 80°, voxel size = 3.0 x 3.0 x

3.0mm3, 35 slices, interslice distance = 0mm. Whole-brain T1-weighted structural

images were also acquired. EEG was also acquired, but here we only focus on the

fMRI data. Seven out of 20 participants were discarded from analyses due to

excessive movement, or only completing one session, leaving 13 subjects for

analysis.

1300

Ayahuasca

The Amazonian beverage ayahuasca is typically used in shamanic religious rituals,

where it is obtained as a tea made from two plants: *Psychotria viridis* and

Banisteriopsis caapi. *Psychotria viridis* contains DMT, which binds to sigma-1 and

serotonin (particularly 2A) receptors[218,219]. *Banisteriopsis caapi* contains beta-

carboline alkaloids, notably harmine, tetrahydroharmine (THH), and harmaline. As

potent monoamine oxidase inhibitors (MAOI), harmine and harmaline prevent the

degradation of DMT by liver MAO that would otherwise render it orally inert;

additionally, they also increase levels of monoamine neurotransmitters; and THH

acts as a mild selective serotonin reuptake inhibitor and a weak MAOI[222–224]. As for

other classic psychedelics, engagement of the 5-HT2A receptor appears to be a

necessary condition for the brain and subjective effects of ayahuasca to manifest [225].

1313

Ayahuasca dataset: Recruitment

The ayahuasca data that we used have been published before[62,144], and we refer the

reader to the original publication for details. Briefly, data were obtained from 9

healthy right-handed adult volunteers (mean age 31.3, from 24 to 47 years), all who

were experienced users of Ayahuasca with at least 5 years use (twice a month) and

at least 8 years of formal education. The experimental procedure was approved by

the Ethics and Research Committee of the University of São Paulo at Ribeirão Preto

(process number 14672/2006). Written informed consent was obtained from all

volunteers, who belonged to the Santo Daime religious organisation. All

experimental procedures were performed in accordance with the relevant guidelines

and regulations. Volunteers were not under medication for at least 3 months prior to
the scanning session and were abstinent from caffeine, nicotine and alcohol prior to
the acquisition. They had no history of neurological or psychiatric disorders, as
assessed by DSM-IV structured interview71. Subjects ingested 120–200 mL (2.2
mL/kg of body weight) of Ayahuasca known to contain 0.8 mg/mL of DMT and 0.21
mg/mL of harmine. Harmaline was not detected via the chromatography analysis, at
the threshold of 0.02 mg/mL7.

1331

Ayahuasca dataset: Study protocol

Volunteers underwent two distinct fMRI scanning sessions: (i) before and (ii) 40
minutes after Ayahuasca intake, when the subjective effects become noticeable (the
volunteers drank 2.2 mL/kg of body weight and the Ayahuasca contained 0.8 mg/mL
of DMT and 0.21 mg/mL of harmine). In both cases, participants were instructed to
close their eyes and remain awake and at rest, without performing any task.

1338

Ayahuasca dataset: MRI Data Acquisition

The fMRI images were obtained in a 1.5 T scanner (Siemens, Magneton Vision),
using an EPI-BOLD like sequence comprising 150 volumes, with the following
parameters: TR = 1700 ms; TE = 66 ms; FOV = 220 mm; matrix 64 ×64; voxel
dimensions of 1.72 mm × 1.72 mm × 1.72 mm. Whole brain high resolution T1-
weighted images were also acquired (156 contiguous sagittal slices) using a
multiplanar reconstructed gradient-echo sequence, with the following parameters:
TR = 9.7 ms; TE = 44 ms; flip angle 12°; matrix 256 × 256; FOV = 256 mm, voxel
size = 1 mm× 1 mm× 1 mm. Data from one volunteer were excluded from analyses
due to acquisition limitations resulting in incomplete brain coverage. The final dataset
included 8 subjects.

1350

MDMA

3,4-methylenedioxymethamphetamine (MDMA) combines the subjective effects of a stimulant and a psychedelic, inducing powerful euphoria and prosociality, but also mild visual hallucinations[226–228]. It inhibits reuptake of noradrenaline, dopamine, and serotonin by acting on their respective transporters, and it also stimulates their release – with preferential effects on 5-HT mediating at least in part its induction of positive mood[226–228].

MDMA dataset: Recruitment

The MDMA dataset has been published before[63,229], and we refer the reader to the original publications for details. The study was approved by the National Research Ethics Service West London Research Ethics Committee, Joint Compliance and Research Office of Imperial College London, Research Ethics Committee of Imperial College London, Head of the Department of Medicine of Imperial College London, Imanova Centre for Imaging Science, and Faculty of Medicine of Imperial College London. The study was conducted in accordance with Good Clinical Practice guidelines. A Home Office Licence was obtained for the storage and handling of a Schedule 1 drug. Imperial College London sponsored the research.

The study included 25 healthy participants (mean age, 34±11 years; 7 females) with at least one previous experience with MDMA. None of the participants had used MDMA for at least 7 days or other drugs for at least 48 hours, which was confirmed by a urine screen. An alcohol breathalyzer test confirmed that none of the participants had recently consumed alcohol. Volunteers were screened for good physical and mental health, and magnetic resonance imaging compatibility. All subjects were deemed physically and mentally healthy, and none had any history of drug or alcohol dependence or diagnosed psychiatric disorder.

MDMA dataset: Study protocol

The study design was within-subjects, and placebo-controlled. Participants were scanned twice, once after MDMA (100 mg encapsulated MDMA-HCl) and once after placebo (encapsulated ascorbic acid/vitamin-C), in a double-blind, randomized,

counterbalanced order. The MDMA and placebo study days were separated by 7

days, and on each occasion, participants underwent two arterial spin labelling (ASL;

not included in the present analysis) and two resting-state BOLD fMRI scans within a

90-minute scan session. Biochip Array Technology (Randox Laboratories Ltd., Co.,

Antrim, United Kingdom) was used to detect MDMA from plasma samples obtained

shortly after each participant's MDMA scanning session (i.e., 2 hours after capsule

ingestion).

1390

The first resting-state BOLD scan took place 60 minutes after capsule ingestion and

the second resting-state BOLD scan occurred 113 minutes after capsule ingestion.

Participants relaxed with their eyes closed during the ASL and BOLD resting state

scans. Peak subjective effects were reported ~100 minutes post administration of

MDMA, consistent with the plasma t-max of MDMA. Therefore, here we focus on the

second MDMA and placebo rs-fMRI scans.

1397

MDMA dataset: MRI Data Acquisition

MR images were acquired on a 3T Siemens Tim Trio (Siemens Healthcare,

Erlangen, Germany) using a 32-channel phased array head coil. Anatomical

reference images were acquired using the ADNI-GO recommended MPRAGE

parameters (1 mm isotropic voxels, TR = 2300 ms, TE = 2.98 ms, 160 sagittal slices,

256 x 256 in-plane resolution, flip angle = 9 degrees, bandwidth = 240 Hz/pixel,

GRAPPA acceleration = 2). T2*-weighted echo-planar images (EPI) were acquired

for the resting state BOLD functional scan using 3 mm isotropic voxels in a 192 mm

in-plane FOV, TR = 2 s, echo time = 31 ms, 80 degree flip angle, 36 axial slices in

each TR, bandwidth = 2298 Hz/pixel, and a GRAPPA acceleration of 2. Each rs-fMRI

scan lasted for 6 minutes, corresponding to 180 functional volumes.

1409

Modafinil

Modafinil is a wakefulness promoting drug used for the treatment of sleep disorders

such as narcolepsy (under the commercial name Provigil), as well as finding use as

a cognitive enhancer for attention and memory[6,7,65], and to combat the cognitive

symptoms of Attention Deficit/Hyperactivity Disorder (ADHD), and mood disorders,

owing to its lower addiction risk in comparison with amphetamine-like
psychostimulants[6–9,230]. This drug has a broad neurotransmitter profile: it acts as a
blocker of the dopamine and noradrenaline transporters, as well as modulating locus
coeruleus noradrenergic firing, and acting on the wake-promoting hypothalamic
neuropeptide orexin to activate the histamine system; it also influences both the
glutamate and GABA systems[8,231–235].

1421

Modafinil dataset: Recruitment

The modafinil dataset has been published before[65,236]. As reported in the original
publication, the study was approved by the ethics committee of University of Chieti
(PROT 2008/09 COET on 14/10/2009) and conducted in accordance with the
Helsinki Declaration. The study design was explained in detail and written informed
consent was obtained from all participants involved in our study. Recruitment was
performed throughout February 2011, drug/placebo administration and fMRI
acquisitions started on March 2011, went on until January 2012, and the study was
completed with the last fMRI session in January 2012. After securing financial
coverage for costs related to the analysis of the study, the trial was registered on
10/09/2012   (NCT01684306http://clinicaltrials.gov/ct2/show/NCT01684306).   After
obtaining registration, the double-blind study was opened and analyzed.

This dataset was obtained from the OpenfMRI database. Its accession number is
ds000133. A total of twenty six young male right-handed (as assessed by the
Edinburgh Handedness inventory) adults (age range: 25–35 years) with comparable
levels of education (13 years) were enrolled. All subjects had no past or current
signs of psychiatric, neurological or medical (hypertension, cardiac disorders,
epilepsy) conditions as determined by the Millon test and by clinical examination.
Subjects showing visual or motor impairments were excluded as well as individuals
taking psychoactive drugs or having a history of alcohol abuse. All volunteers were
instructed to maintain their usual amount of nicotine and caffeine intake and avoid
alcohol consumption in the 12h before the initiation of the study.

1444

Modafinil dataset: Study protocol

Study subjects received, in a double blind fashion, either a single dose of modafinil

(100 mg) (modafinil group; N=13) or a placebo (placebo group; N=13) pill identical to

the drug. Randomization of study subjects was obtained by means of a random

number generator. Here, we only considered data from the modafinil group. The day

after drug/placebo assumption, subjects were asked about perceived side effects

and, in particular, sleep disturbances. All but one reported no modafinil-induced side-

effects or alterations in the sleep-wake cycle. Rs-fMRI BOLD data were separated in

three runs lasting four minutes each followed by high resolution T1 anatomical

images. Two scanning sessions took place: one before ingesting the drug/placebo,

and one 3 hours later, to account for pharmacokinetics. Subjects were asked to relax

while fixating the central point in the middle of a grey-background screen that was

projected on an LCD screen and viewed through a mirror placed above the subject's

head. Subject head was positioned within an eight-channel coil and foam padding

was employed to minimize involuntary head movements.

1460

Modafinil dataset: MRI Data Acquisition

BOLD functional imaging was performed with a Philips Achieva 3T Scanner (Philips

Medical Systems, Best, The Netherlands), using T2*-weighted echo planar imaging

(EPI) free induction decay (FID) sequences and applying the following parameters:

TE 35 ms, matrix size 64x64, FOV 256 mm, in-plane voxel size 464 mm, flip angle

75 degrees, slice thickness 4 mm and no gaps. 140 functional volumes consisting of

30 transaxial slices were acquired per run with a volume TR of 1,671 ms. High-

resolution structural images were acquired at the end of the three rs-fMRI runs

through a 3D MPRAGE sequence employing the following parameters: sagittal,

matrix 256x256, FOV 256 mm,slice thickness 1 mm, no gaps, in-plane voxel size 1

mm x 1 mm,flip angle 12 degrees, TR = 9.7 ms and TE = 4 ms. Two subjects from

the modafinil group were excluded from analysis due to acquisition limitations,

leaving N=11 subjects for analysis.

1474

Methylphenidate

Methylphenidate is used as a cognitive enhancer to treat the cognitive symptoms of ADHD and narcolepsy (under the name Ritalin)[4,11]. Pharmacologically, it inhibits the reuptake of both dopamine and noradrenaline by blocking their transporters; although yet to be conclusively confirmed, there is also in vitro evidence suggesting an additional minor affinity of methylphenidate for the 5-HT1A receptor[237–243].

Methylphenidate dataset: Recruitment

The methylphenidate dataset used here has been published before[10,51]. Unlike the other datasets included in this study, the methylphenidate data were not obtained from healthy controls, but rather from a cohort of patients suffering from traumatic brain injury (TBI). Volunteers with a history of moderate to severe traumatic brain injury (inclusion criteria: age 18–60 years and not recruited to more than three research studies within the calendar year) were referred from the Addenbrooke's Neurosciences Critical Care Unit Follow-Up Clinic, Addenbrooke's Traumatic Brain Injury Clinic and The Royal London Hospital Intensive Care Unit (see the original publication for details of patient injuries). The patients were sent a written invitation to take part in the study. All volunteers gave written informed consent before participating in the study.

Thirty-eight volunteers were recruited to the study; 17 (12 male, 5 female) into the TBI arm of the study and 21 (13 male, 8 female) into the healthy control (HC) arm of the study. Exclusion criteria included National Adult Reading Test (NART) <70, Mini Mental State Examination (MMSE) <23, left-handedness, history of drug/alcohol abuse, history of psychiatric or neurological disorders, contraindications for MRI scanning, medication that may affect cognitive performance or prescribed for depression, and any physical handicap that could prevent the completion of testing.

Our sample contains mostly patients with diffuse axonal injuries and small lesions. The patients were at least 6 months post TBI. Four sustained moderate TBI with a score of between 9 and 12 on the Glasgow Coma Scale (GCS) and 11 sustained

1504 severe TBI with a GCS score of 8 or below on presentation. The mean age of the
1505 patient group was 36 years (±13 years).

1506

Methylphenidate dataset: Study protocol

The study consisted of two visits (separated by 2–4 weeks) for both groups of
participants. The TBI volunteers were randomly allocated in a Latin square design to
receive one of the two interventions on the first visit (a placebo tablet or 30 mg tablet
of methylphenidate), and the alternate intervention on the second visit. The decision
to use 30 mg of methylphenidate was based on comparable doses used in previous
studies in healthy participants, as well as NICE guidelines for medication in adults
(www.nice.org.uk) which stipulate that when methylphenidate is titrated for side
effects and responsiveness in each individual subject, the dose should range from a
minimum of 15 mg to a maximum dose of 100 mg. As the dose of methylphenidate
was not calculated by the participant's body weight, an interventional dose at the
lower end of the dose range was chosen. After a delay of 75 min to ensure that peak
plasma levels of methylphenidate were reached, the volunteers completed an MRI
scan which included both fMRI and structural image acquisition. The healthy controls
attended their two fMRI assessments at the same time interval as the patients, but
without any pharmacological intervention. Therefore, here we only considered the
patient data.

1524

Methylphenidate dataset: MRI Data Acquisition

MRI data were acquired on a Siemens Trio 3-Tesla MR system(Siemens AG,
Munich, Germany). MRI scanning started with the acquisition of a localizer scan and
was followed by a 3D high resolution MPRAGE image [TR 2,300 ms, Echo Time
(TE) 2.98 ms, Flip Angle 9∘, FOV 256×256 mm2]. Diffusion Tensor Imaging (DTI)
data (63non-collinear directions,b=1,000 s/mm2with one volume acquired without
diffusion weighting (b=0), echo time 106ms, repetition time 1,700 ms, field of view
192×192 mm,2 mm3isotropic voxels) were also collected to investigate white matter
integrity. Here, we did not analyse the DTI data.

Functional imaging data were acquired using an echo-planar imaging (EPI)
sequence with parameters TR = 2,000 ms, TE = 30 ms, Flip Angle = 78∘, FOV
192×192 mm2, in-plane resolution 3.0×3.0 mm, 32 slices 3.0 mm thick with a gap of
0.75 mm between slices. Two patients were excluded from the analysis (one patient
only attended one of the study sessions and the other had excessive movement
artifacts in their fMRI scan), leaving N=15 patients for analysis.

1540

1541 Functional MRI preprocessing and denoising

1542 Preprocessing

To maximise the uniformity between the different datasets included here, we elected
to preprocess all datasets using the same pipeline, as opposed to relying on the
different pipelines originally employed by the group that collected each dataset. For
the same reason, we also applied the same denoising procedure across all datasets.

For each dataset and condition, we applied a standard preprocessing pipeline, which
we have previously employed with pharmaco-MRI datasets comprising both
anaesthetics and psychedelics[18,21,145,196], demonstrating its suitability for the present
analysis. Preprocessing was performed using the CONN toolbox, version 17f
(CONN; http://www.nitrc.org/projects/conn)[244] based on Statistical Parametric
Mapping 12 (http://www.fil.ion.ucl.ac.uk/spm), implemented in MATLAB 2016a. The
pipeline involved the following steps: removal of the first 10s, to achieve steady-state
magnetization; motion correction; slice-timing correction; identification of outlier
volumes for subsequent scrubbing by means of the quality assurance/artifact
rejection software *art* (http://www.nitrc.org/projects/artifact_detect); normalisation to
Montreal Neurological Institute (MNI-152) standard space (2 mm isotropic
resampling resolution), using the segmented grey matter image from each
volunteer's T1-weighted anatomical image, together with an *a priori* grey matter
template.

1561

1562 Denoising

Denoising was also performed using the CONN toolbox, using the same approach as in our previous publications with pharmaco-MRI datasets[18,21,145,196], which has also been adopted with similar pharmaco-MRI datasets in publications by independent groups[245].

Pharmacological agents can induce alterations in physiological parameters (heart rate, breathing rate, motion) or neurovascular coupling[246]. The anatomical CompCor (aCompCor) method removes physiological fluctuations by extracting principal components from regions unlikely to be modulated by neural activity; these components are then included as nuisance regressors[247]. Following this approach, five principal components were extracted from white matter and cerebrospinal fluid signals (using individual tissue masks obtained from the T1-weighted structural MRI images)[245]; and regressed out from the functional data together with six subject-specific realignment parameters (three translations and three rotations) as well as their first-order temporal derivatives; followed by scrubbing of outliers identified by ART, using Ordinary Least Squares regression[244]. Finally, the denoised BOLD signal timeseries were linearly detrended and band-pass filtered to eliminate both low-frequency drift effects and high-frequency noise, thus retaining frequencies between 0.008 and 0.09 Hz.

The step of global signal regression (GSR) has received substantial attention in the literature as a denoising method[248–250]. GSR mathematically mandates that approximately 50% of correlations between regions will be negative[251]; however, the proportion of anticorrelations between brain regions has been shown to vary across states of consciousness, including anaesthesia and psychedelics[18,145]. Furthermore, recent work has demonstrated that the global signal contains information about states of consciousness, across pharmacological and pathological perturbations[252]. Therefore, in line with ours and others' previous studies, here we avoided GSR in favour of the aCompCor denoising procedure, which is among those recommended[249].

Summarising pharmacological effects on brain function

For each subject at each condition, the denoised regional BOLD signals were parcellated into 100 cortical regions according to the local-global functional

parcellation of Schaefer and colleagues[112]. The parcellated regional BOLD signals

were then correlated ("functional connectivity"); after removing negative-valued

edges, the regional strength of functional connectivity (node strength) was measured

for each region. The regional change in FC strength was then quantified for each

subject (for the methylphenidate dataset, this was quantified with respect to the

mean of controls' node strength values). Finally, for each dataset, we computed the

mean across subjects of the FC strength deltas. Therefore, each pharmacological

intervention was summarised as one vector of regional FC strength deltas (Figure 1).

1603

Receptor maps from Positron Emission Tomography

1604

Receptor densities were estimated using PET tracer studies for a total of 19

receptors and transporters, across 9 neurotransmitter systems, recently made

available by Hansen and colleagues

at https://github.com/netneurolab/hansen_receptors[31]. These include dopamine

(D1[77], D2[78–81], DAT[82]), norepinephrine (NET[83–86]), serotonin (5-HT1A[87], 5-HT1B[87–

90,90–92], 5-HT2A[93], 5-HT4[93], 5-HT6[94,95], 5-HTT[93]), acetylcholine ($\alpha 4\beta 2$[96,97], M1[98],

VAChT[99,100]), glutamate (mGluR5[101,102], NMDA[103,104]), GABA (GABA-A[105]), histamine

(H3[106]), cannabinoid (CB1[107–110]), and opioid (MOR[111]). Volumetric PET images were

registered to the MNI-ICBM 152 nonlinear 2009 (version c, asymmetric) template,

averaged across participants within each study, then parcellated and

receptors/transporters with more than one mean image of the same tracer (5-HT1b,

D2, VAChT) were combined using a weighted average. See the dedicated article by

Hansen et al[31] for detailed information about each PET dataset and their respective

acquisition and limitations.

1619

Partial Least Squares Analysis

1620

1621

PLS analysis (implemented using MATLAB code from Hansen et al., 2021)[59]

available at https://github.com/netneurolab/hansen_genescognition/tree/master/PLS)

was used to relate regional neurotransmitter density to pharmacologically-induced

functional connectivity changes. PLS analysis is an unsupervised multivariate

statistical technique that decomposes relationships between two datasets (in our

case, neurotransmitter density with *n* regions and *r* neurotransmitters, X_{nxr}, and drug-

induced functional connectivity changes, Y_{nxd} with *n* regions and *d* drugs) into

orthogonal sets of latent variables with maximum covariance, which are linear

combinations of the original data[119,120]. In other words, PLS finds components from

the predictor variables (100 × 19 matrix of regional neurotransmitter receptor and

transporter density scores) that have maximum covariance with the response

variables (100 × 15 matrix of regional changes in FC induced by different drugs). The

PLS components (i.e., linear combinations of the weighted neurotransmitter density)

are ranked by covariance between predictor and response variables, so that the first

few PLS components provide a low-dimensional representation of the covariance

between the higher dimensional data matrices. Thus, the first PLS component

(PLS1) is the linear combination of the weighted neurotransmitter density scores that

have a brain expression map that covaries the most with the map of regional FC

changes.

1641

1642   This is achieved by z-scoring both data matrices column-wise and applying singular

1643   value decomposition on the matrix *Y'X*, such that:

1644

1645   *(Y'X)' = USV'*

1646   (1)

1647

where U_{gxt} and V_{txt} are orthonormal matrices consisting of left and right singular

vectors and S_{txt} is a diagonal matrix of singular values. The i[th] columns of *U* and *V*

constitute a latent variable, and the i[th] singular value in *S* represents the covariance

between singular vectors. The i[th] singular value is proportional to the amount of

covariance between neurotransmitter density, and drug-induced FC changes

captured by the i[th] latent variable, where the effect size can be estimated as the ratio

of the squared singular value to the sum of all squared singular values. In the

present study, the left singular vectors (that is, the columns of *U*) represent the

degree to which each neurotransmitter contributes to the latent variable and

demonstrate the extracted association between neurotransmitter density and drug-

induced FC changes (neurotransmitter weights). The right singular vectors (that is,

the columns of *V*) represent the degree to which the FC changes contribute to the

same latent variable (term weights). Positively weighed neurotransmitters covary

with positively weighed drug-induced changes, and negatively weighed neurotransmitters covary with negatively weighed drug-induced changes.

Scores at each brain region for each latent variable can be computed by projecting the original data onto the singular vector weights. Positively scored brain regions are regions that demonstrate the covariance between the prevalence of positively weighted neurotransmitters and positively weighted drug-induced effects (and vice versa for negatively scored brain regions). Loadings for each variable were computed as the Pearson's correlation between each individual variable's activity (neurotransmitter density and drug-induced FC changes) and the PLS analysis-derived neurotransmitter score pattern. Squaring the loading (a correlation) equals the percentage variance shared between an original variable and the PLS analysis-derived latent variable. Variables with high absolute loadings are highly correlated to the score pattern, indicating a large amount of shared variance between the individual variable and the latent variable. We confirmed that PLS1 explained the largest amount of variance by testing across a range of PLS components (between 1 and 12) and quantifying the relative variance explained by each component.

Hierarchical organisation

We quantified the spatial similarity of each pharmacologically-induced pattern of change in FC strength, with several canonical maps of hierarchical brain organisation ("canonical brain hierarchies") derived from multimodal neuroimaging. We considered the anatomical gradient of intracortical myelination obtained from T1w/T2w MRI ratio[58]; evolutionary cortical expansion obtained by comparing human and macaque[126]; the principal component of variation in gene expression from the Allen Human Brain Atlas transcriptomic database (AHBA; https://human.brain-map.org/), referred to as "AHBA PC1"[54,59,127]; the principal component of variation in task activation from NeuroSynth, (https://github.com/neurosynth/neurosynth), an online meta-analytic tool that synthesizes results from more than 15,000 published fMRI studies by searching for high-frequency key words that are published alongside fMRI voxel coordinates, using the volumetric association test maps (referred to as

1694 "NeuroSynth PC1")[54,59,128]; the map of cerebral blood flow[54]; the principal gradient of

1695 variation in functional connectivity[57]; and a recently derived gradient of regional

1696 prevalence of different kinds of information, from redundancy to synergy[129].

1697

ENIGMA cortical vulnerability data

1698

1699

Patterns of cortical thickness were collected for the available 11 neurological,

neurodevelopmental, and psychiatric disorders from the ENIGMA (Enhancing

Neuroimaging Genetics through Meta-Analysis) consortium and the *enigma* toolbox

(https://github.com/MICA-MNI/ENIGMA)[130]: 22q11.2 deletion syndrome[131], attention-

deficit/hyperactivity disorder[132], autism spectrum disorder[133], idiopathic generalized

epilepsy[134], right temporal lobe epilepsy[134], left temporal lobe epilepsy[134],

depression[135], obsessive-compulsive disorder[136], schizophrenia[137], bipolar

disorder[138], and Parkinson's disease[139]. The ENIGMA consortium is a data-sharing

initiative that relies on standardized image acquisition and processing pipelines, such

that disorder maps are comparable[253]. Altogether, over 21,000 patients were

scanned across the thirteen disorders, against almost 26,000 controls. The values

for each map are z-scored effect sizes (Cohen's *d*) of cortical thickness in patient

populations versus healthy controls. Imaging and processing protocols can be found

at http://enigma.ini.usc.edu/protocols/.

For every brain region, we constructed an 11-element vector of disorder abnormality,

where each element represents a disorder's cortical abnormality at the region. For

every pair of brain regions, we correlated the abnormality vectors to quantify how

similarly two brain regions are affected across disorders. This results in a region-by-

region matrix of "disorder co-susceptibility"[113].

1719

Gradients from diffusion map embedding

1720

We used the *BrainSpace* toolbox[140] (http://brainspace.readthedocs.io) to obtain

gradients from diffusion map embedding. A joint network of disease- and drug- co-

susceptibility was obtained from the network fusion procedure of Paquola *et al*

(2020)[141], through horizontal concatenation of matrices and production of a node-to-

node affinity matrix using row-wise normalised angle similarity.

We then employed diffusion map embedding, a nonlinear manifold learning

technique based on the graph Laplacian[140,254], to obtain a low dimensional

representation of the joint drug- and disorder-susceptibility. A single parameter α

controls the influence of the sampling density on the manifold (α = 0, maximal

influence; α = 1, no influence). Following extensive previous work using this

approach with neuroimaging data[57,140,141,255], we set α = 0.5, to retain the global

relations between data points in the embedded space. The ability to combine global

and local geometry differentiates diffusion maps from global-only methods, such as

PCA and multidimensional scaling[141]. A small number of components can be

identified based on decreasing eigenvalues. The decay of each eigenvector obtained

from the diffusion map embedding provides an overall measure of the connectivity

between nodes along the axis delineated by its spatial distribution on the brain

(gradient).

Statistical Analysis

1739

The statistical significance of the variance explained by each PLS model was tested

by permuting the response variables 1,000 times, while considering the spatial

dependency of the data by using spatial autocorrelation-preserving permutation

tests, termed spin tests[121–124]. Parcel coordinates were projected onto the spherical

surface and then randomly rotated and original parcels were reassigned the value of

the closest rotated parcel (10,000 repetitions). The procedure was performed at the

parcel resolution rather than the vertex resolution to avoid upsampling the data. In

PLS analysis, the spin test is applied to the singular values (or equivalently, the

covariance explained) of the latent variables, producing a null distribution of singular

values. This is done applying PLS analysis to the original *X* matrix and a spun *Y*

matrix. The spin test embodies the null hypothesis that neurotransmitter density and

drug-induced FC changes are spatially correlated with each other only because of

inherent spatial autocorrelation. The p-value is computed as the proportion of null

singular values that are greater in magnitude than the empirical singular values.

Thus, these p-values represent the probability that the observed spatial

correspondence between neurotransmitter density and drug-induced FC changes could occur by randomly correlating maps with comparable spatial autocorrelation. Likewise, spatial similarity between brain maps was quantified in terms of Spearman correlation, and statistical significance was assessed against a spin-based null model with preserved spatial autocorrelation, as described above[121–124].

Data and code availability

Pharmacological-fMRI data are available upon request from the corresponding authors of the original publications referenced herein. The neurotransmitter receptor and transporter PET maps are available at https://github.com/netneurolab/hansen_receptors. The Allen Human Brain Atlas transcriptomic database is available at https://human.brain-map.org/; NeuroSynth is available at https://github.com/neurosynth/neurosynth. The ENIGMA toolbox and data are available at https://github.com/MICA-MNI/ENIGMA. MATLAB code for Partial Least Squares analysis is freely available at https://github.com/netneurolab/hansen_genescognition/tree/master/PLS.

Acknowledgements

This work was carried out thanks to support from the Gates Cambridge Trust (OPP 1144) [to AIL]; the Wellcome Trust Research Training Fellowship (grant no. 083660/Z/07/Z), Raymond and Beverly Sackler Studentship, and the Cambridge Commonwealth Trust [to RA]; the Canadian Institute for Advanced Research (CIFAR; grant RCZB/072 RG93193) [to DKM and EAS]; the Cambridge Biomedical Research Centre and NIHR Senior Investigator Awards and the British Oxygen Professorship of the Royal College of Anaesthetists [to DKM]; the Stephen Erskine Fellowship at Queens' College, Cambridge [to EAS]. BM is supported by the Natural Sciences and Engineering Research Council of Canada (NSERC Discovery Grant RGPIN #017-04265) and Canada Research Chairs Program. JYH is supported by the Helmholtz International BigBrain Analytics & Learning Laboratory, the Natural Sciences and Engineering Research Council of Canada, and Fonds de reserches de Québec. AMO is supported by the Canada Excellence Research Chairs program (215063); LN acknowledges support by the L'Oreal-Unesco for Women in Science Excellence Research Fellowship. LR acknowledges support of the Imperial College President's Scholarship. PC and NLNA are supported by the Belgian National Funds for Scientific Research (F.R.S-FNRS). PC is also

supported by the GIGA-Doctoral School for Health Sciences (University of Liège). NLNA also acknowledges the support of the Human Brain Project. SLS is supported by funds from the Italian Department of Education [Fondo per gli Investimenti della Ricerca di Base (FIRB) 2003; Programmi di Ricerca di Rilevante Interesse nazionale (PRIN) 2008]. RLC-H was supported by the Alex Mosley Charitable Trust and supporters of the Centre for Psychedelic Research during the period of data collection and now holds the Ralph Metzner Distinguished Professorship at UCSF. The original LSD study received support from a Crowd Funding Campaign and the Beckley Foundation, as part of the Beckley-Imperial Research Programme. The MDMA research was supported by funds provided by the British public service broadcast station Channel 4 and was performed as part of the Beckley Foundation–Imperial College research program. The DMT study was funded via a donation by Patrick Vernon. The Beckley Foundation mediated Patrick Vernon's support. The psilocybin study received support from the Beckley Foundation and financial support from the Neuropsychoanalysis Foundation, Multidisplinary Association for Psychedelic Studies, and the Heffter Research Institute. The Cambridge ketamine study was funded by the Bernard Wolfe Health Neuroscience Fund and the Wellcome Trust. Data acquisition for the sevoflurane dataset was funded by the Departments of Anesthesiology, Neurology, and Neuroradiology of the Klinikum rechts der Isar of the Technical University Munich. The anaesthetic ketamine study was supported by the Belgian Society of Anesthesiology, Resuscitation, Perioperative medicine and Pain management (BeSARPP, Brussels, Belgium), the Belgian National Funds for Scientific Research (Brussels, Belgium), the European Commission (Brussels, Belgium), the James McDonnell Foundation (Saint Louis, Missouri), the European Space Agency (Brussels, Belgium), the Mind Science Foundation (San Antonio, Texas), the French Speaking Community Concerted Research Action (ARC - 06/11 – 340, Brussels, Belgium), the Public Utility Foundation "Université Européenne du Travail" (Brussels, Belgium), "Fondazione Europea di Ricerca Biomedica" (Milan, Italy), and the University and University Hospital of Liege (Liege, Belgium). This work was performed using resources provided by the Cambridge Service for Data Driven Discovery (CSD3) operated by the University of Cambridge Research Computing Service (www.csd3.cam.ac.uk), provided by Dell EMC and Intel using Tier-2 funding from the Engineering and Physical Sciences Research Council (capital grant EP/T022159/1), and DiRAC funding from the Science and Technology Facilities Council (www.dirac.ac.uk). Computing infrastructure at the Wolfson Brain Imaging Centre (WBIC-HPHI) was funded by the MRC research infrastructure award (MR/M009041/1). We acknowledge the contribution of Santo Daime members for volunteering and for providing the Ayahuasca. We also thank members of the Cognition and Consciousness Imaging Group, and of the Network Neuroscience Lab, for many helpful discussions.

Conflicts of Interest

RCH reports receiving scientific advisory fees in the last 2 years from: Entheon Biomedical and Beckley Psytech. D.B.A. serves as scientific advisor of Biomind Labs. VB has had financial relationships with the following companies: Orion Pharma, Medtronic, Edwards, and Elsevier. All other authors report no conflicts of interest.

Author Contributions

AIL, JYH, EAS, BM conceived the study. AIL, JYH, EAS, BM designed the methodology and the analysis. AIL analysed the data. JYH, ARDP contributed to data analysis. RA, DKM, EAS, LR, RLCH, CT, DG, AR, RI, DJ, VB, AV, AD, OJ, MAB, NLNA, PC, AEM, DBdA, SLS, AMO, LN were involved in design, execution, and data acquisition and curation for the original studies for which the present data were collected. AIL, EAS, BM wrote the manuscript with feedback from co-authors.

References

1. Wandschneider, B. & Koepp, M. J. Pharmaco fMRI: Determining the functional anatomy of the effects of medication. *NeuroImage Clin.* **12**, 691–697 (2016).
2. Brown, E. N., Lydic, R. & Schiff, N. D. General Anesthesia, Sleep, and Coma. *N. Engl. J. Med.* **27**, 2638–50 (2010).
3. Cortese, S. *et al.* Comparative efficacy and tolerability of medications for attention-deficit hyperactivity disorder in children, adolescents, and adults: a systematic review and network meta-analysis. *Lancet Psychiatry* **5**, 727–738 (2018).
4. Faraone, S. V. The pharmacology of amphetamine and methylphenidate: Relevance to the neurobiology of attention-deficit/hyperactivity disorder and other psychiatric comorbidities. *Neurosci. Biobehav. Rev.* **87**, 255–270 (2018).
5. Moldofsky, H., Broughton, R. J. & Hill, J. D. A randomized trial of the long-term, continued efficacy and safety of modafinil in narcolepsy. *Sleep Med.* **1**, 109–116 (2000).
6. Turner, D. C., Robbins, T. W., Clark, L., Aron, A. R., Dowson, J. & Sahakian, B. J. Cognitive enhancing effects of modafinil in healthy volunteers. *Psychopharmacology (Berl.)* **165**, 260–269 (2003).
7. Minzenberg, M. J. & Carter, C. S. Modafinil: a review of neurochemical actions and effects on cognition. *Neuropsychopharmacol. Off. Publ. Am. Coll. Neuropsychopharmacol.* **33**, 1477–1502 (2008).
8. Minzenberg, M. J., Watrous, A. J., Yoon, J. H., Ursu, S. & Carter, C. S. Modafinil shifts human locus coeruleus to low-tonic, high-phasic activity during functional MRI. *Science* **322**, 1700–1702 (2008).
9. Kahbazi, M., Ghoreishi, A., Rahiminejad, F., Mohammadi, M.-R., Kamalipour, A. & Akhondzadeh, S. A randomized, double-blind and placebo-controlled trial of modafinil

in children and adolescents with attention deficit and hyperactivity disorder. *Psychiatry*
Res. **168**, 234–237 (2009).

10. Manktelow, A. E., Menon, D. K., Sahakian, B. J. & Stamatakis, E. A. Working Memory
after Traumatic Brain Injury: The Neural Basis of Improved Performance with
Methylphenidate. *Front. Behav. Neurosci.* **11**, (2017).

11. Storebø, O. J. *et al.* Methylphenidate for attention deficit hyperactivity disorder (ADHD)
in children and adolescents – assessment of adverse events in non-randomised
studies. *Cochrane Database Syst. Rev.* **2018**, CD012069 (2018).

12. Carhart-Harris, R. *et al.* Trial of Psilocybin versus Escitalopram for Depression. *N. Engl.*
J. Med. **384**, 1402–1411 (2021).

13. de Gregorio, D., Aguilar-Valles, A., Preller, K. H., Heifets, B. D., Hibicke, M., Mitchell, J.
& Gobbi, G. Hallucinogens in mental health: Preclinical and clinical studies on LSD,
psilocybin, MDMA, and ketamine. *J. Neurosci.* **41**, 891–900 (2021).

14. Mitchell, J. M. *et al.* MDMA-assisted therapy for severe PTSD: a randomized, double-
blind, placebo-controlled phase 3 study. *Nat. Med. 2021 276* **27**, 1025–1033 (2021).

15. Davis, A. K., Barrett, F. S., May, D. G., Cosimano, M. P., Sepeda, N. D., Johnson, M.
W., Finan, P. H. & Griffiths, R. R. Effects of Psilocybin-Assisted Therapy on Major
Depressive Disorder: A Randomized Clinical Trial. *JAMA Psychiatry* (2020)
doi:10.1001/jamapsychiatry.2020.3285.

16. Vollenweider, F. X. & Preller, K. H. Psychedelic drugs: neurobiology and potential for
treatment of psychiatric disorders. *Nat. Rev. Neurosci.* **21**, 611–624 (2020).

17. Vargas, M. V., Meyer, R., Avanes, A. A., Rus, M. & Olson, D. E. Psychedelics and
Other Psychoplastogens for Treating Mental Illness. *Front. Psychiatry* **12**, 1691 (2021).

18. Luppi, A. I. *et al.* Consciousness-specific dynamic interactions of brain integration and
functional diversity. *Nat. Commun.* **10**, (2019).

19. Huang, Z., Zhang, J., Wu, J., Mashour, G. A. & Hudetz, A. G. Temporal circuit of
macroscale dynamic brain activity supports human consciousness. *Sci. Adv.* **6**, 87–98
(2020).

20. Campbell, J. M., Huang, Z., Zhang, J., Wu, X., Qin, P., Northoff, G., Mashour, G. A. &
Hudetz, A. G. Pharmacologically informed machine learning approach for identifying
pathological states of unconsciousness via resting-state fMRI. *NeuroImage* **206**,
116316 (2020).

21. Luppi, A. I. *et al.* Connectome Harmonic Decomposition of Human Brain Dynamics
Reveals a Landscape of Consciousness. *bioRxiv* (2020)
doi:10.1101/2020.08.10.244459.

22. Sarasso, S. *et al.* Consciousness and complexity during unresponsiveness induced by
propofol, xenon, and ketamine. *Curr. Biol.* **25**, 3099–3105 (2015).

23. Schartner, M. M., Carhart-Harris, R. L., Barrett, A. B., Seth, A. K. &
Muthukumaraswamy, S. D. Increased spontaneous MEG signal diversity for
psychoactive doses of ketamine, LSD and psilocybin. *Sci. Rep.* **7**, 46421 (2017).

24. Li, D. & Mashour, G. A. Cortical dynamics during psychedelic and anesthetized states
induced by ketamine. *NeuroImage* **196**, 32–40 (2019).

25. Johnson, M. W., Hendricks, P. S., Barrett, F. S. & Griffiths, R. R. Classic psychedelics:
An integrative review of epidemiology, therapeutics, mystical experience, and brain
network function. *Pharmacol. Ther.* **197**, 83–102 (2019).

26. Carhart-Harris, R. L. & Goodwin, G. M. The Therapeutic Potential of Psychedelic
Drugs: Past, Present, and Future. *Neuropsychopharmacol. Off. Publ. Am. Coll.*
Neuropsychopharmacol. **42**, 2105–2113 (2017).

27. Goulas, A., Changeux, J. P., Wagstyl, K., Amunts, K., Palomero-Gallagher, N. &
Hilgetag, C. C. The natural axis of transmitter receptor distribution in the human
cerebral cortex. *Proc. Natl. Acad. Sci. U. S. A.* **118**, (2021).

28. Zilles, K. & Palomero-Gallagher, N. Multiple transmitter receptors in regions and layers
of the human cerebral cortex. *Front. Neuroanat.* **11**, 78 (2017).

29. Suárez, L. E., Markello, R. D., Betzel, R. F. & Misic, B. Linking Structure and Function
in Macroscale Brain Networks. *Trends Cogn. Sci.* **24**, 302–315 (2020).

30.   Shine, J. M. Neuromodulatory Influences on Integration and Segregation in the Brain.
Trends Cogn. Sci. **23**, 572–583 (2019).

31.   Hansen, J. Y. *et al.* Mapping neurotransmitter systems to the structural and functional
organization of the human neocortex. *Nat. Neurosci.* **(accepted)**, 1–26 (2022).

32.   Trapani, G., Altomare, C., Sanna, E., Biggio, G. & Liso, G. Propofol in Anesthesia.
Mechanism of Action, Structure-Activity Relationships, and Drug Delivery. *Curr. Med.*
Chem. **7**, 249–271 (2000).

33.   Jurd, R. *et al.* General anesthetic actions *in vivo* strongly attenuated by a point mutation
in the GABA $_A$ receptor β3 subunit. *FASEB J.* **17**, 250–252 (2003).

34.   Yip, G. M. S. *et al.* A propofol binding site on mammalian GABA A receptors identified
by photolabeling. *Nat. Chem. Biol.* **9**, 715–720 (2013).

35.   Nishikawa, K. & Harrison, N. L. The Actions of Sevoflurane and Desflurane on the γ-
Aminobutyric Acid Receptor Type A: Effects of TM2 Mutations in the α and β Subunits.
Anesthesiology **99**, 678–684 (2003).

36.   Wu, J., Harata, N. & Akaike, N. Potentiation by sevoflurane of the gamma-aminobutyric
acid-induced chloride current in acutely dissociated CA1 pyramidal neurones from rat
hippocampus. *Br. J. Pharmacol.* **119**, 1013–1021 (1996).

37.   Hirota, K. & Roth, S. H. Sevoflurane modulates both GABAA and GABAB receptors in
area CA1 of rat hippocampus. *Br. J. Anaesth.* **78**, 60–65 (1997).

38.   Jenkins, A., Franks, N. P. & Lieb, W. R. Effects of temperature and volatile anesthetics
on GABA(A) receptors. *Anesthesiology* **90**, 484–491 (1999).

39.   Campagna, J. A., Miller, K. W. & Forman, S. A. Mechanisms of actions of inhaled
anesthetics. *N. Engl. J. Med.* **348**, 2110–2124 (2003).

40.   Domino, E. F. Taming the ketamine tiger. *Anesthesiology* **113**, 678–684 (2010).

41.   Domino, E. F., Chodoff, P. & Corssen, G. Pharmacologic effects of CI-581, a new
dissociative anesthetic, in man. *Clin. Pharmacol. Ther.* **6**, 279–291 (1965).

42.   Krystal, J. H., Karper, L. P., Seibyl, J. P., Freeman, G. K., Delaney, R., Bremner, J. D.,
Heninger, G. R., Bowers, M. B. & Charney, D. S. Subanesthetic Effects of the
Noncompetitive NMDA Antagonist, Ketamine, in Humans: Psychotomimetic,
Perceptual, Cognitive, and Neuroendocrine Responses. *Arch. Gen. Psychiatry* **51**,
199–214 (1994).

43.   Mashour, G. A. Top-down mechanisms of anesthetic-induced unconsciousness. *Front.*
Syst. Neurosci. **8**, (2014).

44.   Olney, J. W., Newcomer, J. W. & Farber, N. B. NMDA receptor hypofunction model of
schizophrenia. *J. Psychiatr. Res.* **33**, 523–533 (1999).

45.   Li, L. & Vlisides, P. E. Ketamine: 50 Years of Modulating the Mind. *Front. Hum.*
Neurosci. **10**, (2016).

46.   Nichols, D. E. Psychedelics. *Pharmacol. Rev.* **68**, 264–355 (2016).

47.   Glennon, R. A., Titeler, M. & McKenney, J. D. Evidence for 5-HT2 involvement in the
mechanism of action of hallucinogenic agents. *Life Sci.* **35**, 2505–2511 (1984).

48.   Nichols, D. E. Dark Classics in Chemical Neuroscience: Lysergic Acid Diethylamide
(LSD). *ACS Chem. Neurosci.* **9**, 2331–2343 (2018).

49.   Nichols, D. E. Hallucinogens. *Pharmacol. Ther.* **101**, 131–181 (2004).

50.   Sleigh, J., Harvey, M., Voss, L. & Denny, B. Ketamine – More mechanisms of action
than just NMDA blockade. *Trends Anaesth. Crit. Care* **4**, 76–81 (2014).

51.   Spindler, L. R. B. *et al.* Dopaminergic brainstem disconnection is common to
pharmacological and pathological consciousness perturbation. *Proc. Natl. Acad. Sci. U.*
S. A. **118**, e2026289118 (2021).

52.   Lawn, T., Dipasquale, O., Vamvakas, A., Tsougos, I., Mehta, M. A. & Howard, M. A.
Differential contributions of serotonergic and dopaminergic functional connectivity to the
phenomenology of LSD. *Psychopharmacology (Berl.)* (2022) doi:10.1007/S00213-022-
06117-5.

53.   Ballentine, G., Friedman, S. F. & Bzdok, D. Trips and neurotransmitters: Discovering
principled patterns across 6850 hallucinogenic experiences. *Sci. Adv.* **8**, eabl6989
(2022).

54.   Sydnor, V. J. *et al.* Neurodevelopment of the association cortices: Patterns,
1978   mechanisms, and implications for psychopathology. *Neuron* (2021)
1979   doi:10.1016/j.neuron.2021.06.016.
1980   55.   Huntenburg, J. M., Bazin, P. L., Goulas, A., Tardif, C. L., Villringer, A. & Margulies, D.
1981   S. A Systematic Relationship Between Functional Connectivity and Intracortical Myelin
1982   in the Human Cerebral Cortex. *Cereb. Cortex* **27**, 981–997 (2017).
1983   56.   Valk, S. L. *et al.* Genetic and phylogenetic uncoupling of structure and function in
1984   human transmodal cortex. *bioRxiv* **5**, 2021.06.08.447522 (2021).
1985   57.   Margulies, D. S. *et al.* Situating the default-mode network along a principal gradient of
1986   macroscale cortical organization. *Proc. Natl. Acad. Sci. U. S. A.* **113**, 12574–12579
1987   (2016).
1988   58.   Burt, J. B., Demirtaş, M., Eckner, W. J., Navejar, N. M., Ji, J. L., Martin, W. J.,
1989   Bernacchia, A., Anticevic, A. & Murray, J. D. Hierarchy of transcriptomic specialization
1990   across human cortex captured by structural neuroimaging topography. *Nat. Neurosci.*
1991   **21**, 1251–1259 (2018).
1992   59.   Hansen, J. Y., Markello, R. D., Vogel, J. W., Seidlitz, J., Bzdok, D. & Misic, B. Mapping
1993   gene transcription and neurocognition across human neocortex. *Nat. Hum. Behav.* 1–
1994   11 (2021) doi:10.1038/s41562-021-01082-z.
1995   60.   Carhart-Harris, R. L. *et al.* Neural correlates of the LSD experience revealed by
1996   multimodal neuroimaging. *Proc. Natl. Acad. Sci.* **113**, 201518377 (2016).
1997   61.   Carhart-Harris, R. L. *et al.* Neural correlates of the psychedelic state as determined by
1998   fMRI studies with psilocybin. *Proc. Natl. Acad. Sci.* **109**, 2138–2143 (2012).
1999   62.   Viol, A., Palhano-Fontes, F., Onias, H., De Araujo, D. B. & Viswanathan, G. M.
2000   Shannon entropy of brain functional complex networks under the influence of the
2001   psychedelic Ayahuasca. *Sci. Rep.* **7**, (2017).
2002   63.   Carhart-Harris, R. L. *et al.* The Effects of Acutely Administered 3,4-
2003   Methylenedioxymethamphetamine on Spontaneous Brain Function in Healthy
2004   Volunteers Measured with Arterial Spin Labeling and Blood Oxygen Level-Dependent
2005   Resting State Functional Connectivity. *Biol. Psychiatry* **78**, 554–562 (2015).
2006   64.   Dandash, O., Harrison, B. J., Adapa, R., Gaillard, R., Giorlando, F., Wood, S. J.,
2007   Fletcher, P. C. & Fornito, A. Selective Augmentation of Striatal Functional Connectivity
2008   Following NMDA Receptor Antagonism: Implications for Psychosis.
2009   *Neuropsychopharmacology* **40**, 622–631 (2015).
2010   65.   Esposito, R. *et al.* Acute Effects of Modafinil on Brain Resting State Networks in Young
2011   Healthy Subjects. *PLoS ONE* **8**, (2013).
2012   66.   Ranft, A. *et al.* Neural Correlates of Sevoflurane-induced Unconsciousness Identified
2013   by Simultaneous Functional Magnetic Resonance Imaging and
2014   Electroencephalography. *Anesthesiology* **125**, 861–872 (2016).
2015   67.   Stamatakis, E. A., Adapa, R. M., Absalom, A. R. & Menon, D. K. Changes in resting
2016   neural connectivity during propofol sedation. *PloS One* **5**, e14224 (2010).
2017   68.   Naci, L., Sinai, L. & Owen, A. M. Detecting and interpreting conscious experiences in
2018   behaviorally non-responsive patients. *NeuroImage* **145**, 304–313 (2017).
2019   69.   Dipasquale, O. *et al.* Unravelling the effects of methylphenidate on the dopaminergic
2020   and noradrenergic functional circuits. doi:10.1038/s41386-020-0724-x.
2021   70.   Dipasquale, O., Selvaggi, P., Veronese, M., Gabay, A. S., Turkheimer, F. & Mehta, M.
2022   A. Receptor-Enriched Analysis of functional connectivity by targets (REACT): A novel,
2023   multimodal analytical approach informed by PET to study the pharmacodynamic
2024   response of the brain under MDMA. (2019) doi:10.1016/j.neuroimage.2019.04.007.
2025   71.   Kringelbach, M. L., Cruzat, J., Cabral, J., Knudsen, G. M., Carhart-Harris, R., Whybrow,
2026   P. C., Logothetis, N. K. & Deco, G. Dynamic coupling of whole-brain neuronal and
2027   neurotransmitter systems. *Proc. Natl. Acad. Sci. U. S. A.* **117**, 9566–9576 (2020).
2028   72.   Deco, G., Cruzat, J., Cabral, J., Whybrow, P. C., Logothetis, N. K. & Kringelbach, M. L.
2029   Whole-Brain Multimodal Neuroimaging Model Using Serotonin Receptor Maps Explains
2030   Non-linear Functional Effects of LSD. *Curr. Biol.* **28**, 3065–3074 (2018).

73. Burt, J., Preller, K., Demirtaş, M., Ji, J. L., Krystal, J., Vollenweider, F., Anticevic, A. &
Murray, J. Transcriptomics-informed large-scale cortical model captures topography of
pharmacological neuroimaging effects of LSD. *eLife* **10**, (2021).
74. Parker Singleton, S. A., Luppi B, A. I., Carhart-Harris D, R. L., Cruzat, J. F., Roseman,
L. D., Deco, G. G., Kringelbach K, M. L., Stamatakis B, E. A. & Kuceyeski, A. A.
Psychedelics Flatten the brain's energy landscape: evidence from receptor-informed
network control theory. *bioRxiv* 2021.05.14.444193 (2021)
doi:10.1101/2021.05.14.444193.
75. Bonhomme, V. *et al.* Resting-state Network-specific Breakdown of Functional
Connectivity during Ketamine Alteration of Consciousness in Volunteers.
Anesthesiology **125**, 873–888 (2016).
76. Lee, U., Ku, S., Noh, G., Baek, S., Choi, B. & Mashour, G. A. Disruption of frontal-
parietal communication by ketamine, propofol, and sevoflurane. *Anesthesiology* **118**,
1264–75 (2013).
77. Kaller, S. *et al.* Test–retest measurements of dopamine D1-type receptors using
simultaneous PET/MRI imaging. *Eur. J. Nucl. Med. Mol. Imaging* **44**, (2017).
78. Sandiego, C. M., Gallezot, J.-D., Lim, K., Ropchan, J., Lin, S., Gao, H., Morris, E. D. &
Cosgrove, K. P. Reference region modeling approaches for amphetamine challenge
studies with [11C]FLB 457 and PET. *J. Cereb. Blood Flow Metab. Off. J. Int. Soc.*
Cereb. Blood Flow Metab. **35**, 623–629 (2015).
79. Slifstein, M. *et al.* Deficits in prefrontal cortical and extrastriatal dopamine release in
schizophrenia: a positron emission tomographic functional magnetic resonance imaging
study. *JAMA Psychiatry* **72**, 316–324 (2015).
80. Smith, C. T. *et al.* Partial-volume correction increases estimated dopamine D2-like
receptor binding potential and reduces adult age differences. *J. Cereb. Blood Flow*
Metab. Off. J. Int. Soc. Cereb. Blood Flow Metab. **39**, 822–833 (2019).
81. Zakiniaeiz, Y. *et al.* Sex differences in amphetamine-induced dopamine release in the
dorsolateral prefrontal cortex of tobacco smokers. *Neuropsychopharmacol. Off. Publ.*
Am. Coll. Neuropsychopharmacol. **44**, 2205–2211 (2019).
82. Dukart, J. *et al.* Cerebral blood flow predicts differential neurotransmitter activity. *Sci.*
Rep. **8**, 4074 (2018).
83. Sanchez-Rangel, E., Gallezot, J.-D., Yeckel, C. W., Lam, W., Belfort-DeAguiar, R.,
Chen, M.-K., Carson, R. E., Sherwin, R. & Hwang, J. J. Norepinephrine transporter
availability in brown fat is reduced in obesity: a human PET study with [11C] MRB. *Int.*
J. Obes. **44**, 964–967 (2020).
84. Ding, Y.-S. *et al.* PET imaging of the effects of age and cocaine on the norepinephrine
transporter in the human brain using (S,S)-[(11)C]O-methylreboxetine and HRRT.
Synap. N. Y. N **64**, 30–38 (2010).
85. Belfort-DeAguiar, R., Gallezot, J.-D., Hwang, J. J., Elshafie, A., Yeckel, C. W., Chan,
O., Carson, R. E., Ding, Y.-S. & Sherwin, R. S. Noradrenergic Activity in the Human
Brain: A Mechanism Supporting the Defense Against Hypoglycemia. *J. Clin.*
Endocrinol. Metab. **103**, 2244–2252 (2018).
86. Li, C. R. *et al.* Decreased norepinephrine transporter availability in obesity: Positron
Emission Tomography imaging with (S,S)-[(11)C]O-methylreboxetine. *NeuroImage* **86**,
306–310 (2014).
87. Savli, M. *et al.* Normative database of the serotonergic system in healthy subjects using
multi-tracer PET. *NeuroImage* **63**, 447–459 (2012).
88. Gallezot, J.-D. *et al.* Kinetic modeling of the serotonin 5-HT(1B) receptor radioligand
[(11)C]P943 in humans. *J. Cereb. Blood Flow Metab. Off. J. Int. Soc. Cereb. Blood*
Flow Metab. **30**, 196–210 (2010).
89. Matuskey, D. *et al.* Reductions in brain 5-HT1B receptor availability in primarily
cocaine-dependent humans. *Biol. Psychiatry* **76**, 816–822 (2014).
90. Murrough, J. W. *et al.* The effect of early trauma exposure on serotonin type 1B
receptor expression revealed by reduced selective radioligand binding. *Arch. Gen.*
Psychiatry **68**, 892–900 (2011).

91. Pittenger, C. *et al.* OCD is associated with an altered association between sensorimotor
gating and cortical and subcortical 5-HT1b receptor binding. *J. Affect. Disord.* **196**, 87–
96 (2016).
92. Saricicek, A. *et al.* Test-retest reliability of the novel 5-HT1B receptor PET radioligand
[11C]P943. *Eur. J. Nucl. Med. Mol. Imaging* **42**, 468–477 (2015).
93. Beliveau, V., Ganz, M., Feng, L., Ozenne, B., Højgaard, L., Fisher, P. M., Svarer, C.,
Greve, D. N. & Knudsen, G. M. A High-Resolution In Vivo Atlas of the Human Brain's
Serotonin System. *J. Neurosci.* **37**, 120 (2017).
94. Radhakrishnan, R. *et al.* In vivo 5-HT6 and 5-HT2A receptor availability in antipsychotic
treated schizophrenia patients vs. unmedicated healthy humans measured with
[11C]GSK215083 PET. *Psychiatry Res. Neuroimaging* **295**, 111007 (2020).
95. Radhakrishnan, R. *et al.* Age-Related Change in 5-HT6 Receptor Availability in Healthy
Male Volunteers Measured with 11C-GSK215083 PET. *J. Nucl. Med.* **59**, 1445–1450
(2018).
96. Hillmer, A. T. *et al.* Imaging of cerebral α4β2* nicotinic acetylcholine receptors with (-)-
[(18)F]Flubatine PET: Implementation of bolus plus constant infusion and sensitivity to
acetylcholine in human brain. *NeuroImage* **141**, 71–80 (2016).
97. Baldassarri, S. R. *et al.* Use of Electronic Cigarettes Leads to Significant Beta2-
Nicotinic Acetylcholine Receptor Occupancy: Evidence From a PET Imaging Study.
Nicotine Tob. Res. Off. J. Soc. Res. Nicotine Tob. **20**, 425–433 (2018).
98. Naganawa, M. *et al.* First-in-Human Assessment of 11C-LSN3172176, an M1
Muscarinic Acetylcholine Receptor PET Radiotracer. *J. Nucl. Med. Off. Publ. Soc. Nucl.*
Med. **62**, 553–560 (2021).
99. Bedard, M.-A., Aghourian, M., Legault-Denis, C., Postuma, R. B., Soucy, J.-P.,
Gagnon, J.-F., Pelletier, A. & Montplaisir, J. Brain cholinergic alterations in idiopathic
REM sleep behaviour disorder: a PET imaging study with 18F-FEOBV. *Sleep Med.* **58**,
35–41 (2019).
100. Aghourian, M., Legault-Denis, C., Soucy, J.-P., Rosa-Neto, P., Gauthier, S., Kostikov,
A., Gravel, P. & Bédard, M.-A. Quantification of brain cholinergic denervation in
Alzheimer's disease using PET imaging with [18F]-FEOBV. *Mol. Psychiatry* **22**, 1531–
1538 (2017).
101. Smart, K., Cox, S. M. L., Scala, S. G., Tippler, M., Jaworska, N., Boivin, M., Séguin, J.
R., Benkelfat, C. & Leyton, M. Sex differences in [11C]ABP688 binding: a positron
emission tomography study of mGlu5 receptors. *Eur. J. Nucl. Med. Mol. Imaging* **46**,
1179–1183 (2019).
102. DuBois, J. M. *et al.* Characterization of age/sex and the regional distribution of mGluR5
availability in the healthy human brain measured by high-resolution [(11)C]ABP688
PET. *Eur. J. Nucl. Med. Mol. Imaging* **43**, 152–162 (2016).
103. Galovic, M. *et al.* In vivo NMDA receptor function in people with NMDA receptor
antibody encephalitis. 2021.12.04.21267226 Preprint at
https://doi.org/10.1101/2021.12.04.21267226 (2021).
104. Galovic, M. *et al.* Validation of a combined image derived input function and venous
sampling approach for the quantification of [18F]GE-179 PET binding in the brain.
NeuroImage **237**, 118194 (2021).
105. Nørgaard, M., Beliveau, V., Ganz, M., Svarer, C., Pinborg, L., Keller, S., Jensen, P.,
Greve, D. & Knudsen, G. A High-Resolution In Vivo Atlas of the Human Brain's
Benzodiazepine Binding Site of GABA A Receptors. *NeuroImage* **232**, 117878 (2021).
106. Gallezot, J.-D. *et al.* Determination of receptor occupancy in the presence of mass
dose: [11C]GSK189254 PET imaging of histamine H3 receptor occupancy by PF-
03654746. *J. Cereb. Blood Flow Metab. Off. J. Int. Soc. Cereb. Blood Flow Metab.* **37**,
1095–1107 (2017).
107. D'Souza, D. C. *et al.* Rapid Changes in CB1 Receptor Availability in Cannabis
Dependent Males after Abstinence from Cannabis. *Biol. Psychiatry Cogn. Neurosci.*
Neuroimaging **1**, 60–67 (2016).

108. Neumeister, A. *et al.* Positron emission tomography shows elevated cannabinoid CB1
receptor binding in men with alcohol dependence. *Alcohol. Clin. Exp. Res.* **36**, 2104–
2109 (2012).
109. Normandin, M. D. *et al.* Imaging the cannabinoid CB1 receptor in humans with
[11C]OMAR: assessment of kinetic analysis methods, test-retest reproducibility, and
gender differences. *J. Cereb. Blood Flow Metab. Off. J. Int. Soc. Cereb. Blood Flow
Metab.* **35**, 1313–1322 (2015).
110. Ranganathan, M. *et al.* Reduced Brain Cannabinoid Receptor Availability in
Schizophrenia. *Biol. Psychiatry* **79**, 997–1005 (2016).
111. Kantonen, T. *et al.* Interindividual variability and lateralization of µ-opioid receptors in
the human brain. *NeuroImage* **217**, 116922 (2020).
112. Schaefer, A., Kong, R., Gordon, E. M., Laumann, T. O., Zuo, X.-N., Holmes, A. J.,
Eickhoff, S. B. & Yeo, B. T. T. Local-Global Parcellation of the Human Cerebral Cortex
from Intrinsic Functional Connectivity MRI. *Cereb. Cortex* **28**, 3095–3114 (2018).
113. Hansen, J. Y. *et al.* Molecular and connectomic vulnerability shape cross-disorder
cortical abnormalities. *bioRxiv* 2022.01.21.476409 (2022)
doi:10.1101/2022.01.21.476409.
114. Betzel, R. F. & Bassett, D. S. Specificity and robustness of long-distance connections in
weighted, interareal connectomes. *Proc Nati Acad Sci USA* (2018)
doi:10.1073/pnas.1720186115.
115. Roberts, J. A., Perry, A., Lord, A. R., Roberts, G., Mitchell, P. B., Smith, R. E.,
Calamante, F. & Breakspear, M. The contribution of geometry to the human
connectome. *NeuroImage* **124**, 379–393 (2016).
116. Stiso, J. & Bassett, D. S. Spatial Embedding Imposes Constraints on Neuronal Network
Architectures. *Trends Cogn. Sci.* **22**, 1127–1142 (2018).
117. Wold, S., Josefson, M., Gottfries, J. & Linusson, A. The utility of multivariate design in
PLS modeling. *J. Chemom.* **18**, 156–165 (2004).
118. McIntosh, A. R. & Lobaugh, N. J. Partial least squares analysis of neuroimaging data:
applications and advances. *NeuroImage* **23 Suppl 1**, S250-263 (2004).
119. Krishnan, A., Williams, L. J., McIntosh, A. R. & Abdi, H. Partial Least Squares (PLS)
methods for neuroimaging: A tutorial and review. *NeuroImage* **56**, 455–475 (2011).
120. McIntosh, A. R. & Mišić, B. Multivariate Statistical Analyses for Neuroimaging Data.
Annu. Rev. Psychol. **64**, 499–525 (2013).
121. Markello, R. D. & Misic, B. Comparing spatial null models for brain maps. *NeuroImage*
236, 118052 (2021).
122. Váša, F. & Miši´c, B. M. *Null Models in Network Neuroscience.*
123. Váša, F. *et al.* Adolescent tuning of association cortex in human structural brain
networks. *Cereb. Cortex* **28**, 281–294 (2018).
124. Alexander-Bloch, A. F., Shou, H., Liu, S., Satterthwaite, T. D., Glahn, D. C., Shinohara,
R. T., Vandekar, S. N. & Raznahan, A. On testing for spatial correspondence between
maps of human brain structure and function. *NeuroImage* **178**, 540–551 (2018).
125. Yeo, B. T. T. *et al.* The organization of the human cerebral cortex estimated by intrinsic
functional connectivity. *J. Neurophysiol.* **106**, 1125–1165 (2011).
126. Hill, J., Inder, T., Neil, J., Dierker, D., Harwell, J. & Van Essen, D. Similar patterns of
cortical expansion during human development and evolution. *Proc. Natl. Acad. Sci.*
107, 13135–13140 (2010).
127. Hawrylycz, M. J. *et al.* An anatomically comprehensive atlas of the adult human brain
transcriptome. *Nature* **489**, 391–399 (2012).
128. Yarkoni, T., Poldrack, R. A., Nichols, T. E., Van Essen, D. C. & Wager, T. D. Large-
scale automated synthesis of human functional neuroimaging data. *Nat. Methods* **8**,
665–670 (2011).
129. Luppi, A. I. *et al.* A synergistic core for human brain evolution and cognition. *Nat.
Neurosci.* **25**, 771–782 (2022).
130. Larivière, S. *et al.* The ENIGMA Toolbox: multiscale neural contextualization of multisite
neuroimaging datasets. *Nat. Methods 2021 187* **18**, 698–700 (2021).

131. Sun, D. *et al.* Large-scale mapping of cortical alterations in 22q11.2 deletion syndrome: Convergence with idiopathic psychosis and effects of deletion size. *Mol. Psychiatry* **25**, 1822–1834 (2020).

132. Hoogman, M. *et al.* Brain Imaging of the Cortex in ADHD: A Coordinated Analysis of Large-Scale Clinical and Population-Based Samples. *Am. J. Psychiatry* **176**, 531–542 (2019).

133. van Rooij, D. *et al.* Cortical and Subcortical Brain Morphometry Differences Between Patients With Autism Spectrum Disorder and Healthy Individuals Across the Lifespan: Results From the ENIGMA ASD Working Group. *Am. J. Psychiatry* **175**, 359–369 (2018).

134. Whelan, C. D. *et al.* Structural brain abnormalities in the common epilepsies assessed in a worldwide ENIGMA study. *Brain J. Neurol.* **141**, 391–408 (2018).

135. Schmaal, L. *et al.* Cortical abnormalities in adults and adolescents with major depression based on brain scans from 20 cohorts worldwide in the ENIGMA Major Depressive Disorder Working Group. *Mol. Psychiatry* **22**, 900–909 (2017).

136. Boedhoe, P. S. W. *et al.* Cortical Abnormalities Associated With Pediatric and Adult Obsessive-Compulsive Disorder: Findings From the ENIGMA Obsessive-Compulsive Disorder Working Group. *Am. J. Psychiatry* **175**, 453–462 (2018).

137. van Erp, T. G. M. *et al.* Cortical Brain Abnormalities in 4474 Individuals With Schizophrenia and 5098 Control Subjects via the Enhancing Neuro Imaging Genetics Through Meta Analysis (ENIGMA) Consortium. *Biol. Psychiatry* **84**, 644–654 (2018).

138. Hibar, D. P. *et al.* Cortical abnormalities in bipolar disorder: an MRI analysis of 6503 individuals from the ENIGMA Bipolar Disorder Working Group. *Mol. Psychiatry* **23**, 932–942 (2018).

139. Laansma, M. A. *et al.* International Multicenter Analysis of Brain Structure Across Clinical Stages of Parkinson's Disease. *Mov. Disord.* **36**, 2583–2594 (2021).

140. Vos de Wael, R. *et al.* BrainSpace: a toolbox for the analysis of macroscale gradients in neuroimaging and connectomics datasets. *Commun. Biol.* **3**, (2020).

141. Paquola, C. *et al.* A multi-scale cortical wiring space links cellular architecture and functional dynamics in the human brain. *PLOS Biol.* **18**, e3000979 (2020).

142. Cammoun, L., Gigandet, X., Meskaldji, D., Thiran, J. P., Sporns, O., Do, K. Q., Maeder, P., Meuli, R. & Hagmann, P. Mapping the human connectome at multiple scales with diffusion spectrum MRI. *J. Neurosci. Methods* **203**, 386–397 (2012).

143. Holze, F., Vizeli, P., Müller, F., Ley, L., Duerig, R., Varghese, N., Eckert, A., Borgwardt, S. & Liechti, M. E. ARTICLE Distinct acute effects of LSD, MDMA, and D-amphetamine in healthy subjects. doi:10.1038/s41386-019-0569-3.

144. Viol, A., Palhano-Fontes, F., Onias, H., de Araujo, D. B., Hövel, P. & Viswanathan, G. M. Characterizing complex networks using entropy-degree diagrams: Unveiling changes in functional brain connectivity induced by Ayahuasca. *Entropy* **21**, (2019).

145. Luppi, A. I., Carhart-Harris, R. L., Roseman, L., Pappas, I., Menon, D. K. & Stamatakis, E. A. LSD alters dynamic integration and segregation in the human brain. *NeuroImage* **227**, 117653 (2021).

146. Atasoy, S., Roseman, L., Kaelen, M., Kringelbach, M. L., Deco, G. & Carhart-Harris, R. L. Connectome-harmonic decomposition of human brain activity reveals dynamical repertoire re-organization under LSD. *Sci. Rep.* **7**, 1–18 (2017).

147. Lord, L. D. *et al.* Dynamical exploration of the repertoire of brain networks at rest is modulated by psilocybin. *NeuroImage* **199**, 127–142 (2019).

148. Schartner, M., Seth, A., Noirhomme, Q., Boly, M., Bruno, M. A., Laureys, S. & Barrett, A. Complexity of multi-dimensional spontaneous EEG decreases during propofol induced general anaesthesia. *PLoS ONE* **10**, (2015).

149. Toker, D. *et al.* Consciousness is supported by near-critical cortical electrodynamics. doi:10.1073/pnas.XXXXXXXXX.

150. Carhart-Harris, R. L. & Friston, K. J. REBUS and the anarchic brain: Toward a unified model of the brain action of psychedelics. *Pharmacol. Rev.* **71**, 316–344 (2019).

151. Varley, T. F., Luppi, A. I., Pappas, I., Naci, L., Adapa, R., Owen, A. M., Menon, D. K. & Stamatakis, E. A. Consciousness & Brain Functional Complexity in Propofol Anaesthesia. *Sci. Rep.* **10**, 1–13 (2020).

152. Varley, T. F., Craig, M. M., Adapa, R., Finoia, P., Williams, G., Allanson, J., Pickard, J., Menon, D. K. & Stamatakis, E. A. Fractal dimension of cortical functional connectivity networks & severity of disorders of consciousness. *PLoS ONE* **15**, 1–20 (2020).

153. Varley, T. F., Carhart-Harris, R., Roseman, L., Menon, D. K. & Stamatakis, E. A. Serotonergic psychedelics LSD & psilocybin increase the fractal dimension of cortical brain activity in spatial and temporal domains. *NeuroImage* **220**, (2020).

154. Lebedev, A. V., Kaelen, M., Lövdén, M., Nilsson, J., Feilding, A., Nutt, D. J. & Carhart-Harris, R. L. LSD-induced entropic brain activity predicts subsequent personality change. *Hum. Brain Mapp.* **37**, 3203–3213 (2016).

155. Boveroux, P., Vanhaudenhuyse, A. & Phillips, C. Breakdown of within- and between-network Resting State during Propofol-induced Loss of Consciousness. *Anesthesiology* **113**, 1038–1053 (2010).

156. Wu, H. *et al.* Anterior precuneus related to the recovery of consciousness. *NeuroImage Clin.* **33**, 102951 (2022).

157. Palhano-Fontes, F., Andrade, K. C., Tofoli, L. F., Jose, A. C. S., Crippa, A. S., Hallak, J. E. C., Ribeiro, S. & De Araujo, D. B. The psychedelic state induced by Ayahuasca modulates the activity and connectivity of the Default Mode Network. *PLoS ONE* **10**, (2015).

158. Girn, M., Roseman, L., Bernhardt, B., Smallwood, J., Carhart-Harris, R. & Spreng, R. N. LSD flattens the functional hierarchy of the human brain. *bioRxiv* 2020.05.01.072314 (2020) doi:10.1101/2020.05.01.072314.

159. Demirtaş, M. *et al.* Hierarchical Heterogeneity across Human Cortex Shapes Large-Scale Neural Dynamics. *Neuron* **101**, 1181-1194.e13 (2019).

160. Beul, S. F. & Hilgetag, C. C. Neuron density fundamentally relates to architecture and connectivity of the primate cerebral cortex. *NeuroImage* **189**, 777–792 (2019).

161. Baum, G. L. *et al.* Development of structure–function coupling in human brain networks during youth. *Proc. Natl. Acad. Sci. U. S. A.* **117**, 771–778 (2020).

162. Doss, M. K., May, D. G., Johnson, M. W., Clifton, J. M., Hedrick, S. L., Prisinzano, T. E., Griffiths, R. R. & Barrett, F. S. The Acute Effects of the Atypical Dissociative Hallucinogen Salvinorin A on Functional Connectivity in the Human Brain. *Sci. Rep.* **10**, 16392 (123AD).

163. Akeju, O. *et al.* Disruption of thalamic functional connectivity is a neural correlate of dexmedetomidine-induced unconsciousness. *eLife* **3**, (2014).

164. Guldenmund, P. *et al.* Brain functional connectivity differentiates dexmedetomidine from propofol and natural sleep. *BJA Br. J. Anaesth.* **119**, 674–684 (2017).

165. Nelson, L. E., Lu, J., Guo, T., Saper, C. B., Franks, N. P. & Maze, M. The α2-adrenoceptor agonist dexmedetomidine converges on an endogenous sleep-promoting pathway to exert its sedative effects. *Anesthesiology* **98**, 428–436 (2003).

166. Van de Bittner, G. C., Ricq, E. L. & Hooker, J. M. A philosophy for CNS radiotracer design. *Acc. Chem. Res.* **47**, 3127–3134 (2014).

167. Elsinga, P. H., Hatano, K. & Ishiwata, K. PET tracers for imaging of the dopaminergic system. *Curr. Med. Chem.* **13**, 2139–2153 (2006).

168. Redinbaugh, M. J., Phillips, J. M., Kambi, N. A., Mohanta, S., Andryk, S., Dooley, G. L., Afrasiabi, M., Raz, A. & Saalmann, Y. B. Thalamus modulates consciousness via layer-specific control of cortex. *Neuron* **106**, 66-75.e12 (2020).

169. Tasserie, J., Uhrig, L., Sitt, J. D., Manasova, D., Dupont, M., Dehaene, S. & Jarraya, B. Deep brain stimulation of the thalamus restores signatures of consciousness in a nonhuman primate model. *Sci Adv* **8**, 5547 (2022).

170. Barttfeld, P., Bekinschtein, T. A., Salles, A., Stamatakis, E. A., Adapa, R., Menon, D. K. & Sigman, M. Factoring the brain signatures of anesthesia concentration and level of arousal across individuals. *NeuroImage Clin.* **9**, 385–391 (2015).

171. Preller, K. H. *et al.* Changes in global and thalamic brain connectivity in LSD-induced altered states of consciousness are attributable to the 5-HT2A receptor. *eLife* **7**, (2018).

172. Preller, K. H., Razi, A., Zeidman, P., Stämpfli, P., Friston, K. J. & Vollenweider, F. X. Effective connectivity changes in LSD-induced altered states of consciousness in humans. *Proc. Natl. Acad. Sci. U. S. A.* **116**, 2743–2748 (2019).

173. Shine, J. M. The thalamus integrates the macrosystems of the brain to facilitate complex, adaptive brain network dynamics. *Prog. Neurobiol.* 101951 (2020) doi:10.1016/j.pneurobio.2020.101951.

174. Kringelbach, M. L. & Deco, G. Brain States and Transitions: Insights from Computational Neuroscience. *Cell Rep.* **32**, 108128 (2020).

175. Cofré, R., Herzog, R., Mediano, P. A. M., Piccinini, J., Rosas, F. E., Perl, Y. S. & Tagliazucchi, E. Whole-brain models to explore altered states of consciousness from the bottom up. *Brain Sci.* **10**, 1–29 (2020).

176. Deco, G. & Kringelbach, M. L. Great expectations: Using whole-brain computational connectomics for understanding neuropsychiatric disorders. *Neuron* **84**, 892–905 (2014).

177. Shine, J. M., Müller, E. J., Munn, B., Cabral, J., Moran, R. J. & Breakspear, M. Computational models link cellular mechanisms of neuromodulation to large-scale neural dynamics. *Nat. Neurosci.* 1–12 (2021) doi:10.1038/s41593-021-00824-6.

178. Herzog, R., Mediano, P. A. M., Rosas, F. E., Carhart-Harris, R., Perl, Y. S., Tagliazucchi, E. & Cofre, R. A mechanistic model of the neural entropy increase elicited by psychedelic drugs. *Sci. Rep.* **10**, 17725 (2020).

179. Luppi, A. I. *et al.* Whole-brain modelling identifies distinct but convergent paths to unconsciousness in anaesthesia and disorders of consciousness. *Commun. Biol.* **5**, 384 (2022).

180. Veselis, R. A., Feshchenko, V. A., Reinsel, R. A., Beattie, B. & Akhurst, T. J. Propofol and thiopental do not interfere with regional cerebral blood flow response at sedative concentrations. *Anesthesiology* **102**, 26–34 (2005).

181. Johnston, A. J., Steiner, L. A., Chatfield, D. A., Coleman, M. R., Coles, J. P., Al-Rawi, P. G., Menon, D. K. & Gupta, A. K. Effects of propofol on cerebral oxygenation and metabolism after head injury. *Br. J. Anaesth.* **91**, 781–786 (2003).

182. Hemmings, H. C., Akabas, M. H., Goldstein, P. A., Trudell, J. R., Orser, B. A. & Harrison, N. L. Emerging molecular mechanisms of general anesthetic action. *Trends Pharmacol. Sci.* **26**, 503–510 (2005).

183. Bademosi, A. T. *et al.* Trapping of Syntaxin1a in Presynaptic Nanoclusters by a Clinically Relevant General Anesthetic. *Cell Rep.* **22**, 427–440 (2018).

184. Wang, Y., Yu, T., Yuan, C., Yuan, J., Luo, Z., Pan, Y., Zhang, Y., Zhang, Y. & Yu, B. *Effects of propofol on the dopamine, metabolites and GABAA receptors in media prefrontal cortex in freely moving rats. Am J Transl Res* vol. 8 2301–2308 www.ajtr.org/ISSN:1943-8141/AJTR0024580 (2016).

185. Guo, J., Xu, K., Yin, J., Zhang, H., Yin, J. & Li, Y. Dopamine transporter in the ventral tegmental area modulates recovery from propofol anesthesia in rats. *J. Chem. Neuroanat.* **121**, 102083 (2022).

186. Naci, L. *et al.* Functional diversity of brain networks supports consciousness and verbal intelligence. *Sci. Rep.* **8**, 1–15 (2018).

187. Coppola, P. *et al.* Network dynamics scale with levels of awareness. *NeuroImage* 119128 (2022) doi:10.1016/J.NEUROIMAGE.2022.119128.

188. Kandeepan, S., Rudas, J., Gomez, F., Stojanoski, B., Valluri, S., Owen, A. M., Naci, L., Nichols, E. S. & Soddu, A. Modeling an auditory stimulated brain under altered states of consciousness using the generalized ising model. *NeuroImage* **223**, 117367 (2020).

189. Solt, K., Eger, E. I. & Raines, D. E. Differential modulation of human N-methyl-D-aspartate receptors by structurally diverse general anesthetics. *Anesth. Analg.* **102**, 1407–1411 (2006).

190. Hollmann, M. W., Liu, H. T., Hoenemann, C. W., Liu, W. H. & Durieux, M. E. Modulation of NMDA receptor function by ketamine and magnesium. Part II: interactions with volatile anesthetics. *Anesth. Analg.* **92**, 1182–1191 (2001).

191. Scheller, M., Bufler, J., Schneck, H., Kochs, E. & Franke, C. Isoflurane and sevoflurane interact with the nicotinic acetylcholine receptor channels in micromolar concentrations. *Anesthesiology* **86**, 118–127 (1997).

192. Alkire, M. T., McReynolds, J. R., Hahn, E. L. & Trivedi, A. N. Thalamic microinjection of nicotine reverses sevoflurane-induced loss of righting reflex in the rat. *Anesthesiology* **107**, 264–272 (2007).

193. Palanca, B. J. A., Avidan, M. S. & Mashour, G. A. Human neural correlates of sevoflurane-induced unconsciousness. *BJA Br. J. Anaesth.* **119**, 573–582 (2017).

194. Golkowski, D. *et al.* Coherence of BOLD signal and electrical activity in the human brain during deep sevoflurane anesthesia. *Brain Behav.* **7**, 1–8 (2017).

195. Golkowski, D. *et al.* Changes in Whole Brain Dynamics and Connectivity Patterns during Sevoflurane- and Propofol-induced Unconsciousness Identified by Functional Magnetic Resonance Imaging. *Anesthesiology* **130**, 898–911 (2019).

196. Luppi, A. I., Golkowski, D., Ranft, A., Ilg, R., Jordan, D., Menon, D. K. & Stamatakis, E. A. Brain network integration dynamics are associated with loss and recovery of consciousness induced by sevoflurane. *Hum. Brain Mapp.* (2021) doi:10.1002/hbm.25405.

197. Basso, L., Bönke, L., Aust, S., Gärtner, M., Heuser-Collier, I., Otte, C., Wingenfeld, K., Bajbouj, M. & Grimm, S. Antidepressant and neurocognitive effects of serial ketamine administration versus ECT in depressed patients. *J. Psychiatr. Res.* **123**, 1–8 (2020).

198. Phillips, J. L., Norris, S., Talbot, J., Hatchard, T., Ortiz, A., Birmingham, M., Owoeye, O., Batten, L. A. & Blier, P. Single and repeated ketamine infusions for reduction of suicidal ideation in treatment-resistant depression. *Neuropsychopharmacol. Off. Publ. Am. Coll. Neuropsychopharmacol.* **45**, 606–612 (2020).

199. Corlett, P. R., Honey, G. D. & Fletcher, P. C. Prediction error, ketamine and psychosis: An updated model. *J. Psychopharmacol. (Oxf.)* **30**, 1145–1155 (2016).

200. Moore, J. W. *et al.* Ketamine administration in healthy volunteers reproduces aberrant agency experiences associated with schizophrenia. *Cognit. Neuropsychiatry* **16**, 364–381 (2011).

201. Studerus, E., Gamma, A. & Vollenweider, F. X. Psychometric evaluation of the altered states of consciousness rating scale (OAV). *PLoS ONE* **5**, (2010).

202. Corlett, P. R. *et al.* Frontal responses during learning predict vulnerability to the psychotogenic effects of ketamine: Linking cognition, brain activity, and psychosis. *Arch. Gen. Psychiatry* **63**, 611–621 (2006).

203. Marona-Lewicka, D., Kurrasch-Orbaugh, D. M., Selken, J. R., Cumbay, M. G., Lisnicchia, J. G. & Nichols, D. E. Re-evaluation of lisuride pharmacology: 5-hydroxytryptamine1A receptor-mediated behavioral effects overlap its other properties in rats. *Psychopharmacology (Berl.)* **164**, 93–107 (2002).

204. Passie, T., Halpern, J. H., Stichtenoth, D. O., Emrich, H. M. & Hintzen, A. The pharmacology of lysergic acid diethylamide: A review. *CNS Neurosci. Ther.* **14**, 295–314 (2008).

205. Kraehenmann, R., Pokorny, D., Vollenweider, L., Preller, K. H., Pokorny, T., Seifritz, E. & Vollenweider, F. X. Dreamlike effects of LSD on waking imagery in humans depend on serotonin 2A receptor activation. *Psychopharmacology (Berl.)* 1–16 (2017) doi:10.1007/s00213-017-4610-0.

206. Tagliazucchi, E. *et al.* Increased Global Functional Connectivity Correlates with LSD-Induced Ego Dissolution. *Curr. Biol.* **26**, 1043–1050 (2016).

207. Vollenweider, F. X., Vollenweider-Scherpenhuyzen, M. F., Bäbler, A., Vogel, H. & Hell, D. Psilocybin induces schizophrenia-like psychosis in humans via a serotonin-2 agonist action. *Neuroreport* **9**, 3897–3902 (1998).

208. Passie, T., Seifert, J., Schneider, U. & Emrich, H. M. The pharmacology of psilocybin. *Addict. Biol.* **7**, 357–364 (2002).

209. González-Maeso, J. *et al.* Hallucinogens recruit specific cortical 5-HT(2A) receptor-
mediated signaling pathways to affect behavior. *Neuron* **53**, 439–452 (2007).
210. McKenna, D. J., Repke, D. B., Lo, L. & Peroutka, S. J. Differential interactions of
indolealkylamines with 5-hydroxytryptamine receptor subtypes. *Neuropharmacology* **29**,
193–198 (1990).
211. Halberstadt, A. L., Koedood, L., Powell, S. B. & Geyer, M. A. Differential contributions
of serotonin receptors to the behavioral effects of indoleamine hallucinogens in mice. *J.*
Psychopharmacol. Oxf. Engl. **25**, 1548–1561 (2011).
212. Christian, S. T., Harrison, R., Quayle, E., Pagel, J. & Monti, J. The in vitro identification
of dimethyltryptamine (DMT) in mammalian brain and its characterization as a possible
endogenous neuroregulatory agent. *Biochem. Med.* **18**, 164–183 (1977).
213. Smythies, J. R., Morin, R. D. & Brown, G. B. Identification of dimethyltryptamine and O-
methylbufotenin in human cerebrospinal fluid by combined gas chromatography/mass
spectrometry. *Biol. Psychiatry* **14**, 549–556 (1979).
214. Strassman, R. J. Human psychopharmacology of N,N-dimethyltryptamine. *Behav. Brain*
Res. **73**, 121–124 (1995).
215. Strassman, R. J., Qualls, C. R., Uhlenhuth, E. H. & Kellner, R. Dose-Response Study
of N,N-Dimethyltryptamine in Humans: II. Subjective Effects and Preliminary Results of
a New Rating Scale. *Arch. Gen. Psychiatry* **51**, 98–108 (1994).
216. Alamia, A., Timmermann, C., Vanrullen, R. & Carhart-Harris, R. L. DMT alters cortical
travelling waves. *eLife* **9**, 1–16 (2020).
217. Timmermann, C. *et al.* Neural correlates of the DMT experience assessed with
multivariate EEG. *Sci. Rep.* **9**, 16324 (2019).
218. Callaway, J. C., McKenna, D. J., Grob, C. S., Brito, G. S., Raymon, L. P., Poland, R. E.,
Andrade, E. N., Andrade, E. O. & Mash, D. C. Pharmacokinetics of Hoasca alkaloids in
healthy humans. *J. Ethnopharmacol.* **65**, 243–256 (1999).
219. Fontanilla, D., Johannessen, M., Hajipour, A. R., Cozzi, N. V., Jackson, M. B. & Ruoho,
A. E. The hallucinogen N,N-dimethyltryptamine (DMT) is an endogenous sigma-1
receptor regulator. *Science* **323**, 934–937 (2009).
220. Pierce, P. A. & Peroutka, S. J. Hallucinogenic drug interactions with neurotransmitter
receptor binding sites in human cortex. *Psychopharmacology (Berl.)* **97**, 118–122
(1989).
221. Ray, T. S. Psychedelics and the Human Receptorome. *PLoS ONE* **5**, e9019 (2010).
222. McKenna, D. J., Towers, G. H. & Abbott, F. Monoamine oxidase inhibitors in South
American hallucinogenic plants: tryptamine and beta-carboline constituents of
ayahuasca. *J. Ethnopharmacol.* **10**, 195–223 (1984).
223. Buckholtz, N. S. & Boggan, W. O. Monoamine oxidase inhibition in brain and liver
produced by β-carbolines: structure-activity relationships and substrate specificity.
Biochem. Pharmacol. **26**, 1991–1996 (1977).
224. Mckenna, D., Riba, J., Mckenna, D. & Riba, J. New World Tryptamine Hallucinogens
and the Neuroscience of Ayahuasca. *Curr Top. Behav Neurosci* **36**, 283–311 (2016).
225. Valle, M. *et al.* Inhibition of alpha oscillations through serotonin-2A receptor activation
underlies the visual effects of ayahuasca in humans. *Eur. Neuropsychopharmacol.* **26**,
1161–1175 (2016).
226. van Wel, J. H. P., Kuypers, K. P. C., Theunissen, E. L., Bosker, W. M., Bakker, K. &
Ramaekers, J. G. Effects of acute MDMA intoxication on mood and impulsivity: role of
the 5-HT2 and 5-HT1 receptors. *PloS One* **7**, e40187 (2012).
227. Liechti, M. E. & Vollenweider, F. X. Which neuroreceptors mediate the subjective
effects of MDMA in humans? A summary of mechanistic studies. *Hum.*
Psychopharmacol. **16**, 589–598 (2001).
228. Schmid, Y., Gasser, P., Oehen, P. & Liechti, M. E. Acute subjective effects in LSD- and
MDMA-assisted psychotherapy. *J. Psychopharmacol. (Oxf.)* **35**, (2020).
229. Walpola, I. C., Nest, T., Roseman, L., Erritzoe, D., Feilding, A., Nutt, D. J. & Carhart-
Harris, R. L. Altered Insula Connectivity under MDMA. *Neuropsychopharmacology*
4235, 2152–2162 (2017).

230. Ghahremani, D. G., Tabibnia, G., Monterosso, J., Hellemann, G., Poldrack, R. A. &
London, E. D. Effect of Modafinil on Learning and Task-Related Brain Activity in
Methamphetamine-Dependent and Healthy Individuals. *Neuropsychopharmacology* **36**,
950–959 (2011).
231. Ishizuka, T., Murotani, T. & Yamatodani, A. Modanifil activates the histaminergic
system through the orexinergic neurons. *Neurosci. Lett.* **483**, 193–196 (2010).
232. Volkow, N. D. *et al.* Effects of Modafinil on Dopamine and Dopamine Transporters in
the Male Human Brain: Clinical Implications. *JAMA J. Am. Med. Assoc.* **301**, 1148–
1154 (2009).
233. Gass, J. T. & Foster Olive, M. Glutamatergic substrates of drug addiction and
alcoholism. *Biochem. Pharmacol.* **75**, 218–265 (2008).
234. Wisor, J. P., Nishino, S., Sora, I., Uhl, G. H., Mignot, E. & Edgar, D. M. Dopaminergic
role in stimulant-induced wakefulness. *J. Neurosci. Off. J. Soc. Neurosci.* **21**, 1787–
1794 (2001).
235. Madras, B. K. *et al.* Modafinil Occupies Dopamine and Norepinephrine Transporters in
Vivo and Modulates the Transporters and Trace Amine Activity in Vitro. *J. Pharmacol.*
Exp. Ther. **319**, 561–569 (2006).
236. Cera, N., Tartaro, A., Sensi, S. L. & Jan Stam, C. Modafinil Alters Intrinsic Functional
Connectivity of the Right Posterior Insula: A Pharmacological Resting State fMRI Study.
PLOS ONE (2014) doi:10.1371/journal.pone.0107145.
237. Hannestad, J., Gallezot, J.-D., Planeta-Wilson, B., Lin, S.-F., Williams, W. A., van Dyck,
C. H., Malison, R. T., Carson, R. E. & Ding, Y.-S. Clinically relevant doses of
methylphenidate significantly occupy norepinephrine transporters in humans in vivo.
Biol. Psychiatry **68**, 854–860 (2010).
238. Rowley, H. L., Kulkarni, R. S., Gosden, J., Brammer, R. J., Hackett, D. & Heal, D. J.
Differences in the neurochemical and behavioural profiles of lisdexamfetamine
methylphenidate and modafinil revealed by simultaneous dual-probe microdialysis and
locomotor activity measurements in freely-moving rats. *J. Psychopharmacol. Oxf. Engl.*
28, 254–269 (2014).
239. Koda, K., Ago, Y., Cong, Y., Kita, Y., Takuma, K. & Matsuda, T. Effects of acute and
chronic administration of atomoxetine and methylphenidate on extracellular levels of
noradrenaline, dopamine and serotonin in the prefrontal cortex and striatum of mice. *J.*
Neurochem. **114**, 259–270 (2010).
240. Williard, R. L., Middaugh, L. D., Zhu, H.-J. B. & Patrick, K. S. Methylphenidate and its
ethanol transesterification metabolite ethylphenidate: brain disposition, monoamine
transporters and motor activity. *Behav. Pharmacol.* **18**, 39–51 (2007).
241. Markowitz, J. S., DeVane, C. L., Pestreich, L. K., Patrick, K. S. & Muniz, R. A
comprehensive in vitro screening of d-, l-, and dl-threo-methylphenidate: an exploratory
study. *J. Child Adolesc. Psychopharmacol.* **16**, 687–698 (2006).
242. Volkow, N. D., Fowler, J. S., Wang, G., Ding, Y. & Gatley, S. J. Mechanism of action of
methylphenidate: insights from PET imaging studies. *J. Atten. Disord.* **6 Suppl 1**, S31-
43 (2002).
243. Volkow, N. D., Wang, G. J., Fowler, J. S., Gatley, S. J., Logan, J., Ding, Y. S.,
Hitzemann, R. & Pappas, N. Dopamine transporter occupancies in the human brain
induced by therapeutic doses of oral methylphenidate. *Am. J. Psychiatry* **155**, 1325–
1331 (1998).
244. Whitfield-Gabrieli, S. & Nieto-Castanon, A. Conn: A Functional Connectivity Toolbox for
Correlated and Anticorrelated Brain Networks. *Brain Connect.* **2**, 125–141 (2012).
245. Müller, F., Holze, F., Dolder, P., Ley, L., Vizeli, P., Soltermann, A., Liechti, M. E. &
Borgwardt, S. MDMA-induced changes in within-network connectivity contradict the
specificity of these alterations for the effects of serotonergic hallucinogens.
Neuropsychopharmacology (2020) doi:10.1038/s41386-020-00906-2.
246. Power, J. D., Barnes, K. A., Snyder, A. Z., Schlaggar, B. L. & Petersen, S. E. Spurious
but systematic correlations in functional connectivity MRI networks arise from subject
motion. *NeuroImage* **59**, 2142–2154 (2012).

247. Behzadi Y, Restom K, Liau J, & Liu TT. A component based noise correction method
(CompCor) for BOLD and perfusion based fMRI. *NeuroImage* **37**, 90–101 (2007).
248. Andellini, M., Cannatà, V., Gazzellini, S., Bernardi, B. & Napolitano, A. Test-retest
reliability of graph metrics of resting state MRI functional brain networks: A review. *J.*
Neurosci. Methods **253**, 183–192 (2015).
249. Lydon-Staley, D. M., Ciric, R., Satterthwaite, T. D. & Bassett, D. S. Evaluation of
confound regression strategies for the mitigation of micromovement artifact in studies of
dynamic resting-state functional connectivity and multilayer network modularity. *Netw.*
Neurosci. **3**, 427–454 (2019).
250. Power, J. D., Mitra, A., Laumann, T. O., Snyder, A. Z., Schlaggar, B. L. & Petersen, S.
E. Methods to detect, characterize, and remove motion artifact in resting state fMRI.
NeuroImage **84**, 320–341 (2014).
251. Murphy, K. & Fox, M. D. Towards a consensus regarding global signal regression for
resting state functional connectivity MRI. *NeuroImage* **154**, 169–173 (2017).
252. Tanabe, S. *et al.* Altered Global Brain Signal during Physiologic, Pharmacologic, and
Pathologic States of Unconsciousness in Humans and Rats. *Anesthesiology* 1392–
1406 (2020) doi:10.1097/ALN.0000000000003197.
253. Thompson, P. M. *et al.* ENIGMA and global neuroscience: A decade of large-scale
studies of the brain in health and disease across more than 40 countries. *Transl.*
Psychiatry **10**, 1–28 (2020).
254. Lioi, G., Gripon, V., Brahim, A., Rousseau, F. & Farrugia, N. Gradients of connectivity
as graph Fourier bases of brain activity. *Netw. Neurosci.* **5**, 322–336 (2021).
255. Paquola, C. *et al.* Microstructural and functional gradients are increasingly dissociated
in transmodal cortices. *PLoS Biol.* **17**, e3000284 (2019).
2545