Informing plasmid compatibility with bacterial hosts using protein-protein interaction data

Research Article

Tim Downing[1,2], Min Jie Lee[1], Conor Archbold[1], Adam McDonnell[1], Alexander Rahm[3].

[1] School of Biotechnology, Dublin City University, Dublin, Ireland.
[2] The Pirbright Institute, UK. Corresponding author. Email: Tim.Downing@pirbright.ac.uk Orcid: 0000-0002-8385-6730.
[3] GAATI Lab, University of French Polynesia, Tahiti, French Polynesia. Orcid: 0000-0002-5534-2716.

Highlights

- We tested for protein interactions between key plasmids and 4,363 bacterial hosts
- 23 host species had high rates of protein interactions with four *bla*OXA-48 plasmids
- *Pseudomonas* species showed high rates of interactions with these plasmids
- Host-plasmid protein-protein interaction rates may be associated with compatibility

Abstract

The compatibility of plasmids with new host cells is significant given their role in spreading antimicrobial resistance (AMR)[1] and virulence factor genes. Evaluating this using *in vitro* screening is laborious and can be informed by computational analyses of plasmid-host compatibility through rates of protein-protein interactions (PPIs) between plasmid and host cell proteins. We identified large excesses of such PPIs in eight important plasmids, including pOXA-48, using most known bacteria (n=4,363). 23 species had high rates of interactions with four *bla*OXA-48-positive plasmids. We also identified 48 species with high interaction rates with plasmids common in *Escherichia coli*. We found a strong association between one plasmid and the fimbrial adhesin operon *pil*, which could enhance host cell adhesion in aqueous environments. An excess rate of PPIs could be a sign of host-plasmid compatibility, which is important for AMR control given that plasmids like pOXA-48 move between species with ease.

Keywords: Fimbriae, plasmid, adhesin, protein-protein interaction, network.

---

1. Introduction

Plasmids can provide new genes to host cells, enhancing properties like antimicrobial resistance (AMR) and adhesion to host molecules [1,2]. Plasmids are short extrachromosomal DNA elements that are usually circular, such that conjugative plasmids can move between different host cells sharing the same ecological niche [3,4]. DNA transformation, extracellular vesicles and bacteriophages also permit plasmid gene transfer, albeit less often than conjugation [5-8]. Collectively, these horizontal gene transfer (HGT) processes meaning that the role of plasmids in enabling adaptation to new environments is a key concern in addressing the spread of infectious bacteria. Additionally, plasmids typically induce a small fitness cost in their host cells [9,10]. To overcome this, many plasmids possess genes to parasitise host cells via post-segregation killing and stable inheritance: this ensures their propagation in daughter cells [11] and facilitates vertical plasmid transmission. Many plasmids also endow their host cells with enhanced AMR [12], thereby enhancing their fitness and subsequently the plasmid's survival. Consequently, clonal radiations of emerging pathogens are often associated with the HGT-mediated acquisition of plasmids encoding key AMR genes (like *bla*OXA-48) that enable survival in antibiotic-rich niches.

Plasmid-encoded AMR genes help drive the transmission of carbapenemase-producing *Enterobacteriaceae* (CPE) including *Escherichia coli* and *Klebsiella*. These genes are concerning because they can be transmitted to other bacteria. These two priority pathogens impair human health according to the World Health Organisation (WHO) [13] and were the most common CPEs in England in 2014-16 at 27% (*E. coli*) and 52% (*Klebsiella*) where the most common carbapenemase gene was *bla*OXA-48 [14]. More recent surveys of CPEs in England in 2020-22 confirmed this pattern, showing *E. coli* and *Klebsiella* were responsible for 26% and 33% of cases, respectively [15]. Such plasmid-driven AMR gain by extraintestinal pathogenic *E. coli* (ExPEC) has enhanced their capacity to infect epithelial cells beyond their normal gut niche [16-25].

Highly transmissible pOXA-48-like plasmids are a compelling example of challenges driven by mobile genetic elements [26]. The *bla*OXA-48 gene encodes an Ambler class D beta-lactamase first identified in *Klebsiella pneumoniae* from Turkey [27]. It is usually on a highly mobile 62 Kb conjugative plasmid named pOXA-48a [28] that has spread the *bla*OXA-48 gene globally, causing substantial outbreaks in many countries, including Ireland [29-32]. An estimated 87% of *bla*OXA-48-positive isolates are MDR, and about half of CPE isolated in Europe have this group II OXA carbapenemase gene [33], including *Serratia maracescens* (since 2021). This gene has been found regularly in community and hospital settings across diverse host species, most typically in the *Enterobacterales* order. Host-pOXA-48a co-evolution has occurred, for example through host cell wall permeability changes to alleviate an initial low level of *bla*OXA-48 carbapenemase activity [34].

Beyond pOXA-48, IncF plasmids' roles in spreading AMR genes are well known [12,35-36], including non-conjugative plasmids pEK499 and pEK516 in MDR *E. coli* [37-40]. These plasmids were first identified in England in *E. coli* and have ~75% similarity to each other [37]. Additionally, a distinct conjugative IncI1 *bla*CTX-M-3-*bla*TEM-1b-positive plasmid pEK204 was found in epidemiologically related hosts in Belfast [37-38]. Plasmid pEK204 encodes a type IV pilus (T4P) through its *pil* operon which encodes a thin pilus for epithelial cell adhesion or biofilm formation [41]. Plasmids encoding such fimbrial adhesins could allow bacterial cells to survive in new environments because adherence to human epithelial cells facilitates bacterial colonisation and the formation of biofilms [42]. These adhesins include type 1, P, F1C/S and AFA fimbriae/pili that are typically assembled by the chaperone-usher secretion pathway in gram-negative bacteria [43-47]. Certain pathogens retain multiple different fimbriae that can be expressed depending on the environmental conditions [48-49]. This indicates that possessing a repertoire of fimbriae could be adaptively beneficial. T4P are thin connections that can be elongated or retracted by ATPases [50] and are associated with biofilm formation [51], virulence [52] and bacteriophage infection [53]. The *pil* operon (particulary *pilS*) may allow differential epithelial cell or abiotic surface adhesion with a thin flexible pilus [41] that can elevate conjugation rates in liquid environments [54]. It can also permit biofilm formation such that a T4P-negative cell can adhere to

other T4P-expressing cells bound with a host cell [41]. Although the *pil* operon is somewhat rare, its widespread nature across bacteria implies a role in human infection [55].

These and other findings imply that a plasmid's capacity to interact successfully with a host cell is important and stems more from its genetic profile more so than the host genome [4]. This suggests host fitness increases following plasmid gain depend not only on the environment, but also on plasmid-host protein-protein interactions (PPIs) arising from their collective genetic profiles. These PPIs must be beneficial for plasmids to persist because plasmid-encoded genes must be replicated and expressed in the host cell. This presents the hypothesis that higher rates of plasmid-host PPIs are a signature of higher compatibility between the bacterial cell and the plasmid.

Limited work has examined plasmid-host PPIs across all bacteria. Previously, PPI network analysis has identified AMR genes in *E. coli* [55,56], but has not been applied at the level of whole plasmids. Here, we examined PPIs between plasmid and bacterial hosts to explore the potential compatibility between the host cell and plasmid. We focused on four pOXA-48-related isoforms and four *E. coli* plasmids (pEK204, pEK499, pEK516, pEC958). We assessed these features initially in 4,363 samples representative of most bacteria. We found that it was possible that elevated sample-plasmid PPI levels could be symptomatic of host-plasmid compatibility, and that this observation may stem from operons associated with environments where those bacterial samples have been found.

2. Methods

2.1 Protein-protein interaction data extraction for 24 *Enterobacterales* and all bacteria

We retrieved PPI information for two key datasets from STRING database v12 [57] using R packages Rentrez v1.2.2 [58], STRINGdb v2.0.1 [59], and StringR v1.4.0 [60] (Figure 1), in a working environment using Rv4.0.1 [61], RStudio v2022.2.3.492 [62], Bioconductor v3.11 [63]. The dataset first was for three genera from the *Enterobacterales* order containing: *Escherichia* and *Shigella* (n=11), *Klebsiella* (n=2) and *Serratia* (n=11) (Table S1). *Serratia* belongs to the *Yersiniaceae* family, whereas the others are all *Enterobacteriaceae*. Three other potential samples in StringDB belonging to these genera were excluded because they lacked any pOXA-48-related plasmid-sample PPIs and so had no information. The second dataset was composed of 4,363 StringDB bacterial samples (including the 24 *Enterobacterales* from above) with valid PPI data as outlined previously [79] (Figure 1). Using a score threshold of 400 as per previous work [79], the number of proteins and their PPIs were recorded. These StringDB scores come from a range of sources, including: text mining, pathway databases, co-expression research, small-scale and high-throughput experiments, and computational predictions (chromosomal proximity, phylogenetic context and gene fusion) [57]. The data was processed, collated and visualised using Dplyr v1.0.8 [64], Forcats v0.5.1 [65], Ggplot2 v3.3.5 [66], Ggrepel v0.9.1 [67], ReadR v2.1.2 [68], Readxl v1.4.0 [69], Tibble v3.1.6 [70], TidyR v1.2.0 [71], Tidyverse v1.3.0 [72] and VennDiagram v1.7.3 [73].

Fig 1. The data sources and methods summary, read from right to left. We used PPI data from StringDB related to two datasets: one with 24 *Enterobacterales* (middle) and another with 4,363 diverse bacterial samples (top and bottom). Top: We compared the interacting host proteins and associated PPIs for four plasmids pEK204, pEK499, pEK516 and pEC958 compared to the 24 *Enterobacterales* samples' PPI data to assess their PPIs. These plasmids were also compared to the 4,363 samples' PPI data. Bottom: We examined 31 initial pOXA-48-related plasmids that resolved as four main isoforms whose interacting host proteins and associated PPIs we compared to the 24 *Enterobacterales'* PPI and the 4,363 samples in the same way. Created with BioRender.com.

2.2 Plasmid data retrieval and quality control

The IncL pOXA-48a plasmid has been strongly associated with *E. coli* and *K. pneumoniae* [28]. To explore this plasmid type further, we downloaded accession data for all (n=31) NCBI NucCore entries containing "pOXA-48". Plasmid pKP112 possessed a *bla*OXA-48 gene and therefore was considered a valid pOXA-48 plasmid in this case. These came from the three *Enterobacterales* species: *E. coli* (n=7), *K. pneumoniae* (n=22), and *Serratia marcescens* (n=2). Among these 31 sequences, there were four genetically distinct pOXA-48 isoforms, all isolated from *K. pneumoniae* (strains named Kp11978, KP112, Kpn-30715/15 and K8) (Table 1). Among these four isoforms, pOXA-48_3071 had a *bla*OXA-48 gene that was not properly annotated in the GenBank entry, so it was considered as a pOXA-48-positive plasmid here. We also obtained the unique proteins of four important plasmids linked to antibiotic resistance in *E. coli*: pEK204 (IncI1), pEK499 (IncFII-FIA), pEK516 (IncF) and pEC958 (IncF) (Table S2). Plasmids pEC958 and pEK499 have high similarity (~92%) [74].

Plasmid group	Plasmid name	Strain of origin	Accession number	Length (bp)	Genes	CDSs
pOXA-48	pOXA-48	Kp11978	JN626286.1	61,881	40	74
	pKP112	KP112	LN864819.1	84,252	62	102
	pOXA-48_30715	Kpn-30715/15	NZ_KX523901	65,488	84	84
	pOXA-48_k8	K8	MT441554.1	65,499	87	87
Entero-bacterales	pEK204	Strain C	EU935740	93,732	89	95
	pEK499	Strain A	EU935739	117,536	103	141
	pEK516	Strain D	EU935738	64,471	65	79
	pEC958	EC958	NZ_HG941719	135,602	153	153

Table 1. The plasmids examined came from two groups. The four pOXA-48-related plasmids were originally identified in *K. pneumoniae* (Strain of origin), and the four linked to *Enterobacterales* were initially isolated in *E. coli*.

To assess plasmid protein similarity, we measured the amino acid sequence similarity of the translated genes of the eight plasmids in a pairwise manner using R package Biostrings v2.60.2 [75]. Among the proteins shared by the pOXA-48-related plasmid pairs, 94% (132 out of 138) had a match (*bitscore < −100*). Among the proteins shared by pEK499, pEK516 and pEC958, 94% (130 out of 140) had a match, but pEK204's comparisons with pEK499, pEK516 and pEC958 showed less similarity (18%). Comparisons between each group (pOXA-48-related vs pEK204, pEK499, pEK516 and pEC958) showed only 22% of protein-level similarity for shared proteins.

The PPI information for these eight plasmids was compared with StringDB PPI data for the 24 *Enterobacterales* and 4,363 bacterial samples above using the above packages in R v4.0.1 and Genbankr v1.20.0 [76]. We computed the number of host proteins interacting with each plasmid protein, the number of host-plasmid PPIs, and the Jaccard index of the fraction of interacting host proteins compared to the combined total of sample and plasmid proteins. The number of genes per plasmid was recorded and analysed using R package UpSetR [77]. All p values reported in this study were corrected for multiple testing using the Benjamini-Hochberg method in R.

2.3 Plasmid comparison and estimation of an excess of protein-protein interactions

2,021 samples (out of 4,363) had PPIs with at least one of these four pOXA-48 isoforms, of which 650 had PPIs with all four isoforms. Samples without PPIs were not examined because these provided no information. There were 6,181 sample-plasmid pairs in total with ≥10 interacting proteins and ≥10 PPIs, coming from: pOXA-48 (1,820 pairs), pKP112 (1,868), pOXA-48_30715 (673) and pOXA-48_k8 (1,820). For the four *E. coli* plasmids (pEK204, pEK499, pEK516 and pEC958), all 24 *Enterobacterales*

and 1,604 other samples had ≥1 PPI with at least one of these plasmids. There were 3,828 sample-plasmid combinations with ≥10 interacting host proteins and ≥10 PPIs across these samples, coming from: pEK204 (856), pEK499 (1,324), pEK516 (852) and pEC958 (796).

To quantify instances of high host-plasmid PPI rates for each plasmid set above, we examined on the association between the numbers of interacting host proteins per sample and the number of sample-plasmid PPIs using a linear model predicting the number of PPIs per sample-plasmid pair. We based the prediction on the numbers of: interacting host proteins, plasmid proteins, proteins per sample, and associated pairwise interactions for each plasmid set (Table S3). This showed that the only informative variables were the number of interacting proteins and the number of PPIs per sample and not the other variables. Thus, we used these parameters in a linear model to get the residual numbers of PPIs per sample-plasmid pair, reflecting instances where the observed number of PPIs was far in excess of the expected number. These residuals from the resid R function were normalised such that the average excess of PPIs was zero. We did not explore deficits in the observed PPI numbers.

2.4 Interactions between fimbrial adhesin operons

We focused further on pEK204's *pil* operon, which is associated with adhesion and has 14 genes, *pilI-V* [37]. We used pEK499 as a baseline because it had the most samples with PPIs (1,324 out of 4,363) among the other three plasmids. We found 1,060 samples had detected PPIs with both pEK204 and pEK499. We examined them using a normalised log2-scaled ratio of the Jaccard indices of pEK204 versus pEK499. The pairwise Pearson correlation between 14 *pil* genes was determined using ppcor v1.1 [78] based on presence-absence data for 4,377 samples from across bacteria [79]. Genes with multiple synonyms were collapsed into a single representative. These 14 *pil* genes were also compared to 14 *fim* genes and four *pap* genes in the same manner. We retrieved the *pil, fim* and *pap* gene synonyms and associated Pil, Fim and Pap PPIs in the gene name and PPI data for the 4,363 samples.

3 Results

217 To test for excess PPI rates with different hosts and plasmids, we compared the PPIs of four pOXA-48
218 plasmid isoforms and four key *E. coli* plasmids with two sample collections whose PPI data was
219 retrieved from StringDB. The first sample collection consisted of 24 *Escherichia*, *Klebsiella*, *Serratia*
220 and *Shigella* samples to examine genus-level plasmid links, and the second collection had 4,363 samples
221 from across bacteria to inform on wider patterns.

3.1 No genus-level association of pOXA-48 within *Enterobacterales*

We examined the PPI levels of four known pOXA-48 isoforms (pOXA-48, pKP112, pOXA-48_30715,
pOXA-48_k8, Table 1) with 24 *Enterobacterales* from *Escherichia* and *Shigella* (n=11), *Klebsiella*
(n=2) and *Serratia* (n=11) (Table S1). There was a wide range of PPI levels but no genus-level
association for any of these plasmids, consistent with previous work [80]. To resolve this finding, we
examined the genetic composition of these plasmids. These pOXA-48 isoforms shared just four genes
including *ssb*, *pemI/K* and *bla*OXA-48 (Figure 2A) (*bla*OXA-48 was present but not annotated on
pOXA-48_3071). PemI is the antitoxin of a type II toxin-antitoxin (TA) system for post-segregational
killing that can neutralise its cognate toxin PemK to buffer
plasmids' effects on host cel

is not annotated properly on pOXA-48_3071 and so the intersection of all four plasmids
...es but contains *bla*OXA-48 too). Bottom: The numbers of genes per sample (left, lime-
...with the corresponding sets (right: cyan circles). The most divergent isoform was
...had 22 unique genes. (B) The overlap of genes between the four common *E. coli*
245 plasmids (pEK204, pEK499, pEK516, pEC958). Top: The intersection sizes (y-axis) and the genes per
246 set (purple) showed that nine genes were shared by all four plasmids: *ssb*, *psiB* and *traA/E/L/K/R/V/Y*.
247 Bottom: The numbers of genes per sample (left, pink, set size) with the corresponding sets (right: purple
248 circles). The most divergent isoform was pEK204 (51 unique genes).

3.2 Incomplete associations of pEK204, pEK499, pEK516 and pEC958 with *Escherichia* species

Next, we assessed the level of PPIs between these 24 *Enterobacterales* and four major plasmids (pEK204, pEK499, pEK516, pEC958, Table 1). These plasmids possess numerous AMR genes and are prevalent, particularly in certain *E. coli* phylogroups. As above, the numbers of interacting host proteins per sample and PPIs had no clear association (Table S1). Again, there were elevated rates of interacting proteins and PPIs for nine of ten *Escherichia* samples, but *Klebsiella* and *Serratia* were more heterogeneous (Table S1). Consequently, the plasmids' proteomes appeared to be the major factors, like pOXA-48. The gene content of pEK204, pEK499, pEK516 and pEC958 showed nine shared genes: *ssb*, *psiB* and *traA/E/L/K/R/V/Y* (Figure 2B). The *tra* region is associated with conjugation, *psiB* encodes a plasmid SOS inhibition protein that stops induction of the SOS pathway by binding RecA [82], and *ssb* encodes a ssDNA-binding protein involved in DNA replication, recombination and repair [83].

3.3 Varied patterns of pOXA-48 interactions across bacteria

To shed light on these complex patterns, we extended our analysis of the four pOXA-48 plasmids to all bacterial samples with valid PPI data from StringDB (n=4,363), focusing on 6,181 sample-plasmid combinations with ≥10 interacting host proteins and ≥10 PPIs. Plasmid pOXA-48_30715 had the most interacting host proteins (53±23, median±SD) and PPIs (53±26), followed by pKP112 (interacting proteins, 39±45; PPIs, 43±65), pOXA-48 (interacting proteins, 35±45; PPIs, 37±58), and then pOXA-48_k8 (interacting proteins, 30±33; PPIs, 31±37) (Figure S1). Notably, the interacting host proteins per plasmid had the largest association with the number of PPIs per pair ($0.957 \leq r \leq 0.995$, Figure S2).

Only 0.6% (40 out of 6,401) of the sample-plasmid pairs had high levels of association, measured as an excess of plasmid-linked PPIs ≥100, ≥100 interacting host proteins and ≥20% excess PPIs after using a model to correct for the number of interacting host proteins per sample-plasmid pair (Figure S3, Table S4). 22 of these 40 were linked to pKP112, 13 to pOXA-48, three to pOXA-48_k8 and two to pOXA-48_30715 (Figure 3). *P. aeruginosa* had the most interacting host proteins of these 40: it shared 19 proteins on pKP112 (Figure S4) that had PPIs with 450 other *P. aeruginosa* proteins (Figure S5) with 684 PPIs, an excess of 27%. *P. aeruginosa* also had >300 interacting host proteins, >340 PPIs and an excess of PPIs for pOXA-48. The sample with the 2nd highest number of interacting host proteins was *Aromatoleum aromaticum* EbN1, which had 32 proteins on pOXA-48 (Figure S6) interacting with 252 *A. aromaticum* proteins with 400 PPIs, an excess of 34% (Figure S7). This sample also had 297 interacting proteins linked to pKP112 with 452 PPIs, an excess of 28%. *Enterobacter cloacae subsp. cloacae* ATCC 13047 had 125 other host proteins interacting with pKP112 (Figure S8) with 173 PPIs (an excess of 20%, Figure S9). Two *Legionella* and a *Citrobacter* species also had high associations with pKP112 and pOXA-48. *Klebsiella oxytoca* had 173 interacting host proteins with pKP112 that had 221 PPIs (an excess of 9%), but no such association with the other three plasmids.

3.4 Varied protein-level interactions of bacteria with pEK204, pEK499, pEK516 and pEC958

Assessing pEK204, pEK499, pEK516 and pEC958 in the same manner with the larger collection of 4,363 samples, we identified 1,604 samples with ≥1 PPI with at least one of these plasmids. We focused further on 3,828 sample-plasmid pairs with ≥10 interacting host proteins and ≥10 PPIs in 1,410 samples, of which 48.4% (683 out of 1,410) had PPIs with all four plasmids. The number of interacting host proteins per sample was comparable across all four plasmids, and highest for pEK204 (median of 64 vs 54 for the others, Figure S10). The distributions of numbers of PPIs per sample were similar, but with a much higher variance for pEK204 relative to the other three (13.8 vs 8.2), highlighting that samples had elevated PPI rates (Figure S11).

Here, 1.5% (56 out of 3,828) of sample-plasmid pairs had strong associations, measured as an excess of plasmid-linked PPIs ≥300, ≥100 interacting host proteins and an excess of PPIs of ≥20% (Table S5). 48 were associated with pEK204, five with pEC958, two with pEK499, and one with pEK516 (Figure 4). In particular, the number of pEK204 PPIs per pair varied much more than the other plasmids (Figure S12). Nine of the total of 56 sample-plasmid pairs were from the *Pseudomonas* genus, a pathogen

existing in diverse environments [84]. *P. fluorescens* F113 (aka *P. ogarae* F113) had the most interacting host proteins with pEK204 (500) linked to 1,215 PPIs, an excess PPI rate of 22%. As with pKP112 and pOXA-48, *P. aeruginosa* had 24 pEK204-encoded proteins (including 12 from *pil*) (Figure S13) interacting with 596 other *P. aeruginosa* proteins (Figure S14) with 1,187 PPIs (an excess of 4%). Like pKP112 and pOXA-48 too, *A. aromaticum* EbN1 had 44 proteins on pEK204 including those from the *pil* operon and *tra* region (Figure S15) interacting with 582 other *A. aromaticum* EbN1 proteins (Figure S16) with 1,191 PPIs (an excess of 6%).

Fig 3. The numbers of interacting host proteins per sample (x-axis) versus the excess fraction of PPIs (y-axis) for each sample-plasmid pair for the four pOXA-48 isoforms (A) pOXA-48 (red-green), (B) pKP112 (red-yellow), (C) pOXA-48_30715 (brown-blue) and (D) pOXA-48_k8 (mauve-purple) using data from 6,181 samples' plasmids with >10 interacting host proteins and >10 PPIs each. The number of PPIs per sample-plasmid pair is denoted by the point size and by the shade based on the scale: larger means fewer PPIs, smaller means more PPIs. Note this scale differs for each plasmid. In (B), *P. aeruginosa* had the most interacting proteins with pKP112 (yellow).

Fig 4. The numbers of interacting host proteins per sample (x-axis) versus the excess fraction of PPIs (y-axis) for each sample-plasmid pair for (A) pEK204 (brown-green), (B) pEK499 (brown-blue), (C) pEK516 (red-mauve) and (D) pEC958 (orange-brown) using data from 3,828 samples' plasmids with 10+ interacting proteins and 10+ PPIs each. The number of PPIs per sample-plasmid pair is denoted by the point size and by the shade based on the scale: larger means fewer PPIs, smaller means more PPIs. Note this scale differs for each plasmid. In (A) for pEK204, the five samples with the most interacting proteins per sample were *P. fluorescens* F113 (aka *P. ogarae* F113), *P. aeruginosa, Aromatoleum aromaticum* EbN1, *Rubrivivax gelatinosus* IL144 and *Cellvibrio* sp. BR were the six rightmost samples (all green). *E. coli* K12 MG1655 had the most interacting proteins with pEK499 (B, blue) and pEK516 (C, mauve). In (D), *Ralstonia eutropha* H16 had the most interacting proteins with pEC958 (brown).

3.5 The *pil* operon caused elevated protein interaction signals in pEK204

We examined the conjugative plasmid pEK204 further given the *pil* operon's role in conjugation. 316 samples out of 1,060 with PPIs for both pEK204 and pEK499 had ≥1 of pEK204's 14 *pilI-pilV* genes: the most common were *pilQ* (encoding a T4P secretin) and *pilT* (encoding a T4P twitching motility

ATPase that helps retract the pilus). We scaled the Jaccard index for pEK204 by that for pEK499 to create a log2-scaled ratio that was positively correlated with the number of interacting host proteins for pEK204. This ratio identified samples with a higher potential for pEK204-specific PPIs, including several *Pseudomonas* samples (*P. aeruginosa, P. fluorescens* F113 as above) (Figure 5, Table S6). Some samples had >11 *pil* operon genes, including *Cellvibrio sp. BR*, whose 13 pEK204 proteins (Figure S17) had the highest number of pEK204-related PPIs of all samples (1,383, an excess of 7.4%), interacting with 660 other *Cellvibrio sp. BR* proteins (Figure S18). Samples with the highest pEK204-linked PPIs were associated with urine or water environments based on environmental niche information from BacDive [85], such as *Aromatoleum aromaticum* EbN1, a Betaproteobacterium from a river source. Similarly, water-associated *Rubrivivax gelatinosus* IL144 had 14 pEK204-matching proteins (Figure S19) that had PPIs with 507 other *Rubrivivax gelatinosus* IL144 proteins with 1,127 PPIs (an excess of 2%, Figure S20).

Fig 5. A plot of the normalised log2-scaled ratio of the Jaccard indices of pEK204 divided by that for pEK499 (x-axis) versus the number of sample proteins interacting with pEK204 proteins (y-axis). The data plotted is for 1,060 bacterial samples, of which 316 had ≥1 pEK204-linked *pil* gene(s) (14 genes, *pilIJKLMNOPQRSTUV*. The 12 labelled samples had ≥400 interacting host proteins and a log2-scaled ratio for pEK204 >0. A larger point size and a redder colour signifies the presence of more pEK204-related *pil* genes: blue indicates fewer genes. The histograms indicate the marginal densities of the scatterplots for both axes.

3.6 Interactions of *pil* operon gene products with other fimbrial adhesin proteins

The pairwise correlations of the 14 pEK204 *pil* genes' frequencies across the 4,363 bacterial samples (Table S7) showed that most were positively or negatively correlated with one another, with the

exceptions of *pilL*, *pilQ* and *pilV*. Previous work has shown that PilK-PilV, but not PilI or PilJ, are essential for R64 pilus generation [86]. Adding four *pap* genes and 14 *fim* genes to this identified five sets of genes, of which three involved *pil* genes. The first two sets were negatively correlated within their own groups, but were positively correlated with the other set (*pilO/P/R/T* and *pilA/I/J/M/N/S/U/W/*V). The 3rd gene set was largely uncorrelated and stemmed from from FimT's (a T4P biogenesis protein) and FimV's (a motility protein) interactions with PilQ (a ComQ homolog), PilL and PilV (for FimT) and PilF/G/J (for FimV) in *P. aeruginosa* PAO1 and *Acinetobacter baumannii* A118 [87]. Testing for PPIs between *pil, fim* and *pap* proteins in the 4,363 samples found that *Ralstonia eutropha* H16 was the single sample with a Fim-Pil PPI across its 13 *pil* genes and four *fim* genes (though major structural protein FimA with PilM-O and pilQ, Figure S22A). Notably, both *E. coli* K12 MG1655 (Figure S22B) CFT073 (Figure S22C) had Fim and Pap PPIs, suggesting scope for Pil proteins to interact or evolve such PPIs.

4. Discussion

To explore the hypothesis that an elevated rate of plasmid-host PPIs might be associated with higher compatibility between a plasmid and a host cell, we tested for an excess of host-plasmid PPIs using eight key plasmids in representatives from most bacteria (n=4,363). We achieved this by modelling data from StringDB. We saw that these plasmids had extensive heterogeneity in their PPI levels across 24 *Enterobacterales* (*Escherichia*, *Klebsiella*, *Serratia* and *Shigella*) samples. This absence of a genus-level association was consistent with previous work [80].

To illuminate this issue, we explored the four genetically distinct *bla*OXA-48-positive plasmids in a larger set of 4,363 samples and found that these plasmids had ≥1 PPI with most bacteria, allowing us to identify instances where there was a large excess of PPIs in 40 (0.6%) sample-plasmid pairs. This identified *P. aeruginosa, A. aromaticum* EbN1 and 21 other common animal and plant pathogens as more likely to have PPIs with ≥1 of these *bla*OXA-48-positive plasmids, particularly with the pKP112 and pOXA-48 isoforms, suggesting those plasmids could become common in these species. Consistent with these predictions, *bla*OXA-48-positive human clinical isolates from Bangladesh [88] and Sudan [89] have been found in *P. aeruginosa*, even though previous *in vitro* experiments suggested incompatibility [34]. This is concerning because *bla*OXA-48 is the most common AMR gene in CPE and is prevalent in *Pseudomonas* based on carbapenemase gene rates in England from 2020-22 [15]. Given their already frequent retention of these plasmids, the high levels of carbapenems in clinical environments may allow these hosts to obtain, co-evolve with and compensate for the metabolic cost of possessing these plasmids [90].

We used the same approach to assess the other group of four genetically related plasmids (pEK204, pEK499, pEK516, pEC958) that likely originated in ExPEC [37]. ExPEC pose a significant public health problem and change their plasmid composition to improve their fitness [91]. Like the pOXA-48-related plasmids, these plasmids had ≥1 PPI with many bacteria, and we found large excess rates of PPIs in 48 species. 19% of these were from the *Pseudomonas* genus, where *P. fluorescens* F113 and *P. aeruginosa* had strong associations with pEK204. This was concerning because pEK204 has spread across pathogenic bacteria [26], and IncI1 plasmids like pEK204 promote AMR due to the presence of *bla*CTX-M, *bla*TEM, *bla*CMY and other ESBL genes [92-93].

P. fluorescens F113 and *P. aeruginosa* were among 12 samples with high rates of PPIs with pEK204, particularly in samples whose proteins interacted with *pil* operon proteins. This operon encodes a T4P with two main known isoforms [94], had 14 genes on pEK204, and is a virulence factor for *P. aeruginosa* [95-96]. A ten-gene *pil* operon is involved in conjugation of pathogenicity island PAPI-1 in *P. aeruginosa* PA14 independently of pEK204 [95] that has been linked to *P. aeruginosa* ST309 infection outbreaks [97], and *pil* genes form part of an integrative conjugative element Pac_ICE1 in *P. syringae* pv actinidiae [98]. Notably, *pil* may be more associated with aqueous environments, and the

loss of *pil* genes from PAPI-1 has been associated with better acute human lung [99] and mouse gut
[100] infection, perhaps due to Pil proteins acting as cell surface antigens [101]. Consistent with this
connection to water, our approach identified bacteria associated with rivers like *Cellvibrio sp. BR* and
others found in wastewater, sludge, sewage and freshwater *Rubrivivax gelatinosus* IL144 [100]. We
identified three distinct groups of correlated *pil* gene rates in these 4,363 samples, suggesting that
pEK204's *pil* repertoire differed from other *pil* operon compositions. We found evidence of PPIs
between *pil, fim* and *pap* gene products in *E. coli* and *R. eutropha*, which is important because ExPEC
can possess type 1, IV and P fimbriae and tend to express a single fimbrial type at any time depending
on the host niche, indicating coordinated expression [103-105]. In this way, the gain of *pil* via pEK204
could facilitate superior host survival, conjugation or biofilm formation in water environments [41].

A limitation of this study was the inconsistent nomenclature of plasmid and samples, which meant that
certain protein matches were not detected or misclassified, and affecting PPI identification. For instance,
functions were assigned to ~46% of genes in previous work [106]. In addition, we did not map reads or
align sequences here to detect genes and relied on the accuracy of existing gene annotation tools based
on protein sequence patterns. Such DNA sequence analysis would yield diverse and likely ambiguous
gene cluster alignment scores that also lead to matches being unclear or imprecisely categorised. Future
work is needed to improve genome and plasmid annotation, explore the strength and specificity of PPIs,
and assess other highly transmissible plasmids.

Conclusions

Given that no study yet has explored PPI between key plasmids across all bacteria to date, so we
developed a model testing for excess rates of PPIs between 4,363 bacterial proteomes and the proteomes
of eight plasmids of public health concern. 23 potential bacterial host cells had high PPI rates with four
*bla*OXA-48-positive plasmids, including *P. aeruginosa*. Similarly, 48 species had elevated PPI rates
with four common *E. coli* plasmids, including nine from the *Pseudomonas* genus. One of the latter
plasmids (pEK204), is associated with host cell binding and conjugative transfer in aqueous
environments, and appeared to be a major driver of host-plasmid PPIs in the species with high rates of
PPIs with pEK204. These results overcome a limitation that the gain of a new plasmid may have
heterogenous effects on different host cells, and this can depend on genetic differentiation levels, as has
been illustrated for pOXA-48 [80]. Consequently, our approach using excess rates of PPIs between
plasmid and host cell proteins added results on the potential compatibility of plasmids with particular
species. Our study has implications for controlling emerging problems, such as *bla*OXA-48-carrying
plasmids that can spread among numerous diverse host genera.

Author Contributions

162 TD – Conceptualization, Methodology, Software, Formal analysis, Investigation, Data Curation, Writing
163 - Original Draft, Writing - Review & Editing, Visualisation, Project administration. MJL –
164 Methodology, Formal analysis, Investigation, Data Curation, Visualisation. CA – Formal analysis,
165 Investigation, Data Curation, Visualisation. AMcD – Formal analysis, Investigation, Data Curation,
166 Visualisation. AR – Conceptualization, Methodology, Writing - Review & Editing.

Acknowledgements

170 The authors acknowledge School of Biotechnology (Dublin City University) student funding. We thanks
171 the reviewers and others with comments for their constructive input.

Conflicts of interest

175 The authors declare that there are no conflicts of interest.

Data Summary

Processing files and relevant code associated with this paper are available on Github at
https://github.com/downingtim/plasmids_2022.

S. Kurtz, A. Phillippy, A.L. Delcher, M. Smoot, M. Shumway, C. Antonescu, S. L. Salzberg, Versatile
and open software for comparing large genomes, Genome Biol. 5 (2004) 1–9.

References

1. S. Partridge, S. R., Kwong, S. M., Firth, N. & S. O. Jensen, Mobile genetic elements associated with antimicrobial resistance, Clin. Microbiol. Rev. 31 (2018) e00088-17.
2. A. San Millan, Evolution of plasmid-mediated antibiotic resistance in the clinical context, Trends Microbiol. 26 (2018) 978–985.
3. Y. Che, et al, Conjugative plasmids interact with insertion sequences to shape the horizontal transfer of antimicrobial resistance genes, Proc Natl Acad Sci U S A, 118(6) (2021) e2008731118. doi: 10.1073/pnas.2008731118
4. C. S. Smillie, et al, Ecology drives a global network of gene exchange connecting the human microbiome, Nature 480 (2011) 241–244.
5. U. Canosi, G. Lüder, T. A. Trautner, SPP1-mediated plasmid transduction, J Virol. 44(2) (1982) 431-6. doi: 10.1128/JVI.44.2.431-436.1982
6. S. Erdmann, B. Tschitschko, L. Zhong, M. J. Raftery, R. Cavicchioli, A plasmid from an Antarctic haloarchaeon uses specialized membrane vesicles to disseminate and infect plasmid-free cells, Nat Microbiol. 2(10) (2017) 1446-1455. doi: 10.1038/s41564-017-0009-2
7. X. Zhang, et al, Improvement in the efficiency of natural transformation of *Haemophilus parasuis* by shuttle-plasmid methylation, Plasmid, 98 (2018) 8-14. doi: 10.1016/j.plasmid.2018.07.001
8. T. Wein, T. Dagan, Plasmid evolution, Curr Biol, 30(19) (2020) R1158-R1163. Doi: 10.1016/j.cub.2020.07.003
9. A. San Millan, R. C. MacLean, Fitness Costs of Plasmids: a Limit to Plasmid Transmission. Microbiol Spectr. 5(5) (2017). doi: 10.1128/microbiolspec.MTBP-0016-2017
10. J. P. J. Hall, et al, Plasmid fitness costs are caused by specific genetic conflicts enabling resolution by compensatory mutation, PLoS Biol. 19(10) (2021) e3001225. doi: 10.1371/journal.pbio.3001225
11. M. H. Nicolas-Chanoine, X. Bertrand, J. Y. Madec. *Escherichia coli* ST131, an intriguing clonal group. Clin Microbiol Rev 27 (2014) 543–574.
12. L. Neffe, L. Abendroth, W. Bautsch, S Häussler, J. Tomasch, High plasmidome diversity of extended-spectrum beta-lactam-resistant *Escherichia coli* isolates collected during one year in one community hospital, Genomics 114(3) (2022) 110368. doi: 10.1016/j.ygeno.2022.110368
13. E. Tacconelli, et alm WHO Pathogens Priority List Working Group. Discovery, research, and development of new antibiotics: the WHO priority list of antibiotic-resistant bacteria and tuberculosis, Lancet Infect Dis. 18(3) (2018) 318-327. doi: 10.1016/S1473-3099(17)30753-3
14. K. L. Hopkins, et al, Diversity of carbapenemase-producing *Enterobacterales* in England as revealed by whole-genome sequencing of isolates referred to a national reference laboratory over a 30-month period, J Med Microbiol. 71(5) (2022). doi: 10.1099/jmm.0.001518
15. UK Health Security Agency, Quarterly laboratory surveillance of acquired carbapenemase-producing Gram-negative bacteria in England: October 2020 to March 2022, Health Protection Report 16(5) (2022).
16. J. J. Jacob, A. K. Pragasam, K. Vasudevan, B. Veeraraghavan, G. Kang, J. John, V. Nagvekar, A. Mutreja, *Salmonella* Typhi acquires diverse plasmids from other Enterobacteriaceae to develop cephalosporin resistance, Genomics 113(4) (2021) 2171-2176. doi: 10.1016/j.ygeno.2021.05.003.
17. V. Calhau, G. Ribeiro, N. Mendonça, G. J. Da Silva, Prevalent combination of virulence and plasmidic-encoded resistance in ST 131 *Escherichia coli* strains, Virulence 4 (2013) 726–729.
18. J. T. Poolman, M. Wacker, Extraintestinal pathogenic *Escherichia coli*, a common human pathogen: challenges for vaccine development and progress in the field, J Infect Dis 213 (2016) 6–13.

19. N. L. Ben Zakour, et al, Sequential acquisition of virulence and fluoroquinolone resistance has
shaped the evolution of *Escherichia coli* ST131, mBio 7 (2016) e00347.
20. A. McNally, et al, Combined analysis of variation in core, accessory and regulatory genome
regions provides a super-resolution view into the evolution of bacterial populations, PLoS Genet
12 (2016) 1006280.
21. J. R. Johnson, et al, Household Clustering of *Escherichia coli* Sequence Type 131 Clinical and
Fecal Isolates According to Whole Genome Sequence Analysis, Open Forum Infect Dis 3 (2016)
ofw129.
22. T. Kallonen, et al, Systematic longitudinal survey of invasive *Escherichia coli* in England
demonstrates a stable population structure only transiently disturbed by the emergence of ST131,
Genome Res 27 (2017) 1437–1449.
23. C. Goswami, et al, Genetic analysis of invasive *Escherichia coli* in Scotland reveals determinants
of healthcare-associated versus community-acquired infections. Microb Genom 4 (2018).
24. A. G. Decano, et al, Complete assembly of *Escherichia coli* ST131 genomes using long reads
demonstrates antibiotic resistance gene variation within diverse plasmid and chromosomal
contexts, mSphere 4 (2019) e00130–19.
25. S. Ny, L. Sandegren, M. Salemi, C. G. Giske. Genome and plasmid diversity of extended-
spectrum β-lactamase-producing *Escherichia coli* ST131 – tracking phylogenetic trajectories with
Bayesian inference, Sci Rep 9 (2019) 10291.
26. R. León-Sampedro, et al, Pervasive transmission of a carbapenem resistance plasmid in the gut
microbiota of hospitalized patients, Nat Microbiol. 6(5) (2021) 606-616. doi: 10.1038/s41564-
021-00879-y.
27. L. Poirel, T. Naas, P. Nordmann, Diversity, epidemiology, and genetics of class D beta-
lactamases, Antimicrob Agents Chemother. 54(1) (2010) 24-38. doi: 10.1128/AAC.01512-08
28. L. Poirel, C. Héritier, V. Tolün, P. Nordmann P., Emergence of oxacillinase-mediated resistance
to imipenem in *Klebsiella pneumoniae*, Antimicrob Agents Chemother. 48(1) (2004) 15-22. doi:
10.1128/AAC.48.1.15-22.2004
29. D. J. O'Brien, et al, First isolation and outbreak of OXA-48-producing *Klebsiella pneumoniae* in
an Irish hospital, March to June 2011, Euro Surveill. 16(29) (2011) 19921.
30. C. Wrenn, et al, Investigation of the first outbreak of OXA-48-producing *Klebsiella pneumoniae*
in Ireland, J Hosp Infect 87(1) (2014) 41-6. doi: 10.1016/j.jhin.2014.03.001.
31. D. Morris, et al, Dissemination of clonally related multidrug-resistant *Klebsiella pneumoniae* in
Ireland, Epidemiol Infect. 144(2) (2016) 443-8. doi: 10.1017/S0950268815001041.
32. C. Brehony, et al, Molecular Epidemiology of an Extended Multiple-Species OXA-48 CPE
Outbreak in a Hospital Ward in Ireland, 2018–2019, Antimicrobial Stewardship & Healthcare
Epidemiology 1(1) (2021) e54. doi: https://doi.org/10.1017/ash.2021.206
33. B. D. Johnston, et al, Global molecular epidemiology of carbapenem-resistant *Escherichia coli*
(2002-2017), Eur J Clin Microbiol Infect Dis. 2021 doi: 10.1007/s10096-021-04310-6.
34. L. Poirel, R. A. Bonnin, P. Nordmann, Genetic features of the widespread plasmid coding for the
carbapenemase OXA-48, Antimicrob Agents Chemother. 56(1) (2012) 559-62. doi:
10.1128/AAC.05289-11
35. T. Downing, Tackling Drug Resistant Infection Outbreaks of Global Pandemic *Escherichia coli*
ST131 Using Evolutionary and Epidemiological Genomics, Microorganisms 3(2) (2015) 236-67.
doi: 10.3390/microorganisms3020236
36. C. Goswami, et al, Origin, maintenance and spread of antibiotic resistance genes within plasmids
and chromosomes of bloodstream isolates of *Escherichia coli,* Microb Genom. 6(4) (2020)
e000353. doi: 10.1099/mgen.0.000353
37. N. Woodford, et al, Complete nucleotide sequences of plasmids pEK204, pEK499, and pEK516,
encoding CTX-M enzymes in three major *Escherichia coli* lineages from the United Kingdom, all
belonging to the international O25:H4-ST131 clone, Antimicrob Agents Chemother 53 (2009)
4472–4482.
38. H. Dhanji, et al, Molecular epidemiology of fluoroquinolone-resistant ST131 *Escherichia coli*
producing CTX-M extended-spectrum beta-lactamases in nursing homes in Belfast, UK, J
Antimicrob Chemother. 66(2) (2011) 297-303. doi: 10.1093/jac/dkq463.

39. M. Doumith, et al, Characterization of plasmids encoding extended-spectrum β-lactamases and
their addiction systems circulating among *Escherichia coli* clinical isolates in the UK, J
Antimicrob Chemother. 67(4) (2012) 878-85. doi: 10.1093/jac/dkr553

40. A. G. Decano, T. Downing, An *Escherichia coli* ST131 pangenome atlas reveals population
structure and evolution across 4,071 isolates, Sci Rep. 9(1) (2019) 17394. doi: 10.1038/s41598-
019-54004-5

41. E. G. Dudley, et al, An IncI1 plasmid contributes to the adherence of the atypical
enteroaggregative *Escherichia coli* strain C1096 to cultured cells and abiotic surfaces, Infect
Immun 74 (2006) 2102–211.

42. S. Sarkar, D. Vagenas, M. A. Schembri, M. Totsika, Biofilm formation by multidrug resistant
Escherichia coli ST131 is dependent on type 1 fimbriae and assay conditions, Pathog Dis 74
(2016) ftw013.

43. M. J. Kuehn, J. Heuser, S. Normark, S. J. Hultgren, P pili in uropathogenic *E. coli* are composite
fibres with distinct fibrillar adhesive tips, Nature 356 (1992) 252–255.

44. I. Connell, et al, Type 1 fimbrial expression enhances *Escherichia coli* virulence for the urinary
tract, Proc Natl Acad Sci U S A 93 (1996) 9827–9832.

45. J. J. Martinez, M. A. Mulvey, J. D. Schilling, J. S. Pinkner, S. J. Hultgren, Type 1 pilus-mediated
bacterial invasion of bladder epithelial cells, Embo J 19 (2000) 2803–2812.

46. R. Selvarangan, et al, Interaction of Dr adhesin with collagen type IV is a critical step in
Escherichia coli renal persistence, Infect Immun 72 (2004) 4827–4835.

47. S. P. Nuccio, A. J. Baumler, Evolution of the chaperone/usher assembly pathway: fimbrial
classification goes Greek, Microbiol Mol Biol Rev 71(4) (2007) 551-575.

48. G. T. Werneburg, D. G. Thanassi, Pili Assembled by the Chaperone/Usher Pathway in
Escherichia coli and *Salmonella*, EcoSal Plus 8(1) (2018).

49. D. J. Wurpel, S. A. Beatson, M. Totsika, N. K. Petty, M. A. Schembri, Chaperone-usher fimbriae
of *Escherichia coli*, PLoS One 8(1) (2013) e52835.

50. L. Craig, K. T. Forest, B. Maier, Type IV pili: dynamics, biophysics and functional consequences,
Nat. Rev. Microbiol. 17 (2019) 429–440.

51. G. A. O'Toole, R. Kolter, Flagellar and twitching motility are necessary for *Pseudomonas
aeruginosa* biofilm development, Mol Microbiol 30 (1998) 295–304.

52. J. C. Comolli, et al, *Pseudomonas aeruginosa* gene products PilT and PilU are required for
cytotoxicity in vitro and virulence in a mouse model of acute pneumonia, Infect Immun 67 (1999)
3625–3630.

53. A. K. Schmidt, et al, A Filamentous Bacteriophage Protein Inhibits Type IV Pili To Prevent
Superinfection of *Pseudomonas aeruginosa*, mBio e0244121 (2022). doi: 10.1128/mbio.02441-21

54. J. L. Cottell, M. A. Webber, L. J. L. Piddock, Persistence of transferable extended-spectrum-β-
lactamase resistance in the absence of antibiotic pressure, Antimicrob. Agents Chemother. 56(9)
(2012) 4703-4706.

55. A. G. Decano, et al, Plasmids shape the diverse accessory resistomes of *Escherichia coli* ST131,
Access Microbiol. 3(1) (2020) acmi000179. doi: 10.1099/acmi.0.000179

56. S. K. Miryala, S. Ramaiah, Exploring the multi-drug resistance in *Escherichia coli* O157:H7 by
gene interaction network: A systems biology approach, Genomics 111 (2019) 958–965. doi:
10.1016/j.ygeno.2018.06.002

57. D. Szklarczyk, et al, The STRING database in 2021: customizable protein-protein networks, and
functional characterization of user-uploaded gene/measurement sets, Nucleic Acids Res. 49(D1)
(2021) D605-D612. doi: 10.1093/nar/gkaa1074.

58. D. J. Winter., rentrez: an R package for the NCBI eUtils API, The R Journal 9(2) (2017) 520-526.

59. D. Szklarczyk, et al, STRING v11: protein-protein association networks with increased coverage,
supporting functional discovery in genome-wide experimental datasets, Nucleic acids research
47.D1 (2018) D607-D613.

60. H. Wickham, stringr: Simple, Consistent Wrappers for Common String Operations, R package
version 1.4.0 (2019) https://CRAN.R-project.org/package=stringr

61. R Core Team, R: A language and environment for statistical computing, R Foundation for
Statistical Computing, Vienna, Austria (2021), URL https://www.R-project.org

540   62.   RStudio Team, RStudio: Integrated Development Environment for R, RStudio, PBC, Boston, MA
541   (2022), URL http://www.rstudio.com

542   63.   R. Ihaka, R. Gentleman, R: A Language for Data Analysis and Graphics. Journal of
543   Computational and Graphical Statistics, (5)3 (1996) 299-314.

544   64.   H. Wickham, R. François, L. Henry, K. Müller., dplyr: A Grammar of Data Manipulation, R
545   package version 1.0.8 (2022), https://CRAN.R-project.org/package=dplyr

546   65.   H. Wickham, forcats: Tools for Working with Categorical Variables (Factors), R package version
547   0.5.1 (2021), https://CRAN.R-project.org/package=forcats

548   66.   H. Wickham, ggplot2: Elegant Graphics for Data Analysis, Springer-Verlag New York (2016).

549   67.   K. Slowikowski, ggrepel: Automatically Position Non-Overlapping Text Labels with 'ggplot2', R
550   package version 0.9.1 (2021), https://CRAN.R-project.org/package=ggrepel

551   68.   H. Wickham, J. Hester, J. Bryan, readr: Read Rectangular Text Data, R package version 2.1.2
552   (2022), https://CRAN.R-project.org/package=readr

553   69.   H. Wickham, J. Bryan, readxl: Read Excel Files, R package version 1.4.0 (2022),
554   https://CRAN.R-project.org/package=readxl

555   70.   K. Müller, H. Wickham, tibble: Simple Data Frames, R package version 3.1.6 (2021),
556   https://CRAN.R-project.org/package=tibble

557   71.   H. Wickham, M. Girlich, tidyr: Tidy Messy Data, R package version 1.2.0 (2022),
558   https://CRAN.R-project.org/package=tidyr

559   72.   H. Wickham, et al, Welcome to the tidyverse, Journal of Open Source Software 4(43) (2019) 1686
560   doi: https://doi.org/10.21105/joss.01686.

561   73.   H. Chen, VennDiagram: Generate High-Resolution Venn and Euler Plots, R package version 1.7.3
562   (2022), https://CRAN.R-project.org/package=VennDiagram

563   74.   M. Totsika, et al, Insights into a multidrug resistant *Escherichia coli* pathogen of the globally
564   disseminated ST131 lineage: genome analysis and virulence mechanisms, PLoS One 6(10) (2011)
565   e26578. doi: 10.1371/journal.pone.0026578

566   75.   H. Pagès, P. Aboyoun, R. Gentleman, S. DebRoy, Biostrings: Efficient manipulation of biological
567   strings, R package version 2.60.2 (2021), https://bioconductor.org/packages/Biostrings

568   76.   G. Becker, M. Lawrence, genbankr: Parsing GenBank files into semantically useful objects, R
569   package version 1.24.0 (2022).

570   77.   J. R. Conway, A. Lex, N. Gehlenborg, UpSetR: an R package for the visualization of intersecting
571   sets and their properties, Bioinformatics 33(18) (2017) 2938-2940 doi:
572   10.1093/bioinformatics/btx364

573   78.   S. Kim, ppcor: An R Package for a Fast Calculation to Semi-partial Correlation Coefficients,
574   Communications for Statistical Applications and Methods 22(6) (2015) 665-674.

575   79.   T. Downing, A. D. Rahm, Bacterial plasmid-associated and chromosomal proteins have
576   fundamentally different properties in protein interaction networks, BioRxiv (2022) doi:
577   https://doi.org/10.1101/2022.04.29.490008

578   80.   A. Alonso-del Valle, et al, Variability of plasmid fitness effects contributes to plasmid persistence
579   in bacterial communities, Nat Commun. 12(1) (2021) 2653, doi: 10.1038/s41467-021-22849-y.

580   81.   S. Tsuchimoto, H. Ohtsubo, E. Ohtsubo, Two genes, pemK and pemI, responsible for stable
581   maintenance of resistance plasmid R100, J Bacteriol. 170(4) (1988) 1461-6, doi:
582   10.1128/jb.170.4.1461-1466.1988.

583   82.   A. Bailone, et al, PsiB polypeptide prevents activation of RecA protein in *Escherichia coli*, Mol
584   Gen Genet. 214(3) (1988) 389-95, doi: 10.1007/BF00330471.

585   83.   R. R. Meyer, P. S. Laine, The single-stranded DNA-binding protein of *Escherichia coli*, Microbiol
586   Rev. 54(4) (1990) 342-80, doi: 10.1128/mr.54.4.342-380.1990.

587   84.   A. Peix, M. H. Ramírez-Bahena, E. Velázquez. The current status on the taxonomy of
588   *Pseudomonas* revisited: An update, Infect Genet Evol. 57 (2018) 106-116, doi:
589   10.1016/j.meegid.2017.10.026

590   85.   L. C. Reimer, et al, BacDive in 2019: bacterial phenotypic data for High-throughput biodiversity
591   analysis., Nucleic Acids Research database issue (2019).

592   86.   T. Yoshida, S. R. Kim, T. Komano, Twelve pil genes are required for biogenesis of the R64 thin
593   pilus, J Bacteriol. 181(7) (1999) 2038-43, doi: 10.1128/JB.181.7.2038-2043.1999.

87. R. A. Alm, J. S. Mattick, Identification of two genes with prepilin-like leader sequences involved
in type 4 fimbrial biogenesis in *Pseudomonas aeruginosa*, J Bacteriol. 178(13) (1996) 3809-17,
doi: 10.1128/jb.178.13.3809-3817.1996

88. D. O. Ali, M. H. Arbab, N. M. Admed, Molecular Detection of bla OXA-48 Gene Encoding
Carbapenem Resistance Pseudomonas aeruginosa Clinical Isolates from Khartoum State
Hospitals, Sudan, Gezira Journal of Health Sciences 2(16) (2021) 18-26, doi:
http://journals.uofg.edu.sd/index.php/gjhs/article/view/1690

89. N. Hoque, et al, Genomic diversity and molecular epidemiology of a multidrug resistant
Pseudomonas aeruginosa DMC30b isolated from hospitalized burn patient in Bangladesh,
BioRxiv (2022), doi: https://doi.org/10.1101/2022.07.06.498939

90. A. Newbury, et al, Fitness effects of plasmids shape the structure of bacteria-plasmid interaction
networks, Proc Natl Acad Sci U S A. 119(22) (2022) e2118361119, doi:
10.1073/pnas.2118361119

91. C. Ludden, A. G. Decano, et al, Genomic surveillance of *Escherichia coli* ST131 identifies local
expansion and serial replacement of subclones, Microb Genom. 6(4) (2020) e000352, doi:
10.1099/mgen.0.000352

92. S. L. Foley, P. R. Kaldhone, S. C. Ricke, J. Han, Incompatibility Group I1 (IncI1) Plasmids: Their
Genetics, Biology, and Public Health Relevance, Microbiol Mol Biol Rev. 85(2) (2021) e00031-
20, doi: 10.1128/MMBR.00031-20

93. A. Carattoli, L. Villa, D. Fortini, A. García-Fernández, Contemporary IncI1 plasmids involved in
the transmission and spread of antimicrobial resistance in *Enterobacteriaceae*, Plasmid 118 (2021)
102392. doi: 10.1016/j.plasmid.2018.12.001

94. A. Siryaporn, et al, Surface attachment induces *Pseudomonas aeruginosa* virulence, Proc. Natl
Acad. Sci. USA. 111 (2014) 16860–16865 .

95. A. Persat, et al, Type IV pili mechanochemically regulate virulence factors in *Pseudomonas
aeruginosa*, Proc Natl Acad Sci U S A 112(24) (2015) 7563-8, doi: 10.1073/pnas.1502025112

96. M. Q. Carter, J. Chen, S. Lory, The *Pseudomonas aeruginosa* pathogenicity island PAPI-1 is
transferred via a novel type IV pilus, J Bacteriol. 192(13) (2010) 3249-58, doi: 10.1128/JB.00041-
10.

97. E. L. Fonseca, et al, Emergence of XDR high-risk *Pseudomonas aeruginosa* ST309 in South
America: a global comparative genomic analysis, BioRxiv (2021) doi:
https://doi.org/10.1101/2021.01.21.427610.

98. R. T. M. Poulter, J. Ho, T. Handley, G. Taiaroa, M. I. Butler, Comparison between complete
genomes of an isolate of *Pseudomonas syringae* pv. actinidiae from Japan and a New Zealand
isolate of the pandemic lineage, Sci Rep. 8(1) (2018) 10915. doi: 10.1038/s41598-018-29261-5

99. A. Lorenz, et al, Importance of flagella in acute and chronic *Pseudomonas aeruginosa* infections,
Environ Microbiol. 21(3) (2019) 883-897, doi: 10.1111/1462-2920.14468.

100. D. Skurnik, et al, A comprehensive analysis of in vitro and in vivo genetic fitness of *Pseudomonas
aeruginosa* using high-throughput sequencing of transposon libraries, PLoS Pathog. 9(9) (2013)
e1003582, doi: 10.1371/journal.ppat.1003582.

101. C. S. Arlehamn, T. J. Evans, *Pseudomonas aeruginosa* pilin activates the inflammasome, Cell
Microbiol. 13(3) (2011) 388-401, doi: 10.1111/j.1462-5822.2010.01541.x

102. S. Nagashima, et al, Complete genome sequence of phototrophic betaproteobacterium *Rubrivivax
gelatinosus* IL144, J Bacteriol. 194(13) (2012) 3541-2, doi: 10.1128/JB.00511-12

103. Y. Xia, D. Gally, K. Forsman-Semb, B. E. Uhlin, Regulatory cross-talk between adhesin operons
in *Escherichia coli*: inhibition of type 1 fimbriae expression by the PapB protein, Embo J 19
(2000) 1450–1457.

104. N. J. Holden, B. E. Uhlin, D. L. Gally, PapB paralogues and their effect on the phase variation of
type 1 fimbriae in *Escherichia coli*, Mol Microbiol 42 (2001) 319–330.

105. N. J. Holden, D. L. Gally, Switches, cross-talk and memory in *Escherichia coli* adherence, J Med
Microbiol 53 (2004) 585–593.

106. H. Zhou H, et al, Functions predict horizontal gene transfer and the emergence of antibiotic
resistance, Sci Adv. 7(43) (2021) eabj5056, doi: 10.1126/sciadv.abj5056

3,303 samples had no *pil*-related PPIs

Get pEK204-related pil genes & PPIs

Quality control

1,060 samples had *pil*-related PPIs

4,363 samples with PPI data

4,445 potential samples from StringDB

PPI data inferred for pEK204, pEK499, pEK516 & pEC958 from StringDB

Get sample-plasmid pair's PPI & interacting host proteins

24 Enterobacterales **PPI data:** *Escherichia* / Shigella *(n=11),* **Klebsiella *(n=2)* &** *Serratia* (n=11)

Quality control

27 *Enterobacterales* samples from StringDB

4 pOXA-48 isoforms

Check gene content

31 plasmids matching "pOXA-48"

2,396 samples with pOXA-48-related PPI data

Get pOXA-48-related genes & PPIs

Quality control

4,363 samples with PPI data

4,445 potential samples from StringDB

2,203 samples had no pOXA-48-related PPIs