A glycine zipper motif governs translocation of type VI secretion toxic effectors across the cytoplasmic membrane of target cells

Jemal Ali[1,2,3], Manda Yu[1#], Li-Kang Sung[1], Yee Wai Cheung[1], and Erh-Min Lai[1,2,4*]

[1] Institute of Plant and Microbial Biology, Academia Sinica, Taipei, Taiwan

[2] Molecular and Biological Agricultural Sciences Program, Taiwan International Graduate Program, National Chung-Hsing University and Academia Sinica, Taipei, Taiwan

[3] Graduate Institute of Biotechnology, National Chung-Hsing University, Taichung, Taiwan

[4] Biotechnology Center, National Chung-Hsing University, Taichung, Taiwan

Running title: Glycine zipper mediated effector entry

*Correspondence author

Institute of Plant and Microbial Biology, Academia Sinica

128, Sec. 2, Academia Road, Nankang, Taipei, Taiwan 11529

Tel: 886-2-27871158

E-mail: emlai@gate.sinica.edu.tw

#Current address

Department of Biological Sciences

University of Wisconsin-Milwaukee

Milwaukee, Wisconsin, USA

ABSTRACT

Type VI secretion systems (T6SSs) can deliver diverse toxic effectors into eukaryotic and

bacterial cells. Although much is known about the regulation and assembly of T6SS, the

translocation mechanism of effectors into the periplasm and/or cytoplasm of target cells

remains elusive. Here we used the *Agrobacterium tumefaciens* DNase effector Tde1 to

unravel the mechanism of translocation from attacker to prey. We demonstrate that Tde1

loading onto the secretion machinery is mediated via binding to its adaptor through the

N-terminus, which harbours continuous copies of GxxxG motifs resembling the glycine

zipper structure found in proteins involved in the membrane channel formation. By amino

acid substitutions on a conserved glycine zipper motif, we showed $G^{39}xxxG^{43}$ motif

governs the delivery into target cells by permeabilizing the cytoplasmic membrane. The

findings demonstrate effector itself mediates target cell entry. Considering the widespread

presence of GxxxG motifs in bacterial effectors and pore-forming toxins, we proposed

that glycine zipper mediated permeabilization is a conserved mechanism used by

bacterial effectors for translocation across target cell membranes.

Key words: Type VI secretion system, DNase effector, antibacterial, glycine zipper,

translocation, *Agrobacterium tumefaciens*

INTRODUCTION

In a complex microbial community, bacteria have evolved versatile secretion systems for

export or import of substrates across their membranes in response to different

environmental cues. Each of specialized protein secretion systems (type I to X secretion

system [TISS to TXSS]) [1,2,3] can recognize specific substrates for secretion and

translocation across one or multiple membranes. The type VI secretion system (T6SS) is

a molecular weapon deployed by many Proteobacteria for pathogenesis, antagonism, or

nutrient acquisition [4]. The T6SS effectors discovered so far exert functions in antibacterial,

anti-eukaryotic, and metal acquisition [5-8]. The most established T6SS effectors are

bacterial toxins, in which bacteria also produce cognate immunity proteins to prevent self-
intoxication and toxicity in the sibling cells.

T6SS is a multiprotein complex, composed of at least 13 conserved core proteins
resembling a phage tail structure, that extends from the cytoplasm to the outer membrane
of the attacker cell [9,10]. The T6SS machine consists of the Tss(J)LM membrane complex
(MC), Tss(A)EFGK base plate (BP), TssBC contractile sheath, and Hcp-VgrG-PAAR
puncturing device. The MC interacts with the BP [11 ,12], which serves as a docking site of
VgrG-PAAR effector complex to initiate polymerization of the tail [13]. The tail is composed
of the Hcp inner tube and TssBC outer sheath, and when triggered, the sheath contracts
and ejects out the effector decorated puncturing device into extracellular milieu or target
cells [14,15].

The T6SS has multiple strategies for delivering diverse effectors. On the basis of the
known effectors and their transport mechanisms, effectors can be classified as
"specialized" or "cargo" effectors [9,16]. Specialized effectors are fused to either of the C-
termini of three core structural proteins (Hcp, VgrG, or PAAR) while cargo effectors
interact directly or require a specific chaperone/adaptor to be loaded into the lumen of the
Hcp tube or onto the VgrG spike prior to secretion. Antibacterial effectors can act in the
cytoplasm, membrane, or periplasm of the target cells [5,6,8]. It is not known how effectors
that target the cytoplasm breach both the outer and inner membranes of target cells.

Glycine zipper structure consisting of repetitive GxxxG motifs is commonly found in
membrane-associated proteins [17] and bacterial toxins [18,19]. Glycine zipper motifs are
known be involved in toxicity of some bacterial effectors for membrane channel formation.
For examples, the transmembrane domain (TMD) of a vacuolating toxin, VacA of
Helicobacter pylori encodes three GxxxG motifs forming helix-helix packing interactions
[19], which are required for the vacuolation and membrane channelling contributing to VacA
toxicity [20]. Type I secretion effectors CdzC and CdzD of *Caulobacter crescentus* and
T6SS effector Tse4 of *Pseudomonas aeruginosa,* also possess glycine zipper motifs
involved in antibacterial activity [21,22]. Expression of Tse4 disrupted the proton motive force
of the inner membrane while CdzC and CdzD form surface aggregation for contact-

dependent killing of target cells. However, how glycine zipper motifs of Tse4 and CdzCD

involved in toxicity remain unknown.

A T6SS-encoding locus is highly conserved in the genome of plant pathogenic bacterium

Agrobacterium tumefaciens and the apparatus functions as an antibacterial weapon [23-26].

We previously revealed that *A. tumefaciens* strain C58 deploys two <u>T</u>ype VI <u>D</u>Nase

<u>e</u>ffectors (Tde1 and Tde2) as the major antibacterial weapons, in which the cognate

immunity proteins (namely Tdi1 and Tdi2) prevent autointoxication [23]. Both Tde1 and

Tde2 harbour a novel C-terminal Ntox15 toxin domain [27] containing an HxxD catalytic

motif required for its DNase activity [23]. Tde1 requires its cognate chaperone/adaptor Tap1

for loading onto VgrG1 for secretion [28].

By obtaining an uncoupling Tde1 variant that remains capable for binding to Tap1 for

export but deficient in membrane permeability, we revealed the secretion and

translocation mechanism of Tde1 from the attacker cell to the target cell. We showed that

the N-terminal region of Tde1 harbouring repetitive glycine zipper motifs is sufficient for

interacting with Tap1 for secretion. Once secreted, a conserved glycine zipper motif is

necessary for translocation across target cell membranes. The findings demonstrate a

new role of glycine zipper motif(s) in effector delivery into target cells.

RESULTS

Tde1 can cause DNase-independent growth inhibition of prey cells

Our previous study showed that overexpression of Tde1 in *Agrobacterium tumefaciens*

C58 caused growth inhibition, and the immunity protein Tdi1 only partially protected

against this cytotoxicity [23]. We hypothesized that Tde1 has domains apart from the DNase

domain that contribute to its toxicity. In addition to the C-terminal Ntox15 DNase domain

(amino acid 99-247) [23], Tde1 has a predicted transmembrane domain (TMD, 22-42)

(Fig.1A). Thus, three fragments of Tde1, that being the N-terminal, N-Tde1(1-97) and two

C-terminal regions, C1-Tde1(49-278), and C2-Tde1(98-278) were tested for toxicity. To

avoid confounding effects by the DNase activity, substitutions of catalytic residues

(H190A, D193A) were introduced in the C1-Tde1 and the full length wide-type Tde1 to

become C1-Tde1(M) and Tde1(M) respectively (Fig.1A). Ectopic expression in
Escherichia coli (DH10B) under an IPTG inducible promoter showed that N-Tde1 was
sufficient to inhibit growth (Fig.1B). Tde1(M), but not the C1-Tde1(M), is growth inhibitory.
Surprisingly, C2-Tde1(WT), which retains the wild type DNase catalytic residues, was not
sufficient to inhibit growth. Evidence suggests the N-terminal region of Tde1 is sufficient
to confer toxicity under conditions tested and that the C-terminal DNase domain requires
the N-terminus for it to cause toxicity.

To test Tdi1, the immunity protein for the DNase toxicity of Tde1 [23], can also neutralize
the N-Tde1 toxicity, the Tde1 variants were co-expressed with the Tdi1. The result
showed that Tdi1 could not rescue the growth inhibition caused by the N-Tde1 and
Tde1(M) (Fig 1C). This indicates that Tdi1 cannot neutralize the N-terminus-mediated
toxicity.

A glycine zipper motif in N-terminus of Tde1 is required for toxicity and enhanced
membrane permeability

To get an insight into the cause of growth inhibition by N-terminus of Tde1, we used N-
Tde1 region as a query to search against the NCBI non redundant (nr) database and
identified Tde1 homologues encoded in the T6SS gene clusters of
Agrobacterium/Rhizobium as well as tape measure proteins (TMP) encoded in genomes
of *Paraburkholderia/Burkholderia* (Fig 1D, Fig S1). We noticed the striking conservation
of continuous copies of GxxxG motifs (12-51), which resembles the glycine zipper motifs
overrepresented in membrane proteins and reported to be involved in the membrane
channel formation [17]. Thus, we hypothesized that these repetitive glycine zipper motifs are
necessary for membrane permeability and are the cause for N-Tde1 toxicity.

To test this, two highly conserved glycine residues at position 39 and 43 of a glycine
zipper motif were substituted with leucine (G39L and G43L), and the resulting N-Tde1
and Tde1(M) variants were named as N-Tde1^{GLGL}, Tde1(M)^{GLGL} respectively. The growth
analysis of *E. coli* DH10B cells showed that both N-Tde1^{GLGL} and Tde1(M)^{GLGL} lost the
ability to cause growth inhibition (Fig.2A). Similar results were also observed when they
were overexpressed in *A. tumefaciens* Δ*tde1* mutant (Fig. S2), indicating that the
$G^{39}xxxG^{43}$ glycine zipper motif of Tde1 is responsible for the observed toxicity.

Next, we tested whether N-Tde1 can increase *E. coli* inner membrane permeability. To
do so, we used the β-galactosidase activity assay to measure the entry of ortho
nitrophenyl galactopyranoside (ONPG) (301 Da) into the cytosol. ONPG normally
requires a functional permease LacY to enter into the cytosol but can enter if the inner
membrane is permeabilized/compromised [29,30]. The N-Tde1 and Tde1(M) as well as their
glycine zipper substitution variants were expressed in *E. coli* BW25113Δ*lacY* [31] carrying
β-galactosidase (pYTA-*lacZ*). The BW25113Δ*lacY*(pYTA-*lacZ*) complemented with *lacY*
was used as a positive control.

The results showed that cells expressing either N-Tde1 or Tde1(M) had similar β-
galactosidase activity as LacY expressing cells. In contrast, cells expressing N-Tde1^{GLGL}
and Tde1(M)^{GLGL} only exhibited background level activity as the negative controls (Fig
2B). These results indicate that the N-Tde1 and Tde1(M) are able to increase membrane
permeability depending on the $G^{39}xxxG^{43}$ motif. The data also suggest that the N-
terminus-mediated growth inhibition is caused by its ability to enhance inner membrane
permeability through glycine zipper motifs.

To further analyse the extent of enhanced membrane permeabilization, cells from the
same experiment were normalized to the same OD_{600} and stained with Hoechst and
propidium iodide (PI). Hoechst (616 Da) is a nucleic acid staining dye which is permeable
to live Gram-negative bacterial cells while PI (668.4 Da) can enter through a compromised
inner membrane or dead cells. The PI/Hoechst staining showed strong PI signals in cells
expressing N-Tde1 and Tde1 (M) but no or weak signals were detected in cells expressing
N-Tde1^{GLGL}, Tde1(M)^{GLGL}, or vector control, indicating that N-Tde1 is able to enhance
membrane permeability to allow molecules with size 668.4 Da to pass (Fig 2C). Hoechst
signals were detected in all cells but stronger in cells expressing N-Tde1 or Tde1(M).

The cells used for membrane permeability assays were also collected for counting viable
cells. Accordingly, there was no obvious difference on the number of viable cells among
the strains tested (Fig 2D), suggesting that the detected enhanced PI fluorescence in cells
expressing N-Tde1 and Tde1(M) is due to compromised membrane permeabilization but
not cell death.

We next determined whether the N-terminus of Tde1 is bacteriostatic or bacteriolytic. *E.*

coli cells were induced with IPTG to express N-Tde1 or Tde1(M) and after one hr, washed

with fresh media without IPTG for continuous cultivation. We found that growth was

recovered when cells were washed of the IPTG inducer, in contrast to the growth inhibition

of cells with continuous IPTG induction (Fig 2E). The combined data suggest that the N-

terminus of Tde1 is sufficient to facilitate membrane permeability for bacteriostatic toxicity,

and such activity requires the conserved $G^{39}xxxG^{43}$ glycine zipper motif.

The N-terminus of Tde1 is necessary and sufficient for Tap1 interaction

Tap1 is the adaptor for loading Tde1 onto VgrG1 [23,28]. However, the mechanism that Tde1

and Tap1 interact remains undefined. We first performed co-immunoprecipitation (co-IP)

assay to identify the specific region of Tde1 that can interact with Tap1 in *A. tumefaciens*.

The HA-tagged Tde1 variants were expressed in Δ*tde1* and anti-HA agarose bead was

used to co-precipitate the interacting proteins followed by western blotting to detect Tde1

variants and Tap1. The results showed that the N-Tde1 and Tde1(M) interact with Tap1

but not the C-terminal variants, C1-Tde1(M) and C2-Tde1 (Fig.3A). N-Tde1^{GLGL} and

Tde1(M)^{GLGL} remain capable of interacting with Tap1 (Fig.3A).

Next, we determined whether the Tap1 can interact with Tde1 in *E. coli* where T6SS is

absent. Similar to the results in *A. tumefaciens,* the N-Tde1 and Tde1(M) interacted with

Tap1 but not the C-terminal variants (Fig. 3B). Tap1 interaction with N-Tde1^{GLGL} was also

detected, but since Tde1(M)^{GLGL} was unstable in *E. coli*, intact Tde1^{GLGL} was barely

detected and the interaction with Tap1 was similarly diminished (Fig. 3B). The co-IP

results in *A. tumefaciens* and *E. coli* together suggest that Tap1 interacts with Tde1

through the N-terminus and that the G39L and G43L substitution does not affect Tde1-

Tap1 interaction. However, $G^{39}xxxG^{43}$ motif is crucial to maintain integrity of Tde1(M)

when expressed in *E. coli* (Fig. 3B)*.*

The N-terminus of Tde1 is necessary and sufficient for secretion

Because N-Tde1 interacts with Tap1, we hypothesized that this region is required for Tde1

secretion. Thus, we performed secretion assay by expressing the various HA-tagged

Tde1 variants in Δ*tdei,* a deletion mutant lacking both *tde1-tdi1* and *tde2-tdi2* toxin

immunity pairs. Both cellular and extracellular fractions were collected to determine their

expression and secretion, respectively. The results showed that indeed all Tde1 variants

containing N-terminus is secreted but not the C-terminus, C1-Tde1(M). The secretion was

in a T6SS-dependent manner as secretion was abrogated in *ΔtdeiΔtssK,* which lacks both

tdei and *tssk* encoding the baseplate component. N-Tde1^{GLGL} and Tde1(M)^{GLGL} are also

stably expressed and secreted (Fig. 3C). The data suggest that N-terminus of Tde1 is

necessary and sufficient for secretion and that the G39L and G43L substitution did not

interfere with the secretion capacity of Tde1. Accordingly, Hcp secretion levels are highly

correlated with Tap1-Tde1 interaction and secretion of Tde1 variants (Fig. 3C). The data

also confirmed the requirement of the Tap1-Tde1 interaction for Tde1 secretion and

supported our previous finding that Tde loading onto VgrG is critical for active T6SS

secretion [32].

G^{39}xxxG^{43} motif of Tde1 is required for target cell delivery

Because the G^{39}xxxG^{43} glycine zipper motif located in N-Tde1 increased the membrane

permeability but was not required for secretion, we hypothesized that G^{39}xxxG^{43} is

responsible for inserting Tde1 into the inner membrane and delivering it into the

cytoplasm of target cells. Here, we engineered each of Tde1 variants fused to super-

folder green fluorescence protein (sfGFP) with a flexible (GGGS) linker between Tde1

and sfGFP to avoid the Tde1 functional/structural interference by GFP. The sfGFP fused

Tde1 variants were expressed in *A. tumefaciens Δtdei* and *ΔtdeiΔtssK* mutants. The

secretion assay results showed that N-Tde1-sfGFP and Tde1(M)-sfGFP but not C1-

Tde1(M)-sfGFP are secreted and the G39L and G43L substitution did not compromise

the secretion (Fig 4A). No proteins were observed in the extracellular fractions of

Δtdei1Δtssk mutants, demonstrating that T6SS-dependent secretion.

To investigate the translocation of Tde1 variants, we mixed *A. tumefaciens Δtdei,*

expressing sfGFP-fused Tde1 variants, with *E. coli* cells expressing mCherry. *A.*

tumefaciens expressing sfGFP only (Vector-sfGFP) was used as a negative control. After

co-culture, we imaged populations for mCherry (false coloured in blue) and GFP (green)

to detect *E. coli* and *A. tumefaciens*, respectively. We merged images to identify cyan

colored cells (overlayed blue and green signals), which represent *E. coli* cells with

translocated Tde1 variants carrying sfGFP. We were able to observe cyan fluorescence
when *A. tumefaciens* expressing N-Tde1-sfGFP and Tde1(M)-sfGFP was co-cultured
with *E. coli* mCherry whereas the GFP and mCherry signals were not overlapped in the
E. coli cells co-cultured with *A. tumefaciens* strains expressing GFP only or sfGFP-fused
C1-Tde1(M), N-Tde1^{GLGL}, Tde1(M)^{GLGL} respectively (Fig. 4B). No cyan fluorescence was
observed when N-Tde1-sfGFP and Tde1(M)-sfGFP were expressed in the Δ*tdei1*Δ*tssk*
mutant as the attacker (Fig. S3).

The data suggested that Tde1 is translocated into target cells in a T6SS- and G^{39}xxxG^{43}-
dependent manner. Because N-Tde1^{GLGL}-sfGFP and Tde1(M)^{GLGL}-sfGFP could be
secreted but not translocated into target cells, G^{39}xxxG^{43} motif is necessary for delivering
Tde1 into the target cell.

G^{39}xxxG^{43} is required for interbacterial competition but dispensable for DNase
activity

To assess the role of the G^{39}xxxG^{43} motif for target cell intoxication in the context of cell-
cell contact, we further performed interbacterial competition assays. *A. tumefaciens* C58
Δ*tdei* expressing either Tde1-Tdi1 or its glycine zipper variant, Tde1^{GLGL}-Tdi1, was
competed with target *E. coli* (DH10B) cells. *A. tumefaciens* Δ*tdei* (Tde1-Tdei1*)* exhibited
a strong killing activity (~10^3-fold reduction of target cell survival) as compared to the
negative controls, the secretion deficient mutants Δ*tssL* and Δ*tdei*Δ*tssK* (Tde1-Tdei1*)* (Fig
5A). In contrast, no killing activity could be observed for *A. tumefaciens* Δ*tdei* expressing
Tde1(M)-Tdi1. The killing activity of Δ*tdei*(Tde1^{GLGL}-Tdi1*)* was largely compromised,
similar to that of negative controls. We also performed interbacterial competition assays
using *A. tumefaciens* strain 1D1609, which is susceptible to T6SS killing by C58 [33]. The
intra-species killing activity was weaker (~10^2-fold reduced target cell survival), but the
killing activity of Δ*tdei* (Tde1^{GLGL}-Tdi1*)* against 1D1609 was not detectable (Fig 5B). The
results together indicated that G^{39}xxxG^{43} motif is required for interbacterial competition at
both inter- or intra-species levels. We also performed a secretion assay of these *A.*
tumefaciens attacker strains and as expected, Tde1^{GLGL} proteins were secreted (Fig. 5C).
It is notable that Tde1^{GLGL} proteins accumulate at slightly lower levels to that of Tde1 and
Tde1(M), which also led to reduced secretion levels of Tde1^{GLGL} and Hcp.

Although the reduced secretion level may partly contribute to compromised killing effect

of Δ*tdei*(Tde1^{GLGL}-Tdi1*)*, the most plausible explanation is that Tde1^{GLGL} is unable to

mediate its translocation across target cell membranes. To exclude the possibility that

G39L and G43L substitution may influence its DNase activity, we perform *in vivo* plasmid

DNA degradation assay. Tde1, Tde1^{GLGL}, and Tde1(M) were each expressed by tightly

controlled arabinose-inducible promoter for *in vivo* plasmid DNA degradation assay in *E.*

coli BW25113 as described [23]. As expected, plasmid DNA was completely degraded in

cells expressing Tde1, but not in the negative controls, the cells without arabinose

induction or cells expressing Tde1(M). As for Tde1^{GLGL}, plasmid DNA was also degraded

but not as complete as Tde1 (Fig. 5D). This is not surprising because Tde1(M)^{GLGL} is

prone for degradation in *E. coli* (Fig. 3B). Indeed, no intact Tde1^{GLGL} could be detected,

in contrast to the abundance of Tde1 and Tde1(M) (Fig. 5D). We also generated single

G39L and G43L substitution variants (Tde1^{G39L} and Tde1^{G43L}) and found that both

Tde1^{G39L} and Tde1^{G43L} are stably expressed and exhibit wild-type level DNase activity.

The degree of plasmd DNA degration is also correlated with the growth inhibition effect

(Fig. 5E and Fig. S2C). The slightly recovery of Tde1^{GLGL} in growth inhibition is consistent

with the instability of Tde1^{GLGL}. The evidence that G39L and G43L substitution of Tde1

abolished inner membrane permeability, translocation, and interbacterial competition but

did not affect DNase activity and secretion of Tde1 suggested the G^{39}xxxG^{43} motif is

necessary for delivering Tde1 across the inner membrane into the cytoplasm of target

cells.

DISCUSSION

Through the dissection of Tde1 DNase effector, we provide strong evidence for a role of

the N-terminal glycine-zipper motif(s) of Tde1 in delivering the T6SS effector into target

cells. Here, we proposed a model explaining the loading, firing, and translocation of Tde1

(Fig 6). In *A. tumefaciens,* Tde1 DNase activity is neutralized by Tdi1 by binding to C-

terminal DNase domain while its N-terminal domain interacts with Tap1 for loading onto

VgrG1 (Step 1). The VgrG1-Tap1-Tde1-Tdi1 complex is then recruited onto membrane-

associated baseplate, which serves as a docking site for polymerization of Hcp tube and

TssBC sheath (Step 2). Upon TssBC sheath contraction (Step 3), Tap1 and Tdi1 may fall

off and Hcp-VgrG-Tde1 puncturing device is then ejected for secretion. In contact with a

target cell, Tde1 may be delivered to periplasm of the target cell where Tde1

permeabilises the inner membrane in a $G^{39}xxxG^{43}$ motif-dependent manner (Step 5).

Once delivered, Tde1 exerts its toxicity by attacking DNA for degradation (Step 6).

T6SS cargo effectors often require the specific chaperone/adaptor for loading onto

puncturing device for secretion. Our previous findings demonstrated that Tap1, a

DUF4123-containing protein, specifically interacts with Tde1 for loading onto VgrG1 for

secretion [23,28]. We now showed that N-terminal region of Tde1 is necessary and sufficient

for interaction with Tap1 for secretion and delivery into target cells. The evidence that

Tde1^{GLGL} variant remains capable for binding to Tap1 for export but deficient in membrane

permeability and translocation demonstrated a distinct role of this $G^{39}xxxG^{43}$ motif in

target cell delivery. Among the nine classes of the Ntox15-containing proteins, the

majority of them including Tde1 belong to class I without detectable N-terminal domains

(Fig. S1). We identified the presence of glycine zipper motifs overlapping with

transmembrane domain (TMD) not only in N-terminal region of all Tde1 orthologs

encoded in Rhizobiaceae but also in C-terminal region of tape measure proteins (TMP)

encoded in genomes of *Paraburkholderia/Burkholderia,* likely as a prophage*.* TMP is a

phage protein suggested to have a channel forming activity [34,35] and as a determinant in

connecting host inner membrane proteins for injecting phage genome into bacterial host

cells [36]. Such conservation in Tde1 orthologs suggested that this glycine zipper-mediated

delivery could be a common strategy deployed by these bacterial effectors for

translocation across target cell membranes. To this end, whether other more distantly

related Ntox15-containing proteins or more broadly bacterial toxins harbouring C-terminal

toxin domain may also use N-terminal glycine zipper to mediate effector translocation

await future investigations. It would be also interesting to investigate whether TMP also

employs its C-terminal glycine zipper to mediate phage genome entry into host cells.

Previous study in *P. aeruginosa* showed that VgrG-loaded Tse6-EgaT6 complex is

sufficient to translocate across a lipid bilayer *in vitro* [37], suggesting a role of VgrG-effector

complex itself in inserting across the inner membrane of target cells. A recent study further

uncovered a widespread prePAAR motif in N-terminal TMDs of T6SS effectors involved

in interaction with Eag family chaperone for export [38]. The findings from the Tap1 and Eag

chaperone-mediated T6SS toxins led us to propose that the bacterial toxins harbouring a

N-terminal TMD may be protected by its cognate chaperone/adaptor from insertion into

membranes in the attacker cell. However, once the effector is secreted in the periplasm

of the target cell, specific motifs (such as glycine zippers or perhaps prePAAR) may

mediate TMD to insert into the inner membrane for the delivery into the cytoplasm.

A few membrane permeabilizing T6SS toxins have been reported. The *Vibrio cholerae*

VasX caused dissipation of membrane potential, leading to membrane permeabilization

of target bacterial cells similar to the Tme effectors of *V. parahaemolyticus,* which

represents a widespread family of T6SS effectors harbouring C-terminal TMD for

membrane disruption [39,40]. On the other hand, Tse4 disrupts the membrane potential and

forms a cation selective pore without membrane permeabilization where the pore cannot

even allow the permeability of a relatively smaller molecular weight (ONPG, 300 Da) [21].

Distinct from these toxins in which they confer pore forming activity for toxicity, the role of

glycine zipper(s) of Tde1 appears to enhance membrane permeability for bringing the

toxin domain into target cell cytoplasm.

To date, no structural information is available for Ntox15 superfamily proteins where Tde1

belongs. While N-terminus of Tde1 lacks sequence similarity to any of those known pore-

forming toxins, structural modelling showed structural similarity of two helixes containing

consecutive glycine zipper motifs to the pore forming domain of pyocin S5 [41] (Fig S4).

Pyocin S5 can cause ATP leakage and PI permeability [42] potentially to inner membrane

after translocation through FptA and TonB1 [41]. Tde1 allows the passage of relatively

larger molecule, PI (668 Da), suggesting that its N-terminal glycine zipper(s) may form

larger pores similar to pyocin S5. $G^{39}xxxG^{43}$ motif plays no role in DNase activity of Tde1

but is crucial for its protein stability. Tde1 with substitution of one of the two glycine

residues to leucine retains stability of intact proteins but Tde1 is prone for truncations and

degradation when both glycine residues are substituted to leucine. The instability is most

evident when ectopically expressed in *E. coli* and when retaining DNase activity. Single

glycine substitution (Tde1^{G39L} and Tde1^{G43L} variants) does not influence protein stability

may suggest that the adjacent glycine residues (G^{35} or G^{47}) are sufficient to compensate

the loss of one glycine of $G^{39}xxxG^{43}$ motif for structural integrity in both variants. The

importance of $G^{39}xxxG^{43}$ motif in Tde1 protein stability is consistent with the role of glycine

zippers in structural impact [17].

With the knowledge of effector translocation mechanisms, the bacterial protein secretion

apparatus also offers a strategy for use of effector translocation domain for delivering

heterologous proteins to specific cells. T6SS is a promising vehicle for such purpose

because effectors or secreted proteins appear to be delivered with their folded form,

unlike those to be transported as unfolded forms in most of other specialized secretion

systems [1]. Engineering T6SS carriers such as VgrG spikes to carry exogenous effectors

proteins into target cells are feasible but challenging [43-45]. By using a truncated variant of

PAAR, a recent study showed delivering exogenous T6SS effectors and Cre recombinase

for genetic modification in the target cells [46]. Our success in using N-Tde1 in the delivery

of sfGFP proteins into target *E. coli* cells where they exert fluorescence also suggested

potential applications of N-Tde1 for the delivery of proteins of interests such as genetic

modifiers. This strategy provides advantages over transforming foreign DNA for

expressing a protein of interest from creating undesired genome manipulation.

MATERIALS AND METHODS

Strains and growth conditions

The strains and plasmids used in this study are listed in the supplementary table (S1)

and (S2). The *E. coli* strains used in this study are BW25113 and DH10B. All the *A.*

tumefaciens strains were cultured on 523 medium [47] at 28 °C unless stated. The *E. coli*

strains were cultured on Luria Bertani (LB) medium (10 g L−1 NaCl, 10 g L−1 tryptone,

and 5 g L−1 yeast extract) at 37°C unless stated. Where appropriate, the media were

supplemented with 100 µg mL^{-1} spectinomycin, gentamycin 25 µg mL^{-1} (for *E. coli*) and

50 µg mL^{-1} (for *A. tumefaciens*)*,* 50 µg mL^{-1} ampicillin, 50 µg mL^{-1} kanamycin, 1 mM

Isopropyl β- d-1-thiogalactopyranoside (IPTG).

Growth inhibition assay

For growth inhibition assay of IPTG-inducible expression of Tde1 and its variants, *E. coli*

(DH10B) harbouring pTrc200 vector or the derivatives expressing Tde1 variants were

grown overnight in LB medium supplemented with spectinomycin prior to 1:30 dilution in

a fresh medium and incubated for 2 hrs at 37°C with 250 rpm. After 2 hrs, the cultures

were normalized to OD_{600} 0.1 in LB with 1 mM IPTG for continuous culture in the same

growth condition. The growth of *E. coli* was monitored for OD_{600} every one hr using

ULTROSPEC® 10 cell density meter (Biochrom, UK). For growth inhibition assay of

arabinose-inducible expression, *E. coli* BW25113 harbouring pJN105 vector or the

derivatives expressing Tde1 variants were used. Overnight cultures of *E. coli* cells were

adjusted to an OD_{600} of 0.1 in 200 µL LB with 0.2% L-arabinose in a 96-well plate. The

OD_{600} values were measured by the Synergy H1 microplate reader (Agilent

Technologies, USA) with agitation at 37 °C. The OD_{600} values of final time points were

used to calculate mean ± SD of three repeats. One way Analysis of Variance (ANOVA)

was used for the analysis of statistical significance followed by the Tukey's multiple

comparison.

***In vivo* plasmid DNA degradation assay.** The *in vivo* plasmid DNA degradation assay

was performed as described [23] with minor modifications. Briefly, overnight cultures of *E.*

coli BW25113 carrying pJN105 vector or the derivatives expressing Tde1 variants were

adjusted to an OD_{600} of 0.3 in 4 mL LB with 0.5% D-glucose or 0.2% L-arabinose. After

induction for 3 hrs, bacterial cells normalized by OD_{600} were collected for plasmid DNA

extraction and western blot analysis. The plasmids were then extracted and applied to

0.6% agarose gel electrophoresis to detect the DNA degradation. The OD_{600} values

were measured by DEN-600 photometer (Biosan, Latvia) every hr.

β-galactosidase and viability assays for ONPG update

β-galactosidase assay was performed as described [48] with minor modifications.

BW25113 wild type, BW25113Δ*lacY*(pYTA-lacZ), or BW25113Δ*lacY* harbouring

pTrc200 vector or the derivatives expressing Tde1 variants were grown overnight and

refreshed to a fresh medium as stated for growth inhibition assay. After subculture for 2

hrs, the cells were induced with 1 mM IPTG, and incubated for one more hr. Part of the

culture was adjusted to OD_{600} = 0.3 in Z-buffer and the Intracellular β-galactosidase

activity was measured by mixing 100 µl of 4 mg/ml ONPG with 900 µL of the cells and

incubation at room temperature for 10 mins prior to measurement at OD_{420}. The

remaining cells were normalized to OD_{600} 0.3 in 0.9 % sterile saline and after serial

dilution, 10 µL was spotted on LB plate without antibiotics to recover the viable cells.

Data of OD_{420} were used to calculate mean ± SD of three independent experiments.

One way ANOVA was used for the analysis of statistical significance followed by the

Tukey's multiple comparison.

Co-immunoprecipitation

The Co-immunoprecipitation (co-IP) was performed according to the manufacturer's

recommendations of EZview red Anti-HA agarose (Sigma-E6779) with minor

modifications. To identify Tap1 interacting domain of Tde1, the HA tagged Tde1 variants

were expressed on pTrc200 plasmid. For co-IP in *A. tumefaciens,* C58 Δ*tde1* cells

expressing the Tde1 variants grown in 523 medium overnight were resuspended in a 1:30

ratio to a fresh medium and incubated at 25°C for 3 hrs followed by 1 mM IPTG induction

for additional 3 hrs. After 6 hrs post incubation, the cells were normalized to OD_{600} of 5

per mL in ice-cold PBS buffer (pH 7.4). After cell lysis by lysozyme treatment and

sonication, the lysate was centrifuged and a 100 µL aliquot of the lysate was saved for

the input fraction. The remaining 900 µL lysate was mixed with 25 µL of pre-equilibrated

EZview red Anti-HA agarose and incubated at 4 °C for 1 hr. The beads were then washed

3 times with ice-cold PBS buffer and the proteins bound to the beads were eluted with

100 µL of 2X SDS sample loading buffer. Similarly, the aliquoted input fraction was mixed

with equal volume of 2X SDS sample loading buffer for analysis by western blotting.

For Co-IP in *E. coli*, the HA tagged Tde1 variants were expressed in DH10B cells from

pTrc200 plasmid and Tap1 was expressed from pRL662 plasmid. Overnight culture was

resuspended to fresh medium with 1:30 ratio and incubated for two hrs, followed by

induction with 1 mM IPTG and incubated at 37°C with 250 rpm for two hrs. The cells were

lysed followed by co-IP as stated earlier.

Secretion assay

Type VI Secretion assay was performed in 523 as described previously [32]. Briefly, *A.*

tumefaciens strain was cultured overnight in 523 medium and normalized to OD_{600} of 0.2

in a fresh medium. After 6 hrs of culture, the secreted proteins were collected by

centrifuging at 10,000 g for 5 mins. The resulting pellet was adjusted to OD_{600} of 10 as a

cellular fraction. The culture supernatant was filtered with 0.22 µm Millipore filter

membrane and resulting filtrate was subjected for TCA precipitation [49] and referred as an

extracellular fraction.

Western blotting

448   Western blot analyses were done as previously described [50]. The following primary

449   antibody titres used were: HA epitope (1:4,000), Tap1 (1:3,000) [50], Strep (1:4,000), EF-

450   Tu (1:6,000), and C-terminal Tde1 (1:4,000) [23,28], Hcp (1:2,500) [49].

Interbacterial competition assays

For interbacterial competition with *E. coli* (target), *A. tumefaciens* strain (attacker) was

grown overnight at 28°C in 523 broth with appropriate antibiotics if needed. *E. coli* DH10B

harbouring pRL662 plasmid was grown at 37°C in LB broth with gentamycin. After

harvesting and washing the cells in 0.9% saline, the attacker to target cell density was

adjusted to 30:1 (OD_{600} =3: 0.1) and the mix was spotted on Agrobacterium kill-triggering

(AK) medium containing 1.5% (wt/vol) agar [24]. Similar procedure is used when using *A.*

tumefaciens strain 1D1609 as a target, which was grown at 28°C in 523 broth prior to

competition. After incubation of the mixed strains for 16 hrs at 28°C, the surviving cells

were selected on a gentamycin-containing LB agar square plate.

Fluorescence microscopy

For propidium iodide and Hoechst staining, *E. coli* cells (BW25113) harbouring Trc200

vector or derivatives expressing Tde1 variants were grown overnight and refreshed to a

fresh medium as stated for growth inhibition assay. After subculture for 2 hrs, the cells

were induced with 1 mM IPTG for one hr and OD_{600} equivalent to 0.3 was collected in 1

ml PBS and stained with Hoechst 33342 (H3570) to a final concentration of 12.3 µg/mL

and 1 µg/ml Propidium iodide (2208511) and incubated for 2 minutes in dark. Finally, 2

µL was spotted on 2.5% agarose pad.

For the translocation experiment, the sfGFP fused Tde1 variants were expressed in *A.*

tumefaciens Δtdei cells (attacker). *E. coli* (target) cells were labelled with mCherry (false

colour blue) expressed from pBBRMCS2. *A. tumefaciens* attacker cells were cultured in

523 broth overnight, and *E. coli* target cells were separately cultured on LB broth.

Overnight cultured attacker and target cells were mixed at 5:1 ratio (OD_{600}= 1.0: 0.2), and

10 μL of the mix was cultured on LB-agar plate without IPTG. After 20 hrs of co-culture,

the cells were washed with 100 μL PBS and 2 μL of suspension was spotted on the 2.5%

agarose pad on a microscopic slide. The merged GFP and mCherry signals was based

on a cyan fluorescence.

Fluorescence microscopy was performed using Axio Observer 7 (Zeiss, Germany)

microscope equipped with an Axiochem 702 digital camera and a Plan-Apochromat

100x/1.4 Oil DIC H objective. Exposure times were adjusted to 20 ms for Phase, 20 ms

for Hoechst, 150 ms for PI, 200 ms for GFP and 5000 ms for mCherry. Multiple images

were taken from different fields and all the experiments were performed at least in

triplicate and a representative image is shown. Images were analyzed by using ZEN 2.3

(blue edition) software.

Domain prediction and analysis

Full-length Tde1 (1-278) was used as a query for conserved domain search on the

conserved domains database (CDD) [51] of the National Center for Biotechnology

Information (NCBI). Prediction of transmembrane domain was done using the PRED-

TMR2 [52]. The Tde1 homologues and tape measure proteins (TMPs) for the multiple

sequence alignment were obtained by BLAST search of N-Tde1 (1-97) against the

NCBI non redundant database (nr) with representative sequences selected for multiple

sequence alignment. The domain architectures of the Ntox15 domain containing

proteins were obtained using the full length Tde1 against Conserved Domain

Architectural retrieval tool (CDART) of NCBI. The information of gene clusters encoding

Tde1 homologues and TMPs including upstream and downstream three genes was

retrieved from their respective genomes. Full-length Tde1 (1-278) was used as a query

for structural prediction on a Phyre2 [53]. Three-dimensional structure modelling was done

using Phyre2 in intensive modelling mode. Crystal structure served as best template for

the N-terminus and percentage of confidence for three-dimensional structure modelling

are indicated in the legends of corresponding figures. The structural graphics were

generated by using ChimeraX 1.1 [54].

ACKNOWLEDGEMENT

We would like to thank the Lai lab members for their help and fruitful discussion

throughout this study, Yun-Wei Lien for generating the Δ*tdei*Δ*tssK* strain, and Chih-

Horng Kuo for discussion and suggestion on BLAST analysis. The authors highly

appreciate Jeff Chang, See-Yeun Ting, Lay-Sun Ma, and Chih-Feng Wu for critically

reading the manuscript and their valuable comments. We also acknowledge the

technical assistance of fluorescence microscope provided by Mei-Jane Fang from Live

Cell Imaging Division of Cell Biology Core and Sanger DNA sequencing service

provided by Genomic Technology Core, both located at the Institute of Plant and

Microbial Biology, Academia Sinica. The authors also thank Ying Wang for illustration of

proposed model. The funding was supported by Ministry of Science and Technology of

Taiwan (MOST 110-2311-B-001-032-MY3) and Academia Sinica Investigator Award

(AS-IA-107-L01) to EML. YWC was supported by the postdoctoral fellowship (MOST

110-2811-B-001-524). The funders had no role in study design, data collection and

interpretation, or the decision to submit the work for publication.

AUTHOR CONTRIBUTION

Conceptualization: JA, MY, EML; Investigation: JA, MY, LKS, YWC; Project Funding

acquisition and administration: EML; Supervision: EML; Writing of original draft: JA,

EML; Writing – methodology, review & editing: JA, MY, LKS, YWC, EML.

FIGURE LEGENDS

Figure 1. Schematic domain organization, sequence alignment, growth inhibition

assay. (A) Schematic domain organization of Tde1 protein and its variants. The N-

terminal repeated glycine zipper motifs (12-51) overlapping a predicted transmembrane

domain (22-42) and Ntox15 DNase domain (99-247) are indicated. Tde1 and its variants

with truncation or amino acid substitutions were illustrated. (B) Growth inhibition assay of

E. coli DH10B cells harbouring pTrc200 vector or each of its derivatives expressing Tde1

variants with IPTG induction. (C) Growth inhibition assay of *E. coli* DH10B cells co-

expressing the Tde1 variants expressed from pTrc200 plasmid and Tdi1 immunity gene

expressed from pRL662 plasmid. Growth curve was determined at $OD_{600.}$ Graphs of

panels B and C show mean ± SD of three independent experiments. The significant

differences were shown by the different letters (p value <0.01). (D) Multiple sequence

alignment of N-Tde1 homologues were presented with highly conserved amino acid

residues highlighted in yellow. The bacterial species, strain name, and locus number of

Tde1 orthologs (*Agrobacterium/Rhizobium*) or tape measure proteins

(*Paraburkholderia/Burkholderia*) are indicated on the left and right of aligned sequences.

Two conserved glycine residues (G^{39}, G^{43}) subjected for mutagenesis were indicated by

the arrows above the sequences.

Figure 2. Membrane permeabilization assays. (A) Growth inhibition assay of *E. coli*

DH10B cells harbouring pTrc200 vector or each of its derivatives expressing Tde1

variants with IPTG inducible expression. BW25113 WT or BW25113 (Δ*lacY)* cells

harbouring pTrc200 vector or each of its derivatives expressing Tde1 variants were

carried out for (B) β-galactosidase activity assay to determine ONPG uptake, (C)

propidium iodide permeability with cells treated with Propidium iodide and Hoechst for

detection by fluorescence microscope, and (D) viability assay for *E. coli* cells derived from

the ONPG uptake assay. (E) Bacteriostatic activity assay. *E. coli* DH10B cells harbouring

pTrc200 vector or each of its derivatives expressing Tde1 variants were cultured with or

without IPTG induction for 1 hr. The IPTG-induced cells were further centrifuged and

resuspended in the fresh medium with or without IPTG. Cell density was measured again

before continuous growth for additional 1 hr. Graphs of panels A, B, and E show mean ±

SD of three independent experiments. The significant differences were shown by the

different letters (p value <0.01). Panels C and D are representative results of three
independent experiments.

Figure 3. The N-terminus of Tde1 is sufficient for interaction with Tap1. (A) Co-
immunoprecipitation (Co-IP) in *A. tumefaciens. A. tumefaciens* C58 Δ*tde1* harbouring
pTrc200 vector or its derivatives expressing HA-tagged Tde1 variants. Anti-HA resin was
used to co-IP the Tde1 variants and Tap1. (B) Co-IP in *E. coli* DH10B. DH10B cells co-
expressing HA-tagged Tde1 variants from pTrc200 and Tap1 from pRL662 were collected
for co-IP. Anti-HA resin was used to co-IP the Tde1 variants and Tap-1. Representative
western blot results of three independent experiments were shown with antibody against
HA for Tde1 variants and Tap1 respectively. (C). Secretion assay for HA-tagged Tde1
variants. Western blot for the cellular and extracellular fractions of *A. tumefaciens* C58
Δ*tdei* and Δ*tdei1*Δ*tssK* expressing the HA-tagged Tde1 variants. Hcp secretion was
detected as a positive control for active T6SS secretion. Representative western blot
results of three independent experiments were shown with antibody against HA, Hcp, or
EF-Tu serving as a loading and non-secreted protein control. Protein markers are
indicated in kDa.

Figure 4. The roles of N-Tde1 and G^{39}xxxG^{43} glycine zipper motif for secretion and
translocation. (A) Secretion assay for Tde1 variants fused with sfGFP. Western blot for
the cellular and extracellular fractions of *A. tumefaciens* C58 Δ*tdei* and Δ*tdei1*Δ*tssK*
expressing the Tde1 variants fused with sfGFP were detected by anti-GFP antibody.
Representative western blot results of three independent experiments were shown with
antibody against GFP, Hcp, or EF-Tu serving as a loading and non-secreted protein
control. Hcp secretion served as a positive control for active T6SS secretion. Protein
markers are indicated in kDa. (B) Fluorescence microscopy for Tde1 translocation. *A.*
tumefaciens C58 Δ*tdei* expressing Tde1 variants fused with sfGFP (in green) and *E. coli*
DH10B carrying mCherry (false coloured in blue) were co-cultured for 20 hrs. A cyan
fluorescence with merged blue and green signals represented the translocation of Tde1
variants from *A. tumefaciens* to *E. coli*.

Figure 5. G^{39}xxxG^{43} glycine zipper motif of Tde1 is required for DNase-mediated killing of target cells during interbacterial competition. (A) Inter-species Interbacterial competition. *A. tumefaciens* C58 Δ*tdei* and Δ*tdei1*Δ*tssK* expressing the Tde1 variants were co-incubated with *E. coli* (DH10B) cells on AK medium followed by selecting target cell survival with serial dilution on a gentamycin containing LB agar at 37°C. (B) Intra-species Interbacterial competition between *A. tumefaciens* C58 *(Δtdei)* and Δ*tdei1*Δ*tssK* expressing the Tde1 variants and *A. tumefaciens* 1D1609 target cell. The competition was done as in (B) except 1D1609 was recovered at 28°C. (C) Secretion assay for Tde1 and its variants co-expressed with its immunity protein Tdi1. The toxin immunity pairs Tde1-Tdi1, Tde1^{GLGL}-Tdi1, and Tde1(M)-Tdi1 were expressed in *A. tumefaciens* C58 Δ*tdei* and Δ*tdei*Δ*tssK*. (D) *In vivo* plasmid DNA degradation assay. *E. coli* BW25113 carrying pJN105 empty vector or the derivatives expressing different variants of Tde1 were supplemented with 0.5% glucose ("-") or 0.2% L-arabinose ("+") for 3 hrs to either repress or induce Tde1 production. The plasmids were then extracted to observe the DNA degradation and bottom panel showed western blots of specific Tde1 protein bands indicated by arrows. (E) Growth inhibition assay of Tde1 and its variants. *E. coli* BW25113 cells were induced by adding 0.2% L-arabinose for Tde1 production. The OD$_{600}$ values were measured every 15 minutes. The OD$_{600}$ values of the 4 h post L-arabinose induction were analysed for statistical analysis. Graphs show mean ± SD of three biological repeats. The significant differences were shown by the different letters (p value <0.01). All data were shown with representative results of three independent experiments. Western blots were detected with specific antibody against Tde1, Hcp, or EF-Tu serving as a loading and non-secreted protein control. Protein markers are indicated in kDa.

Figure 6. Proposed model of the loading, firing, and translocation of Tde1. The Tde1 translocation is proposed through six steps. Step 1: Tde1 forms a complex with Tdi1 and Tap1 in the attacker cell. Step 2: Tap1-Tde1-Tdi1 complex binds to the VgrG and the Hcp-VgrG-PAAR puncturing device carrying Tde1 effector complex is loaded onto

membrane-associated baseplate. Step 3: Hcp tube and TssB/C sheath polymerize on the

Tde1-loaded VgrG/baseplate while Tdi1 and Tap1 fall off with unknown mechanisms

before or upon firing. Step 4: TssBC sheath contracts and ejects Tde1 into the target cell

periplasm or cytoplasm. Step 5: The glycine zipper(s) on the N-terminus of Tde1

permeabilize the target cell membrane. Step 6: Intact or truncated Tde1 proteins attack

DNA for degradation in the target cell.

Figure S1. Domain architecture and genetic organizations of Tde homologues. (A)

Domain organization of the Ntox15-containing proteins. Top 10 classes of the Ntox15-

containing proteins are shown with the identifiable domains (not to scale). The number of

proteins in each class were indicated on the left based on the information on June 29,

2022. The *A. tumefaciens* Tde1 belonged to the first class where the N-terminal region

lacks an identifiable domain. (B) Genetic organizations of genes encoding Tde1

orthologues and Tape Measure Proteins (TMPs) with sequence similarity to N-terminus

of Tde1. The proteins encoded from the upstream and downstream of *tde1* and *tmp* genes

are shown with their identified domain organizations.

Figure S2. Growth inhibition assays in *A. tumefaciens* and *E. coli*. Growth curve (A)

and western blot analyses (B) of *A. tumefaciens* C58 Δ*tde1* carrying pTrc200 or its

derivatives expressing HA-tagged Tde1 variants. The growth curve was detected every

2 hrs in 523 medium supplemented with 1 mM IPTG. Graphs show mean ± SD of three

independent experiments. The significant differences were shown by the different letters

(p value <0.01). The proteins collected at end point (6 hr) were analysed for western

blotting with antibody against HA. Representative results of three independent

experiments were shown. Protein markers are indicated in kDa. (C) The growth curve

analysis of *E. coli* cells used for *in vivo* plasmid DNA degradation assay. The turbidity of

E. coli BW25113 expressing Tde1 and its variants carried out for the *in vivo* plasmid DNA

degradation assay were measured. The *E. coli* cells were supplemented with 0.5%

glucose (glu) or 0.2% L-arabinose (ara) for the repression or induction of Tde1 and its

variants. The OD_{600} values were measured by DEN-600 photometer (Biosan, Latvia)
every hr.

Figure S3. Fluorescence microscopy for negative controls of translocation assay.
A. tumefaciens C58 Δ*tdei*Δ*tssK* expressing N-Tde1-sfGFP or Tde1(M)-sfGFP (in green)
and *E. coli* DH10B carrying mCherry (false coloured in blue) were co-cultured for 20 hrs.
No cyan fluorescence with merged blue and green signals could be detected when
attacker cells are T6SS-inactive, which served as negative controls for the translocation
assay.

Figure S4. Structural prediction of the Tde1 N-terminus with similarity to pyocin S5
and colicins. (A) Predicted results of N-terminal Tde1 (1-97) as a query. (B) N-terminal
Tde1 with structural similarity to pore-forming domain of the pyocin S5, colicines and other
membrane perturbing proteins based on Phyre2 prediction. (C) Cartoon model of the
Tde1 (residue 10 – 62) by using on the basis of the crystal structure of Pyocin S5 (PDB
6THK) with 77.3% of identity. Glycine residues of Tde1 were indicated. (D)
Superimposition of N-Tde1 and pore-forming domain of pyocin S5. Tde1 N-terminus is
in red and partially pore-forming domain of pyocin S5 is in teal, G^{39} and G^{43} in the putative
glycine zipper motif are highlighted in green. All data were analyzed by Phyre2 server.

REFERENCES

1   Costa, T. R. *et al.* Secretion systems in Gram-negative bacteria: structural and mechanistic insights.
Nat Rev Microbiol **13**, 343-359, doi:10.1038/nrmicro3456 (2015).
2   Christie, P. J. The Rich Tapestry of Bacterial Protein Translocation Systems. *Protein J* **38**, 389-408,
doi:10.1007/s10930-019-09862-3 (2019).
3   Palmer, T., Finney, A. J., Saha, C. K., Atkinson, G. C. & Sargent, F. A holin/peptidoglycan hydrolase-
dependent protein secretion system. *Mol Microbiol*, doi:10.1111/mmi.14599 (2020).
4   Coulthurst, S. The Type VI secretion system: a versatile bacterial weapon. *Microbiology (Reading)*
165, 503-515, doi:10.1099/mic.0.000789 (2019).
5   Russell, A. B., Peterson, S. B. & Mougous, J. D. Type VI secretion system effectors: poisons with a
purpose. *Nat Rev Microbiol* **12**, 137-148, doi:10.1038/nrmicro3185 (2014).
6   Lien, Y. W. & Lai, E. M. Type VI Secretion Effectors: Methodologies and Biology. *Front Cell Infect*
Microbiol **7**, 254, doi:10.3389/fcimb.2017.00254 (2017).
7   Hachani, A., Wood, T. E. & Filloux, A. Type VI secretion and anti-host effectors. *Curr Opin Microbiol*
29, 81-93, doi:10.1016/j.mib.2015.11.006 (2016).

8   Jurenas, D. & Journet, L. Activity, delivery, and diversity of Type VI secretion effectors. *Mol*
Microbiol, doi:10.1111/mmi.14648 (2020).
9   Cherrak, Y., Flaugnatti, N., Durand, E., Journet, L. & Cascales, E. Structure and Activity of the Type
VI Secretion System. *Microbiol Spectr* **7**, doi:10.1128/microbiolspec.PSIB-0031-2019 (2019).
10   Wang, J., Brodmann, M. & Basler, M. Assembly and Subcellular Localization of Bacterial Type VI
Secretion Systems. *Annu Rev Microbiol* **73**, 621-638, doi:10.1146/annurev-micro-020518-115420
(2019).
11   Durand, E. *et al.* Biogenesis and structure of a type VI secretion membrane core complex. *Nature*
523, 555-560, doi:10.1038/nature14667 (2015).
12   Cherrak, Y. *et al.* Biogenesis and structure of a type VI secretion baseplate. *Nat Microbiol* **3**, 1404-
1416, doi:10.1038/s41564-018-0260-1 (2018).
13   Zoued, A. *et al.* Priming and polymerization of a bacterial contractile tail structure. *Nature* **531**,
59-63, doi:10.1038/nature17182 (2016).
14   Basler, M., Pilhofer, M., Henderson, G. P., Jensen, G. J. & Mekalanos, J. J. Type VI secretion requires
a dynamic contractile phage tail-like structure. *Nature* **483**, 182-186, doi:nature10846 (2012).

15   Vettiger, A. & Basler, M. Type VI Secretion System Substrates Are Transferred and Reused among
Sister Cells. *Cell* **167**, 99-110 e112, doi:10.1016/j.cell.2016.08.023 (2016).
16   Cianfanelli, F. R., Monlezun, L. & Coulthurst, S. J. Aim, Load, Fire: The Type VI Secretion System, a
Bacterial Nanoweapon. *Trends Microbiol* **24**, 51-62, doi:10.1016/j.tim.2015.10.005 (2016).
17   Kim, S. *et al.* Transmembrane glycine zippers: physiological and pathological roles in membrane
proteins. *Proc Natl Acad Sci U S A* **102**, 14278-14283, doi:10.1073/pnas.0501234102 (2005).
18   Fonte, V. *et al.* A glycine zipper motif mediates the formation of toxic beta-amyloid oligomers in
vitro and in vivo. *Mol Neurodegener* **6**, 61, doi:10.1186/1750-1326-6-61 (2011).
19   Kim, S., Chamberlain, A. K. & Bowie, J. U. Membrane channel structure of *Helicobacter pylori*
vacuolating toxin: role of multiple GXXXG motifs in cylindrical channels. *Proc Natl Acad Sci U S A*
101, 5988-5991, doi:10.1073/pnas.0308694101 (2004).
20   McClain, M. S. *et al.* Essential role of a GXXXG motif for membrane channel formation by
Helicobacter   pylori   vacuolating   toxin.   *J   Biol   Chem*   **278**,   12101-12108,
doi:10.1074/jbc.M212595200 (2003).
21   LaCourse, K. D. *et al.* Conditional toxicity and synergy drive diversity among antibacterial effectors.
Nat Microbiol **3**, 440-446, doi:10.1038/s41564-018-0113-y (2018).
22   Garcia-Bayona, L., Guo, M. S. & Laub, M. T. Contact-dependent killing by *Caulobacter crescentus*
via cell surface-associated, glycine zipper proteins. *Elife* **6**, doi:10.7554/eLife.24869 (2017).
23   Ma, L. S., Hachani, A., Lin, J. S., Filloux, A. & Lai, E. M. *Agrobacterium tumefaciens* deploys a
superfamily of type VI secretion DNase effectors as weapons for interbacterial competition in
planta. *Cell Host Microbe* **16**, 94-104, doi:10.1016/j.chom.2014.06.002 (2014).
24   Yu, M., Wang, Y. C., Huang, C. J., Ma, L. S. & Lai, E. M. *Agrobacterium tumefaciens* Deploys a
Versatile   Antibacterial   Strategy   to   Increase   its   Competitiveness.   *J   Bacteriol*,
doi:10.1128/JB.00490-20 (2020).
25   Chou, L. *et al.* Modular evolution of secretion systems and virulence plasmids in a bacterial species
complex. *BMC Biol* **20**, 16, doi:10.1186/s12915-021-01221-y (2022).
26   Wu, C. F. *et al.* Diversification of the Type VI Secretion System in Agrobacteria. *mBio* **12**, e0192721,
doi:10.1128/mBio.01927-21 (2021).
27   Zhang, D., de Souza, R. F., Anantharaman, V., Iyer, L. M. & Aravind, L. Polymorphic toxin systems:
Comprehensive   characterization   of   trafficking   modes,   processing,   mechanisms   of   action,
immunity and ecology using comparative genomics. *Biol Direct* **7**, 18, doi:1745-6150-7-18 (2012).

28   Bondage, D. D., Lin, J. S., Ma, L. S., Kuo, C. H. & Lai, E. M. VgrG C terminus confers the type VI
effector transport specificity and is required for binding with PAAR and adaptor-effector complex.
Proc Natl Acad Sci U S A **113**, E3931-3940, doi:10.1073/pnas.1600428113 (2016).
29   Epand, R. F., Sarig, H., Mor, A. & Epand, R. M. Cell-wall interactions and the selective bacteriostatic
activity of a miniature oligo-acyl-lysyl. *Biophys J* **97**, 2250-2257, doi:10.1016/j.bpj.2009.08.006
(2009).
30   Casteels, P., Ampe, C., Jacobs, F. & Tempst, P. Functional and chemical characterization of
Hymenoptaecin, an antibacterial polypeptide that is infection-inducible in the honeybee (Apis
mellifera). *J Biol Chem* **268**, 7044-7054 (1993).
31   Baba, T. *et al.* Construction of *Escherichia coli* K-12 in-frame, single-gene knockout mutants: the
Keio collection. *Mol Syst Biol* **2**, 2006 0008 (2006).
32   Wu, C. F. *et al.* Effector loading onto the VgrG carrier activates type VI secretion system assembly.
Embo Rep **21**, doi:ARTN e47961DOI 10.15252/embr.201947961 (2020).

33   Santos, M. N. M. *et al.* Redundancy and Specificity of Type VI Secretion vgrG Loci in Antibacterial
Activity of *Agrobacterium tumefaciens* 1D1609 Strain. *Frontiers in Microbiology* **10**, doi:ARTN
3004 10.3389/fmicb.2019.03004 (2020).

34   Roessner, C. A. & Ihler, G. M. Formation of transmembrane channels in liposomes during injection
of lambda DNA. *J Biol Chem* **261**, 386-390 (1986).
35   Roessner, C. A. & Ihler, G. M. Proteinase sensitivity of bacteriophage lambda tail proteins gpJ and
pH in complexes with the lambda receptor. *J Bacteriol* **157**, 165-170, doi:10.1128/jb.157.1.165-
170.1984 (1984).
36   Cumby, N., Reimer, K., Mengin-Lecreulx, D., Davidson, A. R. & Maxwell, K. L. The phage tail tape
measure protein, an inner membrane protein and a periplasmic chaperone play connected roles
in the genome injection process of E. coli phage HK97. *Mol Microbiol* **96**, 437-447,
doi:10.1111/mmi.12918 (2015).
37   Quentin, D. *et al.* Mechanism of loading and translocation of type VI secretion system effector
Tse6. *Nat Microbiol* **3**, 1142-1152, doi:10.1038/s41564-018-0238-z (2018).
38   Ahmad, S. *et al.* Structural basis for effector transmembrane domain recognition by type VI
secretion system chaperones. *Elife* **9**, doi:10.7554/eLife.62816 (2020).
39   Fridman, C. M., Keppel, K., Gerlic, M., Bosis, E. & Salomon, D. A comparative genomics
methodology reveals a widespread family of membrane-disrupting T6SS effectors. *Nat Commun*
11, 1085, doi:10.1038/s41467-020-14951-4 (2020).
40   Miyata, S. T., Unterweger, D., Rudko, S. P. & Pukatzki, S. Dual expression profile of type VI
secretion system immunity genes protects pandemic *Vibrio cholerae*. *PLoS Pathog* **9**, e1003752,
doi:10.1371/journal.ppat.1003752 (2013).
41   Behrens, H. M. *et al.* Pyocin S5 Import into *Pseudomonas aeruginosa* Reveals a Generic Mode of
Bacteriocin Transport. *mBio* **11**, doi:10.1128/mBio.03230-19 (2020).
42   Ling, H., Saeidi, N., Rasouliha, B. H. & Chang, M. W. A predicted S-type pyocin shows a bactericidal
activity against clinical *Pseudomonas aeruginosa* isolates through membrane damage. *FEBS Lett*
584, 3354-3358, doi:10.1016/j.febslet.2010.06.021 (2010).
43   Wettstadt, S. & Filloux, A. Manipulating the type VI secretion system spike to shuttle passenger
proteins. *PLoS One* **15**, e0228941, doi:10.1371/journal.pone.0228941 (2020).
44   Wettstadt, S., Lai, E. M. & Filloux, A. Solving the Puzzle: Connecting a Heterologous *Agrobacterium*
tumefaciens T6SS Effector to a *Pseudomonas aeruginosa* Spike Complex. *Front Cell Infect*
Microbiol **10**, 291, doi:10.3389/fcimb.2020.00291 (2020).

45   Ho, B. T., Fu, Y., Dong, T. G. & Mekalanos, J. J. *Vibrio cholerae* type 6 secretion system effector
trafficking in target bacterial cells. *Proc Natl Acad Sci U S A* **114**, 9427-9432,
doi:10.1073/pnas.1711219114 (2017).
46   Hersch, S. J., Lam, L. & Dong, T. G. Engineered Type Six Secretion Systems Deliver Active Exogenous
Effectors and Cre Recombinase. *mBio* **12**, e0111521, doi:10.1128/mBio.01115-21 (2021).
47   Kado, C. I., and Heskett, M.G. . Selective media for isolation of *Agrobacterium, Carynebacterium,*
Erwinia, Pseudomonas, and *Xanthomonas*. . *Phytopathology* **60**, 969-976 (1970).
48   Saint Jean, K. D. *et al.* Effects of Hydrophobic Amino Acid Substitutions on Antimicrobial Peptide
Behavior. *Probiotics Antimicrob Proteins* **10**, 408-419, doi:10.1007/s12602-017-9345-z (2018).
49   Wu, H. Y., Chung, P. C., Shih, H. W., Wen, S. R. & Lai, E. M. Secretome Analysis Uncovers an Hcp-
Family Protein Secreted via a Type VI Secretion System in *Agrobacterium tumefaciens*. *J. Bacteriol*
190, 2841-2850 (2008).
50   Lin, J. S., Ma, L. S. & Lai, E. M. Systematic Dissection of the *Agrobacterium* Type VI Secretion
System Reveals Machinery and Secreted Components for Subcomplex Formation. *PLoS One* **8**,
e67647, doi:10.1371/journal.pone.0067647 (2013).

51   Lu, S. *et al.* CDD/SPARCLE: the conserved domain database in 2020. *Nucleic Acids Res* **48**, D265-
D268, doi:10.1093/nar/gkz991 (2020).
52   Pasquier, C., Promponas, V. J., Palaios, G. A., Hamodrakas, J. S. & Hamodrakas, S. J. A novel method
for predicting transmembrane segments in proteins based on a statistical analysis of the SwissProt
database: the PRED-TMR algorithm. *Protein Eng* **12**, 381-385, doi:10.1093/protein/12.5.381
(1999).
53   Kelley, L. A., Mezulis, S., Yates, C. M., Wass, M. N. & Sternberg, M. J. The Phyre2 web portal for
protein modeling, prediction and analysis. *Nat Protoc* **10**, 845-858, doi:10.1038/nprot.2015.053
(2015).
54   Goddard, T. D. *et al.* UCSF ChimeraX: Meeting modern challenges in visualization and analysis.
Protein Sci **27**, 14-25, doi:10.1002/pro.3235 (2018).
55   Ma, L. S., Lin, J. S. & Lai, E. M. An IcmF family protein, ImpLM, is an integral inner membrane
protein interacting with ImpKL, and its walker a motif is required for type VI secretion system-
mediated Hcp secretion in *Agrobacterium tumefaciens*. *J Bacteriol* **191**, 4316-4329, doi:JB.00029-
09 (2009).

56   Schmidt-Eisenlohr, H., Domke, N. & Baron, C. TraC of IncN plasmid pKM101 associates with
membranes and extracellular high-molecular-weight structures in Escherichia coli. *J Bacteriol* **181**,
5563-5571 (1999).
57   Vergunst, A. C. *et al.* VirB/D4-dependent protein translocation from *Agrobacterium* into plant cells.
Science **290**, 979-982, doi:8951 [pii] (2000).
58   Newman, J. R. & Fuqua, C. Broad-host-range expression vectors that carry the L-arabinose-
inducible *Escherichia coli araBAD* promoter and the *araC* regulator. *Gene* **227**, 197-203,
doi:S0378-1119(98)00601-5 [pii] (1999).

814   **Table S1. List of Strains and plasmids**

Strain/plasmid Name	Relevant Characteristics	Source/Reference	EML No
A. tumefaciens			
C58	Wild type, virulent strain containing nopaline-type Ti plasmid pTiC58	Eugene Nester	EML530
ΔtssL	tssL deletion mutant	[55]	EML1073
Δtde1	Deletion of atu4350		
Δtdei	Deletion of tde1-tdi1 (atu4350-atu4351) and tde2-tdi2 (atu3640-atu3639) toxin immunity pairs	[23]	EML3559
ΔtdeiΔtssK	Deletion of tssK in Δtdei	This Study	EML5136
E. coli			
DH10B	F- araDJ39 Δara, leu)7697 ΔlacX74 galU galK rpsL deoR φ80dlacZΔM15 endAI nupG recAI mcrA Δ (mrr hsdRMS mcrBC)	Invitrogen	EML5396
BW25113	rrnB Δ lacZ4787 hsdR514 Δ (araBAD)567 DE (rhaBAD)568 rph-1 (ME9062)	NBRP, Japan	EML5398
BW25113 (ΔlacY)	lacY deletion mutant of the BW25113	NBRP, Japan	
Plasmids			
pTrc200	Sm^R, Sp^R, pVS1 origin lacIq, trc promoter expression vector	[56]	EML5481
pRL662	Gm^R, non-conjugal transferable broad host range vector derived from pBBR1MCS-2,	[57]	EML5452
pRL662-Tap1	Constitutive expression of Tap1 on pRL662 plasmid	[23]	EML4269
pRL662-Tdi1-Strep	Constitutive expression of Tdi1-Strep by lac promoter on the pRL662 plasmid	[23]	EML3398
pTrc-Tde1(M)-HA	Sp^R, C-terminal HA-tagged Tde1(M) driven by trc promoter on pTrc200	This Study	EML5445
pRLTdi1-Strep	Gm^R, constitutive expression of C-terminal Strep-tagged Tdi1 by lac promoter on pRL662	This Study	EML5453
pTrc-N-Tde1-HA	Sp^R, N-terminus of Tde1 (1-49) with C-terminal HA tag driven by trc promoter on pTrc200	This Study	EML5482
pTrc-N-Tde1^{GLGL}-HA	Sp^R, N-terminus of Tde1 (1-49) with amino acid substitutions (G39L, G43L) and HA tag on C-terminus driven by trc promoter on pTrc200	This Study	EML5464
pTrc-Tde1(M)^{GLGL}-HA	Sp^R, C-terminal HA-tagged Tde1(M) with amino acid substitutions (G39L, G43L) driven by trc promoter on pTrc200	This study	EML5465
pTrc-C1-Tde1(M)-HA	Sp^R, C-terminus of Tde1(M) (49-278) with C-terminal HA tag driven by trc promoter on pTrc200.	This study	EML5479

pTrc-C2-Tde1(WT)-HA	Sp[R], C-terminus of Tde1 (98-278) with C-terminal HA tag driven by *trc* promoter on pTrc200.	This Study	EML5478
pYTA_LacZ	Amp[R], full length *lacZ* gene was cloned on to the YTA vector by TA cloning	This Study	EML1661
pBBRMCS2-mCherry	Km[R], full length mCherry gene was cloned on to pBBRMCS2 plasmid	This Study	EML3022
pTrc200-LacY	Sp[R], *lacY* from BW25113 with C-terminal HA tag driven by *trc* promoter on pTrc200 plasmid.	This Study	EML6350
pTrc-N-Tde1-sfGFP	Sp[R], N-terminus of Tde1 (1-49) fused to the sfGFP on the C-terminal with a GGGS linker driven by *trc* promoter on pTrc200 plasmid	This Study	EML5489
pTrc-N-Tde1[GLGL]-sfGFP	Sp[R], N-terminus of Tde1 (1-49) with point mutation (G39L, G43L) fused to the sfGFP on the C-terminal with a GGGS linker driven by *trc* promoter on pTrc200 plasmid	This Study	EML5490
pTrc-Tde1(M)-sfGFP	Sp[R], full length of Tde1(M) fused to the sfGFP on the C-terminal with a GGGS linker driven by *trc* promoter on pTrc200 plasmid	This Study	EML5493
pTrc-Tde1(M)[GLGL]-sfGFP	Sp[R], full length of Tde1(M) with point mutation on (G39L, G43L) fused to the sfGFP on the C-terminal with a GGGS linker driven by *trc* promoter on pTrc200 plasmid	This Study	EML5494
pTrc-C1-Tde1(WT)-HA	Sp[R], C-terminus of Tde1 (49-278) fused to the sfGFP on the C-terminal with a GGGS linker driven by *trc* promoter on pTrc200 plasmid	This Study	EML5492
pTrc-C2-Tde1(WT)-HA	Sp[R], C-terminus of Tde1 (98-278) fused to the sfGFP on the C-terminal with a GGGS linker driven by *trc* promoter on pTrc200 plasmid	This Study	EML5491
pTrc200-sfGFP	Sp[R], sfGFP was cloned in to the pTrc200 plasmid	This Study	EML5495
pJN105	Gm[R], arabinose-inducible gene expression vector derived from pBBRMCS-1, araC-P$_{BAD}$	[58]	
pJN105_Tde1	Gm[R], pJN105 expressing C-terminal 6xHis tagged Tde1	This study	
pJN105_Tde1[G39L]	Gm[R], pJN105 expressing C-terminal 6xHis tagged Tde1 with the substitution of G39L	This study	
pJN105_Tde1[G43L]	Gm[R], pJN105 expressing C-terminal 6xHis tagged Tde1 with the substitution of G43L	This study	
pJN105_Tde1[GLGL]	Gm[R], pJN105 expressing C-terminal 6xHis tagged Tde1 with the substitutions of G39L and G43L	This study	
pJN105_Tde1(M)	Gm[R], pJN105 expressing C-terminal 6xHis tagged Tde1 with the substitutions of H190A and D193A	This study	

817 **Table S2. List of primers used to make constructs**

Plasmid/Construct made with the primers	Primer names and Sequence (restriction site is indicated underline)
pTrc200_sfGFP	NcoI_sfGFP_F: TGGAC<u>CCATGG</u>ACCGTAAAGGCGAAGAGCTGTTCA HindIII_sfGFP_R:TCGAC<u>AAGCTT</u>TTTGTACAGTTCATCCATACCATG
pTrc200_Tde1(M)-HA pTrc200_N-Tde1-HA pTrc200_N-Tde1^G-HA	NCoI_Tde1_F: AAAAA<u>CCATGG</u>ACATGAGTGCGACGACAACTGTCG XbaI-Tde1_R: AAAAA<u>TCTAGA</u>CACCGGGACGTCAGGATTGCC XbaI_N-tde1_R: AAAAA<u>TCTAGA</u> A GCT CCC TCA GGC GGA CTA AAG CAA T
pTrc200_C1Tde1(M)-HA pTrc200_C2-Tde1(M)-HA	NCoI_C2-Tde1_F: AAAAA<u>CCATGG</u>ACACGCCTAAGCAAGTTGAAGAGTTTAGG Ncol_C1_Tde1_F: AAAAA<u>CCATGG</u>ACGCGGGTAGAGCGGCTGCCTCCGCAA
pTrc200_N-Tde1^G-HA pTrc200_Tde1(M)^G-HA	G39L_G43L_F: TGGAGGTACAGCGGTTTTAGCATGGATACTTGGGCAAGTCGGCGCTAT G39L_G43L_R: ATAGCGCCGACTTGCCCAAGTATCCATGCTAAAACCGCTGTACCTCCAG
pTrc200_lacY-HA	Ncol_LacY_F: AAAAA<u>CCATGG</u>ACATGTACTATTTAAAAAACACAAAC XbaI_LacY_R: AAAAA<u>TCTAGA</u>AGCGACTTCATTCACCTGACGAC
pTrc200_N-Tde1-sfGFP	N-tde_GGGS_sfGFP_F: TTAGTCCGCCTGAGGGAGCT<u>TCTAGA</u>GGTGGTGGTTCTCGTAAAGGCGAAGAGCTGTT sfGFP_GGGS_Ntde1_R: ACAGCTCTTCGCCTTTACGAGAACCACCACC<u>TCTAGA</u>AGCTCCCTCAGGCGGACTAA
pTrc200_sfGFP	Ncol_sfGFP_F: TGGAC<u>CCATGG</u>ACCGTAAAGGCGAAGAGCTGTTCA HindIII_sfGFP_R:TCGAC<u>AAGCTT</u>TTTGTACAGTTCATCCATACCATG
pJN105_Tde1	pET-rbs-F1-XbaI: CCCC<u>TCTAGA</u>AATAATTTTGTTTAAC pET-R1-SacI: GGAA<u>GAGCTC</u>CCTTTCGGGCTTTGTTAG

Figure 1. Schematic domain organization, sequence alignment, growth inhibition assay. (A) Schematic domain organization of Tde1 protein and its variants. The N-terminal repeated glycine zipper motifs (12-51) overlapping a predicted transmembrane domain (22-42) and Ntox15 DNase domain (99-247) are indicated. Tde1 and its variants with truncation or amino acid substitutions were illustrated. (B) Growth inhibition assay of *E. coli* DH10B cells harbouring pTrc200 vector or each of its derivatives expressing Tde1 variants with IPTG induction. (C) Growth inhibition assay of *E. coli* DH10B cells co-expressing the Tde1 variants expressed from pTrc200 plasmid and Tdi1 immunity gene expressed from pRL662 plasmid. Growth curve was determined at OD_{600}. Graphs of panels B and C show mean ± SD of three independent experiments. The significant differences were shown by the different letters (p value <0.01). (D) Multiple sequence alignment of N-Tde1 homologues were presented with highly conserved amino acid residues highlighted in yellow. The bacterial species, strain name, and locus number of Tde1 orthologs (*Agrobacterium/Rhizobium*) or tape measure proteins (*Paraburkholderia/Burkholderia*) are indicated on the left and right of aligned sequences. Two conserved glycine residues (G^{39}, G^{43}) subjected for mutagenesis were indicated by the arrows above the sequences.

Figure 2. Membrane permeabilization assays. (A) Growth inhibition assay of *E. coli* DH10B cells harbouring pTrc200 vector or each of its derivatives expressing Tde1 variants with IPTG inducible expression. BW25113 WT or BW25113 (Δ*lacY)* cells harbouring pTrc200 vector or each of its derivatives expressing Tde1 variants were carried out for (B) β-galactosidase activity assay to determine ONPG uptake, (C) propidium iodide permeability with cells treated with Propidium iodide and Hoechst for detection by fluorescence microscope, and (D) viability assay for *E. coli* cells derived from the ONPG uptake assay. (E) Bacteriostatic activity assay. *E. coli* DH10B cells harbouring pTrc200 vector or each of its derivatives expressing Tde1 variants were cultured with or without IPTG induction for 1 hr. The IPTG-induced cells were further centrifuged and resuspended in the fresh medium with or without IPTG. Cell density was measured again before continuous growth for additional 1 hr. Graphs of panels A, B, and E show mean ± SD of three independent experiments. The significant differences were shown by the different letters (p value <0.01). Panels C and D are representative results of three independent experiments.

Figure 3. The N-terminus of Tde1 is sufficient for interaction with Tap1. (A) Co-immunoprecipitation (Co-IP) in *A. tumefaciens*. *A. tumefaciens* C58 Δ*tde1* harbouring pTrc200 vector or its derivatives expressing HA-tagged Tde1 variants. Anti-HA resin was used to co-IP the Tde1 variants and Tap1. (B) Co-IP in *E. coli* DH10B. DH10B cells co-expressing HA-tagged Tde1 variants from pTrc200 and Tap1 from pRL662 were collected for co-IP. Anti-HA resin was used to co-IP the Tde1 variants and Tap-1. Representative western blot results of three independent experiments were shown with antibody against HA for Tde1 variants and Tap1 respectively. (C). Secretion assay for HA-tagged Tde1 variants. Western blot for the cellular and extracellular fractions of *A. tumefaciens* C58 Δ*tdei* and Δ*tdei1*Δ*tssK* expressing the HA-tagged Tde1 variants. Hcp secretion was detected as a positive control for active T6SS secretion. Representative western blot results of three independent experiments were shown with antibody against HA, Hcp, or EF-Tu serving as a loading and non-secreted protein control. Protein markers are indicated in kDa.

A.

B.

	Phase	mCherry	GFP	Merge1	Merge2
Vector-sfGFP					
C1-Tde1(M)					
N-Tde1					
N-Tde1^{GLGL}					
Tde1(M)					
Tde1(M)^{GLGL}					

Figure 4. The roles of N-Tde1 and G^{39}xxxG^{43} glycine zipper motif for secretion and translocation. (A) Secretion assay for Tde1 variants fused with sfGFP. Western blot for the cellular and extracellular fractions of *A. tumefaciens* C58 Δ*tdei* and Δ*tdei1*Δ*tssK* expressing the Tde1 variants fused with sfGFP were detected by anti-GFP antibody. Representative western blot results of three independent experiments were shown with antibody against GFP, Hcp, or EF-Tu serving as a loading and non-secreted protein control. Hcp secretion served as a positive control for active T6SS secretion. Protein markers are indicated in kDa. (B) Fluorescence microscopy for Tde1 translocation. *A. tumefaciens* C58 Δ*tdei* expressing Tde1 variants fused with sfGFP (in green) and *E. coli* DH10B carrying mCherry (false coloured in blue) were co-cultured for 20 hrs. A cyan fluorescence with merged blue and green signals represented the translocation of Tde1 variants from *A. tumefaciens* to *E. coli*.

Figure 5. G^{39}xxxG^{43} glycine zipper motif of Tde1 is required for DNase-mediated killing of target cells during interbacterial competition. (A) Inter-species Interbacterial competition. *A. tumefaciens* C58 Δ*tdei* and Δ*tdei1*Δ*tssK* expressing the Tde1 variants were co-incubated with *E. coli* (DH10B) cells on AK medium followed by selecting target cell survival with serial dilution on a gentamycin containing LB agar at 37°C. (B) Intra-species Interbacterial competition between *A. tumefaciens* C58 (Δ*tdei*) and Δ*tdei1*Δ*tssK* expressing the Tde1 variants and *A. tumefaciens* 1D1609 target cell. The competition was done as in (B) except 1D1609 was recovered at 28°C. (C) Secretion assay for Tde1 and its variants co-expressed with its immunity protein Tdi1. The toxin immunity pairs Tde1-Tdi1, Tde1^{GLGL}-Tdi1, and Tde1(M)-Tdi1 were expressed in *A. tumefaciens* C58 Δ*tdei* and Δ*tdei*Δ*tssK.* (D) *In vivo* plasmid DNA degradation assay. *E. coli* BW25113 carrying pJN105 empty vector or the derivatives expressing different variants of Tde1 were supplemented with 0.5% glucose ("-") or 0.2% L-arabinose ("+") for 3 hrs to either repress or induce Tde1 production. The plasmids were then extracted to observe the DNA degradation and bottom panel showed western blots of specific Tde1 protein bands indicated by arrows. (E) Growth inhibition assay of Tde1 and its variants. *E. coli* BW25113 cells were induced by adding 0.2% L-arabinose for Tde1 production. The OD$_{600}$ values were measured every 15 minutes. The OD$_{600}$ values of the 4 h post L-arabinose induction were analysed for statistical analysis. Graphs show mean ± SD of three biological repeats. The significant differences were shown by the different letters (p value <0.01). All data were shown with representative results of three independent experiments. Western blots were detected with specific antibody against Tde1, Hcp, or EF-Tu serving as a loading and non-secreted protein control. Protein markers are indicated in kDa.

Figure 6. Proposed model of the loading, firing, and translocation of Tde1. The Tde1 translocation is proposed through six steps. Step 1: Tde1 forms a complex with Tdi1 and Tap1 in the attacker cell. Step 2: Tap1-Tde1-Tdi1 complex binds to the VgrG and the Hcp-VgrG-PAAR puncturing device carrying Tde1 effector complex is loaded onto membrane-associated baseplate. Step 3: Hcp tube and TssB/C sheath polymerize on the Tde1-loaded VgrG/baseplate while Tdi1 and Tap1 fall off with unknown mechanisms before or upon firing. Step 4: TssBC sheath contracts and ejects Tde1 into the target cell periplasm or cytoplasm. Step 5: The glycine zipper(s) on the N-terminus of Tde1 permeabilize the target cell membrane. Step 6: Intact or truncated Tde1 proteins attack DNA for degradation in the target cell.

A.

B.

A. Phage tail tube protein FII   C. Phage tail tape-measure protein   E. Phage_tail_X super family
B. Phage_TAC_7   D. Phage_P2_GpU

Glycine zipper | DNase domain | Uncharacterized protein | Transposase | PhageMin_Tail

Figure S1. Domain architecture and genetic organizations of Tde homologues. (A) Domain organization of the Ntox15-containing proteins. Top 10 classes of the Ntox15-containing proteins are shown with the identifiable domains (not to scale). The number of proteins in each class were indicated on the left based on the information on June 29, 2022. The *A. tumefaciens* Tde1 belonged to the first class where the N-terminal region lacks an identifiable domain. (B) Genetic organizations of genes encoding Tde1 orthologues and Tape Measure Proteins (TMPs) with sequence similarity to N-terminus of Tde1. The proteins encoded from the upstream and downstream of *tde1* and *tmp* genes are shown with their identified domain organizations.

A.

B.

C.

Figure S2. Growth inhibition assays in *A. tumefaciens* and *E. coli*. Growth curve (A) and western blot analyses (B) of *A. tumefaciens* C58 Δ*tde1* carrying pTrc200 or its derivatives expressing HA-tagged Tde1 variants. The growth curve was detected every 2 hrs in 523 medium supplemented with 1 mM IPTG. Graphs show mean ± SD of three independent experiments. The significant differences were shown by the different letters (p value <0.01). The proteins collected at end point (6 hr) were analysed for western blotting with antibody against HA. Representative results of three independent experiments were shown. Protein markers are indicated in kDa. (C) The growth curve analysis of *E. coli* cells used for *in vivo* plasmid DNA degradation assay. The turbidity of *E. coli* BW25113 expressing Tde1 and its variants carried out for the *in vivo* plasmid DNA degradation assay were measured. The *E. coli* cells were supplemented with 0.5% glucose (glu) or 0.2% L-arabinose (ara) for the repression or induction of Tde1 and its variants. The OD_{600} values were measured by DEN-600 photometer (Biosan, Latvia) every hr.

Figure S3. Fluorescence microscopy for negative controls of translocation assay.
A. tumefaciens C58 Δ*tdei*Δ*tssK* expressing N-Tde1-sfGFP or Tde1(M)-sfGFP (in green) and *E. coli* DH10B carrying mCherry (false coloured in blue) were co-cultured for 20 hrs. No cyan fluorescence with merged blue and green signals could be detected when attacker cells are T6SS-inactive, which served as negative controls for the translocation assay.

A.

B.

Tde1 N-terminus
(Tde1: residues 10 – 62)

Pyocin S5
(6THK: residues 446 – 495)

Colicin 1a
(1CII: residues 575 – 620)

Colicin A
(1COL: residues 142 – 199)

Colicin S4
(3FEW: residues 434 – 494)

Colicin B
(1RH1: residues 451 – 509)

C.

D.

Figure S4. Structural prediction of the Tde1 N-terminus with similarity to pyocin S5 and colicins. (A) Predicted results of N-terminal Tde1 (1-97) as a query. (B) N-terminal Tde1 with structural similarity to pore-forming domain of the pyocin S5, colicines and other membrane perturbing proteins based on Phyre2 prediction. (C) Cartoon model of the Tde1 (residue 10 – 62) by using on the basis of the crystal structure of Pyocin S5 (PDB 6THK) with 77.3% of identity. Glycine residues of Tde1 were indicated. (D) Superimposition of N-Tde1 and pore-forming domain of pyocin S5. Tde1 N-terminus is in red and partially pore-forming domain of pyocin S5 is in teal, G^{39} and G^{43} in the putative glycine zipper motif are highlighted in green. All data were analyzed by Phyre2 server.