Title: A systematic method for detecting abnormal mRNA splicing and assessing its clinical

impact in individuals undergoing genetic testing for hereditary cancer syndromes

Authors:

Nick Kamps-Hughes,[1] Victoria E.H. Carlton,[1] Laure Fresard,[1] Steve Osazuwa,[1] Elizabeth

Starks,[1] John J. Vincent,[1] Sarah Albritton,[1] Robert L. Nussbaum,[1] Keith Nykamp[1]*

[1]Invitae Corporation, San Francisco, CA, 94103, USA

*Correspondence:

Keith Nykamp, PhD

1400 16th Street, San Francisco, CA 94103

Ph: (608) 332-9750

Fax: (415) 276-4164

Email: keith.nykamp@invitae.com

Number of text pages: 19

Number of Figures: 4

Number of Tables: 3

Running head: Gauging Impact of altered mRNA splicing

KEYWORDS: RNA sequencing; splicing; variant interpretation; diagnostic testing; splice site

prediction

ABSTRACT

Nearly 14% of disease-causing germline variants result from disruption of mRNA splicing. Most (67%) DNA variants predicted *in silico* to disrupt splicing end up classified as variants of uncertain significance (VUS). We developed and validated an analytic workflow — Splice Effect Event Resolver (SPEER) — that uses mRNA sequencing to reveal significant deviations in splicing, pinpoints the DNA variants potentially responsible, and measures the deleterious effect of the altered splicing on mRNA transcripts, providing evidence to assess the pathogenicity of the variant. SPEER was used to analyze leukocyte RNA encoding 63 hereditary cancer syndrome genes in 20,317 individuals undergoing clinical genetic testing. Among 3,563 (17.5%) individuals with at least one DNA variant predicted to affect splicing, 971 (4.8%) had altered splicing with a deleterious effect on the transcript and 31 had altered splicing due to a DNA variant located outside our laboratory's reportable range. Integrating SPEER results into variant interpretation allowed reclassification of VUS to P/LP in 0.4% and to B/LB in 5.9% of the 20,317 patients. SPEER evidence had a significantly higher impact on allowing P/LP and B/LB interpretations in non-White individuals than in non-Hispanic White individuals, illustrating that evidence derived from RNA splicing analysis may reduce ethnic/ancestral disparities in genetic testing.

INTRODUCTION

Disruption of normal mRNA splicing is a common cause of genetic disease. Approximately 14% of germline pathogenic or likely pathogenic (P/LP) variants cause disease through misplicing[1], with the vast majority (>90%) of these P/LP splicing variants disrupting the canonical donor and acceptor splice sites (first 1-2 bp flanking an exon) (referred to as CSS). In contrast, many DNA variants other than in the CSS dinucleotides are predicted by commonly used algorithms to

affect splicing, but most (69%) of these potential splicing variants (PSpVs) are classified as

variants of uncertain significance (VUS)[1], demonstrating the challenge of classifying non-CSS

PSpVs in the absence of confirmatory RNA testing. Knowing which of these PSpVs have a

clinically significant impact on splicing could inform variant interpretation, resulting in fewer VUS

and increased actionability for genetic testing.

Here, we describe a novel method, the Splice Effect Event Resolver (SPEER), that uses patient

mRNA sequencing to determine whether variants found during DNA sequencing cause aberrant

mRNA splicing and have a deleterious impact on transcript structure and function. SPEER

accomplishes this task in three steps, first by determining whether there is abnormal splicing

compared to controls, then by quantifying by how much normal splicing at a critical splice

junction has been reduced, and finally by examining the consequences of abnormal splicing

events on mRNA structure and function. SPEER begins with short-read sequencing of cDNA

and several overlapping RNA analysis methods[2–6] to analyze exon-exon junctions and

determine whether observed alterations in the mRNA splicing pattern of a gene containing a

PSpV are statistically significant relative to a panel of control individuals. In this way, SPEER

distinguishes alterations in splicing from the naturally occurring alternative splicing that

generates protein diversity and functional specificity across tissue types.[7–9] Next, it examines

the canonical splice junction or junctions most likely impacted by the PSpV and quantifies the

extent of loss of normal splicing in the patient compared to normal controls. Finally, SPEER

assesses how deleterious the impact of the abnormal splicing of particular mRNA transcripts

would be on mRNA stability and function by collecting all the alterations in the transcripts

together into an "abnormal splicing event group", which is then evaluated for reading frame

shifts and potential nonsense mediated decay. By assessing the deleterious impact of a DNA

variant on transcript structure and function, the evidence generated by SPEER can be used for

variant interpretation within a system, such as Sherloc [10], that combines many other types of

evidence to ultimately assess pathogenicity of a variant.

In this manuscript, we report how we validated SPEER's ability to detect aberrant splicing

events and then applied it to a cohort of more than 20,000 individuals undergoing clinical testing

of hereditary cancer syndrome genes to identify which PSpVs detected during DNA sequencing

significantly altered splicing and had a deleterious effect on transcript structure and function.

SPEER also detected abnormal splicing changes in a small number of patients, which allowed

for the detection and confirmation of DNA variants located outside the reportable range of our

next-generation sequencing (NGS) DNA test. When SPEER evidence was used in Sherloc for

variant interpretation, a significant fraction of PSpVs could be interpreted definitively as

Pathogenic/Likely Pathogenic (P/LP) or Benign/Likely Benign (B/LB); most notably, SPEER

allowed an even greater rate of definitive classification of PSpVs among individuals who self-

reported as non-White compared to those who self-reported as non-Hispanic White.

MATERIAL AND METHODS

Cohorts

In phase 1 of this two-phase study, we ascertained a retrospective cohort of 532 research

participants to be used for SPEER validation. Of these 532 participants, 342 had prior genetic

testing for hereditary cancer syndromes performed at Invitae and had a germline DNA variant

that was known or predicted to alter splicing (PSpV), and 190 did not have a PSpV, but had a

personal or family history that was strongly suggestive of a particular hereditary cancer

syndrome, e.g. Familial Adenomatous Polyposis (McK #175100). Informed written consent was

obtained under a protocol approved by the WCG Institutional Review Board (#20190811). As a

reference panel, we used 273 anonymous samples from presumed healthy male and female

blood donors (BioIVT, NewYork) that we oversampled for self-reported ancestries

underrepresented in genetic studies, including African American (51.2%) and Hispanic (30.4%)

individuals. This sampling strategy was designed to guard against mistakenly inferring aberrant

splicing when comparing splicing in non-White versus White individals by failing to take into

account potential variation in splicing of germline transcripts among individuals of different

ancestries, as has been seen in tumor samples in The Cancer Genome Atlas [11].

In Phase 2, a prospective cohort of 20,317 individuals referred to Invitae for germline hereditary

cancer gene testing underwent paired DNA and RNA sequencing between July 2021 and May

2022. SPEER was used to analyze the leukocyte RNA of 63 hereditary cancer genes in these

individuals. Use of de-identified samples and data was approved by an independent institutional

review board (WCG IRB #20161796).

Selection of genes for RNA analysis

A total of 85 hereditary cancer genes were initially considered for RNA analysis (Table S1).

Twenty-two of these were ultimately excluded from the final assay because 1) they are

expressed at too low a level in leukocyte transcripts to allow statistically significant

demonstration of a splicing change, 2) they are genes for which loss-of-function, the usual effect

of abnormal splicing, is not a known mechanism for an increased risk for cancer, or 3) they have

only a single known loss-of-function missense variant associated with an increased risk for

cancer.

DNA Sequencing

NGS of gene panels were performed as previously described [1,12,13] using oligonucleotide baits

(Twist Bioscience, South San Francisco, CA; Integrated DNA Technologies, Coralville, IA) to

capture coding exon sequences ± 20 bases of flanking intronic sequences, and certain non-

coding regions of clinical interest, defined as our reportable range (RR). Targeted regions were

sequenced to a minimum depth of 50x and an average depth of 350x read coverage at each

nucleotide position within the RR. Full gene sequencing, deletion/duplication analysis, and

variant interpretation were performed at Invitae (San Francisco, CA), as previously described

[10,13].

Identification of potential splicing variants (PSpV) in DNA

135   PSpVs were predicted initially using MaxEntScan[14], SpliceSiteFinder-Like[15], and the Alamut

136   Splicing Module (Interactive Biosoftware, Rouen, France, version 1.4.4_2016.02.03) and later

137   using SpliceAI [16].

RNA Sequencing

RNA was extracted from whole blood in PaxGene RNA tubes (762165, BD, New Jersey)

containing additives that inhibit RNA degradation, including nonsense mediated decay (NMD),

using RNAdvance Blood (A35604, Beckman Coulter, Indianapolis) and quantified by

fluorometry. Residual DNA contamination was removed by DNaseI treatment (M0303L, New

England Biolabs, Massachusetts). Indexed cDNA libraries were prepared using the KAPA hyper

RNA kit (KK8541, Roche, Switzerland), substituting proprietary adapters and indexing primers.

Transcripts of interest were enriched via hybridization with custom-designed biotinylated

oligonucleotide baits (Integrated DNA Technologies, Iowa) followed by Streptavidin bead

capture. Illumina compatible P5 P7 sequences were added during post capture amplification

and libraries were submitted for 2 x 150bp paired-end next generation sequencing (Illumina,

San Diego, CA), targeting 15 million clusters per sample. Adapters were trimmed sequencing

reads and UMIs were extracted. Trimmed reads were aligned on the GRCh37 human genome

assembly with STAR [17], using a RefSeq- based transcript annotation. Duplicate reads were

removed using unique molecular identifier sequences (umi-tools,[18]). Raw split read counts were

extracted from the deduplicated BAM files using Leafcutter scripts sam2bed.pl and bed2junc.pl

[2].

Detection of statistically significant splicing changes

SPEER uses targeted sequencing of RNA isolated from blood samples to determine splice

junction counts across 63 genes of interest (Table S1). To assess the statistical significance of

splice junction usage in a patient sample, SPEER (Figure 1 A,B) measured usage of known and

novel splice junctions using "percent spliced in" (PSI)[19,20]. The PSI of each junction is defined

as the ratio of the number of reads supporting the junction divided by the total number of reads

overlapping the junction (i.e., supporting reads plus non-supporting reads). SPEER assesses

the statistical significance of each splicing change in a patient sample by comparing it to the

splice junction usage from the 273-sample reference panel using a beta-binomial test described

below.. PSI changes with $p \leq 10^{-3}$ were considered statistically significant. SPEER uses these

significant junctions as seeds to recursively find other altered ($p \leq 0.05$) junctions that share

breakpoints in order to link all junctions participating in the localized splicing event into an

"abnormal splicing event group" for downstream interpretation.

Beta-binomial model description and parameter estimation

For every junction observed in at least one control, a PSI normal model was calculated using

the beta-binomial distribution. In this model, the number of reads overlapping the junction was

considered the number of trials and the reads supporting the junction was considered the

number of successes. For each junction, the parameters nu (approximately the mean PSI) and

rho (a measure of the increased variance relative to the binomial distribution) were fit to their

maximum likelihood values and stored as a panel of normal splicing bed file. The control

parameters for each junction were then compared to the corresponding PSI for each patient

sample to generate a p-value using a beta-binomial likelihood ratio test. This statistical

approach was inspired by the ShearwaterML algorithm for detecting low frequency somatic DNA

variants from NGS data [21].

Quantifying the loss of the normal transcript directly impacted by a PSpV

Each abnormal splicing event group observed in a patient sample included a reduction in the

PSI for one or more canonical splice junction and an increase in PSI for one or more newly

created or non-canonical splice junction (see Figure 1A). To measure the reduction in normal

splicing associated with a PSpV, we calculated the ratio of the PSI for the canonical junction

determined to be directly impacted by the PSpV over the PSI for that same junction in the

control panel. This metric, which we define as PSI-X, is expected to correlate with loss of the

normal or most commonly expressed transcript. Moreover, a significant reduction in PSI-X is

expected to correlate with pathogenicity for PSpVs in genes known to cause disease with an

autosomal dominant, loss-of-function mechanism.

Directly impacted canonical junctions were defined as the following: for intronic PSpVs, we

chose the canonical junction that was normally created by splicing out the intron that contained

the PSpV; for exonic PSpVs we chose the canonical junction with the highest PSI in the control

panel that also shared a breakpoint with a newly created, non-canonical junction. If the highest

PSI value for a canonical junction was less than 2x other canonical junctions with shared

breakpoints, the PSI values were averaged across all impacted canonical junctions.

Assessing whether abnormal splicing events impact translation and function of the

protein

Abnormal splicing event groups with a reduction in PSI-X that had at least one breakpoint within

50 bp of a PSpV were flagged for review by trained variant scientists (Supplemental Methods).

This was an essential step for assessing whether the transcript created by non-canonical

junctions may compensate for the reduction in normally spliced mRNA. As an example, a PSpV

that abolishes a well annotated acceptor splice may result in substantially decreased PSI-X

while also activating a cryptic acceptor splice site three nucleotides upstream, resulting in a

novel protein with a single amino acid insertion. It's uncertain whether this novel protein will

compensate for the loss of the normally expressed protein, thus warranting a VUS classification

for the PSpV. The first step in this review was to label each splice junction in an event group

with one of the following tags: (1) *canonical*, a junction formed by adjacent exons in the full-

length canonical transcript); (2) *exon skipping*, defined as non-adjacent junctions with one more

exons being skipped; (3) *partial exon exclusion,* in which a portion of a canonical exon was not

included in the transcript; (4) *partial intron inclusion,* in which a portion of an intron was included

in a transcript; (5) *cryptic exon,* when two non-canonical junctions flanked an intronic sequence.

All event groups, other than complex*,* were labeled according to whether the primary non-

canonical junction is expected to cause nonsense-mediated decay (NMD). Event groups with a

single non-canonical junction causing a frameshift and/or premature termination codon

upstream of the last coding exon were labeled as *NMD+.* Those expected to result in in-frame

insertions anywhere in the transcript, or in deletions or premature termination codons

downstream of the last exon junction, were labeled as *NMD-*. If an event group had multiple

non-canonical junctions, with different predicted effects on protein translation, the event group

was labeled as *complex*.

Detection and classification of variants outside our test's reportable range

Rare variants located outside our test's reportable range (all coding exons plus 20 bp flanking

each exon) are missed by standard panel-based NGS tests. When SPEER identifies an

abnormal splicing event group at the variant discovery significance threshold (i.e., $p \leq 10e^{-5}$) *and*

one or more abnormal junctions exhibit a fold-change in PSI ≥10,000, we manually examine the

primary NGS data for a DNA variant that may explain the abnormal splicing event. If none is

found in the primary data, we perform long-read DNA sequencing (PacBio, CA) throughout the

region surrounding the abnormal splicing event. If a PSpV is identified in the region surrounding

the abnormal splicing event, it is flagged for interpretation by the variant scientist team. If a

PSpV could not be identified, the abnormal splicing change was presumed to be a technical

false positive or the result of natural variation in splicing and not included in the clinical report.

Integrating deleterious RNA splicing changes as evidence in DNA variant classification

In order to standardize the application of SPEER evidence for variant classification, we created

a new category of evidence within our semi-quantitative variant classification framework,

Sherloc[10]. We referred to this evidence category as *Observed RNA Effects* (see Table S2).

Each criterion within this new evidence category was assigned pathogenic (P) or benign (B)

point scores. Point scores were assigned and calibrated by applying Observed RNA Effects

criteria to more than 300 PSpVs variants observed in the retrospective cohort of 532 research

participants in Phase 1. Final classifications in Phase 2 of the study were assigned based on the

sum of all available evidence, including the Observed RNA Effects criterion when applicable.

Rates of PSpV upgrades (VUS to LP/P) and downgrades (VUS to LB/B) observed during phase

2 of the study were calculated by removing Observed RNA effect criteria and asking whether

the final classification (and Sherloc score) moved from P/LP or B/LB to a VUS. Differences in

these classification rates among self-reported ethnicities were tested using a one-sided Fisher

Exact Test by comparing the proportion of individuals with downgrades and upgrades for each

ethnicity (Black/African-American, Hispanic, Asian and Other) to the proportion of individuals

from the most common self-reported ethnicity (non-Hispanic White).

RESULTS

SPEER validation

SPEER was validated using 50 positive control samples containing 46 unique DNA variants as

follows: 10 samples with 7 unique variants well documented to alter splicing according to

published literature; 13 samples with 12 unique variants in the essential GT or AG dinucleotides

but without published functional evidence; and 27 samples with 27 unique variants located

within 50 nucleotides of a significant abnormal splicing event ($p \leq 0.001$). False positives were

any significant abnormal splicing event ($p \leq 0.001$) detected throughout the 63-gene panel for a

canonical junction not within 50 bp of a known PSpV. True negatives (TN) were derived from

two sets of samples. The first set were the same 50 samples that had served as *true positives*

for one junction-variant pair, repurposed as a source of *true negatives* by analyzing all other

splice junctions lacking a nearby PSpV. The second set of true negative samples consisted of

16 samples with a PSpV that nonetheless yielded a normal splicing pattern based on a manual

review of RNA sequence alignments to establish the absence of a splicing change. Combining

the two sets yielded 66 true negative (TN) samples suitable for calculating sensitivity, specificity,

accuracy, reproducibility of SPEER's findings between replicates of the same sample analyzed

in parallel and in the same sample over time (Figure 1 C,D and Table S3).

SPEER detected a significant splicing change in 45 of 46 positive control DNA variants in the 50

positive control samples (97.8% sensitivity; Figure 1C) while maintaining a specificity of 99.7%.

We used two of the remaining four positive control samples, along with eight of the negative

control samples, to perform reproducibility and repeatability testing in triplicate across all 10

samples. All 30 (100%) of the reproducibility samples and 29 (96.7%) of 30 repeatability

samples yielded the expected results (Figure 1C). True positive and true negative calls

demonstrated strong separation at the p-value threshold used for detecting splicing effects

(Figure 1D), and the same variants found in multiple samples demonstrated consistent PSI

values (data not shown). SPEER also enabled the detection of variants outside of the reportable

range in four individuals from the phase 1 research cohort, all of whom had presented with a

persuasive personal and/or family history of hereditary cancer syndrome.

Characterization of DNA variants associated with abnormal splicing event groups

SPEER detected abnormal splicing events associated with PSpVs with very high accuracy

(99.7%). The next step was to calculate PSI-X as a measure of the loss of normal splicing at the

canonical junction most relevant to the PSpV compared to controls. We plotted the distribution

of PSI-X for variants previously classified as P/LP due to a splicing abnormality, and therefore

could be confidently considered to have a deleterious effect on the transcript, and those known

to be B/LB and therefore could be considered to have no deleterious effect on splicing (Figure

2A). We computed a receiver operating characteristic (ROC) curve (Figure 2B) for these data

that resulted in an area under the ROC curve (AUROC) of 0.91 for PSI-X, demonstrating that

PSI-X has excellent discriminatory power for separating PSpVs known to be either deleterious

or not, inferred from their status as either P/LP or B/LB.

As another line of evidence to validate the PSI-X metric for quantifying the effect of a variant on

reducing splicing of the junction most relevant to that variant, we assessed PSI-X for samples in

which a heterozygous DNA deletion encompassing one or more exons that would therefore

represented complete loss of junction usage for the flanking splice sites. With these deletion

variants we found a distribution in PSI-X from 0.3 to 0.75 (Figure 2C), suggesting that a PSI-X

<0.8 was strongly correlated with significant reduction in the usage of a canonical splice junction

from one allele.

We then sought to assess how well PSI-X alone could predict what would be the ultimate

clinical interpretation of a PSpV as either P/LP or B/LB. If all PSpVs with a PSI-X <0.8 are

deleterious to mRNA transcript structure and function and therefore might be naively assumed

to be P/LP, and all PSpVs with a PSI-X ≳0.8 were considered non-deleterious and therefore

taken to be B/LB, a comparison of predicted interpretation based on PSI-X with the known

pathogenicity status of these variants would be 100% accurate. This is not the case as PSI-X

alone had an accuracy of 86% for the clinical interpretation of variants. Clearly, not all

deleterious changes in mRNA structure and function must be pathogenic and not all variants

that leave splicing intact are necessarily benign. We therefore included the likelihood that an

abnormal splicing event group would lead to NMD, thereby affecting mRNA stability and

expression of the translated protein product, in the analysis of PSpVs with PSI-X <0.8. In this

way, we recognized that six of nine known B/LB variants with a PSI-X <0.8 affect transcript

structure but are unlikely to affect gene function, since the abnormal transcripts created were

not expected to result in NMD; five event groups resulted in small in-frame (≤ 6 amino acids)

deletions or insertions, while one event group resulted in alternative usage of the first coding

exon. By combining a requirement that the PSI-X be <0.8 *with* the requirement that the

abnormal transcripts lead to NMD, we could achieve a positive predictive value (PPV) for PSI-X

<0.8 of 98.9% for P/LP variants, sufficiently high to be used as evidence for pathogenicity in

Sherloc. In contrast, among 42 PSpVs with PSI-X > 0.8, 9 variants were previously known to be

P/LP presumably because they disrupted the protein directly through coding sequence changes

(i.e., nonsense, missense, in-frame deletion or insertion) and not through abnormal splicing.

Therefore, by combining a PSI-X ≥0.8 with the requirement that the PSpV not be a missense

variant or directly change the coding sequence, we could improve the negative predictive value

(NPV) of PSI-X > 0.8 for the variant being B/LB to 61.2%, which we did not consider sufficiently

high to be used as evidence for the variant being benign in the Sherloc classification framework.

More work needs to be done to increase the specificity of PSI-X ≥0.8 for B/LB variants.

Prospective analysis of paired DNA and RNA testing

In Phase 2 of the study, 20,317 individual blood samples were processed for DNA sequencing

and for RNA extraction and sequencing. Across the entire population, 3,563 (17.5%) of the

patients had at least one DNA variant identified by NGS that was a PSpV (Table 1). Abnormal

splicing, defined by a statistically significant event group, coincided with 605 unique PSpVs

(Table 2, Row 1) in a total of 971 patients (4.8% of all prospective samples) (Table 1). These

605 PSpVs were detected in 54 of the 63 genes analyzed by SPEER (Table S4). We observed

normal splicing associated with a PSpV in 2,487 patients (73% of patients with a PSpV),

illustrating the high false positive rate for the algorithms and heuristics used to identify PSpVs.

By evaluating the location of PSpVs in relation to the exonic sequence, we found that the largest

number of P/LP variants relative to VUS were within the CSS (-1, -2, +1, +2; Figure 3A). In

addition, there was an enrichment at positions +3, +4, and +5 from the canonical donor splice

site, and a cluster at -11, -10 and -9, upstream of the canonical acceptor site that most often

creates a novel acceptor splice site. Overall, we observed abnormal splicing events associated

with 605 PSpVs (Table 2), with 93% of these PSpVs located within our test's reportable range of

+/- 20 bp flanking the exons.

Classification of abnormal splicing events in the prospective cohort

Each PSpV associated with an abnormal splicing event group identified by SPEER was

classified using Observed RNA Effects evidence within the Sherloc interpretation framework.

Overall, half (56.7%) dof predicted splicing variants (PSpVs) observed in this study have been

classified as pathogenic or likely pathogenic (P/LP; 20.2%) and benign or likely benign (B/LB;

36.5%), while the remaining (43.3%) were classified as variants of uncertain significance (VUS)

(Table 2, row 3). As expected, abnormal splicing correlated with a much higher rate of P/LP

variants (42.9%) and much lower rate of B/LB variants (11.6%) (Table 2, row 1). Conversely,

normal splicing correlated with a much lower rate of P/LP variants (9.4%) and much higher rate

of B/LB variants (48.3%) (Table 2, row 2). Importantly, we found that 6.3% of all patients from

the phase 2 cohort had a P/LP or B/LB variant that would have been a VUS by our interpretation

framework if it were not for the added RNA splicing data from our study: 75 patients (0.4% of all

samples) would have received a VUS instead of a P/LP classification, and 1,204 patients (5.9%

of all samples) would have received a VUS instead of a B/LB classification (Table 1). When

comparing these data across self-reported ancestries, we found that individuals with self-

reported Hispanic (9.4%, p=8.2e-12), Black/African American (10.4%, p=1.8e-16), or Asian

(9.1%, p=1.4e-5) ancestries had statistically significantly higher rates of definitive classifications

(P/LP or B/LB) than Non-Hispanic White populations (4.8%) due to the addition of supplemental

RNA splicing data (Table 3).

The most abundant splicing change in the prospective cohort was exon skipping, with partial

exon exclusion and partial intron inclusion events being the next most abundant events. Partial

exon exclusion, partial intron inclusion, and complex events displayed the largest discrepancies

between PSpV definitive classifications (i.e., P, LP, LB, B) and VUS, indicating that these were

the most challenging event groups to interpret (Figure 3B).

Detection of variants affecting splicing outside the reportable range

Among the 190 individuals ascertained in the initial research cohort in Phase 1 specifically

because no P/LP variants were found through routine clinical testing despite a personal or

family history highly suggestive of a hereditary cancer syndrome, 4 (2.1%) were found to have a

variant outside our reportable range with a deleterious effect on splicing. In the prospective

consecutive cohort of 20,317 patients referred for hereditary cancer testing, 42 (0.2%) patients

carried 35 unique variants outside of the reportable range that could be associated with

abnormal splicing (Table 1). Of these 35 variants, 6 (17.1%), seen in nine patients, were

classified as P/LP, representing a 0.04% yield of P/LP variants not present in the reportable

range of our test in the entire cohort (9 of 20,317 patients). An additional two variants, seen in

two patients (4.8%), were classified as benign, and 27 of the 35 unique variants, present in 31

patients (73.8%), were classified as VUS (Figure 4).

DISCUSSION

We describe SPEER, an accurate and scalable approach for detecting and quantifying splicing

alterations in peripheral blood leukocyte transcripts of 63 hereditary cancer syndrome genes in

patients undergoing genetic testing for these syndromes. We demonstrate that adding RNA

analysis to DNA sequencing of hereditary cancer syndrome genes facilitates the interpretation

of DNA variants expected or predicted to alter splicing. In addition, of the 605 unique variants

identified in the prospective cohort that affected splicing, 35 of them (5.8%) were located outside

the reportable range of +/- 20 bp flanking the exons of our NGS panel tests and were detected

by SPEER.

We found that supplemental RNA analysis led to a definitive classification of DNA variants that

would have been classified as a VUS without the RNA analysis in 6.3% of individuals in a

prospective cohort undergoing testing for hereditary cancer; of these, 0.4% received a P/LP

classification while 5.9% of patients tested had a B/LB variant, thereby effectively reducing VUS

rates. This fraction of patients with reclassifications due to RNA analysis in our 20,317

prospective cohort was higher (6.3%) than what was recently reported in a cohort of 43,524

patients undergoing DNA and RNA testing for hereditary cancer (1.3%) [22]. This difference is

likely because of the larger number of genes (63 genes) tested by RNA in our cohort compared

to the recently published cohort (18 genes). Importantly, the rate of definitive classifications (i.e.,

P, LP, LB, B) dependent on RNA analysis was significantly higher in Black/African American,

Asian, and Hispanic individuals combined than in non-Hispanic White individuals. The

preferential impact of RNA analysis on VUS reclassification in these less-studied populations

probably resulted from RNA analysis providing direct evidence that was not limited by the

insufficient representation of individuals of non-European origin in variant databases or in the

published literature. This indicates that including RNA analysis in the interpretation of DNA

variants predicted to impact splicing may preferentially reduce VUS rate for less-studied

ancestral and ethnic populations, thereby reducing health disparities.

These results also highlight the nuances and difficulties of incorporating sample-based RNA

splicing analysis into variant interpretation frameworks used in routine genetic testing.

Alternative splicing is very common, and it can be challenging to pinpoint specific splicing

alterations associated with a hereditary disease. For example, greater than 96% of all patient

samples in our prospective cohort had at least one statistically significant splicing change at the

$p \leq 0.001$ threshold compared to the reference panel of normal samples. This statistical

threshold was not multiple hypothesis-corrected because it was meant to accurately determine

whether a specific PSpV altered normal splicing. By focusing scientific review on splicing

alterations with a PSpV within 50 bp of the nearest altered junction, we reduced the frequency

of abnormal splicing event groups to 4.8% of all patient samples, while maintaining 99.7%

accuracy for detecting abnormal splicing events previously known to be associated with DNA

variants in our validation of SPEER.

Even with supplemental RNA splicing data, the biological impact and clinical relevance of DNA

variants associated with a significant splicing alteration are difficult to assess, often still resulting

in VUS classifications. This is particularly true for variants found outside of the reportable range

of the test since they are often observed in a single individual and lack the additional

corroborating evidence needed to make a definitive classification, such as population frequency

data in publicly available databases. Only 9 of 20,317 patients (0.04%) in our prospective cohort

had variants outside our +/-20 bp reportable range, caused altered splicing, and would

ultimately be classified as P/LP. This rate of P/LP variants that occur >20bp into the introns is

consistent with the rate (0.06% or 28/43,524) seen in another large RNA cohort undergoing

testing for hereditary cancer [22]. Another 2 of 20,317 patients had variants outside our +/-20 bp

reportable range, caused altered splicing, but still were ultimately classified as B/LB,

underscoring the fact that a variant that has a deleterious effect on splicing may still not cause

disease.

Another complicating factor in connecting a deleterious effect on splicing to pathogenicity is that

abnormal splicing can be incomplete. A heterozygous DNA variant may result in a reduction of

normal mRNA splicing that ranges from 0% (i.e., no reduction) to 50% (i.e., complete loss of

normal splicing from the variant allele), with everything in between considered partial loss, also

referred to as 'leakiness[23,24].Importantly, because a PSpV that leads to a complete loss of

normal splicing is more likely to cause disease than a PSpV that leads to a partial loss, accurate

classification of PSpVs using abnormal splicing data requires a confident measure of the

amount of reduction. We developed such a measurement to quantify the reduction in normal

mRNA splicing — PSI-X — and determined that splicing alterations in blood samples associated

with a PSI-X <0.8 correlate with sufficient loss of normal splicing from one allele to cause

disease and qualify as pathogenic evidence for autosomal dominant hereditary cancer

syndromes within our variant classification framework. Although a PSI-X threshold of 0.8

demonstrates high discriminatory performance (AUROC = 0.91) for PSpVs with abnormal

splicing events, we still find that 15% of B/LB variants fall below the threshold and 14% of P/LP

variants are above the threshold. To achieve a level of confidence in the final classification of

PSpVs associated with abnormal splicing suitable for clinical testing, it is essential that trained

variant scientists review the expected effect of abnormal splicing on translation and protein

function, and then integrate these findings into a DNA variant interpretation framework. In this

way, we demonstrated that by pairing DNA and RNA testing, we can increase the number of

patients who have an actionable finding while achieving a PPV of 98.9%. On the other hand,

even with a careful review of the abnormal event groups associated with PSI-X $\gtrsim$0.8, it was still

difficult to discriminate between P/LP and B/LB variants, as demonstrated by a much lower NPV

(61.2%) when PSI-X is$\gtrsim$0.8. As a result, to limit the likelihood of a false negative result for our

clinical testing patients, we chose not to award benign points in the Sherloc framework even for

statistically significant abnormal splicing event groups when PSI-X $\gtrsim$0.8. PSpVs associated with

these higher PSI-X values will generally be VUS unless complementary evidence (i.e., family

segregation, population frequency, additional splicing studies) support a more definitive

classification.

Although this study clearly demonstrates that RNA splicing data can aid in variant classification

and identification, the results should be interpreted in the context of the following technical and

biological limitations. First, the 63 hereditary cancer genes included in this study are tumor

suppressor genes that are not highly expressed in whole blood leukocyte RNA. We expect each

gene to have tissue-specific expression and alternatively spliced products that differ from what

can be detected in a blood sample. As a result, abnormal splicing observed from whole blood

leukocytes may not reflect a biological change in the tissue of interest, while the absence of a

splicing alteration in whole blood may not always correlate with an absence of abnormal splicing

in the disease-relevant tissue. Second, the SPEER algorithm is designed to evaluate changes in

junction usage. A small fraction of splicing alterations result in complete intron retention, which

escapes detection by the SPEER algorithm in its current form. This likely explains why, in our

study, several known P/LP DNA variants had PSI-X $\gtrsim 0.8$ with transcripts that underwent NMD.

Further development is underway to include a method for evaluating the relative frequency of

complete intron inclusion in the SPEER algorithm, which will better resolve the uncertainty of

variants with high PSI-X. Third, although PSI values and aberrant junctions were highly

consistent for the same variant across multiple samples, a few DNA variants showed high levels

of variability. Most of these correlated with the presence of *cis*-acting elements, such as

secondary DNA variants, suspected of modifying the impact of the primary DNA variant on

splicing. Currently it is not possible to predict *a priori* which splicing changes are impacted by

modifiers, but with more samples sequenced for the same DNA variants we may be able to

identify patterns in the DNA sequence that modulate the PSI and alternative splicing events

within these tumor suppressor genes.

In summary, RNA analysis in whole blood is a valuable tool for finding and assessing the impact

of DNA variants suspected of affecting RNA splicing, thereby altering gene expression. It helps

to reduce VUS rates in a fraction of individuals who have DNA variants predicted to affect

splicing, and supports the identification of deleterious, possibly pathogenic variants outside the

test's reportable range. However, interpretation of the splicing information is challenging and

there are limitations in the techniques and in our biological understanding of splicing. By

combining comprehensive DNA and RNA sequencing with the validated SPEER algorithm, we

will continue to collect data individuals referred for hereditary cancer syndrome testing, develop

deeper insights into the clinical relevance of splicing alterations, and improve the interpretation

of inherited DNA variants — leading to a further reduction in uncertainty for hereditary cancer

gene testing.

REFERENCES

1.   Truty R, Ouyang K, Rojahn S, Garcia S, Colavin A, Hamlington B, Freivogel M, Nussbaum

RL, Nykamp K, Aradhya S. Spectrum of splicing variants in disease genes and the ability of

RNA analysis to reduce uncertainty in clinical interpretation. Am J Hum Genet, 2021,

108:696–708

2.   Li YI, Knowles DA, Humphrey J, Barbeira AN, Dickinson SP, Im HK, Pritchard JK.

Annotation-free quantification of RNA splicing using LeafCutter. Nat Genet, 2018, 50:151–8

3.   Frésard L, Smail C, Ferraro NM, Teran NA, Li X, Smith KS, Bonner D, Kernohan KD,

Marwaha S, Zappala Z, Balliu B, Davis JR, Liu B, Prybol CJ, Kohler JN, Zastrow DB,

Reuter CM, Fisk DG, Grove ME, Davidson JM, Hartley T, Joshi R, Strober BJ, Utiramerur

S, Lind L, Ingelsson E, Battle A, Bejerano G, Bernstein JA, Ashley EA, Boycott KM, Merker

JD, Wheeler MT, Montgomery SB. Identification of rare-disease genes using blood

transcriptome sequencing and large control cohorts. Nat Med, 2019, 25:911–9

4.   Mertes C, Scheller IF, Yépez VA, Çelik MH, Liang Y, Kremer LS, Gusic M, Prokisch H,

Gagneur J. Detection of aberrant splicing events in RNA-seq data using FRASER. Nat

Commun, 2021, 12:529

5.   Jenkinson G, Li YI, Basu S, Cousin MA, Oliver GR, Klee EW. LeafCutterMD: an algorithm

for outlier splicing detection in rare diseases. Bioinformatics, 2020, 36:4609–15

6.   Vaquero-Garcia J, Barrera A, Gazzara MR, González-Vallinas J, Lahens NF, Hogenesch

JB, Lynch KW, Barash Y. A new view of transcriptome complexity and regulation through

the lens of local splicing variations. Elife, 2016, 5:e11752

7.   Sibley CR, Blazquez L, Ule J. Lessons from non-canonical splicing. Nat Rev Genet, 2016,

17:407–21

8.   Baralle FE, Giudice J. Alternative splicing as a regulator of development and tissue identity.

Nat Rev Mol Cell Biol, 2017, 18:437–51

9. Ule J, Blencowe BJ. Alternative Splicing Regulatory Networks: Functions, Mechanisms, and
Evolution. Mol Cell, 2019, 76:329–45

10. Nykamp K, Anderson M, Powers M, Garcia J, Herrera B, Ho Y-Y, Kobayashi Y, Patil N,
Thusberg J, Westbrook M, Invitae Clinical Genomics Group, Topper S. Sherloc: a
comprehensive refinement of the ACMG-AMP variant classification criteria. Genet Med,
2017, 19:1105–17

11. Al Abo M, Hyslop T, Qin X, Owzar K, George DJ, Patierno SR, Freedman JA. Differential
alternative RNA splicing and transcription events between tumors from African American
and White patients in The Cancer Genome Atlas. Genomics, 2021, 113:1234–46

12. Truty R, Paul J, Kennemer M, Lincoln SE, Olivares E, Nussbaum RL, Aradhya S.
Prevalence and properties of intragenic copy-number variation in Mendelian disease genes.
Genet Med, 2019, 21:114–23

13. Kurian AW, Hare EE, Mills MA, Kingham KE, McPherson L, Whittemore AS, McGuire V,
Ladabaum U, Kobayashi Y, Lincoln SE, Cargill M, Ford JM. Clinical evaluation of a
multiple-gene sequencing panel for hereditary cancer risk assessment. J Clin Oncol, 2014,
32:2001–9

14. Yeo G, Burge CB. Maximum entropy modeling of short sequence motifs with applications to
RNA splicing signals. J Comput Biol, 2004, 11:377–94

15. Shapiro MB, Senapathy P. RNA splice junctions of different classes of eukaryotes:
sequence statistics and functional implications in gene expression. Nucleic Acids Res,
1987, 15:7155–74

16. Jaganathan K, Kyriazopoulou Panagiotopoulou S, McRae JF, Darbandi SF, Knowles D, Li
YI, Kosmicki JA, Arbelaez J, Cui W, Schwartz GB, Chow ED, Kanterakis E, Gao H, Kia A,
Batzoglou S, Sanders SJ, Farh KK-H. Predicting Splicing from Primary Sequence with
Deep Learning. Cell, 2019, 176:535–48.e24

17. Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, Batut P, Chaisson M,

Gingeras TR. STAR: ultrafast universal RNA-seq aligner. Bioinformatics, 2012, 29:15–21

18. Smith T, Heger A, Sudbery I. UMI-tools: modeling sequencing errors in Unique Molecular

Identifiers to improve quantification accuracy. Genome Res, 2017, 27:491–9

19. Katz Y, Wang ET, Airoldi EM, Burge CB. Analysis and design of RNA sequencing

experiments for identifying isoform regulation. Nat Methods, 2010, 7:1009–15

20. Schafer S, Miao K, Benson CC, Heinig M, Cook SA, Hubner N. Alternative splicing

signatures in RNA-seq data: Percent spliced in (PSI). Curr Protoc Hum Genet, 2015,

87:11.16.1–11.16.14

21. Martincorena I, Roshan A, Gerstung M, Ellis P, Van Loo P, McLaren S, Wedge DC, Fullam

A, Alexandrov LB, Tubio JM, Stebbings L, Menzies A, Widaa S, Stratton MR, Jones PH,

Campbell PJ. Tumor evolution. High burden and pervasive positive selection of somatic

mutations in normal human skin. Science, 2015, 348:880–6

22. Horton C, Cass A, Conner BR, Hoang L, Zimmermann H, Abualkheir N, Burks D, Qian D,

Molparia B, Vuong H, LaDuca H, Grzybowski J, Durda K, Pilarski R, Profato J, Clayback K,

Mahoney M, Schroeder C, Torres-Martinez W, Elliott A, Chao EC, Karam R. Mutational and

splicing landscape in a cohort of 43,000 patients tested for hereditary cancer. NPJ Genom

Med, 2022, 7:49

23. Schröder S, Wieland B, Ohlenbusch A, Yigit G, Altmüller J, Boltshauser E, Dörk T,

Brockmann K. Evidence of Pathogenicity for the Leaky Splice Variant c.1066-6T>G in ATM

in a Patient with Variant Ataxia Telangiectasia. Abstracts of the 46th Annual Meeting of the

Society for Neuropediatrics, 2021. https://doi.org/10.1055/s-0041-1739584

24. Bergsma AJ, Kroos M, Hoogeveen-Westerveld M, Halley D, van der Ploeg AT, Pijnappel

WW. Identification and characterization of aberrant GAA pre-mRNA splicing in pompe

disease using a generic approach. Hum Mutat, 2015, 36:57–68

FIGURE LEGENDS

Figure 1. SPEER method and validation. A) SPEER concept. SPEER quantifies the

annotated and unannotated splicing variation in the healthy population by modeling the beta-

binomial distribution of junction usage for all junctions observed in the healthy normal samples.

Patient junction usage is then tested against this distribution to identify junctions that are

aberrantly used, filtering out common but unannotated variation. Significant splice aberrations

are then mapped to the canonical transcript to infer protein effects. Canonical junctions are

shown in dashed black lines, common alternatively-spliced junctions modeled by the normal

reference panel are shown in solid blue lines, and the aberrant splice junction detected in the

patient sample is shown with a solid red line. B) Bioinformatic workflow. FASTQ files from RNA-

Seq experiments are aligned and split reads are extracted and converted to raw junction counts.

SPEER uses the raw junction counts from the normal samples to compute the percent spliced in

(PSI) distributions for each junction in the healthy normal samples using a beta binomial count

model. Beta binomial parameters for each junction are saved in a panel of normal splicing

(PONS) file, which is calculated once and reused for every process-matched patient sample.

Patient junction counts are converted to PSI values, which are tested against the panel of

normal samples to identify significantly aberrant junction usage. Significant patient junctions are

positionally linked to assemble splice variant events. C) Validation metrics. SPEER was

validated using a set of positive and negative control blood samples from patients with known

and expected splicing events. Anonymous blood samples were used to create the PONS file.

SPEER showed high accuracy and reproducibility in detecting abnormal splicing events in

patient samples. D) Quantitative features of abnormal splicing events used for validation.

SPEER p-value and PSI fold change versus PONS are plotted for known positive and negative

control splice junctions. Positive and negative calls are strongly separated and inaccurate calls

fall near the p-value threshold. Infinite values are set to 20 on both axes for display purposes

and the top right point contains 11 samples with infinite fold change and a p-value of 0.

Figure 2. Establishing a PSI-X threshold for interpreting abnormal splice events as pathogenic. A) PSI-X (percent spliced in fold change) distribution for PSpVs (potential splicing variants) classified as pathogenic or likely pathogenic (PLP, beige) or benign or likely benign (BLB, blue). Dotted line represents PSI-X of 0.80. B) Receiver operating characteristic (ROC) curve for the data in panel A. The area under the ROC curve is 0.91. C) PSI-X values for heterozygous exon deletions detected by DNA sequencing. Although the PSI-X for deletions expected to cause NMD (nonsense mediated decay, beige circles, NMD+) is statistically different (one-sided t-test, p=0.0026) than the PSI-X for deletions expected to escape NMD (red circles, NMD-), all events are below 0.8, demonstrating the ability to correctly identify them independently of NMD status.

Figure 3. Characterization of splicing events observed in the prospective cohort. A)Position of variants associated with splicing events in regards to the transcript exon (0 is exonic). Although there is a clear enrichment of likely pathogenic and pathogenic variants at +1, +2 and +3, we do observe both benign and pathogenic variants at other positions. Overall, 93% of PSpVs associated with abnormal splicing events are within 20 bps of the exon, corresponding with the reportable range of the NGS DNA test. B) Splicing event types plotted by the number of potential splicing variants (PSpVs) observed. Colors represent classification of PSpVs following interpretation of abnormal splicing events. Most PSpVs classified as pathogenic or likely pathogenic are associated with changes leading to exon-skipping, while PSpVs associated with partial exon exclusion, partial intron inclusion, and complex events show the largest discrepancy between definitive classifications and variants of uncertain significance.

Figure 4. Discovery of intronic variants outside our reportable range with SPEER Thirty-five intronic DNA variants outside our reportable range were identified in this study. Five DNA

variants were classified as pathogenic or likely pathogenic (orange), two DNA variants were

classified as benign or likely benign (purple), and the remaining were classified as variants of

uncertain significance (VUS). The affected status of individuals with VUS is represented by

color: green = unaffected, blue = affected, and red = unknown due to lack of clinical information

provided. The canonical PSI fold change (PSI-X) showed a large distribution for these abnormal

splicing events, ranging from 0.23 to 0.9.

FIGURE LEGENDSS

Figure 1. SPEER method and validation.

Figure 2. Establishing a PSI-X threshold for interpreting abnormal splice events as

pathogenic.

Figure 3. Characterization of splicing events observed in the prospective cohort.

Figure 4. Discovery of intronic variants outside our reportable range with SPEER

TABLES

Table 1: Results from Phase 2

	Unique Individuals	% of All Samples
DNA + RNA sequencing	20,317	100
With a Potential Splicing Variant (PSpV)	3,563	17.5
PSpV & Normal Splicing	2,592	12.8
PSpV & Abnormal Splicing	971	4.8
PSpV upgrade VUS to PLP	75	0.4
PSpV downgrade VUS to BLB	1,204	5.9
Variants outside the reportable range	42	0.2

656   **Table 2: Classification of PSpVs in the Phase 2 study**

	PLP	VUS	BLB	Total PSpVs
PSpV & Abnormal splicing	260 (42.9%)	275 (45.5%)	70 (11.6%)	605 (100%)
PSpV & Normal splicing	120 (9.4%)	541 (42.3%)	617 (48.3%)	1278 (100%)
Total PSpVs	380 (20.2%)	816 (43.3%)	687 (36.5%)	1883 (100%)

659 **Table 3: Reclassification rates among self-reported ethnic populations**

	Unique individuals	% of ethnic population
Non-Hispanic White	608	4.8
Black/African American	158	10.4*
Asian	55	9.12*
Hispanic	137	9.4*
Other**	312	9.7

660 *p-value<0.05 after one-sided Fisher Exact test. **Includes Ashkenazi Jewish, Native American,

661 Multiple and Unknown.