1 **Exploring the Extreme Acid Tolerance of a Dynamic Protein Nanocage**

3 Jesse A. Jones[1], Michael P. Andreas[1], and Tobias W. Giessen[1,2]*

4 [1]Department of Biological Chemistry, University of Michigan Medical School, Ann Arbor, MI, USA

5 [2]Department of Biomedical Engineering, University of Michigan Medical School, Ann Arbor, MI, USA

6 *correspondence: tgiessen@umich.edu

Abstract

Encapsulins are microbial protein nanocages capable of efficient self-assembly and cargo enzyme encapsulation. Due to their favorable properties, including high thermostability, protease resistance, and robust heterologous expression, encapsulins have become popular bioengineering tools for applications in medicine, catalysis, and nanotechnology. Resistance against physicochemical extremes like high temperature and low pH is a highly desirable feature for many biotechnological applications. However, no systematic search for acid stable encapsulins has been carried out, while the influence of pH on encapsulin shells has so far not been thoroughly explored. Here, we report on a newly identified encapsulin nanocage from the acid-tolerant bacterium *Acidopropionibacterium acidopropionici*. Using transmission electron microscopy, dynamic light scattering, and proteolytic assays, we demonstrate its extreme acid tolerance and resilience against proteases. We structurally characterize the novel nanocage using cryo-electron microscopy, revealing a dynamic five-fold pore that displays distinct "closed" and "open" states at neutral pH, but only a singular "closed" state under strongly acidic conditions. Further, the "open" state exhibits the largest pore in an encapsulin shell reported to date. Non-native protein encapsulation capabilities are demonstrated, and the influence of external pH on internalized cargo is explored. Our results expand the biotechnological application range of encapsulin nanocages towards potential uses under strongly acidic conditions and highlight pH-responsive encapsulin pore dynamics.

Introduction

Protein-based compartments are used by many prokaryotes to regulate and optimize their metabolism in space and time.[1] One of the most widespread families of microbial protein compartments are encapsulins. Structurally, encapsulins self-assemble from a single type of HK97 phage-like shell protein to form icosahedral nanocages.[2] Encapsulins can exhibit different triangulation numbers, T=1 (60mer, ca. 24 nm), T=3 (180mer, ca. 32 nm), and T=4 (240mer, ca. 42 nm), with pores of varying sizes located at the 5-, 3-, and 2-fold symmetry axes.[3,4] They are classified into four distinct families, with Family 1 being the first discovered and most studied,[4] and derive their name from their ability to encapsulate specific co-regulated cargo enzymes. In Family 1, encapsulation is mediated by conserved peptide sequences found at the C-terminus of all cargo proteins called targeting peptides (TPs).[5] This feature allows encapsulins to perform a variety of biological functions, such as to act as sequestration chambers for dye-decolorizing peroxidases (DyPs), involved in combating oxidative stress;[5,6] to serve as iron mineralization and storage compartments;[7-10] as well as to sequester desulfurase enzymes, likely involved in sulfur metabolism.[11]

Due to their favorable properties, encapsulins have gained much attention as bioengineering tools.[3,12,13] As such, engineered encapsulins have been used in bacteria, yeast, and human cells for various applications, including as metabolic nanoreactors,[14] cellular imaging systems,[13] and drug delivery platforms.[15,16] A recent increase in the number of studies on encapsulin systems highlights the expanding scope of the field.[2,4,10,11,17] The encapsulin shell in particular has received substantial attention in recent years.[17] Efforts aimed at increasing shell stability,[15] controlling shell assembly,[18] and modulating pore size and dynamics have recently been reported.[19,20] Encapsulin shells efficiently self-assemble under many conditions and display marked resistance against chemical or temperature denaturation, pH, and non-specific proteases. For example, the melting temperature of the T=4 shell from *Quasibacillus thermotolerans* was reported to be nearly 87°C,[10] while the encapsulin from *Brevibacterium linens* was shown to be stable across a broad range of pH values (pH 5 to 11).[21] However, a number of biotechnological and industrial processes – including lignocellulose hydrolysis for biofuel production,[22] breakdown of complex sugars for monosaccharide production,[22] and bioleaching to prevent metal contamination and enable bioremediation in the mining industry[23,24] – would benefit from modular protein cages stable at acidic conditions below pH 5. Even though evolutionary adaptations of thermostable proteins have been well characterized and include oligomerization, large hydrophobic cores, and disulfide bond formation, adaptations that lead to acid-stable proteins are poorly understood.[25,26]

Here, we carry out the first bioinformatic search for acid-stable encapsulin nanocages and subsequently characterize the structure and stability of a Family 1 encapsulin shell from *Acidopropionibacterium acidopropionici* (AaEnc) – one of the top hits identified in our *in silico* analysis. Using a combination of techniques, including cryo-electron microscopy (cryo-EM), we characterize the structure and acid stability of the AaEnc nanocage across a wide range of pH values. Our results highlight the acid stability of AaEnc and its resilience towards protease digestion under different pH conditions. Cryo-EM analysis reveals a pH-dependent dynamic 5-fold pore with defined "closed" and "open" states, with the latter

representing the largest pore in an encapsulin shell reported to date. Further analyses confirm the non-
native cargo loading capabilities of AaEnc and reveal the effects of external pH on internalized cargo.

Results and Discussion

<u>Bioinformatic search for acid-tolerant encapsulin shells</u>

Only Family 1 encapsulins have so far been used as engineering platforms as they are the most studied
and well-understood of the different types of encapsulins – especially with respect to non-native cargo
loading. Therefore, with future engineering applications in mind, we chose to focus our bioinformatic
search for acid-tolerant shells on Family 1.[3,4,27,28] Further arguments for focusing on Family 1 are that the
widespread DyP encapsulins found in Family 1 are known to optimally function under acidic conditions[29]
while a variety of Family 1 systems encoded by acid-tolerant and acidophilic bacteria have already been
previously identified.[4]

As acid-tolerant proteins generally possess a low calculated isoelectric point (pI) caused by a large
number of surface-exposed negatively charged residues,[22,25,26,30] it was hypothesized that encapsulins
with a low pI may exhibit increased acid tolerance. Therefore, the large set of previously identified
Family 1 encapsulins with molecular weights between 26 and 35 kDa – the size range of non-cargo fused
encapsulins – were ranked by pI (**Figure 1a, Figure 1b,** and **Supplementary Data 1**). The lowest observed
pIs ranged from 4.14 to 4.49 with one protein found in the pI bin centered at a pI of 4.2 and 28 proteins
found in the pI bin centered at a pI of 4.4. Of these encapsulins, 18 are encoded by halophiles, four by
acidophiles or acid-tolerant bacteria, and the rest by soil bacteria or putative pathogens (**Table S1**).

We chose to focus on an encapsulin encoded by one of the acid-tolerant species, namely, the DyP
system of the industrially relevant *A. acidopropionici* (ATCC 4875) (AaEnc; **Figure 1c, Table S1,** and
Figure S1). *A. acidopropionici* is a Gram-positive actinobacterium able to tolerate acidic conditions as
low as pH 4.4.[31] It is used in agricultural applications and studied for its biotechnological and industrial
potential due to its production of propionic acid as a primary fermentation product, with acetic acid and
carbon dioxide as secondary products.[31-33]

The pI and sequence composition of AaEnc was compared to previously characterized encapsulin shells,
including those from *Mycobacterium smegmatis*, *Brevibacterium linens*, *Mycolicibacterium hassiacum*,
Haliangium ochraceum,[34] *Thermotoga maritima*, and *Myxococcus xanthus* (**Figure 1**, **Figure S1, Table 1,**
and **Table S1**).[2,7,21,35,36] AaEnc was found to have a lower pI than all of the so far characterized Family 1
encapsulins, and to display the largest ratio of acidic to basic residues (**Table 1** and **Table S1**). This makes
AaEnc a promising test system for exploring the acid stability of encapsulin shells.

Figure 1. Bioinformatic analysis of Family 1 encapsulins. **a**) Histogram of encapsulins based on calculated isoelectric point (pI), binned at 0.2 pH units. The pI bin of interest centered at pI 4.4 and containing the *A. acidopropionici* encapsulin (AaEnc) is highlighted in red. **b**) Scatter plot of encapsulins based on calculated pI and molecular weight (MW). AaEnc (pI: 4.41, MW: 28.5 kDa) is shown in red. Previously well-characterized encapsulins from *Mycobacterium smegmatis*, *Brevibacterium linens*, *Mycolicibacterium hassiacum*, *Haliangium ochraceum*, *Thermotoga maritima*, and *Myxococcus xanthus* are shown in cyan. **c**) The *A. acidopropionici* Family 1 encapsulin operon containing a dye-decolorizing peroxidase (DyP) cargo enzyme (blue) and encapsulin shell (AaEnc, red). Functionally unrelated genes are shown in gray with dashed outlines. Scale bar: 1 kilobase (kb).

Table 1. Comparison of the pI and charge of AaEnc and other characterized encapsulins.

Encapsulin	pI[1]	# Acidic residues	# Basic residues	Acid/base ratio	Charge at pH 7.0[2]
A. acidopropionici	4.41	41	25	1.64	-23.47
M. smegmatis	4.72	41	31	1.32	-16.44
B. linens	4.73	40	30	1.33	-18.77
M. hassiacum	4.78	41	32	1.28	-15.50
H. ochraceum	4.82	40	32	1.25	-13.47
T. maritima	4.90	50	39	1.28	-13.05
M. xanthus	5.45	37	36	1.03	-7.50

[1] Theoretical isoelectric points calculated with Expasy Compute pI/MW tool (https://web.expasy.org/compute_pi/).
[2] Charges at pH 7.0 calculated via Geneious Prime 2020.2.4 (https://www.geneious.com).

Biophysical analysis of the AaEnc protein nanocage

To characterize the AaEnc nanocage, it was first heterologously expressed in *Escherichia coli* and then purified via a combination of polyethylene glycol (PEG) precipitation, anion exchange chromatography (IEC), and size exclusion chromatography (SEC), with the latter serving as an initial verification that the nanocage was assembled at pH 7.5 (**Figure 2a**). Aliquots of the purified sample were then exchanged into various buffers across a wide range of pH values (pH 1.5 to 10.0) while holding the salt concentration constant at a physiological value of 150 mM NaCl. After incubation for 6 h, samples were imaged via negative stain transmission electron microscopy (TEM) to assess the effect of pH on protein aggregation and the assembly state of the AaEnc nanocage. Some assembled AaEnc shells could be observed at pH values as low as 1.9 and as high as 7.5 (**Figure S2**). However, the pH range within which AaEnc was close to fully assembled with only minor aggregation occurring was between pH 2.25 and 7.5. Therefore, subsequent biophysical analyses were carried out at four pH values spanning this pH range, namely at pH 2.25, 3.0, 5.0, and 7.5. Dynamic light scattering (DLS) analyses and negative stain TEM indicated that AaEnc maintained a similar size and appearance across all four tested conditions with Z-average diameters of 30.3 nm at pH 2.25, 30.2 nm at pH 3.0, 29.8 nm at pH 5.0, and 24.4 nm at pH 7.5 (**Figure 2b** and **Figure 2c**). The slight increase in average diameter at acidic pH values is likely due to limited aggregation under these conditions, however, as can be seen in TEM micrographs, individual shells at all tested pH values exhibited diameters of ca. 24 nm. Static light scattering (SLS) further indicated that AaEnc is relatively stable across all tested pH values, with aggregation temperatures (T_{agg}) of 36.0°C at pH 2.25, 38.3°C at pH 3.0, 39.3°C at pH 5.0, and 62.4°C at pH 7.5 (**Figure 2d**). However, a clear trend can be observed with lower pH values leading to decreased T_{agg} values. We next explored the resistance of AaEnc against proteolytic degradation at various pH values. AaEnc proved to be relatively resilient to pepsin degradation at pH 3.0 and 37°C during a 3 h incubation period, whereas the control protein, bovine serum albumin (BSA), was completely degraded under the same conditions (**Figure 2e**). Similarly, AaEnc was relatively resistant to degradation by trypsin and chymotrypsin at pH 7.5 and 37°C over an 8 h period, with BSA being again substantially degraded under the same conditions (**Figure 2f**). Overall, these results highlight the substantial acid stability of AaEnc which is significantly higher than that of any other previously characterized encapsulin.

Figure 2. Biophysical analysis of the AaEnc nanocage. **A)** Size exclusion chromatography (SEC) of the AaEnc nanocage depicting elution at 12 mL suggestive of an assembled T=1 encapsulin (left), along with SDS-PAGE analysis of purified AaEnc (right). **B)** Dynamic light scattering (DLS) analysis of AaEnc at pH 7.5 (purple), pH 5.0 (green), pH 3.0 (orange), and pH 2.25 (red). **C)** Transmission electron microscopy (TEM) analysis of AaEnc after 6 h incubation at different pH values: pH 2.25 (red, far left), pH 3.0 (orange, middle left), pH 5.0 (green, middle right), and pH 7.5 (purple, far right). Scale bars: 50 nm. **D)** Representative thermal unfolding curves for AaEnc at different pH values: pH 2.25 (red), pH 3.0 (orange), pH 5.0 (green), and pH 7.5 (purple) with corresponding aggregation temperatures (T_{agg}, vertical dashed lines, respective colors). **e)** Protease stability analysis of AaEnc exposed to pepsin at pH 3.0 for 3 h with bovine serum albumin (BSA) as a control. **f)** Protease stability analysis of AaEnc exposed to trypsin and chymotrypsin (T&C) at pH 7.5 for 8 h with BSA as a control.

Structural characterization of the AaEnc protein nanocage

To further characterize the influence of pH on AaEnc, single particle cryo-EM analysis was carried out, initially at a physiological pH of 7.5. Results revealed the existence of two discrete structural states distinguished by an either all "closed" or all "open" conformation of the 5-fold pores within the AaEnc

shell (**Figure 3** and **Supplementary Video 1**). The "closed" and "open" states were determined to 2.90
(29,056 particles) and 3.32 Å (13,581 particles), respectively (**Figure S3**). About 68% of the used particles
comprise the "closed" state while about 32% exhibit an "open" state. The shell for both states consists
of 60 AaEnc protomers, forming a ca. 1.7 MDa, T=1 icosahedral protein cage with a diameter of 24 nm
and an overall negatively charged exterior surface (**Figure 3a** and **Figure 3c**). Symmetric (icosahedral, I)
and asymmetric (C1) refinements were carried out for both states to investigate if a given pore can be
"closed" or "open" independently of the other pores in the same shell or if all pore states are generally
correlated. Both I and C1 refinements yielded similar all "closed" and all "open" states (**Figure S3**), thus
confirming that under the given experimental conditions, the pore dynamics of all pentameric facets
within a shell appear to be strongly correlated. However, this does not necessarily exclude the possibility
that, under certain environmental conditions or in the presence of cargo, a single shell can contain both
"closed" and "open" 5-fold pores at the same time. This phenomenon – independent dynamic 5-fold
pores – has indeed been observed for the Familiy 1 encapsulin from *H. ochraceum* (**Figure S4**).[37] To
further analyze 5-fold pore dynamics, 3D variability analysis with three components was carried out in
cryoSPARC for both datasets, however, no states could be resolved that would indicate the
simultaneous presence of both "closed" and "open" pores within the shell. The 5-fold pore diameters in
the AaEnc nanocage are 5 Å for the "closed" and 20 Å for the "open" state (**Figure 3b, Figure 3d**, and
Figure S3). Thus, the AaEnc "open" state represents the largest pore found in an encapsulin shell to
date, substantially larger than the previously reported dynamic *H. ochraceum* pore which exhibited an
"open" state diameter of 15 Å.[37]

Detailed examination of the "closed" and "open" state structures indicates that the primary
conformational changes underlying the dynamic nature of the AaEnc pore are located at the apex of the
so-called axial domain (A-domain) of the encapsulin protomer (**Figure 4a**). In addition, an overall
backwards tilt of the "open" state protomer by 10° (**Figure 3b** and **Figure 3d**) also contributes to the
observed size increase of the 5-fold pore (**Figure 3c**, **Figure 3d, Supplementary Video 2**). In the "closed"
state, five pore residues – Asp185, His186, Gly187, Val188, and Pro189 – form a short loop between the
α6 and α7 helices encompassing a predicted α-turn (**Figure S5**); whereas in the "open" state, the same
residues form a tighter turn that loses the predicted α-turn conformation, with Val188 and Pro189
becoming part of and extending the α7 helix (**Figure 4c**, **Figure S5**, and **Figure S6**).[38] His186 undergoes
the most readily observable conformational change. It is located at the apex of the A-domain in the
"closed" state, yet partially buried between two adjacent protomers in the "open" state (**Figure 4b**,
Figure 4c). Furthermore, in the "open" state, intermolecular hydrogen bonding is observed between
His186 and Asp150 as well as Ser181 and Asn157 of adjacent protomers (**Figure 4c**). These two
hydrogen bonds are notably absent in the "closed" state. His186 is not strictly conserved among other
structurally characterized Family 1 encapsulins (**Figure S1**) and cannot be used alone as an indicator for
the presence of dynamic 5-fold pores in encapsulin shells, as in the *H. ochraceum* encapsulin, which also
displays "closed" and "open" pore states, where His186 is substituted with Asp186 (**Figure S1** and **Figure
S4**). Interestingly, in *H. ochraceum* the residue corresponding to Asp150 in AaEnc is Arg150. This could
indicate that analogous to the hydrogen bonding between His186 and Asp150 (AaEnc), bonding
between Asp186 and Arg150 (*H. ochraceum*), with swapped H-bond donors/acceptors, may be possible
under certain conditions. However, this was not observed in the *H. ochraceum* "open" conformation.

Figure 3. Structural overview of the AaEnc nanocage and 5-fold pore. **a**) Electrostatic surface representation of AaEnc in the "closed" conformation viewed down the 5-fold symmetry axis as well as at a 45° rightward turn (red, negative charge; white, neutral; blue, positive charge). **b**) Top-down ribbon and partially transparent surface representation of the "closed" AaEnc pentamer (left; cyan) and solid surface representation rotated 90° and viewed through the frontal axis plane to highlight pore size (right). The angle between protomers and the orthogonal of the 5-fold axis is highlighted. **c**) Electrostatic surface representation of AaEnc in the "open" conformation. **d**) Top-down ribbon and partially transparent surface representation of the "open" AaEnc pentamer (left; purple) and solid surface representation rotated 90° and viewed through the frontal axis plane to highlight pore size (right). The angle between protomers and the orthogonal of the 5-fold axis is highlighted.

Overall, the cryo-EM density for the loop region of the *H. ochraceum* encapsulin in the "open"
conformation was not well defined, whereas for AaEnc, both "closed" and "open" states exhibit strong
and well-defined densities (**Figure S6**). This could in part be due to the additional stabilization of the
"open" state in AaEnc resulting from the hydrogen bonding observed between His186 and Asp150.

Figure 4. Detailed structural analysis of the AaEnc 5-fold pore. **a)** Aligned and overlayed ribbon representation of the "closed"
(cyan) and "open" (purple) AaEnc protomers with dashed box highlighting the A-domain. **b)** Magnified ribbon representation
juxtaposing the dynamic A-domain of the "closed" (cyan; left, solid; right, transparent) and "open" (purple; right, solid; left,
transparent) AaEnc protomer, with the loop residues of interest–Asp185, His186 (labeled), Gly187, Val188, and Pro189–
highlighted (green). **c)** Magnified solid ribbon representation of two adjacent AaEnc A-domains with transparent surface
representation of the AaEnc pentamer highlighting the 5-fold pore. The "closed" state (left, cyan) exhibits a lack of hydrogen
bonds between Asp150 (gray) and His186 (green) as well as Asn157 (gray) with Ser181 (gray), while the "open" state (right,
purple) showcases gained hydrogen bonds between Asp150 (red) with His186 (green), as well as Asn157 (red) with Ser181
(red).

Additional cryo-EM experiments were carried out at pH 3.0 to assess the structure of AaEnc under
strongly acidic conditions. At pH 3.0, only a single "closed" conformational state was observed and was
determined to 2.77 Å resolution (47,164 particles) (**Figure S7** and **Table S3**). It was also found that both
of the "closed" AaEnc states – at pH 7.5 and pH 3.0 – are seemingly identical, with a root-mean-square
deviation (RMSD) of 0.32 between the two aligned protomers (**Figure S8**).[39]

In silico analyses were conducted using the APBS-PDB2PQR software suite to quantitatively assess the
hydrogen bonding and solvent exposure of the His186 and Asp150 residues at pH 7.5 and 3.0 (**Table 2**).[40]
The results further corroborate that His186 is more exposed – calculated as only 10% buried – in the
"closed" states, and more buried – calculated as 39% buried – in the "open" state due to the inter-
protomer hydrogen bonding described above. Furthermore, no hydrogen bonds were predicted
between His186 and Asp150 in either of the "closed" states, while being clearly predicted for the "open"
state.

Based on the results outlined above the "closed" state is clearly favored at low pH. It seems likely that in
addition to global protonation state changes throughout the AaEnc protomer, specifically the
protonation of Asp150 at low pH would preclude any hydrogen bonding with His186, thus making the
conformational change from "closed" to "open" state energetically less favorable at acidic conditions.
The specific molecular and biological functions of dynamic 5-fold pores in encapsulin shells is currently
unknown. However, as the native cargo of AaEnc is a DyP-type peroxidase, generally known to be
optimally active at acidic pH values, the preference of the AaEnc nanocage for the "closed" pore state at
low pH might have significant functional and biological implications.

Table 2. *In silico* analysis of key pore residues. Calculated buriedness, hydrogen bonds, and pKa values are shown.

State and residue	Buried	Sidechain Hydrogen Bond (Partner)[1]	Calculated pKa
Closed (pH 7.5)			
Asp150	44%	-0.57 (Gln147, intramolecular)	5.63
His186	10%	none	6.00
Open (pH 7.5)			
Asp150	12%	-0.53 (His186, intermolecular)	3.50
His186	39%	0.53 (Asp150, intermolecular)	6.63
Closed (pH 3.0)			
Asp150	43%	0.90 (Asp149, intramolecular)	7.64
His186	10%	none	5.85

[1] pKa shift due to respective hydrogen bond.

<u>Computational analysis of continuum electrostatics and solvation of the AaEnc protomer</u>

To gain deeper insights into the stability of the AaEnc nanocage under acidic conditions, an array of
computational analyses focused on continuum electrostatics and solvation of the AaEnc protomer was
carried out. In particular, we thought to investigate if the AaEnc protomer would be predicted to show
increased acid stability by itself outside the context of the encapsulin shell. The APBS-PDB2PQR software
suite was used to assess the surface electrostatics of AaEnc at different pH values ranging from pH 2.0 to
8.0 (**Figure 5a**).[40] The calculated protein electrostatics correlated well with the calculated pI of 4.41,
showing a change from an overall positive surface charge below pH 4.0 to an overall negative surface
charge above pH 5.0. Next, using the Protein-Sol software package, a heatmap depicting the average
charge per residue based on pH and ionic strength was calculated.[41] The results again correlate well with
the calculated pI, with a positive average charge per residue at pH 4.0 and below, and a negative
average charge per residue at pH 4.5 and above, regardless of ionic strength (**Figure 5b**). To investigate if
a discernible increase in folded state protein stability mediated by interactions between ionizable

groups might exist for the AaEnc protomer at acidic pH, further analyses using Protein-sol were carried
out (**Figure 5c**). It was found that at physiological ionic strength (150 mM), per residue energies below
pH 4.5 were positive, indicating decreased stability of the protomer fold below this pH threshold.

Taken together, our computational analysis of the AaEnc protomer suggests that the observed acid
stability of the AaEnc nanocage is not easily attributable to a highly stable protomer building block.
Instead, AaEnc acid stability is likely due to a complex combination of factors at the scale of the
assembled 60mer nanocage. Important factors likely include favorable inter-protomer interactions
within the context of the encapsulin shell, such as the structural dynamics and hydrogen bonding
interactions discussed above.

Figure 5. Computational characterization of the AaEnc protomer. **a**) Surface charge visualization of the AaEnc protomer with
amino acid protonation states calculated by pH with pdb2pqr and protein electrostatics calculated with APBS. **b**) Heatmap of
predicted average charge per residue for the AaEnc protomer at different pH and ionic strength values. **c**) Heatmap of predicted
folded state protein stability as interactions between ionizable groups in joule per residue for the AaEnc protomer at different
pH and ionic strength values.

<u>*In vivo* cargo loading and pH effects on internalized cargo</u>

To investigate if the AaEnc shell has any influence on the acid stability of internalized cargo proteins,
heterologous cargo loading experiments were carried out followed by pH screens. The predicted C-
terminal targeting peptide (TP) of the native AaEnc DyP cargo enzyme, was genetically fused to the C-
terminus of eGFP (eGFP-TP) and cloned immediately upstream of the AaEnc gene for co-expression.[4,42]
eGFP was chosen as a non-native cargo due to its reliable expression, favorable solubility, simple
detection, and predictable and well-reported pH sensitivity profile.[43,44] *In vivo* eGFP-TP cargo loading
was confirmed via its co-purification by SEC with co-expressed AaEnc, negative stain TEM analysis, and
native polyacrylamide gel electrophoresis (PAGE) (**Figure 6a**, **Figure 6b**, and **Figure 6c**). Next, the

fluorescence of equimolar amounts of free eGFP and AaEnc-encapsulated eGFP-TP were compared
across a range of pH values from pH 3.0 to 8.0 using a plate-based fluorescence assay (**Figure 6d**). Both
free eGFP and AaEnc-encapsulated eGFP-TP yielded very similar sigmoidal fluorescence response curves,
demonstrating that the interior pH of AaEnc is not appreciably different from the bulk pH. Further,
encapsulation within AaEnc does apparently not alter cargo pH sensitivity to a significant degree. These
results indicate that the AaEnc shell does not represent an effective diffusion barrier for protons. Thus,
buffer pH changes will result in the rapid equilibration of the external and luminal pH. Similar behavior
has been observed for other protein-based compartments, particularly the carbon-fixing carboxysome
bacterial microcompartment.[45] Stopped-flow pH colorimetry indicated a rapid equilibration of the
luminal carboxysome pH to that of the bulk solvent highlighting the porosity of the carboxysome shell
towards protons. However, considering that one of the likely primary functions of encapsulin and
carboxysome shells is to control the flux of specific small molecules into and out of the shell interior[45,46]
– possibly through the action of dynamic or gated pores – the idea that protein shells could be able to
control the passage of protons is not out of the realm of possibility. A molecular mechanism similar to
that employed by aquaporins, which allow passage of water molecules but block proton flux, could
certainly exist or be engineered in protein shells.[47]

Figure 6. Analysis of eGFP cargo-loaded AaEnc. **a**) Size exclusion chromatography (SEC) analysis of AaEnc-encapsulated eGFP
showing elution at 12-13 mL via protein absorbance at 280 nm and specific eGFP absorbance at 488 nm. **b**) SDS-PAGE analysis
of purified eGFP-loaded AaEnc (left) and negative stain TEM (right). Scale bar: 50 nm. **c**) Native PAGE gel analysis of empty

Conclusion

The large number of encapsulin systems distributed across diverse bacterial and archaeal phyla –
including many extremophiles – represents a largely untapped source of novel biotechnological tools.[4]
Ongoing discoveries and research within the encapsulin field has resulted in the characterization of
many nano-encapsulation systems with a quickly expanding and diverse list of useful molecular
features.[10,11,14,18,28] However, relatively little attention has been focused on systematically exploring
encapsulins from extremophilic bacteria and archaea with unusual molecular characteristics and
stability profiles. With this study, we have taken the first step towards addressing this issue with a focus
on the acid stability of the AaEnc nanocage.

Our results highlight the difficulty of pinpointing specific protein characteristics that lead to increased
acid stability. Beyond the previously reported observation that acid stable proteins contain an increased
number of aspartate and glutamate residues, resulting in a low pI,[48] no other adaptations towards acid
stability are readily apparent for AaEnc. We find that at the protomer level, AaEnc does not display any
obvious properties beyond a low pI, that would indicate exceptionally high acid stability. This analysis is
necessarily purely computational as encapsulin protomers quickly self-assemble to form protein
nanocages and cannot be studied in isolation. It seems likely that the formation of a large 60mer protein
complex plays a role in the acid tolerance of AaEnc with some assembled shells still present after
extended incubation at pH 1.9. It can be speculated that minimizing the number of ionizable groups at
key subunit interfaces within the AaEnc shell would contribute towards its stability at low pH. Beyond its
unusual acid tolerance, AaEnc exhibits the unique feature of seemingly pH-dependent all "closed" or all
"open" 5-fold pore states. At physiological pH, the "closed" and "open" states exist in a ratio of 2:1,
whereas at low pH, the equilibrium is completely shifted towards the "closed" state. This behavior likely
has important functional and biological implications and will require further study. Its stability, pH-
responsive pores, and the fact that the "open" pore state with a diameter of 20 Å represents the largest
encapsulin pore reported to date, make AaEnc an interesting target for future nanocage engineering
applications in catalysis, nanotechnology, and medicine.

Specifically, acid-tolerant protein cages could offer novel opportunities in nanoreactor design and
engineering, particularly in the context of industrial biopolymer degradation which requires acidic
conditions. Enzyme encapsulation or co-localization could improve the performance of enzymes like
chymosins,[49] dye-decolorizing peroxidases,[50] glucoamylases,[51] and proteases,[52] already extensively used
in the food industry, agriculture, and biofuel production.[53-56] Further, bioleaching and bioremediation
approaches aimed at recovering valuable or toxic metals could benefit from acid stable nano-
encapsulation systems able to sequester specific metal-binding enzymes of interest.[24,57] Finally, a
number of biomedical applications of protein nanocages related to drug delivery or intracellular

targeting of acidic compartments could also benefit from robust and easily engineerable nanocages like
AaEnc.[58,59]

In sum, AaEnc is the first characterized highly acid stable encapsulin nanocage with a unique pH-
dependent dynamic pore and, therefore, represents a novel useful tool for the nanocage engineering
community.

Methods

Chemicals and biological materials

All chemicals were used as supplied by vendors without further purification. Imidazole, Invitrogen Novex
WedgeWell 14% tris-glycine Mini Protein Gels, Isopropy-ß-D-thiogalactopyranoside (IPTG), lysozyme,
NativePAGE™ 4 to 16% bis-tris Mini Protein Gels, NativeMark Unstained Protein Standard, Spectra™
Multicolor Broad Range Protein Ladder, Thermo Scientific Pierce 660 nm Protein Assay Reagent, Tris
base, Tris HCl, all restriction enzymes, and all cell culture media and reagents were purchased from
Fisher Scientific, Inc. (USA). Gibson Assembly Master Mix was purchased from NEB (USA). Amicon Ultra-
0.5 mL centrifugal units and Benzonase® nuclease were purchased from MilliporeSigma (USA). BL21
(DE3) Electrocompetent Cells used for *E. coli* expression were also purchased from MilliporeSigma (USA).
Bis-tris propane from Research Products International (USA) was used for the assembly buffer. Ni-NTA
agarose from Gold Biotechnology, Inc. (USA) was used for His-tagged protein purification.

Instrumentation

Cell lysis was conducted via sonication with a Model 120 Sonic Dismembrator from Fisher Scientific, Inc.
(USA). Protein was quantified on a Nanodrop Spectrophotometer from ThermoFisher Scientific, Inc.
(USA). Protein purification was carried out on an AKTA Pure fast liquid protein chromatography system;
size exclusion chromatography (SEC) was carried out with a HiPrep 16/60 Sephacryl S-500 HR and
Superose 6 10/300 GL columns (Cytiva, USA); anion exchange was carried out with a HiTrap Q FF column
(Cytiva, USA). Polyacrylamide gel electrophoresis (PAGE) and NativePAGE were performed in an XCell
SureLock from Invitrogen/ThermoFisher Scientific (USA). Gel images were captured using a ChemiDoc
Imaging System from Bio-Rad Laboratories, Inc. (USA). DLS was carried out on an Uncle from Unchained
Labs (USA). TEM was carried out on a Morgagni 100 keV Transmission Electron Microscope (FEI, USA).
Plate-based fluorescence assays were conducted on the Synergy H1 Microplate Reader from BioTek
Instruments (USA). EM grid glow discharging was conducted with a PELCO easiGlow™ system by Ted
Pella, Inc (USA). A Talos Arctica Cryo Transmission Electron Microscope by ThermoScientific, Inc. (USA)
equipped with a K2 Summit direct electron detector by Gatan, Inc. (USA) located at the University of
Michigan Life Sciences Institute was used for cryo-EM. Smaller materials are listed along with
corresponding methods below.

Software

The following software was used throughout this work: Adobe Illustrator 2021 v25.0.0 (figures),
cryoSPARC v3.3.1[60] (cryo electron microscopy), Fiji/ImageJ v2.1.0/1.53c[61] (densitometric data analysis

and TEM images), GraphPad Prism for Mac OS v9.4.0 (chromatography, melting

temperature/aggregation, and fluorescence graphs), Bio-Rad Image Lab Touch Software (gel imaging),

Microsoft Excel for Mac v16.46 (DLS graphs), Phenix v1.19.2-4158[62] (model building), UCSF Chimera

v1.16[63] and ChimeraX v3[39] (cryo-EM density and model visualization), and UNICORN 7 (FPLC system

control and chromatography). Online software suites or tools are listed along with corresponding

methods below.

<u>Bioinformatic search for acid-stable encapsulins</u>

384　A curated list of Family 1 encapsulins[19] was sorted according to molecular weight, removing any entries

385　falling below 25 kDa or above 35 kDa to remove partial annotations and fusion encapsulins, respectively.

386　Results were then processed via the Expasy Compute pI/MW tool

387　(https://web.expasy.org/compute_pi/). Data was then organized according to calculated pI and binned

388　for histogram analysis or plotted for scatterplot analysis via GraphPad Prism.

<u>Sequence alignments</u>

390　Encapsulin alignments were generated with the ESPript 3 server (http://espript.ibcp.fr/) using a protein

391　sequence alignment produced with Clustal Omega, with secondary structure information based on the

392　TmEnc structure (PDB 3DKT; **Figure S1**) or the "open" and "closed" AaEnc structures (**Figure S5**).[38,64]

<u>Protein production</u>

For all target proteins, plasmids were constructed with target *E. coli* codon-optimized gBlock genes,

synthesized by IDT (USA), inserted into the pETDuet-1 vector via Gibson assembly using the NdeI and

PacI restriction sites (**Table S2**). *E. coli* BL21 (DE3) was transformed with the respective plasmids via

electroporation per protocol and 25% glycerol bacterial stocks were made and stored at -80°C until

further use. Starter cultures were grown in 5 mL LB with 100 mg/mL ampicillin at 37°C overnight. For all

constructs, 500 mL of LB with ampicillin was inoculated with overnight starter cultures and grown at

37°C to an OD600 of 0.4-0.5, then induced with 0.1 mM IPTG and grown further at 30°C overnight for

ca. 18 h. Cells were then harvested via centrifugation at 10,000 rcf for 15 minutes at 4°C and pellets

were frozen and stored at -80°C until further use.

<u>Protein purification</u>

Frozen bacterial pellets were thawed on ice and resuspended in 5 mL/g (wet cell mass) of cold Tris

Buffered Saline (20 mM Tris pH 7.5, 150 mM NaCl). Lysis components were added (0.5 mg/mL lysozyme,

1 mM tris(2-carboxyethyl)phosphine [TCEP], one SIGMA*FAST* EDTA-free protease inhibitor cocktail tablet

per 100 mL, 0.5 mM $MgCl_2$, and 25 units/mL Benzonase® nuclease) and samples were lysed on ice for 10

min. Samples were then sonicated at 60% amplitude for 5 min total (eight seconds on, 16 seconds off)

until no longer viscous. After sonication, samples were centrifuged at 8,000 rcf for 15 minutes at 4°C.

Samples were then subjugated to 10% polyethylene glycol (PEG) 8000 precipitation (lysate brought to

10% PEG 8K and 500 mM NaCl and incubated for 30 minutes on ice, then centrifuged 8,000 rcf for 15

min). Supernatant was discarded and the pellet was resuspended in 5 mL TBS pH 7.5 and filtered using a

0.22 µm syringe filter (Corning, USA). The protein sample was then loaded on an AKTA Pure and purified

via a Sephacryl S-500 column. Sample fractions were pooled and buffer exchanged into AIEX Buffer (20
mM Tris pH 7.5) and loaded onto an AKTA Pure, then purified via HiTrap Q-FF by linear gradient into
AIEX Buffer with 1M NaCl. Sample flow-through was collected and centrifuged at 10,000 rcf for 10 min,
then loaded on an AKTA Pure for final purification via a Superose 6 10/300 GL column pre-equilibrated
with TBS pH 7.5. All proteins were stored at 4°C until use.

For free His-tagged eGFP purification, the sample was lysed as above in NTA Resuspension Buffer (50
mM Tris pH 8.0, 150 mM NaCl, 1 mM TCEP, 10 mM imidazole, and 5% glycerol). Lysate was bound to Ni-
NTA resin pre-equilibrated with NTA Resuspension Buffer via rocking at 4°C for 45 minutes. Supernatant
was discarded and the bound sample was washed once with NTA Resuspension Buffer and a second
time with NTA Resuspension Buffer with 20 mM imidazole. Free His-tagged eGFP was then eluted three
times with Elution Buffer (50 mM Tris pH 8.0, 150 mM NaCl, 1 mM TCEP, 350 mM imidazole, 5%
glycerol) and stored at 4°C for future use.

Transmission electron microscopy

Samples were diluted to 0.1-0.3 mg/mL and buffer exchanged into various buffers ranging from pH 1.5
to pH 10.0 (**Figure 2c** and **Figure S2**) via five successive exchanges in 100 kDa MWCO Amicon Ultra-0.5
mL centrifugal units. Buffers used consisted of 50 mM sodium phosphate and 150 mM NaCl, pH 1.5; 50
mM sodium phosphate and 150 mM NaCl, pH 1.9; 50 mM sodium phosphate and 150 mM NaCl, pH 2.1;
50 mM sodium phosphate and 150 mM NaCl, pH 2.25; 50 mM sodium phosphate and 150 mM NaCl, pH
3.0; 50 mM sodium citrate and 150 mM NaCl, pH 5.0; 50 mM MES and 150 mM NaCl, pH 6.0; 50 mM bis-
tris propane and 150 mM NaCl, pH 9.0; and 50 mM CHES and 150 mM NaCl, pH 10.0. Samples were
incubated at 4°C for six hours, and then immediately stained and imaged. Additional pH 2.25 and pH 3.0
samples were stored at 4°C for two days and stained and imaged. Negative stain transmission electron
microscopy (TEM) was carried out on the various samples with 200-mesh gold grids coated with extra
thick (25-50 nm) formvar-carbon film (EMS, USA) made hydrophilic by glow discharging at 5 mA for 60 s.
Briefly, 3.5 µL of sample was added to the grid and incubated for 30 seconds, wicked with filter paper,
and washed once with distilled water and once with 0.75% (w/v) uranyl formate before staining with 8.5
µL of uranyl formate for 30 seconds. TEM images were captured using a Morgagni transmission electron
microscope at 100 keV at the University of Michigan Life Sciences Institute. For all TEM experiments,
samples were roughly 0.2 mg/mL of AaEnc monomer in appropriate buffer.

Dynamic and static light scattering analyses

All sizing and polydispersity measurements were carried out on an Uncle by Unchained Labs (USA) at 30
°C in triplicate. Purified AaEnc samples were adjusted to 0.4 mg/mL of monomer in the appropriate
corresponding buffers and centrifuged at 10,000 rcf for 10 min, then immediately analyzed via DLS
(**Figure 2b**). Static light scattering aggregation temperature (T_{agg}) analysis was then conducted on
similarly prepared samples over a 25°C to 95°C ramp at 1°C per minute (**Figure 2d**).

Protease assays

AaEnc and bovine serum albumin (BSA; ThermoScientific Pierce, USA) were individually buffer
exchanged into Pepsin Assay Buffer (50 mM Na_2PO_4, 150 mM NaCl, pH 3.0) and mixed in a 40:1 molar

ratio with commercially purchased pepsin protease (Promega, USA), then incubated at 37°C for 3 h and
frozen until later use. Purified AaEnc and BSA were individually buffer exchanged into TBS pH 7.5 and
mixed in a 40:1:1 molar ratio with commercially purchased trypsin (Promega, USA) and chymotrypsin
(Promega, USA), then incubated at 37°C for 8 h and frozen until later use. All samples were then rapidly
thawed and examined via PAGE analysis.

Cryo-electron microscopy

Sample preparation

The purified protein samples were concentrated to 3 mg/mL in 150 mM NaCl, 20 mM Tris pH 7.5 or 150
mM NaCl, 50 mM Na_2PO_4 pH 3.0. 3.5 µL of protein samples were applied to freshly glow discharged
Quantifoil R1.2/1.3 grids and plunged into liquid ethane using an FEI Vitrobot Mark IV (100% humidity,
22°C, blot force 20, blot time 4 seconds, drain time 0 seconds, wait time 0 seconds). The frozen grids
were clipped and stored in liquid nitrogen until data collection.

Data collection

Cryo-electron microscopy movies were collected using a ThermoFisher Scientific Talos Arctica operating
at 200 keV equipped with a Gatan K2 Summit direct electron detector. Movies were collected at 45,000x
magnification using the Leginon[65] software package with a pixel size of 0.91 Å/pixel and an exposure
time of 5 or 8 s, frame time of 200 ms, and total dose of 42 e^-/A^2 for the pH 3.0 sample and 41 e^-/A^2 for
the pH 7.5 sample. 1,357 movies were collected for the pH 3 sample and 975 movies were collected for
the pH 7.5 sample.

Data processing

pH 7.5 sample: All data processing was performed using cryoSPARC v3.3.1.[60] 975 Movies were imported
and motion corrected using Patch Motion Correction and CTF fits were refined using Patch CTF. 821
movies with CTF fits better than 8.0 Å were selected for downstream processing. Roughly 200 particles
were picked manually using Manual Picker and grouped into 10 classes using 2D Classification. Well
resolved classes were selected and used as templates for Template Picker to pick particles with a
specified particle diameter of 240 Å. 56,583 particles with a box size of 384 pixels were extracted and
subjected to 3 rounds of 2D Classification with 100 classes yielding 44,686 particles in good classes. Ab-
Initio Reconstruction with 6 classes and I symmetry was carried out next. The two main classes were
selected representing the all "closed" (29,056 particles) and all "open" (13,581 particles) states. Particles
from each respective state were used as inputs for separate Homogenous Refinement jobs (with I or C1
symmetry) with the following settings: optimize per-particle defocus, optimize per-group CTF params,
and Ewald Sphere correction enabled. The I refinements yielded a 2.90 Å density for the "closed" state,
and a 3.32 Å density for the "open" state, whereas the C1 refinements resulted in 4.84 Å and 4.44 Å
maps, respectively (**Figure S3**). 3D Variability Analysis with 3 components was carried out on both
particle sets using the C1 Refinement results as inputs, however, no components could be resolved
corresponding to "open" and "closed" states within the same density.

pH 3.0 sample: The same preprocessing procedure was used as for the pH 7.5 sample yielding 52,210
extracted particles with a box size of 384 pixels. 3 rounds of 2D classification with 100 classes resulted in
47,599 good particles. Ab-Initio Reconstruction with 6 classes and I symmetry was carried out yielding
one dominant class containing 47,164 particles. This was followed by I and C1 Homogenous Refinement
jobs using the following parameters: optimized per-particle defocus, optimize per-group CTF params,
and Ewald Sphere correction. The I refinement resulted in a density of 2.77 Å while the C1 refinement
yielded a 4.12 Å map (**Figure S7**).

Model building

A homology model was generated using RoseTTAFold[66] on the Robetta server and was used as a starting
model for all model building efforts. This starting model was manually placed into the respective cryo-
EM maps using Chimera v1.16,[63] and was further fit using the Fit to Volume command. The placed
monomeric models were then manually refined against the respective cryo-EM maps using Coot
v8.9.6.[67] The resulting models were further refined using Real Space Refine in Phenix v 1.19.2-4158[68]
with default settings and three iterations. After inspecting the refined models in Coot, symmetry
restraints were pulled from the maps using the Phenix.Find_NCS_from_Map command with I symmetry.
Complete shell models were assembled using the Phenix.Build_from_NCS command. These shell models
were then used as inputs for a final round of Real Space Refine with NCS restraints, 3 iterations, and all
other settings set to default. The models were deposited to the PDB under PDB ID 8DN9, 8DNL, and
8DNA; and the EMDB under EMD-27558, EMD-27573, and EMD-27560.

<u>Computational electrostatics and solvation analyses</u>

In silico hydrogen bonding and buriedness analyses were conducted using the APBS-PDB2PQR software
suite (https://server.poissonboltzmann.org/pdb2pqr) with PROPKA v3.2 to predict pKa values and assign
protonation states at the provided pH values. Analyses were conducted using the "closed" and "open"
state AaEnc structures at pH 7.5 as well as the "closed" AaEnc structure at pH 3.0. The AaEnc protomer
was further analyzed using APBS-PDB2PQR to assess the calculated surface charge of AaEnc across
various pH values from pH 2.0 to pH 8.0 (**Figure 5a**).[40] Calculated protomer charge and stability
heatmaps were generated with the Protein-Sol webtool (https://protein-
sol.manchester.ac.uk/heatmap) using the AaEnc protomer from the "closed" state at pH 7.5 as the input
(**Figure 5b** and **Figure 5c**).[41] Monomer structure alignments were carried out in ChimeraX.[39]

<u>*In vivo* cargo loading, native PAGE, and fluorescence analysis</u>

The AaEnc encapsulated eGFP-TP sample was co-expressed, purified, and analyzed via TEM in the same
manner as AaEnc as described above (**Figure 6a and Figure 6b**). The free His-tagged eGFP and the
AaEnc-encapsulated eGFP-TP samples were concentrated to equimolar concentrations as determined by
densitometric analysis via SDS-PAGE using Fiji/ImageJ.[61] Empty AaEnc and eGFP-TP-loaded AaEnc were
similarly concentrated to equimolar concentrations for comparative NativePAGE analysis.

All NativePAGE analyses were conducted in an Invitrogen XCell SureLock using NativePAGE™ 4 to 16%
bis-tris mini protein gels and NativeMark Unstained Protein Standard from Fisher Scientific (USA) with 1x
running buffer made from 10x Tris/Glycine Buffer from Bio-Rad Laboratories, Inc. (USA). 20 µg of protein

was loaded per well, with effort to maintain equivalent amounts across all lanes for comparative analysis. NativePAGE gels were run overnight at 65 V for 16.5 hours at 4°C. The following day, gels were imaged via fluorescence imaging on a ChemiDoc Imaging System by Bio-Rad Laboratories, Inc. (USA), then stained with ReadyBlue™ Protein Gel Stain from Sigma-Aldrich (USA) and imaged and analyzed.

His-tagged eGFP and AaEnc-encapsulated eGFP-TP were buffer exchanged into varying pH buffers from pH 3.0 to pH 8.0 via five successive exchanges in 100 kDa MWCO Amicon Ultra-0.5 mL centrifugal units and incubated at 22°C for three hours. Endpoint eGFP fluorescence (488 nm/507 nm) was then measured in a BioTek Synergy H1 microplate reader at a final volume of 100 µL in Corning® 96-well flat clear bottom black polystyrene microplates (**Figure 6c**).

References

Cornejo, E., Abreu, N. & Komeili, A. Compartmentalization and organelle formation in bacteria. *Curr Opin Cell Biol* **26**, 132-138 (2014). https://doi.org:10.1016/j.ceb.2013.12.007

Sutter, M. *et al.* Structural basis of enzyme encapsulation into a bacterial nanocompartment. *Nat Struct Mol Biol* **15**, 939-947 (2008). https://doi.org:10.1038/nsmb.1473

Jones, J. A. & Giessen, T. W. Advances in encapsulin nanocompartment biology and engineering. *Biotechnol Bioeng* (2020). https://doi.org:10.1002/bit.27564

Andreas, M. P. & Giessen, T. W. Large-scale computational discovery and analysis of virus-derived microbial nanocompartments. *Nat Commun* **12**, 4748 (2021). https://doi.org:10.1038/s41467-021-25071-y

Contreras, H. *et al.* Characterization of a Mycobacterium tuberculosis nanocompartment and its potential cargo proteins. *J Biol Chem* **289**, 18279-18289 (2014). https://doi.org:10.1074/jbc.M114.570119

Giessen, T. W. & Silver, P. A. Widespread distribution of encapsulin nanocompartments reveals functional diversity. *Nat Microbiol* **2**, 17029 (2017). https://doi.org:10.1038/nmicrobiol.2017.29

McHugh, C. A. *et al.* A virus capsid-like nanocompartment that stores iron and protects bacteria from oxidative stress. *EMBO J* **33**, 1896-1911 (2014). https://doi.org:10.15252/embj.201488566

He, D. *et al.* Structural characterization of encapsulated ferritin provides insight into iron storage in bacterial nanocompartments. *Elife* **5** (2016). https://doi.org:10.7554/eLife.18972

He, D. *et al.* Conservation of the structural and functional architecture of encapsulated ferritins in bacteria and archaea. *Biochem J* **476**, 975-989 (2019). https://doi.org:10.1042/BCJ20180922

Giessen, T. W. *et al.* Large protein organelles form a new iron sequestration system with high storage capacity. *Elife* **8** (2019). https://doi.org:10.7554/eLife.46070

Nichols, R. J. *et al.* Discovery and characterization of a novel family of prokaryotic nanocompartments involved in sulfur metabolism. *Elife* **10** (2021). https://doi.org:10.7554/eLife.59288

12   Giessen, T. W. & Silver, P. A. Converting a Natural Protein Compartment into a
Nanofactory for the Size-Constrained Synthesis of Antimicrobial Silver Nanoparticles.
ACS Synth Biol **5**, 1497-1504 (2016). https://doi.org:10.1021/acssynbio.6b00117
13   Sigmund, F. *et al.* Bacterial encapsulins as orthogonal compartments for mammalian cell
engineering. *Nat Commun* **9**, 1990 (2018). https://doi.org:10.1038/s41467-018-04227-3
14   Jenkins, M. C. & Lutz, S. Encapsulin Nanocontainers as Versatile Scaffolds for the
Development of Artificial Metabolons. *ACS Synth Biol* **10**, 857-869 (2021).
https://doi.org:10.1021/acssynbio.0c00636
15   Moon, H., Lee, J., Min, J. & Kang, S. Developing genetically engineered encapsulin
protein cage nanoparticles as a targeted delivery nanoplatform. *Biomacromolecules* **15**,
3794-3801 (2014). https://doi.org:10.1021/bm501066m
16   Choi, B. *et al.* Effective Delivery of Antigen-Encapsulin Nanoparticle Fusions to
Dendritic Cells Leads to Antigen-Specific Cytotoxic T Cell Activation and Tumor
Rejection. *ACS Nano* **10**, 7339-7350 (2016). https://doi.org:10.1021/acsnano.5b08084
17   Altenburg, W. J., Rollins, N., Silver, P. A. & Giessen, T. W. Exploring targeting peptide-
shell interactions in encapsulin nanocompartments. *Sci Rep* **11**, 4951 (2021).
https://doi.org:10.1038/s41598-021-84329-z
18   Jones, J. A., Cristie-David, A. S., Andreas, M. P. & Giessen, T. W. Triggered Reversible
Disassembly of an Engineered Protein Nanocage*. *Angew Chem Int Ed Engl* **60**, 25034-
25041 (2021). https://doi.org:10.1002/anie.202110318
19   Adamson, L. S. R. *et al.* Pore structure controls stability and molecular flux in engineered
protein cages. *Sci Adv* **8**, eabl7346 (2022). https://doi.org:10.1126/sciadv.abl7346
20   Williams, E. M., Jung, S. M., Coffman, J. L. & Lutz, S. Pore Engineering for Enhanced
Mass Transport in Encapsulin Nanocompartments. *ACS Synth Biol* **7**, 2514-2517 (2018).
https://doi.org:10.1021/acssynbio.8b00295
21   Putri, R. M. *et al.* Structural Characterization of Native and Modified Encapsulins as
Nanoplatforms for in Vitro Catalysis and Cellular Uptake. *ACS Nano* **11**, 12796-12804
(2017). https://doi.org:10.1021/acsnano.7b07669
22   Sharma, A., Kawarabayasi, Y. & Satyanarayana, T. Acidophilic bacteria and archaea:
acid stable biocatalysts and their potential applications. *Extremophiles* **16**, 1-19 (2012).
https://doi.org:10.1007/s00792-011-0402-3
23   Rohwerder, T., Gehrke, T., Kinzler, K. & Sand, W. Bioleaching review part A: progress
in bioleaching: fundamentals and mechanisms of bacterial metal sulfide oxidation. *Appl*
Microbiol Biotechnol **63**, 239-248 (2003). https://doi.org:10.1007/s00253-003-1448-7
24   Pande, V., Pandey, S. C., Sati, D., Bhatt, P. & Samant, M. Microbial Interventions in
Bioremediation of Heavy Metal Contaminants in Agroecosystem. *Front Microbiol* **13**,
824084 (2022). https://doi.org:10.3389/fmicb.2022.824084
25   Reed, C. J., Lewis, H., Trejo, E., Winston, V. & Evilia, C. Protein adaptations in archaeal
extremophiles. *Archaea* **2013**, 373275 (2013). https://doi.org:10.1155/2013/373275
26   Dumorné, K., Córdova, D. C., Astorga-Eló, M. & Renganathan, P. Extremozymes: A
Potential Source for Industrial Applications. *J Microbiol Biotechnol* **27**, 649-659 (2017).
https://doi.org:10.4014/jmb.1611.11006
27   Giessen, T. W. Encapsulins. *Annu Rev Biochem* (2022). https://doi.org:10.1146/annurev-
biochem-040320-102858

28   Rodríguez, J. M., Allende-Ballestero, C., Cornelissen, J. J. L. M. & Castón, J. R.
Nanotechnological Applications Based on Bacterial Encapsulins. *Nanomaterials (Basel)*
11 (2021). https://doi.org:10.3390/nano11061467
29   Shrestha, R., Huang, G., Meekins, D. A., Geisbrecht, B. V. & Li, P. Mechanistic Insights
into Dye-Decolorizing Peroxidase Revealed by Solvent Isotope and Viscosity Effects.
ACS Catal **7**, 6352-6364 (2017). https://doi.org:10.1021/acscatal.7b01861
30   Baker-Austin, C. & Dopson, M. Life in acid: pH homeostasis in acidophiles. *Trends*
Microbiol **15**, 165-171 (2007). https://doi.org:10.1016/j.tim.2007.02.005
31   Guan, N. *et al.* Understanding of how Propionibacterium acidipropionici respond to
propionic acid stress at the level of proteomics. *Sci Rep* **4**, 6951 (2014).
https://doi.org:10.1038/srep06951
32   Parizzi, L. P. *et al.* The genome sequence of Propionibacterium acidipropionici provides
insights into its biotechnological and industrial potential. *BMC Genomics* **13**, 562 (2012).
https://doi.org:10.1186/1471-2164-13-562
33   Hsu, S. T. & Yang, S. T. Propionic acid fermentation of lactose by Propionibacterium
acidipropionici: effects of pH. *Biotechnol Bioeng* **38**, 571-578 (1991).
https://doi.org:10.1002/bit.260380603
34   Ross, J. *et al.* Pore dynamics and asymmetric cargo loading in an encapsulin
nanocompartment. *Sci Adv* **8**, eabj4461 (2022). https://doi.org:10.1126/sciadv.abj4461
35   Tang, Y. *et al.* Cryo-EM structure of Mycobacterium smegmatis DyP-loaded encapsulin.
Proc Natl Acad Sci U S A **118** (2021). https://doi.org:10.1073/pnas.2025658118
36   Loncar, N., Rozeboom, H. J., Franken, L. E., Stuart, M. C. A. & Fraaije, M. W. Structure
of a robust bacterial protein cage and its application as a versatile biocatalytic platform
through enzyme encapsulation. *Biochem Biophys Res Commun* **529**, 548-553 (2020).
https://doi.org:10.1016/j.bbrc.2020.06.059
37   Ross, J. *et al.* Pore dynamics and asymmetric cargo loading in an encapsulin
nanocompartment. *Sci Adv* **8**, eabj4461 (2022). https://doi.org:10.1126/sciadv.abj4461
38   Robert, X. & Gouet, P. Deciphering key features in protein structures with the new
ENDscript server. *Nucleic Acids Res* **42**, W320-324 (2014).
https://doi.org:10.1093/nar/gku316
39   Goddard, T. D. *et al.* UCSF ChimeraX: Meeting modern challenges in visualization and
analysis. *Protein Sci* **27**, 14-25 (2018). https://doi.org:10.1002/pro.3235
40   Jurrus, E. *et al.* Improvements to the APBS biomolecular solvation software suite.
Protein Sci **27**, 112-128 (2018). https://doi.org:10.1002/pro.3280
41   Hebditch, M. & Warwicker, J. Web-based display of protein surface and pH-dependent
properties for assessing the developability of biotherapeutics. *Sci Rep* **9**, 1969 (2019).
https://doi.org:10.1038/s41598-018-36950-8
42   Kamiyama, D. *et al.* Versatile protein tagging in cells with split fluorescent protein. *Nat*
Commun **7**, 11046 (2016). https://doi.org:10.1038/ncomms11046
43   Shinoda, H., Shannon, M. & Nagai, T. Fluorescent Proteins for Investigating Biological
Events in Acidic Environments. *Int J Mol Sci* **19** (2018).
https://doi.org:10.3390/ijms19061548
44   Shinoda, H. *et al.* Acid-Tolerant Monomeric GFP from Olindias formosa. *Cell Chem Biol*
25, 330-338.e337 (2018). https://doi.org:10.1016/j.chembiol.2017.12.005

45   Menon, B. B., Heinhorst, S., Shively, J. M. & Cannon, G. C. The carboxysome shell is
permeable to protons. *J Bacteriol* **192**, 5881-5886 (2010).
https://doi.org:10.1128/JB.00903-10
46   Penrod, J. T. & Roth, J. R. Conserving a volatile metabolite: a role for carboxysome-like
organelles in Salmonella enterica. *J Bacteriol* **188**, 2865-2874 (2006).
https://doi.org:10.1128/JB.188.8.2865-2874.2006
47   Burykin, A. & Warshel, A. What really prevents proton transport through aquaporin?
Charge self-energy versus proton wire proposals. *Biophys J* **85**, 3696-3706 (2003).
https://doi.org:10.1016/S0006-3495(03)74786-9
48   Tokmakov, A. A., Kurotani, A. & Sato, K. I. Protein pI and Intracellular Localization.
Front Mol Biosci **8**, 775736 (2021). https://doi.org:10.3389/fmolb.2021.775736
49   Lambré, C. *et al.* Safety evaluation of the food enzyme rennet paste from the abomasum
of suckling goats, lambs and calves. *EFSA J* **19**, e07006 (2021).
https://doi.org:10.2903/j.efsa.2021.7006
50   Salvachúa, D., Prieto, A., Martínez, Á. & Martínez, M. J. Characterization of a novel
dye-decolorizing peroxidase (DyP)-type enzyme from Irpex lacteus and its application in
enzymatic hydrolysis of wheat straw. *Appl Environ Microbiol* **79**, 4316-4324 (2013).
https://doi.org:10.1128/AEM.00699-13
51   Parashar, D. & Satyanarayana, T. Engineering a chimeric acid-stable α-amylase-
glucoamylase (Amy-Glu) for one step starch saccharification. *Int J Biol Macromol* **99**,
274-281 (2017). https://doi.org:10.1016/j.ijbiomac.2017.02.083
52   Razzaq, A. *et al.* Microbial Proteases Applications. *Front Bioeng Biotechnol* **7**, 110
(2019). https://doi.org:10.3389/fbioe.2019.00110
53   Mohanty, A. K., Mukhopadhyay, U. K., Grover, S. & Batish, V. K. Bovine chymosin:
production by rDNA technology and application in cheese manufacture. *Biotechnol Adv*
17, 205-217 (1999). https://doi.org:10.1016/s0734-9750(99)00010-5
54   Xu, Q. S., Yan, Y. S. & Feng, J. X. Efficient hydrolysis of raw starch and ethanol
fermentation: a novel raw starch-digesting glucoamylase from. *Biotechnol Biofuels* **9**, 216
(2016). https://doi.org:10.1186/s13068-016-0636-5
55   Johnston, D. B. & McAloon, A. J. Protease increases fermentation rate and ethanol yield
in dry-grind ethanol production. *Bioresour Technol* **154**, 18-25 (2014).
https://doi.org:10.1016/j.biortech.2013.11.043
56   Li, L. *et al.* Revealing two important tryptophan residues with completely different roles
in a dye-decolorizing peroxidase from Irpex lacteus F17. *Biotechnol Biofuels* **14**, 128
(2021). https://doi.org:10.1186/s13068-021-01978-y
57   Wiltschi, B. *et al.* Enzymes revolutionize the bioproduction of value-added compounds:
From enzyme discovery to special applications. *Biotechnol Adv* **40**, 107520 (2020).
https://doi.org:10.1016/j.biotechadv.2020.107520
58   Theillet, F. X. *et al.* Physicochemical properties of cells and their effects on intrinsically
disordered proteins (IDPs). *Chem Rev* **114**, 6661-6714 (2014).
https://doi.org:10.1021/cr400695p
59   Li, S. S. *et al.* Monitoring the Changes of pH in Lysosomes during Autophagy and
Apoptosis by Plasmon Enhanced Raman Imaging. *Anal Chem* **91**, 8398-8405 (2019).
https://doi.org:10.1021/acs.analchem.9b01250

60 Punjani, A., Rubinstein, J. L., Fleet, D. J. & Brubaker, M. A. cryoSPARC: algorithms for
rapid unsupervised cryo-EM structure determination. *Nat Methods* **14**, 290-296 (2017).
https://doi.org:10.1038/nmeth.4169
61 Schindelin, J. *et al.* Fiji: an open-source platform for biological-image analysis. *Nat*
Methods **9**, 676-682 (2012). https://doi.org:10.1038/nmeth.2019
62 Adams, P. D. *et al.* PHENIX: a comprehensive Python-based system for macromolecular
structure solution. *Acta Crystallogr D Biol Crystallogr* **66**, 213-221 (2010).
https://doi.org:10.1107/S0907444909052925
63 Pettersen, E. F. *et al.* UCSF Chimera--a visualization system for exploratory research and
analysis. *J Comput Chem* **25**, 1605-1612 (2004). https://doi.org:10.1002/jcc.20084
64 Madeira, F. *et al.* The EMBL-EBI search and sequence analysis tools APIs in 2019.
Nucleic Acids Res **47**, W636-W641 (2019). https://doi.org:10.1093/nar/gkz268
65 Suloway, C. *et al.* Automated molecular microscopy: the new Leginon system. *J Struct*
Biol **151**, 41-60 (2005). https://doi.org:10.1016/j.jsb.2005.03.010
66 Baek, M. *et al.* Accurate prediction of protein structures and interactions using a three-
track neural network. *Science* **373**, 871-876 (2021).
https://doi.org:10.1126/science.abj8754
67 Emsley, P., Lohkamp, B., Scott, W. G. & Cowtan, K. Features and development of Coot.
Acta Crystallographica Section D: Biological Crystallography **66**, 486-501 (2010).
https://doi.org:10.1107/S0907444910007493
68 Adams, P. D. *et al.* PHENIX: A comprehensive Python-based system for macromolecular
structure solution. *Acta Crystallographica Section D: Biological Crystallography* **66**,
213-221 (2010). https://doi.org:10.1107/S0907444909052925

Acknowledgements

We gratefully acknowledge funding from the NIH (R35GM133325). Research reported in this publication
was supported by the University of Michigan Cryo-EM Facility (U-M Cryo-EM). U-M Cryo-EM is grateful for
support from the U-M Life Sciences Institute and the U-M Biosciences Initiative. Molecular graphics and
analyses performed with UCSF ChimeraX, developed by the Resource for Biocomputing, Visualization, and
Informatics at the University of California, San Francisco, with support from National Institutes of Health
R01GM129325 and the Office of Cyber Infrastructure and Computational Biology, National Institute of
Allergy and Infectious Diseases.

Author Contributions

J.A.J. and T.W.G. designed the project. J.A.J. conducted the laboratory experiments and negative stain
transmission electron microscopy, while M.P.A. collected and analyzed cryo-EM data. J.A.J. and M.P.A.
built the AaEnc structural models. J.A.J. wrote the manuscript. T.W.G. processed cryo-EM data, edited the
manuscript and oversaw the project in its entirety.

Competing Interests

The authors declare no competing interests.

Additional Information

Correspondence and requests for materials should be addressed to T.W.G.