<u>Epigenetic markers of adverse lifestyle identified among night shift workers</u>

Paige M. Hulls[1,2], Daniel L. McCartney[3], Yanchun Bao[4], Rosie M. Walker[5], Frank de Vocht[1,6,7], Richard

M. Martin[1,2,8], Caroline L. Relton[1,2], Kathryn L. Evans[3], Meena Kumari[9], Riccardo E. Marioni[3], Rebecca

C. Richmond[1,2]

[1]Population Health Sciences, Bristol Medical School, University of Bristol, Bristol, UK

[2]MRC Integrative Epidemiology Unit at University of Bristol, UK

[3]Centre for Genomic and Experimental Medicine, Institute of Genetics and Cancer, University of

Edinburgh, Edinburgh, UK

[4]Department of Mathematical Science, University of Essex, Essex, UK

[5]Centre for Clinical Brain Sciences, University of Edinburgh, Edinburgh, UK

[6]NIHR School for Public Health Research

[7]NIHR Applied Research Collaboration West (NIHR ARC West)

[8]NIHR Bristol Biomedical Research Centre at the University Hospitals Bristol and Weston NHS

Foundation Trust and the University of Bristol

[9]Institute for Social and Economic Research, University of Essex, Essex, UK

Corresponding Author: Ms Paige M Hulls, Population Health Sciences, Bristol Medical School,

University of Bristol, Oakfield House, Oakfield Grove, Bristol, BS8 2BN. Email:

paige.hulls@bristol.ac.uk

21 **ABSTRACT**

Background: Epigenetic changes in the form of DNA methylation (DNAm) may act as biological
markers of risk factors or adverse health states. We investigated associations between night shift
work and established DNAm predictors of lifestyle, and compared them with those observed
between night shift work and self-reported or conventionally-measured phenotypes.

Methods: In two cohort studies, Generation Scotland (GS) (n=7,028) and Understanding Society
(UKHLS) (n=1,175), we evaluated associations between night shift work and four lifestyle factors
(body mass index, smoking, alcohol, education) using both conventionally-measured phenotypes
and DNA methylation-based scores proxying the phenotypes. DNA methylation-based measures of
biological ageing were also generated using six established "epigenetic clocks". Meta-analysis of GS
and UKHLS results was conducted using inverse-variance weighted fixed effects.

Results: Night shift work was associated with higher BMI (0.79; 95%CI 0.02, 1.56; p=0.04) and lower
education (-0.18; -0.30, -0.07; p=0.002). There was weak evidence of association between night shift
work and DNAm scores for smoking (0.06, -0.03, 0.15; p=0.18) and education (-0.24; -0.49, 0.01;
p=0.06) in fully adjusted models. Two of the epigenetic age measures demonstrated higher age
acceleration among night shift workers (0.80; 0.42, 1.18; p<0.001 for GrimAge and 0.46; 0.00, 0.92;
p=0.05 for PhenoAge).

Conclusions: Night shift work is associated with phenotypic and DNAm-based measures of lower
education. Night shift work was also related to DNAm predictors of smoking and ageing.

Keywords: shift work; smoking; BMI; alcohol; education; biomarkers; DNA methylation;
Understanding Society; Generation Scotland

INTRODUCTION

Shift work has been referred to as *"a work activity scheduled outside standard daytime hours, where*
there may be a handover of duty from one individual or work group to another"[1]. Typically, shift work
has been associated with industries that require 24-hour operation, such as essential public services
or for practical purposes. In recent years there has been an increase in the number of shift workers
in other industries[1] with approximately 19% of the working population engaged in shift work as their
main job[2] in the UK.

It has been argued that the introduction of shift work within wider industries does not consider the
health and wellbeing costs to the individual shift worker[3]. Much of the research that investigates the
health impacts of shift work has previously centred around circadian disruption, which can result in
disturbed sleep and excessive sleepiness during the work shift[4]. However, adverse health behaviours
have been found among shift workers which also put them at higher risk of disease [5-7] and shift work
has been associated with higher risk of diseases[8-11] than non-shift workers.

Recent studies have investigated epigenetic changes in the form of circulating DNA methylation
(DNAm) as an objective measure for evaluating the potential health impact of shift work[12]. While
these studies have typically focused on assessing individual sites in the genome (cytosine-
phosphate-guanine "CpG" sites), recently DNAm scores derived from methylation levels at
numerous CpG sites across the epigenome have been developed which can act as proxies for
lifestyle exposures and may predict health outcomes[13]. Self-reported health behaviours (especially
one-off measures) are subject to measurement error and bias [14], and using more objective measures
of long-term exposure could identify more robust associations with shift work.

One group of DNAm scores aims to capture the epigenetic clock. DNA methylation age (DNAm Age)
has been derived to provide an accurate estimate of biological age across a range of tissues, and at
different life stages[15]. Estimated DNAm Age exceeding true chronological age is known as 'age
acceleration' and studies suggest that DNAm age acceleration is associated with age-related health
outcomes independent of chronological age[16]. As well as the clocks developed based on age, more
recent 'second generation' epigenetic clocks have been developed based on lifestyle factors and
biomarkers which have been found to be highly predictive of both health and lifespan[17,18]. Other
DNAm scores which predict modifiable health, lifestyle and socio-economic factors include scores
developed for alcohol consumption, smoking status, BMI and education[13].

This study aimed to investigate associations of night shift work participation and a series of blood
based DNAm predictors of ageing, BMI, smoking, alcohol and education within the *Generation*
Scotland (GS) and *Understanding Society* (UKHLS) studies.

MATERIALS AND METHODS

1.1. Generation Scotland (GS)

The Generation Scotland: Scottish Family Health Study is a prospective cohort study comprising participants from the general population across five regions of Scotland. The recruitment protocol and cohort characteristics are described in detail elsewhere[19] and in the **Supplementary Methods**.

Blood DNAm was profiled using the Infinium MethylationEPIC BeadChip (Illumina Inc.) in two sample sets from GS: Set 1 and Set 2 (**Supplementary Methods**).

Methylation data were available for 777,193 CpGs measured in Set 1 (n=2,578 unrelated individuals) and 773,860 CpGs measured in Set 2 (n=4,450 unrelated individuals). The data sets had DNAm profiled at separate time points and quality control and normalisation was carried out separately. Details on the variables and how they were derived can be found in the **Supplementary Methods**.

1.2. Understanding Society (UKHLS)

The UK Household Longitudinal Study (UKHLS) (also known as Understanding Society) is a longitudinal panel survey of 40,000 UK households from England, Scotland, Wales and Northern Ireland. The recruitment protocol and cohort characteristics are described in detail elsewhere[20,21] and in the **Supplementary Methods.**

Blood DNAm was profiled using the same Infinium MethylationEPIC BeadChip (Illumina Inc.) as used by GS. Methylation data was available for 837,487 CpGs in 1,175 individuals (**Supplementary Methods**). Details on the variables and how they were derived can be found in the **Supplementary Methods**.

1.3. DNA methylation scores

We derived four DNAm scores related to BMI, smoking, alcohol consumption and education which were based on CpG sites identified in a previous study[13]. Details of the DNAm scores are shown in Table 1. For each individual, DNAm scores were calculated as the sum of methylation values at each CpG multiplied by the effect sizes obtained in a previous study[13].

We also derived six epigenetic biomarkers of ageing using previously published approaches[17,20-22]. (**Supplementary Methods**). In each case, age acceleration was defined as the residual obtained from regressing predicted age, as estimated by the epigenetic clock, on chronological age. This measure of age acceleration is independent of chronological age.

All methylation scores were standardized (mean = 0, standard deviation = 1) in both GS and UKHLS.

1.4. Statistical Analyses

We first assessed whether lifestyle factors were associated with night shift work. We performed linear regression of each phenotype (as the outcome variable) in relation to night shift work (as the exposure variable). This was with the exception for education, which was treated as the exposure variable given that education precedes night shift work. In GS, two models were run: Model 1 with adjustment for age and sex, and Model 2 with additional adjustment for other self-reported phenotypes (e.g., for smoking, the model was also adjusted for BMI, alcohol, and education). In

UKHLS, two models were run: Model 1 with adjustment for age, sex, blood processing day and batch
and Model 2 with additional adjustment for other self-reported phenotypes.

We subsequently assessed whether the lifestyle factors could be proxied with DNAm scores within
GS and UKHLS. We performed linear regression of the phenotypes (exposure variable) and lifestyle
DNAm scores for alcohol, smoking, education, and BMI (outcome variable). For these models, a
logistic regression was performed. In GS, two models were run: Model 1 with adjustment for age,
sex and 20 methylation principal components (PCs) and Model 2 with additional adjustment for
other self-reported phenotypes (e.g., for smoking related DNAm scores, the model was also adjusted
for BMI, alcohol, and education). In UKHLS, two models were run: Model 1 with adjustment for age,
sex, blood processing day and batch and Model 2 with additional adjustment for other self-reported
phenotypes.

We finally assessed whether night shift work was related to the lifestyle DNAm scores. We
performed linear regression of each lifestyle methylation score (outcome) in relation to night shift
work (exposure). This again was with the exception for education, which was treated as the
exposure. In GS, three models were run: Model 1 with adjustment for age, sex and 20 methylation
PCs; Model 2 had additional adjustments for other self-reported phenotypes (e.g., for smoking
related DNAm scores, model adjusted for BMI, alcohol, and education); and Model 3 with further
adjustment for the corresponding phenotype (e.g., for smoking related DNAm scores, the model was
also adjusted for smoking). In UKHLS, three models were run: Model 1 with adjustment for age, sex,
blood processing day and batch; Model 2 with additional adjustment of other self-reported; and
Model 3 with further adjustment for the corresponding phenotype.

We assessed whether night shift work was related to any of the six epigenetic ageing measures. We
performed linear regression of each epigenetic age acceleration (EEA) measure (as the outcome
variable) in relation to night shift work.

In GS, two models were run: Model 1 with adjustment for sex and 20 methylation PCs and Model 2
with additional adjustment for smoking, BMI, education, and alcohol. In UKHLS, two models were
run: Model 1 with adjustment for sex, blood processing day and batch and Model 2 with additional
adjustment for smoking, BMI, education, and alcohol.

We conducted a series of meta-analyses of GS Sets 1 and 2 and UKHLS using an inverse-variance
weighted fixed effects approach. For this, we used the binary measure of night shift work derived in
GS and the current night shift work variable derived in UKHLS. The I^2 statistic was used to assess
heterogeneity across the studies.

We conducted meta-analyses of: i) the self-reported or conventionally measured (in the case of BMI)
lifestyle factors in relation to night shift work, ii) the lifestyle DNAm scores in relation to night shift
work and iii) the epigenetic age acceleration measures in relation to night shift work.

RESULTS

2.1. Baseline characteristics

Summary characteristics of the participants from GS and UKHLS are presented in **Table 2**. 7,028
individuals with DNAm data were included from GS (n=2,578 in Set 1 and n=4,450 in Set 2) and 1,175
individuals from UKHLS. Set 1 and 2 of GS had a younger mean age (50.0±12.5 and 51.4±13.2)
compared to UKHLS (58.0±15.0). There was a comparable balance of men and women across the
three datasets: 38.6% males in GS Set 1, 43.7% in GS Set 2 and 41.6% in UKHLS. BMI was also broadly

comparable: 27.4±5.5 kg/m^2, 26.8±5.0 kg/m^2 and 28.1±6.2 kg/m^2, in GS Set 1, GS Set 2 and UKHLS,
respectively. GS Set 1 had more current smokers whilst UKHLS had more former smokers and GS Set
2 had more never smokers. There was a higher proportion of daily drinkers in UKHLS (16.0%)
compared with GS (12.4% in Set 1 and 13.2% in Set 2). There was also a higher proportion of less
than monthly drinkers in UKHLS (25.2%) compared with GS (17.2% in Set 1 and 16.2% in Set 2). Years
of full-time education was comparable between Set 1 and 2 of GS (13.6±3.4 and 13.8±3.4,
respectively) whilst in UKHLS it was slightly lower (12.3±5.1). In UKHLS, 1.6% of participants (n=18)
were currently working night shifts while 8.8% (n=103) had reported working night shifts over the
previous 11 years. In GS Set 1, 8.1% (n=127) of participants reported working at night for >20 hours
per week at the time of sampling, compared with 7.9% (n=193) in GS Set 2.

2.2. Lifestyle factors

2.2.1. How does night shift work relate to conventionally measured lifestyle factors?

In GS, night shift work was inversely associated with alcohol frequency in both GS Set 1 and GS Set 2
(Model 1: Effect= -0.056; 95% CI -0.095, -0.017; p=0.006 category change per hour for GS Set 1, and -
0.039; -0.072, -0.006; p=0.019 for GS Set 2), although associations attenuated with adjustment for
the other self-reported phenotypes in GS Set 2 (Model 2: -0.024; -0.057, 0.009; p=0.151) (**Table S1**).
Night shift work was positively associated with smoking status in GS Set 1 (Model 1: 0.024; 0.006,
0.042; p=0.009 category change per hour) but were more weakly associated in GS Set 2 (Model 2:
0.012; -0.006, 0.030; p=0.216), and when adjusted for the other phenotypes (**Table S1**). There were
positive associations between night shift work and BMI in both GS Set 1 (Model 1: 0.15; 0.03, 0.28;
p=0.02 kg/m^2 per hour) and GS Set 2 (Model 2: 0.23; 0.14, 0.32; p=1x10^{-6}), although these
associations attenuated when adjusted for the other phenotypes. There was an inverse association
between night shift work and education in GS Set 2 (Model 1: -0.07; -0.13, -0.02; p=0.01 hour per
year) which was less apparent in GS Set 1 (Model 1: -0.05; -0.12, 0.03; p=0.21), although there was
stronger evidence of association of the binary night shift work variable with education, based on in
both datasets (**Table S1**).

In UKHLS, no strong associations were found between current, ever, or previous night shift work and
any of the lifestyle phenotypes, although effect estimates were typically in the same direction as in
GS (**Table S2**).

In the GS-UKHLS combined meta-analysis there was evidence of a positive association between night
shift work and BMI (0.79; 0.02, 1.56; p=0.04 kg/m^2 difference between those who did and did not
work night shifts) and an inverse association between night shift work and education (-0.18; -0.30, -
0.07; p=0.002 log odds per year of education) in the fully adjusted models (Model 2) (**Figure 1**).
There was little evidence for an association between night shift work and either alcohol intake or
smoking status in a meta-analysis across the three datasets (0.00; -0.19, 0.20; p=0.97 and 0.04; -
0.19, 0.27; p=0.73 category change between those who did and did not work night shifts,
respectively). There was little evidence for heterogeneity between the study estimates (I^2 < 5%).
Results of the minimally adjusted models (Model 1) are shown in **Figure S1**).

2.2.2. How are DNA methylation biomarkers associated with lifestyle factors in GS and UKHLS?

There were positive associations between each DNAm score and its respective lifestyle phenotype in
both GS and UKHLS (**Table S3** and **S4**, respectively).

2.2.3. Is night shift work associated with DNA methylation biomarkers?

In GS, there were no clear associations between night shift work and the alcohol DNAm score (**Table**
S5). Night hours were positively associated with the smoking DNAm score in both GS Set 1 and GS
Set 2 (Model 1: 0.04 SD; 0.02, 0.06; p=1.28×10^{-4} and 0.02 SD; 0.00, 0.04; p=0.01), although
associations attenuated with further covariate adjustment in Models 2 and 3. Number of night hours
was also positively associated with the BMI DNAm score in both GS Set 1 and GS Set 2 (Model 1: 0.03
SD; 0.01, 0.05; p=0.01 and 0.03 SD; 0.01, 0.04; p=0.04). Associations attenuated with further
covariate adjustment in Models 2 and 3 (**Table S5**). Number of night hours was inversely associated
with the education DNAm score in GS Set 1 (Model 1: -0.20 hours; -0.33, -0.08; p=0.001) and more
weakly in GS Set 2 (Model 1: -0.09 hours; -0.19, 0.01; p=0.08). With further covariate adjustment,
the association persisted in GS Set 1 but was attenuated in GS Set 2 (**Table S5**).

None of the night shift work measures were strongly associated with the lifestyle DNAm scores
when UKHLS was assessed independently (**Table S6**). When estimates from GS were combined in a
meta-analysis with UKHLS, there was little evidence of night shift work associations with the alcohol
and BMI DNAm scores in the fully adjusted models (Model 3) (**Figure 2**). However, there was some
evidence of an inverse association between night shift work and the education DNAm score in all
three datasets (-0.24; -0.49, 0.01; p=0.06 log odds per year) (**Figure 2**). There was also weak evidence
for a positive association between night shift work and the smoking DNAm score (0.06 SD, -0.03,
0.15; p=0.18). There was little evidence for heterogeneity between the study estimates in the meta-
analysis (I^2 = 0%). Results of the minimally adjusted models (Models 1 and 2) are shown in **Figures S2**
and **S3**.

2.3. Epigenetic ageing

There was evidence of an association between night shift work and GrimAge in GS, but not for the
other epigenetic clocks. For GrimAge, the number of night hours was associated with higher age
acceleration in both GS Set 1 (Model 1: 0.19 years, 0.11, 0.27; p=1.18×10^{-5}) and GS Set 2 (Model 1:
0.10 years, 0.04, 0.16; p=9.45×10^{-4}) (**Table S7**). Associations with GrimAge acceleration remained,
although were partially attenuated on adjustment for smoking, BMI, education and alcohol *(Table*
S7). None of the night shift work measures were strongly associated with epigenetic age acceleration
in UKHLS **(Table S8)**.

In the GS-UKHLS meta-analysis, night shift work was associated with a 0.80 year (0.42, 1.18; p<0.001)
increase in GrimAge acceleration (**Figure 3**). There was also weak evidence of association with
PhenoAge acceleration (0.46 years; 0.00, 0.92; p=0.05). The other four epigenetic clocks showed
limited evidence of association. There was low heterogeneity between the study estimates in the
meta-analysis (I^2 < 50%).

DISCUSSION

We conducted analyses to investigate associations between night shift work and both phenotypic
and DNAm markers in two cohorts. When we assessed phenotypic traits, we found that night shift
work was associated with higher BMI and lower education. When assessing DNAm predictors of the
same traits, there was similar evidence of association of night shift work with BMI and education
DNAm scores. While the association with the BMI score was attenuated after adjusting for the
corresponding phenotype, night shift work was nominally associated with education and smoking in
fully adjusted models. Furthermore, two of the epigenetic age measures, GrimAge and PhenoAge,
demonstrated higher age acceleration among night shift workers.

The observational associations of night shift work with lower education and higher BMI have been
previously reported with comparable effect sizes[23]. While we did not find evidence of a phenotypic
association with reported smoking, the association between night shift work and the smoking
methylation score is consistent with previous studies reporting that smoking behaviour is more
common with shift work comparison to day work[8,24-27]. One study also found that shift workers were
also more likely to start smoking in comparison to their counterpart day workers[28]. We also found
little evidence of an association between night shift work and either self-reported or methylation
measures of alcohol; however, previous research has found that shift workers were more likely to
drink heavily[7].

There is growing evidence to suggest that DNAm-based measures are useful for health and lifestyle
profiling[29]. Furthermore, several studies have shown that methylation predictors can provide a more
accurate measurement of exposure than those based on self-report[30]. For example, previous studies
have shown that smoking methylation scores may provide a more accurate measure of true
exposure compared with self-reported smoking[18,30], possibly due to erroneous self-reporting, the
broad categories for reporting exposure, or because DNAm is able to capture long-term biological
changes as a result of smoking as well as secondary smoking. This is supported in part by our findings
that the education and smoking scores were still weakly associated with night shift work even after
adjusting for the corresponding self-reported exposures.

Night shift work was also associated with higher epigenetic age acceleration, as measured by two
epigenetic clocks: one predictive of mortality (GrimAge)[18] and the other predictive of physiological
dysregulation (PhenoAge)[17]. White *et al* (2019)[12] also found associations between the length of time
working night shifts and increased PhenoAge[31], although GrimAge was not investigated. Circadian
oscillators have been found to contribute to epigenetic ageing[32] and there is emerging evidence that
DNAm age estimators relate to circadian rhythm[33]. However, is should be noted that we did not find
that night shift work was associated with the Hannum and Horvath clocks, which were designed to
estimate chronological age[20,22]. This absence of association is in accordance with findings from White
et al[31] and suggests that intrinsic circadian processes are unlikely to underlie the associations
observed[17,18].

Strengths and limitations

One of the main strengths of the study is the use of GS, an epidemiological cohort study with a large
sample size with DNAm data which has also captured information on night shift work. Meta-
analysing associations with those from the UKHLS dataset also improved power to detect
associations between night shift work, lifestyle factors and DNAm predictors and indicated
consistency in associations across studies. Furthermore, we have the use of both self-reported and
DNAm markers for the same exposure, so were able to directly compare.

There were some limitations to the study. The DNAm scores used in this study were developed in GS
Set 1, which might lead to overfitting of the models. However, by using the second set of
participants in GS and the independent UKHLS datasets, we hope that this should minimise any
potential overfitting issues. There was also limited evidence for heterogeneity in the associations
observed between GS set 1 and the other studies.

We specifically assessed night shift work rather than other forms of shift work (such as
morning/evening only or rotating shift work) in relation to DNA methylation. This is because of the
previous evidence suggesting that night shift work is likely to be particularly disruptive to biological
processes and to have implications for adverse health. Whilst rotating shift work has also been
linked with circadian disruption, we did not specifically investigate this group of workers. There is
limited information on the intensity of shift work per night, e.g. whether the participant works 7pm-

7am three days a week or 7pm-10pm every day, or the direction of shift rotation (forward or
backward rotating) which may have a differential biological impact[34].

We are unable to make conclusions regarding causality of the associations observed. We cannot
exclude reverse causation since the analysis in GS and that based on current night shift work in
UKHLS was assessed cross-sectionally. Given education preceeds shift work status, the association
implies that those with less education engage in occupations that include night shift work. This is in
line with a previous study which found an inverse association between a genetic (rather than
epigenetic) risk score for higher education and shift work participation[35].

To support our findings, similar analysis should be performed in larger cohorts with DNAm data.
Future studies should also look at evaluating whether these biomarkers could provide insights into
the potential effects of night shift work on subsequent health outcomes, e.g., cardiometabolic
diseases and cancer.

CONCLUSIONS

In over 8,000 participants from two cohort studies, night shift work was associated with both
phenotypic and DNA methylation-based measures of higher BMI and lower education. DNAm
predictors of smoking and ageing were also related to night shift work. Epigenetic measures may
provide insights into the health and lifestyle profiles of night shift workers.

Table 1: Origins of lifestyle DNA methylation scores employed in the current analysis

Phenotype	DNA methylation scores	Original publication	No. of CpG sites
Alcohol	Alcohol	Epigenetic prediction of complex traits and death[13]	450
Smoking	Smoking	Epigenetic prediction of complex traits and death[13]	233
BMI	BMI	Epigenetic prediction of complex traits and death[13]	1109
Education	Education	Epigenetic prediction of complex traits and death[13]	373
Ageing	AgeAccelHorvath	DNA methylation age of human tissues and cell types[22]	353
	IEAA	DNA methylation age of human tissues and cell types[22]	353
	AgeAccelHannum	Genome-wide methylation profiles reveal quantitative views of human ageing rates[20]	71
	EEAA	DNA methylation-based measures of biological age: meta-analysis predicting time to death[21]	71
	AgeAccelPheno	An epigenetic biomarker of ageing for lifespan and healthspan[17]	513
	AgeAccelGrim	DNA methylation GrimAge strongly predicts lifespan and healthspan[18]	1,030

309 CpG: cytosine-phosphate-guanine; IEAA: Intrinsic epigenetic age acceleration; EEAA: Extrinsic
310 epigenetic age acceleration

311 **Table 2: Baseline characteristics of the participants in Generation Scotland and Understanding**
312 **Society**

Variables	Categories	Generation Scotland					Understanding Society	
		Set 1 (n=2578)		Set 2 (n=4450)			Overall (n=1175)	
		Mean	*SD*	*Mean*	*SD*		*Mean*	*SD*
Age (years)		50.02	12.5	51.39	13.2		57.97	15.0
Body mass index (kg/m^2)		27.24	5.4	26.76	4.9		28.11	6.2
Education (years of full-time education)		13.62	3.4	13.75	3.4		12.31	5.1
		N	*%*	*N*	*%*		*N*	*%*
Sex	*Male*	1583	38.6	1944	43.7		489	41.6
	Female	995	61.4	2506	56.3		686	58.4
Smoking status	*Current*	460	18.3	675	15.5		186	15.9
	Former	771	30.7	1398	32.1		487	41.6
	Never	1276	50.9	2276	52.3		498	42.5
Alcohol	*Daily*	167	12.4	334	13.2		164	16.0
	More than weekly	728	54	1402	55.6		475	43.6
	More than monthly	222	16.5	386	15.2		166	15.2
	Less than monthly	231	17.2	411	16.2		275	25.2
Night shift work	*0 hours per week*	1015	64.9	1592	65.6	*Never*	1072	91.2
	1-19 hours per week	422	27.0	643	26.5	*Ever*	103	8.8
	>20 hours per week	127	8.1	193	7.9	*Previous*	79	6.9
						Current	18	1.6

Figure 1: Associations between night shift work and phenotypes in Generation Scotland (GS) and
Understanding Society (UKHLS)

*Model 2: adjusted for age, sex, other self-reported phenotypes (e.g., for smoking, model adjusted
for alcohol, body mass index and education)

For these models, education was treated as the exposure and shift work was the outcome; effect
estimates are log odds ratios

Figure 2: Associations between night shift work and methylation scores in Generation Scotland
(GS) and Understanding Society (UKHLS)

*Model 3: adjusted for age, sex, blood processing day, rack barcode, corresponding other self-
reported phenotypes (e.g., for smoking DNAm, model adjusted for smoking, alcohol, body mass
index and education)

For these models, education was treated as the exposure and shift work was the outcome; effect
estimates are log odds ratios

Figure 3: Associations between night shift work and epigenetic age acceleration in Generation Scotland (GS)and Understanding Society (UKHLS)

Model 2: Adjusted for sex, 20 methylation PCs, smoking, alcohol, body mass index and education

For these models, education was treated as the exposure and shift work was the outcome; effect estimates are log odds ratios

SUPPLEMENTARY MATERIALS

Supplementary Table 1: Associations between night shift work and phenotypes in Generation Scotland

Supplementary Table 2: Associations between night shift work and phenotypes in Understanding Society

Supplementary Table 3: Associations between phenotypes and methylation scores in Generation Scotland

Supplementary Table 4: Associations between phenotypes and methylation scores in Understanding Society

Supplementary Table 5: Associations between night shift work and methylation scores in Generation Scotland

Supplementary Table 6: Associations between night shift work and methylation scores in Understanding Society

Supplementary Table 7: Associations between night shift work and measures of epigenetic age acceleration in Generation Scotland

Supplementary Figure 1: Associations between night shift work and phenotypes in Understanding Society and Generation Scotland – Model 1

Supplementary Figure 2: Associations between night shift work and methylation scores in Understanding Society and Generation Scotland – Model 1

Supplementary Figure 3: Associations between night shift work and methylation scores in Understanding Society and Generation Scotland – Model 2

FUNDING

GS received core support from the Chief Scientist Office of the Scottish Government Health Directorates (CZD/16/6) and the Scottish Funding Council (HR03006). Genotyping and DNA methylation profiling of the GS samples was carried out by the Genetics Core Laboratory at the Edinburgh Clinical Research Facility, Edinburgh, Scotland and was funded by the Medical Research Council UK and the Wellcome Trust (Wellcome Trust Strategic Award STratifying Resilience and Depression Longitudinally (STRADL; Reference 104036/Z/14/Z). The DNA methylation data assayed for Generation Scotland was partially funded by a 2018 NARSAD Young Investigator Grant from the Brain & Behavior Research Foundation (Ref: 27404; awardee: Dr David M Howard) and by a JMAS SIM fellowship from the Royal College of Physicians of Edinburgh (Awardee: Dr Heather C Whalley).

For UKHLS, the epigenetics methylation data were analysed by the University of Exeter Medical School (MRC grant K013807) and further facilitated by the University of Essex School of Biological Sciences.

PMH is funded by a Wellcome Trust 4-year studentship (108902/Z/15/Z). RCR is a de Pass VC research fellow at the University of Bristol. RCR, RMM and CLR are members of the MRC Integrative Epidemiology Unit at the University of Bristol funded by the Medical Research Council (MC_UU_00011/5). RMM, CLR and RCR are supported by a Cancer Research UK (C18281/A20919) programme grant (the Integrative Cancer Epidemiology Programme). RMM and CLR are supported by the National Institute for Health Research (NIHR) Bristol Biomedical Research

Centre which is funded by the National Institute for Health Research (NIHR) and is a partnership
between University Hospitals Bristol and Weston NHS Foundation Trust and the University of Bristol.
Department of Health and Social Care disclaimer: The views expressed are those of the authors and
not necessarily those of the NHS, the NIHR or the Department of Health and Social Care. FdV is
supported by the NIHR School for Public Health Research and NIHR Applied Research Collaboration
West.

Understanding Society is an initiative funded by the Economic and Social Research Council
(ES/N00812X/1) and various Government Departments, with scientific leadership by the Institute for
Social and Economic Research, University of Essex, and survey delivery by NatCen Social Research
and Kantar Public.

AUTHOR CONTRIBUTIONS

Conceptualization, Rebecca Richmond; Data curation, Daniel L. McCartney and Yanchun Bao; Formal
analysis, Paige Hulls, Daniel L. McCartney and Rebecca Richmond; Supervision, Frank De Vocht and
Richard Martin; Visualization, Richard Martin; Writing – original draft, Paige Hulls and Rebecca
Richmond; Writing – review & editing, Paige Hulls, Daniel L. McCartney, Yanchun Bao, Rosie Walker,
Frank De Vocht, Caroline Relton, Kathryn Evans, Meena Kumari, Riccardo Marioni and Rebecca
Richmond.

INSTITUTIONAL REVIEW BOARD STATEMENT

Data governance was provided by the METADAC data access committee, funded by ESRC, Wellcome,
and MRC (2015–2018: Grant Number MR/N01104X/1 2018–2020: Grant Number ES/S008349/1).
Ethical approval for Understanding Society was obtained from the National Research Service
(Understanding Society – UK Household Longitudinal Study: A Biosocial Component, Oxfordshire A
REC, Reference: 10/H0604/2). All our consents can be found here:
https://www.understandingsociety.ac.uk/documentation/health-assessment/fieldwork-documents.

INFORMED CONSENT STATEMENT

Informed consent was obtained from all subjects involved in the study.

DATA AVAILABILITY STATEMENT

According to the terms of consent for GS participants, access to data must be reviewed by the GS
Access Committee. Applications should be made to access@generationscotland.org.

CONFLICT OF INTERESTS

REM has received speaker fees from Illumina and is an advisor to the Epigenetic Clock Development
Foundation.

REFERENCES

1.   Health and Safety Executive. Managing Shiftwork. In:2006.
2.   NHS Digital. Health Survey for England - 2013. 2013.
3.   Folkard S, Tucker P. Shift work, safety and productivity. *Occupational Medicine.*
416   2003;53(2):95-101.
417   4.   Wright KP, Jr., Bogan RK, Wyatt JK. Shift work and the assessment and management of shift
418   work disorder (SWD). *Sleep Med Rev.* 2013;17(1):41-54.
419   5.   Suwazono Y, Dochi M, Sakata K, et al. A longitudinal study on the effect of shift work on
420   weight gain in male Japanese workers. *Obesity (Silver Spring).* 2008;16(8):1887-1893.
421   6.   Loef B, van der Beek AJ, Holtermann A, Hulsegge G, van Baarle D, Proper KI. Objectively
422   measured physical activity of hospital shift workers. *Scand J Work Environ Health.*
423   2018;44(3):265-273.
424   7.   Dorrian J, Heath G, Sargent C, Banks S, Coates A. Alcohol use in shiftworkers. *Accid Anal Prev.*
425   2017;99(Pt B):395-400.
426   8.   Knutsson A, Bøggild H. Gastrointestinal disorders among shift workers. *Scand J Work Environ*
427   *Health.* 2010;36(2):85-95.
428   9.   Kamdar BB, Tergas AI, Mateen FJ, Bhayani NH, Oh J. Night-shift work and risk of breast
429   cancer: a systematic review and meta-analysis. *Breast Cancer Res Treat.* 2013;138(1):291-
430   301.
431   10.   Torquati L, Mielke GI, Brown WJ, Kolbe-Alexander T. Shift work and the risk of cardiovascular
432   disease. A systematic review and meta-analysis including dose-response relationship. *Scand*
433   *J Work Environ Health.* 2018;44(3):229-238.
434   11.   Wang F, Zhang L, Zhang Y, et al. Meta-analysis on night shift work and risk of metabolic
435   syndrome. *Obes Rev.* 2014;15(9):709-720.
436   12.   White AJ, Kresovich JK, Xu Z, Sandler DP, Taylor JA. Shift work, DNA methylation and
437   epigenetic age. *Int J Epidemiol.* 2019;48(5):1536-1544.
438   13.   McCartney DL, Hillary RF, Stevenson AJ, et al. Epigenetic prediction of complex traits and
439   death. *Genome Biol.* 2018;19(1):136.
440   14.   Prince SA, Cardilli L, Reed JL, et al. A comparison of self-reported and device measured
441   sedentary behaviour in adults: a systematic review and meta-analysis. *International Journal*
442   *of Behavioral Nutrition and Physical Activity.* 2020;17(1):31.
443   15.   Fransquet PD, Wrigglesworth J, Woods RL, Ernst ME, Ryan J. The epigenetic clock as a
444   predictor of disease and mortality risk: a systematic review and meta-analysis. *Clin*
445   *Epigenetics.* 2019;11(1):62.
446   16.   Marioni RE, Shah S, McRae AF, et al. DNA methylation age of blood predicts all-cause
447   mortality in later life. *Genome Biol.* 2015;16(1):25.
448   17.   Levine ME, Lu AT, Quach A, et al. An epigenetic biomarker of aging for lifespan and
449   healthspan. *Aging-Us.* 2018;10(4):573-591.
450   18.   Lu AT, Quach A, Wilson JG, et al. DNA methylation GrimAge strongly predicts lifespan and
451   healthspan. *Aging (Albany N Y).* 2019;11(2):303-327.
452   19.   Smith BH, Campbell A, Linksted P, et al. Cohort Profile: Generation Scotland: Scottish Family
453   Health Study (GS:SFHS). The study, its participants and their potential for genetic research
454   on health and illness. *Int J Epidemiol.* 2013;42(3):689-700.
455   20.   Hannum G, Guinney J, Zhao L, et al. Genome-wide methylation profiles reveal quantitative
456   views of human aging rates. *Mol Cell.* 2013;49(2):359-367.
457   21.   Chen BH, Marioni RE, Colicino E, et al. DNA methylation-based measures of biological age:
458   meta-analysis predicting time to death. *Aging (Albany NY).* 2016;8(9):1844-1865.
459   22.   Horvath S. DNA methylation age of human tissues and cell types. *Genome Biol.*
460   2013;14(10):R115.

23.   van Drongelen A, Boot CRL, Merkus SL, Smid T, van der Beek AJ. The effects of shift work on
body weight change — a systematic review of longitudinal studies. *Scand J Work Environ*
Health. 2011;37(4):263-275.
24.   Bae M-J, Song Y-M, Shin J-Y, Choi B-Y, Keum J-H, Lee E-A. The Association Between Shift
Work and Health Behavior: Findings from the Korean National Health and Nutrition
Examination Survey. *Korean J Fam Med.* 2017;38(2):86-92.
25.   Knutsson A, Akerstedt T, Jonsson BG. Prevalence of risk factors for coronary artery disease
among day and shift workers. *Scand J Work Environ Health.* 1988;14(5):317-321.
26.   Knutsson A, Nilsson T. Tobacco use and exposure to environmental tobacco smoke in
relation to certain work characteristics. *Scand J Soc Med.* 1998;26(3):183-189.
27.   Bushnell PT, Colombi A, Caruso CC, Tak S. Work schedules and health behavior outcomes at
a large manufacturer. *Ind Health.* 2010;48(4):395-405.
28.   van Amelsvoort LG, Jansen NW, Kant I. Smoking among shift workers: More than a
confounding factor. *Chronobiol Int.* 2006;23(6):1105-1113.
29.   Hillary RF, Marioni RE. MethylDetectR: a software for methylation-based health profiling.
Wellcome open research. 2021;5:283.
30.   Zhang Y, Elgizouli M, Schöttker B, Holleczek B, Nieters A, Brenner H. Smoking-associated
DNA methylation markers predict lung cancer incidence. *Clin Epigenetics.* 2016;8:127-127.
31.   White AJ, Kresovich JK, Xu Z, Sandler DP, Taylor JA. Shift work, DNA methylation and
epigenetic age. *Int J Epidemiol.* 2019.
32.   Oh G, Ebrahimi S, Carlucci M, et al. Cytosine modifications exhibit circadian oscillations that
are involved in epigenetic diversity and aging. *Nat Commun.* 2018;9.
33.   Horvath S, Raj K. DNA methylation-based biomarkers and the epigenetic clock theory of
ageing. *Nat Rev Genet.* 2018;19(6):371-384.
34.   Stevens RG, Hansen J, Costa G, et al. Considerations of circadian impact for defining 'shift
work' in cancer studies: IARC Working Group Report. *Occupational and environmental*
medicine. 2011;68(2):154-162.
35.   Daghlas I, Richmond RC, Lane JM, et al. Selection into shift work is influenced by educational
attainment and body mass index: a Mendelian randomization study in the UK Biobank. *Int J*
Epidemiol. 2021;50(4):1229-1240.