Uncovering cellular processes in HeLa cells using unsupervised machine learning

Tom Maimon[1], Yaron Trink[1], Jacob Goldberger[1], and Tomer Kalisky [1,2]

[1]Faculty of Engineering and Bar-Ilan Institute of Nanotechnology and Advanced

Materials (BINA), Bar-Ilan University, Ramat Gan, Israel 5290002

[2] **Correspondence to:**

Tomer Kalisky

Faculty of engineering, Bar-Ilan University, Ramat Gan, Israel 52900

Office: +972-3-738-4656

Email: tomer.kalisky@biu.ac.il

ABSTRACT

Gene expression measurements taken over multiple time points are useful for

describing dynamic biological phenomena such as tissue development, regeneration,

and cancer. However, since these phenomena involve multiple processes occurring in

parallel, for example differentiation and proliferation, it is difficult to discern their

respective contributions to a measured gene expression profile at any given point in

time. Here, we demonstrate the use of un-supervised machine learning techniques to

identify and "de-convolve" processes occurring in parallel in a simple model system. We

first downloaded a published dataset of RNAseq measurements from synchronized HeLa

cells that were sampled at 14 consecutive time points. We then used Fourier analysis

and Topic modeling to identify two concurrent processes: a periodic process,

corresponding to cell division, and a transient process related to HeLa cell identity (e.g.

cervical cancer), that is presumably required for recovery from cell cycle arrest. This

study demonstrates the use of un-supervised machine learning techniques to identify

hidden states and processes in the cell.

KEYWORDS

Cell cycle, Fourier transform, Topic modeling, machine learning, HeLa cells

INTRODUCTION

Many biological phenomena such as development, regeneration, and cancer, involve thousands of cells dynamically changing their transcriptional state over time, as multiple genes are activated and repressed through a complex network of interactions. One way to characterize cell state dynamics is to perform RNA sequencing at multiple time points, either at the "bulk" or single-cell level, and obtain a series of gene expression profiles that create continuous trajectories in a latent gene expression "space". However, cell states are often a result of multiple processes occurring in parallel, for example, differentiation and proliferation, and the relative contributions of each process to the measured gene expression profiles are often difficult to "deconvolve" from each other.

Here, we demonstrate the use of an unsupervised machine learning technique for identifying and "de-convolving" processes in a simple model system. We first downloaded a dataset of bulk RNAseq measurements performed by Dominguez *et al.* on synchronized HeLa cells at 14 consecutive time points, which corresponds to approximately two cell cycles [1,2]. Then, we performed Fourier analysis and found that the majority of genes have either a periodic or a transient behavior over time. Finally, we used Topic modeling - an unsupervised machine learning technique - to show that the series of gene expression profiles can be represented as a mixture of three hidden topics, two of which are periodic, corresponding to the G1-S and G2-M phases of the cell cycle, and one with a transient behavior that is related to HeLa cell identity (e.g. cervical cancer), presumably required for recovering for cell cycle arrest. We believe that this study demonstrates the use of machine learning algorithms to deconvolve complex biological processed and identify their hidden components.

MATERIALS AND METHODS

Data:

The RNAseq dataset generated by Dominguez *et al.* [1,2] was downloaded from GEO

(accession number GSE81485) using SRA tools.

Preprocessing:

STAR [3] was used to align the reads to the human reference genome (hg38) and to

obtain the raw counts matrix. DESeq2 [4] was used to obtain the normalized counts

matrix. Then, genes whose count sum across all 14 samples was below 10 were

removed.

RNA velocity:

Velocyto [5] was used to obtain the read counts matrices of the spliced and unspliced

mRNA. DESeq2 normalization was performed for the two matrices combined.

Fourier Transform:

For each gene, Fourier analysis was performed using the R function periodogram() from

the "TSA" (Time Series Analysis) R package. This function returns a vector of frequencies

and a vector of scores (spectral densities) for each frequency. For each gene we found

the "dominant frequency", that is, the frequency with the highest score, and calculated

a "dominant frequency score" which measures the degree to which the score of this

frequency is higher than the scores of the other frequencies. Specifically, if for a

particular gene the highest score is S1 and its matching frequency is F1, the second

highest score is S2 and its frequency is F2, the third ranking score is S3 and its frequency

is F3 etc., then the "dominant frequency" for that gene is F1, and the "dominant

frequency score" is: $S1/(S2 + S3)$.

Topic modeling:

Topic modelling was performed on the normalized counts matrix using the R functions

fit_topic_model(), diff_count_analysis(), and structure_plot() from the "fastTopics" R

package [6] with k=3 topics, which we labeled as "k1", "k2", and "k3". Using these

functions, we calculated the three probability distributions $p(gene|k1)$, $p(gene|k2)$,

and $p(gene|k3)$ for every gene in each one of the three topics, as well as the

probabilities $p(k1|sample)$, $p(k2|sample)$, and $p(k3|sample)$ for each topic in every

one of the fourteen samples. We also tested higher numbers of topics but the results

did not change significantly.

One way to visualize the association between a specific gene and each of the three

topics is to calculate the "posterior probabilities" $p(k1|gene)$, $p(k2|gene)$, and

$p(k3|gene)$ and compare them. In order to do this, we used Bayes' theorem:

97
$$p(topic|gene) = \frac{p(gene|topic) \cdot p(topic)}{\sum_{topic} p(gene|topic) \cdot p(topic)}$$

where, for simplicity, we set the prior $p(topic) = 1/3$ for each topic.

RESULTS

RNA velocity analysis of periodically expressed genes reveals a time lag between

spliced and un-spliced mRNA:

We downloaded a published dataset of RNAseq measurements that were performed by

Dominguez *et al.* [1,2]. In that study, HeLa cells were first synchronized by a double

thymidine block and then RNA was collected and sequenced at 14 consecutive time

points, which corresponds to approximately two cell cycles (Figure 1A, Table S1). The

authors also identified a set of 67 "core" cell cycle related genes with periodic behavior,

and categorized them as "G1-S related" or "G2-M related" according to the stage at

which their expression is maximal (Table S1).

We started by manually inspecting the expression levels of spliced and un-spliced mRNA

of "core" cell cycle genes such as CCNE2 and UNG (G1-S related) and MKI67 and TOP2A

(G2-M related, Figure 1B) [5]. We observed that the yet-unspliced mRNA typically

precedes the spliced mRNA by 2-3 hours, as expected. We confirmed this by performing

a cross-correlation between the spliced and un-spliced mRNA levels averaged over the

67 "core" cell cycle genes (Figure 1C).

Fourier analysis identifies sets of genes with periodic and transient behaviors over time:

In order to test the periodicity of every gene in the transcriptome, we performed Fourier analysis. For each gene we plotted its "dominant frequency", that is, the frequency with the highest spectral density, versus its "dominant frequency score", which measures the degree to which the score of this frequency is higher than the scores of the other frequencies (see Methods and Figure 2A). We found that the first or second dominant frequencies - that correspond to either one or two whole cycles – contain genes with higher scores than the other dominant frequencies, whose genes have scores that are similar to those of randomized datasets (Figure S1). This indicates that most of the information in our dataset is contained in the expression levels of the genes having one of the first two dominant frequencies.

We next selected genes from the second dominant frequency (with scores > 3) and performed PCA. We found that the samples form a circular trajectory in latent space that corresponds to almost two complete cell cycles (Figure 2B). Moreover, separating the spliced and yet-unspliced mRNA expression profiles results in two circular trajectories in latent space with a phase difference between them (Figure S2). When we selected the genes from the first and second dominant frequencies we found that the samples form a trajectory in latent space with two patterns of behavior: a transient pattern along PC1 (Figure 2C) and an oscillatory pattern along PC2 and PC3 (Figure 2D-E).

Topic modeling shows that the time-dependent series of gene expression profiles can be represented as a mixture of three hidden topics, two of which are periodic, corresponding to the G1-S and G2-M phases of the cell cycle, and one with a transient behavior that is related to HeLa cell identity (i.e. cervical cancer):

"Topic models" are used in computer science to model documents that contain words

from different "topics" [7,8]. Given a set of documents and an assumed number of

topics k, the model assumes that each document can be described as a mixture of these

k latent topics with different proportions. By fitting a topic model to that set of

documents, it is possible to discover both the k latent topics and their proportions in

each individual document. In our case we assume that each cell state (=gene expression

profile) is a weighted sum of k hidden "processes" (that is, latent components), and that

these weights vary at each time point. Therefore, if each gene expression profile

corresponds to a document and each process corresponds to a topic, fitting a topic

model to our data can help us discover both the k latent processes and their proportions

at each individual time point.

We therefore fitted a topic model with K=3 topics to the 14 gene expression profiles and

found three processes with distinct behaviors, which we labeled as "k1", "k2" and "k3"

(Figure 3A and Table S2). We observed that topics k1 and k3 have an oscillatory

behavior and are periodically expressed at specific phases of the cell cycle throughout

the entire duration of the experiment (approximately two cell cycles) (Figure 3B),

whereas topic k2 is transiently over-expressed mainly during the first cycle following

release from the thymidine block.

Next, we used ToppGene [9] to perform Gene Ontology (GO) enrichment analysis on

genes that are significantly over-expressed in each one of the three topics in order to

identify them (Figure 4A-B, Tables S3-S5). We found that topic k1 over-expresses genes

that are known to be associated with the G1-S phase of the cells cycle (e.g. DNA

replication), topic k3 over-expresses genes that are known to be associated with the G2-

M phase of the cells cycle (e.g. mitosis), and topic k2 over-expresses genes related to

HeLa cell identity (e.g. cervical carcinoma).

We confirmed this by calculating the posterior probabilities p(k1|gene), p(k2|gene), and

p(k3|gene) for selected genes with specific GO terms (Figure 4C). These probabilities

represent the association between these genes and each of the three topics. Indeed, we

observed that within the 67 "core" cell cycle genes, the G1-S related genes were

associated with topic k1 and the G2-M related genes were associated with topic k3.

Similarly, genes related to HeLa cell identity and Cervical Carcinoma were more

associated with topic k2.

DISCUSSION

To summarize, we applied Fourier analysis and Topic modeling to a simple *in-vitro*

model system in order to identify its latent components. We identified two periodic

processes corresponding to the G1-S and G2-M phases of the cell cycle and one

transient process related to HeLa cell identity and cervical cancer. Our results suggest

that, following release from cell cycle arrest induced by the double thymidine block, the

cells enter a transient state lasting approximately twelve hours (samples D1-D7) in

which they activate molecular mechanisms in order to exit cell cycle arrest and resume

cycling. Since each tissue regulates cell division in a unique way, it is logical that many of

these mechanisms are specific to the tissue from which HeLa cells originated from (i.e.

cervical cancer).

We note that for some of the 67 "core" cell cycle genes we observed that the spliced

and unspliced mRNA levels arrive at their first peak (maximum or minimum) at the same

time (Figure 1B), which indicates that initially, during the few hours after release from

the thymidine block, the spliced and unspliced transcripts are synchronized in these

genes, and only after a few hours the typical time lag is acquired. This also indicates the

existence of a transient cell state in which the transcriptional timing mechanisms

operate in a way that is different from the normal cycling state. One possible

explanation is that during the first few hours after release from cell cycle arrest, the

primary mRNA transcribed from these genes is spliced, processed, and translated with

minimal delay in order to rebuild the necessary mechanisms for cell cycle resumption as
soon as possible. Further experimental studies are required to fully resolve and validate
these mechanisms.
205

ACKNOWLEDGEMENTS

We wish to thank Shahar Alon and all the member of our lab for useful comments and
suggestions.

DECLARATION OF INTEREST STATEMENT

The authors have declared that no competing interests exist.

AUTHOR CONTRIBUTIONS

Study initiation and conception – T.M. and T.K.; Data analysis - T.M. and T.K.; Topic
modeling – T.M, J.G., and T.K.; Other intellectual contribution – Y.T.; Manuscript writing
– T.M. and T.K.

FUNDING

T.M., Y.T., and T.K. were supported by the Israel Science Foundation (ICORE no. 1902/12
and Grants no. 1634/13 and 2017/13), the Israel Cancer Association (Grant no.
20150911), the Israel Ministry of Health (Grant no. 3-10146), the EU-FP7 (Marie Curie
International Reintegration Grant no. 618592), and the ICRF (Grant no. 19-101-PG). The
funders had no role in study design, data collection and analysis, decision to publish, or
preparation of the manuscript.

REFERENCES

1. Dominguez D, Tsai YH, Weatheritt R, Wang Y, Blencowe BJ, Wang Z. An extensive program of periodic alternative splicing linked to cell cycle progression. Elife. 2016;5: 1–19. doi:10.7554/eLife.10288

2. Dominguez D, Tsai YH, Gomez N, Jha DK, Davis I, Wang Z. A high-resolution transcriptome map of cell cycle reveals novel connections between periodic genes and cancer. Cell Res. 2016;26: 946–962. doi:10.1038/cr.2016.84

3. Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, et al. STAR: Ultrafast universal RNA-seq aligner. Bioinformatics. 2013;29: 15–21. doi:10.1093/bioinformatics/bts635

4. Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. Genome Biol. 2014;15: 550. doi:10.1186/s13059-014-0550-8

5. La Manno G, Soldatov R, Zeisel A, Braun E, Hochgerner H, Petukhov V, et al. RNA velocity of single cells. Nature. 2018;560: 494–498. doi:10.1038/s41586-018-0414-6

6. Carbonetto P, Luo K, Dey K, Hsiao J, Stephens M. fastTopics: fast algorithms for fitting topic models and non-negative matrix factorizations to count data. R package version 0.4-11. 2021. Available: https://github.com/stephenslab/fastTopics

7. Dey KK, Hsiao CJ, Stephens M. Visualizing the structure of RNA-seq expression data using grade of membership models. PLoS Genet. 2017;13: 1–23. doi:10.1371/journal.pgen.1006599

8. Blei DM, Ng AY, Jordan MI. Latent Dirichlet Allocation. J Mach Learn Res. 2003;3: 993–1022. Available: https://www.jmlr.org/papers/volume3/blei03a/blei03a.pdf

9. Chen J, Bardes EE, Aronow BJ, Jegga AG. ToppGene Suite for gene list enrichment analysis and candidate gene prioritization. Nucleic Acids Res. 2009. doi:10.1093/nar/gkp427

APPENDICES

Supplementary information: Supplementary text and figures.

Table S1: The gene expression dataset from Dominguez *et al.*, including the sampling time points and the list of 67 "core" cell cycle genes.

Table S2: The three topics discovered and their proportions at each time point.

Table S3: GO enrichment analysis for genes that are over-expressed in topic k1.

Table S4: GO enrichment analysis for genes that are over-expressed in topic k2.

Table S5: GO enrichment analysis for genes that are over-expressed in topic k3.

Program: A compressed directory containing programs and datasets for data visualization.

FIGURES:

Figure 1: RNA velocity analysis of periodically expressed genes reveals a time lag

between spliced and un-spliced mRNA. (A) Shown is a sketch of the experiment

performed by Dominguez *et al.* from which the RNAseq dataset was produced. HeLa

cells were synchronized, sampled at 14 consecutive time points, and their RNA was

sequenced. (B) Shown are expression levels over time of spliced and un-spliced mRNA

for selected genes that are known to be over-expressed at the G1-S phases (CCNE2,

UNG) or G2-M phases (MKI67, TOP2A) of the cell cycle. These indicate that the cells

completed two cell divisions during the experiment. It can be seen that the yet un-

spliced mRNA again precedes the spliced mRNA, as expected. Expression levels were

normalized to their maximum and spline curves were plotted through the data points to

assist visibility. (C) A cross-correlation plot between the spliced and un-spliced mRNA

levels averaged over the 67 "core" cell cycle genes, that is, periodically expressed genes

identified by Dominguez *et al.* to be related to the cell cycle in this dataset, shows a

significant time lag between the spliced and yet un-spliced mRNA.

Figure 2: Fourier analysis identifies sets of genes with periodic and transient behaviors over time. (A) For each gene, a periodogram was calculated (e.g., PLK2 and CCNE2, right panels) and its "dominant frequency" vs. "dominant frequency score" were plotted (left panel, see Methods). It can be seen that almost all of the "core" cell cycle genes (65 out

of 67 red dots) have a dominant frequency corresponding to two whole cycles. Likewise,

the first and second dominant frequencies contain genes with higher scores than the

other frequencies, whose genes have scores that are similar to those of randomized

datasets (Figure S1), indicating that the genes within the first and second dominant

frequencies contain most of the periodic information in our dataset. (B) When we select

genes from the second dominant frequency with scores > 3 and perform PCA, the

samples form a circular trajectory in latent space that represents almost two complete

cell cycles. (C-E) When we select genes from the first and second dominant frequencies

and perform PCA, the samples form a trajectory in latent space with two patterns of

behavior: a transient pattern (along PC1 axis) and a periodic pattern (in the PC2 vs. PC3

plane).

Figure 3: Topic modeling shows that the time-dependent series of gene expression profiles can be represented as a mixture of three hidden topics, two of which are periodic and one with a transient behavior. (A) Shown is a structure plot of the proportions of each topic within each sample. Topics k1 and k3 are periodically expressed during both cycles (samples D1-D14), whereas k2 is transiently over-

308 expressed during the first cycle (samples D1-D7). (B) Shown is a visualization of the

309 proportions of each topic along the circular trajectory in latent space representing the

310 two cell cycles. It can be seen that topics k1 and k3 are over-expressed in specific phases

311 during both cycles, whereas topic k2 is over-expressed during the first cycle only. Note

312 that topic k1 is expressed mainly in the upper half of the circular trajectory, whereas

313 topic k3 is expressed mainly in its lower half. (C) Shown are the expression levels of

314 three representative genes that appear at high probability in topic k2. These genes are

315 transiently over-expressed in the first cycle.

A

B

Name	Source	ID	Bonferroni	Hit count	Total list
Genes whose expression peaked at 480 min after stimulation of HeLa cells with EGF [GeneID=1950].	MSigDB C2: CGP Curated Gene Sets (v7.3)	M8729	6.13E-26	26	164
Genes modulated in HeLa cells (cervical carcinoma) by TNF [GeneID=7124] via NFKB pathway.	MSigDB C2: CGP Curated Gene Sets (v7.3)	M14435	1.80E-10	9	26
TCGA-Cervix-Primary_Tumor-Cervical_Carcinoma-Cervical_Squamous_Cell_Carcinoma-6|TCGA-Cervix / Sample_Type by Project: Shred V9	TCGA OncoMap	45a8e069369fdc7 358135bf2229def bee1ac2600	7.44E-06	9	111
TCGA-Cervix-Primary_Tumor-Cervical_Carcinoma-Non-keratinizing_Cervical_Squamous_Cell_Carcinoma-6|TCGA-Cervix / Sample_Type by Project: Shred V9	TCGA OncoMap	10955b54e4a87d0 35093a2756291ef 1d6a14ab63	1.30E-03	8	143

C

Figure 4: Gene Ontology (GO) enrichment analysis shows that the three hidden topics are related to the G1-S/G2-M phases of the cell cycle and to HeLa cell identity (i.e. cervical cancer). (A) Volcano plots showing the over-expression of each gene in each of the three topics. It can be seen that genes known to be over-expressed in the G1-S phases of the cell cycle are over-expressed in topic k1 (left panel), genes known to be over-expressed in the G2-M phases of the cell cycle are over-expressed in topic k3 (middle panel), and genes that are transiently over-expressed in the first cycle are over-expressed in topic k2 (right panel). (B) GO enrichment analysis of genes over-expressed

in topic k2 (Z-score>50) shows that topic k2 is related to HeLa cell identity and cervical

cancer (see also Tables S3-S5). (C) Shown are the posterior probabilities $p(topic|gene)$

for selected gene sets with specific GO terms. Each point represents a gene, and its

components along the k1, k2, and k3 axes are the posterior probabilities $p(k1|gene)$,

$p(k2|gene)$, and $p(k3|gene)$, which represent the association between that gene and

each of the three topics. It can be seen that topic k1 is highly associated with genes that

are known to be over-expressed in the G1-S phases of the cell cycle (left panel), whereas

topic k3 is highly associated with genes that are known to be over-expressed in the G2-

M phases of the cell cycle. Topic k2 is associated with genes that are related to HeLa cell

identity and Cervical Carcinoma (middle and right panels). Note that all points (=genes)

are located on the triangle-shaped two dimensional simplex since the sum of the

posterior probabilities for each gene equals 1.