The influence of extrachromosomal elements in the anthrax "cross-over" strain *Bacillus cereus* G9241.

Grace Taylor-Joyce[1], Shathviga Manoharan[1], Thomas Brooker[1], Carmen Sara Hernandez-Rodrıguez[2], Les Baillie[3], Petra C. F. Oyston[4], Alexia Hapeshi[1], and Nicholas R. Waterfield[♦][1].

[1]Division of Biomedical Sciences, Warwick Medical School, University of Warwick, Gibbet Hill Road, Coventry, CV4 7AL, United Kingdom

[2]Dpto. Microbiología y Ecología, Instituto BIOTECMED, Universitat de València, 46100 Burjassot, Spain.

[3]School of Pharmacy and Pharmaceutical Sciences, Cardiff University, CF10 3AT, Cardiff, United Kingdom

[4]CBR Division, Dstl Porton Down, Salisbury, SP4 0JQ, United Kingdom

♦ **corresponding author (n.r.waterfield@warwick.ac.uk)**

ABSTRACT

Bacillus cereus G9241 was isolated from a welder who survived a pulmonary anthrax-like disease. Strain G9241 carries two virulence plasmids, pBCX01 and pBC210, as well as an extrachromosomal prophage, pBFH_1. pBCX01 has 99.6% sequence identity to pXO1 carried by *Bacillus anthracis* and encodes the tripartite anthrax toxin genes and *atxA,* a mammalian virulence transcriptional regulator. This work looks at how the presence of pBCX01 and temperature may affect the lifestyle of *B. cereus* G9241 using a transcriptomic analysis and by studying spore formation, an important part of the *B. anthracis* lifecycle. Here we report that pBCX01 has a stronger effect on gene transcription at the mammalian infection relevant temperature of 37˚C in comparison to 25˚C. At 37˚C, the presence of pBCX01 appears to have a negative effect on genes involved in cell metabolism, including biosynthesis of amino acids, whilst positively affecting the transcription of many transmembrane proteins. The study of spore formation showed *B. cereus* G9241 sporulated rapidly in comparison to the *B. cereus* sensu stricto type strain ATCC 14579, particularly at 37˚C. The carriage of pBCX01 did not affect this phenotype suggesting that other genetic elements were driving rapid sporulation. An unexpected finding of this study was that pBFH_1 is highly expressed at 37˚C in comparison to 25˚C and pBFH_1 expression leads to the production of Siphoviridae-like phage particles in the supernatant of *B. cereus* G9241. This study provides an insight on how the extrachromosomal genetic elements in *B. cereus* G9241 has an influence in bacterial phenotypes.

INTRODUCTION

The *Bacillus cereus* sensu lato group contains bacterial species from diverse ecological niches and life styles (Jensen *et al.*, 2003). All members of the group, which includes *Bacillus anthracis*, *Bacillus cereus* and *Bacillus thuringiensis* among other ecologically diverse species, have a similar chromosomal genetic composition. It is clear that the distinct phenotypes and life cycles of various species are primarily defined by the different extrachromosomal elements they carry (Rasko *et al.*, 2005). The insect pathogenic *B. thuringiensis* strains encode insecticidal toxins, such as the delta endotoxins, on plasmids such as the large pBtoxis plasmid (Stein *et al.*, 2006). Members of the *B. cereus* sensu stricto species can carry a range of different plasmids that define the severity of opportunistic human infections they can cause. A good example being the pCER270 plasmid which encodes the cereulide toxin biosynthesis gene cluster (Rasko *et al.*, 2007), responsible for severe food poisoning. In the case of *B. anthracis*, the causative agent of anthrax, carriage of the anthrax toxin plasmid pXO1 and the capsule plasmid pXO2 facilitate a lethal mammalian infection life cycle. Interestingly, all *B. anthracis* strains have a frame shift mutation in the gene for the quorum sensing global regulator, PlcR, which controls the secretion of toxins and enzymes in other members of the *B. cereus* sensu lato group (Agaisse *et al.*, 1999).

B. cereus strain G9241 (from here on referred to as *Bc*G9241) was previously identified as the causative agent of an anthrax-like pneumonia infection of a metal worker in Louisiana in 1994. Phenotypic and whole genome sequencing analyses confirmed that the isolate was a *B. cereus* sensu stricto species which carries a close homologue of the *B. anthracis* plasmid pXO1, named pBCX01, encoding the tripartite anthrax toxins and their mammalian host responsive regulator AtxA1 (Hoffmaster *et al.*, 2004). *Bc*G9241 also carries a plasmid named pBC210 which, while unrelated to the *B. anthracis* plasmid pXO2, also encodes capsule genes, albeit distinct from those on pXO2 and a second *atxA1* like regulator gene designated *atxA2*. The third extrachromosomal element found in *Bc*G9241 is pBFH_1, a linear phagemid (Oh *et al.*, 2011). Although the sequence is available for pBFH_1, it is not known if/how it contributes to the lifestyle of *Bc*G9241. While *Bc*G9241 carries these three plasmids and produces anthrax toxins, *Bc*G9241 is resistant to γ-phage and penicillin and carries an intact copy of the *plcR* gene, hence why it is referred to as a *B. cereus* strain (Hoffmaster *et al.*, 2004). A small number of other *B. cereus* strains that are capable of causing an anthrax-like disease have also been isolated and are collectively referred to as 'cross-over' strains, an in-depth description of these strains given in a recent review (Baldwin, 2020). Since this review was published, two further 'cross-over' strains have been isolated from patients with an anthrax like disease (Dawson *et al.*, 2021).

B. cereus and *B. anthracis* differ from each other in the disease states they cause and also
have very different lifestyles. *B. cereus* is an opportunistic pathogen causing primarily
gastrointestinal infections in humans. *B. cereus* is also often found in the guts of many
invertebrates, which both facilitates its dissemination in the environment and provides a niche
for a potential necromonic lifestyle (Margulis *et al.*, 1998; Swiecicka and Mahillon, 2006). In
contrast, *B. anthracis* is primarily adapted to cause acute mammalian infections abandoning
the more generalist lifestyle of its recent *B. cereus* ancestor. *B. anthracis* infects its mammalian
host as a spore, often picked up from the soil by grazing livestock or wildlife through inhalation
or ingestion (Bellan *et al.*, 2013). The animal then dies from the acute anthrax disease.
Vegetative *B. anthracis* cells present in the carcass sporulate as nutrients become limited and
re-enter the soil as a spore, completing the life cycle (Lindeque and Turnbull, 1994). This life
cycle relies on *B. anthracis* being able to successfully kill its host and sporulate once its host
is deceased. Current studies of *Bc*G9241 have focused on the contribution of the anthrax
toxins and the tetrasaccharide and hyaluronic acid capsules to the ability of *Bc*G9241 to cause
a fatal anthrax-like disease (Scarff *et al.*, 2018). The contribution of chromosomal regulation
by pBCX01 or AtxA1 to virulence is less well understood and the sporulation of *Bc*G9241 has
so far not been investigated.

This study looks at how closely life cycle related phenotypic traits of *Bc*G9241 resemble those
of *B. anthracis* and how pBCX01 may influence this. To do this we have carried out RNAseq
experiments to elucidate how carriage of pBCX01 influences the transcriptome of *Bc*G9241
at the mammalian relevant temperature of 37 ˚C compared to the more environmentally
appropriate temperature of 25 ˚C. We have also assessed the sporulation phenotype of
*Bc*G9241. Sporulation is an essential part of the *B. anthracis* life cycle and is intrinsically linked
to other cellular processes, including virulence, through a complex regulatory network (Tan
and Ramamurthi, 2014). The results presented here, as well as other work done by our group
(Manoharan *et al.*, 2022), show that *Bc*G9241 phenotypes have a strong temperature
dependence allowing it to phenocopy *B. anthracis*-like behavior at 37 °C, while conversely
behaving like *B. cereus* sensu stricto at 25 ˚C. pBCX01 has a strong influence on the
transcriptome at 37 ˚C, including the suppression of the transcription of genes involved in
metabolic and biosynthetic pathways whilst simultaneously upregulating the expression of
genes involved in the import of nutrients, an effect which is not seen at 25 ˚C. We propose
that this may facilitate mammalian virulence, allowing *Bc*G9241 to have a growth advantage
whilst simultaneously restricting nutrients available to the host. In addition, an unexpected
finding of our study showed a significant increase in the pBFH_1 lysogenic phagemid
expression at 37 ˚C compared to 25 ˚C. Interestingly, close homologues of this phagemid are

also carried by other 'cross-over' strains, despite no direct phylogenetic relationships between
the chromosomes, suggesting a common role in their life cycles. We further demonstrated
that pBFH_1 produces Siphoviridae-like phage particles in the supernatant of *Bc*G9241. Our
observations regarding sporulation show that *Bc*G9241 has a temperature dependent
sporulation phenotype and sporulates rapidly compared to *B. cereus* ATCC 14579 (from here
on referred to as *Bc*ATCC 14579), particularly at 37 ˚C, which is independent of the presence
of pBCX01.

In summary, our results show that pBCX01 has allowed *Bc*G9241 to adopt a life cycle that
resembles that of *B. anthracis* and that the response of *Bc*G9241 to temperature is crucial to
this. We have also demonstrated that other genetic elements contribute to this temperature
dependent phenotype independently of pBCX01. To allow us to examine the influence of
pBCX01 on bacterial phenotypes, we constructed a plasmid cured strain designated *Bc*G9241
ΔpBCX01.

RESULTS

The influence of pBCX01 on global *Bc*G9241 transcription during
exponential growth

To determine if pBCX01 influences BcG9241 at a transcriptional level, mRNA was extracted
from cultures of *Bc*G9241 WT (WT) and *Bc*G9241 ΔpBCX01 (ΔpBCX01) growing
exponentially (OD_{600}=0.5) at both 25 °C and 37 °C. An RNAseq analysis was performed to
compare the transcriptomes of WT and ΔpBCX01. A comparison of the transcriptomes of each
strain at 25 °C and 37 °C was also performed. The full datasets generated can be seen in the
Supplementary Datasets 1-4.

The transcriptional response of *Bc*G9241 to temperature is influenced by pBCX01.
The number of genes found to be differentially expressed between 37 ˚C and 25 ˚C during
exponential growth of the WT strain was 250 (Figure 1A, shown in green), while for the
ΔpBCX01 strain the number was considerably higher, exhibiting 1,184 differentially expressed
genes (DEGs) (Figure 1A, shown in orange). Furthermore, there were only 177 genes that are
differentially expressed between 25 ˚C and 37 ˚C shared between WT and ΔpBCX01 (Figure
1A). This demonstrates that the pBCX01 plasmid has a potent influence on temperature
dependent expression of the chromosomal genes. To compare the functions of temperature
dependent DEGs of WT and ΔpBCX01, we performed gene enrichment analyses using the
STRING online tool (Szklarczyk *et al.*, 2019) and focused on the KEGG pathways that were
found to be enriched. Unfortunately, this database does not recognize the *Bc*G9241 locus

tags, therefore all genes of the *Bc*G9241 genome were assigned a *B. anthracis* Sterne locus
tag, in cases where there was amino acid homology of 90% or more. As such, *Bc*G9241 genes
that had no homologue in the *B. anthracis* genome have been excluded in this analysis.
148
In ΔpBCX01, genes significantly differentially expressed between 37 ˚C and 25 ˚C showed a
significant enrichment for genes involved in KEGG pathways associated with an increased
growth rate. These include genes involved in pathways relating to metabolism, the ribosome
and biosynthesis of secondary metabolites and amino acids (Table 1). The majority of these
DEGs had a higher relative transcript abundance at 37 ˚C compared to 25 ˚C which is
consistent with *Bc*G9241 having an increased protein production and higher growth rate at 37
˚C compared to 25 ˚C (Figure S1). In contrast, the genes with temperature dependent
differential expression in the WT strain did not show the same profile of functions associated
with an increased growth rate. In the WT there were 16 genes involved in metabolic pathways
that had a higher relative transcript abundance at 37 ˚C compared to 25 ˚C but surprisingly,
no genes involved in amino acid biosynthesis or ribosomal genes. This is unexpected as both
strains exhibit an increased growth rate at 37 ˚C in comparison to that at 25 ˚C (Figure S1).
An analysis of the genes that had a higher relative transcript abundance at 25 ˚C compared
to 37 ˚C in the WT showed there was a significant enrichment for genes involved in bacterial
chemotaxis. Five of the six of these differentially expressed bacterial chemotaxis genes were
not found to be differentially expressed in ΔpBCX01.
165

166
167
Figure 1. Venn diagrams of genes that are differentially expressed between **A)** 37˚ C and 25˚
C in *Bc*G9241 WT (green) and BcG9241 ΔpBCX01 (orange) and **B)** *Bc*G9241 WT and
ΔpBCX01 when cultured at either 25˚ C (green) or 37˚ C (red).

Table 1: KEGG pathway enrichment analysis results of genes with a significantly higher
transcript abundance at 37˚C relative to 25˚C at exponential phase in *Bc*G9241 WT and
*Bc*G9241 ΔpBCX01. A false discovery rate cut off 0.01 was used to accept a significant
enrichment of DEG in each group.

KEGG pathway	Observed gene count	No. of genes in pathway	False discovery rate
Bc G9241 WT, 37˚C > 25˚C			
Chloroalkane and chloroalkene degradation	3	11	0.0095
Glycolysis / Gluconeogenesis	5	50	0.0095
Pyruvate metabolism	5	55	0.0095
Microbial metabolism in diverse environments	9	216	0.0095
Metabolic pathways	16	657	0.0095
Bc G9241 ΔpBCX01, 37˚C > 25˚C			
Metabolic pathways	161	657	1.26E-29
Biosynthesis of secondary metabolites	92	320	1.89E-19
Biosynthesis of antibiotics	76	235	2.18E-18
Microbial metabolism in diverse environments	54	216	3.22E-09
Ribosome	25	56	3.30E-08
Carbon metabolism	36	117	3.37E-08
Biosynthesis of amino acids	39	140	6.44E-08
Pyrimidine metabolism	21	56	4.45E-06
Glycolysis / Gluconeogenesis	19	50	1.21E-05
Purine metabolism	21	68	4.63E-05
Aminoacyl-tRNA biosynthesis	14	35	0.00016
Citrate cycle (TCA cycle)	12	27	0.00026
Propanoate metabolism	13	33	0.00031
Oxidative phosphorylation	15	46	0.00045
Pyruvate metabolism	16	55	0.00074

At 37 ˚C pBCX01 has a much greater influence on the transcriptome than at 25 ˚C.
To look at the direct effect of pBCX01 on the transcriptome at both 37 ˚C and 25 ˚C a list of
genes that were differentially expressed between WT and ΔpBCX01 was generated for
cultures grown at the two temperatures. The number of genes found to be differentially
expressed between WT and ΔpBCX01 when cultures were grown at 37 ˚C was 694 (Figure
1B, shown in red), whereas when cultures were grown at 25 ˚C the number of DEG was lower
at 84 (Figure 1B, shown in green). Of the 84 genes differentially expressed between WT and
ΔpBCX01 at 25 ˚C, 62 of them were also differentially expressed at 37 ˚C between WT and

ΔpBCX01. In addition to this, many of the genes with the highest differential expression
between WT and ΔpBCX01 at 25 ˚C are also seen to be differentially expressed at 37 ˚C
(Table S1-S2). This indicates that at 37 ˚C pBCX01 has a greater influence on transcription
than it does at 25 ˚C.
188
pBCX01 suppresses metabolic and biosynthetic processes at 37 ˚C, while conversely
having a positive effect on the transcription of many transmembrane proteins.
To take a closer look at the DEGs between WT and ΔpBCX01 at 37 ˚C the list of DEGs was
split into those that had a higher relative transcript abundance in either the WT or ΔpBCX01.
As before, STRING was used to perform an enrichment analysis for each list of DEGs. Genes
with higher relative transcript abundance in WT compared to ΔpBCX01 at 37 ˚C had a
significant enrichment of genes encoding transmembrane proteins (Table 2). Many of these
transmembrane proteins have predicted transport functions, including transporters of cobalt,
formate/nitrite, phosphonates and amino acids. Four of the top 15 most differentially
expressed genes showing higher transcription in WT have a predicted transporter function
(Table S1). One of these encoded an EcsB family protein involved in regulation and secretion
of extracellular proteins. A gene encoding InhA1 also had a higher transcription level in WT
compared to ΔpBCX01 at 37 ˚C. InhA1 is a metalloprotease involved in breaking down host
proteins for nutrients (Terwilliger *et al.*, 2015) as well as mediating escape of *B. cereus* from
macrophage (Haydar *et al.*, 2018). Interestingly, pBCX01 also has a positive effect on genes
predicted to encode transcriptional regulators as well as the gene encoding the transition state
regulator AbrB.
206
The analysis of genes with higher relative transcript abundance in ΔpBCX01 compared to WT
showed that there was a significant enrichment for genes involved in metabolism, biosynthesis
of secondary metabolites and amino acids. There is a strong similarity between the DEGs
seen between 37 ˚C and 25 ˚C in ΔpBCX01 and the DEGs with higher transcript abundance
in ΔpBCX01 when grown at 37 ˚C. Both have significant enrichments in genes involved in
metabolism, biosynthesis of secondary metabolites and amino acids. This pattern of DEGs
suggests that at 37 ˚C pBCX01 is suppressing the expression of groups of genes involved in
metabolism, biosynthesis of secondary metabolites and biosynthesis of amino acids,
processes that would normally be associated with an increase in growth rate.
216
Interestingly, the five transcripts with the highest relative abundance in ΔpBCX01 compared
to WT when grown at 37 °C were all around 110 nucleotides in length and encoded adjacent
to each other in the genome (Table S1). Homologues of these five genes can be found in the

same arrangement in other *B. cereus* and *B. anthracis* genomes. The five genes are all
homologues of each other exhibiting a DUF3948, but with no known function.

To compare the overall transcriptional profiles for all four conditions (WT grown at 37 ˚C and
25 ˚C and ∆pBCX01 grown at 37 ˚C and 25 ˚C) a principal component analysis (PCA) plot
was generated using data from each biological replicate (Figure 2). Principal component one
describes 55% of the variance and principal component two describes 20% of the variance
between all samples. All biological replicates generally cluster together. There is some overlap
between the WT and ∆pBCX01 samples cultured at 25 ˚C, supporting the hypothesis that
pBCX01 has a stronger influence on the transcriptome at 37 ˚C than at 25 ˚C. Interestingly,
the PCA plot shows that the WT 37 ˚C samples cluster closer to all the 25 ˚C samples than
they do to the ∆pBCX01 37 ˚C samples. This may be due to activity of pBCX01 at 37 ˚C and
its suppressive effect on genes that would otherwise be upregulated in response to a more
optimal growth temperature.

Table 2: KEGG pathway enrichment analysis results of genes significantly affected by the
presence of pBCX01 at 37˚C exponential phase. A false discovery rate cut off or 0.01 was
used to accept a significant enrichment of DEG in each group.

KEGG pathway	Observed gene count	No. of genes in pathway	False discovery rate
Positively affected by the presence of pBCX01			
Transmembrane	98	984	3.02E-08
Transmembrane helix	96	973	4.14E-08
Negatively affected by the presence of pBCX01			
Metabolic pathways	72	657	4.14E-12
Biosynthesis of antibiotics	41	235	4.51E-12
Biosynthesis of secondary metabolites	47	320	1.09E-11
Carbon metabolism	22	117	5.36E-07
Biosynthesis of amino acids	23	140	1.85E-06
Propanoate metabolism	10	33	7.42E-05
Microbial metabolism in diverse environments	25	216	0.00013
Fatty acid metabolism	9	32	0.00027
Valine, leucine and isoleucine degradation	7	21	0.00076
Butanoate metabolism	8	29	0.00076
Fructose and mannose metabolism	5	16	0.0086
Terpenoid backbone biosynthesis	5	16	0.0086

Figure 2: A principal component analysis generated from a DEseq2 analysis of the RNAseq data from *Bc*G9241 WT grown at either 37˚C or 25˚C and *Bc*G9241 ΔpBCX01 grown at either 37˚C or 25˚C. The x axis, showing principal component 1, describes 55% of the variance in the data and the y axis, showing principal component 2, describes 20% of the variance in the data.

The impact of pBCX01 upon the nutritional requirements of *Bc*G9241

The results of the transcriptome analysis showing that pBCX01 had a negative effect on genes that are associated with an increase growth rate could suggest that at 37 ˚C, WT has a slower growth rate than ΔpBCX01. Nevertheless, this does not appear to be the case when grown in nutrient rich LB, although at 37 ˚C the growth rate of ΔpBCX01 is slightly higher than WT, the difference is negligible (Figure S1). We therefore aimed to investigate whether pBCX01 had any effect on growth rate when cultured in a media designed to resemble the nutrient availability of the mammalian host. Additionally, we wanted to specifically assess the effect of pBCX01 on the ability of *Bc*G9241 to synthesize amino acids as many amino acid biosynthesis genes were seen to be negatively affected by pBCX01. A negative effect of AtxA on amino acid genes has been previously observed in *B. anthracis* where either the *atxA* gene had been removed, or when the bacteria were grown under conditions where AtxA is known to be less active (Bourgogne *et al.*, 2003; Panda *et al.*, 2014; Raynor *et al.*, 2018). This strongly suggests that it may be the pBCX01 encoded AtxA1 activity which causes the observed down regulation of amino acid biosynthesis genes.

A minimal media designed to mimic the nutrient availability of the blood, referred to as blood serum mimic (BSM), was used (Terwilliger *et al.*, 2015) and individual amino acids were

omitted to assess the ability of WT and ΔpBCX01 to synthesize specific amino acids. *Bc*ATCC
14579 was also included in these experiments to see how the growth of WT and ΔpBCX01
compare to a typical *B. cereus* sensu stricto strain. It should be noted that for *Bc*G9241 to
grow in BSM we had to supplement this medium with 0.04 µM biotin and 0.056 mM thiamine,
or no growth was observed. It was anticipated that if WT exhibited a lower expression of the
relevant amino acid biosynthesis genes than ΔpBCX01, then it may have a slower growth rate
in media in which any of the relevant amino acids were limiting. To predict which amino acids
might be most affected by the influence of pBCX01 a KEGG metabolic pathway analysis was
used to construct a map of amino acid biosynthesis genes that were differentially expressed
between WT and ΔpBCX01 (Figure S2). Based on this analysis; arginine, histidine, isoleucine,
leucine, lysine, methionine, tryptophan and valine were selected as candidates to be
individually removed from the defined growth media for testing.
288
Our findings indicated that both WT and ΔpBCX01 were unable to grow without leucine or
valine being present in the media and that *Bc*ATCC 14579 could not grow without methionine
or valine. In media with either, arginine, isoleucine, lysine or tryptophan removed or with all
amino acids present, WT grew slightly better than ΔpBCX01 and although *Bc*ATCC 14579
initially grew faster, WT reached a higher OD_{600} than either *Bc*ATCC 14579 or ΔpBCX01. In
media with multiple amino acids removed, *Bc*ATCC 14579 was unable to grow, presumably
because of the lack of methionine. While WT grew slightly better than ΔpBCX01, there was
very little growth of either strains over 24 hours. Growth curves can be seen in Figure 3.

Figure 3: Growth in AA limiting media. Growth curves showing optical density measured at 600mn wavelength (OD$_{600}$) plotted against time of *Bc*G9241 WT (blue line), *Bc*G9241 ΔpBCX01 (red line) and *Bc*ATCC 14579 (green line). The bars show standard error. Cultures were grown at 37˚C with shaking in blood serum mimic (BSM) media with a certain amino acid omitted. The amino acid written above each graph indicates which amino acid was omitted from the growth media.

Growth at 37°C induces pBFH_1 phagemid encoded virion production in
_Bc_G9241

The transcriptomic comparison of both WT and ΔpBCX01 grown at 25 °C and 37 °C during exponential growth phase also revealed that the pBFH_1 phagemid genes become highly transcribed during growth at 37 °C compared to 25 °C. The most highly induced genes at 37 °C in WT included two encoding subunits of the predicted phage terminase protein, presumably involved in packing the phage genome into the maturing capsid structure. The large (AQ16_5899) and small ATPase subunit (AQ16_5898) genes were log2-fold 3.86 and 4.14 times higher at 37 °C compared to 25 °C respectively (Table S3). Transcription of 12 genes encoding capsid or tail proteins were also highly induced at 37 °C (Figure 4- red squares). Perhaps not unexpectedly a phage anti-repressor encoding gene (AQ16_5855) was seen to be induced at 37 °C. Furthermore, many transcriptional activators carried by pBFH_1 showed higher transcription at 37 °C, including two Xre superfamily like transcriptional regulators (AQ16_ 5850 and 5853). Interestingly a phagemid autolysin regulatory protein was also induced at 37 °C. Taken together this suggested that the lysogenic pBFH_1 phagemid enters a lytic cycle upon exposure to the higher temperature, as evidenced by the increased transcription of virion structural protein genes. The transcriptomic comparison of ΔpBCX01 grown at 25 °C and 37 °C also showed that the majority of pBFH_1 genes were more highly expressed at 37 °C in comparison to 25 °C. Of the 15 genes most highly expressed at 37 °C in comparison to 25 °C, 10 were pBFH_1 encoded genes (Table S5)

A mass spectrometry proteomic analysis of supernatants of mid-exponentially phase WT culture grown at 37 °C and 25 °C was carried out by our group (Manoharan _et al._, 2022). In agreement with the transcriptomic analysis, the proteomic analysis showed that the 8 highest proteins and 40% of the total proteins higher at 37 °C compared to 25 °C were encoded on the pBFH_1 phagemid (Table 3).

The identification of pBFH_1 structural proteins in the supernatants of WT grown at 37 °C, led us to investigate if intact phage particles were present. Particulate material was filtered from supernatants and visualised using negative stain transmission electron microscopy. Figure 5 shows the presence of a phage with Siphoviridae like morphology. The average tail dimensions are 193 x 10 nm and the average head diameter is 60 nm which is consistent with the morphology of the Siphoviridae family (Hendrix _et al._, 2012). Like other Siphoviridae the phage particles appear to have icosahedral shaped heads and a tail made up of stacked discs. This classification is consistent with the analysis of the pBFH_1 sequence by Phaster, which assigns the structural capsid proteins as Siphoviridae (Zhou _et al._, 2011). Furthermore, PCR

analysis of the phage preparations confirmed that intact phage particles carrying pBFH_1 DNA
were present. Interestingly we were not able to detect phage proteins in the cellular proteome
at 37 °C (Manoharan *et al.*, 2022) potentially indicating that the phage had been rapidly
induced and released from only a sub-population of the culture. Indeed, this is consistent with
the observation that there was no reduction in growth rate or apparent lysis of the whole culture
at 37 °C, and a testament to the sensitivity of the proteomic analysis.

Nucleotide BLAST analysis was carried out to identify any other pBFH_1-like phagemids
carried by other *B. cereus* or *B. anthracis* strains. All strains identified that exhibited homology
to the pBFH_1 genome with a percentage coverage of 61% and above were *B. cereus* strains
that were also carrying pXO1/pBCX01, pXO2 or pBC210 plasmids (Table 4). Other strains
with sequences homologous to the pBFH_1 phagemid only had a sequence coverage of 43%
or below in *B. cereus* strains and 17% or below in *B. anthracis* strains. There were two 'cross-
over' strains that did not show any sequence homology to pBFH_1, 03BB102 and 03BB108.
Interestingly neither 03BB102 nor 03BB108 carry pXO2 or pBC210-like plasmids, and
furthermore their pBCX01 plasmids have only partial homology to pXO1.

Figure 4. Genes on the pBFH_1 phagemid are more highly transcribed when *Bc*G9241 is
grown at 37 °C, compared to at 25 °C. *Bc*G9241 was grown at 25 °C and 37 °C to exponential
phase. RNA was extracted and RNA-seq performed. RNA-seq reads were processed, normalised,
compared using DESEQ2 and plotted as the log2-fold change in the transcriptional level of each
gene at 25 °C compared to at 37 °C. Almost all genes were more highly transcribed during growth
at 37 °C (red circles/squares), with the exception of a gene encoding a Fur-regulated protein (blue
circle). All known genes encoding capsid and tail proteins (red squares) were higher at 37 °C. All
points plotted are significant (p value < 0.05; n=3).

423 **Figure 5.** Transmission electron micrograph of the pBFH_1 phagemid virion isolated from
424 supernatant of a *Bc*G9241 WT culture grown at 37 ˚C. Average tail length: 192.57 nm and average
425 head diameter: 60.21 nm.

426
427

428 **Table 3.** Supernatant proteins showing the largest increase in abundance at 37 °C compared to
429 25 °C during exponential growth of *Bc*G9241 WT.

Log2-Fold Change	Protein	Gene Loci (AQ16_)
5.27	(Gp49) Phage family protein	5822
4.97	(Gp34) Putative phage major capsid protein	5824
4.53	Prophage minor structural protein	5836
4.32	(Gp14) Putative gp14-like protein	5832
4.32	N-acetylmuramoyl-L-alanine amidase family protein	5839
3.66	Phage tail family protein	5835
3.41	(GpP) Putative major capsid protein	5831
2.90	Uncharacterized protein	5823

436 **Table 4. Presence of *Bc*G9241 pBFH_1 phagemid homologues in other *B. cereus* strains**
437 **carrying pBCXO1 or pBCXO2.** Presence or absence of pXO1/pBCX01, pXO2 and pBC210 is
438 also indicated.
439

Strain name	pXO1/pBCXO1 Present	pXO2 present	pBC210 Present	G9241 pBFH_1 homologues		Sample source	NCBI accession no.	Submitted by
G9241	Yes	No	Yes	yes	-	Human, USA	CP009590.1	Hoffmaster, 2004
FL2013	Yes (2.5 kb deletion)	No	only partial sequence present	yes	99.26% identity, 73% coverage	Human, USA	JHQN01000000	Marston, 2016
BcLA2007	Yes	No	Yes	No	No pBFH_1 homologues	Human, USA	MUBB00000000	Pena-Gonzalez, 2017
03BB87	No	No	Yes	yes	100% identity and coverage	Human, USA	CP009941.1	Johnson, 2015
03BB102	Only partial homology	No	No	No	No pBFH_1 homologues	Human, USA	CP001407	Hoffmaster, 2006
Elc2	Yes	No	No		unknown	Human, USA	?	Wright, 2011
03BB108	Only partial homology. No Anthax tripartite toxin genes.	No	No	No	No pBFH_1 homologues (pBFI_4: predicted phagemid, 8% coverage. pBFI_5: predicted phagemid, 3% coverage)	Soil, USA	GCA_000832865.1	Hoffmaster, 2006
BC-AK	Yes	Yes	No	yes	95.28% identity, 61% coverage	Kangaroo, China Zoo	NZ_CP020937	Dupke, 2019

440
441

*Bc*G9241 sporulates rapidly in comparison to *Bc*ATCC 14579 and shows a temperature dependant phenotype.

B. anthracis sporulates rapidly under laboratory conditions (Koch, 1876), a phenotype
necessary for its obligate infective lifecycle (Ross and Billing, 1957). To determine if the WT
strain shares this sporulation phenotype, it was visualised using microscopy after 24 and 48
hours growth on LB agar at both 25 °C and 37 °C. For comparison, *Bc*ATCC 14579 was also
imaged under the same conditions (Figure 6A). After 24 hours growth at 25 °C WT cells
remained mainly vegetative with little or no endospore formation. However, after 48 hours
growth, a limited number of endospores could be observed. In contrast, after 24 hours growth
at 37 °C, the WT culture was almost 100% mature spores and we could observe no remaining
vegetative cells. Conversely even after 48 hours growth on LB agar at 37 ˚C, *Bc*ATCC1479
displayed no endospore formation at either temperature. These findings show a striking
difference in the sporulation tendency of WT compared to *Bc*ATCC1479, and also that WT
sporulated more rapidly at 37 °C compared to 25 °C.
456

To quantify the temperature dependant sporulation of WT, a sporulation assay was carried
out in LB broth at 37 ˚C and 25 ˚C. Representative microscopic images were taken to confirm
results of the sporulation assay (Figure S3). After 24 hours growth there was a significant
difference in percentage of spores between the WT cultures grown at 37 °C and those at 25
°C. After 24 hours at 37 °C, measurements suggested 114% of cells had formed spores,
compared to a negligible percentage at 25 °C (Figure 5B). The phenomenon of a >100% spore
count (using this assay) has been observed in previous publications (Dale *et al.*, 2018), and it

is believed to arise from the germination of spores and cell division, after boiling but before

plating. At 48 hours there was still a significantly higher percentage of spores formed at 37 ˚C

than at 25 ˚C. The results of the sporulation assay are consistent with the microscopy analysis

of samples (Figure S3). When comparing the proportion of spores formed in liquid and solid

media we can see that WT sporulated around 24 hours earlier on the LB agar (Figure 5A,

Figure S3).

Figure 6. Temperature dependent sporulation of *Bc*G9241 and *Bc*ATCC 14579 in LB. A) *Bc*G9241 WT and *Bc*ATCC 14579 were grown for 24 or 48 hrs on LB agar at either 25 °C or 37 °C. A loop of bacterial lawn was taken and used to inoculate 25 µl of PBS. 5 µl of this suspension was mounted and images were taken at 100x magnification. No spore formation is seen under any condition for *Bc*ATCC 14579. While *Bc*G9241 WT had not formed any spores within 24 hrs at 25 °C, by 48 hrs growth endospores have begun to form. In contrast, after only 24 hrs growth at 37 °C, fully mature spores had formed and few if any vegetative cells remained. **B**) *Bc*G9241 WT was cultured in LB broth at 25 °C and 37 °C with aeration. At 24 and 48 hours samples were collected and a sporulation assay was performed to quantify percentage of spores. Data is plotted as the % of the culture that had formed spores. After 24 hours at 37 °C, 100% of the cell population had formed spores. However, at 25 °C, only 0.06% of cells have formed spores. * indicates significance to a p-value < 0.05 using Welch's t test. **indicates significance to a p-value of 0.0014 using an ordinary one-way ANOVA. Error bars are one standard deviation and all tests are in triplicate (n=3). See text for an explanation of readings of >100 %.

pBCX01 is not involved in the rapid sporulation phenotype of *Bc*G9241.

A sporulation assay comparing the percentage spore formation of WT, ΔpBCX01 and *Bc*ATCC 14579 at 37 °C and 25 °C was carried out to determine whether the presence of pBCX01 contributes to the rapid sporulation phenotype of *Bc*G9241 when grown under sporulation inducing conditions. Plasmid pBCX01 carries a Rap-Phr system with 100% identity to that encoded on pXO1 from *B. anthracis* (Bongiorni *et al.*, 2007). This system is thought to prevent sporulation of *B. anthracis* under infection relevant conditions when optimal toxin production and rapid vegetative growth is advantageous. Therefore, these experiments were carried in the sporulation inducing media, modified G medium (MGM), to allow the effect of pBCX01 to be studied under conditions that don't inhibit spore formation.

WT, ΔpBCX01 and *Bc*ATCC 14579 were grown in MGM at 37°C for 8 hours and 25°C for 20 hours before the percentage spore formation was assessed. These assays showed that the sporulation phenotype of *Bc*G9241 is not significantly affected by the presence of pBCX01 at either 37 °C or 25 °C (Figure 7). Both WT and ΔpBCX01 form a significantly higher percentage of spores than *Bc*ATCC 14579 at 37 °C or 25 °C at the time points sampled (Figure 7).

As *Bc*G9241 sporulates rapidly at 37 °C compared to *Bc*ATCC 14579 we decided to compare the sporulation processes of each strain. We constructed fluorescent reporter strains, designed to express GFP at two critical transcriptional check-points in the sporulation cascade. The aim was to investigate which point in the sporulation cascade is influenced by a temperature difference. Two promoters were selected to create the GFP reporters that would be specifically activated at (i) the initiation of the sporulation cascade and (ii) completion of DNA engulfment into the forespore. These two promoters are activated by the regulators; Spo0A, in its phosphorylated form, (Strauch *et al.*, 1992) and Sigma G (Regan *et al.*, 2012) respectively. These reporter strains showed that both WT and ΔpBCX01 activated the sporulation cascade at a slightly earlier time point than *Bc*ATCC 14579 and reached the late forespore stage at around 3 hours before *Bc*ATCC 14579 when grown at 37°C in MGM (Figure S4-S5). For brevity the findings of this more detailed analysis have been included in the supplementary data.

Figure 7. Percentage spore formation in MGM. *Bc*G9241 WT, *Bc*G9241 ΔpBCXO1 and

*Bc*ATCC 14579 were grown in modified G medium (MGM) for 8 hours at 37°C (**A**) or 20 hours

at 25°C (**B**) with shaking at 200 rpm. Percentage of spores in each samples was calculated

using a sporulation assay. The black and grey bars show the average percentage of spores

from three biological replicates with the standard error of the mean as a line above the bar.

The horizontal lines indicate the statistical comparisons made between two samples using an

unpaired, Welch's t-test where 'not significant' (ns) = P > 0.05, * = P ≤ 0.05, ** = P ≤ 0.01, ***

= P ≤ 0.001.

DISCUSSION

An analysis of the influence of both pBCX01 and temperature on the transcriptomic landscape

of *Bc*G9241 clearly demonstrates that pBCX01 has a strong influence on the transcriptome of

*Bc*G9241 at the mammalian infection relevant temperature of 37 °C. Notably there is

significantly less influence of this plasmid at the more environmentally appropriate

temperature of 25 °C. Our findings are consistent with experiments in *B. anthracis* where

pXO1 has been shown to contribute to both plasmid and chromosomal gene regulation under

mammalian infection relevant conditions (McKenzie *et al.*, 2014).

Comparing the differential expression at 37 °C and 25 °C between WT and ΔpBCX01, we see
that under 37 °C growth conditions removal of pBCX01 had a positive effect on genes one
would associate with an increase growth rate in comparison to 25 °C. Unexpectedly, when the
plasmid is present (in the WT strain) the same pattern was not seen even though both strains
exhibit a similar increase in growth rate at 37 °C compared to 25 °C. This suggests that the
rich growth medium used is able to provide sufficient resources to support this increased
biomass production irrespective of changes in the transcriptome. The influence of pBCX01 on
the transcriptome results in the suppression of transcription of genes involved in various
aspects of central metabolism and the biosynthesis of secondary metabolites and amino
acids. This effect is likely more important during an actual infection process. Consistent with
our findings, previous studies have shown that in *B. anthracis,* AtxA has a negative effect on
the expression of amino acid biosynthesis genes (Bourgogne *et al.*, 2003; Panda *et al.*, 2014).
Taken together these results suggest that at 37 °C pBCX01 is having a large effect on gene
expression and is acting to suppress the metabolic and biosynthetic processes associated
with a higher growth rate. Whereas at 25 °C, the plasmid is having a minimal effect on the
transcriptome and there is no evidence of a similar suppression of metabolic and biosynthetic
genes we see at 37 °C. Another temperature dependent effect of pBCX01 on the
transcriptome appears to be a suppression of the transcription of bacterial chemotaxis genes
at 37 °C but not at 25 °C.
580
Although pBCX01 has a negative effect on the expression of genes encoding amino acid
biosynthesis proteins, the WT was still able to synthesize most amino acids not made available
in the defined growth media, and indeed even had a growth advantage over ΔpBCX01. It
therefore appears that the level of amino acid biosynthesis enzymes in WT, is still sufficient to
service the need for biomass production *in vitro*. The suppressive effect of pBCX01 may
represent an adaptation to favor amino acid import during a mammalian infection, rather than
expend unnecessary energy in *de novo* synthesis. In support of this, we note the expression
of amino acid transporters was enhanced by the presence of pBCX01, again suggesting that
pBCX01 promotes the uptake of available nutrients in favor of *de novo* biosynthesis.
590
Another striking effect of the carriage of pBCX01 at 37 °C is an increase in the transcription of
many diverse transmembrane proteins. Furthermore, both pBCX01 and temperature had an
effect on the expression of multiple genes associated with maintenance of the cell membrane
of *Bc*G9241. This implies that both the virulence plasmid and temperature both effect how
*Bc*G9241 interacts with its environment. We note that pBCX01 encodes the *hasACB* operon
for the synthesis of a hyaluronic acid capsule as well as other surface associated proteins

such as BslA and an S-layer associated protein, which has been shown to mediate adhesion
to host cells (Wang *et al.*, 2013; Scarff *et al.*, 2018). It is possible that one or more of the
uncharacterized transmembrane proteins showing a higher transcript abundance in the WT
strain might be involved in the secretion or assembly of the hyaluronic acid capsule or other
plasmid encoded surface proteins. Another notable consequence of the carriage of pBCX01
is an increase in transcription of several genes encoding proteins containing helix-turn-helix
domains. Indeed, two of these genes were in the top three most differentially expressed genes
showing higher levels when pBCX01 is present. This suggests that the presence of pBCX01,
and possibly AtxA, can activate the expression of other DNA binding proteins that likely
regulate chromosomal genes.

Modulation of metabolism has been shown to be a strategy used by many bacterial pathogens
to aid successful infection (Olive and Sassetti, 2016). We propose that the influence of AtxA,
or other pBCX01 encoded factors, on the transcription of chromosomal genes may facilitate
virulence in *Bc*G9241. This may act by promoting the use of imported host nutrients, while at
the same time suppressing unnecessary biosynthesis of required nutrients and metabolites
for growth and biomass production. This strategy may give *Bc*G9241 a growth advantage
during infection and therefore increasing pathogenic potential. Indeed, higher numbers of
tissue resident bacilli has previously been associated with an increased likelihood of fatal
outcomes in anthrax infections (Guarner *et al.*, 2003). Furthermore, depriving and weakening
the host cells of nutrients through increased import has also been proposed to constitute a
specific pathogenic strategy in a phenomenon named "nutritional virulence" (Abu Kwaik and
Bumann, 2013). Finally, certain metabolic products produced by a pathogen can also have
the potential to activate an immune response (Wynosky-Dolfi *et al.*, 2014; Gaudet *et al.*, 2015).
Therefore, the modulation of the metabolism of *Bc*G9241 by pBCX01 may serve to limit the
production of potential immune reactive metabolic products.

Sporulation is an essential part of the *B. anthracis* lifestyle and it has been shown to sporulate
rapidly under favorable conditions (Davies, 1960). Analysis of *Bc*G9241 sporulation showed
that it is also able to sporulate rapidly compared to *Bc*ATCC 14579 in both nutrient rich and
nutrient poor sporulation media. This difference is also more pronounced at 37 °C than at 25
°C. Whilst there is a clear difference between the sporulation phenotypes of *Bc*G9241 and
*Bc*ATCC 14579, our findings demonstrate that it is not facillitated by AtxA or the presence of
pBCX01 as ΔpBCX01 exhibits the same rapid sporulation phenotype. This suggests that there
are other genetic elements driving the rapid sporulation phenotype in *Bc*G9241. Interestingly,
this work has identified that the pBFH_1 plasmid is highly expressed at 37 ˚C in comparison

to 25 °C and leads to the production of Siphoviridae-like phage particles. There have been multiple examples of bacteriophages enhancing the sporulation efficiency of their host and it is possible that higher expression and production of pBFH_1 proteins at 37°C is influencing the regulation of sporulation (Silver-Mysliwiec and Bramucci, 1990; Peng and Yuan, 2018). As shown in Table 4, pBFH_1-like plasmids are maintained by multiple 'cross-over' strains suggesting that it could play an essential role in the biology of a *B. cereus* strain causing an anthrax-like disease.

AUTHORS AND CONTRIBUTORS

Grace Taylor-Joyce[1]: Planned and performed experiments and wrote much of the manuscript

Shathviga Manoharan[1]: Assisted in experiments and wrote parts of the manuscript

Thomas Brooker[1]: Planned and performed some of the experiments.

Carmen Sara Hernandez-Rodrıguez[2]: Planned and performed some of the experiments.

Les Baillie[3]: Provided certain bacterial strains and provided advice on handling them.

Petra Oyston[4]: Provided advice on handling the pathogenic strains and assisted in interpreting the results.

Alexia Hapeshi[1]: Assisted in some experimental work and in interpreting certain results.

Nicholas R. Waterfield[1]: Experimental planning, secured funding, assisted in interpreting results and provided guidance and edits for writing the manuscript.

[1]Division of Biomedical Sciences, Warwick Medical School, University of Warwick, Gibbet Hill Road, Coventry, CV4 7AL, United Kingdom

[2]Institut Universitari de Biotecnologia i Biomedicina, Departament de Genètica, Facultad de Ciències Biològiques, University of Valencia, 46100 Burjassot, Valencia, Spain

[3]School of Pharmacy and Pharmaceutical Sciences, Cardiff University, CF10 3AT, Cardiff, United Kingdom

[4]CBR Division, Dstl Porton Down, Salisbury, SP4 0JQ, United Kingdom

CONFLICTS OF INTEREST

The authors declare no conflicts of interest.

FUNDING INFORMATION

This research was funded in whole or in part by the funders and grant numbers below. For the purpose of open access, the author has applied a Creative Commons Attribution (CC BY) licence (where permitted by UKRI, 'Open Government Licence' or 'Creative Commons Attribution No-derivatives (CC BY-ND) licence' may be stated instead) to any Author Accepted Manuscript version arising from this submission.

GTJ was funded by the BBSRC MIBTP doctoral training programme at Warwick University,
UK. **SM** and **TB** were funded by the WCPRS scholarship programme provided by Warwick
University, with funding contributions from DSTL (MoD) at Porton Down, UK (DSTL project
references; DSTLX1000093952 and DSTLX-1000128995). **CSHR** was funded by an EU Marie
Curie fellowship awarded while at the University of Bath, UK (FP7-PEOPLE-2010-IEF project
273155). **AH** was funded by a start-up financial package awarded to **NRW** upon starting at
Warwick University Medical School, UK. **LB** is funded by Cardiff school of biological sciences,
UK. **PO** is funded by the DSTL at Porton Down, UK. **NRW** is funded by Warwick University,
UK.

ACKNOWLEDGEMENTS
We would like to thank the Petra Oyston at the DSTL for advice throughout this project.

MATERIALS AND METHODS

Bacterial strains and culture conditions. The bacterial strains used in this study were
*Bc*G9241 (Hoffmaster *et al.*, 2004), *Bc*G9241 ΔpBCX01 and *B. cereus* reference strain ATCC
14579 (American Type Culture Collection, Manassas, Va.). Overnight cultures were grown in
5 ml of Luria Bertani (LB) broth with appropriate antibiotics at either 25 °C or 37 °C. Before
Bacillus cultures were seeded, pre-cultures were used to synchronise bacterial cell growth by
diluting overnight cultures 1 in 100 in 5 ml LB and grown to exponential phase. Pre-cultures
were then diluted to an $OD_{600} = 0.005$ in a specified media and volume.

RNA extraction for RNAseq. Overnight cultures of *Bc*G9241 and *Bc*G9241 DpBX01 were
grown at 25 °C or 37 °C. Cultures were diluted to $OD_{600} = 0.05$ and incubated at the
corresponding temperature until they had grown to $OD_{600} = 0.5$. This pre-culture was diluted to
$OD_{600} = 0.005$ and grown in 50 ml of LB broth. Cells were cultured to mid exponential phase
and collected by centrifugation at 10,000g for 1 minute and pellets were resuspended in 5x
volume of Qiagen RNA-protect. Resuspended pellets were stored at -20 °C or used
immediately. 1 ml of QIAzol (Qiagen) was added to each pellet suspension before being
transferred to Lysing Matrix B tubes (MP Biomedicals). Cells were lysed using the FastPrep®-
24 Classic instrument with a COOLPREP™ adapter (MP Biomedicals). Bead beating was
conducted at 6 ms^{-1} for 40 s for 2 cycles, with a 300 s pause between cycles. Lysates were
centrifuged for 1 minute at 10,000 g and supernatant extracted. Cell lysates were processed
for total RNA extraction using the RNeasy Micro Kit (Qiagen) and treatment with RNase-free
DNase using the Ambion™ DNase I protocol (ThermoFisher Scientific), following the

manufacturer's protocols. Details of RNAseq library preparation, sequencing and analysis
methods can be found in supplementary methods.

Generation of pBCX01 plasmid-cured *Bc*G9241. For the generation of strain *Bc*G9241
ΔpBCX01 (pBCX01$^-$/pBC210$^+$/pBFH1$^+$), a liquid culture of *Bc*G9241 was grown overnight in
LB broth at 42 ⁰C with shaking at 200 rpm. On each of six consecutive days, the culture was
diluted 1/1000 into fresh LB broth and grown as above. Dilutions of the final culture were plated
onto NBY-Bicarbonate agar (Wilson *et al.*, 2011) and growth at 37 ⁰C in 5 % CO_2 to promote
capsule production. Individual colonies were passaged several times onto fresh NBY-
bicarbonate plates and incubated as described above. Since AtxA, positive regulator of the
capsule operon (Uchida *et al.*, 1997), is encoded in pBCX01 plasmid, pBCX01$^-$ mutant
colonies show a rough appearance, distinguishable from the smooth colonies producing the
capsule. For PCR screening of candidate strains, individual rough colonies were used to
conduct colony-PCR using GoTaq Green master mix (Promega) and a primer set unique to
different regions on each of the *Bc*G9241 plasmids pBCX01 and pBC210. Control reactions
designed to amplify a region of *Bc*G9241 chromosome (*plcR* and ribosomal *16S* genes) and
pBFH1 plasmid were included (Table S5). PCR assays were done under the following
conditions: an initial step of 10 min at 95 ⁰C followed by 30 cycles of 95 ⁰C for 5 min, 45 ⁰C for
1 min, and 72 ⁰C for 1.5 min; final extension for 7 min. As a result of this screening *Bc*G9241
ΔpBCX01 was selected for further experiments. This strain was sequenced on an illumina
MiSeq platform to confirm the loss of pBCX01 and a lack of any other genomic alterations (not
shown).

Extraction of pBFH1 phage particles and electron microscopy.
Exponential pre-cultures of *Bc*G9241 were diluter to OD 0.005 in 50 ml of LB broth and grown
to exponential phase at 37 ˚C. The culture was then spun at 6,000 g for 10 min. The
supernatant was removed and filtered through an 100K Amicon Ultra-15 centrifugal filter
column (Millipore) by centrifugation at 4,500 g until all supernatant had been filtered. 500 µl of
SM buffer (100mM NaCl, 8mM $MgSO_4$ and 50mM Tris HCl) was added to the column filter
and vortexed vigorously to resuspend any bacteriophage caught by the filter. The sample was
removed from the column and filtered through a 0.2 µm filter to remove any bacteria. Samples
were stored at 4 ˚C. To confirm the presence of pBFH_1 phagemid particles, samples were
first treated with DNase I (NEB) then heated to 75˚C for 10 min to inactivate the DNase. To
break open any phage particles, proteinase K was added and incubated at 37˚C for 1 hour
before boiling at 100˚C for 1 hour. The samples were then screened for the presence of
phagemid DNA using two sets of primers (pBFH1_5877_Foward, pBFH1_5877_Reverse,
pBFH1_5899_Foward and pBFH1_5899_Reverse) each targeting a pBFH_1 phagemid gene.

A primer pair targeting the *fusA* gene was also used to control for the presence of non-phage
particle DNA. Phage samples were negatively stained using 2% uranyl acetate for 4 minutes.
Micrographs were collected on the JEOL2011 electron microscope using a US1000 CCD
camera (Gatan Inc) at various levels of magnification.

Bacterial nutritional studies. Pre-cultures of *Bc*G9241 WT, *Bc*G9241 ΔpBCX01 and
*Bc*ATCC 14579 grown in LB broth were first washed in BSM media (Terwilliger *et al.*, 2015)
(Table S6), where all amino acids of interest had been omitted, three times before diluting to
an OD600 = 0.1 in media with the appropriate amino acid omitted. Samples were then
aliquoted into a 96 well plate with 100 ml per well with three technical replicates and grown at
37°C with 700rpm shaking. OD600 readings were taken every 15 min for 24 hours. Three
biological replicates were carried out for each sample with one biological replicate per plate.
All OD600 readings were normalized to a blank media reading and plotted.

***Bacillus* sporulation assay.** All cultures were grown at either 37 °C or 25 °C. Pre-cultures
were diluted to an OD = 0.005 into 50 ml LB broth or 20 ml of MGM broth (Table S6) and
grown with 200 rpm shaking before samples were taken at specified time points. Samples
were diluted and plated on to LB agar to calculate total number of CFU before heat shocking
at 65°C for 30 min to kill any *Bc*G9241 vegetative cells and 70 °C for 15 min to kill any *Bc*ATCC
14579 vegetative cells and plating again onto LB agar to calculate number of spores able to
form colonies. Percentage of spores was then calculated.

Microscopy. 2 µl of sample was applied to a prepared agarose pad and a cover slip placed
over them. Images were captured on a Leica DMi8 premium-class modular research
microscope with a Leica EL6000 external light source, using an ORCA-Flash4.0 V2 Digital
CMOS Hamamatsu Camera at 100x magnification.

REFERENCES

Abu Kwaik, Y. and Bumann, D. (2013) 'Microbial quest for food *in vivo*: "Nutritional virulence" as
an emerging paradigm', *Cellular Microbiology*, 15(6), pp. 882–890.

Agaisse, H., Gominet, M., Økstad, O.A., Kolsto, A.-B. and Lereclus, D. (1999) 'PlcR is a pleiotropic
regulator of extracellular virulence factor gene expression in *Bacillus thuringiensis*', *Molecular
Microbiology*, 32(5), pp. 1043–1053.

Arantes, O. and Lereclus, D. (1991) 'Construction of cloning vectors for *Bacillus thuringiensis*',
Gene, 108, pp. 115–119.

Baldwin, V.M. (2020) 'You Can't B. cereus – A Review of *Bacillus cereus* Strains That Cause

Anthrax-Like Disease', *Frontiers in Microbiology*, 11(1731).

Bellan, S.E., Turnbull, P.C.B., Beyer, W. and Getzd, W.M. (2013) 'Effects of Experimental
Exclusion of Scavengers from Carcasses of Anthrax-Infected Herbivores on *Bacillus anthracis*
Sporulation, Survival, and Distribution', *Applied and Environmental Microbiology*, 79(12), pp.
3756–3761.

Bongiorni, C., Stoessel, R. and Perego, M. (2007) 'Negative regulation of *Bacillus anthracis*
sporulation by the Spo0E family of phosphatases', *Journal of Bacteriology*, 189(7), pp. 2637–2645.

Bourgogne, A., Drysdale, M., Hilsenbeck, S.G., Peterson, S.N. and Koehler, T.M. (2003) 'Global
Effects of Virulence Gene Regulators in a *Bacillus anthracis* Strain with Both Virulence Plasmids',
Infection and Immunity, 71(5), pp. 2736–2743.

Dale, J.L., Raynor, M.J., Ty, M.C., Hadjifrangiskou, M. and Koehler, T.M. (2018) 'A dual role for
the *Bacillus anthracis* master virulence regulator AtxA: Control of sporulation and anthrax toxin
production', *Frontiers in Microbiology*, 9(482).

Davies, D.G. (1960) 'The influence of temperature and humidity on spore formation and
germination in *Bacillus anthracis*', *Journal of Hygiene*, 58(2), pp. 177–186.

Dawson, P., Schrodt, C.A., Feldmann, K., Traxler, R.M., Gee, J.E., Kolton, C.B., Marston, C.K.,
Gulvik, C.A., Antonini, J.M., Negron, M.E., McQuiston, J.R., Hendricks, K., Weiner, Z., Balsamo,
G.A., Sokol, T., Byers, P., Kathryn, T., Zaheer, S., Long, S., O'Sullivan, B., de Perio, M.A.,
Hoffmaster, A.R., Salzer, J.S. and Bower, W.A. (2021) 'Fatal Anthrax Pneumonia in Welders and
Other Metalworkers Caused by *Bacillus cereus* Group Bacteria Containing Anthrax Toxin Genes
— U.S. Gulf Coast States, 1994–2020', *Morbidity and Mortality Weekly Report*, 70(41), pp. 1453–
1454.

Dupke, S., Barduhn, A., Franz, T., Leendertz, F.H., Couacy-Hymann, E., Grunow, R. and Klee,
S.R. (2019) 'Analysis of a newly discovered antigen of *Bacillus cereus* biovar *anthracis* for its
suitability in specific serological antibody testing', *Journal of Applied Microbiology*, 126(1), pp.
311–323.

Gaudet, R.G., Sintsova, A., Buckwalter, C.M., Leung, N., Cochrane, A., Li, J., Cox, A.D., Moffat,
J. and Gray-Owen, S.D. (2015) 'Cytosolic detection of the bacterial metabolite HBP activates TIFA-
dependent innate immunity', *Science*, 348(6240), pp. 1251–1255. doi:10.1126/science.aaa4921.

Guarner, J., Jernigan, J.A., Shieh, W.J., Tatti, K., Flannagan, L.M., Stephens, D.S., Popovic, T.,
Ashford, D.A., Perkins, B.A. and Zaki, S.R. (2003) 'Pathology and pathogenesis of bioterrorism-
related inhalational anthrax', *American Journal of Pathology*, 163(2), pp. 701–709.

Haydar, A., Tran, S.L., Guillemet, E., Darrigo, C., Perchat, S., Lereclus, D., Coquet, L., Jouenne,
T. and Ramarao, N. (2018) 'InhA1-mediated cleavage of the metalloprotease NprA allows *Bacillus*
cereus to escape from macrophages', *Frontiers in Microbiology*, 9(1063).

Hendrix, R.W., Casjens, S.R. and Lavigne, R. (2012) 'Family - Siphoviridae.', *Virus Taxonomy,*
Ninth Report of the International Committee on Taxonomy of Viruses.

Hoffmaster, A.R., Hill, K.K., Gee, J.E., Marston, C.K., De, B.K., Popovic, T., Sue, D., Wilkins, P.P.,

Avashia, S.B., Drumgoole, R., Helma, C.H., Ticknor, L.O., Okinaka, R.T. and Jackson, P.J. (2006)
'Characterization of *Bacillus cereus* isolates associated with fatal pneumonias: Strains are closely
related to *Bacillus anthracis* and Harbor *B. anthracis* virulence genes', *Journal of Clinical*
Microbiology, 44(9), pp. 3352–3360.

Hoffmaster, A.R., Ravel, J., Rasko, D.A., Chapman, G.D., Chute, M.D., Marston, C.K., De, B.K.,
Sacchi, C.T., Fitzgerald, C., Mayer, L.W., Maiden, M.C.J., Priest, F.G., Barker, M., Jiang, L., Cer,
R.Z., Rilstone, J., Peterson, S.N., Weyant, R.S., Galloway, D.R., Read, T.D., Popovic, T. and
Fraser, C.M. (2004) 'Identification of anthrax toxin genes in a *Bacillus cereus* associated with an
illness resembling inhalation anthrax', *PNAS*, 101(22), pp. 8449–8454.

Jensen, G.B., Hansen, B.M., Eilenberg, J. and Mahillon, J. (2003) 'The hidden lifestyles of *Bacillus*
cereus and relatives', *Environmental Microbiology*, 5(8), pp. 631–640.

Johnson, S.L., Daligault, H.E., Davenport, K.W., Jaissle, J., Frey, K.G., Ladner, J.T., Broomall,
S.M., Bishop-lilly, K.A., Bruce, D.C., Gibbons, H.S., Coyne, S.R., Lo, C., Meincke, L., Munk, A.C.,
Koroleva, G.I., Rosenzweig, C.N., Palacios, G.F., Redden, C.L., Minogue, T.D., Chain, P.S., Wt,
S., Mj, H., Taylor, L., Bd, C., Gm, K., Ta, S., C-s, C., Dh, A., Marks, P., Aa, K., Drake, J., Heiner,
C., Clum, A., Copeland, A., Huddleston, J., Ee, E., Sw, T. and Korlach, J. (2015) 'Complete
Genome Sequences for 35 Biothreat Assay-Relevant *Bacillus* Species', *Genome Announcements*,
3(2), pp. 10–11.

Kanehisa, M. and Sato, Y. (2019) 'KEGG Mapper for inferring cellular functions from protein
sequences', *Protein Science*, 29(1), pp. 28–35.

Kim, U. and Goepfert, J.M. (1971) 'Enumeration and Identification of *Bacillus cereus* in Foods',
Applied Microbiology, 22(4), pp. 581–587.

Koch, R. (1876) 'The etiology of anthrax , based on the life history of *Bacillus anthracis*', *Beitrage*
zur Biologie der Pflanzen, 2(2), pp. 277–310.

Lane DL. (1991) 16S/23S rRNA sequencing. *Nucleic Acid Techniques in Bacterial Systematics*,
pp. 115–75.

Lindeque, P.M. and Turnbull, P.C. (1994) 'Ecology and epidemiology of anthrax in the Etosha
National Park, Namibia.', *The Onderstepoort journal of veterinary research*, 61(1), pp. 71–83.

Manoharan, S., Taylor-Joyce, G., Brooker, T., Hernández-Rodriguez, C.S., Hapeshi, A., Baldwin,
V., Baillie, L., Oyston, P. and Waterfield, N.R. (2022) From cereus to anthrax and back again: The
role of the PlcR regulator in the anthrax 'crossover' strain *Bacillus cereus* G9241. [Unpublished
Manuscript], University of Warwick.

Margulis, L., Jorgensen, J.Z., Dolan, S., Kolchinsky, R., Rainey, F.A. and Lo, S.C. (1998) 'The
Arthromitus stage of *Bacillus cereus*: Intestinal symbionts of animals', *PNAS,* 95(3), pp. 1236–
1241.

Marston, C.K., Ibrahim, H., Lee, P., Churchwell, G., Gumke, M., Stanek, D., Gee, J.E., Boyer, A.E.,
Gallegos-Candela, M., Barr, J.R., Li, H., Boulay, D., Cronin, L., Quinn, C.P. and Hoffmaster, A.R.
(2016) 'Anthrax toxin-expressing *Bacillus cereus* isolated from an anthrax-like eschar', *PLoS ONE*,
11(6), pp. 1–7.

McKenzie, A.T., Pomerantsev, A.P., Sastalla, I., Martens, C., Ricklefs, S.M., Virtaneva, K., Anzick,
S., Porcella, S.F. and Leppla, S.H. (2014) 'Transcriptome analysis identifies *Bacillus anthracis*
genes that respond to CO_2 through an AtxA-dependent mechanism', *BMC Genomics*, 15(1).
886

887 Oh, S.-Y., Budzik, J.M., Garufi, G. and Schneewind, O. (2011) 'Two capsular polysaccharides
888 enable *Bacillus cereus* G9241 to cause anthrax-like disease.', *Molecular Microbiology*, 80(2), pp.
889 455–70.
890

891 Olive, A.J. and Sassetti, C.M. (2016) 'Metabolic crosstalk between host and pathogen: Sensing,
892 adapting and competing', *Nature Reviews Microbiology*, 14(4), pp. 221–234.
893

894 Panda, G., Basak, T., Tanwer, P., Sengupta, S., dos Santos, V.A.P.M. and Bhatnagar, R. (2014)
895 'Delineating the effect of host environmental signals on a fully virulent strain of *Bacillus anthracis*
896 using an integrated transcriptomics and proteomics approach', *Journal of Proteomics*, 105, pp.
897 242–265.
898

899 Pena-Gonzalez, A., Marston, C.K., Rodriguez-R, L.M., Kolton, C.B., Garcia-Diaz, J., Theppote, A.,
900 Frace, M., Konstantinidis, K.T. and Hoffmaster, A.R. (2017) 'Draft genome sequence of *Bacillus*
901 *cereus* LA2007, a human-pathogenic isolate harboring anthrax-like plasmids', *Genome*
902 *Announcements*, 5(16).
903

904 Peng, Q. and Yuan, Y. (2018) 'Characterization of a novel phage infecting the pathogenic
905 multidrug-resistant *Bacillus cereus* and functional analysis of its endolysin', *Applied Microbiology*
906 *and Biotechnology*, 102(18), pp. 7901–7912.
907

908 Rasko, D.A., Altherr, M.R., Han, C.S. and Ravel, J. (2005) 'Genomics of the *Bacillus cereus* group
909 of organisms', *FEMS Microbiology Reviews*, 29(2), pp. 303–329.
910

911 Rasko, D.A., Rosovitz, M.J., Økstad, O.A., Fouts, D.E., Jiang, L., Cer, R.Z., Kolstø, A.B., Gill, S.R.
912 and Ravel, J. (2007) 'Complete sequence analysis of novel plasmids from emetic and periodontal
913 *Bacillus cereus* isolates reveals a common evolutionary history among the *B. cereus*-group
914 plasmids, including *Bacillus anthracis* pXO1', *Journal of Bacteriology*, 189(1), pp. 52–64.
915

916 Raynor, M.J., Roh, J.H., Widen, S.G., Wood, T.G. and Koehler, T.M. (2018) 'Regulons and
917 protein–protein interactions of PRD-containing *Bacillus anthracis* virulence regulators reveal
918 overlapping but distinct functions', *Molecular Microbiology*, 109(1), pp. 1–22.
919

920 Regan, G., Itaya, M. and Piggot, P.J. (2012) 'Coupling of σG Activation to Completion of
921 Engulfment during Sporulation of *Bacillus subtilis* Survives Large Perturbations to DNA
922 Translocation and Replication', *Journal of Bacteriology*, 194(22), pp. 6264–6271.
923

924 Ross, K.F. and Billing, E. (1957) 'The water and solid content of living bacterial spores and
925 vegetative cells as indicated by refractive index measurements', *J Gen Microbiol*, 16(2), pp. 418–
926 25.
927

928 Scarff, J.M., Seldina, Y.I., Vergis, J.M., Ventura, C.L. and O'Brien, A.D. (2018) 'Expression and
929 contribution to virulence of each polysaccharide capsule of *Bacillus cereus* strain G9241', *PLoS*
930 *ONE*, 13(8).
931

Silver-Mysliwiec, T.H. and Bramucci, M.G. (1990) 'Bacteriophage-Enhanced Sporulation:
Comparison of Spore-Converting Bacteriophages PMB12 and SP10', *Journal of Bacteriology*,
172(4), pp. 1948–1953.

Stein, C., Jones, G.W., Chalmers, T. and Berry, C. (2006) 'Transcriptional analysis of the toxin-
coding plasmid pBtoxis from *Bacillus thuringiensis* subsp. israelensis', *Applied and Environmental*
Microbiology, 72(3), pp. 1771–1776.

Strauch, M.A., Trach, K.A., Day, J. and Hoch, J.A. (1992) 'Spo0A activates and represses its own
synthesis by binding at its dual promoters', *Biochimie*, 74(7–8), pp. 619–626.

Swiecicka, I. and Mahillon, J. (2006) 'Diversity of commensal *Bacillus cereus sensu lato* isolated
from the common sow bug (*Porcellio scabe*r, Isopoda)', *FEMS Microbiology Ecology*, 56(1), pp.
132–140.

Szklarczyk, D., Gable, A.L., Lyon, D., Junge, A., Wyder, S., Huerta-Cepas, J., Simonovic, M.,
Doncheva, N.T., Morris, J.H., Bork, P., Jensen, L.J. and Von Mering, C. (2019) 'STRING v11:
Protein-protein association networks with increased coverage, supporting functional discovery in
genome-wide experimental datasets', *Nucleic Acids Research*, 47(D1), pp. D607–D613.

Tan, I.S. and Ramamurthi, K.S. (2014) 'Spore formation in *Bacillus subtilis*', *Environmental*
Microbiology Reports, 6(3), pp. 212–225.

Terwilliger, A., Swick, M.C., Pflughoeft, K.J., Pomerantsev, A., Lyons, C.R., Koehler, T.M. and
Maresso, A. (2015) '*Bacillus anthracis* overcomes an amino acid auxotrophy by cleaving host
serum proteins', *Journal of Bacteriology*, 197(14), pp. 2400–2411.

Uchida, I., Makino, S.I., Sekizaki, T. and Terakado, N. (1997) 'Cross-talk to the genes for *Bacillus*
anthracis capsule synthesis by *atxA*, the gene encoding the trans-activator of anthrax toxin
synthesis', *Molecular Microbiology*, 23(6), pp. 1229–1240.

Wang, Y., Oh, S., Hendrickx, A.P.A., Lunderberg, J.M. and Schneewind, O. (2013) '*Bacillus cereus*
G9241 S-Layer Assembly Contributes to the Pathogenesis of Anthrax-Like Disease in Mice',
Journal of Bacteriology, 195(3), pp. 596–605.

Wilson, M.K., Vergis, J.M., Alem, F., Palmer, J.R., Keane-Myers, A.M., Brahmbhatt, T.N., Ventura,
C.L. and O'Brien, A.D. (2011) '*Bacillus cereus* G9241 makes anthrax toxin and capsule like highly
virulent B. anthracis Ames but behaves like attenuated toxigenic nonencapsulated B. anthracis
Sterne in rabbits and mice', *Infection and Immunity*, 79(8), pp. 3012–3019.

Wright, A.M., Beres, S.B., Consamus, E.N., Long, S.W., Flores, A.R., Barrios, R. and Richter, G.S.
(2011) 'Rapidly Progressive, Fatal, Inhalation Anthrax-like Infection in a Human: Case Report,
Pathogen Genome Sequencing, Pathology, and Coordinated Response', *Archives of Pathology &*
Laboratory Medicine, pp. 1447–1459.

Wynosky-Dolfi, M.A., Snyder, A.G., Philip, N.H., Doonan, P.J., Poffenberger, M.C., Avizonis, D.,
Zwack, E.E., Riblett, A.M., Hu, B., Strowig, T., Flavell, R.A., Jones, R.G., Freedman, B.D. and
Brodsky, I.E. (2014) 'Oxidative metabolism enables *Salmonella* evasion of the NLRP3
inflammasome', *Journal of Experimental Medicine*, 211(4), pp. 653–668.

Zhou, Y., Liang, Y., Lynch, K.H., Dennis, J.J. and Wishart, D.S. (2011) 'PHAST: A Fast Phage
Search Tool', *Nucleic Acids Research*, 39, pp. W347–W352.
984