1 **Metformin Improves Ovarian Function and Increases Egg Production in**
2 **Broiler Breeder Hens**
3

4 **Authors:**   Evelyn A. Weaver
5   Center for Reproductive Biology and Health
6   Department of Animal Science
7   The Pennsylvania State University
8   320 AVBS Building,
9   University Park, PA 16802
10
11   Ramesh Ramachandran
12   Center for Reproductive Biology and Health
13   Department of Animal Science
14   The Pennsylvania State University
15   306 AVBS Bldg.,
16   University Park, PA 16802
17
18
19 **Affiliation:**   Center for Reproductive Biology and Health
20   Department of Animal Science
21   The Pennsylvania State University
22   University Park, PA
23

Short title: Metformin and Ovarian Function

In Brief: The pathophysiology of the ovarian dysfunction encountered in broiler breeder hens

remains poorly understood but is similar to a condition in women known as PCOS. This study

reveals that metformin may provide a cheap and effective method of improving ovarian function

in broiler breeder hens.

Corresponding   Dr. Ramesh Ramachandran
author: Center for Reproductive Biology and Health
31   Department of Animal Science
32   The Pennsylvania State University
33   306 AVBS Bldg.,
34   University Park, PA 16802
35   Phone (814) 865-5202
36   Fax (814) 865-5691
37   Email: RameshR@psu.edu
38
39

KEYWORDS: ovary, body weight, adipose tissue, progesterone, testosterone, androstenedione, estrogen

Number of words: 4,195
Number of Figures/Tables: 10
Number of References: 51

85   **ABSTRACT**
86

Broiler breeder hens, the parent stock of commercial broiler chickens, have poor

reproductive efficiency associated with aberrant and excessive recruitment of ovarian follicles

which results in sub-optimal egg production, fertility, and hatchability. The reproductive

dysfunction observed in these hens resembles polycystic ovary syndrome in women, a condition

wherein metformin is prescribed as a treatment. The main objectives of this study were to

determine the effect of metformin on body weight, abdominal fat pad weight, ovarian function,

and plasma steroid hormone concentrations. Broiler breeder hens were treated with 0, 25, 50, or

75 mg/kg body weight of metformin mixed in the diet for 40 weeks (n= 45 hens/treatment; 25-65

weeks of age). At 65 weeks of age, hens that received the highest dose of metformin had

significantly lesser body and abdominal fat pad weights ($p < 0.05$) than the control. Metformin

treatment, at all levels, normalized the preovulatory and prehierarchical ovarian follicular

hierarchy and significantly increased the total number of eggs laid per hen during the entire

production period in hens that received 50 or 75 mg/kg body weight compared to the control.

Fertility and hatchability of eggs were not affected by metformin treatment. Metformin

treatment at all levels altered the plasma profile of reproductive hormones, with significantly

lower plasma testosterone concentrations and a decreased testosterone to androstenedione ratio

in hens that received metformin ($p < 0.05$). Future studies should focus on the mechanisms

underlying the beneficial effects of metformin in improving the reproductive efficiency of broiler

breeder hens.

INTRODUCTION

Broiler breeder hens, the parent stock of commercial broiler chickens, have poor reproductive efficiency due to the selection pressure for superior growth-related traits in broiler progeny. Decades of intensive genetic selection for growth-related traits negatively associated with reproductive function have led to severe ovarian dysfunction in the broiler breeder hen (De Jong and Guémené, 2011; Zuidhof et al., 2014). Full-fed broiler breeder hens accumulate excessive amounts of visceral adipose tissue and often have heavier ovaries, excessive follicular recruitment, double ovulation, or increased incidence of ovarian regression (Yu et al., 1992). Consequently, broiler breeder hens have decreased egg production, sub-optimal fertility and hatchability, and decreased viability of embryos (Yu et al., 1992).

The pathophysiology underlying the ovarian dysfunction in broiler breeder hens is still unknown, but previous studies have shown that it may be related to obesity-induced metabolic dysfunction and lipotoxicity (Cassy et al., 2004; Chen et al., 2006; Pan et al., 2012; Walzem and Chen, 2014) . In studies where broiler breeder hens were allowed to consume feed *ad libitum*, significantly higher body weights, increased abdominal fat pad weights, and increased liver weights were observed (Chen et al., 2006; Pan et al., 2012; Taherkhani et al., 2010). Such increases in the accumulation of abdominal adipose tissue and increased weight gain were associated with increased plasma glucose, insulin, non-esterified fatty acids, triacylglycerol, cholesterol, phospholipid, and ceramide levels (Chen et al., 2006; Pan et al., 2012; Taherkhani et al., 2010). An increased incidence of ovarian dysfunction, follicle atresia, early ovarian regression, and altered plasma progesterone and estradiol levels was also observed in these hens (Cassy et al., 2004; Chen et al., 2006; Pan et al., 2012; Taherkhani et al., 2010). In a study where broiler breeder hens were treated with testosterone, higher rates of internal ovulation and

increased disruption of the reproductive cycle was observed (Navara, Kristen J et al., 2015).

These findings suggest that increased adiposity and disrupted steroid hormone signaling may

play a central role in developing ovarian dysfunction. Although previous studies have focused on

the effects of *ad libitum* feeding, the hyper-recruitment of prehierarchical ovarian follicles and a

deranged preovulatory follicular hierarchy are frequently encountered even when the hens are

fed a restricted amount of feed.

Interestingly, the abnormal ovarian changes and disturbed plasma profiles of reproductive

hormones observed in broiler breeder hens appear similar to a condition in women known as

polycystic ovary syndrome (PCOS). PCOS is most commonly associated with obesity, metabolic

syndrome, irregular steroid hormone production, irregular ovulation, and poor fertility, all of

which have been previously observed in broiler breeder hens (Pasquali et al., 2003; Walzem and

Chen, 2014). Metformin, a biguanide compound, is the most commonly prescribed drug to treat

polycystic ovary syndrome. In women with PCOS, metformin has been shown to reduce insulin

resistance and correct metabolic disturbances commonly observed in PCOS. Metformin has also

been associated with increased ovulation, fertilization, and pregnancy rates (Fleming et al., 2002;

Morley et al., 2017; Nestler et al., 2002; Tang et al., 2006b; Tso et al., 2020; Velazquez et al.,

1997), normalization of the endocrine profile, as well as a return to normal menstrual cyclicity

(Moghetti et al., 2000; Morin-Papunen et al., 1998; van Santbrink et al., 2005; Velazquez et al.,

1994; Xing et al., 2020). While this improvement in reproductive function has previously been

associated with the correction of insulin resistance (Jakubowicz et al., 2001; Perriello et al.,

1994; Prager et al., 1986) and improved energy metabolism, other studies have shown that

metformin can exert direct effects on the ovary (Bertoldo et al., 2014; Ehrmann et al., 1997;

Vrbikova et al., 2001). In agreement with previous studies, we found that metformin attenuates

steroidogenesis in the granulosa cells isolated from broiler breeder hen ovarian follicles (Weaver

and Ramachandran, 2020).

Based on the preceding, we hypothesized that dietary metformin supplementation would

improve reproductive efficiency in the broiler breeder hen. The main objectives of the present

study are to determine the effects of metformin on (i) body weight and abdominal fat pad weight,

(ii) ovarian prehierarchical and preovulatory follicular hierarchy, (iii) egg production, fertility,

and hatchability, and (iv) plasma concentrations of steroid hormones.

MATERIALS AND METHODS

166
Animal Husbandry and Metformin Treatment

The Pennsylvania State University's Institutional Animal Care and Use Committee

approved all animal procedures described herein (protocol number PRAMS200746656). The

overall experimental design and sample collection timeline are provided in Figure 1. Broiler

breeder female (Cobb 500; Cobb-Vantress Inc.) and male (Hubbard M99) chickens were

maintained from 1 day-old until the completion of the study at the Poultry Education and

Research Center at The Pennsylvania State University (University Park, PA). Broiler breeder

hens and roosters were housed individually in battery cages from 21 weeks of age and were feed-

restricted according to the Cobb Breeder and Hubbard management guide. Water was provided

ad libitum. The chickens were photo stimulated with a 16h light:8h dark (4:00 to 20:00 hours)

cycle for the study duration. At 25 weeks of age, the broiler breeder female chickens were

randomly allocated to four experimental groups (n=45) to receive four metformin levels (0, 25,

50, or 75 mg/kg body weight; Midwest Veterinary Supply, Lakeville, MN) supplemented in the

diet. The metformin treatment continued until the end of the study, at which the chickens were

65 weeks of age. Male chickens received no metformin supplementation in their diets. The

broiler breeder hens were on a time-restricted feeding schedule, where a fixed amount of feed

was given each morning as recommended by Cobb-Vantress Inc. This fixed amount of feed and

once-a-day feeding schedule ensured that each hen consumed the correct dose of metformin

daily. A subset of broiler breeder hens from each treatment group (n=10) was weighed every ten

weeks to adjust the amount of metformin mixed into the feed according to their change in

weights over time. Egg production was recorded daily for each broiler breeder hen starting at 22

weeks of age through the end of the study.

Body weight

A random subset of hens (n=10/treatment group) were weighed every ten weeks starting

at 30 weeks of age for the duration of the study to determine the effect of metformin on body

weight. The body weights were also used to adjust the amount of metformin added to the feed to

maintain the doses at 0, 25, 50, or 75 mg/kg throughout the study. A subset of hens (n=12

hens/treatment group) were weighed at the end of the study (65 weeks of age). At 65 weeks,

broiler breeder hens were euthanized and the abdominal fat pad was collected and weighed (n=

12 hens/treatment group).

Fertility and hatchability

For fertility and hatchability studies, a subset of female chickens from each experimental

group was randomly selected (n=20/treatment group). The same chickens were utilized in all

fertility and hatchability data collections. For artificial insemination (AI), semen was collected

and pooled from 15 broiler breeder male chickens. Immediately following collection, 50 mL of

undiluted, pooled semen was deposited into the oviduct of each broiler breeder female chicken.

Inseminations were done every other day, and eggs were collected daily and stored at 55°F until

the end of a 10-day collection period (n= 100-150 eggs/treatment group). Fertility and

hatchability data collections were conducted at ten different time points throughout the study (22,

25, 30, 35, 40, 45, 50, 55, 60, and 65 weeks of age). Eggs were placed in Chick Master (Medina,

OH) incubators set at 100°F and 82% humidity. Eggs were candled on day 10 of incubation to

determine fertility for each treatment group, represented as the total number of fertile eggs to the

total number of eggs set for each treatment group. On day 19 of incubation, fertile eggs were

transferred to Chick Master hatchers set at 98.2°F with humidity ~65-70%. Hatchability was

determined on day 21 and is represented as the total number of chicks hatched to the total

number of fertile eggs transferred to the hatchers for each treatment group.

Plasma Steroid Hormone Quantification

Blood samples were collected from the wing vein from a subset of fasted hens (n=6/

treatment group) every five weeks to quantify plasma steroid hormone concentrations. Blood

samples were immediately placed on ice and centrifuged within 1 h at 2300 x g for 15 min at

4°C. Plasma was collected and stored at -80°C for quantification of progesterone, total estrogens,

testosterone, and androstenedione concentrations. Plasma progesterone concentration was

determined by a commercial human double-antibody radioimmunoassay (RIA) kit (cat#

0717010, MP Biomedicals, LLC, Irvine, CA) according to the manufacturer's recommendations.

The intraassay coefficient of variation (CV) was 8%, and an interassay CV of 9% with a

sensitivity of 0.2-50 ng/mL for this assay. For the analysis of plasma total estrogens,

testosterone, and androstenedione, an extraction step was necessary prior to running the assay

according to the manufacturer's recommendations. Briefly, 6 mL of ethyl acetate: hexane, mixed

in a ratio of 3:2, was added to 500 mL of plasma. Tubes were vortexed vigorously for 60

seconds, and phases were allowed to separate. Next, 5 ml of the organic phase (top phase) was

withdrawn and evaporated under nitrogen gas. The sample residue was reconstituted in 2.5 mL

of steroid diluent provided by the manufacturer and incubated at room temperature for one hour

while gently mixed on a plate shaker. Extracted samples were stored at -20°C. According to the

manufacturer's recommendations, the total estrogens plasma concentration was determined by a

commercial human double-antibody RIA kit (cat# 07140205, MP Biomedicals). For the total

estrogens assay, the intraassay and interassay CV were 8% and 7%, respectively, while the

sensitivity of this assay was 2.5-100 pg/mL. Plasma testosterone and androstenedione

concentrations were determined by RIA using commercial reagents (cat# 0718910 and cat#

07109202, MP Biomedicals, LLC, Irvine, CA). For testosterone and androstenedione, intraassay

CVs were 8%, and the interassay CVs were 3% and 12%, respectively. The sensitivity of the

testosterone and androstenedione assays were 0.025-10 ng/mL and 0.1-10 ng/mL, respectively.

Statistical Analyses

All data were subjected to a one-way analysis of variance (ANOVA) using GraphPad

Prism version 9.1.0, GraphPad Software, San Diego, CA. Differences among individual means

were determined by using Tukey's HSD multiple comparisons test. To analyze cumulative egg

production per hen between the treatment groups and over time, data were subjected to a two-

way ANOVA using GraphPad Prism version 9.1.0, (GraphPad Software, San Diego, CA).

Differences among individual means were determined by using Tukey's HSD multiple

comparisons test. A chi-square test was performed to determine the statistical significance of the

number of hens still laying at the end of the trial (metformin treatment vs. control). A probability

level of P < 0.05 is considered statistically significant. All data are represented as mean ±

standard error of the mean. All data points are compared within a timepoint and not across

timepoints, except for the radioimmunoassay data, which was also analyzed across time for each

treatment group.

RESULTS

Metformin decreases the body weight and accretion of abdominal adipose tissue

Body weight of all hens at 20- or 30 weeks of age did not differ. Hens receiving 75 mg

metformin/kg body weight had significantly lesser body weights at 40 weeks of age when

compared with hens receiving 25 mg metformin/kg body weight. At 50 and 60 weeks of age,

hens receiving the highest dose of metformin had significantly lesser body weights when

compared to all other treatment groups. However, hens receiving the highest dose only differed

from those receiving 0- or 25 mg/kg metformin/kg body weight at 65 weeks of age (Fig. 2A).

Consistent with the observed decrease in body weight, hens receiving the highest dose of

metformin had significantly lesser abdominal fat pad weights at the end of the study (65 weeks

of age) compared to the control group and the group receiving the lowest dose of metformin (25

mg/kg of body weight) (Fig. 2B).

Metformin affects the ovarian follicular hierarchy and increases overall egg production

Figures 3A and 3B show representative hierarchy of preovulatory and prehierarchical

ovarian follicles from hens (40 weeks of age) that received 0 or 25 mg/kg body weight of

metformin, respectively. As seen in Fig. 3A, there are an excessive number of preovulatory and

prehierarchical follicles in the hens that received no metformin. In contrast, supplementation of

metformin in the diet restores the ovarian follicular hierarchy towards a normal phenotype (Fig.

3B). In addition to the changes in the follicular hierarchy, hens which received 50 or 75 mg/kg

metformin laid a significantly greater number of eggs per hen on average when compared to the

control, starting at 50 weeks of age through the end of the study (Fig 4A). When cumulative egg

production per hen was observed over the entire production period (22-65 weeks of age), hens

that received 50 or 75 mg metformin /kg body weight laid a significantly greater number of eggs

per hen when compared to the control (Fig. 4B), which is reflected in a greater total number of

eggs laid per treatment group over the entire study (Fig. 4B insert). Furthermore, hens that

received the highest dose of metformin had a significantly greater number of hens remaining in

production at the end of the study when compared to the control (Table 1). The supplementation

of metformin into the diet of broiler breeder hens had no effect on the fertility or hatchability of

eggs among all treatment groups (Supplementary Table 1).

Metformin improves the plasma endocrine profile of reproductive hormones

The supplementation of metformin in the diet significantly affected plasma progesterone

concentrations when observed over time (Fig. 5A and 5B). From 30 weeks until 50 weeks of age,

hens that received metformin in the diet tended to have lesser plasma progesterone

concentrations when compared to the control. No difference was observed among treatment

groups at 25 and 55 weeks of age. Interestingly, at 60 and 65 weeks of age, we observed a

significant increase in plasma progesterone concentrations in hens that received metformin in the

diet compared to the control, which is consistent with the prolonged egg production observed in

these groups (Fig. 5A). When plasma progesterone concentrations were observed across time for

each treatment group, we noted a significant decline in plasma progesterone concentrations in

hens that received no metformin in the diet beginning at 50 weeks of age. However, plasma

progesterone concentrations remained elevated and more consistent in hens supplemented with

metformin in the diet at all levels (Fig. 5B). Like plasma progesterone concentrations, hens that

received metformin in the diet had significantly lesser plasma total estrogen concentrations at 35,

40, 45, 50, and 60 weeks of age when compared to the control (Fig. 6A). When plasma total

estrogen concentrations were studied over time, a similar trend was observed in which hens that

received 50- or 75 mg/kg had a more consistent total estrogen profile over time when compared

to the hens which received 0- or 25 mg/kg in the diet (Fig. 6B). Beginning at 30 weeks of age

through 65 weeks of age, hens that received metformin in the diet had significantly lesser plasma

testosterone concentrations when compared to the control (Fig. 7A). When plasma testosterone

concentrations were studied over time, from 25 to 65 weeks of age, we observed consistent

plasma testosterone concentration patterns within each treatment group (Fig. 7B). There was no

effect of metformin on plasma androstenedione concentrations across all time points and over

time (Fig. 8A and 8B). However, when we analyzed the testosterone to androstenedione ratio, a

novel biomarker used to assess the metabolic phenotype in women with PCOS (Dumesic, Daniel

A et al., 2021; Misichronis, G et al., 2011; Munzker, J et al., 2017), we observed that hens that

received 50 or 75 mg/kg showed significantly lesser testosterone to androstenedione ratios at 40,

45, 50, 60 and 65 weeks of age when compared to the control (Fig. 9A). When the testosterone to

androstenedione ratio is observed across time, hens that received metformin in the diet had a

lower ratio that remained consistent compared to the control (Fig. 9B).

DISCUSSION

We report for the first time that dietary metformin supplementation enhances ovarian

function in the broiler breeder hen through the normalization of the ovarian follicular hierarchy

and endocrine profile of reproductive hormones, with no effect on fertility or hatchability. The

improved ovarian function was associated with decreased body weights, decreased accretion of

abdominal adipose tissue, increased and prolonged egg production, and altered plasma steroid

hormone concentrations.

Our findings are consistent with previous studies in which metformin was associated with

a decrease in overall body weight of human subjects (Bray et al., 2012; Chen et al., 2021;

Fleming et al., 2002; Guan et al., 2020; Tokubuchi et al., 2017), rodent (Coll et al., 2020; Jing et

al., 2018; Rencbar et al., 2018) and Leghorn chicken (Chen et al., 2011). It has been previously

reported that 40-80% of women diagnosed with PCOS are overweight or obese (Sam, 2007). In

women with this condition, a decrease in body weight has been positively correlated with

improved reproductive function, although some studies have suggested that weight loss may be

the key to improving reproductive function rather than metformin itself (Tang et al., 2006a). Our

findings also showed a significant decrease in abdominal adipose tissue weight in hens that

received the highest dose of metformin. Consistent with our findings, several studies mentioned

above also showed that metformin leads to a decrease in the hip to waist circumference in

women, which is consistent with a decrease in the accretion of visceral adipose tissue.

We chose to begin metformin supplementation at 25 weeks of age as most hens have

begun to lay eggs by this time. Interestingly, we observed a sharp decline in egg production in

hens that received the highest dose of metformin around 27 weeks through 30 weeks of age (EA

Weaver, R Ramachandran, unpublished data), which was not observed in a previous study

conducted in our laboratory in which we supplemented metformin in the diet from 42-60 weeks

of age (EA Weaver, R Ramachandran, unpublished data). Previous studies have shown that a

certain percentage of body fat is required for hens to begin laying and maintain egg production

(Bornstein et al., 1983; Hurwitz and Plavnik, 1989; Yuan et al., 1994), so this observed decline

in egg production could be due to the effect of metformin on body fat percentage in the hens

which received the highest dose. Although a significant decline in egg production was observed

early on, these hens recovered egg production, which was maintained for the duration of the

study. Further studies are required to optimize the ideal body composition and age to begin

supplementing metformin into the diet of broiler breeder hens.

This is the first to report that dietary metformin supplementation corrects the excessive

recruitment of prehierarchical and preovulatory ovarian follicles, thus normalizing the ovarian

follicular hierarchy. This is still observed even when supplemented at the lowest dose of 25

mg/kg of body weight. For a hen to produce a clutch of eggs, it must maintain a small cohort of

prehierarchical follicles from which one follicle is recruited daily (~ every 26 hours) into the

preovulatory pool of follicles. The ovarian dysfunction encountered in broiler breeder hens has

previously been associated with excessive follicular recruitment, leading to a double

preovulatory follicular hierarchy and increased incidence of double ovulation, which are often

non-viable. Consistent with our findings, previous studies have reported a 'normalization' of the

ovarian follicular profile in both rats (Ali et al., 2020; Di Pietro et al., 2015; Rencbar et al., 2018)

and humans (Fleming et al., 2002; Morley et al., 2017) and has been reported to increase

ovulation rates (Fleming et al., 2002; Nestler et al., 2002, 1998) as well as a return to normal

menstrual cyclicity (Moghetti et al., 2000; Morin-Papunen et al., 1998; van Santbrink et al.,

2005; Velazquez et al., 1994) in women with PCOS.

365   Along with the 'normalization' of the ovarian follicular hierarchy, dietary metformin

366  supplementation was also associated with increased and prolonged egg production, with the

367  highest dose having the most significant effect. Even with strict feed restrictions, broiler breeder

368  hens often have a significant reduction in egg production beginning around 50 weeks of age, and

369  many cease laying by 60 weeks of age. Interestingly, during the last 10 weeks of the study (55-

370  65 weeks of age), we observed maintenance of egg production in hens that received metformin in

371  their diet, even at the lowest dose of 25 mg/kg of body weight, with the two highest doses having

372  the most significant effect. Overall, hens that received metformin laid a greater number of eggs

373  on average which can be attributed to this maintenance of egg production. Based on previous

374  studies, this maintenance of egg production could potentially be attributed to a reduction in body

375  weight, reduction in the accumulation of abdominal adipose tissue, an effect of metformin on

376  steroid hormone production or the direct effect of metformin on the ovary. However, further

377  studies are necessary to determine the mechanisms by which metformin improves egg production

378  in the broiler breeder hen. We observed that metformin had no effect on fertility or hatchability

379  amongst all treatment groups; however, previous studies have shown that metformin can increase

380  fertility and pregnancy rates (Morley et al., 2017; Nestler et al., 2002; Tang et al., 2006b; Tso et

381  al., 2020; Velazquez et al., 1997) in women with PCOS.

382   Finally, we observed that supplementation of metformin in the diet significantly affected

383  the plasma endocrine profile of reproductive hormones in broiler breeder hens. It has been

384  previously reported that metformin can normalize the endocrine profile in both women

385  (Velazquez et al., 1994; Xing et al., 2020) and rats (Zhang et al., 2017). In a previous study

386  conducted in our laboratory, we reported that metformin attenuates steroidogenesis in the

387  granulosa cells isolated from prehierarchical follicles and preovulatory follicles (Weaver and

Ramachandran, 2020); however, this is the first report on the long-term *in vivo* effects of

metformin in broiler breeder chickens. In the hen, the largest preovulatory follicle produces a

large amount of progesterone, while prehierarchical follicles are major producers of estrogens.

We observed increased plasma progesterone and total estrogens concentrations in hens that

received no metformin early on in the study, which is consistent with the hyper-recruitment of

preovulatory and prehierarchical follicles. Interestingly, around 50 weeks of age, we observe a

decline in plasma progesterone concentration in hens that received no metformin in the diet. In

contrast, progesterone concentrations remained consistent and elevated in hens that received

metformin at all doses. As metformin supplementation maintained egg production during this

time, this increase in plasma progesterone is consistent with maintaining a preovulatory follicular

hierarchy. We observed a significant effect of metformin on plasma testosterone concentrations,

consistent with previous studies that reported decreased plasma testosterone in women who

received metformin (Guan et al., 2020; Kurzthaler et al., 2014; Nestler and Jakubowicz, 1996).

Increased testosterone levels have been previously reported in broiler breeder hens allowed to

consume feed *ad libitum* (Singh et al., 2013; Taherkhani et al., 2010); however, we are the first

to report increased plasma testosterone even when restricted-fed. Based on our observations and

similar to women with PCOS, it appears that an increased plasma testosterone profile is

associated with increased body weight and adverse reproductive outcomes in the broiler breeder

hen. In the present study, we found that plasma androstenedione concentrations in the broiler

breeder hen treated with metformin did not differ. However, when comparing plasma

testosterone to androstenedione ratios, we observed a significantly decreased testosterone to

androstenedione ratio beginning at 40 weeks of age through the end of the study in hens that

received metformin in the diet at 50 or 75 mg/kg of body weight. Previous studies have

suggested that the androstenedione to testosterone ratio may be utilized as a biomarker of

metabolic health in women with PCOS. An increased testosterone to androstenedione ratio has

been associated with a higher likelihood of developing metabolic dysfunction, obesity, and

insulin resistance in women with PCOS (Dumesic, Daniel A et al., 2021; Misichronis, G et al.,

2011; Munzker, J et al., 2017).

In summary, we provide evidence that dietary metformin supplementation improves

ovarian function in the broiler breeder hen. The improved ovarian function was associated with

decreased body weights and abdominal fat pad weights, the normalization of the ovarian

follicular hierarchy, increased and prolonged egg production, and an improved endocrine profile

of reproductive hormones. As the broiler breeder hen continues to be the poorest in reproductive

efficiency of all avian species, a cheap and effective method to improve egg production is crucial

to maintain broiler production and feed the growing population. Future studies are needed to

determine the mechanisms that underlie the beneficial effect of metformin in correcting the

ovarian follicular hierarchy and increasing egg production in the chicken.

DECLARATION OF INTEREST

There is no conflict of interest that could be perceived as prejudicing the impartiality of

the research reported.

FUNDING

This project was supported by Agriculture and Food Research Initiative Competitive

Grant no. 2017-67015-26506 from the USDA National Institute of Food and Agriculture and in

part, by NIH Grant T32GM108563.

AUTHOR CONTRIBUTION

EAW designed and conducted the trial and experiments and analyzed all data. EAW and

RR wrote the manuscript.

ACKNOWLEDGEMENTS

The authors wish to thank the staff at Poultry Education and Research Center at Penn

State for their assistance with the trial.

443
444

REFERENCES

Ali, E.M.T., Abdallah, H.I., El-Sayed, S.M., 2020. Histomorphological, VEGF and TGF-β immunoexpression changes in the diabetic rats' ovary and the potential amelioration following treatment with metformin and insulin. J. Mol. Histol. 51, 287–305. https://doi.org/10.1007/s10735-020-09880-x

Bertoldo, M.J., Faure, M., Dupont, J., Froment, P., 2014. Impact of metformin on reproductive tissues: an overview from gametogenesis to gestation. Ann. Transl. Med. 2, 55–55. https://doi.org/10.3978/j.issn.2305-5839.2014.06.04

Bornstein, S., Plavnik, I., Lev, Y., 1983. Body weight and/or fatness as potential determinants of the onset of egg production in broiler breeder hens. Br. Poult. Sci. 25, 323–341. https://doi.org/10.1080/00071668408454873

Bray, G., Edelstein, S., Crandall, J., Aroda, V., Franks, P., Fujimoto, W., Horton, E., Jeffries, S., Montez, M., Mudaliar, S., Pi-Sunyer, F., White, N., Knowler, W., 2012. Long-Term Safety, Tolerability, and Weight Loss Associated With Metformin in the Diabetes Prevention Program Outcomes Study. Diabetes Care 35, 731–737. https://doi.org/10.2337/dc11-1299

Cassy, S., Metayer, S., Crochet, S., Rideau, N., Collin, A., Tesseraud, S., 2004. Leptin receptor in the chicken ovary: potential involvement in ovarian dysfunction of ad libitum-fed broiler breeder hens. Reprod. Biol. Endocrinol. 2, 72–72. https://doi.org/10.1186/1477-7827-2-72

Chen, S.E., McMurtry, J.P., Walzem, R.L., 2006. Overfeeding-Induced Ovarian Dysfunction in Broiler Breeder Hens Is Associated with Lipotoxicity. Poult. Sci. 85, 70–81. https://doi.org/10.1093/ps/85.1.70

Chen, W.-L., Wei, H.-W., Chiu, W.-Z., Kang, C.-H., Lin, T.-H., Hung, C.-C., Chen, M.-C., Shieh, M.-S., Lee, C.-C., Lee, H.-M., 2011. Metformin regulates hepatic lipid metabolism through activating AMP-activated protein kinase and inducing ATGL in laying hens. Eur. J. Pharmacol. 671, 107–112. https://doi.org/10.1016/j.ejphar.2011.09.029

Chen, X., He, S., Wang, D., 2021. Effects of metformin on body weight in polycystic ovary syndrome patients: model-based meta-analysis. Expert Rev. Clin. Pharmacol. 14, 121–130. https://doi.org/10.1080/17512433.2021.1863788

Coll, A.P., Chen, M., Taskar, P., Rimmington, D., Patel, S., Tadross, J.A., Cimino, I., Yang, M.,
    Welsh, P., Virtue, S., Goldspink, D.A., Miedzybrodzka, E.L., Konopka, A.R., Esponda,
    R.R., Huang, J.T.-J., Tung, Y.C.L., Rodriguez-Cuenca, S., Tomaz, R.A., Harding, H.P.,
    Melvin, A., Yeo, G.S.H., Preiss, D., Vidal-Puig, A., Vallier, L., Nair, K.S., Wareham,
    N.J., Ron, D., Gribble, F.M., Reimann, F., Sattar, N., Savage, D.B., Allan, B.B.,
    O'Rahilly, S., 2020. GDF15 mediates the effects of metformin on body weight and
    energy balance. Nature 578, 444–448. https://doi.org/10.1038/s41586-019-1911-y

De Jong, I.C., Guémené, D., 2011. Major welfare issues in broiler breeders. Worlds Poult. Sci. J.
    67, 73–82. https://doi.org/10.1017/S0043933911000067

Di Pietro, M., Parborell, F., Irusta, G., Pascuali, N., Bas, D., Bianchi, M.S., Tesone, M.,
    Abramovich, D., 2015. Metformin regulates ovarian angiogenesis and follicular
    development in a female polycystic ovary syndrome rat model. Endocrinology 156,
    1453–1463. https://doi.org/10.1210/en.2014-1765

Dumesic, Daniel A, Tulberg, Ayli, McNamara, Megan, Grogan, Tristan R, Abbott, David H,
    Naik, Rajanigandha, Lu, Gwyneth, Chazenbalk, Gregorio D, 2021. Serum Testosterone
    to Androstenedione Ratio Predicts Metabolic Health in Normal-Weight Polycystic Ovary
    Syndrome Women. J. Endocr. Soc. 5. https://doi.org/10.1210/jendso/bvab158

Ehrmann, D.A., Cavaghan, M.K., Imperial, J., Sturis, J., Rosenfield, R.L., Polonsky, K.S., 1997.
    Effects of Metformin on Insulin Secretion, Insulin Action, and Ovarian Steroidogenesis
    in Women with Polycystic Ovary Syndrome. J. Clin. Endocrinol. Metab. 82, 524–530.
    https://doi.org/10.1210/jcem.82.2.3722

Fleming, R., Hopkinson, Z.E., Michael Wallace, A., Greer, I.A., Sattar, N., 2002. Ovarian
    function and metabolic factors in women with oligomenorrhea treated with metformin in
    a randomized double blind placebo-controlled trial. J. Clin. Endocrinol. Metab. 87, 569–
    574. https://doi.org/10.1210/jcem.87.2.8261

Guan, Y., Wang, D., Bu, H., Zhao, T., Wang, H., 2020. The Effect of Metformin on Polycystic
    Ovary Syndrome in Overweight Women: A Systematic Review and Meta-Analysis of
    Randomized Controlled Trials. Int. J. Endocrinol. 2020.
    https://doi.org/10.1155/2020/5150684

Hurwitz, S., Plavnik, I., 1989. Severe Feed Restriction in Pullets During the Early Growing
    Period: Performance and Relationships Among Age, Body Weight, and Egg Weight at
    the Onset of Production. Poult. Sci. 68, 914–924. https://doi.org/10.3382/ps.0680914

Jakubowicz, D.J., Seppälä, M., Jakubowicz, S., Rodriguez-Armas, O., Rivas-Santiago, A.,
    Koistinen, H., Koistinen, R., Nestler, J.E., 2001. Insulin reduction with metformin
    increases luteal phase serum glycodelin and insulin-like growth factor-binding protein 1
    concentrations and enhances uterine vascularity and blood flow in the polycystic ovary
    syndrome. J. Clin. Endocrinol. Metab. 86, 1126–1133.
    https://doi.org/10.1210/jc.86.3.1126

Jing, Y., Wu, F., Li, D., Yang, L., Li, Q., Li, R., 2018. Metformin improves obesity-associated
    inflammation by altering macrophages polarization. Mol. Cell. Endocrinol. 461, 256–
    264. https://doi.org/10.1016/j.mce.2017.09.025

Kurzthaler, D., Hadziomerovic-Pekic, D., Wildt, L., Seeber, B.E., 2014. Metformin induces a
    prompt decrease in LH-stimulated testosterone response in women with PCOS
    independent of its insulin-sensitizing effects. Reprod. Biol. Endocrinol. 12, 1–6.
    https://doi.org/10.1186/1477-7827-12-98

Misichronis, G, Georgopoulos, N A, Marioli, D J, Armeni, A K, Katsikis, I, Piouka, A D,
Saltamavros, A D, Roupas, N D, Panidis, D, 2011. The influence of obesity on
androstenedione to testosterone ratio in women with polycystic ovary syndrome (PCOS)
and hyperandrogenemia. Gynecol. Endocrinol. 28, 249–252.
Moghetti, P., Castello, R., C Negri, Tosi, F., Perrone, F., Caputo, M., Zanolin, E., Muggeo, M.,
2000. Metformin effects on clinical features, endocrine and metabolic profiles, and
insulin sensitivity in polycystic ovary syndrome: a randomized, double-blind, placebo-
controlled 6-month trial, followed by open, long-term clinical evaluation. J. Clin.
Endocrinol. Metab. 85, 139–146. https://doi.org/10.1210/jcem.85.1.6293
Morin-Papunen, L.C., Koivunen, R.M., Ruokonen, A., Martikainen, H.K., 1998. Metformin
therapy improves the menstrual pattern with minimal endocrine and metabolic effects in
women with polycystic ovary syndrome. Fertil. Steril. 69, 691–696.
https://doi.org/10.1016/S0015-0282(98)00011-9
Morley, L.C., Tang, T., Yasmin, E., Norman, R.J., Balen, A.H., 2017. Insulin-sensitising drugs
(metformin, rosiglitazone, pioglitazone, D-chiro-inositol) for women with polycystic
ovary syndrome, oligo amenorrhoea and subfertility. Cochrane Database Syst. Rev. 2017.
https://doi.org/10.1002/14651858.CD003053.pub6
Munzker, J, Lindheim, L, Adaway, J, Trummer, C, Lerchbaum, E, Pieber, T R, Keevil, B,
Obermayer-Pietsch, 2017. High salivary testosterone-to-androstenedione ratio and
adverse metabolic phenotypes in women with polycystic ovary syndrome. Clin.
Endocrinol. (Oxf.) 86, 567–575. https://doi.org/10.1111/cen.13299
Navara, Kristen J, Pinson, Sara E, Chary, Parag, Taube, Patrick C, 2015. Higher rates of internal
ovulations occur in broiler breeder hens treated with testosterone. Poult. Sci. 94, 1346–
1352. https://doi.org/10.3382/ps/pev103
Nestler, J.E., Jakubowicz, D.J., 1996. Decreases in Ovarian Cytochrome P450c17a Activity and
Serum Free Testosterone after Reduction of Insulin Secretion in Polycystic Ovary
Syndrome. N. Engl. J. Med. 335, 617–623.
Nestler, J.E., Jakubowicz, D.J., Evans, W.S., Pasquali, R., 1998. Effects of metformin on
spontaneous and clomiphene-induced ovulation in the polycystic ovary syndrome. N.
Engl. J. Med. 338, 1876–1880.
Nestler, J.E., Stovall, D., Akhter, N., Iuorno, M.J., Jakubowicz, D.J., 2002. Strategies for the use
of insulin-sensitizing drugs to treat infertility in women with polycystic ovary syndrome.
Fertil. Steril. 77, 209–215. https://doi.org/10.1016/S0015-0282(01)02963-6
Pan, Y.-E., Liu, Z.-C., Chang, C.-J., Xie, Y.-L., Chen, C.-Y., Chen, C.-F., Walzem, R.L., Chen,
S.-E., 2012. Ceramide accumulation and up-regulation of proinflammatory interleukin-1β
exemplify lipotoxicity to mediate declines of reproductive efficacy of broiler hens.
Domest. Anim. Endocrinol. 42, 183–194.
https://doi.org/10.1016/j.domaniend.2011.12.001
Pasquali, R., Pelusi, C., Genghini, S., Cacciari, M., Gambineri, A., 2003. Obesity and
reproductive disorders in women. Hum. Reprod. Update 9, 359–372.
https://doi.org/10.1093/humupd/dmg024
Perriello, G., Misericordia, P., Volpi, E., Santucci, A., Santucci, C., Ferrannini, E., Ventura,
M.M., Santeusanio, F., Brunetti, P., Bolli, G.B., 1994. Acute antihyperglycemic
mechanisms of metformin in NIDDM: Evidence for suppression of lipid oxidation and
hepatic glucose production. Diabetes 43, 920–928. https://doi.org/10.2337/diab.43.7.920

Prager, R., Schernthaner, G., Graf, H., 1986. Effect of metformin on peripheral insulin sensitivity in non insulin dependent diabetes mellitus. Diabete Metab. 12, 346–350.

Rencbar, S., Kurnaz Ozbek, S., Eraldemır, C., Sezer, Z., Kum, T., Ceylan, S., Guzel, E., 2018. Effect of resveratrol and metformin on ovarian reserve and ultrastructure in PCOS: an experimental study. J. Ovarian Res. 11, 55–55. https://doi.org/10.1186/s13048-018-0427-7

Sam, S., 2007. Obesity and Polycystic Ovary Syndrome. Obes. Manag. 3, 69–73. https://doi.org/10.1089/obe.2007.0019

Singh, R.P., Moudgal, R.P., Agarwal, R., Sirajuddin, M., Mohan, J., Sastry, K.V.H., Tyagi, J.H., 2013. Ovarian morphology and internal vis-à-vis non internal laying in relation to triacylglycerol, hormones and their receptors concentration around the age of sexual maturity in broiler breeder hens. Br. Poult. Sci. 54, 157–163. https://doi.org/10.1080/00071668.2013.764491

Taherkhani, R., Zaghari, M., Shivazad, M., Zare Shahneh, A., 2010. A twice-a-day feeding regimen optimizes performance in broiler breeder hens. Poult. Sci. 89, 1692–1702. https://doi.org/10.3382/ps.2009-00488

Tang, T., Glanville, J., Hayden, C.J., White, D., Barth, J.H., Balen, A.H., 2006a. Combined lifestyle modification and metformin in obese patients with polycystic ovary syndrome. A randomized, placebo-controlled, double-blind multicentre study. Hum. Reprod. 21, 80–89. https://doi.org/10.1093/humrep/dei311

Tang, T., Glanville, J., Orsi, N., Barth, J.H., Balen, A.H., 2006b. The use of metformin for women with PCOS undergoing IVF treatment. Hum. Reprod. 21, 1416–1425. https://doi.org/10.1093/humrep/del025

Tokubuchi, I., Tajiri, Y., Iwata, S., Hara, K., Wada, N., Hashinaga, T., Nakayama, H., Mifune, H., Yamada, K., 2017. Beneficial effects of metformin on energy metabolism and visceral fat volume through a possible mechanism of fatty acid oxidation in human subjects and rats. PLoS ONE 12, 1–16. https://doi.org/10.1371/journal.pone.0171293

Tso, L.O., Costello, M.F., Albuquerque, L.E.T., Andriolo, R.B., Macedo, C.R., 2020. Metformin treatment before and during IVF or ICSI in women with polycystic ovary syndrome. Cochrane Database Syst. Rev. 2020. https://doi.org/10.1002/14651858.CD006105.pub4

van Santbrink, E.J.P., Hohmann, F.P., Eijkemans, M.J.C., Laven, J.S.E., Fauser, B.C.J.M., 2005. Does metformin modify ovarian responsiveness during exogenous FSH ovulation induction in normogonadotrophic anovulation? A placebo-controlled double-blind assessment. Eur. J. Endocrinol. 152, 611–617. https://doi.org/10.1530/eje.1.01866

Velazquez, E.M., Acosta, A., Mendoza, S.G., 1997. Menstrual Cyclicity After Metformin Therapy in Polycystic Ovary Syndrome. Obstet. Gynecol. 90, 392–395. https://doi.org/10.1016/S0029-7844(97)00296-2

Velazquez, E.M., Mendoza, S., Hamer, T., Sosa, F., Glueck, C.J., 1994. Metformin therapy in polycystic ovary syndrome reduces hyperinsulinemia, insulin resistance, hyperandrogenemia, and systolic blood pressure, while facilitating normal menses and pregnancy. Metabolism 43, 647–654. https://doi.org/10.1016/0026-0495(94)90209-7

Vrbikova, J., Hill, M., Starka, L., Cibula, D., Bendlova, B., Vondra, K., Sulcova, J., Snajderova, M., 2001. The effects of long-term metformin treatment on adrenal and ovarian steroidogenesis in women with polycystic ovary syndrome. Eur. J. Endocrinol. 619–628. https://doi.org/10.1530/eje.0.1440619

Walzem, R.L., Chen, S., 2014. Obesity-Induced Dysfunctions in Female Reproduction: Lessons
from Birds and Mammals. Adv. Nutr. 5, 199–206. https://doi.org/10.3945/an.113.004747
Weaver, E.A., Ramachandran, R., 2020. Metformin attenuates steroidogenesis in ovarian
follicles of the broiler breeder hen. Reproduction 160, 659–672.
https://doi.org/10.1530/REP-20-0066
Xing, C., Li, C., He, B., 2020. Insulin Sensitizers for Improving the Endocrine and Metabolic
Profile in Overweight Women with PCOS. J. Clin. Endocrinol. Metab. 105, 2950–2963.
https://doi.org/10.1210/clinem/dgaa337
Yu, M.W., Robinson, F.E., Charles, R.G., Weingardt, R., 1992. Effect of Feed Allowance
During Rearing and Breeding on Female Broiler Breeders: 2. Ovarian Morphology and
Production. Poult. Sci. 71, 1750–1761. https://doi.org/10.3382/ps.0711750
Yuan, T., Lien, R.J., McDaniel, G.R., 1994. Effects of Increased Rearing Period Body Weights
and Early Photostimulation on Broiler Breeder Egg Production. Poult. Sci. 73, 792–800.
https://doi.org/10.3382/ps.0730792
Zhang, Y., Hu, M., Meng, F., Sun, X., Xu, H., Zhang, J., Cui, P., Morina, N., Li, X., Li, W., Wu,
X.K., Brännström, M., Shao, R., Billig, H., 2017. Metformin Ameliorates Uterine Defects
in a Rat Model of Polycystic Ovary Syndrome. EBioMedicine 18, 157–170.
https://doi.org/10.1016/j.ebiom.2017.03.023
Zuidhof, M.J., Schneider, B.L., Carney, V.L., Korver, D.R., Robinson, F.E., 2014. Growth,
efficiency, and yield of commercial broilers from 1957, 1978, and 2005. Poult. Sci. 93,
2970–2982. https://doi.org/10.3382/ps.2014-04291

Figure Legends

Figure 1. Experimental design and data collection timeline. Broiler breeder hens were raised from day-old chicks and randomly assigned to a treatment group (n=45 hens/treatment group) to receive metformin (0, 25, 50 or 75 mg/kg body weight) in the diet starting from 25-65 weeks of age. Blood samples were collected from a subset of hens (n=6 hens/treatment group) every 5 weeks starting from 30 weeks of age. A subset of hens (n=10 hens/treatment group) were weighed every 10 weeks. At the end of the study, a subset of hens (n=12 hens/treatment group) were euthanized to collect tissues.

Figure 2. Effect of metformin supplementation on body weight (A) and abdominal fat pad weight (B). Broiler breeder hen diet was supplemented with metformin (0, 25, 50, or 75 mg/kg bw) from 25 to 65 weeks of age. All hens (n=45 hen/treatment group) were weighed at 25 weeks of age, and a random subset (n=10 hens/treatment group) were weighed every 10 weeks from 30 to 65 weeks of age. The entire abdominal fat pad was collected during necropsy done at the end of study. Values are means ± SEM. * $P \leq 0.05$, ** $P \leq 0.01$, *** $P \leq 0.001$, **** $P \leq 0.0001$.

Figure 3. Effect of metformin supplementation on the ovarian follicular hierarchy. Broiler breeder hen diet was supplemented with metformin (0, 25, 50, or 75 mg/kg bw) from 25 to 65 weeks of age. Representative photographs of the ovarian follicular hierarchy from a broiler breeder hen that received 0 or 25 mg/ kg body weigh metformin (A and B, respectively). A double hierarchy of preovulatory follicles (denoted F1-F5) and hyper-recruitment of 3-5 mm and 6-8 mm sized prehierarchical follicles are seen in **(A)** while hens which received 25 mg

metformin/kg body weight display a normal hierarchy of preovulatory and prehierarchical

follicles in **(B)**.

Figure 4. Effect of metformin supplementation on egg production. Broiler breeder hen diet

was supplemented with metformin (0, 25, 50, or 75 mg/kg bw) from 25 to 65 weeks of age. Egg

production was recorded daily for each hen from 22 weeks of age through 65 weeks of age

(n=40-45 hens/treatment group). Average cumulative egg production per hen every 5 weeks over

entire course of the study **(A)** and the overall average cumulative egg production per hen **(B).**

Values are means ± SEM. * $P \leq 0.05$, ** $P \leq 0.01$, *** $P \leq 0.001$, **** $P \leq 0.0001$.

Figure 5. Effect of metformin supplementation on plasma progesterone concentration.

Broiler breeder hen diet was supplemented with metformin (0, 25, 50, or 75 mg/kg bw) from 25

to 65 weeks of age. Progesterone concentrations (ng/mL) were determined by

radioimmunoassay in blood samples collected from a subset of hens (n=6 hens/treatment group)

every 5 weeks from 25-65 weeks of age **(A)**. Plasma progesterone concentrations (ng/ml) within

each treatment group over the course of the study is shown in **(B)**.

[A, B, C, D, E, F] P < 0.05 by ANOVA; n= 6 hens/treatment group for each time point.

Figure 6. Effect of metformin supplementation on plasma total estrogen concentration.

Broiler breeder hen diet was supplemented with metformin (0, 25, 50, or 75 mg/kg bw) from 25

to 65 weeks of age. Total estrogen concentrations (ng/mL) were determined by

radioimmunoassay in blood samples collected from a subset of hens (n=6 hens/treatment group)

every 5 weeks from 25-65 weeks of age **(A)**. Plasma total estrogen concentrations (ng/ml)

within each treatment group over the course of the study is shown in **(B)**.

[A, B,C, D] $P < 0.05$ by ANOVA; n= 6 hens/treatment group for each time point.

Figure 7. Effect of metformin supplementation on plasma testosterone concentration.

Broiler breeder hen diet was supplemented with metformin (0, 25, 50, or 75 mg/kg bw) from 25

to 65 weeks of age. Testosterone concentrations (ng/mL) were determined by

radioimmunoassay in blood samples collected from a subset of hens (n=6 hens/treatment group)

every 5 weeks from 25-65 weeks of age **(A)**. Plasma testosterone concentrations (ng/ml) within

each treatment group over the course of the study is shown in **(B).**

[A, B,] $P < 0.05$ by ANOVA; n= 6 hens/treatment group for each time point.

Figure 8. Effect of metformin supplementation on plasma androstenedione concentration.

Broiler breeder hen diet was supplemented with metformin (0, 25, 50, or 75 mg/kg bw) from 25

to 65 weeks of age. Androstenedione concentrations (ng/mL) were determined by

radioimmunoassay in blood samples collected from a subset of hens (n=6 hens/treatment group)

every 5 weeks from 25-65 weeks of age **(A)**. Plasma androstenedione concentrations (ng/ml)

within each treatment group over the course of the study is shown in **(B)**.

[A, B,] $P < 0.05$ by ANOVA; n= 6 hens/treatment group for each time point.

Figure 9. Effect of metformin supplementation on the plasma testosterone to

androstenedione ratio. Broiler breeder hen diet was supplemented with metformin (0, 25, 50,

or 75 mg/kg bw) from 25 to 65 weeks of age. The plasma testosterone to androstenedione ratio

was calculated from the plasma androstenedione and testosterone concentrations (see Fig. 6 and

7) and represented 25-65 weeks of age **(A)** or within each treatment group over the course of the

study **(B).**

[A, B] $P < 0.05$ by ANOVA; n= 6 hens/treatment group for each time point.

Weaver et al., 2022- Figure 1

A

B

Weaver et al., 2022- Figure 3

A

0 mg/kg metformin

B

25 mg/kg metformin

Weaver et al., 2022- Figure 4

A

B

Weaver et al., 2022- Figure 9