1

Selective loss of CD107a TIGIT+ memory HIV-1-specific CD8+ T cells in PLWH over a decade of ART

Oscar Blanch-Lombarte[1,2], Dan Ouchi[1], Esther Jiménez-Moyano[1], Julieta Carabelli[1],

Miguel Angel Marin[1], Ruth Peña[1], Adam Pelletier[4], Aarthi Talla[4], Ashish Sharma[4],

Judith Dalmau[1], José Ramón Santos[5,6], Rafik-Pierre Sékaly[4],

Bonaventura Clotet[1,3,5,6,7] and Julia G. Prado[1,3,8*]

1. IrsiCaixa AIDS Research Institute, Badalona, Barcelona, Spain
2. Universitat Autònoma de Barcelona, Cerdanyola del Vallès, Barcelona, Spain
3. Germans Trias i Pujol Research Institute (IGTP), Badalona, Spain.
4. Pathology Department, Case Western Reserve University, Cleveland, OH, United States.
5. Lluita contra les Infeccions Foundation, Hospital Universitari Germans Trias i Pujol, Badalona, Spain
6. Infectious Diseases Department, Hospital Universitari Germans Trias i Pujol, Badalona, Spain
7. Faculty of Medicine, University of Vic - Central University of Catalonia (UVic-UCC), Vic, Spain
8. CIBER Enfermedades Infecciosas (CIBERINFEC), Instituto de Salud Carlos III, Madrid, Spain

*Corresponding author: Julia G. Prado (jgarciaprado@irsicaixa.es). IrsiCaixa AIDS Research Institute, Germans Trias i Pujol Research Institute (IGTP), Crta. Canyet s/n; 08916, Badalona, Barcelona, Spain. Tel: +34 934 656 374.

Abstract

The co-expression of inhibitory receptors (IRs) is a hallmark of CD8+ T-cell exhaustion (Tex) in people living with HIV-1 (PLWH). Understanding alterations of IRs expression in PLWH on long-term antiretroviral treatment (ART) remains elusive but is critical to overcoming CD8+ Tex and designing novel HIV-1 cure immunotherapies. To address this, we combine high-dimensional supervised and unsupervised analysis of IRs concomitant with functional markers across the CD8+ T-cell landscape on 24 PLWH over a decade on ART. We define irreversible alterations of IRs co-expression patterns in CD8+ T cells not mitigated by ART and identify negative associations between the frequency of TIGIT+ and TIGIT+TIM-3+ and CD4+ T-cell levels. Moreover, changes in total, SEB-activated, and HIV-1-specific CD8+ T-cells delineate a complex reshaping of memory and effector-like cellular clusters on ART. Indeed, we identify a selective reduction of HIV-1 specific-CD8+ T-cell memory-like clusters sharing TIGIT expression and low CD107a that can be recovered by mAb TIGIT blockade independently of IFNγ and IL-2. Collectively, these data characterize with unprecedented detail the patterns of IRs expression and functions across the CD8+ T-cell landscape and indicate the potential of TIGIT as a target for Tex precision immunotherapies in PLWH at all ART stages.

Introduction

The ART introduction has been the most successful strategy to control viral replication, transforming HIV-1 into a chronic condition. However, ART does not cure the infection, and treatment is required lifelong due to a stable viral reservoir, raising the need to find a cure for people living with HIV-1 (PLWH). A sterilizing or functional cure aims to eliminate or control HIV-1 in the absence of ART. In both scenarios, HIV-1-specific CD8+ T cells are likely to play an essential role as they have been widely recognized as a critical factor in the natural control of viral replication (1–5). Proliferative capacity, polyfunctionality, and ex vivo antiviral potency are features of HIV-1-specific CD8+ T cells associated with spontaneous viral control (6–9).

Although ART in PLWH normalizes the levels of CD8+ T cells and potentially preserves their functional characteristics (9–12), microbial translocation and continuous immune activation lead to long-term CD8+ T-cell dysfunction and exhaustion (Tex), a critical barrier for HIV-1 curative interventions (13–15). CD8+ Tex is defined by the persistent co-expression of inhibitory receptors (IRs) and the progressive loss of immune effector functions linked to transcriptional, epigenetic and metabolic changes (16–19). In HIV-1 infection, IRs are continuously expressed despite long-term suppressive ART (20–26) and have been associated with disease progression and immune status in PLWH (25,27–32). Thus, the expression of IRs is linked to diminished functionality and is a hallmark of CD8+ Tex in PLWH.

The interest in evaluating the blocking of IRs or immune checkpoint blockade (ICB) in PLWH as a therapeutic strategy to reverse CD8+ Tex increases (26,27). Over the last years, several studies supported the recovery of proliferative capacity, cell survival, and

cytokine production of HIV-1-specific CD8+ T cells through ICB (33,34). The blockade

of the PD-1/PDL-1 axis has been extensively studied, demonstrating the functional

recovery of HIV-1-specific CD8+ T cells in PLWH (26,27). Moreover, alternative

pathways to PD-1/ PDL-1, including LAG-3, TIGIT, and TIM-3, have been explored as

candidates for ICB therapies for HIV-1 infection (33,35–37). Also, recent data support

the combinatorial use of ICB to favour synergistic effects on the recovery of HIV-1-

specific CD4+ and CD8+ T-cell function (38,39).

The unprecedented success of the clinical use of ICB in the cancer field (40,41) has

prompted the clinical evaluation of ICB in PLWH (42) to boost immunity to reduce or

eliminate the viral reservoir. However, clinical evidence on the impact of ICB as an HIV-

1 cure intervention continues to be controversial (34,43–47). In this context, the

simultaneous characterization of IRs co-expression and functional patterns across CD8+

T cells is critical to understanding Tex regulation in PLWH. This information is essential

to identify novel targets for precise immunotherapies in PLWH on ART (48).

To address these questions, we performed supervised and unsupervised

immunophenotypic analyses of IRs (PD-1, TIGIT, LAG-3, TIM-3, CD39), and functional

markers (CD107a, IFNγ, and IL-2) across the landscape of CD8+ T cells over a decade

of ART in PLWH and compared to PLWH with early infection and healthy individuals.

We profile changes of bulk, SEB-activated, and HIV-1-specific CD8+ T-cells in PLHW

and unfold a selective decrease of memory-like HIV-1-specific CD8+ clusters sharing

TIGIT expression and low CD107a in PLWH on ART. Moreover, TIGIT blockade

rescues CD107a expression without changes in IFNγ or IL-2 production on HIV-1-

specific CD8+ T-cells. Of note, the response to TIGIT, TIM-3, or TIGIT+TIM-3

blockade was heterogeneous across HIV-1-specific CD8+ T-cell differentiation stages

and functions, indicating the plasticity and complexity of the IRs pathways as targets for

immune-base cure interventions.

Results
114
Alterations in CD8+ T-cell IRs frequencies and expression patterns in PLWH are

not mitigated by ART

Although the co-expression of IRs is a hallmark of CD8+ Tex in HIV-1 infection (20–

26), a detailed characterization of the combinatorial expression of IRs across CD8+ T-

cell lineages in PLWH on long-term ART is still missing. To do this, we combined the

analyses of IRs (PD-1, TIGIT, LAG-3, TIM-3, and CD39) and lineage markers

(CD45RA, CCR7, and CD27) in CD8+ T cells from longitudinal samples in PLWH on

ART by flow-cytometry (**Supplemental Figure 1**). We compare three groups; healthy

controls (HC), PLWH with early infection (Ei), and PLWH on ART (S) with longitudinal

samples available a median period of 2.2 (S1) and 10.1 (S2) years on fully suppressive

ART (**Figure 1A, Source data Figure 1A**). The epidemiological and clinical

characteristics of the study groups are detailed in **Supplemental Table 1** and **Source**

data Figure 1A.

As shown in **Figure 1B (Source data Figure 1 B-C)**, we found persistent alterations in

the expression and co-expression of IRs in naïve, central memory (CM), and transitional

memory (TM) CD8+ T cells in PLWH on ART. These perturbations were maintained

despite prolonged ART (S2) in naïve and CM, but not TM cells compared to HC.

Deconvolution of IRs co-expression patterns by the number of receptors expressed (0, 1,

2, >3) further delineates a significant reduction of Naïve, CM, and TM CD8+ T cells

lacking IRs expression concomitant with a significant increase of CD8+ T cells

expressing one (Naïve and CM) or >3 IRs (TM) on ART (**Figure 1C**). Moreover, we

observed an augment in effector memory (EM), TM, and effector (EFF) CD8+ T cells

co-expressing >3 IRs in Ei and S1 that normalized in S2 (**Figure 1C**). Of note, out of the

32 possible combinations of IRs expression studied in CD8+ T-cell subsets

(**Supplemental Figure 2C, Source data Figure S2**), single TIGIT$^+$ expression in CM

and dual TIGIT$^+$TIM-3$^+$ co-expression in EFF CD8+ T cells accounted for continuous

increases in frequency under suppressive ART (**Figure 1D, Source data Figure 1D**). We

confirmed similar increases in the frequency of TIGIT in total, CM, and TM CD8+ T

cells on ART. These data contrast with transient changes in other IRs upon infection

normalizing with ART (**Supplemental Figure 2A-B, Source data Figure S2**).

These initial findings led us to postulate associations between the expression of IRs in

CD8+ T cells, persistent immune activation, and the degree of CD4+ T-cell immune

recovery in PLWH on ART. For this purpose, we performed correlation analyses that

revealed several negative associations between the frequency of CD8+ T cells expressing

IRs and CD4+ T-cell counts across study groups (**Figure 1, E-G, Source data Figure**

1E-G). Focusing on S1 (**Figure 1F**), we found significant negative correlations between

CD4+ T-cell counts and frequencies of CM CD8+ T cells expressing 2 (p=0.0054, r=-

0.64) and > 1 IRs (p=0.0087, r=-0.61). Focusing on S2 (**Figure 1G**), we observed

significant negative correlations between CD4+ T-cell counts and the frequency of total

CD8+ T cells expressing TIGIT$^+$ (p=0.0157, r=-0.58), expressing 1 IRs (p=0.0386, r=-

0.54) or >1 IRs (p=0.0386, r=-0.51). At the level of CD8+ T-cell subsets, the expression

of 2 IRs in CM and >1 IR in TM negatively correlated with CD4+ T-cell counts

(p=0.0072, r=-0.64; p=0.0346, r=-0.52, respectively) (**Figure 1G**). These correlations

further indicate a negative relationship between IRs expression patterns and immune

status in PLWH on long-term suppressive ART.

In summary, these data support changes in IRs expression not mitigated by long-term

ART in total and CD8+ T-cell subsets expressing one or >1 IRs, particularly TIGIT$^+$ and

TIGIT$^+$TIM-3$^+$. These findings also uncover negative associations between IRs

expression in CD8+ T cells and CD4+ T cell levels in PLWH on ART.

Unsupervised phenotypic analyses of IRs across the CD8+ T-cell landscape in

PLWH on ART

Next, to further characterize IRs expression across CD8+ T cells in PLWH on ART, we

performed an unsupervised net-SNE analysis of flow-cytometry data. We concatenated

1,988,936 total CD8+ T cells and analyzed the phenotypes with the topographical regions

of each surface marker tested (**Figure 2, A-B, Source data Figure 2A-D**). CD8+ T cells

were classified into 38 cellular clusters distributed according to the relative marker

expression of 14 parameters and represented using net-SNE and heatmaps (**Figure 2, C-**

D).

Out of the 38 clusters identified, we found eight cellular clusters (#1, #2, #6, #7, #9, #10,

#11, and #12) with significant differences by inter- and intragroup comparisons (**Figure**

2D-E). Most of the differentially expressed clusters shared memory-like phenotypes. Of

note, clusters #6, #7, and #12 shared memory-like phenotypes and low expression of IRs.

Meanwhile, clusters #9 and #10 shared effector-like phenotypes and co-expression of IRs,

including TIGIT, LAG-3, and low TIM-3 (**Figure 2D**). Briefly, intergroup analyses

demonstrated significant changes in composition and frequency with a decrease of #1,

#6, and an increase of #10 in Ei compared with HC. Also, clusters #1 and #2 decreased

in S1 and S2, respectively, and clusters #9, #10, #11, and #12 increased in S1 while tended

to normalize in S2 compared with HC (**Figure 2E, Source data Figure 2 E-F**).

Intragroup analyses further supported changes in cluster frequency and composition

during long-term ART. Meanwhile, clusters #6 and #7 increased, and #9 and #11

decreased over time on ART (**Figure 2F**). These data support an expansion of memory-

like clusters with low IR expression (#6 and #7) and a contraction of effector-like clusters

sharing TIGIT expression (#9 and #11) during ART. Thus, unsupervised analyses support

cellular clusters' continuous expansion and contraction in frequency and composition

across the landscape of CD8+ T cells in PLWH on ART.

Unsupervised phenotypic characterisation of SEB-activated CD8+ T cells in PLWH

on ART

Then, we evaluate CD8+ T-cell responses by bacterial superantigen activation with

Staphylococcal enterotoxin B (SEB). Using SEB can provide complementary information

on T-cell activation in response to pathogens involved in the disease by stimulating TCR-

VB clonotypes (31,49,50). In this context, we analysed IRs expression and functional

markers using unsupervised net-SNE analysis. We defined SEB-activated CD8+ T cells

by the expression of at least one functional marker (CD107a, IFNγ, IL-2) upon incubation

with SEB, as previously described (51-55) (**Figure 3, A-B, Source data Figure 3A-D**).

We concatenated 253,021 SEB-activated CD8+ T cells and identified 29 unique clusters

represented by net-SNE and heatmaps (**Figure 3, C-D**). Only six of the 29 clusters

showed statistical differences by inter- and intragroup comparisons (**Figure 3D**). All

differential clusters shared memory-like phenotypes (#2, #3, #5, #8, and #14) except

cluster #6, with effector-like phenotype and low TIGIT expression. In addition, we

observed a functional exclusion of clusters expressing IL-2 (#5 and #6) and those

expressing CD107a and IFNγ (#2, #3, and #8) (**Figure 3D**). Intergroup comparisons

identified increases in cluster #2 and a reduction of #14 and #5 with HIV-1 infection

compared to HC. Additionally, ART was linked to the decrease of #5, #6, and #14 in S1

and #3 in S2 when compared with HC (**Figure 3E, Source data Figure 3 E-F**).

Moreover, intragroup analyses identified an increase in clusters #6 and #14 and a

reduction in #3 and #8 on ART. Of note, #6 characterizes by IL-2 expression in the

without TIGIT expression. Meanwhile, clusters #3 and #8 express CD107a, IFNγ and

variable expression of IRs (**Figure 3F**). In agreement with unsupervised clustering

analyses, classical supervised analyses identified an augment of CD107a and IFNγ SEB-

activated CD8+ T cells with Ei that normalized over time on ART. Also, we delineate

significant increases of IL-2 SEB-activated CD8+ T cells across subsets and time on ART

(**Supplemental Figure S3 A-C, Source data Figure S3**).

These data support plasticity in the composition of SEB-activated CD8+ T-cell clusters

with HIV-1 infection and ART. We observed the dominance of memory-like clusters with

changes in composition and frequency in PLWH on ART, particularly IL-2 expression.

Reduction of HIV-1-specific CD8+ T-cell clusters sharing memory-like phenotypes,

TIGIT expression and low CD107a

Next, we characterize HIV-1-specific CD8+ T cells in PLWH on long-term ART

compared to Early infected individuals (Ei), aiming to identify signatures of cellular

dysfunction. We performed unsupervised net-SNE analyses in 53,751 cells concatenated,

based on the production of at least one functional marker (CD107a, IFNγ, IL-2) in

response to HIV-1-Gag peptides combined with lineage markers and IRs (**Figure 4, A-**

B, Source data Figure 4 A-D). We defined 26 HIV-1-specific clusters by net-SNE

analysis. Of note, only three showed significant differences between groups (#1, #2, and

#3) (**Figure 4, C-D**). All three clusters decreased in frequency on ART and shared

memory-like features (**Figure 4E, Source data Figure 4E**). Clusters #1 and #2 shared

co-expression of IRs, mainly PD-1 and TIGIT, low CD107a, IFNγ, and higher expression

of IL-2 than #3. Meanwhile, cluster #3 co-express (TIGIT, PD-1, and TIM-3) low

expression of IL-2 and higher expression of CD107a and IFNγ (**Figure 4D**). Although

CD107a, IFNγ, and IL-2 expression differed between clusters #1, #2, and #3, all shared

TIGIT expression in the context of variable levels of TIM-3. These findings in memory-

like clusters, together with the initial ones, accounting for increases in the frequency of

TIGIT$^+$ and TIGIT$^+$TIM-3$^+$ in CM and EFF cells on ART, led us to postulate then as

potential markers of HIV-1-specific CD8+ T-cell dysfunction in PLWH on ART.

Despite no significant changes observed in the total frequency of CD107a, IFNγ and IL-

2 HIV-1-specific CD8+ T-cell responses between groups (**Supplemental Figure 4A-B**,

Source data Figure S4). The analyses of TIGIT and TIGIT+TIM-3 HIV-1-specific

memory subsets revealed decreases in the frequency of CD107a TIGIT HIV-1-specific

CD8+ T cells limited to the CM compartment (**Figure 4F, Source data Figure 4F-G**).

No changes in IFNγ and IL-2 were observed. Furthermore, polyfunctional analysis of

TIGIT HIV-1-specific CM CD8+s identified a decrease in monofunctional CD107a$^+$ as

well as in bifunctional CD107a$^+$IFNy$^+$ and CD107$^+$IL-2$^+$ cells overtime on ART (**Figure**

4G). Overall, these data support a reduction of HIV-1-specific CD8+ T-cell clusters

sharing memory-like phenotypes, TIGIT expression and low CD107a in PLWH on ART.

TIGIT blockade restores CD107a expression but not IFNγ or IL-2 production in

HIV-1-specific CD8+ T cells

Then, we decided to explore the blockade of TIGIT and TIM-3 pathways as targets for

the recovery of CD107a and potentially IFNγ and IL-2 production in HIV-1-specific

CD8+ T-cells. We performed short-term ICB experiments using monoclonal antibodies

αTIGIT, αTIM-3, and αTIGIT+ αTIM-3 in PBMC from PLWH on suppressive ART (S)

with previous immunophenotype.

After short-term ICB, we monitored changes in CD107, IFNγ, and IL-2 in total and

subsets of HIV-1-specific CD8+ T cells by flow cytometry (**Figure 5A)**. The Net-SNE

TIGIT and TIM-3 projections are represented in **Figure 5B (Source data Figure 5B)**. At

the level of total HIV-1-specific CD8+ T-cells, short-term ICB experiments demonstrate

a specific increase of CD107a expression by αTIGIT (isotype vs. αTIGIT; $p<0.05$) and

αTIGIT + αTIM-3 (isotype vs. αTIGIT + αTIM-3; $p<0.05$) blockade (**Figure 5B, left**).

Moreover, the recovery of CD107a by αTIGIT was consistent across HIV-1-specific

CD8+ T-cell subsets, being particularly marked for CM (isotype vs. αTIGIT; in CM

$p<0.005$) (**Figure 5C**) in agreement with our previous findings. Of note, αTIM-3

blockade did not show any effect, and dual blockade of αTIM-3+αTIGIT did not reveal

an additive effect (**Figure 5C**). No changes in IFNγ and IL-2 production were observed

for any conditions tested (**Figure 5, D-E, Source data Figure 5 D-E**).

These data support heterogeneity in the functional recovery of HIV-1-specific CD8+ T-

cells by differentiation stage based on αTIGIT, αTIM-3, and αTIGIT+αTIM-3 blockade.

Overall, these results identify the targeting of TIGIT to recover the degranulation in HIV-

1-specific CD8+ T cells, particularly within the CM compartment in PLWH on ART.

Discussion

CD8+ Tex displays a range of functional defects in PLWH early in HIV-1 infection and during ART (16–19). The expression of IRs is a hallmark of Tex (20,22,23) (14), and co-expression of IRs has been associated with HIV-1 disease progression (15,25,29,30,32) and cancer severity (56–58). Although ICB has demonstrated promising results in cancer remission (40,41), its applicability in HIV-1 as a cure intervention remains unclear (34,43–47). Therefore, it is essential to understand the patterns of IRs expression and function across the CD8+ T-cell landscape to identify targets for ICB broadly applicable to PLWH on ART (48).

Here, we immunophenotype CD8+ T cells using five IRs and three functional markers in PLWH over the ten years of ART. In this way, we overcome previous study limitations based on single IRs expression, bulk CD8+ T cells, and cross-sectional data (20,23,25,29,32). Our results demonstrated a marked and significant increase in TIGIT CD8+ T cells, particularly within the central memory compartment, not ameliorated by long-term ART. Also, TIGIT CD8+ T cells negatively correlated with CD4+ counts in PLWH on ART. These data support continuous expression of TIGIT despite ART in agreement with previous studies (24,25,30,57,59) and uncover novel associations between TIGIT expression in CD8+ T cells and poorer immune status in PLWH on ART. Thus, these data indicate a specific contribution of TIGIT expression to persistent immune activation and poor CD4+ recovery on ART.

In contrast, we observed transient increases of PD-1, LAG3, TIM3 and CD39 expression in total, memory and effector CD8+ T cells that normalize over time on ART. The potential biological implications of such a difference may relate to the nature of each

receptor and the specific immune regulatory pathway activated during HIV-1 infection

and ART. Also, these divergences may be influenced by the presence of γ-chain

cytokines, such us IL-2, IL-15 and IL-21, in plasma able to upregulate the TIGIT

expression in CD8+ T cells (25). A previous study demonstrated an association between

high levels of IL-15 and high TIGIT expression in CD4+ T cells with suboptimal CD4+

recovery in PLWH on ART (60).

Our study combined high-dimensional supervised and unsupervised analysis providing

an unprecedented deep immunophenotype of the CD8+ T-cell landscape in PLWH over

a decade on ART. We explored complementary levels of complexity for the

characterization of CD8+ T cells, including an absence of stimuli (bulk), the presence of

antigen-independent stimuli (SEB), and the presence of antigen-specific stimuli (HIV-1).

In the absence of stimuli, supervised analyses confirmed heterogeneous and complex

patterns of IRs co-expression across CD8+ T cell lineages altered by HIV-1 infection and

shaped by ART (16,21,30,61,62). Furthermore, unsupervised analyses added complexity

to previous data by delineating in profound detail early and continuous changes of

contraction and expansion of cellular clusters with HIV-1 infection and time on ART. We

tracked memory-like expansion and effector-like cluster contraction over time on ART

according to the establishment of memory responses. Similarly, in the presence of

antigen-independent stimuli, we identified continuous changes in cellular cluster

composition by contraction and expansion of CD8+ cellular cluster with infection and

treatment. We observed a dominance of memory-like clusters with changes in

composition and frequency tracked by an augment of IL-2 expressing clusters on ART

both by supervised and unsupervised analyses. The IL-2 expression regulates

proliferation and homeostasis and contributes to the generation of long-term memory

responses (31,49,50), suggesting a partial functional remodelling of CD8+ T-cell

populations independent of antigen in PLWH over time on ART. Thus, our findings

support the contribution of IRs co-expression in CD8+ Tex and T-cell activation

favouring a continuous reshaping of memory and effector-like CD8+ cellular clusters.

These findings indicate the enormous plasticity and constant homeostasis of CD8+ T cells

in PLWH during a decade of ART (63).

In the presence of HIV-1-specific stimuli, supervised and unsupervised analyses delineate

a reduction of HIV-1-specific CD8+ T cells sharing memory phenotypes, TIGIT

expression and low CD107a. Our findings focused on the memory compartment of TIGIT

expressing HIV-1-specific CD8+ T cells demonstrating a decrease in monofunctional

$CD107a^+$ and bifunctional $CD107a^+IFN\gamma^+$ and $CD107a^+IL-2^+$ cells. Previous studies

support a direct correlation between monofunctional $CD107a^+$ and Eomes intensity in

exhausted HIV-1-specific CD8+ T cells (17). Although this study did not include the

expression of TIGIT across analyses, it may suggest an association between TIGIT

expression and the T-bet /Eomes axis in charge of regulating the exhaustion and memory

cell fate of CD8+ T cells (64).

Our findings further support the role of TIGIT as a signature of dysfunctional and Tex

antiviral responses (25). Indeed, the blockade of the TIGIT pathway restored CD107a

expression in HIV-1-specific CD8+ T cells across cellular compartments with a marked

effect in the central memory compartment according to our findings from HIV-1-specific

immunophenotype of TIGIT CD8+ T cells. To our knowledge, our study is the first to

demonstrate the recovery of CD107a expression in HIV-1-specific CD8+ T cells by

TIGIT blockade. These data contrast with Chew et al., which observed the recovery of

IFNγ production by TIGIT blockade. However, they also reported a reduction in CD107a

expression in TIGIT+ CD8 T cells in response to aCD3/aCD28 activation, supporting a

dysfunctional profile of TIGIT+ CD8+ T cells. Differences between study groups

accounting for time on ART, samples tested and interindividual variability of in vitro ICB

experiments may account for some of the differences observed.

Although the mechanistic behind TIGIT signalling and CD107a expression are not fully

understood, low CD107a expression has been linked to the terminal T-bet^{dim}Eomes^{hi}

exhausted phenotype, and HIV-1-specific CD8+ T cells expressing TIGIT can

degranulate to a certain extent (17,24). Moreover, our data did not support an additive

effect recovering HIV-1-specific CD8+ T cells function of TIGIT and TIM-3

combinatorial blockade over TIGIT blockade. The redundancy and promiscuity of TIM-

3 for several ligands, including Gal-9, CEACAM-1, PtdSer, and HMGB-1 (65,66), may

be associated with these results. We cannot exclude the impact of TIGIT and TIM-3

blockade in other cell types (57,58,67).

We acknowledge several study limitations; First, the sample size of study groups and the

use of peripheral blood samples underestimate the potential contribution of TIGIT

expression to Tex in lymphoid tissues. Second, the use of only Gag as stimuli for the

characterization of HV-1-specific CD8+ T cell responses in the absence of TCR

sequencing. Using alternative HIV-1 antigens such as Nef, Env or Pol may provide

additional information on the profile of CD8+ T-cell functional responses against early

and late-expressed viral proteins in PLWH on ART (68, 69). Third, limited ICB

experiments to CD107a, INFγ and IL-2 functional markers without complementary

cytotoxic markers (perforin, granzyme B). Forth, complementary transcriptomic,

epigenetic, and metabolic markers are needed for a complete description of Tex's immune

signatures linked to TIGIT expression in HIV-1 specific CD8+ T cells in PLWH on ART.

In summary, our study profile with unprecedented detail continuous reshaping of

memory-like and effector-like of CD8+ T cellular clusters in PLWH over a decade of

ART. The study identifies the TIGIT as a critical target for T-cell dysfunction, and Tex

associated the loss of CD107a expression in HIV-1-specific CD8+ T cells. These findings

support targeting the TIGIT/CD155 axis for Tex precision immune-base curative

interventions in PLWH at all ART stages.

Materials and Methods

Study groups

This retrospective study analyzed clinical data and biological sample availability from

3,000 patients assigned to the HIV-1 clinical unit of the Germans Trias i Pujol University

Hospital. We included individuals with cryopreserved PBMCs available in our collection.

We identified 24 chronically HIV-1-infected individuals who had been treated mainly

with a combination of NNRTI and NRTI for more than ten years with sustained

virological suppression (<50 HIV-1-RNA copies/ml) and with longitudinal biological

samples at timepoint 1 (S1), 2.2 (1.8 - 2.8) years undetectable on ART, and at time point

2 (S2), 10.1 (7.4 – 12.9) years undetectable on ART (**Supplemental Table 1, Figure 1A,**

Source data Figure 1A). We excluded individuals with integrase inhibitors, ART as

monotherapy, and treatments with mitochondrial toxicity, including Trizivir, d4T, ddI,

AZT and blips over the ART period (S1- S2) to ensure homogeneous treatment over time.

For comparative purposes, we included 24 early HIV-1-infected individuals (Ei) defined

in a window of 1.3 (0.77 - 17.8) weeks after seroconversion in the absence of ART and

24 healthy controls (HC). The groups were balanced by age to the S2 samples to avoid

confounding effects on IR expression.

CD8+ T cell immunophenotype

Cryopreserved PBMCs from the study groups were thawed and rested overnight at 37ºC

in a 5% CO_2 incubator. The following day, PBMCs were incubated for six hours at 37ºC

in a 5% CO_2 incubator under RPMI complemented medium 10% FBS in the presence of

CD28/49d co-stimulatory molecules (1µl/ml, BD), Monensin A (1µl/ml, BD Golgi

STOP), and anti-human antibody for CD107a (PE-Cy5, clone H4A3, Thermo Fisher

Scientific). PBMCs were left unstimulated, stimulated with SEB (1µg/ml, Sigma-

Aldrich), and stimulated with HIV-1-Gag peptide pool (2µg/peptide/ml, EzBiolab). After

six hours of stimulation, cells were rested overnight at 4ºC as previously described (34).

The next day, PBMCs were washed with PBS 1X and stained for 25 minutes with the

Live/Dead probe (APC-Cy7, Thermo Fisher Scientific) at RT to discriminate dead cells.

Cells were washed with PBS 1X and surface stained with antibodies for 25 minutes at

RT. We used CD3 (A700, clone UCHT1, BD), CD4 (APC-Cy7, clone SK3, BD), CD8

(V500, clone RPA-T8, BD), CD45RA (BV786, clone HI100, BD), CCR7 (PE-CF594,

clone 150503, BD), CD27 (BV605, clone L128, BD), TIGIT (PE-Cy7, clone MBSA43,

Labclinics SA), PD-1 (BV421, clone EH12.1, BD), LAG-3 (PE, clone T47-530, BD),

TIM-3 (A647, clone 7D3, BD) and CD39 (FITC, clone TU66, BD) antibodies.

Afterwards, cells were washed twice in PBS 1X, fixed, and permeabilized with Fix/Perm

kit (A and B solutions, Thermo Fisher Scientific) for intracellular cytokine staining with

anti-human antibodies of IFNγ (BV711, clone B27, BD) and IL-2 (BV650, clone MQ1-

17H12, BD). Finally, stained cells were washed twice with PBS 1X and fixed in

formaldehyde 1%.

TIGIT, TIM-3 and TIGIT+TIM-3 short-term checkpoint blockade

We selected cryopreserved PBMCs from S1 (n=10) and S2 (n=10). Samples were

previously characterized by the expression of TIGIT and TIM-3 on total CD8+ T cells.

PBMCs were thawed and rested for four hours at 37ºC in a 5% CO_2 incubator. Next, cells

were incubated under RPMI complemented medium 10% FBS with 1μl/ml of anti-

CD28/CD49d and 1μl/ml of Monensin A overnight at 37ºC in a 5% CO_2. PBMCs are

divided in the following conditions; 1) unstimulated, 2) SEB (1μg/ml, Sigma-Aldrich)

and 3) HIV-1-Gag peptide pool (2μg/peptide/ml) in the absence or presence of αTIGIT

and/or αTIM-3, and its respective isotype antibodies. For the single blockade of TIGIT

(αTIGIT), we included Ultra-LEAF™ purified anti-human TIGIT antibody (10μg/ml,

clone A15153G, Biolegend) or its control isotype Ultra-LEAF™ purified mouse IgG2a

antibody (10μg/ml, MOPC-173, Biolegend). For single TIM-3 blockade (αTIM-3), we

used Ultra-LEAF™ purified anti-human TIM-3 antibody (10μg/ml, clone F38-2E2,

Biolegend) or its respective isotype Ultra-LEAF™ purified mouse IgG1 antibody

(10μg/ml, MOPC-21, Biolegend). Finally, we included αTIGIT+αTIM-3 or their

respective IgG2+IgG1 isotypes for a combinational blockade. The next day, PBMCs were

surface and intracellularly stained with the panel of antibodies and the methodology

described in the section above.

Supervised immunophenotype data analysis

Stained PBMCs were acquired on an LSR Fortessa cytometer using FACSDiVa software

(BD). Approximately 1,000,000 events of PBMCs were recorded per specimen. Antibody

capture beads (BD) were used for single-stain compensation controls. Flow cytometry

data were analyzed with FlowJo software v10.6.1, and fluorescence minus one (FMO)

was used to set manual gates. We analyzed CD8+ T cells by excluding dump and CD4+

T cells. We excluded patients with <20% viability in lymphocytes and total CD8+ T cells

(**Source data Figure 1A**). As previously described, we measured by supervised and

classical analyses the IRs expression in CD8+ T-cell subsets, including Naïve, central

memory, transitional memory, effector memory and effector CD8+ T cells (20,34). We

performed two technical replicates for SEB-activated and HIV-1-specific CD8+ T-cell

cytokine production. We considered the cytokine response positive after background

subtraction (mean of two technical replicates) used as the cut-off value. For each

independent sample, we recorded a median of 1,000 events and 50 events positive for

cytokines for total and CD8+ T cell subsets, respectively.

Unsupervised immunophenotype data analysis

The phenotypic and functional characterization of cellular populations was analyzed by

using t-Distributed Stochastic Neighbor Embedding (t-SNE) (70) and net-SNE (71)

dimensionality reduction algorithms to visualize single-cell distributions in two-

dimensional maps. Briefly, cell intensity was z-normalized, and a randomly selected

subset of cells, at least 1,000 cells per sample, was passed through the t-SNE algorithm.

The resulting t-SNE dimension was then used to predict the position of all remaining

CD8+ T cells acquired per sample from each group using the net-SNE algorithm based

on neural networks. For functional analysis, we selected polyclonally activated and HIV-

1-specific CD8+ T cells producing at least one cytokine CD107a, IFNγ, or IL-2 under

SEB or HIV-1 conditions, respectively. In parallel, we discovered cell communities using

the Phenograph clustering technique. It operates by computing the Jaccard coefficient

between nearest neighbours, which was set to 30 in all executions, and then locating cell

communities (or clusters) using the Louvain method. The method creates a network

indicating phenotypic similarities between cells. The netSNE maps included

representations of the identified cell communities, and additionally, we built a heatmap

with the clusters in the columns and the markers of interest in the rows to better

comprehend the phenotypical interpretation of each cluster. The colour scale displays

each marker's median intensity on a biexponential scale. We calculated quantitative

assessments of cellular clusters in the percentage of cells for each sample to analyze and

compare the distribution between HC, Ei, S1, and S2 groups, similar to the classical flow

cytometry analysis.

Statistics

Bivariate analysis was conducted using nonparametric methods as follows: Mann-

Whitney U test for independent median comparison between groups, Wilcoxon signed-

rank test for paired median changes over time, permutation test for composition

distribution between groups, Kruskal-Wallis test for comparison between more than two

groups, and spearman linear correlation coefficient to study the association between

continuous variables. Holm's method was used when appropriate to adjust statistical tests

for multiple comparisons with a significance level of 0.05. All statistics and single-cell

analyses were conducted using the R statistical package (72). The selection of clusters

was performed through the significant differences obtained by inter- and intragroup

comparisons between groups. Moreover, pattern distribution and graphical

representations of all possible Boolean combinations for IRs co-expression and functional

markers were conducted using the data analysis program Pestle v2.0 and SPICE v6.0

software (73). Graph plotting was performed by GraphPad Prism v8.0 software and R

packages. The data represents the mean of two technical replicates.

Study Approval

The study was conducted according to the principles of the Declaration of Helsinki

(Fortaleza, 2013). The Hospital Germans Trias i Pujol Ethics Committee approved all

experimental protocols (PI14-084). For the study, subjects provided written informed

consent for research purposes of biological samples taken from them.

Author contributions

OB-L and JGP conceptualized and designed the experiments. OB-L, EJ-M, and RP

developed immunophenotypic analyses. OB-L, MAM, and JC developed short-term

antibody blockade experiments. OB-L, DO, MAM, AP, AT, AS, R-PS, and JGP

performed supervised and unsupervised analyses. JD, JRS, BC, and JGP recruited the

study participants. OB-L and JGP wrote the manuscript. All authors revised it critically

for important intellectual content and have approved the final version submitted for

publication.

Acknowledgements

We thank the Flow Cytometry Core Facility from the Germans Trias i Pujol Research

Institute (IGTP). This research was supported by National Health Institute Carlos III grant

PI17/00164. Oscar Blanch-Lombarte was funded by an AGAUR-FI_B 00582 PhD

fellowship from Catalan Government and the European Social Fund. Redes Temáticas

de Investigación en SIDA (ISCIII RETIC RD16/0025/0041) funded Esther Jiménez-

Moyano. This study has received partial funding from Grifols. The funders had no role

in study design, data collection, analysis, or the decision to publish or draft the

manuscript.

References

1. Borrow P, Lewicki H, Hahn BH, Shaw GM, Oldstone MB. Virus-specific CD8+ cytotoxic T-lymphocyte activity associated with control of viremia in primary human immunodeficiency virus type 1 infection. J Virol. 1994;68:6103–10.

2. Collins DR, Gaiha GD, Walker BD. CD8+ T cells in HIV control, cure and prevention. Nat Rev Immunol. 2020;20:471–82.

3. Mcbrien JB, Kumar NA, Silvestri G. Mechanisms of CD8+ T cell-mediated suppression of HIV/SIV replication. Eur J Immunol. 2018;48:898–914.

4. Cartwright EK, Spicer L, Smith SA, Lee D, Fast R, Lawson BO, et al. CD8+ lymphocytes are required to maintain virus suppression in SIV-infected macaques treated with short-term antiretroviral therapy. Immunity. 2016;45:656–68.

5. Goulder PJR, Walker BD. HIV and HLA Class I: an evolving relationship. Immunity. 2012;37:426–40.

6. Sáez-Cirión A, Lacabaratz C, Lambotte O, Versmisse P, Urrutia A, Boufassa F, et al. HIV controllers exhibit potent CD8 T cell capacity to suppress HIV infection ex vivo and peculiar cytotoxic T lymphocyte activation phenotype. Proc Natl Acad Sci U S A. 2007 Apr;104(16):6776–81.

7. Migueles SA, Osborne CM, Royce C, Compton AA, Joshi RP, Weeks KA, et al. Lytic Granule Loading of CD8+ T Cells Is Required for HIV-Infected Cell Elimination Associated with Immune Control. Immunity. 2008;29(6):1009–21.

8. Nguyen S, Deleage C, Darko S, Ransier A, Truong DP, Agarwal D, et al. Elite control of HIV is associated with distinct functional and transcriptional signatures in lymphoid tissue CD8+ T cells. Sci Transl Med. 2019;11:eaax4077.

9. Perdomo-Celis F, Taborda NA, Rugeles MT. CD8+ T-cell response to HIV infection in the era of antiretroviral therapy. Front Immunol. 2019;10:1–19.

10. Perdomo-Celis F, Velilla PA, Taborda NA, Rugeles MT. An altered cytotoxic program of CD8 + T-cells in HIV-infected patients despite HAART-induced viral suppression. PLoS One. 2019;14:1–24.

11. Tavenier J, Langkilde A, Haupt TH, Henriksen JH, Jensen FK, Petersen J, et al. Immunosenescence of the CD8+ T cell compartment is associated with HIV-infection, but only weakly reflects age-related processes of adipose tissue, metabolism, and muscle in antiretroviral therapy-treated HIV-infected patients and controls. BMC Immunol. 2015;16:1–14.

12. Serrano-Villar S, Sainz T, Lee SA, Hunt PW, Sinclair E, Shacklett BL, et al. HIV-Infected Individuals with Low CD4/CD8 Ratio despite Effective Antiretroviral Therapy Exhibit Altered T Cell Subsets, Heightened CD8+ T Cell Activation, and Increased Risk of Non-AIDS Morbidity and Mortality. PLoS Pathog. 2014;10:e1004078.

13. Ruiz A, Blanch-Lombarte O, Jimenez-Moyano E, Ouchi D, Mothe B, Peña R, et al. Antigen production after latency reversal and expression of inhibitory receptors in CD8+ T cells limit the killing of HIV-1 reactivated cells. Front Immunol. 2019;10:1–16.

14. Jiang W, He Y, He W, Wu G, Zhou X, Sheng Q, et al. Exhausted CD8+T Cells in the Tumor Immune Microenvironment: New Pathways to Therapy. Front Immunol. 2021;11:1–11.

15. Hoffmann M, Pantazis N, Martin GE, Hickling S, Hurst J, Meyerowitz J, et al. Exhaustion of Activated CD8 T Cells Predicts Disease Progression in Primary HIV-1 Infection. PLoS Pathog. 2016;12(7):1–19.

16. Sekine T, Perez-Potti A, Nguyen S, Gorin JB, Wu VH, Gostick E, et al. TOX is expressed by exhausted and polyfunctional human effector memory CD8+ T cells. Sci Immunol. 2020;5:1–15.

17. Buggert M, Tauriainen J, Yamamoto T, Frederiksen J, Ivarsson MA, Michaëlsson J, et al. T-
bet and Eomes Are Differentially Linked to the Exhausted Phenotype of CD8+ T Cells in
HIV Infection. PLoS Pathog. 2014;10:1–15.
18. Bengsch B, Johnson AL, Kurachi M, Odorizzi PM, Pauken KE, Attanasio J, et al.
Bioenergetic insufficiencies due to metabolic alterations regulated by PD-1 are an early
driver of CD8+T cell exhaustion. Immunity. 2016;45:358–73.
19. Sen DR, Kaminski J, Barnitz RA, Kurachi M, Gerdemann U, Yates KB, et al. The epigenetic
landscape of T cell exhaustion. Science (80-). 2016;354(6316):1165–9.
20. Breton G, Chomont N, Takata H, Fromentin R, Ahlers J, Filali-Mouhim A, et al.
Programmed Death-1 Is a Marker for Abnormal Distribution of Naive/Memory T Cell
Subsets in HIV-1 Infection. J Immunol. 2013;191:2194–204.
21. Yamamoto T, Price DA, Casazza JP, Ferrari G, Nason M, Chattopadhyay PK, et al. Surface
expression patterns of negative regulatory molecules identify determinants of virus-specific
CD8+ T-cell exhaustion in HIV infection. Blood. 2011;117:4805–15.
22. Cockerham LR, Jain V, Sinclair E, Glidden D V., Hartogenesis W, Hatano H, et al.
Programmed death-1 expression on CD4+ and CD8+ T cells in treated and untreated HIV
disease. Aids. 2014;28(12):1749–58.
23. Rutishauser RL, Hartogensis W, Deguit CD, Krone M, Hoh R, Hecht FM, et al. Early and
Delayed Antiretroviral Therapy Results in Comparable Reductions in CD8+ T Cell
Exhaustion Marker Expression. AIDS Res Hum Retroviruses. 2017;33:658–67.
24. Tauriainen J, Scharf L, Frederiksen J, Naji A, Ljunggren HG, Sönnerborg A, et al. Perturbed
CD8+ T cell TIGIT/CD226/PVR axis despite early initiation of antiretroviral treatment in
HIV infected individuals. Sci Rep. 2017;7:1–14.
25. Chew GM, Fujita T, Webb GM, Burwitz BJ, Wu HL, Reed JS, et al. TIGIT Marks Exhausted
T Cells, Correlates with Disease Progression, and Serves as a Target for Immune Restoration
in HIV and SIV Infection. PLoS Pathog. 2016;12:1–28.
26. Trautmann L, Janbazian L, Chomont N, Said EA, Gimmig S, Bessette B, et al. Upregulation
of PD-1 expression on HIV-specific CD8+ T cells leads to reversible immune dysfunction.
Nat Med. 2006;12:1198–202.
27. Day CL, Kaufmann DE, Kiepiela P, Brown JA, Moodley ES, Reddy S, et al. PD-1 expression
on HIV-specific T cells is associated with T-cell exhaustion and disease progression. Nature.
2006;443:350–4.
28. Graydon CG, Balasko AL, Fowke KR. Roles, function and relevance of LAG3 in HIV
infection. PLoS Pathog. 2019;15(1):1–14.
29. Gupta PK, Godec J, Wolski D, Adland E, Yates K, Pauken KE, et al. CD39 Expression
Identifies Terminally Exhausted CD8+ T Cells. PLoS Pathog. 2015;11:1–22.
30. Jones RB, Ndhlovu LC, Barbour JD, Sheth PM, Jha AR, Long BR, et al. Tim-3 expression
defines a novel population of dysfunctional T cells with highly elevated frequencies in
progressive HIV-1 infection. J Exp Med. 2008;205:2763–79.
31. Teigler JE, Zalinskyy G, Eller MA, Bolton DL, Marovich M, Gordon AD, et al. Differential
Inhibitory Receptor Expression on T Cells Delineates Functional Capacities in Chronic Viral
Infection. J Virol. 2017;91:e01263-17.
32. Tian X, Zhang A, Qiu C, Wang W, Yang Y, Qiu C, et al. The Upregulation of LAG-3 on T
Cells Defines a Subpopulation with Functional Exhaustion and Correlates with Disease
Progression in HIV-Infected Subjects. J Immunol. 2015;194:3873–82.
33. Chen H, Moussa M, Catalfamo M. The Role of Immunomodulatory Receptors in the
Pathogenesis of HIV Infection: A Therapeutic Opportunity for HIV Cure? Front Immunol.
2020;11:1–20.
34. Blanch-Lombarte O, Gálvez C, Revollo B, Jiménez-Moyano E, Llibre JM, Manzano JL, et
al. Enhancement of Antiviral CD8+ T-Cell Responses and Complete Remission of

Metastatic Melanoma in an HIV-1-Infected Subject Treated with Pembrolizumab. J Clin
Med. 2019;8:2089.

35. Sakuishi K, Apetoh L, Sullivan JM, Blazar BR, Kuchroo VK, Anderson AC. Targeting Tim-3 and PD-1 pathways to reverse T cell exhaustion and restore anti-tumor immunity. J Exp Med. 2010;207:2187–94.

36. Harjunpää H, Guillerey C. TIGIT as an emerging immune checkpoint. Clin Exp Immunol. 2020;200(2):108–19.

37. Maruhashi T, Sugiura D, Okazaki IM, Okazaki T. LAG-3: from molecular functions to clinical applications. J Immunother cancer. 2020;8:e001014.

38. Attanasio J, Wherry EJ. HHS Public Access. 2017;44(5):1052–68.

39. Chiu CY, Chang JJ, Dantanarayana AI, Solomon A, Evans VA, Pascoe R, et al. Combination Immune Checkpoint Blockade Enhances IL-2 and CD107a Production from HIV-Specific T Cells Ex Vivo in People Living with HIV on Antiretroviral Therapy. J Immunol. 2022;208:54–62.

40. Vaddepally RK, Kharel P, Pandey R, Garje R, Chandra AB. Review of indications of FDA-approved immune checkpoint inhibitors per NCCN guidelines with the level of evidence. Cancers (Basel). 2020;12:1–19.

41. Twomey JD, Zhang B. Cancer Immunotherapy Update: FDA-Approved Checkpoint Inhibitors and Companion Diagnostics. AAPS J. 2021;23:1–11.

42. Gonzalez-Cao M, Morán T, Dalmau J, Garcia-Corbacho J, Bracht JWP, Bernabe R, et al. Assessment of the Feasibility and Safety of Durvalumab for Treatment of Solid Tumors in Patients with HIV-1 Infection: The Phase 2 DURVAST Study. JAMA Oncol. 2020;6:1063–7.

43. Fromentin R, DaFonseca S, Costiniuk CT, El-Far M, Procopio FA, Hecht FM, et al. PD-1 blockade potentiates HIV latency reversal ex vivo in CD4+ T cells from ART-suppressed individuals. Nat Commun. 2019;10:1–7.

44. Le Garff G, Samrib A, Lambert-Niclot S, Even S, Lavole A, Cadranel J, et al. Transient HIV-specific T cells increase and inflammation in an HIV-infected patient treated with nivolumab. Aids. 2017;31:1047–8.

45. Gay CL, Bosch RJ, Ritz J, Hataye JM, Aga E, Tressler RL, et al. Clinical trial of the anti-PD-L1 antibody BMS-936559 in HIV-1 infected participants on suppressive antiretroviral therapy. J Infect Dis. 2017;215:1725–33.

46. Guihot A, Marcelin AG, Massiani MA, Samri A, Soulié C, Autran B, et al. Drastic decrease of the HIV reservoir in a patient treated with nivolumab for lung cancer. Ann Oncol. 2017;29:517–8.

47. Uldrick TS, Adams S V., Fromentin R, Roche M, Fling SP, Gonçalves PH, et al. Pembrolizumab induces HIV latency reversal in people living with HIV and cancer on antiretroviral therapy. Sci Transl Med. 2022;14.

48. Deeks SG, Archin N, Cannon P, Collins S, Jones RB, de Jong MAWP, et al. Research priorities for an HIV cure: International AIDS Society Global Scientific Strategy 2021. Nat Med. 2021;27:2085–98.

49. Kou Z C, Halloran M, Lee-Parritz, Shen L, Simon M, Sehgal P K, et al. In vivo effects of a bacterial superantigen on macaque TCR repertoires. J Immunol. 1998; 160(10): 5170-80.

50. Chiu Chris Y., Chang Judy J., Dantanarayana Ashanti I., Solomon Ajantha, Evans Vanessa A., Pascoe Rachel, et al. I Combination Immune Checkpoint Blockade Enhances IL-2 and CD107a Production from HIV-Specific T Cells Ex Vivo in People Living with HIV on Antiretroviral Therapy. J Immunol. 2022; 208(1): 54-62.

51. Gaffen S. L. and Liu K. D., Overview of interleukin-2 function, production and clinical applications. Cytokine. 2004; 28:109-123.

52. Akdis Mübeccel, Aab Alar, Altunbulakli Can, Azkur Kursat, Costa Rita A, Crameri Reto et al., Interleukins (from IL-1 to IL-38), interferons, transforming growth factor b, and TNF-a:

Receptors, functions, and roles in diseases. J. Allergy Clin Immunol. 2016. 138 (4): 984-
1010.

53. Voskoboinik I., Whisstock J. C., and Trapani J. A., Perforin and granzymes: Function,
dysfunction and human pathology. Nat. Rev. Immunol. 2015. 15: 388–400.

54. Aktas Esin, Kucuksezer Umut Can, Bilgic Sema, Erten Gaye, Deniz Gunnur. Relationship
between CD107a expression and cytotoxic activity. Cell Immunol. 2009. 1254 (2): 149-54.

55. Bhat Purnima, Leggatt Graham, Waterhouse Nigel, Frazer Ian H. R Interferon-γ derived
from cytotoxic lymphocytes directly enhances their motility and cytotoxicity. Cell Death
Dis. 2017. 8(6): e2836.

56. Chauvin JM, Pagliano O, Fourcade J, Sun Z, Wang H, Sander C, et al. TIGIT and PD-1
impair tumor antigen-specific CD8+ T cells in melanoma patients. J Clin Invest.
2015;125(5):2046–58.

57. Anderson AC, Joller N, Kuchroo VK. Lag-3, Tim-3, and TIGIT: Co-inhibitory Receptors
with Specialized Functions in Immune Regulation. Immunity. 2016;44:989–1004.

58. Joller N, Kuchroo VK. Tim-3, Lag-3, and TIGIT. Curr Top Microbiol Immunol.
2017;410:127–56.

59. Schildberg FA, Klein SR, Freeman GJ, Sharpe AH. Coinhibitory Pathways in the B7-CD28
Ligand-Receptor Family. Immunity. 2016;44:955–72.

60. Pino M, Pereira Ribeiro S, Pagliuzza A, Ghneim K, Khan A, Ryan E, et al. Increased
homeostatic cytokines and stability of HIV-infected memory CD4 T-cells identify
individuals with suboptimal CD4 T-cell recovery on-ART. Plos Pathogens. 2021;
17(8):e1009825.

61. Noyan K, Nguyen S, Betts MR, Sönnerborg A, Buggert M. Human immunodeficiency virus
type-1 elite controllers maintain low co-expression of inhibitory receptors on CD4+ T cells.
Front Immunol. 2018;9:1–11.

62. Avery L, Filderman J, Szymczak-Workman AL, Kane LP. Tim-3 co-stimulation promotes
short-lived effector T cells, restricts memory precursors, and is dispensable for T cell
exhaustion. PNAS. 2018;115:2455–60.

63. Warren JA, Clutton G and Goonetilleke N. Harnessing CD8+ T Cells Under HIV
Antiretroviral Therapy. Front Immunol. 2019;10:291.

64. Doering TA, Crawford A, Angelosanto JM, Paley MA, Ziegler CG, Wherry EJ. Network
analysis reveals centrally connected genes and pathways involved in CD8+ T cell exhaustion
versus memory. Immunity. 2012;37(6):1130-44.

65. Andrews LP, Yano H, Vignali DAA. Inhibitory receptors and ligands beyond PD-1, PD-L1
and CTLA-4: breakthroughs or backups. Nat Immunol. 2019;20:1425–34.

66. Sabatos-Peyton CA, Nevin J, Brock A, Venable JD, Tan DJ, Kassam N, et al. Blockade of
Tim-3 binding to phosphatidylserine and CEACAM1 is a shared feature of anti-Tim-3
antibodies that have functional efficacy. Oncoimmunology. 2018;7:1–9.

67. Pende D, Castriconi R, Romagnani P, Spaggiari GM, Marcenaro S, Dondero A, et al.
Expression of the DNAM-1 ligands, Nectin-2 (CD112) and poliovirus receptor (CD155), on
dendritic cells: Relevance for natural killer-dendritic cell interaction. Blood. 2006;107:2030–
6.

68. Kløverpris Henrik K, Payne Rebecca P, Sacha Jonah B, Rasaiyaah Jane T, Chen Fabian,
Takiguchi Masafumi, et al. Early antigen presentation of protective HIV-1 KF11Gag and
KK10Gag epitopes from incoming viral particles facilitates rapid recognition of infected
cells by specific CD8+ T cells. J Virol. 2013. 87(5): 2628-38.

69. Stevenson Eva M., Ward Adam R., Truong Ronald, Thomas Allison S., Huang Szu-Han, et
al. HIV-specific T cell responses reflect substantive in vivo interactions with antigen despite
long-term therapy. JCI insight. 2021;6(3):e142640.

70. Maaten L van der. Visualizing Data using t-SNE. J Mach Learn Res. 2008;219:187–202.

71. Hyunghoon Cho BB and JP. Generalizable and Scalable Visualization of Single-Cell Data
Using Neural Networks. Cell Syst. 2018;7(2):185–91.

72. Core Team R development. R: a Language and Environment for Statistical Computing. Vol.
2, http://www.R-project.org/. 2008. 1–2630 p.

73. Roederer M, Nozzi JL, Nason MC. SPICE: Exploration and analysis of post-cytometric
complex multivariate datasets. Cytom Part A. 2011;79 A:167–74.

Figure Legends

Figure 1. Patterns of IRs co-expression and correlations with CD4+ T -cell counts in PLWH. (A) Overview of study design and study groups, healthy controls (HC), PLWH in early HIV-1 infection (Ei), and PLWH on fully suppressive ART (S) in S1 and S2 time points. **(B)** The expression of IRs summarized in the pie chart is none, one, two, or more than three IRs expressed in CD8+ T-cell subsets. For statistical analysis, we used permutation tests using SPICE software. **(C)** Scatter plots showing the median and interquartile ranges of IR combinations in CD8+ T-cell subsets. **(D)** Scatter plots of the frequencies of single TIGIT$^+$ expression in CM and TIGIT$^+$TIM-3$^+$ expression in EM and effector EFF CD8+ T cells. **(E-G)** Correlations between CD4+ T-cell counts as a function of TIGIT$^+$, TIGIT$^+$TIM-3$^+$, and combinations of IRs from total CD8+ T cells and subsets in Ei **(E)**, S1 **(F)**, and S2 **(G)**. The data in B to D represent the mean of two technical replicates. We used the Mann-Whitney U test for intergroup comparison (HC, Ei, S1, and S2) and the signed-rank test for intragroup comparison (S1 and S2). Holm's method was used to adjust statistical tests for multiple comparisons. All possible correlations of the 32 Boolean IRs combinations are not shown.   P-values: *<0.05, **<0.005 and ***<0.0005.

Figure 2. Unsupervised net-SNE analyses of total CD8+ T cells. (A) Gating strategy for selecting total CD8+ T cells (top), net-SNE plots of HC, Ei, S1, S2 and all merge groups. **(B)** Representative net-SNE visualization of surface markers. The colour gradient displays the relative marker expression. **(C)** Unsupervised KNN algorithm of 38 clusters coloured according to the legend. Only clusters with statistical differences are represented in the legend. **(D)** Heatmap of the median biexponential-transformed marker expression normalized to a -3 to 3 range of respective markers. Asterisks represent the clusters with

statistical differences. **(E-F)** Scatter plots of intergroup (HC, Ei, S1 and S2) and

intragroup (S1 and S2) cluster comparisons. Data represent the median and interquartile

ranges of cluster cell frequency. We used the Mann-Whitney U test for intergroup

analyses and the signed-rank test for intragroup analyses. Holm's method was used to

adjust statistical tests for multiple comparisons. P-values: *<0.05, **<0.005 and

***<0.0005.

Figure 3. Unsupervised net-SNE analyses of SEB-activated CD8+ T cells. (A) Gating

representation of CD107a, IFNγ and IL-2 expression in HIV-1-specific CD8+ T cells

(top), net-SNE plots of HC, Ei, S1, S2 and merge groups of SEB-activated CD8+ T cells

(bottom). **(B)** Representative net-SNE visualization of IR expression, lineage, and

functional markers. The colour gradient displays relative marker expression. **(C)**

Unsupervised KNN algorithm for 29 polyclonal clusters colour-coded according to the

legend. Clusters with statistical differences between groups are represented in the legend.

(D) Heatmap of the median biexponential-transformed marker expression normalized to

a -3 to 3 range of respective markers. Asterisks represent the clusters with intergroup

statistical differences. **(E-F)** Scatter plots of intergroup (HC, Ei, S1 and S2) and

intragroup (S1 and S2) cluster comparisons. Data represent the median and interquartile

ranges of cluster cell frequency. We used the Mann-Whitney U test for intergroup

analyses and the signed-rank test for intragroup analyses. Holm's method was used to

adjust statistical tests for multiple comparisons. P-values: *<0.05, **<0.005, ***<0.0005.

Figure 4. Unsupervised and supervised analyses of HIV-1-specific CD8+ T cells. (A)

Gating representation of CD107a, IFNγ and IL-2 expression in HIV-1-specific CD8+ T

cells (top), net-SNE plots of Ei, S1, S2 and merge groups for HIV-1-specific CD8+ T

cells (bottom). **(B)** Representative net-SNE plots for surface and functional markers. The

colour gradient displays relative marker expression. **(C)** Unsupervised KNN algorithm

for 26 HIV-1-specific clusters colour-coded according to the legend. Only clusters with

statistical differences are represented in the legend. **(D)** Heatmap of the median

biexponential-transformed marker expression normalized to a -3 to 3 range of respective

markers. Asterisks represent the clusters with intergroup statistical differences. **(E)**

Scatter plots of intergroup (Ei, S1 and S2) cluster comparisons with significant statistical

differences. Data represent the median and interquartile ranges of cluster cell frequency.

(F) CD107a, IFNγ, and IL-2 frequency of expression in TIGIT+ (upper panel) and

TIGIT+TIM-3+ (bottom panel) HIV-1-specific memory CD8+ T cell subsets. Scatter

plots represent the median and interquartile ranges. **(G)** Polyfunctional analyses of

CD107a IFNγ and IL-2 expression in CM TIGIT HIV-1-specific CD8+ T cells. Scatter

plots represent median and interquartile ranges. We used the Mann-Whitney U test for

intergroup analyses and the signed-rank test for intragroup analyses. Holm's method was

used to adjust statistical tests for multiple comparisons. P-values: *<0.05, ***<0.0005,

and ****<0.0001.

Figure 5. **Effect of TIGIT, TIM-3 and TIGIT+TIM-3 mAb blockade in HIV-1-**

specific CD8+ T cell responses in PLWH on ART. **(A)** Representative flow cytometry

plots gated on CD8+ T-cells, in the absence of HIV-1 Gag stimulation (basal condition)

and presence of HIV-1 Gag stimulation with isotype control, αTIGIT, αTIM-3, and

αTIGIT+αTIM-3 antibodies for CD107a, IFNγ and IL-2 expression. **(B)** Representative

net-SNE plots for HIV-1-specific CD8+ T cells from PLWH concatenated and merged

according to the condition. **(C-E)** Frequency of CD107a, IFNγ, and IL-2 expression in

total and HIV-1-specific CD8+ T cell subsets for the various conditions tested. The

Wilcoxon matched-pairs signed ranked test calculated statistical differences. The data

represent the mean of two technical replicates. P-values: =0.05, *<0.05, **<0.005 and

***<0.0005.

Supplemental Figures

Supplemental Figure 1. Gating strategy for identifying total CD8+ T-cell subsets and IRs expression levels. **(A)** Zebra dot plots showing the gating strategy used for the identification of total CD8+ and cellular using lineage markers (CD45RA, CCR7, and CD27) subsets as follow; naïve (CD45RA+ CCR7+ CD27+), Eff (CD45RA+ CCR7- CD27-), CM (CD45RA- CCR7+ CD27+), TM (CD45RA- CCR7- CD27+) and EM (CD45RA-, CCR7-, and CD27-). **(B)** Zebra dot plots of FMO and IRs staining demonstrating TIGIT, PD-1, LAG-3, TIM-3, and CD39 expression in cryopreserved PBMCs.

Supplemental Figure 2. Expression profile of IRs in total and CD8+ T cell subsets across study groups. **(A)** Frequency of CD8+ T cells and IRs expression in total CD8+ T cells across study groups (HC, Ei, S1 and S2) **(B)** Frequency of IRs expression in CD8+ T-cell subsets. Scatter plots represent median and interquartile ranges. **(C)** Co-expression profile of IRs across CD8+ T cell subset. Pie charts represent the frequency for the 32 possible combinations for TIGIT, PD-1, LAG-3, TIM-3, and CD39 expression. The pie charts represent the frequency of expression according to the colour code, and the arcs indicate the frequency of each IR. We used permutation tests of SPICE software for statistical analysis of IRs co-expression data. We used the Mann-Whitney U test for intergroup analyses and the signed-rank test for intragroup analyses. Holm's method was used to adjust statistical tests for multiple comparisons. P-values: *<0.05, **<0.005, ***<0.0005, ****<0.00001.

Supplemental Figure 3. Supervised analyses of SEB-activated CD8+ T cells. **(A)** The total frequency of CD107a, IFNγ and IL-2 SEB-activated CD8+ T cells in HC, Ei, S1 and

S2. Data represent median and interquartile ranges. (**B**) Table of median frequencies of

CD107a, IFNγ and IL-2 SEB-activated CD8+ T-cell. (**C**) Frequency of IL-2 SEB-

activated CD8+ T cell subsets in S1 and S2. We used the Mann-Whitney U test for

independent median intergroup comparison and the signed-rank test for paired median

changes between S1 and S2. The Holm's method was used to adjust statistical tests for

multiple comparisons. P-values: *<0.05, **<0.005, ***<0.0005.

Supplemental Figure 4. Supervised analyses of HIV-1-specific CD8+ T cell responses

(**A**) Frequency of CD107a, IFNγ and IL-2 HIV-1-specific CD8+ T cells in HC, Ei, S1

and S2. Data represent median and interquartile ranges. (**B**) Table of median frequencies

of CD107a, IFNγ and IL-2 HIV-1-specific CD8+ T cells. We used the Mann-Whitney U

test for intergroup analyses and the signed-rank test for intragroup analyses. Holm's

method was used to adjust statistical tests for multiple comparisons.

Supplemental Table 1. Epidemiological characteristics of the study groups.

Study Groups	Healthy Control HC	Early HIV-1 infection Ei	HIV-1 on ART S		β
Number of subjects	24	24	24		
Gender (M/F) [n (%)]	16/8 (67/33%)	23/1 (96/4%)	18/6 (75/25%)		0.091
Weeks since seroconversion [median (IQR)]	-	1.3 (0.77 - 17.8)	-	-	
Number of samples	24	24	24 (S1)	24 (S2)	
Age at sample (years) [median (IQR)]	43.1 (38.1 - 52.5)	43.3 (41.8 - 46.2)	40.6 (35.1 - 44.7)	50.0 (43.7 - 55.1)	0.0035
CD4+ T-cell counts at sample per mm^3 [median (IQR)]	-	630 (557 - 757.3)	573 (464 - 992)	910 (799.8 - 1111)	0.0001
Years undetectable at sample on ART [median (IQR)]	-	0.0	2.2 (1.8 - 2.8)	10.1 (7.4 - 12.9)	<0.0001
VL log copies/ml at sample [median (IQR)]	-	4.5 (4.3 - 5.1)	undetectable	undetectable	<0.0001

Abbreviations: IQR, interquartile range; P-values were calculated using Kruskal-Wallis test (Dunn´s correction) or X^2-test.

Figure 1.

Figure 2.

Figure 3.

Figure 4.

Figure 5.

Supplemental Figure 1.

A.

B. FMO

Supplemental Figure 3.

A.

B. **Median of cytokine frequencies in SEB-activated CD8+ T cells**

	HC	Ei	S1	S2
CD107a	6.64	11.46	5.75	6.4
IFNγ	4.45	12.26	5.39	5.69
IL-2	3.08	3.56	1.96	3.28

C.

Supplemental Figure 4.

A.

B. **Median of cytokine frequencies in total HIV-1-specific CD8+ T cells**

	Ei	**S1**	**S2**
CD107a	0.20	0.15	0.05
IFNγ	0.19	0.07	0.06
IL-2	0.44	0.30	0.16