Within-species differences in vocal production learning in a songbird are associated with differences in flexible rhythm pattern perception

Andrew A. Rouse[1], Aniruddh D. Patel[1, 2], Samantha Wainapel[3], and Mimi H. Kao[4, 5]

[1]Department of Psychology, Tufts University; [2]Program in Brain, Mind and Consciousness, Canadian Institute for Advanced Research; [3]Tufts University; [4]Department of Biology, Tufts University; [5]Neuroscience Graduate Program, Tufts University.

*Corresponding author: Andrew A. Rouse

Email: andrew.rouse@tufts.edu

Media summary

When we recognize a song that's played faster or slower than usual, we are detecting timing *relations*
between sounds, not just absolute durations. This ability relies on auditory-motor interactions in the brain.
Auditory-motor processing is more developed in animals that learn their vocalizations so we hypothesize
that such animals are advantaged in recognizing a rhythmic pattern at different rates. We test this idea in
a songbird where only males learn to sing, and find males are slightly better than females at rhythmic
pattern recognition. Thus, songbirds are ideal for investigating individual differences in rhythm pattern
perception and underlying brain mechanisms.

Abstract

Humans readily recognize a familiar rhythmic pattern, such as isochrony (equal timing between events)
across a wide range of rates. This ability reflects a facility with perceiving the relative timing of events, not
just absolute interval durations. Several lines of evidence suggest that this ability is supported by precise
temporal predictions that arise from forebrain auditory-motor interactions. We have shown previously that
male zebra finches, which possess specialized auditory-motor networks and communicate with
rhythmically patterned sequences, share our ability to recognize isochrony independent of rate. To test
the hypothesis that flexible rhythm pattern perception is linked to vocal learning, we ask whether female
zebra finches, which do not learn to sing, can also recognize global temporal patterns. We find that non-
singing females can flexibly recognize isochrony but perform slightly worse than males on
average. These findings are consistent with recent work showing that while females have reduced
forebrain song regions, the overall network connectivity of vocal premotor regions is similar to that in
males and supports predictions of upcoming events. Comparative studies of male and female songbirds
thus offer an opportunity to study how individual differences in auditory-motor connectivity influence
perception of relative timing, a hallmark of human music perception.

1. Introduction

The ability to recognize auditory rhythms is critical for many species [1–3], but the underlying neural
mechanisms are only beginning to be understood. One area of progress is recognition of communication
signals based on tempo. For example, female field crickets are attracted by male calling songs composed
of trains of short sound pulses when the tempo is ~ 30 syllables/sec. This selectivity is hard-wired and
mediated by a small network of interneurons that processes instantaneous pulse rate [4]. While this
preference is genetically fixed, in other animals experience can sculpt neural responses to behaviorally
salient call rates. For example, in the mouse auditory cortex, excitatory cells are innately sensitive to the
most common pup distress call rate (~5 syllables/s), but their tuning can broaden to a wider range of rates
following co-housing with pups producing calls across a range of rates [5].

Much less is known about how the brain recognizes rhythmic *patterns* independently of rate. While
humans can encode and remember the rate of auditory sequences [6], we also readily recognize rhythmic
patterns across a broad range of rates [7]. In music, one of the most widespread temporal patterns is
isochrony, or equal timing between events [8], and the ability to detect and predict periodicity is central to
the positive effects of music-based therapies on a variety of neurological disorders, including normalizing
gait in Parkinson's disease [9,10]. The ability to recognize a rhythmic pattern whether it is played fast or
slow is present in infants [11] and is based on recognition of the relative timing of events more than on
their absolute durations. In humans, there is growing evidence that the neural mechanisms underlying
detection of relative timing are distinct from those involved in encoding absolute timing [12–14]. In
addition, neuroimaging studies have shown that both auditory *and* motor regions are active when people
listen to rhythms, even in the absence of overt movement. Responses in several motor regions are
greater when the stimulus has a strong, periodic pulse, or "beat" [15,16] and transient manipulation of
auditory-motor connections using transcranial magnetic stimulation can disrupt beat perception without
affecting single-interval timing [17].

Based on such findings, we and others have suggested that perception of temporal regularity
(independent of rate) depends on the interaction of motor and auditory regions: the motor planning
system uses information from the auditory system to make predictions about the timing of upcoming
events and communicates these predictions back to auditory regions via reciprocal connections [18,19].
Such predictions could support detection of rhythmic patterns independent of tempo because the relative
duration of adjacent intervals remains the same across different rates (e.g., 1:1 for an isochronous
pattern).

Given that vocal learning species often communicate using rhythmically patterned sequences [20,21] and
have evolved specialized motor planning regions that are reciprocally connected to auditory forebrain
regions, we have hypothesized that vocal learners are advantaged in flexible auditory rhythm pattern
perception [22]. Consistent with this hypothesis, zebra finches and starlings (vocal learning songbirds)
trained to discriminate isochronous from arrhythmic sound sequences robustly generalize this ability to
stimuli at novel tempi, including rates distant from the training tempi [22,23]. This ability, also seen in
humans [7], demonstrates a facility with recognizing a rhythm based on global temporal patterns, since
absolute durations of intervals differ markedly at distant tempi. These findings contrast with similar
research conducted with vocal non-learning species. For example, pigeons can learn to discriminate
sound sequences based on tempo but cannot learn to discriminate isochronous from arrhythmic sound
patterns [24]. Rats can be trained to discriminate isochronous from arrhythmic sounds, but when tested at
novel tempi, they show limited generalization, suggesting a strong reliance on absolute timing for rhythm
perception [25].

Here, we further test the hypothesis that differences in vocal learning abilities correlate with differences in
flexible rhythm pattern perception by taking advantage of the sexual dimorphism in zebra finches [26]. In
this species only males learn to imitate song, and the neural circuitry subserving vocal learning is greatly
reduced in females. Thus we predicted that male zebra finches would exhibit faster learning rates for
discriminating isochronous versus arrhythmic stimuli and/or a greater degree of generalization in
recognizing these categories at novel tempi. By using the same apparatus, stimuli, and methods, we can
meaningfully compare the flexibility of male vs. female rhythmic pattern perception in this species.

Several prior findings, however, support the opposite prediction - either no sex differences in rhythmic
pattern perception or better rhythmic pattern perception in female zebra finches. One reason females may
not be disadvantaged in our task is that they analyze male song when choosing a mate, and a previous
meta-analysis found that female zebra finches are faster than males at learning to discriminate spectro-
temporally complex auditory stimuli [27]. In addition, while the volume of vocal motor regions that
subserve song performance is substantially greater in male zebra finches, a recent anatomical study
found male-typical patterns of connectivity in the vocal premotor region in females, including minimal sex
differences in afferent auditory and other inputs [28]. Moreover, a recent study of antiphonal calling
suggests that female zebra finches may perform better than males on rhythmic pattern processing [29]. In
that study, both male and female zebra finches could predict the timing of calls of a rhythmically calling
vocal partner, allowing them to adjust the timing of their own answers to avoid overlap. This ability to
predictively adjust call timing was enhanced in females and was disrupted by lesions of vocal motor
forebrain regions, suggesting that brain regions associated with singing in males may subserve auditory
perception and/or timing abilities in female zebra finches.

2. Methods and analysis

(a) Subjects

Subjects were 24 experimentally naïve female zebra finches from our breeding colony (mean age=72 ± 8
(SD) days post-hatch (dph) at the start of training; range=61-87 dph). Data from 14 age-matched male

zebra finches were collected previously [22]. All procedures were approved by the Tufts University
Institutional Animal Care and Use Committee.

(b) Auditory Stimuli

All stimuli used in this experiment were the same as those described previously [22]. Briefly, we used
sequences of natural sounds from unfamiliar male conspecifics: either an introductory element that is
typically repeated at the start of a song, or a short harmonic stack [30] (see Fig. S1). For each sound, an
isochronous sequence with equal time intervals between event onsets and an arrhythmic sequence with a
unique temporal pattern (but the same mean inter-onset interval (IOI)) were generated at two base tempi:
120 ms and 180 ms IOI (Fig. 1A). These tempi were chosen based on the average syllable rate in zebra
finch song (~7-9 syllables per second, or 111-142 ms IOI [30]). The number of elements, overall duration,
amplitude, and spectral profile were matched across a pair of isochronous and arrhythmic stimuli. For the
arrhythmic stimuli, the minimum gap between consecutive elements was 0 ms, and the maximum
possible IOI was 1.5 times the base IOI. An additional pair of probe stimuli (one isochronous, one
arrhythmic) was generated at a tempo (144 ms IOI) 20% faster/slower than the training stimuli.

For "rule training" (see below), an isochronous/arrhythmic pair was generated with a novel harmonic stack
(sound D, see Fig. S1) at tempi ranging from 75 ms to 275 ms IOI in 1 ms steps (i.e., at 201 rates). Each
arrhythmic stimulus had a temporally unique pattern. For a subset of these stimulus pairs (n=15/201), the
amplitude was inadvertently higher as described previously [22].

(c) Auditory Operant Training Procedure

Training and probe testing used a go/interrupt paradigm as described previously [22]. Briefly, female birds
were mildly water restricted and worked for water rewards, routinely performing ~520 trials/day. Pecking
the trial switch triggered playback of a stimulus: either rewarded (S+, 50% chance) or unrewarded (S-,
50% chance). For rhythm discrimination experiments, the isochronous patterns were rewarded (S+
stimuli). Trial and response switches were activated 500 ms after stimulus onset, and pecking either
switch stopped playback. "Hits" were correct pecks of the response switch during S+ trials. "False alarms"
were pecks of the response switch on S- trials and resulted in lights out (up to 25s) [31]. If neither switch
was pecked within 5 s of trial onset, the trial would end ('no response'). No response to the S- stimulus
was counted as a "correct rejection", and no response to the S+ stimulus was considered a "miss". During
this 5 s window, a bird could also peck the trial switch again to "interrupt" the current trial, which was
counted as a 'miss' or 'correct rejection' depending on whether a S+ or S- stimulus had been presented.
The use of the trial switch to interrupt trials varied widely among birds and was not analyzed further.
Regardless of response, birds had to wait 100 ms after the stimulus stopped playing before a new trial
could be initiated.

Shaping and Performance Criteria. To learn the go/interrupt procedure, birds were first trained to
distinguish between two unfamiliar conspecific songs (~2.4 s long, "shaping" phase), one acting as the S+
(rewarded or 'go') stimulus, and the other as the S- (unrewarded or 'no-go') stimulus. Lights-out
punishment was not implemented until a bird performed ≥ 100 trials. The criterion for advancing to the
next phase was ≥ 60% hits, ≥ 60% correct rejections, and ≥ 75% overall correct for two of three
consecutive days. Five females did not complete the shaping process within 30 days and were removed
from the study.

Training: rhythm discrimination. Once a bird reached criterion performance on the shaping stimuli, she
was trained to discriminate isochronous versus arrhythmic stimuli (n=13 birds). As described previously,
each bird was trained using multiple sound types and multiple stimulus rates. In the first training phase,
each bird learned to discriminate two isochronous stimuli (120 ms and 180 ms IOI) from two arrhythmic

stimuli (matched for each mean IOI). Once performance criterion was reached, the bird was presented
with a new set of stimuli at the same tempi but with a novel sound element (and a novel irregular temporal
pattern at each tempo; see Fig. 1B). One group of females ('ABC'; n=6) was trained to discriminate sound
A stimuli, followed by sound B and then sound C; a second group ('CAB'; n=7) was trained with sound C,
followed by sounds A and B. One female 'ABC' bird was presented with stimuli at 3 additional tempi, but
those trials were not reinforced and were excluded from analysis.

Probe Testing/Generalization. To test whether birds could generalize the isochronous vs. arrhythmic
classification at a novel tempo, females were tested with probe stimuli at 144 ms IOI, 20% slower/faster
than the training stimuli (120 ms and 180 ms IOI). We previously showed that male zebra finches can
discriminate between two stimuli that differ in tempo by 20% [22] and confirmed this in female zebra
finches (see below and Fig. S2B). Probe sounds were introduced after a bird had successfully completed
two phases of training (see Fig. 1B). Prior to probe testing, the reinforcement rate for training stimuli was
reduced to 80% for at least two days. During probe testing, training stimuli (90% of trials) and probe
stimuli (10% of trials) were interleaved randomly. Probe trials and 10% of the interleaved training stimuli
were not reinforced or punished[22,32].

Rule Training. Following probe testing with all three sound types, subjects (n=5 birds) were presented
with a new set of isochronous and arrhythmic stimuli using a novel sound (sound D). In this phase,
stimulus tempi included every integer rate from 75 ms to 275 ms IOI (201 total rates). Arrhythmic stimuli
were again generated independently so that each rate had a novel, random irregular pattern, with the
same mean IOI as its corresponding isochronous pattern. Trials were randomly drawn with replacement
from this set of 402 stimuli, and all responses were rewarded or punished as during training. This large
stimulus set made it unlikely that subjects could memorize individual temporal patterns.

Discrimination of other acoustic features. To determine whether the 20% difference between the training
and probe stimuli could be detected by female zebra finches, we tested a separate cohort of females on
tempo discrimination (n=6; 71 ± 9.1 (SD) dph at start of shaping). Following shaping, these birds were
trained to discriminate isochronous sequences of sound A based on rate: 120 ms vs. 144 ms IOI. The
rewarded (S+) stimulus was 120 ms IOI for four females and 144 ms IOI for two females. These females
were also tested on their ability to discriminate spectral features using frequency-shifted isochronous
sequences of sound A: one shifted 3 semitones up, and the other shifted 6 semitones up ("Change pitch"
in Audacity v 2.1.2). Four females were tested with the 6-semitone shift as the S+ stimulus, and two
females were tested with the 3-semitone shift as the S+ stimulus.

(d) Data Analysis

All statistical tests, except for the binomial tests for training and generalization, were performed in R (v.
3.6.2) within RStudio (v. 1.2.5033). Binomial logistic regressions were performed with lme4 (glmer)
statistical package for R.

Training and generalization testing. To quantify performance the proportion of correct responses ((Hits +
Correct Rejections) / Total number of trials) was computed for each stimulus pair (isochronous and
arrhythmic patterns of a given sound at a particular tempo). For training phases, proportion correct was
always computed based on the last 500 trials. For probe testing, performance was computed for the 80
probe trials. The proportion of correct responses was compared to chance performance ($p=0.5$) with a
binomial test using $α=0.05/2$ in the training conditions (2 tempi), $α=0.05/3$ for probe trials (3 tempi). A
linear least-squares regression was used to examine the correlation between each subject's average
performance on interleaved training stimuli and probe stimuli.

Performance of females (n=7) during probe trials was compared to that of male zebra finches (n=7)
collected in a prior study [22]. Performance across all probe trials (n=240 probe trials, 80 probe trials per
song element x 3 song elements) was analyzed with a binomial logistic regression using a generalized

linear mixed model with sex, probe phase number, and sex-by-probe phase number interaction as fixed
effects and subject as a random effect.

Rule training. As described previously [22], the first 1000 trials were analyzed for each bird (n=5) to
minimize any potential effect of memorization. Trials were binned in 10 ms increments, and the number of
correct responses was analyzed with a binomial logistic regression using a generalized linear mixed
model (tempo bin as a fixed effect and subject as a random effect). Performance in each bin was
compared to performance in the 75-85 ms IOI bin, where performance fell to chance since the degree of
temporal variation in inter-element intervals was severely limited by the duration of the sound element. To
identify possible sex differences, we performed an additional analysis using data from a range of tempi
(95-215 ms) in which both males and females performed significantly above chance (p<0.005; binomial
test). The number of correct and incorrect trials were calculated per sex and analyzed for significant
group differences with a χ2 test.

Discrimination of Other Acoustic Features. Performance on tempo or frequency discrimination was
analyzed in the same manner as for rhythm discrimination: the proportion of correct responses in the last
500 trials was compared to chance performance using a binomial test with α=0.05.

Reaction Time. For the last 500 trials in a training phase, the time between trial initiation and response
selection (when the bird pecked a switch) was computed and averaged. For the rhythm discrimination
birds, only the first training phase (prior to probe testing) was used. Mean reaction time values were
compared between 14 birds that completed all rhythmic training phases, 9 birds that failed to complete
rhythm discrimination training, and 10 birds that learned the frequency discrimination (data from 6
females in this study and from 4 males collected as part of [22]) using a two-way ANOVA (sex, group, and
sex-by-group interaction) followed by a Tukey multiple comparisons post-hoc test. Data met criteria for
homoscedasticity and normal error distribution.

3. Results

(a) Rhythmic pattern training and generalization testing in female zebra finches

To test the ability of female songbirds to recognize a rhythmic pattern based on the relative timing of
events, 13 female zebra finches were first trained to discriminate isochronous from arrhythmic sequences
using a go/no-go paradigm with three training phases (Figs. 1A–B & S1; age: 61–87 dph on the first day
of training). Seven out of 13 females learned to discriminate these sequences in all three phases, each of
which used a different zebra finch song element. The number of trials for these females to reach criteria
for rhythmic discrimination (see Materials and Methods) was comparable to that of male zebra finches
tested previously (Fig. 1C; data from male birds in [22]). Fig. 2A shows the time course of learning for a
representative female zebra finch (*y7o97*), which gradually learned to withhold her response to the
arrhythmic stimulus. Across the seven successful females, rhythm discrimination performance was ~81%
accurate at the end of the first training phase (Fig. 2B, left column), and a comparable accuracy level was
attained in each training phase (median = 80% proportion correct; see Fig. S2A for learning curves for
each bird on the rhythm discrimination task).

After successful completion of two phases of rhythm discrimination, females were then asked whether
they could generalize the discrimination of isochronous versus arrhythmic stimuli at a novel tempo (Fig.
1B, 'probe testing'). Fig. 2B shows the performance on randomly interleaved training stimuli (*dark gray
bars*; 90% of the trials*) and 80 probe stimuli at a novel tempo (*light gray bars;* 10% of the trials) for the
seven females that successfully completed all three phases of rhythm discrimination training.
Performance on probe stimuli was significantly above chance for 17 of 21 probe tests (n=3 probe
tests/bird x 7 birds, *p*<0.0167, binomial test with Bonferroni correction), indicating that female zebra

finches robustly generalized the discrimination of isochronous vs. arrhythmic stimuli to a novel tempo
distant from the training tempi (Fig. 2B).

(b) Sex differences in rhythm pattern perception

To test the hypothesis that the ability to flexibly perceive rhythmic patterns is linked to vocal learning, we
directly compared performance of female zebra finches, which do not learn to imitate song, with that of
vocal learning males (Fig. 3). Across all generalization tests, discrimination of isochronous versus
arrhythmic stimuli at a novel tempo (probe stimuli) was significantly lower for female zebra finches
compared to males (n=7 males and 7 females; 240 probe trials per bird; mixed-effects logistic regression,
$p = 0.033$). The observed sex difference in the ability to generalize rhythm discrimination did not reflect
differences in motivation to perform the task. On average, female zebra finches performed ~520
trials/day, compared to ~530 trials/day for males in our previous work. Across all subjects (n=14),
performance on interleaved training stimuli (learned discrimination) was positively correlated with
performance on probe stimuli (measure of generalization; Fig. S3; $p = .0375$).

As a second test of the ability to recognize temporal regularity independent of tempo, five females that
completed rhythm discrimination training were tested with sequences of a new sound element (sound D,
see Figs. S1-S2) across a wide range of tempi (75-275 ms IOI; "rule training"). For each trial, one of 402
stimuli was played (with equal odds of an isochronous or arrhythmic sequence), reducing the likelihood
that correct discrimination was based on memorization. Fig. 4 shows the average performance of all birds
over the first 1000 trials, broken into 10 ms bins (~50 trials per bin). For female birds, performance was
best between 95 and 215 ms IOI (tempi ~20% slower to ~25% faster than the original training range:
darker bars with >=2 black asterisks, $p<0.005$; mixed effect logistic regression). Performance fell to
chance at faster tempi (75-85 ms IOI), when temporal variability in IOIs was limited by the length of the
sound element, and at slower tempi (225-275 ms IOI). To directly compare performance to that of male
zebra finches, the number of correct and incorrect trials within the 95-215 ms IOI tempo range (where
both sexes performed well above chance) were grouped by sex. In this range, female performance was
significantly worse than male performance ($\chi^2(1) = 13.919$, $p<0.001$).

(c) Reaction times for discrimination of acoustic features.

Across female birds who succeeded in rhythm discrimination training (n=7), the average reaction time at
the end of the first training phase was ~1.70 s after stimulus onset, or ~75% of the duration of the
stimulus train (Fig. 5). Data from previously tested males that succeeded in rhythm discrimination training
(n=7) shows a similar average reaction time (1.90 s), indicating that both males and females heard 10-15
intervals on average before responding (overall average reaction time of ~1.80 s). In contrast, the
average reaction time for the females that did not reach the criterion for rhythm discrimination during
training (n=6) was ~1.30 s, ~50% of the duration of each stimulus, similar to the average failed male
(n=3) reaction time of ~1.18 s.

Shorter reaction times, however, do not necessarily indicate poor perceptual discrimination. In a separate
cohort of birds that learned to discriminate isochronous sequences that differed in frequency by 3
semitones (i.e., a quarter of an octave; n = 6 females; average proportion correct responses in last 500
trials = 91%; see Fig. S2B and n = 4 males from [22]), the average reaction times for successful
discrimination was ~1.24 s. This suggests that unlike rhythm discrimination, discrimination based on
spectral features does not require hearing long stretches of the sequence.

Comparison of reaction times between successful rhythm discrimination birds, unsuccessful rhythm
discrimination birds, and successful frequency discrimination birds showed a significant effect of group
(Fig. 5; two-away ANOVA, F(2,27), $p = 0.001$). Tukey post-hoc tests showed that the reaction times of the
successful rhythm birds were significantly higher (slower) than both the unsuccessful rhythm birds and the
successful frequency birds. There was no effect of sex or a sex-by-group interaction. Taken together,

these results suggest that success at rhythm discrimination may be related to how long birds listened to
the stimuli before responding. Indeed, even among the birds that successfully generalized the
discrimination of temporal regularity across a wide range of tempi, reaction time was longer for the tempi
where the discrimination was more difficult (i.e., when the proportion of correct responses was closer to
chance performance: 75-85 ms and >235 ms; Fig. S4).

4. Discussion

To test the hypothesis that differences in the capacity for flexible rhythm pattern perception correlate with
differences in vocal learning, we investigated the ability of a sexually dimorphic songbird to recognize a
fundamental rhythmic pattern common in music and vocal communication – isochrony [8,20,21]. In zebra
finches and many other songbirds, only the males learn to sing and possess pronounced forebrain motor
regions for vocal production learning [26]. Previously we showed that male zebra finches can readily
discriminate isochronous versus arrhythmic patterns independent of absolute time intervals [22], and we
predicted that female zebra finches would perform less well when tested with the same rhythm
discrimination and generalization tasks. Using a sequential training paradigm with multiple sound types
and tempi, we found that about half of females tested (n=7/13) could learn to differentiate isochronous
from arrhythmic patterns, and the learning rates for these females were similar to those of males (Fig.
1C). Females that completed rhythm discrimination training readily generalized the discrimination to
stimuli at novel tempi (Fig. 2B & 4). Correct discrimination of isochronous vs. arrhythmic stimuli was
significantly above chance at tempi ranging from 20% slower to ~25% faster than the original training
range, indicating that female zebra finches, like males, can recognize a rhythmic pattern based on the
relative timing of events rather than on absolute durations. Notably, in both sexes, birds that succeeded in
learning to recognize isochrony listened for longer before responding compared to birds who failed (Fig.
5, left two bars), consistent with attention to global temporal patterns.

While female zebra finches can learn to recognize isochrony independent of tempo, we found small but
significant differences in performance between males and females. First, across all three tests of
generalization at a novel tempo, males outperformed females (Fig. 3; average proportion of correct
responses during probe trials: 75% vs. 69%). Second, males consistently slightly outperformed females
when tested with stimuli across a broad range of tempi at which both sexes performed well above chance
(Fig. 4; 95-215 ms IOI; average proportion correct: 76% vs 72%). These differences did not reflect
differences in motivation to perform the task; on average, males and females performed comparable
numbers of trials/day. Finally, the proportion of females that successfully recognized isochrony based on
global temporal patterns (n=7/13) was lower than that of males tested previously (n=7/10), although more
data would be needed to determine if this difference is reliable.

It is important to note that the sex difference we observed, while consistent with our hypothesis, is an
average difference. Modest, but consistent, sex differences are well known in biology (e.g., on average,
men are taller than women, due, in part, to sex-biased gene expression [33]), but these differences
typically pertain to anatomy, not cognition [34]. Prior work on sex differences in the auditory domain has
focused largely on neural mechanisms in the periphery [35–38], although more recent work has
demonstrated hormone-mediated differences in forebrain auditory responses to conspecific songs in birds
[39,40]. Here, we show a small, but consistent sex difference in an auditory cognitive task – flexible
rhythm pattern recognition – in a sexually dimorphic bird.

Yet just as some women are taller than some men, we find that some individual female zebra finches can
outperform individual males in our tasks (e.g., probe tests in Fig. 3). How can this be reconciled with our
hypothesis of a link between the neural circuitry for vocal learning and flexible rhythm pattern perception?
Recent neuroanatomical work found that although female zebra finches possess smaller vocal motor
regions, the overall network connectivity of vocal premotor regions is similar in male and female zebra
finches [28]. Similarities in ascending auditory inputs to premotor regions, in particular, suggest that

auditory processing mechanisms may be conserved across sex. This raises the question of whether
individual variation in auditory-motor circuitry correlates with individual differences in flexibility of rhythmic
pattern perception, irrespective of sex. Indeed, in humans, the strength of cortical auditory-motor
connections predicts individual differences in rhythmic abilities [41,42].

Our finding that female zebra finches perform slightly worse than males in recognizing isochrony
contrasts with a prior finding that female zebra finches outperform males in a task involving temporal
processing of rhythmic signals [29]. In that study, birds (6 males and 6 females) called antiphonally with a
robotic partner that emitted calls at a rate of 1 call/second. Birds quickly learned to adjust the timing of
their calls in order to avoid a jamming signal introduced at a fixed latency after the robot call, with females
showing more pronounced adjustments of call timing than males. While both studies highlight the
importance of prediction for rhythm perception, the underlying mechanisms for predicting upcoming
events may differ between the two studies. In principle, avoidance of the jamming signal could have
resulted from learning a single temporal interval – the time between the robot call and the jamming signal.
In contrast, in our task birds had to learn to recognize the relative timing of successive events and to
respond to the same pattern, even when absolute interval durations changed. Prior work suggests distinct
mechanisms for single-interval versus relative timing [12–14], so a male advantage on a task the depends
on relative timing is not necessarily inconsistent with a female advantage in single interval timing.

Several recent studies have begun to elucidate the neural mechanisms for rhythm perception in
nonhuman animals. For example, Asokan et al. (2021) found that the responses of neurons in the primary
auditory cortex of mice are modulated by the rhythmic structure of a sound sequence. While neurons in
the midbrain and thalamus encode local temporal intervals with a short latency, cortical neurons integrate
inputs over longer a timescale (~1 s), and the timing of cortical spikes differs depending on whether
consecutive sounds are arranged in a repeating rhythmic pattern or are randomly timed. In gerbils,
midbrain neurons have also been shown to exhibit context-dependent responses: on average, neural
responses were greater for noise bursts that occurred on the beat of a complex rhythm compared to
when the same noise bursts occurred off-beat [44]. Similarly, in rhesus monkeys, occasional deviant
sounds in auditory sequences elicit a larger auditory mismatch negativity in electroencephalogram
recordings when those sequences had isochronous vs. arrhythmic event timing [45]. While these studies
found context-dependent modulation of auditory responses, they did not test the ability of the animals to
recognize a learned rhythm independently of tempo. Demonstrating this ability requires behavioral
methods, and an important lesson from prior research is that training methods can strongly influence to
what extent such abilities are revealed (e.g., compare [46] with [22] and [47] with [48]; see also [49]).
More generally, the ability to relate neural activity to perception and behavior is critical for understanding
the contributions of motor regions to detecting temporal periodicity and predicting the timing of upcoming
events, two hallmarks of human rhythm processing that are central to music's positive effect on a variety
of neurological disorders.

Data accessibility. Concatenated trial data for female zebra finches, summarized data files, and female
analysis code are available through Mendeley Data: https://dx.doi.org/10.17632/2r29x6gr7w.1

Published data for male zebra finches as well as stimulus files, circuit diagrams, male analysis code, and
operant chamber setup diagram are available through Mendeley Data:
https://dx.doi.org/10.17632/fw5f2vrf4k.2

The modified Pyoperant code for this experiment is available on GitHub:
https://github.com/arouse01/pyoperant.

Acknowledgments. We thank T. Gardner and his laboratory for equipment and technical assistance with
the operant chambers and T. Gentner for sharing the Pyoperant code. We thank members of the Kao lab
for useful discussions and comments on earlier versions of this manuscript.

Funding. This work was supported by a Tufts University Collaborates grant (M.H.K. and A.D.P.), NIH
Grant R21NS114682 (M.H.K. and A.D.P.), and a Canadian Institute for Advanced Research catalyst
grant (A.D.P.).
378

379
380

References

1.   Clemens J, Schöneich S, Kostarakos K, Hennig RM, Hedwig B. 2021 A small, computationally
flexible network produces the phenotypic diversity of song recognition in crickets. *Elife* **10**, 1–31.
(doi:10.7554/eLife.61475)

2.   Mathevon N, Casey C, Reichmuth C, Charrier I. 2017 Northern Elephant Seals Memorize the
Rhythm and Timbre of Their Rivals' Voices. *Current Biology* **27**, 2352-2356.e2.
(doi:10.1016/j.cub.2017.06.035)

3.   Garcia M, Theunissen F, Sèbe F, Clavel J, Ravignani A, Marin-Cudraz T, Fuchs J, Mathevon N.
2020 Evolution of communication signals and information during species radiation. *Nature
Communications* **11**, 1–15. (doi:10.1038/s41467-020-18772-3)

4.   Schöneich S, Kostarakos K, Hedwig B. 2015 An auditory feature detection circuit for sound pattern
recognition. *Science Advances* **1**. (doi:10.1126/sciadv.1500325)

5.   Schiavo JK, Valtcheva S, Bair-Marshall CJ, Song SC, Martin KA, Froemke RC. 2020 Innate and
plastic mechanisms for maternal behaviour in auditory cortex. *Nature* **587**, 426–431.
(doi:10.1038/s41586-020-2807-6)

6.   Levitin DJ, Cook PR. 1996 Memory for musical tempo: Additional evidence that auditory memory
is absolute. *Perception and Psychophysics* **58**, 927–935. (doi:10.3758/BF03205494)

7.   Espinoza-Monroy M, de Lafuente V. 2021 Discrimination of Regular and Irregular Rhythms
Explained by a Time Difference Accumulation Model. *Neuroscience* **459**, 16–26.
(doi:10.1016/j.neuroscience.2021.01.035)

8.   Savage PE, Brown S, Sakai E, Currie TE. 2015 Statistical universals reveal the structures and
functions of human music. *Proc Natl Acad Sci U S A* **112**, 8987–8992.
(doi:10.1073/pnas.1414495112)

9.   Dalla Bella S, Benoit CE, Farrugia N, Keller PE, Obrig H, Mainka S, Kotz SA. 2017 Gait
improvement via rhythmic stimulation in Parkinson's disease is linked to rhythmic skills. *Scientific
Reports* **7**, 1–11. (doi:10.1038/srep42005)

10.   Krotinger A, Loui P. 2021 Rhythm and groove as cognitive mechanisms of dance intervention in
Parkinson's disease. *PLoS ONE* **16**, 1–20. (doi:10.1371/journal.pone.0249933)

11.   Trehub SE, Thorpe LA. 1989 Infants' perception of rhythm: categorization of auditory sequences
by temporal structure. *Canadian Journal of Psychology* **43**, 217–229. (doi:10.1037/h0084223)

12.   Grube M, Cooper FE, Chinnery PF, Griffiths TD. 2010 Dissociation of duration-based and beat-
based auditory timing in cerebellar degeneration. *Proceedings of the National Academy of
Sciences* **107**, 11597–11601. (doi:10.1073/pnas.0910473107)

13.   Teki S, Grube M, Kumar S, Griffiths TD. 2011 Distinct Neural Substrates of Duration-Based and
Beat-Based Auditory Timing. *Journal of Neuroscience* **31**, 3805–3812.
(doi:10.1523/JNEUROSCI.5561-10.2011)

14.   Breska A, Ivry RB. 2018 Double dissociation of single-interval and rhythmic temporal prediction in
cerebellar degeneration and Parkinson's disease. *Proc Natl Acad Sci U S A* **115**, 12283–12288.
(doi:10.1073/pnas.1810596115)

15.   Grahn JA, Brett M. 2007 Rhythm and beat perception in motor areas of the brain. *Journal of
Cognitive Neuroscience* **19**, 893–906. (doi:10.1162/jocn.2007.19.5.893)

16. Chen JL, Penhune VB, Zatorre RJ. 2008 Listening to musical rhythms recruits motor regions of the
brain. *Cerebral Cortex* **18**, 2844–2854. (doi:10.1093/cercor/bhn042)

17. Ross JM, Iversen JR, Balasubramaniam R. 2018 The Role of Posterior Parietal Cortex in Beat-
based Timing Perception: A Continuous Theta Burst Stimulation Study. *Journal of Cognitive*
Neuroscience **30**, 634–643. (doi:10.1162/jocn_a_01237)

18. Cannon JJ, Patel AD. In press. How beat perception coopts motor neurophysiology. *Trends in*
Cognitive Sciences (doi:10.1101/805838)

19. Patel AD, Iversen JR. 2014 The evolutionary neuroscience of musical beat perception: the Action
Simulation for Auditory Prediction (ASAP) hypothesis. *Frontiers in Systems Neuroscience* **8**, 57.
(doi:10.3389/fnsys.2014.00057)

20. Roeske TC, Tchernichovski O, Poeppel D, Jacoby N. 2020 Categorical Rhythms Are Shared
between Songbirds and Humans. *Current Biology* **30**, 3544-3555.e6.
(doi:10.1016/j.cub.2020.06.072)

21. Norton P, Scharff C. 2016 "Bird song metronomics": Isochronous organization of zebra finch song
rhythm. *Frontiers in Neuroscience* **10**, 97–99. (doi:10.3389/fnins.2016.00309)

22. Rouse AA, Patel AD, Kao MH. 2021 Vocal learning and flexible rhythm pattern perception are
linked: Evidence from songbirds. *Proceedings of the National Academy of Sciences* **118**,
e2026130118. (doi:10.1073/pnas.2026130118)

23. Hulse SH, Humpal J, Cynx J. 1984 Discrimination and Generalization of Rhythmic and Arrhythmic
Sound Patterns by European Starlings (Sturnus vulgaris). *Music Perception* **1**, 442–464.
(doi:10.2307/40285272)

24. Hagmann CE, Cook RG. 2010 Testing meter, rhythm, and tempo discriminations in pigeons.
Behavioural Processes **85**, 99–110. (doi:10.1016/j.beproc.2010.06.015)

25. Celma-Miralles A, Toro JM. 2020 Discrimination of temporal regularity in rats (Rattus norvegicus)
and humans (Homo sapiens). *Journal of Comparative Psychology* (doi:10.1037/com0000202)

26. Nottebohm F, Arnold AP. 1976 Sexual Dimorphism in Vocal Control Areas of the Songbird Brain.
Science (1979) **194**, 211–213. (doi:10.1126/science.959852)

27. Kriengwatana B, Spierings MJ, ten Cate C. 2016 Auditory discrimination learning in zebra finches:
effects of sex, early life conditions and stimulus characteristics. *Animal Behaviour* **116**, 99–112.
(doi:10.1016/j.anbehav.2016.03.028)

28. Shaughnessy DW, Hyson RL, Bertram R, Wu W, Johnson F. 2019 Female zebra finches do not
sing yet share neural pathways necessary for singing in males. *Journal of Comparative Neurology*
527, 843–855. (doi:10.1002/cne.24569)

29. Benichov JI, Benezra SE, Vallentin D, Globerson E, Long MA, Tchernichovski O. 2016 The
forebrain song system mediates predictive call timing in female and Male Zebra finches. *Current*
Biology **26**, 309–318. (doi:10.1016/j.cub.2015.12.037)

30. Zann RA. 1996 *The zebra finch: a synthesis of field and laboratory studies.* Oxford, UK: Oxford
University Press.

31. Gess A, Schneider DM, Vyas A, Woolley SMN. 2011 Automated auditory recognition training and
testing. *Animal Behaviour* **82**, 285–293. (doi:10.1016/j.anbehav.2011.05.003)

32. van der Aa J, Honing H, ten Cate C. 2015 The perception of regularity in an isochronous stimulus
in zebra finches (Taeniopygia guttata) and humans. *Behavioural Processes* **115**, 37–45.
(doi:10.1016/j.beproc.2015.02.018)

33. Naqvi S, Godfrey AK, Hughes JF, Goodheart ML, Mitchell RN, Page DC. 2019 Conservation,
acquisition, and functional impact of sex-biased gene expression in mammals. *Science (1979)*
365. (doi:10.1126/science.aaw7317)

34. Zentner M. 2021 Social bonding and credible signaling hypotheses largely disregard the gap
between animal vocalizations and human music. *Behavioral and Brain Sciences* **44**, e120.
(doi:10.1017/S0140525X2000165X)

35. Krizman J, Bonacina S, Kraus N. 2020 Sex differences in subcortical auditory processing only
partially explain higher prevalence of language disorders in males. *Hearing Research* **398**,
108075. (doi:10.1016/j.heares.2020.108075)

36. Krizman J, Rotondo EK, Nicol T, Kraus N, Bieszczad K. 2021 Sex differences in auditory
processing vary across estrous cycle. *Scientific Reports* **11**, 1–7. (doi:10.1038/s41598-021-02272-
5)

37. Berninger E. 2007 Characteristics of normal newborn transient-evoked otoacoustic emissions: Ear
asymmetries and sex effects. *International Journal of Audiology* **46**, 661–669.
(doi:10.1080/14992020701438797)

38. Gall MD, Salameh TS, Lucas JR. 2013 Songbird frequency selectivity and temporal resolution
vary with sex and season. *Proceedings of the Royal Society B: Biological Sciences* **280**.
(doi:10.1098/rspb.2012.2296)

39. Krentzel AA, Remage-Healey L. 2015 Sex differences and rapid estrogen signaling: A look at
songbird audition. *Frontiers in Neuroendocrinology* **38**, 37–49. (doi:10.1016/j.yfrne.2015.01.001)

40. Brenowitz EA, Remage-Healey L. 2016 It takes a seasoned bird to be a good listener:
Communication between the sexes. *Current Opinion in Neurobiology* **38**, 12–17.
(doi:10.1016/j.conb.2016.01.005)

41. Blecher T, Tal I, Ben-Shachar M. 2016 White matter microstructural properties correlate with
sensorimotor synchronization abilities. *Neuroimage* **138**, 1–12.
(doi:10.1016/j.neuroimage.2016.05.022)

42. Vaquero L, Ramos-Escobar N, François C, Penhune V, Rodríguez-Fornells A. 2018 White-matter
structural connectivity predicts short-term melody and rhythm learning in non-musicians.
Neuroimage **181**, 252–262. (doi:10.1016/j.neuroimage.2018.06.054)

43. Asokan MM, Williamson RS, Hancock KE, Polley DB. 2021 Inverted central auditory hierarchies
for encoding local intervals and global temporal patterns. *Current Biology* **31**, 1762-1770.e4.
(doi:10.1016/j.cub.2021.01.076)

44. Rajendran VG, Harper NS, Garcia-Lazaro JA, Lesica NA, Schnupp JWH. 2017 Midbrain
adaptation may set the stage for the perception of musical beat. *Proceedings of the Royal Society
B: Biological Sciences* **284**. (doi:10.1098/rspb.2017.1455)

45. Honing H, Bouwer FL, Prado L, Merchant H. 2018 Rhesus monkeys (Macaca mulatta) sense
isochrony in rhythm, but not the beat: Additional support for the gradual audiomotor evolution
hypothesis. *Frontiers in Neuroscience* **12**, 475. (doi:10.3389/fnins.2018.00475)

46. van der Aa J, Honing H, ten Cate C. 2015 The perception of regularity in an isochronous stimulus
in zebra finches (Taeniopygia guttata) and humans. *Behavioural Processes* **115**, 37–45.
(doi:10.1016/j.beproc.2015.02.018)

47. Hulse SH, Humpal J, Cynx J. 1984 Discrimination and Generalization of Rhythmic and Arrhythmic
Sound Patterns by European Starlings (Sturnus vulgaris). *Music Perception* **1**, 442–464.
(doi:10.2307/40285272)

48. Samuels B, Grahn J, Henry MJ, Macdougall-Shackleton SA. 2021 European starlings (sturnus
vulgaris) discriminate rhythms by rate, not temporal patterns. *J Acoust Soc Am* **149**, 2546.
(doi:10.1121/10.0004215)

49. Bouwer FL, Nityananda V, Rouse AA, ten Cate C. 2021 Rhythmic abilities in humans and non-
human animals: a review and recommendations from a methodological perspective. *Philosophical
Transactions of the Royal Society B: Biological Sciences* **376**. (doi:10.1098/rstb.2020.0335)

Figures

521
Figure 1. Experiment design for testing the ability to flexibly perceive rhythmic patterns. *(A)*
Amplitude waveforms of isochronous (*Iso*) and arrhythmic (*Arr*) sequences of a repeated song element
(sound B; see Fig. S1) with 120 ms (*top*) and 180 ms (*bottom*) mean inter-onset interval (IOI). *(B)*
Schematic of the protocol. After a pre-training procedure (not shown), birds learned to discriminate
between isochronous and arrhythmic sound sequences at 120 and 180 ms IOI, for sounds A, B, and C
('ABC training', n=6 females) or starting with sound C followed by A and B ('CAB training', n=7 females)
(color indicates sound type; see Fig. S1). To test for the ability to generalize the discrimination to new
tempi, probe stimuli (144 ms IOI) were introduced after birds had successfully completed two training
phases. A subset of birds was then tested with a broader stimulus set using a novel sound element
(sound D) and every integer rate between 75 and 275 ms IOI ("Rule Training"; see *Materials and Methods*
for details). Color conventions indicating sound type are used in all subsequent figures. *(C)* Comparison
of training time, plotted as Trials to Criteria, for male and female birds that completed rhythm
discrimination training for all three phases. Symbols denote sex. For this and subsequent figures, data
from male birds were collected previously [22], and are replotted here for comparison. A Mann-Whitney
test for each training phase showed no sex-based differences in training time for any phase (*p*>0.05).

537
Figure 2. Learning of rhythmic pattern discrimination and generalization to new tempi for female
birds. (*A*) Learning curves for rhythm discrimination for a representative female bird (*y7o97*) across three
training phases (ABC training) in 100-trial bins. Overall proportion of correct responses is shown by *thick*
black line. Proportion correct for isochronous stimuli (S+, "hit rate") is indicated by *thin black* line;
proportion correct for arrhythmic stimuli (S-, "correct rejection") is indicated by *dotted line*. Chance
performance is indicated by dashed horizontal line. Data are plotted until criterion performance was
reached. (*B*) Results for rhythm discrimination training and probe testing for successful female birds
(n=7). Data left of the vertical dashed line show performance in the final 500 trials of the first rhythm
discrimination training phase (no probe testing, see Fig. 1B). Data right of the vertical dashed line show
performance during probe testing with stimuli at an untrained tempo of 144 ms IOI (*light gray*) and for
interleaved training stimuli (*dark gray*). Triangles denote performance for each female (n = 3 probe
tests/bird x 7 birds). All seven females were able to generalize the isochronous versus arrhythmic
discrimination to new tempi (20% different from the training tempi) for at least two sounds. Performance
on probe stimuli was significantly above chance for 17 out of 21 probe tests (*white* triangles; *p*<.0.0167,
binomial test with Bonferroni correction; black triangles denote performance not significantly different from
chance). Average performance across birds in each group is indicated by bars.

554
Figure 3. Comparison of generalization of rhythm discrimination by sex. Mean performance of all
successful males (*circles;* n=7) and females (*triangles*, n=7) during the last 500 trials of the first training
phase (first column) and for probe (*light gray*) and interleaved training stimuli (*dark gray)* in subsequent
phases. Data are collapsed across training order, and interleaved training trials are combined across
tempo. Dashed horizontal line indicates chance performance. Filled black symbols are not significantly
different from chance. Across probe tests, average performance was significantly lower for females than
for males by ~9% (*p*<0.05, mixed-effects logistic regression).

562
Figure 4. Sex differences in generalization of the learned discrimination across a wide range of
tempi. Average performance of 5 female birds (dark bars) during the initial 1000 trials of "Rule training".
Mean performance ± SD is plotted for 10 ms IOI bins. Chance performance is indicated by the horizontal
dashed line. Female performance for binned IOIs from 95-225 ms is significantly above performance in
the 75-85 ms IOI bin, where performance was at chance (***$p<0.001$; **$p<0.005$; *$p<0.05$, mixed-effect
logistic regression, black asterisks). Data from 7 males from [22] are plotted for comparison (light green
bars and white asterisks). IOIs used in rhythm discrimination training phases before rule training are
shown in bold and boxes on the x-axis.

571
Figure 5. Reaction times during rhythmic pattern discrimination and frequency discrimination
tasks. Mean reaction time for the final 500 responses in the first training phase are shown for three
groups of birds. For rhythm discrimination, data are shown for birds which completed all three training
phases (first bar, n=7 males and 7 females) and birds that did not complete rhythm discrimination training
(middle bar, n=3 males and 6 females). For frequency discrimination, data are shown for a separate
cohort of birds (third bar, n=4 males and 6 females). The 500ms period between trial start and activation
of the response switch is indicated by the horizontal dashed line. Symbols indicate the average reaction
time for each bird. A 2-way ANOVA of sex, group, and sex-by-group interaction showed only a significant
effect of group ($F_{(2,27)}$, $p = 0.001$). Successful rhythm discrimination birds responded significantly slower
than the other two groups ($p<0.01$ compared to the failed birds, $p<0.005$ compared to the frequency
discrimination birds; Tukey post-hoc tests).

Supplemental Figures

584
585 **Figure S1. Amplitude waveforms of the four sound elements used in the stimuli.** Sound files are
586 available at https://dx.doi.org/10.17632/fw5f2vrf4k.2

587
Figure S2. Learning curves for discrimination of stimuli based on rhythmic pattern, tempo, or
588
589
frequency. (*A*) Learning curves for 7 female birds that successfully discriminated isochronous vs.
590
arrhythmic stimuli within 30 days for each sound type. (*B*) Learning curves for a separate cohort of
591
females (n=6) trained to discriminate isochronous sequences of sound A that differed in tempo (120 ms
592
IOI vs. 144 ms IOI) or frequency (shifted up 3 semitones vs. 6 semitones). All females tested met the
593
criterion for discriminating stimuli based on tempo or frequency. Conventions as in Fig. 2A**.**

594
Figure S3. Relationship between performance on probe trials and interleaved training trials. Across
birds that successfully completed all three training phases (n=14), there was a significant positive
correlation between the accuracy on interleaved training stimuli (learned discrimination) and probe stimuli
(generalization); least squares regression fit; R^2 =0.31; p = 0.0375). For each bird, data were combined
across all three sound elements; each data point is the average performance across 240 probe stimuli
and ~2000 interleaved training stimuli.

601
602 **Figure S4. Reaction time (RT) across a broad range of tempi.** Mean reaction time for correct trials
603 versus tempo during Rule Training for male (*circles*) and female (*triangles*) birds. RT tended to be longer
604 at the end of the tempo range (<100 ms and >220 ms), when discrimination of isochronous vs. arrhythmic
605 stimuli fell to chance. Fitted quadratic curves are plotted to help illustrate the observed relationship.