1
2
3 # Loop stacking organizes genome folding from TADs to chromosomes
4
5

Antonina Hafner[1], Minhee Park[1,2], Scott E. Berger[3], Elphège P. Nora[4,5], Alistair N. Boettiger[1*]

[1] Department of Developmental Biology, Stanford University, United States

[2] Current address: Department of Biological Sciences, Korea Advanced Institute of Science and

Technology, Republic of Korea

[3] Biophysics Program, Stanford University, Stanford, United States

[4] Cardiovascular Research Institute, University of California San Francisco, United States

[5] Department of Biochemistry and Biophysics, University of California San Francisco, United

States

* Corresponding author. Email: boettiger@stanford.edu

Abstract

While population level analyses reveal significant roles for CTCF and cohesin in mammalian genome organization, their contribution to chromatin structure and gene regulation at the single-cell level remain incompletely understood [1–4]. Here, we use a super-resolution microscopy approach, Optical Reconstruction of Chromatin Architecture (ORCA) [5] to measure the effects of removal of CTCF or cohesin on genome folding across genomic scales. In untreated embryonic stem cells, we observe intricate, frequently stacked loops of chromatin which are largely dissolved upon cohesin removal. The loops compact chromatin at the < 3 Mb scale, increasing proximity between sequences not only within but also between TADs. We find multi-way contacts among loop anchors, preferentially at TAD borders, and these hubs largely dissolve upon CTCF degradation. CTCF-hubs bridge intervening TAD boundaries while keeping border distal regions from neighboring TADs apart outside the hub. Cohesin dependent loops at the < 3 Mb scale impede mixing at larger chromosomal scales through steric effects of loop stacking, dramatically reducing genomic cross-talk. Disruption of this ordered chromosomal structure led to increased cell-cell variability in gene expression, exceeding changes to average expression. Together our data revise the TAD-centric understanding of CTCF and cohesin, and provide a multi-scale, structural picture of how they organize the genome on the single-cell level through distinct contributions to loop stacking.

Main

Introduction

Mammalian genomes are organized in 3D into Topologically Associating Domains (TADs). TADs

were discovered by Hi-C [6–8] and are defined as regions in which within-domain contacts are

substantially more common than between-domain contacts [9–12]. These preferential interactions

within TADs are thought to be important for the specificity of gene regulation as enhancers and

their target promoters frequently lie within the same TAD, and in a multitude of cases mutations

disrupting TAD boundaries alter normal enhancer-promoter contact and gene expression [1,3,13–15].

Cohesin and CTCF are essential for the establishment and maintenance of TADs. Indeed, CTCF

and cohesin binding is enriched at TAD boundaries and acute depletion of these proteins has

been shown by Hi-C and imaging to lead to loss of TADs [16–20]. Polymer modeling work has

proposed that these two proteins work together in forming TADs through distinct roles; cohesin

extrudes loops of chromatin but stalls at sites bound by CTCF [21,22], and considerable evidence

supports these molecular roles [2]. However, a detailed view of the physical 3D structure and how

it is perturbed upon removal of CTCF or cohesin, with resolution from genes to whole

chromosomes, is lacking. Such a high-resolution view may tease apart mechanisms or regimes

in which CTCF and cohesin operate that are not distinguished by ensemble averages alone, and

thus help us better understand how they effect cis-regulation of transcription. Microscopy

approaches are well suited for investigating heterogeneity at the single-cell level.

Our earlier work on cohesin depleted cells imaging megabase-scale regions found 'TAD-like'

domains were preserved in the absence of cohesin, though the position of their borders became

more uniform [20]. Two more recent microscopy studies that labeled TADs and looked at the

consequences of cohesin and CTCF loss between and within TADs came to opposing

conclusions [23,24]. Upon degrading cohesin, Luppino and colleagues found reduced overlap

between adjacently labeled TADs, while the volume of the TADs remained the same. By contrast,

the second imaging study by Szabo and colleagues found that loss of cohesin did not significantly

change the overlap between adjacently labeled TADs but increased the volume of each TAD.

To better understand the role of cohesin and CTCF on genome structure, we imaged genome

structure from the few-kilobase scale of cis-regulatory elements to the scale of whole

chromosomes, using a multi-scale chromosome tracing approach, similar to (refs [25–27]). We

multiplexed this imaging by using single-cell barcodes to combine different genetically edited cell

lines and different treatment conditions into a single experiment, enabling direct measure of 3D

structure in nanometers without risk of batch effects. Our single-chromosome data provide a

detailed view of how cohesin and CTCF shape the 3D structure of chromatin from the kilobase to

the whole chromosome scale; enhancing local contact by forming heterogeneous loops,

preventing distal contact by loop clash, and facilitating border bypass by positioning loop anchors.

Results

We use microscopy to understand how TADs are lost upon cohesin and CTCF depletion

We first wanted to find out *how* TADs disappear; is it through loss of preferential interactions within

TADs or gain of interactions between TADs? (**Fig. 1a**). These two scenarios would lead to distinct

biophysical structures (**Fig. 1a**). We selected two ~2.5 Mb regions, encompassing multiple TADs

and sub-TAD structures (**Fig. 1d**), and containing developmentally important transcription factors,

Hoxa or Sox2.

We used a microscopy chromosome tracing approach, Optical Reconstruction of Chromatin

Architecture (ORCA) [5] to image single chromosomes in mouse embryonic stem cells (mESCs)

with or without CTCF or RAD21, a subunit of the cohesin complex. We used cell lines where

RAD21 and CTCF are tagged with the auxin inducible degron (AID) at their endogenous loci and

can be rapidly depleted by addition of auxin (**Extended Data Fig. 1a**) [19,24]. For all experiments

described hereinafter, auxin-treated cells (+Aux) were treated with auxin for 4 hours, and control

cells were not treated (NT).

To image all conditions (CTCF NT, CTCF +Aux, RAD21 NT, RAD21 +Aux) within the same

experiment, we developed a multiplexed ORCA approach. We adapted a recently published

method from single-cell sequencing [28] to attach distinct oligonucleotide-barcodes to the cell

surface, after which cells can be combined, hybridized with ORCA probes, and imaged within the

same experiment, where cell surface barcodes are read out by sequential hybridization, similar

to ORCA barcodes (**Fig. 1b, 1c**). This approach both accelerates data collection and removes

potential batch effects, intrinsic to high-content, high-resolution assays. After imaging, we

demultiplex cell labels and assign the single chromosome traces from ORCA to individual cells

from each genotype and each treatment condition. In the multiplexed cell image shown in Fig.

1C, we combined the parental E14 cell line with RAD21 and CTCF lines +/- auxin treatment.

Replicates showed high reproducibility (**Extended Data Fig. 1b**). To increase cell numbers and

statistical power, these were merged for all subsequent plots and analyses.

We can compute population-level contact frequency maps with ORCA by calculating, for each

pairwise combination, the fraction of chromosomes that are within a certain distance (we used

200 nm, approximately equal to the mean distance between the centers of adjacent regions).

Plotting contact frequency over all ORCA traces in each condition (**Fig. 1d**) we found good

agreement with Hi-C contact maps (**Extended Data Fig. 2a, b**). Cohesin-depleted contact maps

showed loss of TADs [17,18] in CTCF-depleted cells, TADs were still visible, although with lower

insulation between them (**Fig. 1d**, **Extended Data Fig. 2c**). Published Hi-C data of CTCF

depletion in mESCs [16,19] also showed that, unlike cohesin depletion [18] CTCF loss did not fully

abolish TADs for these regions (**Extended Data Fig. 2a, b**).

Thus, with the ability to 1) rapidly deplete CTCF and cohesin, 2) multiplex multiple cell types and

treatment conditions, and 3) measure genome structure that matches well at the population level

with Hi-C, we set out to examine in detail how these proteins affect genome folding in the

subsequent sections.

CTCF contributes to TAD separation while enabling cross-TAD contacts by organizing

chromosomes radially

In ORCA and Hi-C population contact frequency maps for unperturbed cells, we observed loops

that span CTCF-marked TAD borders, which we called 'bridges' for short (**Fig. 2a, b, Extended**

Data Fig. 3a, b). Hi-C data show such bridges throughout the genome (e.g. [29]), though the

population-average nature of the data does not distinguish between multiple structural origins of

this feature. Such a bridging pattern in contact frequency could arise from a heterogeneous

population in which distinct portions of the population have either the long-range contact or the

separated structure (**Fig. 2a**). To test for this, we turned to our single-chromosome data and

examined the all-to-all contact frequency computed using only those chromosomes which

contained bridging contacts (**Fig. 2b, Extended Data Fig. 3c**). Interestingly, this revealed that the

population of chromosomes with bridging contacts still contained clear TAD borders beneath the

bridges (**Fig. 2b, Fig. Extended Data Fig. 3c**). Thus, a bridge can occur between separated

structures and the two scenarios (**Fig. 2a**) are not mutually exclusive.

To better see how these bridge chromosomes differ from non-bridge chromosomes we subtracted

their contact maps. The difference maps revealed a prominent stripe connecting the bridge

anchors, with strengthened loops among intervening CTCF sites in bridge-containing

chromosomes (**Fig. 2c**, **Extended Data Fig. 3d**). This implies that bridge anchors frequently

contact several additional sites in between, with a preference for intervening CTCF sites, rather

than spanning anchor-to-anchor in a single loop (**Fig. 2d**). Inspection of individual chromosome

traces revealed examples of distal CTCF sites bridged by multiple loops, including intervening

CTCF sites (**Fig. 2e**).

We next computed the frequency of three-way contacts among CTCF sites. We compared these

to distance-matched, non-CTCF interactions using a sliding window across the genomic region.

Three-way contacts were more common among the CTCF sites than non-CTCF sites (**Fig. 2f**).

Notably, CTCF-site-proximal regions also showed elevated rates of three-way contacts, which fell

off as a function of distance from the CTCF-site (**Fig. 2f**). The three-way contact frequency among

CTCF sites was also higher than expected based on the measured pairwise interaction

frequencies (assuming pairs are independent), providing further evidence of preferential

clustering missed in bulk pairwise data (**Fig. 2f**). Depletion of CTCF significantly reduced the

frequency of three-way interactions among both CTCF and CTCF proximal sites, though made

little change for CTCF-distal sites (**Fig. 2f**).

To explore the effects of CTCF loss across the domain, we computed the change in contact

frequency upon CTCF degradation. We note that ORCA contact frequency is in absolute terms

(fraction of chromosomes), rather than normalized within the experiment, enabling a direct

subtraction between contact frequencies. Previous microscopy studies [23,24] and simulations [21,30]

concluded that the distance between TADs decreased upon CTCF depletion. ORCA data from

the *Hoxa* and *Sox2* domains revealed a more complex pattern upon CTCF depletion, where some

pairwise distances increased while others decreased (**Fig. 2g**). In particular, while several of the

CTCF borders exhibited increased interaction between the upstream and downstream TADs, TAD

bridges exhibited reduced interaction, even though the contact frequency increased between

TADs underneath the bridge (**Fig. 2g** red box, blue circle).

Together these single-cell analyses point to a tendency of CTCF to organize chromatin in a radial

pattern, rather than functioning purely as a border between domains or a tether within a domain

(**Fig. 2d**). CTCF-border sites preferentially position on the axial part of this structure (**Fig. 2d** gray

region), where they are more likely to interact, including across boundaries. Intra-TAD regions

(far from CTCF sites) preferentially position away from this axis, where they are generally more

isolated from other intra-TADs. Depletion of CTCF reduced this clustering tendency of the CTCF-

marked borders and led to more disordered positioning of loops (**Fig. 2h, i**), which led to a

weakening of TAD borders on the population-level contact frequency maps (**Fig. 1d**). We note

that although this radial organization is reminiscent of the rosette, single-cell traces show

considerable heterogeneity of the number and identity of sequences at the center, and a relatively

low frequency of multi-way contacts for any particular triplet, even though triplet interactions

among CTCF sites are substantially more common than non-CTCF sites (**Fig. 2f**).

Cohesin loss increases distances both within and between TADs

We next examined the role of cohesin in this radial chromosome organization. Degradation of

cohesin clearly reduced the preference for CTCF and CTCF-proximal regions to form three-way

contacts (**Extended Data Fig. 3e**). In contrast to CTCF depletion, three-way contacts among all

loci decreased, both CTCF-proximal and CTCF distal (**Extended Data Fig. 3e**). Inspecting

individual chromosome traces, we found that depletion of cohesin led to less loopy structures and

expansion of the traces (**Fig. 3a**), unlike CTCF, where loops persisted in CTCF-depleted cells but

with more random positioning of TAD borders (**Fig. 2h, i**).

To quantify the expansion after cohesin depletion across the whole population, we plotted the

median pairwise distances in units of nanometers (**Extended Data Fig. 4a**). We found that

depletion of cohesin led to increased pairwise distances by up to a few hundred nanometers for

both *Hoxa* (2.2 Mb) and *Sox2* (2.8 Mb) domains (**Fig. 3b, Extended Data Fig. 4a**). Distances

increased significantly (ranksum, *p*<0.05) **(Extended Data Fig. 4b)** for all pairwise interactions

both within and between TADs, with the fold change of expansion being greater within TADs

(**Extended Data Fig. 4c**). The only non-significantly changed interactions (**Extended Data Fig.**

4b) we observed were at the Polycomb-bound *Hoxa* region (**Fig. 3b, Extended Data Fig. 4d**

arrow) which remained compact and maintained contact with another upstream Polycomb-bound

region in 19.4% of cells (relative to 29.7% in untreated cells) (**Extended Data Fig. 4d**).

An advantage of the multiplexed microscopy approach is that changes in distance are measured

in absolute units of nanometers, and systematic batch effects between experiments are removed

by analyzing both treated and untreated cells at the same time. It is more challenging to make

direct comparisons between treatment conditions with Hi-C data, as there are no spike-in control

methods available to enable normalization across experiments. This uncertainty may explain why

the untreated/cohesin-depleted ratio of Hi-C maps for *Hoxa* and *Sox2* suggested reduced contact

within and increased contact between TADs (**Extended Data Fig. 4e**), in contrast to the ORCA

data - even though the original, non-subtracted contact frequency maps agree qualitatively for

both conditions (**Fig. 1d**, **Extended Data Fig. 2b**).

To test whether this expansion was dependent in part or in full on the loss of sister chromatid

cohesion, we performed immunofluorescence labeling of Geminin and ORCA in the same cells

(**Extended Data Fig. 4f**). Geminin is a cell-cycle-regulated protein that is not detected in G1 and

progressively accumulates through the S and G2 phases of the cell cycle. We selected cells with

low Geminin levels (lowest 25% of the cells) and found that these also showed expansion upon

loss of cohesin, both within and between TADs (**Extended Data Fig. 4g**). Further, we found no

correlation between Geminin levels and the median pairwise separation among all imaged

domains on a single-cell basis (**Extended Data Fig. 4h**). From this we conclude that the

expansion of chromosomes at the megabase scale is primarily driven by changes in cis-structure,

and not loss of sister cohesion.

In summary, not only do we find that TAD loss is the consequence of increased separation of

sequences within the TAD (and not mixing between TADs), but that it occurs in the context of a

significant general expansion of the genome at the megabase scale (TAD scale) (**Fig. 3c**). Our

findings are consistent with previous measurements at this scale that found an increased

separation of genomic loci in *cis* upon cohesin loss in fixed cells [20,23,24,31–34] and live cells [35,36].

Cohesin loss leads to increased mixing at the chromosome scale

Increase of pairwise distances along the ~2.5 Mb scale upon loss of cohesin made us wonder

what the consequences were on genome organization at the nuclear scale. We first quantified

the nuclear area among untreated and cohesin-depleted cells as well as the E14 parental cell

line. We observed a similar distribution of nuclear sizes in all conditions (**Extended Data Fig. 5a**).

Given the expansion of distances at the 2-3 Mb scale and lack of change in nuclear size, we

wanted to understand what happens to the genome at the tens-of-megabases and whole-

chromosome scales. We designed probes spanning both a ~30 Mb region and the whole

chromosome (~150 Mb). We did this for both chromosomes 3 and 6 that contain the *Sox2* and

Hoxa domains examined previously (**Fig. 4a**).

Previous work using Hi-C found that depletion of cohesin did not significantly alter the

chromosome-scale genome organization other than a slight strengthening of compartments [17,37].

Compartments refer to a plaid pattern commonly seen at the chromosome scale in Hi-C contact

maps, indicating a preferential separation of the genome into two groups, commonly denoted A

and B [38]. The separation is readily visualized through the Pearson map, where each element of

the map reports the cross-correlation of the corresponding row and column after distance

normalization [38]. Examining the Pearson maps for the 30-Mb and full-chromosome ORCA data,

we observed a similar plaid pattern to Hi-C (**Fig. 4b**, **Extended Data Fig. 5b, c**), an agreement

also observed in compartment analysis from earlier tracing experiments [39]. Upon cohesin

depletion, we also observed a slightly stronger plaid pattern and higher Pearson's r, indicating

compartment strengthening (**Fig. 4b, Extended Data Fig. 5b-e**).

We then looked at single-chromosome traces to understand how the expansion we observed at

the 2-3 Mb scale could be reconciled with minor compartment changes at the 30 Mb and 150 Mb

scales. The 30-Mb and whole-chromosome traces looked more disordered after depletion of

cohesin. While in untreated cells, adjacent labels (**Fig. 4c, d** red and blue labels) remained in

physical proximity relative to genomically distant regions (red and cyan), in cohesin-depleted cells,

there was less distinction in terms of physical distance between genomically proximal or distal

labels. To test these effects population wide, we plotted the difference of median distance

between auxin-treated and untreated cells for all pairwise combinations (**Fig. 4e, f**). We found

that at the 30-Mb scale (**Fig. 4e**), pairwise distances less than ~4 Mb apart, increased significantly

(**Fig. 4e, Extended Data Fig. 5f**), consistent with our data showing expansion at the TAD scale

(**Fig. 3a, b**). However beyond the ~4 Mb separation we found a gradual transition from increased

to decreased distance (**Fig. 4e**, blue to white to red). The decrease in distance upon cohesin

depletion became more and more pronounced and significant (**Extended Data Fig. 5f**) as the

genomic distance between two pairs increased, as seen in the chromosome-scale difference

maps (**Fig. 4f**).

Our ORCA measurements at the 30-Mb and chromosomal scale show that cohesin is essential

for maintaining an ordered chromosome structure (**Fig. 4g**). By primarily contributing to a more

linear ordering of the chromosome, cohesin also opposes the demixing of compartment types

(**Fig. 4h, i**), explaining the slight compartment strengthening after cohesin depletion.

Loop extrusion simulations can exhibit multiscale effects on genome folding

Considerable evidence supports the model that cohesin shapes genome structure, including

TADs, through loop extrusion (for review see [2,40]). In this model, cohesin is loaded onto DNA and

extrudes a loop until it is stalled by CTCF. We wanted to know if our ORCA observations of

expansion at the ~2.5 Mb scale (**Fig. 3**) and increased mixing at the 30-Mb and full-chromosome

scales (**Fig. 4**) could also be understood through the process of loop extrusion. We performed

Langevin dynamic simulations of loop extrusion extended from previous polymer models [21,30]

(**Methods**). As expected [21], simulations including cohesin formed TADs between CTCF

boundaries, and loops between convergent CTCF sites, while simulations without cohesin did not

(Extended Data Fig. 6a). Examining individual traces at the 3-Mb scale, we found that without

cohesin loops, the polymer trace expanded (**Fig. 5a**). To follow structural changes at the

chromosome scale, we plotted the whole chromosome structure and colored 3-Mb-sized regions

blue, red and cyan (**Fig. 5b**). With cohesin, adjacent regions (blue and red) could be found

proximal in 3D space (**Fig. 5b**). In contrast, without cohesin, individual regions expanded

significantly, such that the ends of a 3-Mb region could be near opposite sides of the chromosome

territory (e.g. red domain and gray chromosome, **Fig. 5b**). This expansion also facilitated

proximity between linearly distal elements in some chromosomes (**Fig. 5b**, blue and cyan

regions). This expansion at shorter scales and contraction at larger scales could be seen across

the population of replicate simulations, as shown by the difference of the median distance maps

(**Fig. 5c, and Extended Data Fig. 6b**. Thus, cohesin-mediated loops reduce disorder in chromatin

folding by ensuring linearly proximal regions remain proximal in 3D and linearly distal regions

remain distal in 3D.

Not all parameter choices for loop extrusion produced the switch from expansion to contraction

that we observed in our experimental data and in the simulations described above. Simulations

with low-density confinement (confined to 10% of their relaxed volume) exhibited increased

separation at all length scales (**Extended Data Fig. 6c**). Chromosomes with a higher frequency

of chain-crossing behavior (as might be expected for a high level of topoisomerase, see **Methods**)

also failed to exhibit the qualitative switch, expanding at short scales and exhibiting no change at

larger scales (**Extended Data Fig. 6d**). Thus, only with substantial confinement and reduced

chain-crossing did the cohesin-loops significantly hold apart the more distal parts of the

chromosome by steric exclusion – loop-clash. This finding is consistent with recent polymer theory

showing that chromatin loops "experience topological repulsion" [41].

Furthermore, our simulations predict that at length scales less than ~200 kb, the relative effect of

cohesin loss on 3D contacts would be less than observed at the Mb scale (**Fig. 5c**), potentially

because 200 kb is shorter than the typical distance between cohesins (**Fig. 5a,** black boxed

regions). To test if <200 kb regions are indeed less affected by cohesin loss, we labeled at a

higher resolution (5-kb step size) the 170 kb region containing the Sox2 gene and its downstream

enhancer [42,43], as well as more distal regions flanking this domain, 200 kb to 1.5 Mb apart (**Fig.**

5d, e). Upon depletion of cohesin, we found that within the 170 kb region, expansion was

significantly lower (ranksum p=1.03*10-168) than for the more distally spaced probes (**Fig. 5f,**

Extended Data Fig. 7).

Thus we find that the loop extrusion model is consistent with ORCA data collected at all scales

(**Fig. 5c, f**), if sufficiently confined in the nucleus and sufficiently restricted in the ability for chain-

crossing. The model does not quantitatively match the data in all respects, but does show how

loop extrusion, with a few parameter constraints, can explain the key qualitative trends in the data.

The picture that emerges is that extrusion of chromatin loops by cohesin locally compacts DNA

both within TADs and across TAD borders, and that these loops further serve to organize the

chromosome by sterically inhibiting longer-range contacts that would otherwise be promoted by

the dense packaging environment of the nucleus.

Cohesin depletion increases the variability in gene expression

Previous studies have shown that degradation of cohesin has a limited immediate effect on

transcription [17,18,44–46] relative to degradation of transcription factors or co-factors [47–50] or genetic

disruptions of TAD borders [5,51–54]. Given the increased disorder in genome structure, where

proximal regions got farther apart and distal genomic regions became closer, we wondered if at

the single-cell level cells would exhibit increased variability in gene expression.

We performed multiplexed single-molecule FISH to measure expression of a set of

developmentally regulated pluripotency genes. We used cellular barcoding to minimize batch

effects among different treatment conditions and cell lines (RAD21-AID, CTCF-AID, E14) (**Fig.**

6a). We also added mock (DMSO) treated cells as a further comparison for the magnitude of the

effect. The mean mRNA counts showed small fold changes between treatment conditions. Effects

of auxin on median depletion were scarcely stronger than for our mock treatment (**Fig. 6b**). To

see if cohesin depletion led to increased variability between cells, we calculated the coefficient of

variation. We found that the majority of genes tested showed a modestly higher coefficient of

variation after cohesin depletion (**Fig. 6c**). Depletion of CTCF had less effect on the coefficient of

variation (**Fig. 6c**), likely reflecting less change in genome structure after CTCF depletion at either

the TAD scale or chromosome scale (**Extended Data Fig. 8**).

Cluster analysis by TSNE showed that RAD21-depleted cells did not split into distinct groups, nor

cluster separately from untreated cells (**Fig. 6d**). This suggests increased variability was not due

to partial differentiation of the mESCs after auxin treatment for 4 hours. Indeed all the genetic

backgrounds and treatment conditions clustered together (**Fig. 6d**), with variation of cell-cycle-

dependent factors aligning with one major TSNE axis and variation in *Tbx3* defining the other

(**Fig. 6e**), a gene which showed little CTCF or cohesin dependent change (**Fig. 6b**).

Overall our data show that acute loss of cohesin leads to disordered chromosome structures that

result in modestly increased cell-to-cell variability in gene expression, which is missed by bulk

population assays.

Discussion

Here we introduced a multiplexed approach for ORCA, enabling simultaneous measurement of

chromosome folding across multiple treatment conditions. Our approach enables quantification

of physical structure in absolute units, enabling quantitative comparison between conditions,

overcoming a limitation of Hi-C. As the field moves increasingly to the study of perturbations, we

expect the ability to measure absolute frequency differences across genomic length scales

without normalization artifacts or batch effects, the ability to measure variation in single cells, and

the ability to process multiple conditions at a time will facilitate more rapid understanding. In this

work, we characterized the structure of the interphase genome from the kilobase to the

chromosome scale, expanding our understanding of the role of CTCF and cohesin in shaping that

structure, with implications for gene regulation.

CTCF is thought to determine TAD borders by defining the bounds of cohesin-mediated loop

extrusion [1–4]. Consistent with this, 2-color FISH experiments [19,23,24] reported a decrease in

distances between TADs, and an increase in overlap. While supporting these conclusions, our

ORCA data revealed more complex patterns upon CTCF loss. In addition to supporting TAD

boundaries, we identified a role for CTCF in bridging across TAD borders through a multi-loop

structure. This behavior provides a mechanism that would enable contacts between enhancers

and promoters that are separated by one or more TADs. Indeed, one of the CTCF-border-crossing

sites in the *Hoxa* locus has recently been shown to contain enhancer activity for *Hoxa* genes

during hematopoiesis [55] and cranial-facial patterning [56]. Furthermore, several groups recently

inserted CTCF sites between the *Sox2* gene and its embryonic enhancer region, but found

minimal impact on *Sox2* expression, despite establishing a new TAD border [57,58]. As both the

enhancer region and promoter are each already CTCF proximal, the multi-way clustering of CTCF

could explain this insulator bypass as well.

The radial organization arising from the preferential multi-way contacts of CTCF sites resembles

earlier models of chromatin structure based on 3C assays, like the archipelago of enhancers and

promoters described for *Hoxd* [59], the subTAD-clusters described for *Hoxa* [60], or the CTCF hubs

described for the *protocadherin* genes [61], or B-globin genes [62,63] among other regions of the

genome [64,61] Multi-contact 4C and multi-contact-Hi-C experiments have provided further evidence

for three-way interactions among CTCF sites [63,65]. By observing the structure of individual

chromosomes, we provide more direct evidence for these multi-way interactions and how they

maintain intervening borders. We also see they exist within a heterogeneous population, in which

diverse parts of the chromatin fiber alternate at loop bases and that any particular triplet is

individually rare.

Furthermore, CTCF degradation, compared to cohesin degradation, exhibited substantially

weaker impacts on TADs in our data, consistent with recent reports [16,19,66]. Whether this is due to

co-factors at some sites which help retain CTCF locally amidst global depletion or additional

factors contributing to cohesin-dependent boundary activity [57,67] remains an interesting question

for future studies.

Cohesin depletion led to a disordered genome structure with distinct effects at different genomic

scales. At the ~200 kb scale, typical of the majority of enhancer-promoter interactions [68], we found

relatively mild increases in pairwise distances as compared to the ~2.5 Mb scale which showed

a significant expansion after depletion of cohesin. Thus, effects of cohesin loss would be different

for each gene. Genes whose enhancers are within ~200 kb distance, will be less affected in terms

of E-P contact than genes that rely on more distal enhancers (>200 kb - 3 Mb away). At larger

genomic scales (>4 Mb) we found increased mixing. This implies that all genes are more likely to

experience contacts with very distal enhancers (>4 Mb to 150 Mb away) and these ectopic

enhancer-promoter contacts will be different for each cell. While any particular contact occurs at

relatively low frequency, each promoter is now exposed to many more such rare long-range

contacts than before. With this perspective, rather than expecting a global up- or down-regulation

of genes after depletion of cohesin, our ORCA measurements suggested that each gene and

each cell would behave differently. This would manifest as increased cell-cell variation in mRNA

levels, which we observed for most of the developmentally important genes we assayed by single

molecule RNA FISH. Recent studies have similarly reported that disruptions in chromatin 3D

organization increase variability in gene expression, using single-cell RNA-seq [69], RNA-FISH and

flow-cytometry [70], or integrative modeling of omics data [71]. Whether and how these acute changes

in cell-cell variability due to disruption of chromatin structure connect to functional outcomes of

cohesin loss or reduction in development and disease [72–77] will require future investigation.

The structural changes we observe in genome organization at the chromosome scale also allow

us to revisit recent conclusions about the role of cohesin in aspects of genome organization, such

as compartments and polycomb loops. It has been proposed that cohesin opposes demixing of

the genome into A/B compartments directly, through its ability to bridge compartment boundaries

and pull together A and B chromatin [17,30,37]. This proposal is supported by the observed

strengthening of A/B compartments upon cohesin depletion [17,37]. Our data suggest an additional

mechanism. By significantly decreasing the frequency of interactions >4 Mb, through the dense

stacking and clash of loops, cohesin indirectly disrupts compartmentalization by reducing the

opportunity of linearly separated A (or B) domains to contact and mix. While both the direct and

indirect effects of cohesin may operate in tandem, the direct explanation predicts that short-range

compartmentalization should be most affected (where A meets B along the linear genome),

whereas the indirect mechanism suggests long-range interactions should be most affected.

It has been proposed that cohesin processivity can disrupt the Polycomb (Pc) homotypic

interactions which drive formation of ultra-long range contacts among Polycomb domains (1-20+

Mb), "Polycomb loops" [18]. This data is supported by the observation that these loops strengthen

upon cohesin degradation [18]. As we have shown for the 1-Mb scale interaction in the *Hoxa* region,

the shorter Pc-Pc interactions actually decrease in absolute frequency upon cohesin loss, while

>4 Mb interactions increase along the entire chromosome. Thus, the effect of cohesin on Pc loops

we propose is also an indirect effect of cohesin linearizing the chromosome (promoting short <4

Mb contacts and reducing longer range ones). Indeed, shorter range Pc loops were also reported

to show less change or increase interaction by Hi-C [18], an effect readily explained by the loop-

clash model that would not be expected if cohesin linear scanning were to knock apart Pc-Pc

loops.

Together, these data suggest cohesin and CTCF organize the genome through their distinct

effects on chromatin loops and loop stacking. Cohesin increases the density of chromatin loops,

and thus proximal sequences are brought into frequent contact by individual loops and stacking

loops, while distal sequences are held apart by the clash of many loops. When loops stack

together, the loop bases aggregate in a central hub. By stalling cohesin, CTCF sites are more

likely to be found at loop bases, and more likely to end up at the hub. This aggregation of 3 or

more borders facilitates cross-border contacts among CTCF-proximal regions, while maintaining

border function among CTCF-distal regions. Thus, beyond its known functions in enhancer

blocking and looping, CTCF may also facilitate cross-TAD gene regulation.

Methods

Cell culture, treatment and collection

E14 parental cell line as well as the Rad21-AID and CTCF-AID cell lines were a gift from Elphege

Nora. Cells were grown in DMEM/F12 supplemented with 15% FBS, NEAA, Glutamax. A final

concentration of 500uM IAA (Sigma i2886) was used for all auxin treatments. Cells were treated

for 4 hours, media was changed in the untreated condition at the same time as the fresh media

with auxin was added to auxin treated cells.

For cell collection, cells were trypsinized, counted, spun down and after removal of supernatant

media, were fixed in suspension in 4%PFA in 1xPBS on a nutator for 10 minutes, washed 3 times

with 1xPBS, resuspended in 70%EtOH in 1xPBS at a concentration of 1-10 000 cells/ul and stored

at -20C for up to 6 months.

Western blot

Cells were collected by trypsinization, washed in 1xPBS and pellets were frozen on dry ice/EtOH

bath and stored at -80C. Cells were lysed in mRIPA buffer with Protease and Phosphatase

inhibitor cocktail. 20ug of protein was run on a 4-20% Mini-PORTEAN TGX precast gels in Tris-

Glycine-SDS running buffer. After transfer onto the nitrocellulose membrane, and blocking in Li-

Cor Intercept Blocking Buffer (TBS), blots were probed with either CTCF (Abcam 128873, 1:1000)

or Rad21 (Abcam 154769, 1:1000) antibodies as well as beta-Actin (Cell Signaling 3700) in

blocking buffer at 4C overnight. Membranes were then washed 3x 5min in TBST at room

temperature, incubated with secondary antibodies (Li-Cor: IRDye 680RD Donkey anti-Rabbit 926-

68073 and IRDye 800CW Donkey anti-Mouse, 1:10 000 diluted in blocking buffer) for 1h at room

temperature. After 3x 5min washes in TBST, and 1 wash in TBS, samples were imaged on the

Li-Cor scanner.

Multiplexed cell labeling

The multiplexed cell labeling protocol was adapted from Gehring et al. 2020. The detailed protocol

is as follows.

Part 1: Preparation of labelled oligos

1) 3' amine-modifed oligos were ordered from IDT and resuspended to 500 µM in 50 mM sodium borate buffer pH 8.5

2) In 1.5 mL microcentrifuge tube, add 25.0 µL 3' amine-modified oligo in 50 mM sodium borate buffer, 8.2 µL 10 mM Methyltetrazine-NHS ester, 41.8 µL DMSO, incubate protected from light for 30 minutes with agitation

3) After 30 minutes, add another 8.2 µL 10 mM Methyltetrazine-NHS ester, then add once again after an additional 30 minutes

4) After the 90-minute reaction, quench by adding 180 µL 50 mM sodium borate buffer and 30 µL 3 M NaCl, then add 750 µL ice cold 100% EtOH]

5) Precipitate @ -80 °C overnight

6) Transfer contents to ultracentrifuge tubes, balance with analytical balance, and spin at 20,000 xg for 30 minutes at 4 °C (see Barna lab for TLA 120.2 rotor, we have model 343778 tubes)

7) Wash twice with 750 µL 70% EtOH, ensuring not to disrupt the pellet (don't mix pipette). Additionally, let pellet dry for ~10 minutes in fume hood until nearly all ethanol has evaporated

8) Resuspend pellet in 100 µL cold HEPES buffer

9) Quantify recovery with A260 and manufacturer's listed molar extinction coefficient. Typical concentration is around ~80 µM.

<u>Part 2: Cell labeling</u>

1) Between 500K and 5M fixed cells (stored in 70%EtOH at -20C) were washed twice in

1xPBS and resuspended in 100ul of 1xPBS.

2) 4 µL of 1 mM TCO-PEG4-TFP Ester was added to the cells, mixed by pipetting and

incubated for 5 min @RT protected from light

3) 6 µL of each methyltetrazine oligo amide (prepared in Part 1) was added, mixed by

pipetting and incubated for 30 min @RT protected from light

4) Methyltetrazine-PEG4-Amine was added for final concentration of 50 µM and Tris HCl was

added to 10 mM final concentration and incubated for 5 min @RT

5) Cells were then diluted ~twofold with 1X PBS, spun down and washed 3x in 1xPBS

6) Cells were resuspended in 70% EtOH in 1xPBS, and kept -20C for up to 6 months after

the initial cell collection date.

Cell plating for imaging

Cells stored in suspension at -20 C were resuspended by pipetting. Barcoded cells were

combined at desired ratios in a single tube. A small volume of cells (~5ul of cells at 10k/ul-20k/ul

concentration) was spotted on a poly-lysine coated area in the center of the slide and let dry for

3-5 min. 1xPBS was then added and the slide was inspected to check for optimal cell density

(ideally monolayer of cells evenly covering the entire slide). Cells were then used for IF or ORCA.

ORCA primary probe hybridization

For ORCA DNA experiments, we used the protocol described in Mateo et al. 2019 and in detail

in Mateo et al. 2020. Briefly, cells were fixed in 4% PFA in 1x PBS for 10 min. After cell plating,

cells were permeabilized with 0.5% Triton-X in 1xPBS for 10min, followed by 2 washes with

1xPBS. Cells were incubated in 0.1M HCl for 5 min followed by 3 washes in 1xPBS. We then

treated cells for 30min with RNAse A (10ug/ml) at 37C followed by 3 washes in 2xSSC. Cells

were then incubated for 35 min in Hybridization buffer (2xSSC, 50% formamide and 0.1% Tween).

Primary probes (for 2-3 Mb Hoxa and Sox2 probes, we used 15ug of probe, for 30Mb and entire

chromosome tiling we used 3-4 ug of probe) diluted in hybridization solution (2xSSC, 50%

formamide, 10% dextran sulfate, 0.1% Tween) were then added onto cells, covered with a

coverglass, denatured for 3min at 90C and incubated overnight at 42C. The following day, cells

were washed with 2xSSC, postfixed in 2%GA, 8% PFA in 1xPBS for 30min to 1h and set up on

the microscope for sequential labeling and imaging. If ORCA was preceded by IF, we picked

positions to match as closely as possible to cells imaged by IF. Cell barcodes were either imaged

before or after probe barcodes.

RNA labeling was done as described above for DNA with the exception of the HCl, RNAse A or

post fix steps.

ORCA imaging

Samples hybridized with primary probes were imaged on the custom microscopy and microfluidics

setup as previously described [5,78]. Briefly, we used oligos complementary to the readout barcodes

on primary probes carrying either a Cy5 or a 750 dye. We used strand displacement to remove

an imaged barcode. We also used a Cy3 oligo to label all probes and imaged this Cy3 fiducial

channel at each round of imaging. The fiducial spots were subsequently used for spot calling and

registration in the image analysis pipeline.

Immunofluorescence followed by ORCA

Plated cells were permeabilized and blocked in antibody dilution buffer (2%BSA, 0.1% Triton X -

100 in 1xPBS) for 10 min @RT. Cells were then incubated with the Geminin primary antibody

(ab195047, 1:100) diluted in antibody dilution buffer for 1 h @ RT, washed 3x with 1xPBS,

incubated with secondary antibody (anti-rabbit 647 Invitrogen A31573) and DAPI diluted in

antibody dilution buffer for 30 min @RT, washed 3x with 1xPBS after which they were ready to

image.

Loop extrusion simulations

Loop extrusion simulations were performed using the "polychrom" package from "open2c" github

project [79], which was developed from earlier polymer models of loop extrusion written by the Mirny

lab [21,30], and powered by the GPU accelerated molecular simulation toolkit openMM [80]. All

simulations were run on an NVIDIA Titan Xp card. polychrom uses a Langevin approach to

simulate the dynamics of a polymer under several user defined energy constraints. Complete

details of the energetic constraints and other parameters are specified in the python simulation

scripts   included   in   our   github   repository   for   this   project:

https://github.com/BoettigerLab/Hafner2022/, and described in brief below. These can be run

independently using polychrom, which may be downloaded from the polychrom github page:

https://https://github.com/open2c/polychrom.

We simulated the chromosomes as 6 polymers confined in a spherical geometry representing the

nucleus. Each polymer was 5,000 monomers long, corresponding to 30 kb/monomer for 150 Mb

chromosomes, on par with the probe size used at our 3 Mb, 30 Mb, and whole-chromosome (150

Mb) experiments. Each simulation was run in 600 independent replicates, for sufficient time that

the final 3D structures were uncorrelated with the starting structure at all length scales and that

the pattern of TADs and loops could be reproduced from averaging single replicates over time

identically as to averaging over replicates (see below for loop extrusion and TADs). For high

resolution experiments, we conducted additional simulations, consisting of 2 polymers

constrained in a spherical volume, each 50,000 monomers long (3 kb/monomer). These

simulations ran substantially more slowly, and even after running 10 fold longer the structure at

the very longest length scales was not completely uncorrelated from the initialization conditions,

though the structures at shorter length scales had reached a stationary distribution. For these

reasons, we used these high resolution simulations only to model the submegabase and not

whole chromosome organization.

Loop extrusion was simulated using the previously described sequential 1D and 3D simulations,

implemented by *polychrom.* Briefly, the 1D simulations simulate the loading, unloading, and

motion of loop extruders with left and right arms that walk bi-directionally on chromatin and

interactions with extrusion blockers. These are followed by 3D simulations, in which short

harmonic bonds between the left and right arms of the extrusion factors impose the loop. Loop

extrusion blocking sites, simulating CTCF, were distributed in the pattern shown in Fig 6D. This 3

Mb pattern was replicated back to back to produce the 150 Mb chromosome pattern. Cohesins

were loaded with an average spacing of 300 kb, and a lifetime of corresponding to a maximum

extrusion length of 600 kb if not stalled. The confinement radius for the nucleus was chosen so

20% of the nuclear volume was filled with polymer. Confinement volumes 0.02% were used to

represent the unconstrained conditioned and 0.2 the 10% of relaxation volume condition. A hard-

sphere repulsion energy of 6 was used in all simulations except simulating a higher degree of

chain crossing, which used a value of 3. No additional attractive or repulsive interactions between

monomers was used in the simulations, which do not attempt to recapitulate the A/B

compartmentalization. See code available on github link above for details.

It should be emphasized that these are coarse-grained simulations, intended to capture the

qualitative, emergent properties of a small number of physical processes – such as loop extruders

moving on a large polymer in a confined volume. The utility of these coarse grained physics

models is not to capture as many features of reality as possible in a common model, but rather to

reproduce key patterns in the experimental data from a minimal mechanistic hypothesis [81,82]. By

using as few features from reality as possible in the model rather than as many as possible, we

can develop deeper intuition about the underlying processes. As a consequence of coarse-

graining abstraction, the precise relation between simulation units, monomers to kb, cohesin

spacing to kb, monomer radii to nanometer, etc, are approximate. We did not undertake a

comprehensive sweep of the parameter space of the model. The intensive computational time for

the thousands of independent simulations limit such analysis, and showing that the multiscale

qualitative effects of cohesin chromatin folding can be captured by loop extrusion did not require

it.

Image processing and quantification

<u>Image processing and spot calling for ORCA data</u>

Image processing (drift correction and localization of spots) analysis was performed as described

in [5,78].

<u>Cell segmentation</u>

We used the CellPose [83] for segmenting individual nuclei.

<u>Geminin IF and cell size quantification</u>

For Geminin IF quantification and the quantification of nuclear size, we performed all calculations

on the experiments for the Hoxa and Sox2 domains for 2.2 and 2.8 Mb respectively where cells

were stained for Dapi and Geminin prior to the ORCA experiments. Nuclear segmentation was

done on Dapi staining using CellPose. The resulting cell masks were then used to quantify levels

of Geminin immunofluorescence by computing the mean intensity per nucleus. The Geminin low

population was defined as the lowest 25% of the geminin expression level across all stained cells

within the experiment (combined the E14 parental cell line together with CTCF-AID and Rad21-

AID auxin treated and untreated cells). For nuclear area quantification, we used the total number

of pixels in each segmented mask in this experiment.

Demultiplexing cell barcodes

For each fiducial spot, we quantified the intensity of each cell barcode signal. To normalize for

differences in brightness across cell barcodes, we then balanced the intensities across all values.

We used the nearest neighbors approach to classify each spot into groups based on the values

for all cell barcodes for that spot.

RNA FISH data analysis

RNA FISH data was quantified similarly to [5]. Briefly, images were maximum-z-projected and

flatfield corrected. Foci corresponding to single mRNAs were identified by using a local-maxima

search with manually defined thresholds. Foci positions were then overlaid with cell segmentation

masks from CellPose to compute single-cell transcript mRNA counts.

ORCA analyses

Merge and filtering of ORCA data

For the ~3 Mb scale data, we had 2 technical replicates (performed on the same batch of collected

and barcoded cells) and a biological replicate (independently thawed, treated, fixed and barcoded

cells). Replicates showed a high degree of reproducibility and showed identical patterns of TADs

and loops and were comparable in median distance (Supplement 1). We thus merged the data

for all subsequent analyses. Barcodes with low hybridization efficiency (< 10 %) were filtered on

the merged data. An additional filter was done on a dataset basis to remove obvious artifacts such

as failed strand displacement events.

Insulation score

Insulation score was calculated on contact frequency maps. We used a window size of 8 barcodes

(8 * 30 kb steps size = 240 kb). For each position along the diagonal, we calculated the sum of

contacts in the upstream and downstream window. To be able to compare values across

conditions we then divided the values by the mean insulation score in the untreated condition.

<u>Three-way contact analyses</u>

For the three-way CTCF contact analyses in Fig 2, we picked the 3 CTCF sites for *Hoxa* (barcodes

20, 34, 54) and *Sox2 regions* (barcodes 42, 46, 70). We calculated the frequency of

chromosomes where the distance between sites 1-2 and 2-3 were both below 200 nm. This

frequency was normalized for detection efficiency for these pairs. We then shifted barcode

positions 8 steps (with each step= 30 kb) in each direction and performed the same calculation

for each shift. The expected frequencies were generated by multiplying the frequency for the pairs

(frequency(1,2) * frequency (2-3)) and dividing by their detection efficiency.

We performed bootstrapping to estimate the confidence intervals for the contact frequencies.

Bootstrapping was done by randomly sampling with replacement from the data. The height bars

in the barplot (Fig. 2, Supplement 3) correspond to the mean and the error bars correspond to the

25th and 75th quantiles. Thus, errorbars that do not overlap reflect contact-frequencies that were

distinct in 94% of resamplings ($94 = 1- 0.25^2$). We used 200 resampling draws in our

bootstrapping, and noted more resamplings made little difference on the spread.

<u>Compartment analysis using Pearson's Matrix</u>

Compartmentalization was analyzed in the ORCA data starting from the median pairwise distance

matrices, **O**, for both the 30 Mb scale and whole chromosome (150 Mb) data. The "expected"

distance was computed by averaging the 3D distance for all loci pairs which had the same

genomic spacing. This produces a matrix, **E**, with a smooth decay from the main diagonal. We

then divided the observed distances by the expected distances to create a new matrix, **N**, $N_{i,j}$ =

$O_{i,j}/E_i$. Finally, we computed the Pearson Matrix **P**, such that $P_{i,j}$ is Pearson's correlation

coefficient between row *i* and column *j* of matrix **N**. This parallels the original analysis of

compartmentalization described for Hi-C, though using our distance measures in place of the

contact-frequency per bin [38].

We note that the Hi-C and ORCA compartment analysis were computed with an important

difference. The Hi-C compartment analysis at the 30 and 150 Mb scale uses reads from every

part of the 30Mb (150 Mb) interval. In contrast, the ORCA data looks exclusively at particular 30

kb elements that skip through that domain at 1 Mb or 5 Mb intervals. Thus, substantial parts of

the genome driving the correlation signal in the Hi-C data are not included. We note that our

calculation using Hi-C data that focused only on the interactions between the 30 kb windows

covered by the ORCA experiments were too sparsely populated to permit detection of

compartments. Expanding these to 120 kb bins resulted in only minor improvement, and thus we

elect to show the Hi-C compartmentalization computed using all the data. Thus, some qualitative

differences in the position/distribution of the compartment boundaries likely reflects the

contributions of genomic sequences not covered in our imaging experiments, though the sparsity

of the Hi-C data makes this difficult to quantify precisely.

Published genomic data

<u>Hi-C data</u>

We examined Hi-C experiments from [16,19] for CTCF degradation, from [18] for Rad21 degradation,

from [84] for untreated cells as these were all generated in mESCs. For Nora et. al and Rhodes et

al. datasets, we loaded the processed Hi-C .cool files from either GEO (GSE98671) or

ArrayExpress (E-MTAB-7816) respectively. We used Cooler [85] to export normalized tables directly

using the cooler 'dump' command for the coordinates that match our probes. For the Kubo et al.

dataset, we loaded the processed .hic files from GEO (GSE94452), uploaded the .hic files to

Juicebox [86] and exported the normalized tables. For Bonev et al. dataset, we exported the

normalized tables for the regions of interest directly from Juicebox. We then downsampled the

data by averaging bins to match the resolution of the probes used for ORCA (30kb for the ~2.5Mb

scale, 1Mb for the ~30Mb scale and 5Mb for the ~150Mb scale probes).

<u>ChIP data</u>

We used CTCF and Ring1b ChIP-seq data from [84] downloaded from GEO in the bigwig format

(GSE96107_ES_Ring1B.bw and GSE96107_ES_CTCF.bw), converted to bedGraph format

using bigWigToBedGraph program. We then mapped the genomic coordinates with coordinates

for the Hoxa and Sox2 probes. For the CTCF density plot, we averaged the signal for each

barcode (spanning 30kb regions).

Acknowledgements

We thank Max Imakaev and Neil Chowdhury for code sharing and feedback on the polymer simulations; the 4DN community for support and the opportunities to present this work; Andrea Cosolo (4DN) for help with formatting and uploading the data to the 4DN server; S. Murphy for support and comments on the text and Figures; current and former members of the Boettiger lab for comments and discussion. This work was supported by NIH grant U01DK127419, the Packard Foundation, and a Beckman Young Investigator's Award to A.N.B. M.P. was supported by a postdoctoral fellowship from the Damon Runyon Cancer Research Foundation. A.H. was supported by Walter and Idun V. Berry Postdoctoral fellowship. S.E.B. was supported by the Stanford Graduate Fellowship. A.H. and A.N.B. designed the experiments. A.H., M.P. and S.E.B. performed the experiments. E.N. contributed key reagents. A.H. and A.N.B. performed analysis and interpreted the results. A.H. and A.N.B., wrote the manuscript with input from E.N. S.E.B. and M.P.

Conflicts of interest

Authors declare no competing interests.

Data and code availability

All probe coordinates and data tables are in the process of being annotated and formatted for upload on the 4DN public access server. Probe libraries used in this work can be viewed through the UCSC genome browser: https://genome.ucsc.edu/s/toniai/mm10_Hafner2022

Code to reproduce the simulations described in this work is available at https://github.com/BoettigerLab/Hafner2022/. These simulations require the polychrom

754 simulation tools, available at: https://https://github.com/open2c/polychrom. Additional information

755 on the simulations is included in the **Methods**.

References

1. McCord, R. P., Kaplan, N. & Giorgetti, L. Chromosome Conformation Capture and Beyond:

Toward an Integrative View of Chromosome Structure and Function. *Mol. Cell* **77**, 688–708

(2020).

2. Mirny, L. & Dekker, J. Mechanisms of Chromosome Folding and Nuclear Organization:

Their Interplay and Open Questions. *Cold Spring Harb. Perspect. Biol.* (2021)

doi:10.1101/cshperspect.a040147.

3. Cavalheiro, G. R., Pollex, T. & Furlong, E. E. To loop or not to loop: what is the role of

TADs in enhancer function and gene regulation? *Curr. Opin. Genet. Dev.* **67**, 119–129

(2021).

4. Jerković, I., Szabo, Q., Bantignies, F. & Cavalli, G. Higher-Order Chromosomal Structures

Mediate Genome Function. *J. Mol. Biol.* **432**, 676–681 (2020).

5. Mateo, L. J. *et al.* Visualizing DNA folding and RNA in embryos at single-cell resolution.

Nature **568**, 49–54 (2019).

6. Sexton, T. *et al.* Three-dimensional folding and functional organization principles of the

Drosophila genome. *Cell* **148**, 458–472 (2012).

7. Dixon, J. R. *et al.* Topological domains in mammalian genomes identified by analysis of

chromatin interactions. *Nature* **485**, 376–380 (2012).

8. Nora, E. P. *et al.* Spatial partitioning of the regulatory landscape of the X-inactivation

centre. *Nature* **485**, 381–385 (2012).

9. Dixon, J. R., Gorkin, D. U. & Ren, B. Chromatin Domains: The Unit of Chromosome

Organization. *Mol. Cell* **62**, 668–680 (2016).

10. Dekker, J. & Mirny, L. A. The 3D Genome as Moderator of Chromosomal Communication.

Cell **164**, 1110–1121 (2016).

11. Rowley, M. J. & Corces, V. G. Organizational principles of 3D genome architecture. *Nat.*

Rev. Genet. **19**, 789–800 (2018).

12. Dekker, J. & Heard, E. Structural and functional diversity of Topologically Associating

Domains. *FEBS Lett.* **589**, 2877–2884 (2015).

13. Ibrahim, D. M. & Mundlos, S. The role of 3D chromatin domains in gene regulation: a multi-

facetted view on genome organization. *Curr. Opin. Genet. Dev.* **61**, 1–8 (2020).

14. Ibrahim, D. M. & Mundlos, S. Three-dimensional chromatin in disease: What holds us

together and what drives us apart? *Curr. Opin. Cell Biol.* **64**, 1–9 (2020).

15. Jerkovic´, I. & Cavalli, G. Understanding 3D genome organization by multidisciplinary

methods. *Nat. Rev. Mol. Cell Biol.* 1–18 (2021).

16. Kubo, N. *et al.* Promoter-proximal CTCF binding promotes distal enhancer-dependent gene

activation. *Nat. Struct. Mol. Biol.* (2021) doi:10.1038/s41594-020-00539-5.

17. Rao, S. S. P. *et al.* Cohesin Loss Eliminates All Loop Domains. *Cell* **171**, 305–320.e24

(2017).

18. Rhodes, J. D. P. *et al.* Cohesin Disrupts Polycomb-Dependent Chromosome Interactions in

Embryonic Stem Cells. *Cell Rep.* **30**, 820–835.e10 (2020).

19. Nora, E. P. *et al.* Targeted Degradation of CTCF Decouples Local Insulation of

Chromosome Domains from Genomic Compartmentalization. *Cell* **169**, 930–944.e22

(2017).

20. Bintu, B. *et al.* Super-resolution chromatin tracing reveals domains and cooperative

interactions in single cells. *Science* **362**, (2018).

21. Fudenberg, G. *et al.* Formation of Chromosomal Domains by Loop Extrusion. *Cell Rep.* **15**,

2038–2049 (2016).

22. Sanborn, A. L. *et al.* Chromatin extrusion explains key features of loop and domain

formation in wild-type and engineered genomes. *Proc. Natl. Acad. Sci. U. S. A.* **112**,

E6456–65 (2015).

23. Luppino, J. M. *et al.* Cohesin promotes stochastic domain intermingling to ensure proper
regulation of boundary-proximal genes. *Nat. Genet.* **52**, 840–848 (2020).

24. Szabo, Q. *et al.* Regulation of single-cell genome organization into TADs and chromatin
nanodomains. *Nat. Genet.* (2020) doi:10.1038/s41588-020-00716-8.

25. Liu, M. *et al.* Multiplexed imaging of nucleome architectures in single cells of mammalian
tissue. *Nat. Commun.* **11**, 2907 (2020).

26. Su, J.-H., Zheng, P., Kinrot, S. S., Bintu, B. & Zhuang, X. Genome-Scale Imaging of the 3D
Organization and Transcriptional Activity of Chromatin. *Cell* (2020)
doi:10.1016/j.cell.2020.07.032.

27. Takei, Y. *et al.* Integrated spatial genomics reveals global architecture of single nuclei.
Nature (2021) doi:10.1038/s41586-020-03126-2.

28. Gehring, J., Hwee Park, J., Chen, S., Thomson, M. & Pachter, L. Highly multiplexed single-
cell RNA-seq by DNA oligonucleotide tagging of cellular proteins. *Nat. Biotechnol.* **38**, 35–
38 (2020).

29. Rao, S. S. P. *et al.* A 3D map of the human genome at kilobase resolution reveals
principles of chromatin looping. *Cell* **159**, 1665–1680 (2014).

30. Nuebler, J., Fudenberg, G., Imakaev, M., Abdennur, N. & Mirny, L. A. Chromatin
organization by an interplay of loop extrusion and compartmental segregation. *Proceedings*
of the National Academy of Sciences **115**, E6697–E6706 (2018).

31. Nolen, L. D., Boyle, S., Ansari, M., Pritchard, E. & Bickmore, W. A. Regional chromatin
decompaction in Cornelia de Lange syndrome associated with NIPBL disruption can be
uncoupled from cohesin and CTCF. *Hum. Mol. Genet.* **22**, 4180–4193 (2013).

32. Sofueva, S. *et al.* Cohesin-mediated interactions organize chromosomal domain
architecture. *EMBO J.* **32**, 3119–3129 (2013).

33. Zuin, J. *et al.* Cohesin and CTCF differentially affect chromatin architecture and gene
expression in human cells. *Proc. Natl. Acad. Sci. U. S. A.* **111**, 996–1001 (2014).

34. Luppino, J. M. *et al.* NIPBL and WAPL balance cohesin activity to regulate chromatin

folding and gene expression. *bioRxiv* 2022.04.19.488785 (2022)

doi:10.1101/2022.04.19.488785.

35. Gabriele, M. *et al.* Dynamics of CTCF- and cohesin-mediated chromatin looping revealed

by live-cell imaging. *Science* **376**, 496–501 (2022).

36. Mach, P. *et al.* Live-cell imaging and physical modeling reveal control of chromosome

folding dynamics by cohesin and CTCF. doi:10.1101/2022.03.03.482826.

37. Schwarzer, W. *et al.* Two independent modes of chromatin organization revealed by

cohesin removal. *Nature* **551**, 51–56 (2017).

38. Lieberman Aiden, E. *et al.* Comprehensive mapping of long-range interactions reveals

folding principles of the human genome. *Science* **326**, 289–293 (2009).

39. Wang, S. *et al.* Spatial organization of chromatin domains and compartments in single

chromosomes. *Science* **353**, 598–602 (2016).

40. Mirny, L. A. Cells use loop extrusion to weave and tie the genome. *Nature* **590**, 554–555

(2021).

41. Polovnikov, K., Belan, S., Imakaev, M., Brandão, H. B. & Mirny, L. A. Fractal polymer with

loops recapitulates key features of chromosome organization. *bioRxiv* 2022.02.01.478588

(2022) doi:10.1101/2022.02.01.478588.

42. Li, Y. *et al.* CRISPR reveals a distal super-enhancer required for Sox2 expression in mouse

embryonic stem cells. *PLoS One* **9**, e114485 (2014).

43. Zhou, H. Y. *et al.* A Sox2 distal enhancer cluster regulates embryonic stem cell

differentiation potential. *Genes Dev.* **28**, 2699–2711 (2014).

44. Hsieh, T.-H. S. *et al.* Enhancer-promoter interactions and transcription are maintained upon

acute loss of CTCF, cohesin, WAPL, and YY1. doi:10.1101/2021.07.14.452365.

45. Aljahani, A. *et al.* Analysis of sub-kilobase chromatin topology reveals nano-scale

regulatory interactions with variable dependence on cohesin and CTCF. *Nat. Commun.* **13**,

2139 (2022).

46. Liu, N. Q. *et al.* WAPL maintains a cohesin loading cycle to preserve cell-type-specific distal

gene regulation. *Nat. Genet.* **53**, 100–109 (2021).

47. Bates, L. E., Alves, M. R. P. & Silva, J. C. R. Auxin-degron system identifies immediate

mechanisms of OCT4. *Stem Cell Reports* **16**, 1818–1831 (2021).

48. Jaeger, M. G. *et al.* Selective Mediator dependence of cell-type-specifying transcription.

Nat. Genet. **52**, 719–727 (2020).

49. Weber, C. M. *et al.* mSWI/SNF promotes Polycomb repression both directly and through

genome-wide redistribution. *Nat. Struct. Mol. Biol.* **28**, 501–511 (2021).

50. Muhar, M. *et al.* SLAM-seq defines direct gene-regulatory functions of the BRD4-MYC axis.

Science **360**, 800–805 (2018).

51. Lupiáñez, D. G. *et al.* Disruptions of Topological Chromatin Domains Cause Pathogenic

Rewiring of Gene-Enhancer Interactions. *Cell* **161**, 1012–1025 (2015).

52. Flavahan, W. A. *et al.* Insulator dysfunction and oncogene activation in IDH mutant gliomas.

Nature 1–16 (2015).

53. Hnisz, D. *et al.* Activation of proto-oncogenes by disruption of chromosome neighborhoods.

Science **351**, 1454–1458 (2016).

54. Hnisz, D., Day, D. S. & Young, R. A. Insulated Neighborhoods: Structural and Functional

Units of Mammalian Gene Control. *Cell* **167**, 1188–1200 (2016).

55. Wang, X. Q. D. *et al.* Three-dimensional regulation of HOXA cluster genes by a cis-element

in hematopoietic stem cell and leukemia. *bioRxiv* 2020.04.16.017533 (2020)

doi:10.1101/2020.04.16.017533.

56. Wilderman, A. *et al.* A distant global control region is essential for normal expression of

anterior HOXA genes during mouse and human craniofacial development. *bioRxiv*

2022.03.10.483852 (2022) doi:10.1101/2022.03.10.483852.

57. Huang, H. *et al.* CTCF mediates dosage- and sequence-context-dependent transcriptional

insulation by forming local chromatin domains. *Nat. Genet.* 1–11 (2021).

58. Chakraborty, S. *et al.* High affinity enhancer-promoter interactions can bypass

CTCF/cohesin-mediated insulation and contribute to phenotypic robustness. *bioRxiv*

2021.12.30.474562 (2022) doi:10.1101/2021.12.30.474562.

59. Montavon, T. *et al.* A Regulatory Archipelago Controls Hox Genes Transcription in Digits.

Cell **147**, 1132–1145 (2011).

60. Berlivet, S. *et al.* Clustering of Tissue-Specific Sub-TADs Accompanies the Regulation of

HoxA Genes in Developing Limbs. *PLoS Genet.* **9**, e1004018 (2013).

61. Guo, Y. *et al.* CTCF/cohesin-mediated DNA looping is required for protocadherin promoter

choice. *Proceedings of the National Academy of Sciences* **109**, 21081–21086 (2012).

62. Huang, P. *et al.* Comparative analysis of three-dimensional chromosomal architecture

identifies a novel fetal hemoglobin regulatory element. *Genes and Development* **31**, 1704–

1713 (2017).

63. Allahyar, A. *et al.* Enhancer hubs and loop collisions identified from single-allele topologies.

Nat. Genet. 1 (2018).

64. Knoch, T. A. *et al.* The detailed 3D multi-loop aggregate/rosette chromatin architecture and

functional dynamic organization of the human and mouse genomes. *Epigenetics Chromatin*

9, 58 (2016).

65. Oudelaar, A. M. *et al.* Single-allele chromatin interactions identify regulatory hubs in

dynamic compartmentalized domains. *Nat. Genet.* **50**, 1744–1751 (2018).

66. Kubo, N. *et al.* Preservation of Chromatin Organization after Acute Loss of CTCF in Mouse

Embryonic Stem Cells. *bioRxiv* 118737 (2017).

67. Ortabozkoyun, H. *et al.* CRISPR and biochemical screens identify MAZ as a cofactor in

CTCF-mediated insulation at Hox clusters. *Nat. Genet.* **54**, 202–212 (2022).

68. de Laat, W. & Duboule, D. Topology of mammalian developmental enhancers and their

regulatory landscapes. *Nature* **502**, 499–506 (2013).

69. Wang, W., Ren, G., Hong, N. & Jin, W. Exploring the changing landscape of cell-to-cell

variation after CTCF knockdown via single cell RNA-seq. *BMC Genomics* **20**, 1015 (2019).

70. Ren, G. *et al.* CTCF-Mediated Enhancer-Promoter Interaction Is a Critical Regulator of Cell-

to-Cell Variation of Gene Expression. *Mol. Cell* **67**, 1049–1058.e6 (2017).

71. Chiang, M. *et al.* Gene structure heterogeneity drives transcription noise within human

chromosomes. *bioRxiv* 2022.06.09.495447 (2022) doi:10.1101/2022.06.09.495447.

72. Cuartero, S. *et al.* Control of inducible gene expression links cohesin to hematopoietic

progenitor self-renewal and differentiation. *Nat. Immunol.* **19**, 932–941 (2018).

73. Pauli, A. *et al.* Cell-Type-Specific TEV Protease Cleavage Reveals Cohesin Functions in

Drosophila Neurons. *Dev. Cell* **14**, 239–251 (2008).

74. Pauli, A. *et al.* A direct role for cohesin in gene regulation and ecdysone response in

Drosophila salivary glands. *Curr. Biol.* **20**, 1787–1798 (2010).

75. Cummings, C. T. & Rowley, M. J. Implications of Dosage Deficiencies in CTCF and

Cohesin on Genome Organization, Gene Expression, and Human Neurodevelopment.

Genes **13**, (2022).

76. Avagliano, L. *et al.* Chromatinopathies: A focus on Cornelia de Lange syndrome. *Clin.*

Genet. **97**, 3–11 (2020).

77. Di Nardo, M., Pallotta, M. M. & Musio, A. The multifaceted roles of cohesin in cancer. *J.*

Exp. Clin. Cancer Res. **41**, 96 (2022).

78. Mateo, L. J., Sinnott-Armstrong, N. & Boettiger, A. N. Tracing DNA paths and RNA profiles

in cultured cells and tissues with ORCA. *Nat. Protoc.* **16**, 1647–1713 (2021).

79. *polychrom: Polymer simulations of chromosomes and generating 'in silico' Hi-C maps.*

(Github).

80. Eastman, P. *et al.* OpenMM 7: Rapid development of high performance algorithms for

molecular dynamics. *PLoS Comput. Biol.* **13**, e1005659 (2017).

81. A Conversation with Leonid Mirny. *Cold Spring Harb. Symp. Quant. Biol.* **82**, 403–405

(2017).

82. Igoshin, O. A. *et al.* Biophysics at the coffee shop: lessons learned working with George

Oster. *Mol. Biol. Cell* **30**, 1882–1889 (2019).

83. Stringer, C., Wang, T., Michaelos, M. & Pachitariu, M. Cellpose: a generalist algorithm for

cellular segmentation. *Nat. Methods* **18**, 100–106 (2021).

84. Bonev, B. *et al.* Multiscale 3D Genome Rewiring during Mouse Neural Development. *Cell*

171, 557–572.e24 (2017).

85. Abdennur, N. & Mirny, L. A. Cooler: scalable storage for Hi-C data and other genomically

labeled arrays. *Bioinformatics* **36**, 311–316 (2020).

86. Robinson, J. T. *et al.* Juicebox.js Provides a Cloud-Based Visualization System for Hi-C

Data. *Cell Syst* **6**, 256–258.e1 (2018).

Figures and figure legends

Figure 1 | A multiplexed approach for chromosome tracing. a. Schematic depiction of two ways that preferential contacts within TADs can be lost upon degradation of CTCF or cohesin. **b**. Schematic depiction of the multiplexed ORCA workflow. **c**. Example of a field of view with pseudo-colored cell barcode labels for all imaged conditions: mES parental E14 cell line, CTCF-AID and RAD21-AID auxin-treated (+Aux) or untreated (NT) cells. Fiducial signal (labeling all probes in the Sox2 2.8 Mb domain), is shown as well as an example trace of a single chromosome for the Sox2 domain. **d**. Contact frequency maps from the demultiplexed cell conditions for both the *Hoxa* (2.2 Mb) and *Sox2* (2.8 Mb) regions imaged in 30 kb steps resulting in 73 steps for *Hoxa* and 93 steps for *Sox2*.

Figure 2 | CTCF enables cross-TAD interactions. a. Schematic depiction of bridges (left) and schematic hypothesis how these arise from a mixed population containing either the bridge contact or the domain separation. **b**. Contact frequency map from all cells with the bridging contact indicated (black circle). **c**. Subtraction of contact frequency maps from all cells that have the indicated bridging contact minus those without it. **d**. Schematic depiction showing how CTCF interactions create a radially organized chromosome. Near the center axis (gray region) CTCF sites (red) and CTCF-proximal regions make multi-way interactions. **e**. Examples of single chromosomes with CTCF-marked boundary positions colored in red. **f**. Top panels, contact frequency difference map between CTCF Aux and untreated cells, with arrows indicating the positions chosen for the 3-way analysis. Lower panels: Frequency of 3-way contact between the three CTCF-marked positions indicated by black arrows. Frequency of 3-way contact for other distance-matched triplets (black) is shown as a function of how far (in kb) they are shifted from the black triplet. In gray, expected 3-way contact based on the pairwise contact probabilities and in blue, frequencies of 3-way contact for CTCF auxin-treated condition. Error bars show confidence intervals (see Methods). **g**. Contact frequency difference map for CTCF auxin-treated and untreated cells. Black lines show TAD boundaries. **h**. Example chromosome traces for CTCF auxin-treated cells. **I**. Schematic illustrating the effect of CTCF loss on randomizing the position of CTCF-marked boundary sites.

Figure 3 | Cohesin loss leads to increased distances within and between TADs. a. Example single chromosome traces for *Hoxa* and *Sox2* regions from untreated and auxin-treated cells. **b**. Differences in median distance between auxin-treated and untreated cells show that all pairwise distances increase (all blue). TAD outlines are drawn in black. For the *Hoxa* region, the area indicated by the arrow corresponds to the *Hoxa* gene region that is coated by Polycomb and does not expand upon auxin treatment (white). **c**. Schematic illustrating that loss of cohesin leads to general expansion, increasing distances within and between TADs.

Figure 4 | Cohesin loss leads to increased mixing at the chromosome scale. a. Schematic of probes designed to

image folding of ~30 Mb and ~150 Mb regions of the genome, shown relative to the ~2.5 Mb probes imaged previously.

b. ORCA Pearson correlation maps for the 150-Mb scale probes in RAD21 untreated or auxin-treated cells. **c-d.**

Examples of single chromosome traces for ~30 Mb (c) or ~150 Mb (d) regions of chromosome 6 and chromosome 3

with or without auxin. Probes that correspond to consecutive (red and blue) and a more distal (cyan) genomic distances

are represented with larger spheres. Their interprobe distances are shown in the linear colorbar. **e-f.** Differences in

median distance between RAD21 auxin-treated and untreated cells for the ~30 Mb (e) and ~150 Mb (f) regions, show

that distal genomic regions are closer (red) after auxin treatment. The black squares in (e) show the position of the ~2.5

Mb domains imaged previously, while the black squares in (f) show the position of the ~30 Mb domains. **g.** Schematic

illustrating the disordered chromosome structure upon cohesin degradation. **h.** Example chromosome traces in

untreated or RAD21 depleted cells colored by compartment identity, as in (b). **i.** Schematic illustrating the effect of

cohesin degradation on compartments.

Fig. 5 | Loop extrusion model shows the multi-scale effects of cohesin loss. a. Simulated polymer corresponding

to 3 Mb in length with or without loop extrusion factors. The regions shown are a zoom-in view of the blue region inside

the black boxes in (b). **b**. Polymer simulations at the chromosome scale (150 Mb) with or without loop extrusion factors.

The chromosome is shown in gray. Other chromosomes inside the simulated nucleus are not shown to avoid overlap.

Individual 3 Mb domains are colored. Their positions on the linear polymer are indicated by the colorbar. **c.** Difference

in median distance between simulations with and without loop extrusion at different genomic scales: 3 Mb, 30 Mb and

150Mb. Zoom-out of the 3 Mb corresponds to a 200 kb domain and a few more distally spaced regions. **d**. Schematic

of the probe design for a 170 kb region encompassing the *Sox2* gene with probes spaced 5 kb apart, as well as 4

adjacent and more distally spaced regions, relative to the 2.8 Mb probeset used in Figs. 1-3. **e.** ORCA contact frequency

maps for finer-scale Sox2 probes in RAD21 untreated or auxin-treated cells. **f.** Difference in median distances between

RAD21 auxin-treated and untreated cells for the 2.8-Mb sized domain (same as in Fig. 3b and Fig. 4e, f) and the finer-

scale region. Squares on the 2.8 Mb map indicate the regions that were imaged at higher resolution.

1008

1009 **Figure 6 | Cohesin depletion increases the variability in gene expression a.** Example of multiplexed cells for

1010 different treatment conditions labeled by single molecule FISH. **b**. Mean fold change of expression. **c**. Fold change in

1011 coefficient of variation (COV).

1012

Extended Data Fig.1 | CTCF and cohesin can be depleted after auxin addition and reproducibility of ORCA replicates (related to Figure 1) a. Western blot labeling Actin (loading control) and either CTCF or Cohesin in the parental E14 cell line and in the CTCF-AID-eGFP or RAD21-AID-eGFP cells, with or without auxin treatment. **b**. Scatter plots of inter-barcode distances, in nanometers, comparing reproducibility among three replicates labeling the *Hoxa* or *Sox2* domains. Median distance is plotted for each barcode pair against the pairwise median distance in a different replicate. Pearson's correlation coefficient, r, is shown on the plot. The lower left plot shows the axis scale and the same scale was used for all plots.

Extended Data Fig. 2 | Published Hi-C data for the Hoxa and Sox2 domains qualitatively agrees with ORCA (related to Figure 1) a-b. Hi-C data[16,18,19] for the *Hoxa* and *Sox2* domains corresponding to regions imaged with ORCA. **c**. Insulation score calculation for *Hoxa* and *Sox2* domains done on contact frequency maps of CTCF untreated, CTCF auxin-treated and RAD21 auxin-treated conditions.

Extended Data Fig. 3 | "Bridges" in ORCA and Hi-C (related to Figure 2). a. Contact frequency maps from untreated cells. Circles denote visible loops that bridge across TAD boundaries. The heatmap on the left of the contact frequency maps shows the density of CTCF ChIP-seq signal. **b**. Hi-C contact frequency for untreated mESCs from Bonev et al. 2017 [84] showing the same bridging contacts as in (a). **c**. Contact frequency map for cells with different bridging contacts. **d**. Difference in contact frequency maps between cells that have the bridging contacts and cells that do not. **e**. Frequency of the 3-way contact between the positions corresponding to the CTCF-marked boundaries relative to positions not near a boundary, as in Fig. 2f.

Extended Data Fig. 4 | Cohesin depletion leads to expansion at the ~3 Mb scale (related to Figure 3). a. Median distance maps for the *Hoxa* and *Sox2* domains in untreated or auxin treated cells. **b**. Maps of P-values computed using ranksum for each pairwise distance between auxin treated and untreated cells. Arrow indicates the only regions other than the diagonal that did not significantly change upon auxin treatment. **c**. Log$_2$ ratio of median distance between auxin-treated and untreated cells. Black lines indicate TAD outlines and the black arrow indicates the region that did not change upon auxin treatment. **d**. Contact fraction maps for the *Hoxa* region and the corresponding Ring1B (member of the polycomb complex) ChIP signal from [84] as well as the position of *Hoxa* genes. Arrows point to Ring1B coated regions on the heatmaps. **e.** Log$_2$ ratio of Hi-C data between RAD21 auxin-treated and untreated cells shows a decrease in intra-TAD contacts and gain of contacts between TADs. **f.** Example field of view image of DAPI and immunofluorescent staining of Geminin and ORCA fiducial staining for the same cells. **g**. Difference in median distance

1044   between the Geminin-low population for the RAD21 auxin treated cells and all RAD21 untreated cells. **h**. Scatter plot

1045   of median distance (over all pairs) in each chromosome relative to Geminin levels in the same cell. Spearman's

1046   correlation coefficient is indicated for each condition.

1047

Extended Data Fig. 5 | Quantification of nuclear size and compartments after depletion of cohesin (related to

Figure 4) a. Example images of individual cells labeled with DAPI and the cell surface marker for the parental E14 cell

line, RAD21 NT or RAD21 Aux cells and the quantification of cell sizes across all cells (n=1729 cells for RAD21 NT,

n=7172 cells for RAD21 +Aux, n=3435 for E14). **b**. Hi-C [84] Pearson correlation maps for the 30 Mb and 150 Mb regions

corresponding to regions imaged by ORCA and the distribution of Pearson correlation coefficients. **c**. ORCA Pearson

correlation maps for the 30 Mb and 150 Mb probes in RAD21 untreated or auxin treated cells. **d.** Distribution of Pearson

correlation coefficients for Hi-C data. **e**. Distribution of Pearson correlation coefficients for ORCA data. **f**. P-values

computed using ranksum for each pair between auxin-treated and untreated cells.

1056

1057   **Extended Data Fig. 6 | Polymer simulations with lower nuclear confinement and increased strand crossing**

1058   **(related to Figure 5) a.** Contact frequency of simulated polymers with or without loop extrusion factors. **b**. p-values

1059   comparing simulated polymers with or without loop extrusion factors at different genomic scales. **c-d**. Median distance

1060   difference maps and p-values for simulations with or without loop extrusion factors at different scales for simulations

1061   with low density confinement (c) or with more strand crossing (d).

1062

1063   **Extended Data Fig. 7 | Additional ORCA data for the fine-scale Sox2 region (related to Figure 5) a.** Median

1064   distance and **b**. p-values for the *Sox2* 170 kb domain with genomically distant regions imaged by ORCA. **c**.Histogram

1065   of pairwise distance differences in median distance. In cyan are pairs that are <170 kb in genomic distance relative to

1066   >170 kb in dark blue.

1067

1068   **Extended Data Fig. 8 | CTCF depletion leads to less change in physical distance than cohesin depletion (related**

1069   **to Figure 6) a.** Median distance difference maps for CTCF Aux - CTCF NT cells at different scales of imaging. **b**. P-

1070   values computed using ranksum for each pair between CTCF auxin-treated and untreated cells at different genomic

1071   scales. **c**.Comparison of median distance differences between auxin-treated and untreated cells for either CTCF (blue)

1072   or RAD21 (red).

1073

Figure 1

Figure 2

Figure 3

a

RAD21 NT   RAD21 + Aux

Hoxa

Sox2

500 nm

b

Median distance RAD21Aux-NT

Hoxa

50.59   chr 6   52.78Mb

Sox2

33.39   chr 3   36.18Mb

-200

nm

-200

c

Cohesin

less within-TAD
and less
between-TAD
contact

Figure 4

Figure 5

a   Folding at the TAD scale

0.30 Mb   0.30 Mb

b   Folding at the chromosome scale

contracts

500 nm   expands

Linear position

c   Median distance difference: No extruders - with extruders

0.20 Mb   3 Mb   30 Mb   150 Mb

400   0   -200   nm

d

chr 3

34   1Mb   35   36 Mb

2.8 Mb (30kb steps)

20 kb - 20 kb - 170kb - 20 kb - 10 kb
(5 kb steps)

Sox2

f   Median distance RAD21Aux-NT

170 kb   2.8 Mb   32 Mb   151 Mb

200   0   -200   nm

34.60   34.77   33.39   36.18   18   49   5   155 Mb

chr 3

e

RAD21 NT   RAD21 Aux

0.6   0.4   0.2   Contact frequency

Sox2

SCR   n=2251   n=2347

34.60   34.77Mb

chr 3

Figure 6

Extended Data Fig. 1

a

b

Extended Data Fig. 2

Extended Data Fig. 3

Extended Data Fig. 4

a Median distance

RAD21 NT   RAD21 +Aux

Hoxa   n=5505   n=7462   nm   600 400 200
50.59 chr 6 52.78 Mb

Sox2   n=4094   n=8090   600 400 200
33.39 chr 3 36.18 Mb

b p-value

50.59 chr 6 52.78 Mb
0.05 1e-10 1e-20 1e-30 1e-40 <1e-50

33.39 chr 3 36.18 Mb
0.05 1e-10 1e-20 1e-30 1e-40 <1e-50

c ORCA Aux/NT

50.59 chr 6 52.78 Mb
1 0 -1 log2 (ratio)

33.39 chr 3 36.18 Mb
1 0 -1

e Hi-C Aux/NT

50.59 chr 6 52.78Mb
2 0 -2 log2 (ratio)

33.39 chr 3 36.18Mb
2 0 -2

d Contact frequency

Ring1b ChIP   HoxA genes   RAD21 NT   RAD21 +Aux
0.4 0.3 0.2 0.1
50.59 chr 6 52.78 Mb

f Measuring Geminin levels and ORCA in the same cells

Geminin   DAPI

IF   ORCA

g ORCA Geminin low
RAD21 Aux -NT

Hoxa   n=679   200 0 -200 nm
50.59 chr 6 52.78 Mb

Sox2   n=1165   200 0 -200
33.39 chr 3 36.18 Mb

h RAD21 NT n=677   RAD21 Aux n=2866   E14 n=1616
16 14 12 10   rho=-0.08   rho=0.04   rho=-0.01
200 600 1000

log2(Geminin)   RAD21 NT n=1052   RAD21 Aux n=4306   E14 n=1819
16 14 12 10   rho=-0.13   rho=0.03   rho=-0.04
200 600 1000   nm

Extended Data Fig. 5

a

🔬 Median distance

RAD21 NT   RAD21 Aux

n=2251   n=2347

600
400
200

nm

34.60   34.77Mb
chr3

b

p-value

0.05
1e-10
1e-20
1e-30
1e-40
<1e-50

34.60   34.77Mb
chr3

c

pairs ≤ 170 kb   pairs > 170 kb

47.6   187.6

60
40
20
0

# of pairs

0   100   200   300

Aux-NT [nm]

Extended Data Fig. 8

a

Median differences CTCF Aux -NT

b

p-values between CTCF Aux and CTCF NT distances

c

Distributions of median distance differences for CTCF vs RAD21 depletion