Modulatory dynamics of periodic and aperiodic activity in respiration-brain coupling

Daniel S. Kluger*[1,2], Carina Forster[3,4,5], Omid Abbasi[1], Nikos Chalas[1,2], Arno Villringer[3,4,6], Joachim Gross[1,2]

[1] Institute for Biomagnetism and Biosignal Analysis, University of Münster, Münster, Germany

[2] Otto Creutzfeldt Center for Cognitive and Behavioral Neuroscience, University of Münster, Münster, Germany

[3] Department of Neurology, Max Planck Institute for Human Cognitive and Brain Sciences, Leipzig, Germany

[4] Charité – Universitätsmedizin Berlin, Einstein Center for Neurosciences, Berlin, Germany

[5] Charité – Universitätsmedizin Berlin, Bernstein Center for Computational Neuroscience, Berlin, Germany

[6] Humboldt-Universität zu Berlin, Faculty of Philosophy, Berlin School of Mind and Brain, MindBrainBody Institute, Berlin, Germany

* corresponding author

Contact Info

Daniel S. Kluger

Institute for Biomagnetism and Biosignal Analysis, University of Münster

Malmedyweg 15, 48149 Münster

Germany

Telephone: 0049-251-8352543

E-mail: daniel.kluger@uni-muenster.de

Keywords

respiration, excitation/inhibition, 1/f slope, excitability, body-brain coupling, MEG, EEG

Abstract

Bodily rhythms such as respiration are increasingly acknowledged to modulate neural oscillations underlying human action, perception, and cognition. Conversely, the link between respiration and aperiodic brain activity – a non-oscillatory reflection of excitation-inhibition (E:I) balance – has remained unstudied. Aiming to disentangle potential respiration-related dynamics of periodic and aperiodic activity, we applied recently developed algorithms of time-resolved parameter estimation to resting-state MEG and EEG data from two labs (total N = 78 participants). We provide evidence that fluctuations of aperiodic brain activity (1/f slope) are phase-locked to the respiratory cycle, which suggests that spontaneous state shifts of excitation-inhibition balance are at least partly influenced by peripheral bodily signals. Moreover, differential temporal dynamics in their coupling to non-oscillatory and oscillatory activity raise the possibility of a functional distinction in the way each component is related to respiration. Our findings highlight the role of respiration as a physiological influence on brain signalling.

Introduction

Recent years have seen a surge in reports of respiration as a modulator of brain signalling - at least partly through modulation of rhythmic brain activity [1]. These respiration-modulated brain oscillations (RMBOs) have been localised within a widespread network across the cortex [2] and are increasingly being linked to changes in motor function [3,4], cognitive processing [5-7], and fundamental aspects of perception [8,9]. One candidate mechanism by which respiration (and other bodily signals such as the cardiac rhythm; see e.g. [10]) is thought to coordinate brain signalling is cortical excitability, a dynamic brain state indexing neural populations' readiness to become activated. This proposal rests on findings from intracranial animal studies demonstrating that respiration modulates spike rates in a variety of brain regions [11-13]. Moreover, one recent study [9] directly investigated respiratory modulation of excitability during low-level visual perception.

Critically, brain activity is not entirely oscillatory: One can distinguish its genuinely periodic, oscillatory component from an underlying stationary, aperiodic component typically showing a characteristic 1/f dependence of power and frequency (meaning that lower frequencies carry higher power [14]). In fact, dominance of low-frequency power is by no means just an epi-phenomenon of neural power spectra, but does reflect meaningful characteristics of brain function. Most relevantly, the 'steepness' of the aperiodic component has been shown to reflect the dynamics of excitation-inhibition (E:I) balance [15-17], with steeper 1/f slope indicating a stronger influence of inhibitory activity.

While sophisticated algorithms exist to disentangle periodic and aperiodic components of neural power spectra [18], time-resolved estimations of 1/f slope in particular have proven computationally expensive and lacked empirical validation. Consequently, the selective dynamics linking peripheral signals (such as respiration) to periodic and aperiodic neural activity are entirely unstudied. Meanwhile, the recent introduction of the SPRiNT toolbox [19] provides methodological advances that allow us to parameterize stationary and oscillatory brain activity dynamically over time. Here, we apply these algorithms in the realm of human body-brain interactions to test the hypothesis that 1/f slope as a marker of E:I balance is modulated by respiration.

Specifically, we used concurrent resting-state M/EEG and respiratory recordings in populations from two labs (Münster, Leipzig) to investigate distinct respiratory modulations of periodic and aperiodic brain activity. Our rationale was twofold: First, we aimed to characterise respiration phase-locked changes of aperiodic 1/f slope over parieto-occipital cortices to corroborate previous findings of excitability changes in visual perception [9]. Second, we extended the scope for a general investigation of potentially independent modulations of

aperiodic and oscillatory components within the cortical RMBO network [2]. Here, we particularly
aimed to assess whether both components of neural signalling would be modulated by
respiration and, if so, how these modulatory dynamics would be related to one another. Our
results highlight respiration as a slow physiological modulator of both periodic and aperiodic
cortical signalling, strengthening its previous implication in the regulation of excitability states.
Furthermore, the overall pattern of findings strongly suggests differential temporal dynamics
between the coupling of both components.

Results

We computed time-resolved fits of both aperiodic (1/f) and periodic (oscillatory) components
of resting-state brain activity to disentangle their dynamics in respiration-brain coupling. A
moving-window application of the novel SPRiNT algorithm [19] and the underlying specparam
procedure [18] to MEG and EEG data yielded individual time courses of aperiodic slopes and
periodic spectra for a total of N = 78 participants. Concurrent respiration recordings allowed
us to extract respiratory phase vectors that corresponded to each of the moving windows.
Next, we performed binning of respiratory phase and bin-wise averaging of corresponding 1/f
slope and periodic spectra on the group level (see Fig. 1 and Methods for details). We then
used these phase-specific estimates of slope and periodic spectra to statistically test if and
how periodic and aperiodic brain activity is modulated by respiration.

131
Fig. 1. Synopsis of acquired data and applied methods. a, In two labs, we simultaneously acquired nasal
respiration as well as eyes-open, resting-state MEG (IBB, Münster) and EEG data (MPI, Leipzig) in continuous 5-
min recordings. After preprocessing, single-sensor/-channel M/EEG data (middle panel) were subjected to the
SPRiNT algorithm [26] (top). Here, using a moving-window approach (window length = 1 s, 75% overlap between
neighbouring windows), spectral components of neural time series are estimated using a Fast Fourier transform.
These frequency-domain data are then parameterized using the specparam algorithm [18] which yields both aperiodic

and periodic components of neural activity in that time window. Repeating this procedure along the entire recording
thus yields time-resolved fits of the aperiodic 1/f slope as well as time-resolved periodic spectra ranging from 1 -
40 Hz (top right). Respiratory phase was computed using two-way interpolation (int) of the normalised raw
respiration signal (peak-to-trough, trough-to-peak; bottom). **b**, For each time point used as a moving-window centre
in the SPRiNT algorithm, we then extracted the corresponding respiratory phase. This allowed us to sort all time-
resolved slope fits and periodic spectra according to the respiratory phase at which they had been computed. In
keeping with previous work [2,9] we finally partitioned the respiration cycle into n = 60 equidistant, overlapping phase
bins and computed bin-wise averages of slope fits and periodic spectra. This approach thus yielded quasi-
continuous, respiration phase-resolved courses of both periodic and aperiodic components of brain activity for each
sensor/channel within each participant (bottom right).

Respiration relates to aperiodic neural activity over posterior cortices. The first analysis
was closely connected to a recent study showing respiration phase-dependent modulations of
cortical excitability in a visual detection task [9]. In line with a rich literature of visual perception
studies (see [20]), increased excitability was indexed by desynchronised alpha oscillations (i.e.,
lower alpha power corresponding to higher excitability). However, changes in alpha power can
be confounded by changes in the slope of aperiodic background activity [18] - itself a widely-
used measure of cortical excitability and excitation/inhibition balance [17]. Therefore, we aimed
to disambiguate the effects of respiration on periodic and aperiodic brain activity and started
by first testing the hypothesis that respiration-induced changes of excitability are reflected in
cyclic changes of the slope of aperiodic brain activity. For both MEG (Münster) and EEG data
(Leipzig), we set up a linear mixed-effects model (LMEM) to predict individual 1/f slopes of
parieto-occipital average power spectra by respiration phase. A null distribution of the resulting
beta weights was constructed by re-computing the LMEM based on individual surrogate
respiration time series k = 5000 times. Subsequently, we pooled both data sets in order to
capitalise on higher statistical power and to illuminate phase-locked modulations jointly found
across modalities. For both MEG and EEG as well as for the pooled data, the empirical group-
level beta weight for respiratory phase was significantly higher than the respective null beta
weights (all p < .001), strongly indicating a significant overall association between respiratory
phase and aperiodic neural signals over posterior cortices (Fig. 2). In order to characterise the
temporal dynamics of respiration phase-locked changes in 1/f slope, we repeated the binwise
slope estimation as outlined above, but this time on surrogate respiration time courses. For
each participant, we thus generated k = 5000 'null slope estimations' within each phase bin
which we used to extract percentiles for the empirical group-level bin averages relative to
these binwise 'null averages' (Fig. 2a; see Methods for details). For the pooled data from both
labs, we observed a steeper slope (indicating stronger inhibitory activity) around the
expiration-to-inspiration transition (ranging from -156° to 143° and around 131°). Conversely,
the slope was flatter (indicating stronger excitatory currents) during the inspiratory phase

(between -88° and -64°) and shortly after the inspiration-to-expiration transition (9° to 46°, see

Fig. 2). This general pattern was consistent across both data sets, although the flattening of

the slope during the inspiratory phase did not reach significance in the EEG data alone

(presumably due to slightly lower statistical power). Consistency across MEG and EEG data

was supported by a non-parametric two-sample test for circular data (akin to a Kruskal-Wallis

test for linear data), which showed that the circular means of MEG and EEG slope ~ respiration

courses were not significantly different from one another (P(77) = 0.21, p = .651).

Fig. 2. Group-level respiration-locked modulation of parieto-occipital 1/f slope. a, Respiration phase was
extracted for n = 60 overlapping phase bins. For each data set, we show combined LMEM beta weights for the
phase vector norm (light blue) against k = 5000 null betas (top right corners). Coloured bold dots show respiration
phase-dependent parieto-occipital 1/f slope. Radial scatter plots indicate null distributions of k = 5000 bin-wise
group-level mean exponent values. Solid black lines indicate the 5^{th} and 95^{th} percentile of each bin's null
distribution. In the MEG sample, 1/f slope was flattened during inspiration (around -70°) and around the inspiration-
to-expiration transition (-9° and 15° to 21°; blue markings). 1/f slope was steeper during expiration (82° to 88°) and
around the expiration-to-inspiration transition (-174° to 162°; red markings). The EEG sample showed flatter slope
after the inspiration-to-expiration transition (9° to 46° and around 58°) and steeper slope around the expiration-to-
inspiration transition (-162° to 131°). In = inspiration, ex = expiration. **b,** Overlay of ROI-average 1/f slope estimates
(black) and respiratory phase (green) for a single MEG participant. Grey lines show single-sensor slope estimates.
c, Neither breathing rate (top) nor depth (bottom) were significantly correlated with steepness of the aperiodic
component in MEG (left) or EEG (right; two-sided t-test against zero). **d,** Respiratory trace of a single participant
(left) and median group-level breathing depths (right) during normal and deep breathing (reanalysed from [3]).
Participants were breathing deeper during the deep (vs normal) breathing condition (z(27) = 6.19, p < .001, two-
sided Wilcoxon test for n = 28 participants). **e,** Individual ranges of slope ~ respiration phase were consistently
greater for deep (compared to normal) breathing (z(27) = 3.02, p = .003, two-sided Wilcoxon test). **f,** Comparison
of parieto-occipital 1/f slope ~ respiration phase during nasal (left) and oral breathing (right) from a follow-up control
study (see Methods). LMEMs confirmed a significant influence of respiratory phase on 1/f slope for both breathing
conditions (see histogram insets). **g,** Individual ranges of slope ~ respiration) did not differ between conditions
(z(25) = 0.80, p = .426, two-sided Wilcoxon test for n = 26 participants).
205
Aperiodic modulations as a function of respiratory parameters. We conducted different
control analyses to assess whether 1/f slope and its modulation strength covaried with
parameters like breathing rate, depth, or route (i.e., nasal vs oral breathing). First, in the
present data, individual 1/f slopes (averaged across the respiratory cycle) were not
significantly correlated with breathing rates (MEG: r(40) = .-17, p = .297; EEG: r(38) = .13, p
= .418) or breathing depths (MEG: r(40) = .19, p = .230; EEG: r(38) = -.09, p = .591) in either
data set (Fig. 2c). However, the interpretability of these results is somewhat limited by the
small amount of intraindividual variation in the respiratory time series - natural breathing is
simply too consistent to conclusively investigate whether different breathing 'modes' (e.g.,
deep vs normal breathing) would influence respiration-locked modulations in 1/f slope.
Therefore, we next re-analysed a previously published data set [3] in which we had recorded 5
minutes of whole-head resting-state MEG during both deep and normal nasal breathing (see
Fig 2d). In these data, a comparison of deep vs normal nasal breathing revealed that 1/f slope
over the posterior ROI was indeed more strongly modulated during deep (compared to normal)
breathing (z(27) = 3.02, p = .003, see Fig. 2e).
Finally, to investigate the role of breathing route, we compared respiration-locked 1/f slope
modulations at rest during nasal vs oral breathing in a new sample of N = 25 MEG data sets
(see Methods). Repeating the posterior ROI analysis described above, respective LMEMs
confirmed significant coupling of respiratory phase to 1/f slope during both nasal and oral

breathing (both p < .001; Fig. 2f). The modulatory pattern over phase was again similar to the
original findings (with certain limitations due to reduced power): During nasal breathing, 1/f
slope was flattened during the inspiratory phase (-95° to -52°) as well as shortly after the
inspiration-to-expiration transition (around 36°) and steeper around the expiration-to-
inspiration transition (143° to -150°). During oral breathing, 1/f slope was flattened around the
inspiration-to-expiration transition (-64° to 52°) and steeper around the expiration-to-
inspiration transition (131° to -125°). A comparison of individual ranges of slope ~ respiration
(i.e., max_{slope} - min_{slope} over the respiratory cycle) revealed that the strength of respiration-
locked 1/f slope modulations did not significantly differ between nasal and oral breathing (z(24)
= 0.80, p = .426).

236 Respiration-locked slope modulations are wide-spread, but strongest over posterior
237 sites. Overall, the first analysis demonstrated a consistent pattern of respiration-locked 1/f
slope modulations over parieto-occipital sensors: The slope of aperiodic neural activity was
not uniform over the respiratory cycle, but systematically modulated in such a way that 1/f
slope was flattened around the inspiration-to-expiration transition and steeper around the
expiration-to-inspiration transition. In the next step, we aimed to i) broaden the scope to
investigate respiration-locked slope changes across all M/EEG sensors and ii) characterise
both direction and strength of these modulations across the scalp.
For each participant, single-sensor/channel 1/f slope courses across 60 respiratory phase bins
were extracted using the pipeline described above on single-sensor time series. For each of
the N = 275 MEG sensors (Münster) and N = 62 EEG channels (Leipzig), we then repeated
the combined LMEM and permutation approach (see above). Thus, for each sensor, the
percentile of the empirical beta weight (relative to its null distribution constructed from LMEMs
on surrogate respiration) indicated significance of phase-locked 1/f slope changes. Group-
level average slope courses for n = 240 significant MEG sensors (top) and n = 53 significant
EEG channels (bottom) are shown in Fig. 3a.
To quantify whether there was a consistent modulation of 1/f slope across all
sensors/channels, we computed the circular mean of single-sensor/channel slope courses.
The histograms in Fig. 3a illustrate a clear clustering around the expiration-to-inspiration
transition (i.e., phase bins around ± π). Supporting our parieto-occipital findings from above,
Rayleigh tests on sensor-/channel-wise circular means corroborated that single-
sensor/channel slope courses were not uniform across respiratory phases, but instead
showed a consistent mean direction with the steepest slope around ± π for both data sets
(MEG: z = 126.86, p < .001; EEG: z = 14.38, p < .001). A topographic representation of sensor-
/channel-wise circular means is shown in Fig. 3b.

Fig. 3. Single-sensor (MEG) and channel (EEG) modulation of 1/f slope. a, Polar plots shows group-level averages of normalised single-sensor/channel slope modulation courses over the respiration cycle. LMEM revealed 1/f slope over respiration to significantly deviate from a uniform distribution in n = 240 out of 275 MEG sensors and n = 53 out of 62 EEG channels. Only significant MEG sensors (top) and EEG channels (bottom) are shown, colour coding illustrates sensors and channels 1 to n. Polar histograms illustrate circular mean directions of single-sensor/channel slope distributions (colour-coded according to respiration phase). **b**, Topographic representations of significant modulations and their phase direction for MEG sensors (top) and EEG channels (bottom). **c**, Population statistics quantifying the effect size of slope modulations seen in b. For each sensor/channel, we show the vector norm of LMEM beta weights for sine and cosine of the respiratory phase (akin to a harmonic regression approach). These vector norms were normalised using the mean and standard deviation of the null distribution and are thus given in z values (see Methods for details). In = inspiration, ex = expiration.

Finally, in line with previous approaches [2,7], we quantified the strength of single-sensor/channel phase-locked slope modulations relative to their null distributions: For each sensor/channel, we computed z-scored LMEM beta weights by subtracting the mean of its 'null beta' distribution and dividing by the corresponding standard deviation (see Methods). In both MEG and EEG topographies, the strongest modulations of aperiodic neural activity were indeed observed over parieto-occipital cortices (Fig. 3c).

Periodic and aperiodic components of the RMBO network are distinctly coupled to respiration. In a final series of analyses, we aimed to disentangle the dynamics of respiratory coupling to periodic (i.e., oscillatory) and aperiodic (i.e., non-oscillatory) components of neural activity. So far, no investigation of respiration-brain coupling has directly considered either phase-locked changes of E:I balance or its distinction from oscillatory modulations. To target

this question, we computed source-localised time series from the MEG data (mapped onto n
= 230 parcels taken from the HCP atlas [21]) and defined regions of interest according to the
original RMBO network publication [2]. For a total of n = 10 ROIs (consisting of n = 23 HCP
parcels), we extracted respiration phase-locked 1/f slope as well as periodic spectra with the
1/f characteristic removed. For the computation of parcel-specific courses of aperiodic slope
over respiratory phase, we followed the sensor-level approach outlined above. The periodic
component was defined as the accumulated power of oscillatory peaks (between 1 - 40 Hz)
after the 1/f slope was removed from the respective periodic spectrum.
In keeping with our sensor-level analyses, we determined significance of respiration phase-
locked changes in aperiodic slope and oscillatory power by means of LMEMs within each
parcel of interest. Again, each parcel's empirical vector norm of LMEM beta weights for
respiratory sine and cosine was assessed relative to k = 5000 null beta weights from LMEMs
computed on surrogate distributions of slope and power over respiratory phases. Of the 23
parcels of interest, 18 parcels showed a significant association between respiration and 1/f
slope. Significant phase-locked modulations of oscillatory power (within frequencies from 1 -
40 Hz) were found for 19 parcels (see Fig. 4a). Overall, all 23 parcels showed either significant
non-uniformity of slope or accumulated power.
Next, we investigated differences in temporal dynamics between respective slope and power
changes within each parcel. Specifically, we extracted the circular means of each parcel's
slope and power courses (binned into n = 60 overlapping respiratory phase bins) to compute
parcel-wise phase differences between the two courses: In case slope and power were
similarly modulated by respiration, the mean phase difference across parcels would be (close
to) zero. As shown in the polar histogram in Fig. 4b, this was not the case (U^2 = 1.26, p < .001,
5000 permutations). Next, we tested whether slope and power courses of each parcel were
drawn from the same distribution. For all parcels, Watson's U^2 permutation tests confirmed
significant differences between the underlying distributions (all p < .001), providing further
evidence for differential modulatory dynamics of slope and power courses within the RMBO
network. For both slope and accumulated power, Fig. 4c shows ROI-wise distributions of
subject-level U^2 statistics for each parcel as an estimate of respective non-uniformity (centre
panel). Exemplary comparisons between ROIs are shown for slope ~ respiration (left panel)
and power spectra ~ respiration (right panel).
318

Fig. 4. Source-level ROI modulation of 1/f slope and accumulated oscillatory power in the MEG. a, For each cortical node of the RMBO network [2], we show the group-level average courses of 1/f slope (left polar plots) and accumulated spectral power after removal of the aperiodic component (1 - 40 Hz; right polar plots) over the respiratory cycle. For ROIs comprising more than one parcel from the HCP atlas, slope and power courses are superimposed for each individual parcel. In case the respective LMEM did not indicate significant association with either slope or power within any given parcel, the corresponding time series are drawn with reduced opacity. For significantly associated parcels only, red and blue triangles indicate respective maxima and minima over the respiratory cycle (see legend). In = inspiration, ex = expiration. **b**, Polar histogram shows phase differences between the circular means of slope and power courses across parcels. The clear majority of non-zero phase differences were corroborated by Watson's U² statistics indicating distinct temporal dynamics of slope and power modulations within each parcel (see main text). **c**, Centre: For each ROI, we show violin plots of n = 40 single-participant U² statistics (slope/power vs uniform distribution, respectively). Higher U² statistics indicate stronger non-uniformity. Left: Exemplary slope modulations of SMA and OFC (M ± SEM) normalised across respiration phase for illustration purposes). SMA was more strongly non-uniform overall (see centre), which is visible from both stronger maxima (Δmax) and minima (Δmin) compared to OFC. Right: Exemplary power modulations of OFC and l. INS. For both ROIs, we superimpose power spectra (with 1/f removed) from all 60 phase bins. Overall, power varied more strongly with respiratory phase within OFC compared to l.INS (see centre). TPJ = temporo-parietal junction, ITG/STG = inferior/superior temporal gyrus, ANG = angular gyrus, FEF = frontal eye-field, TP = temporal pole, SMA = supplementary motor area, OFC = orbitofrontal cortex, CUN = cuneus, LS = lingual sulcus, ACC/PCC = anterior/posterior cingulate cortex, INS = insular cortex. In = inspiration, ex = expiration.

Finally, single-band modulations of 1/f-removed theta (4-8 Hz), alpha (8-13 Hz), and beta
power (13-30 Hz) within the RMBO network are shown in Fig. 5. Out of 23 parcels, band-
specific LMEMs revealed that theta power was significantly modulated in 17 parcels, alpha
power in 22 parcels, and beta power in 21 parcels (see Fig. 5a and Methods for details). While
band-specific oscillatory modulations were not the main focus of this paper, we provide this
characterisation to motivate hypothesis generation for future work.

Fig. 5. Source-level ROI modulation of band-specific oscillatory power in the MEG. a, For each cortical node
of the RMBO network, we show the group-level average courses of band-specific spectral power after removal of
the aperiodic component over the respiratory cycle. For ROIs comprising more than one parcel from the HCP atlas,
power courses are shown for each individual parcel. In case the band-specific LMEM did not indicate significant
modulation of the periodic component within any given parcel, the corresponding power series are not shown. **b,**
Exemplary ROI-average spectra of anterior cingulate (top) and left insula (bottom), colour-coded according to
respiratory phase. **c,** Group-level median U² statistics of each parcel testing single-band power ~ respiration phase
against a uniform distribution. Higher values indicate stronger deviation from uniform distribution.

Discussion

Our present findings highlight the role of respiration as a physiological modulator of neural
signalling. We provide evidence that fluctuations of aperiodic brain activity are phase‑locked
to the respiratory cycle which strongly suggests that spontaneous state shifts of excitation‑
inhibition balance are at least partly influenced by peripheral bodily signals. Moreover,
differential temporal dynamics in their coupling to non-oscillatory and oscillatory activity raise
the question whether there is a functional distinction in the way each component is related to
respiration.

While the mechanisms underlying the 1/f characteristic of neural power spectra have yet to be fully understood [22], crossmodal evidence has highlighted its functional significance in several domains. Most prominently, the steepness of the aperiodic slope (i.e., its exponent) has been shown to reflect the balance between excitatory (E) and inhibitory (I) synaptic currents in computational [16], pharmaceutical [15], as well as optogenetic work [23]: The larger the exponent (i.e., steeper power spectral slope), the stronger the inhibitory influence and vice versa [24]. In general, constant E:I balancing is essential to maintain neural homeostasis [25] so that the excitability within a particular neural array remains at a critical state [26].

Facilitated by novel methodological advances, our present findings of respiration-locked changes in E:I balance thus illuminate a previously inaccessible aspect of what is emerging as a recurrent motif in respiration-brain coupling across species: Animal studies have shown excitability [27] as well as spike rates [12,13] to covary with the respiratory cycle, prompting non-invasive human work from our lab [9] in which we linked respiratory phase to excitability of visual cortices (measured as parieto-occipital alpha power) during low-level perception. Given that breathing is under volitional control, we adopted the theoretical perspective of predictive processing [28] to propose respiration as a means to actively align the sampling of external sensory information with transient internal states of increased excitability. This interpretation rests on the concept of active sensing [29] in rodents which recently gained further support from a comprehensive demonstration by Karalis and Sirota [30]: In a series of elaborate experiments, they found neocortical up and down states during sleep and quiet rest to be modulated by respiratory phase, strongly suggesting that breathing influences (sub-)cortical excitability in the sense of a pacemaker aligning neural network dynamics. While the spontaneous aperiodic shifts of E:I balance in our paradigm are not to be equated to the reported up and down states in quiescent animals, we propose that the shared observation of respiration phase-locked state changes bears valuable insight into human respiration-brain coupling.

Investigating the respiration phase-locked changes in aperiodic vs periodic activity inevitably entails the more general distinction of 1/f and genuine oscillations per se and how previous works have addressed one or the other. To reiterate, neural oscillations are narrowband, rhythmic components embedded within a background of broadband, aperiodic activity which itself does not require any underlying rhythmicity [31,32]. So far, studies of breath-brain interactions have neglected aperiodic activity entirely, most likely due to the fact that one proposed mechanism of respiration-brain coupling critically relies on oscillatory entrainment to the breathing rhythm: Here, the nasal airstream entrains neural activity within the olfactory tract via mechanoreceptors, after which the phase of this infraslow rhythm is coupled to the amplitude of faster oscillations [33,12]. This propagation through phase-amplitude coupling [34] has largely been quantified by measures of coupling strength such as the modulation index (MI

[35]), a method that – despite its benefits - prohibits any conclusion regarding non-oscillatory

modulations.

In the present study, we consistently observed steepest 1/f slopes around ± π, with strongest

effects localised over posterior sensors, complementing previous reports of a spatial gradient

of aperiodic dynamics [18, 36]. Such a posterior-to-anterior gradient conceivably increases

parieto-occipital SNR, which could facilitate the detection of respiration phase-locked

changes. As for the pattern of preferred respiratory phase around ± π, He and colleagues [37]

demonstrated a rich temporal organisation of aperiodic (or 'scale-free') brain activity in ECoG

recordings from epilepsy patients: Extending the concept of hierarchically nested oscillations

[38], the authors showed nested frequencies in aperiodic activity during quiet wakefulness; in

other words, coupling of low-frequency phase to high-frequency amplitude of 1/f slope.

Intriguingly, the preferred phase clustered around phases 0 and ± π, i.e., peaks and troughs

of low-frequency fluctuations, closely resembling present findings of flatter slopes (i.e.,

increased excitation) towards the end of inspiration as well as steeper slopes (i.e., more

inhibition) towards the end of expiration (see Figs. 2 and 3). These aperiodic shifts of E:I state

further corroborate previous findings of increased excitability during inspiration [9] and improved

cognitive performance for stimuli presented during the inspiratory phase [6,7].

If the proposed functional mechanisms of respiration-brain coupling were indeed to be

extended to include aperiodic brain activity, respiratory modulations of 1/f slope should be

flexibly adapted during cognitive or perceptual tasks to facilitate performance. A study by

Waschke and colleagues [15] recently provided convincing evidence that the aperiodic

component itself not only captured task-relevant attentional changes in E:I balance, but also

covaried with individual behavioural performance. Critically, the authors used anaesthetics to

distinguish between oscillatory (i.e., increased band-specific power) and non-oscillatory,

aperiodic changes in EEG spectra. In a similar vein, Donoghue et al. [18] stressed the

importance of disentangling periodic and aperiodic changes in neural activity in order to

recognise their potentially distinct contributions to observed effects. We argue that the same

is true in order to more comprehensively describe the physiological underpinnings of

respiration-brain coupling.

In the present data, the preferred phase differences between source-level periodic and

aperiodic modulations within the RMBO network show at least partly differential dynamics

between both components, suggesting a more complex coupling of neural signalling to the

respiratory rhythm than previously assumed. Conceivably, changes in periodic and aperiodic

activity could reflect respiration-locked neural modulations arising from two (not necessarily)

independent mechanisms: While the respiratory rhythm itself is generated in the brain stem

[39], oscillatory changes on the one hand are assumed to be induced by mechanosensory

feedback stimulation from the nasal airstream during inspiration and expiration (see above).
Convincing evidence for this hypothesis comes from studies showing that both neural and
behavioural modulations dissipate when there is no nasal airflow stimulating olfactory
mechanoreceptors, i.e. during oral breathing [6,7] or in bulbectomised animals [12]. On the other
hand, CO_2 levels continuously regulate blood flow and respiration rate in a feedforward
fashion, which in turn determine changes in CO_2 levels [40]. Moreover, CO_2 is inversely related
to tissue acidity (pH), which has been linked to neural excitability [41,42]. Lowered pH (due to
increased CO_2) causes an increase in extracellular adenosine [43], a neuromodulator gating
synaptic transmission. Consequently, Dulla and colleagues [44] concluded that CO_2-induced
changes in neural excitability are caused by pH-dependent modulation of adenosine and ATP
levels (also see [45]).

Since CO_2 levels do not differ between nasal and oral breathing, the two mechanisms are
clearly functionally distinct from one another: CO_2 fluctuations alone cannot account for neural
or behavioural differences between nasal and oral breathing. Hence, there are (at least) two
different ways in which neural activity could be coupled to the act of breathing. It may be the
case that the differential coupling of respiration to periodic and aperiodic activity reflects these
distinct mechanisms. However, the evidence so far (including present data from the control
study) neither precludes the possibility that both types of modulation could be rooted in the
same underlying mechanism. Therefore, critical future work is needed to further elucidate the
functional and anatomical underpinnings of respiratory modulations in brain dynamics.
460
The present findings open multiple avenues for future research, of which we want to briefly
sketch three main directions. First, mechanistic advances could be made in a replicative study
which includes a contrast of nasal vs oral respiration. The absence of phase-amplitude
coupling driven by nasal airstreams during oral breathing could conceivably reveal to what
extent aperiodic changes rely on and/or interact with oscillatory changes. While we speculate
that respective changes in 1/f slope and oscillatory power could point to distinct modulatory
pathways, targeted follow-up studies with sufficient statistical power are of critical importance
to extend our initial results.

Second, as outlined above, it would further be highly instructive to complement existing
evidence of task-related 1/f slope changes [23,46] by investigating their link to respiration in
behavioural cognitive or perceptual paradigms. Timed breathing (in healthy participants or
ventilated patients) or contexts involving different brain states, e.g. resting-state vs task, during
attentional or arousal manipulation (including hyperventilation), or during sleep stages, could
potentially unravel the link between breathing and fluctuations of (non-)oscillatory brain
activity.

Finally, a third line of research should focus on potential translational applications of
respiration-brain coupling. While others have made a convincing case for studying respiratory
involvement in clinical contexts before (for a recent review, see [47]), the results presented here
demonstrate a novel link between bodily signals and aperiodic neural activity. Importantly,
changes in 1/f slope itself - an indicator of E:I imbalance - have been implicated in neurological
and psychiatric disorders like Alzheimer's disease [48], schizophrenia [49], autism spectrum
disorder [50], and epilepsy [51]. Linking these known neural alterations to peripheral signals could
provide substantial insight into body-brain interactions in health and disease.
484
Methods

Participants and data acquisition (Münster). Forty right-handed volunteers (21 female, age
25.1 ± 2.7 y [M ± SD]) participated in the study. All participants reported having no respiratory
or neurological disease and gave written informed consent prior to all experimental
procedures. The study was approved by the local ethics committee of the University of
Münster (approval ID 2018-068-f-S). All participants provided written informed consent and
received financial compensation for their participation in the study. Participants were seated
upright in a magnetically shielded room while we simultaneously recorded 5 minutes of MEG
and respiratory data. MEG data was acquired using a 275 channel whole-head system
(OMEGA 275, VSM Medtech Ltd., Vancouver, Canada) at a sampling frequency of 600 Hz.
During recording, participants were to keep their eyes on a fixation cross centred on a projector
screen placed in front of them. To minimise head movement, participants' heads were
stabilised with cotton pads inside the MEG helmet.

Participants were instructed to breathe naturally through their nose while the respiratory signal
was measured as thoracic circumference by means of a respiration belt transducer (BIOPAC
Systems, Goleta, USA) placed around their chest. Continuous monitoring via video ensured
participants were breathing through their nose instead of their mouth. Individual respiration
time courses were visually inspected for irregular breathing patterns such as breath holds or
unusual breathing frequencies, but no such artefacts were detected.
504

Participants and data acquisition (Leipzig). Thirty-eight healthy volunteers (18 female, age
27.1 ± 4.0 y [M ± SD]) were recruited from the database of the Max Planck Institute for Human
Cognitive and Brain Sciences, Leipzig, Germany. Participants reported no history or current
neurological or psychological condition. The study was approved by the Ethical Committee of
the University of Leipzig's Medical Faculty (No. 462-15-28082020). All participants provided
written informed consent and received financial compensation for their participation in the
study. Participants were seated upright in an EEG booth while 5 minutes of EEG and

respiration were recorded. EEG was recorded from 62 scalp positions distributed over both hemispheres according to the international 10–10 system, using a commercial EEG acquisition system (ActiCap Snap, BrainAmp; Brain Products). The mid-frontal electrode (FCz) was used as the reference and a mid-frontal electrode placed on the middle part of the forehead (between FP1 and FP2) as ground. Electrode TP9 was used to measure ECG and electrode TP10 captured eye movements. Electrode impedance was kept below 10 kΩ for all channels. EEG was sampled with a rate of 2.5 kHz and online bandpass-filtered between 0.015 - 1000 Hz. Participants were instructed to keep their eyes open and fixate a cross on the screen in front of them to avoid excess eye movements. Participants were instructed to breathe naturally while respiration was measured with a respiration belt transducer (BIOPAC Systems, Goleta, USA) placed around the chest.

Participants and data acquisition (control MEG study for deep vs normal breathing). The MEG sample for the comparison of deep vs normal breathing has previously been published elsewhere [3]. Twenty-eight volunteers (14 female, age 24.8 ± 2.9 y [M ± SD]) participated in the study conducted at the Institute for Biomagnetism and Biosignal Analysis in Münster. All participants reported having no respiratory or neurological disease and gave written informed consent prior to all experimental procedures. The study was approved by the local ethics committee of the University of Münster (approval ID 2018-068-f-S). MEG recording parameters and procedures were identical to the original recordings described above.

Data were acquired in two 5-minute runs with a short intermediate break (determined by the participants). Within each run, participants were instructed to either breathe normally or voluntarily deeply (with their mouth closed) while maintaining their normal respiration frequency, i.e. not to breathe more slowly during blocks of deep breathing. Continuous monitoring via video ensured participants did indeed keep their mouth closed throughout the recording. The order of deep and normal breathing was counterbalanced across participants. For both runs, we recorded the respiratory signal as thoracic circumference by means of a respiration belt transducer (BIOPAC Systems, Goleta, USA) placed around the participant's chest. Individual respiration time courses were visually inspected for irregular breathing patterns such as breath holds or unusual breathing frequencies, but no such artefacts were detected.

Participants and data acquisition (control MEG study for nasal vs oral breathing). Twenty-five right-handed volunteers (10 female, age 26.3 ± 3.3 y [M ± SD]) participated in the control MEG study conducted at the Institute for Biomagnetism and Biosignal Analysis in Münster. All participants reported having no respiratory or neurological disease and gave written informed consent prior to all experimental procedures. The study was approved by the

local ethics committee of the University of Münster (approval ID 2021-785-f-S). MEG recording
parameters and procedures were identical to the original recordings described above.

For each participant, we recorded two 6-minute runs of resting state activity: In one run,
participants were instructed to breathe naturally through their nose. Continuous monitoring via
video ensured participants did indeed keep their mouth closed throughout the recording. In
the other run, participants were instructed to breathe through their mouth while wearing a nose
clip to prevent nasal breathing. The order of nasal and oral breathing was counterbalanced
across participants. Again, we recorded the respiratory signal as thoracic circumference by
means of a respiration belt transducer (BIOPAC Systems, Goleta, USA) placed around the
participant's chest. Individual respiration time courses were visually inspected for irregular
breathing patterns such as breath holds or unusual breathing frequencies, but no such
artefacts were detected.

MRI acquisition and co-registration (Münster). For MEG source localisation, we obtained
high-resolution structural magnetic resonance imaging (MRI) scans in a 3T Magnetom Prisma
scanner (Siemens, Erlangen, Germany). Anatomical images were acquired using a standard
Siemens 3D T1-weighted whole-brain MPRAGE imaging sequence (1×1×1 mm voxel size,
TR = 2,130ms, TE = 3.51 ms, 256 × 256 mm field of view, 192 sagittal slices). MRI
measurement was conducted in supine position to reduce head movements, and gadolinium
markers were placed at the nasion as well as left and right distal outer ear canal positions for
landmark-based co-registration of MEG and MRI coordinate systems. Data preprocessing was
performed using Fieldtrip [52] running in MATLAB R2021a (The Mathworks, Natick, USA).
Individual raw MEG data were visually inspected for jump artefacts and bad channels, but
neither were detected. Both MEG and respiration data were resampled to 300 Hz prior to
further analyses.

Co-registration of structural T1 MRIs to the MEG coordinate system was done for each
participant by initial identification of three anatomical landmarks (nasion, left and right pre-
auricular points) in their individual MRI. Using the implemented segmentation algorithms in
Fieldtrip and SPM12, individual head models were constructed from anatomical MRIs. A
solution of the forward model was computed using the realistically-shaped single-shell volume
conductor model [53] with a 5 mm grid defined in the Human Connectome Project (HCP)
template brain [21] after linear transformation to the individual MRI.

Respiratory preprocessing (both labs). To obtain continuous respiration phase angles, we
used Matlab's findpeaks function (with minimal peak prominence set to 1) to identify time
points of peak inspiration (peaks) and peak expiration (troughs) in the normalised respiration
time course. Phase angles were linearly interpolated from trough to peak (−π to 0) and peak

to trough (0 to π) in order to yield respiration cycles centred around peak inspiration (i.e.,
phase 0).

In order to assess the potential influence of individual breathing parameters on aperiodic
fluctuations, we computed breathing rates and depths for each participant. Breathing rates
were extracted as the mean distance between inspiratory peaks as defined by the peak
detection algorithm described above. Breathing depths were computed as the integral of the
individual respiration time series, normalised by the number of breathing cycles during the
recording.

Head movement correction (Münster). In order to rule out head movement as a potential
confound in our analyses, we used a correction method established by Stolk and colleagues
[54]. This method uses the accurate online head movement tracking that is performed by our
acquisition system during MEG recordings. This leads to six continuous signals (temporally
aligned with the MEG signal) that represent the x, y, and z coordinates of the head centre (H_x,
H_y, H_z) and the three rotation angles (H_ψ, H_θ, H_ϕ) that together fully describe head movement.
We constructed a regression model comprising these six 'raw' signals as well as their
derivatives and, from these 12 signals, the first-, second-, and third-order non-linear
regressors to compute a total of 36 head movement-related regression weights (using a third-
order polynomial fit to remove slow drifts). This regression analysis was performed on the
power spectra of single-sensor and single-voxel time courses, respectively, removing signal
components that can be explained by translation or rotation of the head with respect to the
MEG sensors.

Respiration-locked computations of 1/f slope and oscillatory power. Our first aim was to
further investigate excitability changes over posterior cortices we previously reported [9]. To this
end, we defined parieto-occipital regions of interest for movement-corrected MEG (k = 41
sensors) as well as EEG data (k = 17 channels). Single-sensor/channel time series within
these ROIs were entered into the SPRiNT algorithm [19] with default parameter settings and
subsequently averaged. In short, SPRiNT is based on the specparam algorithm [18] and uses a
short-time Fourier transform (frequency range 1-40 Hz) to compute aperiodic and periodic
components of neural time series within a moving window (width of 1 s, 75% overlap between
two neighbouring windows). Due to the slow nature of the respiratory signal, we did not
average between neighbouring windows at this point (see below for details).

SPRiNT thus yielded time series of both aperiodic (i.e., 1/f exponent) and periodic signals (i.e.,
oscillatory power) with a temporal resolution of 250 ms and a frequency resolution of 1 Hz. In
order to relate these time series to the respiratory signal, we extracted respiratory phase at all
time points for which slope and power were fitted (i.e., the centres of each moving window).

Following previous work [9,2], we then partitioned the entire respiratory cycle (-π to π) into n =

60 equidistant, overlapping phase bins. Moving along the respiration cycle in increments of

$\Delta\omega = \pi/30$, we collected all SPRiNT outputs (i.e., slope fits and periodic spectra) computed at

a respiration angle of $\omega \pm \pi/10$. At this point, we computed individual bin-wise averages of 1/f

slope and periodic spectra, yielding quasi-continuous 'phase courses' of aperiodic and

periodic neural signals for each participant. For the control MEG studies, we separately

applied this pipeline to the MEG data recorded during deep/normal and nasal/oral breathing,

respectively. The methodological approach is illustrated in Fig. 1.

For the source-level MEG analyses shown in Figs. 4 and 5, SPRiNT computed 1/f slope fits

and periodic spectra based on the source-localised time series of individual parcels from the

HCP atlas.

Statistical analysis of sensor-level 1/f slope modulation. As a first analysis of respiration

phase-locked changes in 1/f slope, we computed the following linear mixed-effects model

(LMEM):

$$SE_j = \beta_0 + (S_{1,j} + \beta_1) * \sin_{resp} + (S_{2,j} + \beta_2) * \cos_{resp} + e_j \qquad (1)$$

For each participant j, the model predicted individual slope (resulting from SPRiNT

computations, see above) as a combination of the intercept (β_0), fixed effects of respiratory

sine and cosine (β_1, β_2), and an error term ($e_j \sim N(0,\sigma^2)$). In line with previous work [9], resulting

beta weights for sine and cosine of the respiratory signal were combined in a respiratory phase

vector norm, i.e.

$$v = \sqrt{\beta_1^2 + \beta_2^2} \qquad (2)$$

To test whether 1/f was significantly modulated over the ROI sensors, we computed k = 5000

random iterations of the LMEM by shuffling bin-wise 1/f slope fits on the individual level. This

way, any meaningful relations between 1/f slope and respiratory phase were removed, thereby

specifically testing the hypothesis that 1/f slope changes significantly with respiratory phase.

For each of these iterations, we once again computed the phase vector norm from the resulting

beta weights, yielding a random distribution of k = 5000 'null vector norms'. Significance of the

empirical vector norm was determined by its percentile relative to this null distribution.

As shown in Figs. 2 and 3, for the sensor-level analyses, SPRiNT parameterisation was

conducted on individual MEG sensors (Münster) and EEG channels (Leipzig). For the analysis

of 1/f slope modulations within the parieto-occipital ROI, we computed mean SPRiNT

estimates across the corresponding sensors or channels, respectively. LMEMs were then
computed on ROI-average estimates for 1/f slope ~ respiration phase for each lab individually
(Münster, Leipzig) as well as for the pooled data from both labs (see Fig. 2). For the whole-
scalp analyses shown in Fig. 3, LMEMs were computed on single-sensor/channel SPRiNT
estimates.

Having established an overall significant influence of respiratory phase on 1/f slope with the
LMEM approach described above, we next aimed to characterise when (i.e., at which
respiratory phases) 1/f slope was significantly steeper or flatter. To this end, we implemented
a permutation approach as follows: For each participant, we constructed a surrogate
respiration time series using the iterated amplitude-adjusted Fourier transform (IAAFT [55]). In
contrast to shuffling the respiration time series, this iterative procedure preserves the temporal
autocorrelation of the signal, which is critical for constructing a fitting null distribution for
permutation testing. From these IAAFT-transformed respiration time series, we extracted the
surrogate respiratory phase values corresponding to each slope estimation from SPRiNT. In
keeping with the approach above, we finally binned all SPRiNT outputs into n = 60 equidistant,
overlapping phase bins covering one entire respiratory cycle (-π to π) and computed the bin-
wise average slope fit to yield a 'null time series' of 1/f slope over surrogate respiration phase.
For each participant, this procedure was repeated 5000 times and resulted in a null distribution
of 5000 surrogate 1/f slope estimates x 60 phase bins. We then computed the group-level
average null distribution of 1/f slope for each of the 60 phase bins which allowed us to extract
the percentiles of empirical bin-wise group means relative to these null distributions. As
illustrated in Fig. 2, empirical group-level means were deemed significant if they exceeded the
95th percentile (or fell below the 5th percentile) of the null distribution.

For single-sensor slope analyses (Fig. 3) as well as single-parcel analyses of 1/f slope and
periodic spectra (Fig. 4), we used the circstat toolbox for Matlab [56] to compute the circular
means of group-level average slope (and power) courses over the respiratory cycle. In the
absence of meaningful slope modulation by respiratory phase, a polar illustration of the binned
slope estimates would be a uniform circle. In contrast, any phase-locked slope changes would
be represented as modulations or peaks within the polar representation. Albeit not for all
distributions, the circular mean provides at least some indication with regard to the direction
of such modulations (see below for alternative measures). Therefore, we used it to
characterise the consistency of slope modulations across the whole scalp both within
MEG/EEG data as well as across both modalities and labs (see Fig. 3).

In addition to potential respiration-related modulations of slope courses, we quantified the
strength of these phase-locked modulations. To this end, we computed z scores from sensor-
specific LMEM beta weights of the respiratory phase vector norm (see above for details) by

subtracting the mean of its null distribution and dividing by the corresponding standard deviation:

$$z = \frac{v - \mu_{v(\text{null})}}{\sigma_{v(\text{null})}}$$ (3)

with v being the vector norm of beta weights for respiratory sine and cosine (see Eq. 2). This way, empirical LMEM beta weights for the respiratory phase vector are given in units of standard deviation relative to their null distribution.

Source reconstruction. Source reconstruction was performed using the linearly constrained minimum variance (LCMV) beamformer approach [57] with the lambda regularisation parameter set to 5%. This approach estimates a spatial filter for each location of the 5 mm grid along the direction yielding maximum power. A single broadband LCMV beamformer was used to estimate voxel-level activities across all frequencies.

ROI-based analyses of source-level 1/f slope and oscillatory power. For the investigation of source-level modulations of 1/f slope and periodic spectra, we first extracted neural time series from a total of n = 230 cortical parcels from the HCP atlas [21]. In order to ground our analyses in previous findings, we then focussed our analyses on those parcels representing the cortical nodes of the RMBO network [2]. This network has been demonstrated to show respiration-modulated brain oscillations (RMBOs) during resting state, which is why it was particularly well-suited to investigate potentially distinct dynamics of periodic and aperiodic modulations. As some RMBO nodes comprised bilateral or wider-spread anatomical sites, a total of N = 10 ROIs consisting of n = 23 parcels were used for source-level analyses (see Fig. 4). For the average time series within these parcels, we applied the SPRiNT analysis described above to yield both 1/f slope and oscillatory power across the respiratory cycle. For the latter, SPRiNT computed periodic spectra between 1 - 40 Hz with the aperiodic component removed, i.e., solely the power of the periodic neural signal. Just as with the 1/f exponents, we sorted all periodic spectra with respect to the corresponding respiratory phase to yield individual bin-average periodic spectra for n = 60 overlapping phase bins. These spectra thus reflect, for each phase of the respiratory cycle, the extent of 'true' oscillatory activity remaining after the underlying 1/f characteristic is removed. In a first step, we quantified the amount of oscillatory activity by computing the accumulated sum over all frequencies (illustrated in Fig. 4). For frequency band-specific power courses, we then computed accumulated power for canonical bands separately (theta: 4 - 8 Hz, alpha: 8 - 13 Hz, beta: 13 - 30 Hz).

Next, we aimed to investigate potentially distinct respiration phase-locked dynamics of periodic

and aperiodic modulations. Therefore, for each HCP parcel, we implemented Watson's U^2

tests with k = 5000 random permutations (watsons_U2_perm_test function [58] for Matlab) to

test the null hypothesis that the group-level average vectors of slope and power were drawn

from the same distribution. We chose the U^2 test as an additional measure as it is sensitive

towards multimodal distributions [59], which is not the case for the circular mean alone. For the

sake of comparability, however, we also computed phase differences between each parcel's

circular means of slope and power courses: If 1/f slope and oscillatory power were similarly

coupled to the respiratory rhythm, their respective courses over respiratory phase would not

differ and the phase difference between their circular means would be (close to) zero. In

contrast, systematically different distributions of slope and power would suggest differential

dynamics in their respective coupling to respiratory phase.

Finally, we characterised the strength of slope and power modulations for each of the n = 10

RMBO ROIs by means of U^2 statistics. For each participant, we conducted Watson's U^2 tests

of the (potentially multimodal) courses of 1/f slope and accumulated power within each ROI

against a uniform distribution. Here, higher U^2 values indicate stronger non-uniformity within a

given parcel, which allowed us to characterise RMBO ROIs by their respective group-level U^2

distributions (as shown in Fig. 4c).

Data availability

The processed data generated in this study will be made publicly available from the Open

Science Framework (https://osf.io/8nw9t/) upon publication of the study.

Code availability

All custom Matlab code to reproduce the central findings of this study will be made publicly

available from the Open Science Framework (https://osf.io/8nw9t/) upon publication of the

study.

References

1. Jelinčić, V., Van Diest, I., Torta, D. M. & von Leupoldt, A. The breathing brain: The

potential of neural oscillations for the understanding of respiratory perception in health

and disease. Psychophysiology e13844 (2021) doi:10.1111/psyp.13844.

2. Kluger, D. S. & Gross, J. Respiration modulates oscillatory neural network activity at rest.

PLoS Biol. **19**, e3001457 (2021).

3. Kluger, D. S. & Gross, J. Depth and phase of respiration modulate cortico-muscular

communication. Neuroimage **222**, 117272 (2020).

4. Rassler, B. & Raabe, J. Co-ordination of breathing with rhythmic head and eye
movements and with passive turnings of the body. Eur. J. Appl. Physiol. **90**, 125–130
(2003).

5. Johannknecht, M. & Kayser, C. The influence of the respiratory cycle on reaction times in
sensory-cognitive paradigms. Sci. Rep. **12**, 2586 (2022).

6. Perl, O. et al. Human non-olfactory cognition phase-locked with inhalation. Nat. Hum.
Behav. **3**, 501–512 (2019).

7. Zelano, C. et al. Nasal respiration entrains human limbic oscillations and modulates
cognitive function. J. Neurosci. **36**, 12448–12467 (2016).

8. Grund, M. et al. Respiration, heartbeat, and conscious tactile perception. J. Neurosci. **42**,
643–656 (2022).

9. Kluger, D. S., Balestrieri, E., Busch, N. A. & Gross, J. Respiration aligns perception with
neural excitability. eLife **10**, (2021).

10. Galvez-Pol, A., McConnell, R. & Kilner, J. M. Active sampling in visual search is coupled
to the cardiac cycle. Cognition **196**, 104149 (2020).

11. Biskamp, J., Bartos, M. & Sauer, J.-F. Organization of prefrontal network activity by
respiration-related oscillations. Sci. Rep. **7**, 45508 (2017).

12. Ito, J. et al. Whisker barrel cortex delta oscillations and gamma power in the awake mouse
are linked to respiration. Nat. Commun. **5**, 3572 (2014).

13. Yanovsky, Y., Ciatipis, M., Draguhn, A., Tort, A. B. L. & Brankačk, J. Slow oscillations in
the mouse hippocampus entrained by nasal respiration. J. Neurosci. **34**, 5949–5964
(2014).

14. He, B. J. Scale-free brain activity: past, present, and future. Trends Cogn Sci (Regul Ed)
18, 480–487 (2014).

15. Waschke, L. et al. Modality-specific tracking of attention and sensory statistics in the
human electrophysiological spectral exponent. eLife **10**, (2021).

16. Gao, R., Peterson, E. J. & Voytek, B. Inferring synaptic excitation/inhibition balance from
field potentials. Neuroimage **158**, 70–78 (2017).

17. Trakoshis, S., Rocchi, F., Canella, C., You, W., Chakrabarti, B., Ruigrok, A. N., ... & MRC
AIMS Consortium. Intrinsic excitation-inhibition imbalance affects medial prefrontal cortex
differently in autistic men versus women. elife, 9, e55684 (2020).

18. Donoghue, T. et al. Parameterizing neural power spectra into periodic and aperiodic
components. Nat. Neurosci. **23**, 1655–1665 (2020).

19. Wilson, L. E., da Silva Castanheira, J., & Baillet, S. Time-resolved parameterization of
aperiodic and periodic brain activity. eLife, 11, e77348 (2022).

20. Samaha, J., Iemi, L., Haegens, S. & Busch, N. A. Spontaneous Brain Oscillations and

Perceptual Decision-Making. Trends Cogn Sci (Regul Ed) **24**, 639–653 (2020).

21. Glasser, M. F. et al. A multi-modal parcellation of human cerebral cortex. Nature **536**,
171–178 (2016).

22. Buzsáki, G., Anastassiou, C. A. & Koch, C. The origin of extracellular fields and currents-
-EEG, ECoG, LFP and spikes. Nat. Rev. Neurosci. **13**, 407–420 (2012).

23. Chini, M., Pfeffer, T., & Hanganu-Opatz, I. An increase of inhibition drives the
developmental decorrelation of neural activity. eLife, 11, e78811 (2022).

24. Voytek, B. et al. Age-Related Changes in 1/f Neural Electrophysiological Noise. J.
Neurosci. **35**, 13257–13265 (2015).

25. Turrigiano, G. G. & Nelson, S. B. Homeostatic plasticity in the developing nervous system.
Nat. Rev. Neurosci. **5**, 97–107 (2004).

26. Xue, M., Atallah, B. V. & Scanziani, M. Equalizing excitation-inhibition ratios across visual
cortical neurons. Nature **511**, 596–600 (2014).

27. Dulla, C. G. et al. Adenosine and ATP link PCO2 to cortical excitability via pH. Neuron **48**,
1011–1023 (2005).

28. Mumford, D. On the computational architecture of the neocortex. II. The role of cortico-
cortical loops. Biol. Cybern. **66**, 241–251 (1992).

29. Wachowiak, M. All in a sniff: olfaction as a model for active sensing. Neuron **71**, 962–973
(2011).

30. Karalis, N. & Sirota, A. Breathing coordinates cortico-hippocampal dynamics in mice
during offline states. Nat. Commun. **13**, 467 (2022).

31. Buzsáki, G. Rhythms of the Brain. (Oxford University Press, 2006).
doi:10.1093/acprof:oso/9780195301069.001.0001.

32. Bullock, T. H., Mcclune, M. C. & Enright, J. T. Are the electroencephalograms mainly
rhythmic? Assessment of periodicity in wide-band time series. Neuroscience **121**, 233–
252 (2003).

33. Fontanini, A. & Bower, J. M. Slow-waves in the olfactory system: an olfactory perspective
on cortical rhythms. Trends Neurosci. **29**, 429–437 (2006).

34. Canolty, R. T. & Knight, R. T. The functional role of cross-frequency coupling. Trends
Cogn Sci (Regul Ed) **14**, 506–515 (2010).

35. Tort, A. B. L. et al. Dynamic cross-frequency couplings of local field potential oscillations
in rat striatum and hippocampus during performance of a T-maze task. Proc Natl Acad
Sci USA **105**, 20517–20522 (2008).

36. Mahjoory, K., Schoffelen, J.-M., Keitel, A. & Gross, J. The frequency gradient of human
resting-state brain oscillations follows cortical hierarchies. eLife **9**, (2020).

37. He, B. J., Zempel, J. M., Snyder, A. Z. & Raichle, M. E. The temporal structures and
functional significance of scale-free brain activity. Neuron **66**, 353–369 (2010).

38. Lakatos, P. et al. An oscillatory hierarchy controlling neuronal excitability and stimulus
processing in the auditory cortex. J. Neurophysiol. **94**, 1904–1911 (2005).

39. Del Negro, C. A., Funk, G. D. & Feldman, J. L. Breathing matters. Nat. Rev. Neurosci. **19**,
351–367 (2018).

40. Feldman, J.L., Mitchell, G.S., & Nattie, E.E. Breathing: rhythmicity, plasticity,
chemosensitivity. Annu. Rev. Neurosci. 26, 239–266 (2003).

41. Chesler, M. Regulation and modulation of pH in the brain. Physiol. Rev. 83, 1183–1221
(2003).

42. Lee, J., Taira, T., Pihlaja, P., Ransom, B.R., & Kaila, K. Effects of CO_2 on excitatory
transmission apparently caused by changes in intracellular pH in the rat hippocampal
slice. Brain Res. 706, 210–216 (1996).

43. Ito, H., Kanno, I., Ibaraki, M., Hatazawa, J., & Miura, S. Changes in human cerebral blood
flow and cerebral blood volume during hypercapnia and hypocapnia measured by positron
emission tomography. J. Cereb. Blood Flow Metab. 23, 665–670 (2003).

44. Dulla, C.G., Dobelis, P., Pearson, T., Frenguelli, B.G., Staley, K.J., & Masino, S.A.
Adenosine and ATP link PCO_2 to cortical excitability via pH. Neuron 48, 1011–1023
(2005).

45. Gourine, A.V., Llaudet, E., Dale, N., & Spyer, K.M. ATP is a mediator of chemosensory
transduction in the central nervous system. Nature 436, 108–111 (2005).

46. Ouyang, G., Hildebrandt, A., Schmitz, F. & Herrmann, C. S. Decomposing alpha and 1/f
brain activities reveals their differential associations with cognitive processing speed.
Neuroimage **205**, 116304 (2020).

47. Heck, D. H. et al. Recent insights into respiratory modulation of brain activity offer new
perspectives on cognition and emotion. Biol. Psychol. **170**, 108316 (2022).

48. Ramírez-Toraño, F. et al. Functional connectivity hypersynchronization in relatives of
alzheimer's disease patients: an early E/I balance dysfunction? Cereb. Cortex **31**, 1201–
1210 (2021).

49. Molina, J. L. et al. Memantine effects on electroencephalographic measures of putative
excitatory/inhibitory balance in schizophrenia. Biol. Psychiatry Cogn. Neurosci.
Neuroimaging **5**, 562–568 (2020).

50. Rubenstein, J. L. R. & Merzenich, M. M. Model of autism: increased ratio of
excitation/inhibition in key neural systems. Genes Brain Behav. **2**, 255–267 (2003).

51. van Heumen, S. et al. Case Report: Aperiodic Fluctuations of Neural Activity in the Ictal
MEG of a Child With Drug-Resistant Fronto-Temporal Epilepsy. Front. Hum. Neurosci.
15, 646426 (2021).

52. Oostenveld, R., Fries, P., Maris, E. & Schoffelen, J.-M. FieldTrip: Open source software
for advanced analysis of MEG, EEG, and invasive electrophysiological data. Comput.

Intell. Neurosci. **2011**, 156869 (2011).

53. Nolte, G. The magnetic lead field theorem in the quasi-static approximation and its use

for magnetoencephalography forward calculation in realistic volume conductors. Phys.

Med. Biol. **48**, 3637–3652 (2003).

54. Stolk, A., Todorovic, A., Schoffelen, J.-M. & Oostenveld, R. Online and offline tools for

head movement compensation in MEG. Neuroimage **68**, 39–48 (2013).

55. Theiler, J., Eubank, S., Longtin, A., Galdrikian, B., & Farmer, J. D. (1992). Testing for

nonlinearity in time series: the method of surrogate data. Physica D: Nonlinear

Phenomena, 58(1-4), 77-94.

56. Berens, P. circstat : a MATLAB toolbox for circular statistics. J. Stat. Softw. **31**, (2009).

57. Van Veen, B. D., van Drongelen, W., Yuchtman, M. & Suzuki, A. Localization of brain

electrical activity via linearly constrained minimum variance spatial filtering. IEEE Trans

Biomed Eng **44**, 867–880 (1997).

58. Mégevand, P. pierremegevand/watsons_u2

https://github.com/pierremegevand/ watsons_u2, GitHub. Retrieved July 5, 2022. (2022)

59. Landler, L., Ruxton, G. D. & Malkemper, E. P. Advice on comparing two independent

samples of circular data in biology. Sci. Rep. **11**, 20337 (2021).

897   **Acknowledgements**

898 We would like to thank Karin Wilken, Ute Trompeter, and Hildegard Deitermann for their

899 invaluable assistance during data collection in Münster. This work was supported by the

900 Interdisciplinary Center for Clinical Research (IZKF) of the medical faculty of Münster

901 (Gro3/001/19, awarded to JG). DSK (KL-3580/1-1), AV (Research Training Group 2386), and

902 JG (GR 2024/ 5-1) were further supported by the DFG.

904   **Author Contributions Statement**

905 Conceptualization, DSK; Methodology, DSK, CF, JG; Investigation, DSK, CF; Writing –

906 Original Draft, DSK, CF; Writing – Review & Editing, DSK, CF, OA, NC, AV, JG; Visualization,

907 DSK; Funding Acquisition, DSK, AV, JG

909   **Competing Interests Statement**

910 The authors declare no competing interests.