Modulatory dynamics of periodic and aperiodic activity in respiration-brain coupling

Daniel S. Kluger*[1,2], Carina Forster[3,4,5], Omid Abbasi[1], Nikos Chalas[1,2], Arno Villringer[3,4,6], Joachim Gross[1,2]

[1] Institute for Biomagnetism and Biosignal Analysis, University of Münster, Münster, Germany

[2] Otto Creutzfeldt Center for Cognitive and Behavioral Neuroscience, University of Münster, Münster, Germany

[3] Department of Neurology, Max Planck Institute for Human Cognitive and Brain Sciences, Leipzig, Germany

[4] Charité – Universitätsmedizin Berlin, Einstein Center for Neurosciences, Berlin, Germany

[5] Charité – Universitätsmedizin Berlin, Bernstein Center for Computational Neuroscience, Berlin, Germany

[6] Humboldt-Universität zu Berlin, Faculty of Philosophy, Berlin School of Mind and Brain, MindBrainBody Institute, Berlin, Germany

* corresponding author

Contact Info

Daniel S. Kluger

Institute for Biomagnetism and Biosignal Analysis, University of Münster

Malmedyweg 15, 48149 Münster

Germany

Telephone: 0049-251-8352543

E-mail: daniel.kluger@uni-muenster.de

Keywords

respiration, excitation/inhibition, 1/f slope, excitability, body-brain coupling, MEG, EEG

Abstract

Bodily rhythms such as respiration are increasingly acknowledged to modulate neural oscillations underlying human action, perception, and cognition. Conversely, the link between respiration and *aperiodic* brain activity – a non-oscillatory reflection of excitation-inhibition (E:I) balance – has remained unstudied. Aiming to disentangle potential respiration-related dynamics of periodic and aperiodic activity, we applied novel algorithms of time-resolved parameter estimation to resting-state M/EEG data from two labs (N = 78).

Our findings highlight the role of respiration as a physiological influence on brain signalling. We provide first evidence that fluctuations of aperiodic brain activity (1/f slope) are phase-locked to the respiratory cycle which strongly suggests that spontaneous state shifts of excitation-inhibition balance are at least partly influenced by peripheral bodily signals. Moreover, differential temporal dynamics in their coupling to non-oscillatory and oscillatory activity raise the possibility of a functional distinction in the way each component is related to respiration.

Introduction

Recent years have seen a surge in reports of respiration as a modulator of brain signalling -

at least partly through modulation of rhythmic brain activity [1]. These *respiration-modulated*

brain oscillations (RMBOs) have been localised within a widespread network across the

cortex [2] and are increasingly being linked to changes in motor function [3,4], cognitive

processing [5-7], and fundamental aspects of perception [8,9]. One candidate mechanism by

which respiration (and other bodily signals such as the cardiac rhythm; see e.g. [10]) is thought

to coordinate brain signalling is *cortical excitability*, a dynamic brain state indexing neural

populations' readiness to become activated. This proposal rests on findings from intracranial

animal studies demonstrating that respiration modulates spike rates in a variety of brain

regions [11-13]. Moreover, one recent study [9] directly investigated respiratory modulation of

excitability during low-level visual perception.

Critically, brain activity is not entirely oscillatory: One can distinguish its genuinely periodic,

oscillatory component from an underlying stationary, aperiodic component typically showing

a characteristic 1/f dependence of power and frequency (meaning that lower frequencies

carry higher power [14]). In fact, dominance of low-frequency power is by no means just an

epi-phenomenon of neural power spectra, but does reflect meaningful characteristics of

brain function. Most relevantly, the 'steepness' of the aperiodic component has been shown

to reflect the dynamics of excitation-inhibition (E:I) balance [15-17], with steeper 1/f slope

indicating a stronger influence of inhibitory activity.

While sophisticated algorithms exist to disentangle periodic and aperiodic components of

neural power spectra [18], *time-resolved* estimations of 1/f slope in particular have proven

computationally expensive and lacked empirical validation. Consequently, the selective

dynamics linking peripheral signals (such as respiration) to periodic and aperiodic neural

activity are entirely unstudied. Meanwhile, the recent introduction of the *SPRiNT* toolbox [19]

provides methodological advances that allow us to parameterize stationary and oscillatory

brain activity dynamically over time. Here, we apply these algorithms in the realm of human

body-brain interactions to test for the first time the hypothesis that 1/f slope as a marker of

E:I balance is modulated by respiration.

Specifically, we used concurrent resting-state M/EEG and respiratory recordings in

populations from two labs (Münster, Leipzig) to investigate distinct respiratory modulations of

periodic and aperiodic brain activity. Our rationale was twofold: First, we aimed to

characterise respiration phase-locked changes of aperiodic 1/f slope over parieto-occipital

cortices to corroborate previous findings of excitability changes in visual perception [9].

Second, we extended the scope for a general investigation of potentially independent
modulations of aperiodic and oscillatory components within the cortical RMBO network [2].
Here, we particularly aimed to assess whether both components of neural signalling would
be modulated by respiration and, if so, how these modulatory dynamics would be related to
one another. Our results highlight respiration as a slow physiological modulator of both
periodic and aperiodic cortical signalling, strengthening its previous implication in the
regulation of excitability states. Furthermore, the overall pattern of findings strongly suggests
differential temporal dynamics between the coupling of both components.

Results

We computed time-resolved fits of both aperiodic (1/f) and periodic (oscillatory) components
of resting-state brain activity to disentangle their dynamics in respiration-brain coupling. A
moving-window application of the novel SPRiNT algorithm [19] and the underlying *specparam*
procedure [18] to MEG and EEG data yielded individual time courses of aperiodic slopes and
periodic spectra for a total of N = 78 participants. Concurrent respiration recordings allowed
us to extract respiratory phase vectors that corresponded to each of the moving windows.
Next, we performed binning of respiratory phase and bin-wise averaging of corresponding 1/f
slope and periodic spectra on the group level (see Fig. 1 and Methods for details). We then
used these phase-specific estimates of slope and periodic spectra to statistically test if and
how periodic and aperiodic brain activity is modulated by respiration.

Fig. 1. Synopsis of acquired data and applied methods. a, In two labs, we simultaneously acquired nasal
respiration as well as eyes-open, resting-state MEG (IBB, Münster) and EEG data (MPI, Leipzig) in continuous 5-
min recordings. After preprocessing, single-sensor/-channel M/EEG data (middle panel) were subjected to the

SPRiNT algorithm [26] (top). Here, using a moving-window approach (window length = 1 s, 75% overlap between
neighbouring windows), spectral components of neural time series are estimated using a Fast Fourier transform.
These frequency-domain data are then parameterized using the *specparam* algorithm [18] which yields both
aperiodic and periodic components of neural activity in that time window. Repeating this procedure along the
entire recording thus yields time-resolved fits of the aperiodic 1/f slope as well as time-resolved periodic spectra
ranging from 1 - 40 Hz (top right). Respiratory phase was computed using two-way interpolation (int) of the
normalised raw respiration signal (peak-to-trough, trough-to-peak; bottom). **b**, For each time point used as a
moving-window centre in the *SPRiNT* algorithm, we then extracted the corresponding respiratory phase. This
allowed us to sort all time-resolved slope fits and periodic spectra according to the respiratory phase at which
they had been computed. In keeping with previous work [2,9] we finally partitioned the respiration cycle into n = 60
equidistant, overlapping phase bins and computed bin-wise averages of slope fits and periodic spectra. This
approach thus yielded quasi-continuous, respiration phase-resolved courses of both periodic and aperiodic
components of brain activity for each sensor/channel within each participant (bottom right).

Respiration relates to aperiodic neural activity over posterior cortices. The first
analysis was closely connected to a recent study showing respiration phase-dependent
modulations of cortical excitability in a visual detection task [9]. In line with a rich literature of
visual perception studies (see [20]), increased excitability was indexed by desynchronised
alpha oscillations (i.e., lower alpha power corresponding to higher excitability). However,
changes in alpha power can be confounded by changes in the slope of aperiodic
background activity [18] - itself a widely-used measure of cortical excitability and
excitation/inhibition balance [17]. Therefore, we aimed to disambiguate the effects of
respiration on periodic and aperiodic brain activity and started by first testing the hypothesis
that respiration-induced changes of excitability are reflected in cyclic changes of the slope of
aperiodic brain activity. For both MEG (Münster) and EEG data (Leipzig), we set up a linear
mixed-effects model (LMEM) to predict individual 1/f slopes of parieto-occipital average
power spectra by respiration phase. A null distribution of the resulting beta weights was
constructed by re-computing the LMEM based on individual surrogate respiration time series
k = 5000 times. Subsequently, we pooled both data sets in order to capitalise on higher
statistical power and to illuminate phase-locked modulations jointly found across modalities.
For both MEG and EEG as well as for the pooled data, the empirical group-level beta weight
for respiratory phase was significantly higher than the respective null beta weights (all $p <$
.001), strongly indicating a significant overall association between respiratory phase and
aperiodic neural signals over posterior cortices (Fig. 2). In order to characterise the temporal
dynamics of respiration phase-locked changes in 1/f slope, we repeated the binwise slope
estimation as outlined above, but this time on surrogate respiration time courses. For each
participant, we thus generated k = 5000 'null slope estimations' within each phase bin which
we used to extract percentiles for the empirical group-level bin averages relative to these
binwise 'null averages' (Fig. 2a; see Methods for details). For the pooled data from both labs,

we observed a steeper slope (indicating stronger inhibitory activity) around the expiration-to-
inspiration transition (ranging from -156° to 143° and around 131°). Conversely, the slope
was flatter (indicating stronger excitatory currents) during the inspiratory phase (between -
88° and -64°) and shortly after the inspiration-to-expiration transition (9° to 46°, see Fig. 2).
This general pattern was consistent across both data sets, although the flattening of the
slope during the inspiratory phase did not reach significance in the EEG data alone
(presumably due to slightly lower statistical power). Consistency across MEG and EEG data
was supported by a non-parametric two-sample test for circular data (akin to a Kruskal-
Wallis test for linear data), which showed that the circular means of MEG and EEG slope ~
respiration courses were not significantly different from one another ($P(77) = 0.21$, $p = .651$).
187

188
Fig. 2. Group-level respiration-locked modulation of aperiodic 1/f slope over parieto-occipital sensors. a,
Respiration phase was extracted for a total of n = 60 overlapping phase bins covering the entire respiration cycle
between subsequent inhalations. Overall significance of respiration phase-locked modulations was assessed by

means of permutation statistics: For each data set, the combined LMEM beta weight for the phase vector norm
(i.e., sqrt(sin² + cos²)) is shown in light blue against a distribution of null betas (k = 5000, top right corners). Within
each polar plot, coloured bold dots show respiration phase-dependent 1/f slope gained from SPRiNT
computations on periodic spectra over parieto-occipital MEG sensors or EEG channels, respectively. Concentric
scatter plots (black dots) indicate null distributions of group-level mean exponent values computed for each
phase bin (k = 5000 iterations). Solid black lines indicate the 5th and 95th percentile of each bin's null distribution.
In the MEG sample, 1/f slope was flattened during the inspiratory phase (around -70°) as well as around the
inspiration-to-expiration transition (-9° and 15° to 21°; blue markings). 1/f slope was steeper during the expiration
phase (82° to 88°) and around the expiration-to-inspiration transition (-174° to 162°; red markings). Similarly, the
EEG sample showed flatter slope after the inspiration-to-expiration transition (9° to 46° and around 58°) and
steeper slope around the expiration-to-inspiration transition (-162° to 131°). In = inspiration, ex = expiration. **b**,
Exemplary overlay of ROI-average 1/f slope estimates and respiratory phase over time for a single participant
from the MEG sample. Transparent black lines show single-sensor slope estimates, solid black line shows mean
slope across ROI sensors. Blueish-green scatter plot shows respiratory phase. **c**, Correlation between mean 1/f
slope and breathing rate (top) as well as breathing depth (bottom) for MEG (left) and EEG data (right). Neither
respiratory parameter was significantly correlated with overall steepness of the aperiodic component during rest.
d, For the analysis of normal vs deep breathing, we revisited a previously published MEG data set [3]. For both
breathing conditions, we show the respiratory trace of a single participant (left) and median group-level breathing
depths (right). As instructed, participants were breathing deeper during the deep (vs normal) breathing condition
($z(27) = 6.19$, $p < .001$). **e**, Using the same preprocessing and ROI analysis pipelines as in the previous analyses,
we tested whether the strength of 1/f slope modulations over posterior sensors differed between normal (i.e.,
automatic) and voluntary deep breathing. Indeed, the individual ranges of slope ~ respiration phase were
consistently greater for deep (compared to normal) breathing ($z(27) = 3.02$, $p = .003$). **f**, Comparison of parieto-
occipital 1/f slope ~ respiration phase during nasal (left) and oral breathing (right) collected in a follow-up control
study (see Methods for details). Using the identical experimental setup and analyses as in the original MEG data,
LMEMs confirmed a significant influence of respiratory phase on 1/f slope for both breathing conditions (see
histogram insets). 1/f slope ~ respiration courses were similar to our previous MEG and EEG results. **g**, Individual
ranges of slope ~ respiration (as a measure of modulation strength) did not differ between conditions ($z(25) =$
0.80, $p = .426$).

Aperiodic modulations as a function of respiratory parameters. We conducted different
control analyses to assess whether 1/f slope and its modulation strength covaried with
parameters like breathing rate, depth, or route (i.e., nasal vs oral breathing). First, in the
present data, individual 1/f slopes (averaged across the respiratory cycle) were not
significantly correlated with breathing rates (MEG: $r(40) = .-17$, $p = .297$; EEG: $r(38) = .13$, p
$= .418$) or breathing depths (MEG: $r(40) = .19$, $p = .230$; EEG: $r(38) = -.09$, $p = .591$) in either
data set (Fig. 2c). However, the interpretability of these results is somewhat limited by the
small amount of intraindividual variation in the respiratory time series - natural breathing is
simply too consistent to conclusively investigate whether different breathing 'modes' (e.g.,
deep vs normal breathing) would influence respiration-locked modulations in 1/f slope.
Therefore, we next re-analysed a previously published data set [3] in which we had recorded 5
minutes of whole-head resting-state MEG during both deep and normal nasal breathing (see

Fig 2d). In these data, a comparison of deep vs normal nasal breathing revealed that 1/f
slope over the posterior ROI was indeed more strongly modulated during deep (compared to
normal) breathing ($z(27) = 3.02$, $p = .003$, see Fig. 2e).
Finally, to investigate the role of breathing route, we compared respiration-locked 1/f slope
modulations at rest during nasal vs oral breathing in a new sample of N = 25 MEG data sets
(see Methods). Repeating the posterior ROI analysis described above, respective LMEMs
confirmed significant coupling of respiratory phase to 1/f slope during both nasal and oral
breathing (both $p < .001$; Fig. 2f). The modulatory pattern over phase was again similar to
the original findings (with certain limitations due to reduced power): During nasal breathing,
1/f slope was flattened during the inspiratory phase (-95° to -52°) as well as shortly after the
inspiration-to-expiration transition (around 36°) and steeper around the expiration-to-
inspiration transition (143° to -150°). During oral breathing, 1/f slope was flattened around
the inspiration-to-expiration transition (-64° to 52°) and steeper around the expiration-to-
inspiration transition (131° to -125°). A comparison of individual ranges of slope ~ respiration
(i.e., max_{slope} - min_{slope} over the respiratory cycle) revealed that the strength of respiration-
locked 1/f slope modulations did not significantly differ between nasal and oral breathing
($z(24) = 0.80$, $p = .426$).
251
Respiration-locked slope modulations are wide-spread, but strongest over posterior
sites. Overall, the first analysis demonstrated a consistent pattern of respiration-locked 1/f
slope modulations over parieto-occipital sensors: The slope of aperiodic neural activity was
not uniform over the respiratory cycle, but systematically modulated in such a way that 1/f
slope was flattened around the inspiration-to-expiration transition and steeper around the
expiration-to-inspiration transition. In the next step, we aimed to i) broaden the scope to
investigate respiration-locked slope changes across all M/EEG sensors and ii) characterise
both direction and strength of these modulations across the scalp.
For each participant, single-sensor/channel 1/f slope courses across 60 respiratory phase
bins were extracted using the pipeline described above on single-sensor time series. For
each of the N = 275 MEG sensors (Münster) and N = 62 EEG channels (Leipzig), we then
repeated the combined LMEM and permutation approach (see above). Thus, for each
sensor, the percentile of the empirical beta weight (relative to its null distribution constructed
from LMEMs on surrogate respiration) indicated significance of phase-locked 1/f slope
changes. Group-level average slope courses for n = 240 significant MEG sensors (top) and
n = 53 significant EEG channels (bottom) are shown in Fig. 3a.
To quantify whether there was a consistent modulation of 1/f slope across all
sensors/channels, we computed the circular mean of single-sensor/channel slope courses.
The histograms in Fig. 3a illustrate a clear clustering around the expiration-to-inspiration

transition (i.e., phase bins around ± π). Supporting our parieto-occipital findings from above,
Rayleigh tests on sensor-/channel-wise circular means corroborated that single-
sensor/channel slope courses were not uniform across respiratory phases, but instead
showed a consistent mean direction with the steepest slope around ± π for both data sets
(MEG: z = 126.86, *p* < .001; EEG: z = 14.38, *p* < .001). A topographic representation of
sensor-/channel-wise circular means is shown in Fig. 3b.

278
Fig. 3. Single-sensor (MEG) and channel (EEG) modulation of 1/f slope. a, Polar plots shows group-level
averages of normalised single-sensor/channel slope modulation courses over the respiration cycle. LMEM
revealed 1/f slope over respiration to significantly deviate from a uniform distribution in n = 240 out of 275 MEG
sensors and n = 53 out of 62 EEG channels. Only significant MEG sensors (top) and EEG channels (bottom) are
shown, colour coding illustrates sensors and channels 1 to n. Polar histograms illustrate circular mean directions
of single-sensor/channel slope distributions (colour-coded according to respiration phase). **b**, Topographic
representations of significant modulations and their phase direction for MEG sensors (top) and EEG channels
(bottom). **c**, Population statistics quantifying the effect size of slope modulations seen in b. For each
sensor/channel, we show the vector norm of LMEM beta weights for sine and cosine of the respiratory phase
(akin to a harmonic regression approach). These vector norms were normalised using the mean and standard
deviation of the null distribution and are thus given in z values (see Methods for details). In = inspiration, ex =
expiration.

Finally, in line with previous approaches [2,7], we quantified the strength of single-
sensor/channel phase-locked slope modulations relative to their null distributions: For each
sensor/channel, we computed z-scored LMEM beta weights by subtracting the mean of its
'null beta' distribution and dividing by the corresponding standard deviation (see Methods).
In both MEG and EEG topographies, the strongest modulations of aperiodic neural activity
were indeed observed over parieto-occipital cortices (Fig. 3c).

Periodic and aperiodic components of the RMBO network are distinctly coupled to respiration. In a final series of analyses, we aimed to disentangle the dynamics of respiratory coupling to periodic (i.e., oscillatory) and aperiodic (i.e., non-oscillatory) components of neural activity. So far, no investigation of respiration-brain coupling has directly considered either phase-locked changes of E:I balance or its distinction from oscillatory modulations. To target this question, we computed source-localised time series from the MEG data (mapped onto n = 230 parcels taken from the HCP atlas [21]) and defined regions of interest according to the original RMBO network publication [2]. For a total of n = 10 ROIs (consisting of n = 23 HCP parcels), we extracted respiration phase-locked 1/f slope as well as periodic spectra with the 1/f characteristic removed. For the computation of parcel-specific courses of aperiodic slope over respiratory phase, we followed the sensor-level approach outlined above. The periodic component was defined as the accumulated power of oscillatory peaks (between 1 - 40 Hz) after the 1/f slope was removed from the respective periodic spectrum.

In keeping with our sensor-level analyses, we determined significance of respiration phase-locked changes in aperiodic slope and oscillatory power by means of LMEMs within each parcel of interest. Again, each parcel's empirical vector norm of LMEM beta weights for respiratory sine and cosine was assessed relative to k = 5000 null beta weights from LMEMs computed on surrogate distributions of slope and power over respiratory phases. Of the 23 parcels of interest, 18 parcels showed a significant association between respiration and 1/f slope. Significant phase-locked modulations of oscillatory power (within frequencies from 1 - 40 Hz) were found for 19 parcels (see Fig. 4a). Overall, all 23 parcels showed either significant non-uniformity of slope or accumulated power.

Next, we investigated differences in temporal dynamics between respective slope and power changes within each parcel. Specifically, we extracted the circular means of each parcel's slope and power courses (binned into n = 60 overlapping respiratory phase bins) to compute parcel-wise phase differences between the two courses: In case slope and power were similarly modulated by respiration, the mean phase difference across parcels would be (close to) zero. As shown in the polar histogram in Fig. 4b, this was not the case ($U^2 = 1.26$, $p < .001$, 5000 permutations). Next, we tested whether slope and power courses of each parcel were drawn from the same distribution. For all parcels, Watson's U^2 permutation tests confirmed significant differences between the underlying distributions (all $p < .001$), providing further evidence for differential modulatory dynamics of slope and power courses within the RMBO network. For both slope and accumulated power, Fig. 4c shows ROI-wise distributions of subject-level U^2 statistics for each parcel as an estimate of respective non-

334 uniformity (centre panel). Exemplary comparisons between ROIs are shown for slope ~

335 respiration (left panel) and power spectra ~ respiration (right panel).

336
Fig. 4. Source-level ROI modulation of 1/f slope and accumulated oscillatory power in the MEG. a, For
each cortical node of the RMBO network [2], we show the group-level average courses of 1/f slope (left polar plots)
and accumulated spectral power after removal of the aperiodic component (1 - 40 Hz; right polar plots) over the
respiratory cycle. For ROIs comprising more than one parcel from the HCP atlas, slope and power courses are
superimposed for each individual parcel. In case the respective LMEM did not indicate significant association with
either slope or power within any given parcel, the corresponding time series are drawn with reduced opacity. For
significantly associated parcels only, red and blue triangles indicate respective maxima and minima over the
respiratory cycle (see legend). **b**, Polar histogram shows phase differences between the circular means of slope
and power courses across parcels. The clear majority of non-zero phase differences were corroborated by
Watson's U² statistics indicating distinct temporal dynamics of slope and power modulations within each parcel
(see main text). **c**, Centre: For each ROI, we show violin plots of single-participant U² statistics (slope/power vs
uniform distribution, respectively). Higher U² statistics indicate stronger non-uniformity. Left: Exemplary slope
modulations of SMA and OFC (normalised across respiration phase for illustration purposes). SMA was more
strongly non-uniform overall (see centre), which is visible from both stronger maxima (Δmax) and minima (Δmin)
compared to OFC. Right: Exemplary power modulations of OFC and l. INS. For both ROIs, we superimpose
power spectra (with 1/f removed) from all 60 phase bins. Overall, power varied more strongly with respiratory
phase within OFC compared to l.INS (see centre). TPJ = temporo-parietal junction, ITG/STG = inferior/superior

temporal gyrus, ANG = angular gyrus, FEF = frontal eye-field, TP = temporal pole, SMA = supplementary motor
area, OFC = orbitofrontal cortex, CUN = cuneus, LS = lingual sulcus, ACC/PCC = anterior/posterior cingulate
cortex, INS = insular cortex. In = inspiration, ex = expiration.

Finally, single-band modulations of 1/f-removed theta (4-8 Hz), alpha (8-13 Hz), and beta
power (13-30 Hz) within the RMBO network are shown in Fig. 5. Out of 23 parcels, band-
specific LMEMs revealed that theta power was significantly modulated in 17 parcels, alpha
power in 22 parcels, and beta power in 21 parcels (see Fig. 5a and Methods for details).
While band-specific oscillatory modulations were not the main focus of this paper, we
provide this characterisation to motivate hypothesis generation for future work.

Fig. 5. Source-level ROI modulation of band-specific oscillatory power in the MEG. a, For each cortical
node of the RMBO network, we show the group-level average courses of band-specific spectral power after
removal of the aperiodic component over the respiratory cycle. For ROIs comprising more than one parcel from
the HCP atlas, power courses are shown for each individual parcel. In case the band-specific LMEM did not
indicate significant modulation of the periodic component within any given parcel, the corresponding power series
are not shown. **b,** Exemplary ROI-average spectra of anterior cingulate (top) and left insula (bottom), colour-
coded according to respiratory phase. **c,** Group-level median U² statistics of each parcel testing single-band
power ~ respiration phase against a uniform distribution. Higher values indicate stronger deviation from uniform
distribution.

Discussion

Our present findings highlight the role of respiration as a physiological modulator of neural
signalling. We provide first evidence that fluctuations of *aperiodic* brain activity are phase-
locked to the respiratory cycle which strongly suggests that spontaneous state shifts of
excitation-inhibition balance are at least partly influenced by peripheral bodily signals.
Moreover, differential temporal dynamics in their coupling to non-oscillatory and oscillatory

activity raise the question whether there is a functional distinction in the way each
component is related to respiration.

Respiratory modulation of excitation-inhibition balance. While the mechanisms
underlying the 1/f characteristic of neural power spectra have yet to be fully understood [22],
crossmodal evidence has highlighted its functional significance in several domains. Most
prominently, the steepness of the aperiodic slope (i.e., its exponent) has been shown to
reflect the balance between excitatory (E) and inhibitory (I) synaptic currents in
computational [16], pharmaceutical [15], as well as optogenetic work [23]: The larger the exponent
(i.e., steeper power spectral slope), the stronger the inhibitory influence and vice versa [24]. In
general, constant E:I balancing is essential to maintain neural homeostasis [25] so that the
excitability within a particular neural array remains at a critical state [26].
Facilitated by novel methodological advances, our present findings of respiration-locked
changes in E:I balance thus illuminate a previously inaccessible aspect of what is emerging
as a recurrent motif in respiration-brain coupling across species: Animal studies have shown
excitability [27] as well as spike rates [12,13] to covary with the respiratory cycle, prompting non-
invasive human work from our lab [9] in which we linked respiratory phase to excitability of
visual cortices (measured as parieto-occipital alpha power) during low-level perception.
Given that breathing is under volitional control, we adopted the theoretical perspective of
predictive processing [28] to propose respiration as a means to actively align the sampling of
external sensory information with transient internal states of increased excitability. This
interpretation rests on the concept of *active sensing* [29] in rodents which recently gained
further support from a comprehensive demonstration by Karalis and Sirota [30]: In a series of
elaborate experiments, they found neocortical *up* and *down states* during sleep and quiet
rest to be modulated by respiratory phase, strongly suggesting that breathing influences
(sub-)cortical excitability in the sense of a pacemaker aligning neural network dynamics.
While the spontaneous aperiodic shifts of E:I balance in our paradigm are not to be equated
to the reported up and down states in quiescent animals, we propose that the shared
observation of respiration phase-locked state changes bears valuable insight into human
respiration-brain coupling.

Differential dynamics of respiration-brain coupling. Investigating the respiration phase-
locked changes in aperiodic vs periodic activity inevitably entails the more general distinction
of 1/f and genuine oscillations per se and how previous works have addressed one or the
other. To reiterate, neural oscillations are narrowband, rhythmic components embedded
within a background of broadband, aperiodic activity which itself does not require any
underlying rhythmicity [31,32]. So far, studies of breath-brain interactions have neglected

aperiodic activity entirely, most likely due to the fact that one proposed mechanism of
respiration-brain coupling critically relies on oscillatory entrainment to the breathing rhythm:
Here, the nasal airstream entrains neural activity within the olfactory tract via
mechanoreceptors, after which the phase of this infraslow rhythm is coupled to the amplitude
of faster oscillations [33,12]. This propagation through phase-amplitude coupling [34] has largely
been quantified by measures of coupling strength such as the modulation index (MI [35]), a
method that – despite its benefits - prohibits any conclusion regarding non-oscillatory
modulations.
In the present study, we consistently observed steepest 1/f slopes around ± π, with strongest
effects localised over posterior sensors, complementing previous reports of a spatial
gradient of aperiodic dynamics [18, 36]. Such a posterior-to-anterior gradient conceivably
increases parieto-occipital SNR, which could facilitate the detection of respiration phase-
locked changes. As for the pattern of preferred respiratory phase around ± π, He and
colleagues [37] demonstrated a rich temporal organisation of aperiodic (or 'scale-free') brain
activity in ECoG recordings from epilepsy patients: Extending the concept of hierarchically
nested oscillations [38], the authors showed nested frequencies in aperiodic activity during
quiet wakefulness; in other words, coupling of low-frequency phase to high-frequency
amplitude of 1/f slope. Intriguingly, the preferred phase clustered around phases 0 and ± π,
i.e., peaks and troughs of low-frequency fluctuations, closely resembling present findings of
flatter slopes (i.e., increased excitation) towards the end of inspiration as well as steeper
slopes (i.e., more inhibition) towards the end of expiration (see Figs. 2 and 3). These
aperiodic shifts of E:I state further corroborate previous findings of increased excitability
during inspiration [9] and improved cognitive performance for stimuli presented during the
inspiratory phase [6,7].
If the proposed functional mechanisms of respiration-brain coupling were indeed to be
extended to include aperiodic brain activity, respiratory modulations of 1/f slope should be
flexibly adapted during cognitive or perceptual tasks to facilitate performance. A study by
Waschke and colleagues [15] recently provided convincing evidence that the aperiodic
component itself not only captured task-relevant attentional changes in E:I balance, but also
covaried with individual behavioural performance. Critically, the authors used anaesthetics to
distinguish between oscillatory (i.e., increased band-specific power) and non-oscillatory,
aperiodic changes in EEG spectra. In a similar vein, Donoghue et al. [18] stressed the
importance of disentangling periodic and aperiodic changes in neural activity in order to
recognise their potentially distinct contributions to observed effects. We argue that the same
is true in order to more comprehensively describe the physiological underpinnings of
respiration-brain coupling.
454

Different mechanisms underlying non-oscillatory versus oscillatory modulations? In the present data, the preferred phase differences between source-level periodic and aperiodic modulations within the RMBO network show at least partly differential dynamics between both components, suggesting a more complex coupling of neural signalling to the respiratory rhythm than previously assumed. Conceivably, changes in periodic and aperiodic activity could reflect respiration-locked neural modulations arising from two (not necessarily) independent mechanisms: While the respiratory rhythm itself is generated in the brain stem [39], oscillatory changes on the one hand are assumed to be induced by mechanosensory feedback stimulation from the nasal airstream during inspiration and expiration (see above). Convincing evidence for this hypothesis comes from studies showing that both neural and behavioural modulations dissipate when there is no nasal airflow stimulating olfactory mechanoreceptors, i.e. during oral breathing [6,7] or in bulbectomised animals [12]. On the other hand, CO_2 levels continuously regulate blood flow and respiration rate in a feedforward fashion, which in turn determine changes in CO_2 levels [40]. Moreover, CO_2 is inversely related to tissue acidity (pH), which has been linked to neural excitability [41,42]. Lowered pH (due to increased CO_2) causes an increase in extracellular adenosine [43], a neuromodulator gating synaptic transmission. Consequently, Dulla and colleagues [44] concluded that CO_2-induced changes in neural excitability are caused by pH-dependent modulation of adenosine and ATP levels (also see [45)]).

Since CO_2 levels do not differ between nasal and oral breathing, the two mechanisms are clearly functionally distinct from one another: CO_2 fluctuations alone cannot account for neural or behavioural differences between nasal and oral breathing. Hence, there are (at least) two different ways in which neural activity could be coupled to the act of breathing. It may be the case that the differential coupling of respiration to periodic and aperiodic activity reflects these distinct mechanisms. However, the evidence so far (including present data from the control study) neither precludes the possibility that both types of modulation could be rooted in the same underlying mechanism. Therefore, critical future work is needed to further elucidate the functional and anatomical underpinnings of respiratory modulations in brain dynamics.

Clinical implications and future directions. The present findings open multiple avenues for future research, of which we want to briefly sketch three main directions. First, mechanistic advances could be made in a replicative study which includes a contrast of nasal vs oral respiration. The absence of phase-amplitude coupling driven by nasal airstreams during oral breathing could conceivably reveal to what extent aperiodic changes rely on and/or interact with oscillatory changes. While we speculate that respective changes in 1/f slope and oscillatory power could point to distinct modulatory pathways, targeted

follow-up studies with sufficient statistical power are of critical importance to extend our initial results.

Second, as outlined above, it would further be highly instructive to complement existing evidence of task-related 1/f slope changes [23,46] by investigating their link to respiration in behavioural cognitive or perceptual paradigms. Timed breathing (in healthy participants or ventilated patients) or contexts involving different brain states, e.g. resting-state vs task, during attentional or arousal manipulation (including hyperventilation), or during sleep stages, could potentially unravel the link between breathing and fluctuations of (non-)oscillatory brain activity.

Finally, a third line of research should focus on potential translational applications of respiration-brain coupling. While others have made a convincing case for studying respiratory involvement in clinical contexts before (for a recent review, see [47]), the results presented here demonstrate a novel link between bodily signals and aperiodic neural activity. Importantly, changes in 1/f slope itself - an indicator of E:I imbalance - have been implicated in neurological and psychiatric disorders like Alzheimer's disease [48], schizophrenia [49], autism spectrum disorder [50], and epilepsy [51]. Linking these known neural alterations to peripheral signals could provide substantial insight into body-brain interactions in health and disease.

Acknowledgements

We would like to thank Karin Wilken, Ute Trompeter, and Hildegard Deitermann for their invaluable assistance during data collection in Münster. This work was supported by the Interdisciplinary Center for Clinical Research (IZKF) of the medical faculty of Münster (Gro3/001/19). DSK (KL-3580/1-1), AV (Research Training Group 2386), and JG (GR 2024/5-1) were further supported by the DFG.

Author Contributions

Conceptualization, DSK; Methodology, DSK, CF, JG; Investigation, DSK, CF; Writing – Original Draft, DSK, CF; Writing – Review & Editing, DSK, CF, OA, NC, AV, JG; Visualization, DSK; Funding Acquisition, DSK, AV, JG

Declaration of Interest

The authors declare no competing interests.

References

1. Jelinčić, V., Van Diest, I., Torta, D. M. & von Leupoldt, A. The breathing brain: The

potential of neural oscillations for the understanding of respiratory perception in health
and disease. *Psychophysiology* e13844 (2021) doi:10.1111/psyp.13844.

2. Kluger, D. S. & Gross, J. Respiration modulates oscillatory neural network activity at
rest. *PLoS Biol.* **19**, e3001457 (2021).

3. Kluger, D. S. & Gross, J. Depth and phase of respiration modulate cortico-muscular
communication. *Neuroimage* **222**, 117272 (2020).

4. Rassler, B. & Raabe, J. Co-ordination of breathing with rhythmic head and eye
movements and with passive turnings of the body. *Eur. J. Appl. Physiol.* **90**, 125–130
(2003).

5. Johannknecht, M. & Kayser, C. The influence of the respiratory cycle on reaction times
in sensory-cognitive paradigms. *Sci. Rep.* **12**, 2586 (2022).

6. Perl, O. *et al.* Human non-olfactory cognition phase-locked with inhalation. *Nat. Hum.*
Behav. **3**, 501–512 (2019).

7. Zelano, C. *et al.* Nasal respiration entrains human limbic oscillations and modulates
cognitive function. *J. Neurosci.* **36**, 12448–12467 (2016).

8. Grund, M. *et al.* Respiration, heartbeat, and conscious tactile perception. *J. Neurosci.*
42, 643–656 (2022).

9. Kluger, D. S., Balestrieri, E., Busch, N. A. & Gross, J. Respiration aligns perception with
neural excitability. *eLife* **10**, (2021).

10. Galvez-Pol, A., McConnell, R. & Kilner, J. M. Active sampling in visual search is coupled
to the cardiac cycle. *Cognition* **196**, 104149 (2020).

11. Biskamp, J., Bartos, M. & Sauer, J.-F. Organization of prefrontal network activity by
respiration-related oscillations. *Sci. Rep.* **7**, 45508 (2017).

12. Ito, J. *et al.* Whisker barrel cortex delta oscillations and gamma power in the awake
mouse are linked to respiration. *Nat. Commun.* **5**, 3572 (2014).

13. Yanovsky, Y., Ciatipis, M., Draguhn, A., Tort, A. B. L. & Brankačk, J. Slow oscillations in
the mouse hippocampus entrained by nasal respiration. *J. Neurosci.* **34**, 5949–5964
(2014).

14. He, B. J. Scale-free brain activity: past, present, and future. *Trends Cogn Sci (Regul Ed)*
18, 480–487 (2014).

15. Waschke, L. *et al.* Modality-specific tracking of attention and sensory statistics in the
human electrophysiological spectral exponent. *eLife* **10**, (2021).

16. Gao, R., Peterson, E. J. & Voytek, B. Inferring synaptic excitation/inhibition balance from
field potentials. *Neuroimage* **158**, 70–78 (2017).

17. Trakoshis, S., Rocchi, F., Canella, C., You, W., Chakrabarti, B., Ruigrok, A. N., ... &
MRC AIMS Consortium. Intrinsic excitation-inhibition imbalance affects medial prefrontal
cortex differently in autistic men versus women. *elife*, *9*, e55684 (2020).

18. Donoghue, T. *et al.* Parameterizing neural power spectra into periodic and aperiodic
components. *Nat. Neurosci.* **23**, 1655–1665 (2020).

19. Wilson, L. E., da Silva Castanheira, J., & Baillet, S. Time-resolved parameterization of
aperiodic and periodic brain activity. *eLife*, *11*, e77348 (2022).

20. Samaha, J., Iemi, L., Haegens, S. & Busch, N. A. Spontaneous Brain Oscillations and
Perceptual Decision-Making. *Trends Cogn Sci (Regul Ed)* **24**, 639–653 (2020).

21. Glasser, M. F. *et al.* A multi-modal parcellation of human cerebral cortex. *Nature* **536**,
171–178 (2016).

22. Buzsáki, G., Anastassiou, C. A. & Koch, C. The origin of extracellular fields and
currents--EEG, ECoG, LFP and spikes. *Nat. Rev. Neurosci.* **13**, 407–420 (2012).

23. Chini, M., Pfeffer, T., & Hanganu-Opatz, I. An increase of inhibition drives the
developmental decorrelation of neural activity. *eLife*, *11*, e78811 (2022).

24. Voytek, B. *et al.* Age-Related Changes in 1/f Neural Electrophysiological Noise. *J.*
Neurosci. **35**, 13257–13265 (2015).

25. Turrigiano, G. G. & Nelson, S. B. Homeostatic plasticity in the developing nervous
system. *Nat. Rev. Neurosci.* **5**, 97–107 (2004).

26. Xue, M., Atallah, B. V. & Scanziani, M. Equalizing excitation-inhibition ratios across
visual cortical neurons. *Nature* **511**, 596–600 (2014).

27. Dulla, C. G. *et al.* Adenosine and ATP link PCO2 to cortical excitability via pH. *Neuron*
48, 1011–1023 (2005).

28. Mumford, D. On the computational architecture of the neocortex. II. The role of cortico-
cortical loops. *Biol. Cybern.* **66**, 241–251 (1992).

29. Wachowiak, M. All in a sniff: olfaction as a model for active sensing. *Neuron* **71**, 962–
973 (2011).

30. Karalis, N. & Sirota, A. Breathing coordinates cortico-hippocampal dynamics in mice
during offline states. *Nat. Commun.* **13**, 467 (2022).

31. Buzsáki, G. *Rhythms of the Brain.* (Oxford University Press, 2006).
doi:10.1093/acprof:oso/9780195301069.001.0001.

32. Bullock, T. H., Mcclune, M. C. & Enright, J. T. Are the electroencephalograms mainly
rhythmic? Assessment of periodicity in wide-band time series. *Neuroscience* **121**, 233–
252 (2003).

33. Fontanini, A. & Bower, J. M. Slow-waves in the olfactory system: an olfactory
perspective on cortical rhythms. *Trends Neurosci.* **29**, 429–437 (2006).

34. Canolty, R. T. & Knight, R. T. The functional role of cross-frequency coupling. *Trends*
Cogn Sci (Regul Ed) **14**, 506–515 (2010).

35. Tort, A. B. L. *et al.* Dynamic cross-frequency couplings of local field potential oscillations
in rat striatum and hippocampus during performance of a T-maze task. *Proc Natl Acad*
Sci USA **105**, 20517–20522 (2008).

36. Mahjoory, K., Schoffelen, J.-M., Keitel, A. & Gross, J. The frequency gradient of human
resting-state brain oscillations follows cortical hierarchies. *eLife* **9**, (2020).

37. He, B. J., Zempel, J. M., Snyder, A. Z. & Raichle, M. E. The temporal structures and
functional significance of scale-free brain activity. *Neuron* **66**, 353–369 (2010).

38. Lakatos, P. *et al.* An oscillatory hierarchy controlling neuronal excitability and stimulus
processing in the auditory cortex. *J. Neurophysiol.* **94**, 1904–1911 (2005).

39. Del Negro, C. A., Funk, G. D. & Feldman, J. L. Breathing matters. *Nat. Rev. Neurosci.*
19, 351–367 (2018).

40. Feldman, J.L., Mitchell, G.S., & Nattie, E.E. Breathing: rhythmicity, plasticity,
chemosensitivity. *Annu. Rev. Neurosci. 26*, 239–266 (2003).

41. Chesler, M. Regulation and modulation of pH in the brain. *Physiol. Rev. 83*, 1183–1221
(2003).

42. Lee, J., Taira, T., Pihlaja, P., Ransom, B.R., & Kaila, K. Effects of CO_2 on excitatory
transmission apparently caused by changes in intracellular pH in the rat hippocampal
slice. *Brain Res. 706*, 210–216 (1996).

43. Ito, H., Kanno, I., Ibaraki, M., Hatazawa, J., & Miura, S. Changes in human cerebral
blood flow and cerebral blood volume during hypercapnia and hypocapnia measured by
positron emission tomography. *J. Cereb. Blood Flow Metab. 23*, 665–670 (2003).

44. Dulla, C.G., Dobelis, P., Pearson, T., Frenguelli, B.G., Staley, K.J., & Masino, S.A.
Adenosine and ATP link PCO_2 to cortical excitability via pH. *Neuron 48*, 1011–1023
(2005).

45. Gourine, A.V., Llaudet, E., Dale, N., & Spyer, K.M. ATP is a mediator of chemosensory
transduction in the central nervous system. *Nature 436*, 108–111 (2005).

46. Ouyang, G., Hildebrandt, A., Schmitz, F. & Herrmann, C. S. Decomposing alpha and 1/f
brain activities reveals their differential associations with cognitive processing speed.
Neuroimage **205**, 116304 (2020).

47. Heck, D. H. *et al.* Recent insights into respiratory modulation of brain activity offer new
perspectives on cognition and emotion. *Biol. Psychol.* **170**, 108316 (2022).

48. Ramírez-Toraño, F. *et al.* Functional connectivity hypersynchronization in relatives of
alzheimer's disease patients: an early E/I balance dysfunction? *Cereb. Cortex* **31**,
1201–1210 (2021).

49. Molina, J. L. *et al.* Memantine effects on electroencephalographic measures of putative
excitatory/inhibitory balance in schizophrenia. *Biol. Psychiatry Cogn. Neurosci.*
Neuroimaging **5**, 562–568 (2020).

50. Rubenstein, J. L. R. & Merzenich, M. M. Model of autism: increased ratio of
excitation/inhibition in key neural systems. *Genes Brain Behav.* **2**, 255–267 (2003).
51. van Heumen, S. *et al.* Case Report: Aperiodic Fluctuations of Neural Activity in the Ictal
MEG of a Child With Drug-Resistant Fronto-Temporal Epilepsy. *Front. Hum. Neurosci.*
15, 646426 (2021).

642
643
644
645

646   **Methods**

Participants and data acquisition (Münster). Forty right-handed volunteers (21 female,
age 25.1 ± 2.7 y [mean ± SD]) participated in the study. All participants reported having no
respiratory or neurological disease and gave written informed consent prior to all
experimental procedures. The study was approved by the local ethics committee of the
University of Münster (approval ID 2018-068-f-S). Participants were seated upright in a
magnetically shielded room while we simultaneously recorded 5 minutes of MEG and
respiratory data. MEG data was acquired using a 275 channel whole-head system (OMEGA
275, VSM Medtech Ltd., Vancouver, Canada) at a sampling frequency of 600 Hz. During
recording, participants were to keep their eyes on a fixation cross centred on a projector
screen placed in front of them. To minimise head movement, participants' heads were
stabilised   with   cotton   pads   inside   the   MEG   helmet.
Participants were instructed to breathe naturally through their nose while the respiratory
signal was measured as thoracic circumference by means of a respiration belt transducer
(BIOPAC Systems, Goleta, USA) placed around their chest. Continuous monitoring via video
ensured participants were breathing through their nose instead of their mouth. Individual
respiration time courses were visually inspected for irregular breathing patterns such as
breath holds or unusual breathing frequencies, but no such artefacts were detected.

Participants and data acquisition (Leipzig). Thirty-eight healthy volunteers (18 female,
age 27.1 ± 4.0 y [mean ± SD]) were recruited from the database of the Max Planck Institute
for Human Cognitive and Brain Sciences, Leipzig, Germany. Participants reported no history
or current neurological or psychological condition. The study was approved by the Ethical
Committee of the University of Leipzig's Medical Faculty (No. 462-15-28082020). All
participants provided written informed consent and received financial compensation for their
participation in the study. Participants were seated upright in an EEG booth while 5 minutes
of EEG and respiration were recorded. EEG was recorded from 62 scalp positions

distributed over both hemispheres according to the international 10–10 system, using a
commercial EEG acquisition system (ActiCap Snap, BrainAmp; Brain Products). The mid-
frontal electrode (FCz) was used as the reference and a mid-frontal electrode placed on the
middle part of the forehead (between FP1 and FP2) as ground. Electrode TP9 was used to
measure ECG and electrode TP10 captured eye movements. Electrode impedance was kept
below 10 kΩ for all channels. EEG was sampled with a rate of 2.5 kHz and online bandpass-
filtered between 0.015 - 1000 Hz. Participants were instructed to keep their eyes open and
fixate a cross on the screen in front of them to avoid excess eye movements. Participants
were instructed to breathe naturally while respiration was measured with a respiration belt
transducer (BIOPAC Systems, Goleta, USA) placed around the chest.

Participants and data acquisition (control MEG study for deep vs normal breathing).
The MEG sample for the comparison of deep vs normal breathing has previously been
published elsewhere [3]. Twenty-eight volunteers (14 female, age 24.8 ± 2.9 y [mean ± SD])
participated in the study conducted at the Institute for Biomagnetism and Biosignal Analysis
in Münster. All participants reported having no respiratory or neurological disease and gave
written informed consent prior to all experimental procedures. The study was approved by
the local ethics committee of the University of Münster (approval ID 2018-068-f-S). MEG
recording parameters and procedures were identical to the original recordings described
above.
Data were acquired in two 5-minute runs with a short intermediate break (determined by the
participants). Within each run, participants were instructed to either breathe normally or
voluntarily deeply (with their mouth closed) while maintaining their normal respiration
frequency, i.e. not to breathe more slowly during blocks of deep breathing. Continuous
monitoring via video ensured participants did indeed keep their mouth closed throughout the
recording. The order of deep and normal breathing was counterbalanced across participants.
For both runs, we recorded the respiratory signal as thoracic circumference by means of a
respiration belt transducer (BIOPAC Systems, Goleta, USA) placed around the participant's
chest. Individual respiration time courses were visually inspected for irregular breathing
patterns such as breath holds or unusual breathing frequencies, but no such artefacts were
detected.

Participants and data acquisition (control MEG study for nasal vs oral breathing).
Twenty-five right-handed volunteers (10 female, age 26.3 ± 3.3 y [mean ± SD]) participated
in the control MEG study conducted at the Institute for Biomagnetism and Biosignal Analysis
in Münster. All participants reported having no respiratory or neurological disease and gave
written informed consent prior to all experimental procedures. The study was approved by

the local ethics committee of the University of Münster (approval ID 2021-785-f-S). MEG

recording parameters and procedures were identical to the original recordings described

above.

For each participant, we recorded two 6-minute runs of resting state activity: In one run,

participants were instructed to breathe naturally through their nose. Continuous monitoring

via video ensured participants did indeed keep their mouth closed throughout the recording.

In the other run, participants were instructed to breathe through their mouth while wearing a

nose clip to prevent nasal breathing. The order of nasal and oral breathing was

counterbalanced across participants. Again, we recorded the respiratory signal as thoracic

circumference by means of a respiration belt transducer (BIOPAC Systems, Goleta, USA)

placed around the participant's chest. Individual respiration time courses were visually

inspected for irregular breathing patterns such as breath holds or unusual breathing

frequencies, but no such artefacts were detected.

MRI acquisition and co-registration (Münster). For MEG source localisation, we obtained

high-resolution structural magnetic resonance imaging (MRI) scans in a 3T Magnetom

Prisma scanner (Siemens, Erlangen, Germany). Anatomical images were acquired using a

standard Siemens 3D T1-weighted whole-brain MPRAGE imaging sequence (1×1×1 mm

voxel size, TR = 2,130ms, TE = 3.51 ms, 256 × 256 mm field of view, 192 sagittal slices).

MRI measurement was conducted in supine position to reduce head movements, and

gadolinium markers were placed at the nasion as well as left and right distal outer ear canal

positions for landmark-based co-registration of MEG and MRI coordinate systems. Data

preprocessing was performed using Fieldtrip[52] running in MATLAB R2021a (The Mathworks,

Natick, USA). Individual raw MEG data were visually inspected for jump artefacts and bad

channels, but neither were detected. Both MEG and respiration data were resampled to 300

Hz prior to further analyses.

Co-registration of structural T1 MRIs to the MEG coordinate system was done for each

participant by initial identification of three anatomical landmarks (nasion, left and right pre-

auricular points) in their individual MRI. Using the implemented segmentation algorithms in

Fieldtrip and SPM12, individual head models were constructed from anatomical MRIs. A

solution of the forward model was computed using the realistically-shaped single-shell

volume conductor model[53] with a 5 mm grid defined in the Human Connectome Project

(HCP) template brain[21] after linear transformation to the individual MRI.

Respiratory preprocessing (both labs). To obtain continuous respiration phase angles, we

used Matlab's *findpeaks* function (with minimal peak prominence set to 1) to identify time

points of peak inspiration (peaks) and peak expiration (troughs) in the normalised respiration

time course. Phase angles were linearly interpolated from trough to peak ($-\pi$ to 0) and peak

to trough (0 to π) in order to yield respiration cycles centred around peak inspiration (i.e.,

phase 0).

In order to assess the potential influence of individual breathing parameters on aperiodic

fluctuations, we computed breathing rates and depths for each participant. Breathing rates

were extracted as the mean distance between inspiratory peaks as defined by the peak

detection algorithm described above. Breathing depths were computed as the integral of the

individual respiration time series, normalised by the number of breathing cycles during the

recording.

Head movement correction (Münster). In order to rule out head movement as a potential

confound in our analyses, we used a correction method established by Stolk and colleagues

[54]. This method uses the accurate online head movement tracking that is performed by our

acquisition system during MEG recordings. This leads to six continuous signals (temporally

aligned with the MEG signal) that represent the x, y, and z coordinates of the head centre

(H_x, H_y, H_z) and the three rotation angles (H_ψ, H_θ, H_φ) that together fully describe head

movement. We constructed a regression model comprising these six 'raw' signals as well as

their derivatives and, from these 12 signals, the first-, second-, and third-order non-linear

regressors to compute a total of 36 head movement-related regression weights (using a

third-order polynomial fit to remove slow drifts). This regression analysis was performed on

the power spectra of single-sensor and single-voxel time courses, respectively, removing

signal components that can be explained by translation or rotation of the head with respect

to the MEG sensors.

Respiration-locked computations of 1/f slope and oscillatory power. Our first aim was

to further investigate excitability changes over posterior cortices we previously reported [9]. To

this end, we defined parieto-occipital regions of interest for movement-corrected MEG (k =

41 sensors) as well as EEG data (k = 17 channels). Single-sensor/channel time series within

these ROIs were entered into the SPRiNT algorithm [19] with default parameter settings and

subsequently averaged. In short, SPRiNT is based on the *specparam* algorithm [18] and uses

a short-time Fourier transform (frequency range 1-40 Hz) to compute aperiodic and periodic

components of neural time series within a moving window (width of 1 s, 75% overlap

between two neighbouring windows). Due to the slow nature of the respiratory signal, we did

not average between neighbouring windows at this point (see below for details).

SPRiNT thus yielded time series of both aperiodic (i.e., 1/f exponent) and periodic signals

(i.e., oscillatory power) with a temporal resolution of 250 ms and a frequency resolution of 1

Hz. In order to relate these time series to the respiratory signal, we extracted respiratory

phase at all time points for which slope and power were fitted (i.e., the centres of each

moving window). Following previous work [9,2], we then partitioned the entire respiratory cycle

($-\pi$ to π) into n = 60 equidistant, overlapping phase bins. Moving along the respiration cycle

in increments of $\Delta\omega = \pi/30$, we collected all SPRiNT outputs (i.e., slope fits and periodic

spectra) computed at a respiration angle of $\omega \pm \pi/10$. At this point, we computed individual

bin-wise averages of 1/f slope and periodic spectra, yielding quasi-continuous 'phase

courses' of aperiodic and periodic neural signals for each participant. For the control MEG

studies, we separately applied this pipeline to the MEG data recorded during deep/normal

and nasal/oral breathing, respectively. The methodological approach is illustrated in Fig. 1.

For the source-level MEG analyses shown in Figs. 4 and 5, SPRiNT computed 1/f slope fits

and periodic spectra based on the source-localised time series of individual parcels from the

HCP atlas.

Statistical analysis of sensor-level 1/f slope modulation. As a first analysis of respiration

phase-locked changes in 1/f slope, we computed the following linear mixed-effects model

(LMEM):

802                 $SE_j = \square_0 + (S_{1,j} + \square_1) * \sin_{resp} + (S_{2,j} + \square_2) * \cos_{resp} + e_j$

803   (1)

805   For each participant *j*, the model predicted individual slope (resulting from SPRiNT

806   computations, see above) as a combination of the intercept ($\square_0$), fixed effects of respiratory

807   sine and cosine ($\square_1$, $\square_2$), and an error term ($e_j \sim N(0,\sigma^2)$). In line with previous work [9],

808   resulting beta weights for sine and cosine of the respiratory signal were combined in a

809   respiratory phase vector norm, i.e.

811                    $v = \sqrt{\beta_1^2 + \beta_2^2}$ (2)

To test whether 1/f was significantly modulated over the ROI sensors, we computed k =
5000 random iterations of the LMEM by shuffling bin-wise 1/f slope fits on the individual
level. This way, any meaningful relations between 1/f slope and respiratory phase were
removed, thereby specifically testing the hypothesis that 1/f slope changes significantly with
respiratory phase. For each of these iterations, we once again computed the phase vector
norm from the resulting beta weights, yielding a random distribution of k = 5000 'null vector
norms'. Significance of the empirical vector norm was determined by its percentile relative to
this null distribution.

As shown in Figs. 2 and 3, for the sensor-level analyses, SPRiNT parameterisation was
conducted on individual MEG sensors (Münster) and EEG channels (Leipzig). For the
analysis of 1/f slope modulations within the parieto-occipital ROI, we computed mean
SPRiNT estimates across the corresponding sensors or channels, respectively. LMEMs
were then computed on ROI-average estimates for 1/f slope ~ respiration phase for each lab
individually (Münster, Leipzig) as well as for the pooled data from both labs (see Fig. 2). For
the whole-scalp analyses shown in Fig. 3, LMEMs were computed on single-sensor/channel
SPRiNT   estimates.
Having established an overall significant influence of respiratory phase on 1/f slope with the
LMEM approach described above, we next aimed to characterise *when* (i.e., at which
respiratory phases) 1/f slope was significantly steeper or flatter. To this end, we implemented
a permutation approach as follows: For each participant, we constructed a surrogate
respiration time series using the iterated amplitude-adjusted Fourier transform (IAAFT [55]). In
contrast to shuffling the respiration time series, this iterative procedure preserves the
temporal autocorrelation of the signal, which is critical for constructing a fitting null
distribution for permutation testing. From these IAAFT-transformed respiration time series,
we extracted the surrogate respiratory phase values corresponding to each slope estimation
from SPRiNT. In keeping with the approach above, we finally binned all SPRiNT outputs into
n = 60 equidistant, overlapping phase bins covering one entire respiratory cycle (-π to π)
and computed the bin-wise average slope fit to yield a 'null time series' of 1/f slope over
surrogate respiration phase. For each participant, this procedure was repeated 5000 times
and resulted in a null distribution of 5000 surrogate 1/f slope estimates x 60 phase bins. We
then computed the group-level average null distribution of 1/f slope for each of the 60 phase
bins which allowed us to extract the percentiles of empirical bin-wise group means relative to
these null distributions. As illustrated in Fig. 2, empirical group-level means were deemed
significant if they exceeded the 95[th] percentile (or fell below the 5[th] percentile) of the null
distribution.
848

For single-sensor slope analyses (Fig. 3) as well as single-parcel analyses of 1/f slope and
periodic spectra (Fig. 4), we used the *circstat* toolbox for Matlab [56] to compute the circular
means of group-level average slope (and power) courses over the respiratory cycle. In the
absence of meaningful slope modulation by respiratory phase, a polar illustration of the
binned slope estimates would be a uniform circle. In contrast, any phase-locked slope
changes would be represented as modulations or peaks within the polar representation.
Albeit not for all distributions, the circular mean provides at least some indication with regard
to the direction of such modulations (see below for alternative measures). Therefore, we
used it to characterise the consistency of slope modulations across the whole scalp both
within MEG/EEG data as well as across both modalities and labs (see Fig. 3).
In addition to potential respiration-related modulations of slope courses, we quantified the
strength of these phase-locked modulations. To this end, we computed z scores from
sensor-specific LMEM beta weights of the respiratory phase vector norm (see above for
details) by subtracting the mean of its null distribution and dividing by the corresponding
standard   deviation:

865
$$z = \frac{v - \mu_{v(null)}}{\sigma_{v(null)}}$$
(3)

with *v* being the vector norm of beta weights for respiratory sine and cosine (see Eq. 2). This
way, empirical LMEM beta weights for the respiratory phase vector are given in units of
standard deviation relative to their null distribution.

Source reconstruction. Source reconstruction was performed using the linearly
constrained minimum variance (LCMV) beamformer approach [57] with the lambda
regularisation parameter set to 5%. This approach estimates a spatial filter for each location
of the 5 mm grid along the direction yielding maximum power. A single broadband LCMV
beamformer was used to estimate voxel-level activities across all frequencies.

ROI-based analyses of source-level 1/f slope and oscillatory power. For the
investigation of source-level modulations of 1/f slope and periodic spectra, we first extracted
neural time series from a total of n = 230 cortical parcels from the HCP atlas [21]. In order to
ground our analyses in previous findings, we then focussed our analyses on those parcels
representing the cortical nodes of the RMBO network [2]. This network has been
demonstrated to show respiration-modulated brain oscillations (RMBOs) during resting state,
which is why it was particularly well-suited to investigate potentially distinct dynamics of
periodic and aperiodic modulations. As some RMBO nodes comprised bilateral or wider-

spread anatomical sites, a total of N = 10 ROIs consisting of n = 23 parcels were used for
source-level analyses (see Fig. 4). For the average time series within these parcels, we
applied the SPRiNT analysis described above to yield both 1/f slope and oscillatory power
across the respiratory cycle. For the latter, SPRiNT computed periodic spectra between 1 -
40 Hz with the aperiodic component removed, i.e., solely the power of the periodic neural
signal. Just as with the 1/f exponents, we sorted all periodic spectra with respect to the
corresponding respiratory phase to yield individual bin-average periodic spectra for n = 60
overlapping phase bins. These spectra thus reflect, for each phase of the respiratory cycle,
the extent of 'true' oscillatory activity remaining after the underlying 1/f characteristic is
removed. In a first step, we quantified the amount of oscillatory activity by computing the
accumulated sum over all frequencies (illustrated in Fig. 4). For frequency band-specific
power courses, we then computed accumulated power for canonical bands separately
(theta: 4 - 8 Hz, alpha: 8 - 13 Hz, beta: 13 - 30 Hz).

Next, we aimed to investigate potentially distinct respiration phase-locked dynamics of
periodic and aperiodic modulations. Therefore, for each HCP parcel, we implemented
Watson's U^2 tests with k = 5000 random permutations (*watsons_U2_perm_test* function [58] for
Matlab) to test the null hypothesis that the group-level average vectors of slope and power
were drawn from the same distribution. We chose the U^2 test as an additional measure as it
is sensitive towards multimodal distributions [59], which is not the case for the circular mean
alone. For the sake of comparability, however, we also computed phase differences between
each parcel's circular means of slope and power courses: If 1/f slope and oscillatory power
were similarly coupled to the respiratory rhythm, their respective courses over respiratory
phase would not differ and the phase difference between their circular means would be
(close to) zero. In contrast, systematically different distributions of slope and power would
suggest differential dynamics in their respective coupling to respiratory phase.

Finally, we characterised the strength of slope and power modulations for each of the n = 10
RMBO ROIs by means of U^2 statistics. For each participant, we conducted Watson's U^2 tests
of the (potentially multimodal) courses of 1/f slope and accumulated power within each ROI
against a uniform distribution. Here, higher U^2 values indicate stronger non-uniformity within
a given parcel, which allowed us to characterise RMBO ROIs by their respective group-level
U^2 distributions (as shown in Fig. 4c).

919   **Data availability**

920   The anonymised data supporting the findings of this study will be made publically available
921   on the Open Science Framework (https://osf.io/8nw9t/) upon acceptance of the manuscript.
922

Code availability

924   All custom Matlab code to reproduce the central findings of this study will be made publically
925   available on the Open Science Framework (https://osf.io/8nw9t/) upon acceptance of the
926   manuscript.
927

References

52. Oostenveld, R., Fries, P., Maris, E. & Schoffelen, J.-M. FieldTrip: Open source software
for advanced analysis of MEG, EEG, and invasive electrophysiological data. *Comput.*
Intell. Neurosci. **2011**, 156869 (2011).

53. Nolte, G. The magnetic lead field theorem in the quasi-static approximation and its use
for magnetoencephalography forward calculation in realistic volume conductors. *Phys.*
Med. Biol. **48**, 3637–3652 (2003).

54. Stolk, A., Todorovic, A., Schoffelen, J.-M. & Oostenveld, R. Online and offline tools for
head movement compensation in MEG. *Neuroimage* **68**, 39–48 (2013).

55. Theiler, J., Eubank, S., Longtin, A., Galdrikian, B., & Farmer, J. D. (1992). Testing for
nonlinearity in time series: the method of surrogate data. *Physica D: Nonlinear*
Phenomena, *58*(1-4), 77-94.

56. Berens, P. circstat□: a *MATLAB* toolbox for circular statistics. *J. Stat. Softw.* **31**, (2009).

57. Van Veen, B. D., van Drongelen, W., Yuchtman, M. & Suzuki, A. Localization of brain
electrical activity via linearly constrained minimum variance spatial filtering. *IEEE Trans*
Biomed Eng **44**, 867–880 (1997).

58. Mégevand, P. pierremegevand/watsons_u2
https://github.com/pierremegevand/ watsons_u2, GitHub. Retrieved July 5, 2022. (2022)

59. Landler, L., Ruxton, G. D. & Malkemper, E. P. Advice on comparing two independent
samples of circular data in biology. *Sci. Rep.* **11**, 20337 (2021).