Local fat content determines global and local stiffness in fatty livers

[Fat accumulation locally softens liver]

David Li[1,2], Paul A. Janmey[2,3,4,5], and Rebecca G. Wells[1,2,3]

[1]Division of Gastroenterology and Hepatology, Department of Medicine
[2]NSF Science and Technology Center for Engineering MechanoBiology
[3]Department of Bioengineering
[4]Institute for Medicine and Engineering
[5]Department of Physiology
University of Pennsylvania, Philadelphia, PA

Grant Support:
This work was supported by NIH grant R01EB017753 to PAJ and RGW and by the Center for Engineering MechanoBiology (CEMB), an NSF Science and Technology Center, under grant agreement CMMI1548571.

Abbreviations:
NAFLD, non-alcoholic fatty liver disease; NASH, non-alcoholic steatohepatitis; DAPI, 4',6'-diamidino-2-phenylindole

Correspondence:
Rebecca G. Wells
905 BRB II/III
421 Curie Boulevard
Philadelphia, PA 19104-6140
Email: rgwells@pennmedicine.upenn.edu
Tel.: +1-215-573-1860

Disclosures:
The authors disclose no conflicts.

Data Transparency:
All study data are included in the article and/or supplementary material, and all analytic methods are available upon request from the authors.

Abstract

Nonalcoholic fatty liver disease (NAFLD) refers to a spectrum of pathology characterized by steatosis and other changes to the liver not associated with heavy alcohol use. NAFLD is the most common form of liver disease globally, affecting over 80 million people in the United States. The contribution of mechanical changes resulting from fat accumulation to NAFLD progression is unclear and the effect of steatosis on the mechanical properties of the liver has yet to be defined. We studied the livers of ob/ob and wild-type littermate mice using parallel plate rheometry as well as a sub-millimeter scale microindentation technique to define the stiffness of steatotic livers. The microindentation technique was adapted to enable determination of local fat content and identification of other local features in the indented region. We found that steatotic livers and steatotic regions of fatty livers were softer than non-steatotic livers/regions. Higher lipid content in a region correlated with decreased stiffness. The finding that steatosis is associated with both global and local liver softening has significant implications for understanding the progression of early stage NAFLD, given that stiffening is a mechanical stimulus for the development of fibrosis.

Manuscript

Nonalcoholic fatty liver disease (NAFLD) refers to a spectrum of pathology characterized by liver steatosis and other changes to the liver not associated with heavy alcohol use.[1] NAFLD is the most common form of liver disease globally, affecting over 80 million people in the United States.[1,2] In NAFLD, simple steatosis can progress to non-alcoholic steatohepatitis (NASH), fibrosis, cirrhosis and end-stage liver disease, and hepatocellular carcinoma.[3] Inflammation, as in NASH, promotes fibrosis.[4] However, NAFLD progression is mechanistically complex and poorly understood.[4]

The contribution of mechanical changes resulting from fat accumulation to NAFLD progression is unclear. Liver stiffening precedes fibrosis in rodent models of liver injury and promotes hepatic stellate cell myofibroblast differentiation and fibrogenesis.[5,6] The effect of steatosis itself on the mechanical properties of the liver has yet to be defined. Clinical elastography methods to determine liver stiffness reflect confounding factors such as venous pressure and have yielded contradictory results,[7] while previous ex vivo rheology studies have focused on stiffening associated with fibrosis.[8]

To determine the effect of fat accumulation on the mechanical properties of the liver as a whole, we compared steatotic and non-steatotic mouse livers ex vivo by parallel plate rheometry. We used ob/ob mice as a model of liver steatosis without fibrosis, with wild-type littermates as controls. Livers

were removed at 8, 12, and 36 weeks of age, and H&E staining demonstrated an increase in lipid

droplets in ob/ob mice at all ages compared to controls (Figure 1A). Picrosirius red staining confirmed

the absence of fibrosis in both ob/ob and control livers (Figure 1B and Supplementary Figure 1). Shear

rheometry was performed on liver cores[5] to determine the shear storage modulus G', the shear loss

modulus G", and Young's modulus E, showing that livers from ob/ob mice were significantly softer and

had less pronounced compression stiffening than those from age-matched controls (Figure 1C). Thus,

steatosis results in overall liver softening.

Figure 1. Simple steatosis induces liver softening in ob/ob mice. (A) Representative H&E stains of
livers from ob/ob mice and wild type littermates at 8w, 12w, and 36w of age. (B) Representative
picrosirius red stains of ob/ob and wt livers taken under polarized light microscopy. Scale bar = 250 μm.
(C) Shear elastic modulus G', shear loss modulus G", and Young's modulus E of ob/ob (blue) and wt
(black) livers at 8w, 12w, and 36w under varied tension and compression. Under all metrics, ob/ob livers
were significantly softer than wt livers. n ≥ 4 animals for each condition. Error bars indicate standard
error. * P < 0.005 by two-way ANOVA.

       Fat accumulation is heterogeneous in NAFLD, preferentially localizing to regions around the

central veins.[9] We therefore next asked whether localized lipid accumulation induces localized

softening.

To measure regions of fat accumulation at the submillimeter scale, we adapted a previously

described microindentation technique[10] in order to both measure stiffness and examine cell structures

such as lipid droplets within the same submillimeter regions of the liver (Figure 2A). 4',6'-diamidino-2-

phenylindole (DAPI) was applied to liver cores after maximal indentation, staining around but not under

the indenter and resulting in a ring demarcating the region where stiffness was measured. After release

of the indenter, whole mount Oil Red O staining was used to label lipids. Finally, epifluorescence

microscopy was performed to capture demarcation, lipid accumulation, and autofluorescence of the

liver, the latter of which was used to avoid large vasculature structures (Figure 2B and Supplementary

Figure 2).

97
Figure 2. Increased local lipid accumulation in steatosis is associated with local softening. (A)
Schematic of technique combining demarcation of microindented regions with visualization of tissue
features. (B) Whole mount epifluorescence microscopy images of a 12w wt liver showing DAPI (blue)
demarcating the microindented region (left), autofluorescence (green) and heterogeneous lipid
accumulation using Oil Red O (red, middle), and the three channels merged (right) showing low lipid
within the indented region (white dashed line). Scale bar 500 μm. (C) Representative fluorescent images
of indented wt and ob/ob livers with increasing lipid accumulation within the indented region. Scale bar
100 μm. (D) Scatter plots of Young's Modulus E vs. lipid accumulation within microindented regions of
ob/ob (blue) and wt (black) livers taken at 8w, 12w, and 36w. Submillimeter regions of ob/ob livers with
greater local steatosis had lower local stiffness at all tested ages. n ≥ 11 measurements from ≥ 3 animals
for each condition. * r_s < -0.7 and P < 0.005 by Spearman's rank correlation.

The amount of fat within the demarcated regions was greater in ob/ob livers than controls, and there was significant variation in different demarcated regions in ob/ob livers (Figure 2C). To examine the relationship between local stiffness and local fat accumulation, we quantified the amount of lipid staining in the demarcated microindented regions and found a significant inverse correlation between fat content and local stiffness through Spearman's rank correlation, indicating that local fat accumulation of simple steatosis results in associated local softening (Figure 2D). The 12w samples, which had the highest lipid accumulation, demonstrated a plateau (percolation threshold) at approximately 50-60% lipid, beyond which there was no further softening observed.

The finding that steatosis is associated with both global and local liver softening has significant implications for understanding the progression of early stage NAFLD, given that stiffening is a mechanical stimulus for the development of fibrosis. In the absence of bulk stiffening, locally soft areas are in particular contrast to other features of the liver microarchitecture, such as vasculature. Uncovering the relationship between these microscale mechanical features may yield mechanistic insights into the progression of NAFLD. The ability to correlate submillimeter-scale stiffness measurements with histological features can also be used to study other pathological states, such as steatohepatitis and fibrosis, and to investigate development and disease in other tissues and organ systems.

Acknowledgements

The authors thank Jessica Llewellyn, Dongning Chen, Xuechen Shi, and Jeff Byfield for training, technical support, and feedback, the Center for Molecular Studies in Digestive and Liver Diseases (P30DK050306) for histology and microscopy support, Guy Genin for thought-provoking discussions, and Yang Liu for assistance with the illustration.

References

1. Chalasani N, et al. *Hepatology*. 2018; **67**: 328-357

2. Estes C, et al. *Hepatology*. 2018; **67**: 123-133

3. Matteoni CA, et al. *Gastroenterology*. 1999; **116**: 1413-1419

4. Friedman SL, et al. *Nat Med*. 2018; **24**: 908-922

5. Perelpelyuk M, et al. *Am J Physiol Gastrointest Liver Physiol*. 2013; **304**: G605-G614

6. Olsen AL, et al. *Am J Physiol Gastrointest Liver Physiol*. 2011; **301**: G110-G118

7. Mueller S, et al. *World J Gastroenterol*. 2010; **16**: 966-972

8. Chin L, et al. Am J Physiol Gastrointest Liver Physiol. 2020; **319**: G11-G22

9. Takahashi Y, et al. *World J Gastroenterol*. 2014; **20**: 15539-15548

10. Levental I, et al. *J Phys Condens Matter*. 2010; **22**: 194120-194120

Supplementary information

Steatotic livers soften with local stiffness determined by local fat content

[Fat accumulation locally softens liver]

David Li[1,2], Paul A. Janmey[2,3,4,5,6], and Rebecca G. Wells[1,2,3,4]

Materials and Methods

Animal studies

All animal work was carried out in strict accordance with the recommendations in the Guide for the Care and Use of Laboratory Animals of the National Institutes of Health. Animal protocols were approved by the Institutional Animal Care and Use Committee of the University of Pennsylvania (protocol #804031). Ob/ob mice were obtained from the Jackson Laboratories (strain #000632). Animals were housed in a temperature-controlled environment with appropriate enrichment, ad libitum feeding of standard rodent chow and water, and 12h light/dark cycles. Euthanasia was carried out by CO_2 inhalation followed by exsanguination. Livers were harvested from ob/ob mice and wild type littermates at 8w, 12w, and 36w and stored in PBS at 4°C for up to 5h before being analyzed; rodent livers are adequately preserved under these conditions and rheological properties are maintained.[1,2]

Shear rheometry

Shear rheometry was performed as described previously.[1] Briefly, liver samples were prepared using an 8 mm punch (MP0144, Alabama R&D) with all punches taken in the same orientation from the same lobe. The height of the slices ranged from 2.0 to 3.9 mm in the uncompressed state. Samples were kept hydrated during all experiments with PBS. Parallel plate shear rheometry was carried out on a Kinexus PRO rheometer (Kinexus series, Malvern Instruments) at room temperature. Samples were attached to rheometer platforms with fibrin glue by applying 5 μL each of 20 mg/mL bovine plasma fibrinogen (341573, Sigma-Aldrich) and 100 U/mL bovine thrombin (T4648, Sigma-Aldrich) to both the top and bottom side of the sample. The upper platen was quickly lowered until 0.02 N of nominal initial

force was applied to ensure adhesive contact of the sample with the metal surfaces of the rheometer, and

the sample was allowed to sit for 10 min for the fibrin glue to polymerize fully before performing

measurements. Previous literature indicates that the fibrin glue does not affect the mechanics of the

system.[1] Mechanics were measured with a dynamic time sweep test (2% constant strain, oscillation

frequency 1rad/s, measurements taken for 120s). These measurements were carried out under the

uncompressed condition, then increasing uniaxial tension (10% and 20%), then the uncompressed

condition followed by increasing uniaxial compression (10, 15, 20, and 25%). Tension and compression

were applied by changing the gap between the platform and upper platen of the rheometer. The

following correction was applied to account for the change in cross-sectional area during testing under

the assumption that total tissue volume is conserved, where G' is the storage modulus and λ is axial

strain:

$$G'_{actual} = G'_{measured}(1 + \lambda)^2$$

A similar correction was applied to the loss modulus G'' and normal stress δ, and Young's

modulus E was determined by calculating the slope of stress (Pa) vs axial strain[1].

Microindentation and demarcation of microindented regions

Microindentation was performed as described previously[3], with additional modifications to

permit the demarcation and visualization of mesoscale features within the microindented region of the

tissue. Briefly, the microindentation device consists of a stepping motor (L4018S1204-M6, Nanotec)

attached to a μN-resolution tensiometric probe adapted from the surface tension measurement apparatus

of a Langmuir monolayer trough (MicroTrough X, Kibron Inc), consisting of a 0.510 mm diameter

blunt-ended cylindrical tungsten alloy wire hung from a digital microbalance. The force-displacement

relationship was found to behave as a Hookean spring (force linearly related to displacement - k_{probe} ~

4.5N/m) with the spring constant calibrated prior to each measurement.

Liver samples 3 mm thick were prepared from liver tissue cores obtained using an 8 mm punch,

removing the Glisson's capsule. A ring of Parafilm was prepared using a 2 mm disposable biopsy punch

and placed on top of the liver sample. Samples were then manually positioned under the free-hanging

probe, followed by downward displacement of the probe until contact occurred between the probe and

the sample in the center of the Parafilm ring. Following establishment of contact between probe and

sample, the probe was translated downward at a continuous rate of 0.0125 mm/s. These translations

resulted in decreases in measurable force from the probe, which were converted into local Young's

modulus as described previously.[3] After the probe was fully indented into the sample (as determined by

no measurable force from the probe), 0.5 µL of 100 µg/mL DAPI (D1306, Invitrogen) solution was

pipetted around the probe and allowed to selectively stain within the Parafilm ring for 15s before

washing with PBS. Multiple measurements were obtained per liver sample using this approach. After

microindentation and demarcation, liver samples were stored in PBS at 4°C for up to 5h before being

stained and imaged for lipid accumulation.

Visualization and quantification of lipid accumulation of microindented regions

Liver samples were stained for lipid using an adaptation of whole mount Oil Red O staining.[4] All

incubation steps were performed at room temperature on a shaker. Briefly, samples were incubated with

0.5% Tween 20 (#170-6531, Bio-Rad) in PBS for 15min, changing the solution every 5min, before

incubating with 0.5% Oil Red O in propylene glycol (D1306, Poly Scientific R&D Corp) for 15min.

Afterwards, the samples were washed once with Tween solution before incubating with Tween solution

for 15min, changing the solution every 5min.

Visualization of lipid accumulation within microindented regions was performed using whole

mount epifluorescence microscopy on a Nikon Eclipse Ti widefield microscope equipped with a 4x/0.31

N.A. PlanFluor dry objective lens and motorized stage. Lipid[5], DAPI demarcation, and liver

autofluorescence were visualized using Cy3, DAPI, and FITC filters respectively. Quantification of lipid

accumulation within demarcated DAPI rings was performed by thresholding the Cy3 channel in ImageJ.

Statistical analysis

Statistical significance of differences between strain-dependent shear rheometry curves (G', G'',

and E) of ob/ob mice and wild type mice were determined using two-way ANOVA.[1] Statistical

significance of the decreasing monotonic trend between increasing local lipid accumulation and local

stiffness within sub-mm regions of the ob/ob and wild type mouse liver was determined using

Spearman's rank correlation coefficient.[6] For all statistical analyses, a P value of ≤ 0.05 was considered

significant.

Supplementary Figures

226
227 **Supplementary Figure 1: Ob/ob mice do not show fibrosis.** Representative picrosirius red stains of
228 ob/ob and wt livers taken at 8w, 12w, and 36w under brightfield microscopy. Images are of the same
229 regions from Figure 1B. Scale bars = 250 μm.

Supplementary Figure 2: Whole mount fluorescent staining and microscopy shows submillimeter features within ob/ob livers. (A) Representative fluorescent images of autofluorescence (left) and intracellular lipid accumulation within hepatocytes show spatial heterogeneity of lipid accumulation at the submillimeter scale within 12w ob/ob livers. (B) Epifluorescent composite image of a 12w ob/ob liver with region of indentation from a 1 mm indenter marked with DAPI and stained for lipid accumulation (Dapi, blue; autofluorescence, green; lipid, red). * indicates an intrahepatic blood vessel within the indented region. Scale bars = 500 μm.

References

1. Perepelyuk M, et al. *PLoS One.* 2016; **11**: e0146588.

2. Tan K, et al. *J Biomech.* 2013; **46**: 1060-1066.

3. Levental I, et al. *J Phys Condens Matter.* 2010; **22**: 194120.

4. Kim S-H, et al. *PLoS One.* 2015; **10**: e0125980.

5. Koopman R, et al. *Histochem Cell Biol.* 2001; **116**: 63-68.

6. Sünderhauf A, et al. *Cell Mol Gastroenterol Hepatol.* 2021; **12**: 229-250.